Published in final edited form as:

J Pediatr. 2019 March ; 206: 105–112. doi:10.1016/j.jpeds.2018.10.053.

Exposure to Diabetes in Utero Is Associated with Earlier Pubertal Timing and Faster Pubertal Growth in the Offspring: The EPOCH Study

Christine W Hockett¹, Edward J. Bedrick², Philip Zeitler³, Tessa L. Crume¹, Stephen Daniels³, and Dana Dabelea¹

¹Department of Epidemiology, Colorado School of Public Health, Aurora, CO 80045

²Department of Epidemiology and Biostatistics, University of Arizona, Tucson, AZ 85721

³Department of Pediatrics, University of Colorado School of Medicine, Aurora, CO 80045

Abstract

Objective: To examine the associations of *in utero* exposure to maternal diabetes with surrogate measures of offspring pubertal timing (age at peak height velocity [APHV]) and speed of pubertal growth (peak height velocity [PHV]).

Study design: Data from 77 exposed and 340 unexposed youth followed from age 2 to 19 years (51% non-Hispanic white, 50% females) were analyzed using the Exploring Perinatal Outcomes among Children study, a historical prospective cohort. Maternal diabetes status was collected from obstetric records, and child heights from 2 years to current age from pediatric records. Other covariates were collected during research visits. The Superimposition by Translation and Rotation method, using height measurements (4–52 per participant), modeled APHV and PHV. Accelerated failure time analyses were used to test whether exposure to maternal diabetes was associated with younger APHV and faster PHV.

Results: Adjusting for child's sex and race/ethnicity, median APHV was reached ~3 months earlier in youth exposed to maternal diabetes compared with unexposed youth ($P < .03$). Youth exposed to maternal diabetes had a faster PHV than unexposed youth: exposed females had 10.5% greater median PHV compared with unexposed females and exposed males had a 4.0% greater median PHV compared with unexposed males ($p < 0.001$ for exposure by sex interaction).

Conclusions: Our findings provide evidence that exposure to maternal diabetes *in utero* is associated with earlier pubertal timing and faster pubertal growth. Whether earlier puberty or

Corresponding Author: Christine Hockett, PhD, University of Colorado Anschutz Medical Campus, 13001 E. 17th Place, Mail Stop B119, Historic Building 500, Room W3110, Aurora, CO 80045, Tel: 605-691-1274, Fax: 303 724-4491, Christine.Hockett@ucdenver.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Portions of this study were presented as an oral presentation at the American Diabetes Association Conference, June 10–14, 2016, New Orleans, Louisiana.

faster speed of pubertal growth mediates the association between maternal diabetes exposure and later chronic disease risk remains to be studied.

Keywords

age at peak height velocity; peak height velocity; lifecourse epidemiology; perinatal exposure; SITAR

Puberty is a complex physiological process defined by intense hormonal changes and rapid physical growth, leading to psychological and physical maturation. A population-level trend towards earlier pubertal onset has been reported over the last several decades in both boys and girls(1,2). Numerous studies have investigated the long-term effects of earlier pubertal timing on chronic disease risk in adulthood and have found an association with type 2 diabetes(3), cardiovascular disease(4), breast cancer(5), and all-cause mortality(6), specifically in females. In males, earlier pubertal timing has been associated with higher adiposity(7), increased blood pressure(8), and a higher risk of cardio-metabolic diseases in adulthood(9).

Exposure to maternal diabetes during intrauterine life has been shown to result in fetal over-nutrition and endocrine dysfunction(10,11). Exposure to maternal hyperglycemia causes the developing fetal pancreas to respond by producing additional anabolic hormones and growth factors, which promote growth(10,12). This exposure has also been linked to increased obesity and cardio-metabolic outcomes in the offspring later in life (11,13–17). It is possible that such exposure also plays a role in the programming of puberty onset and maturation. Given the population-level increase in the prevalence of diabetes during pregnancy and the decrease in the age of pubertal onset(2,18–21), understanding if exposure to maternal diabetes influences puberty in the offspring is important. However, few researchers have investigated the impact of maternal diabetes during pregnancy on offspring pubertal timing and growth velocity(22–24).

When measuring pubertal timing and speed of puberty, the gold standard is assessment of Tanner stages by a pediatric endocrinologist(25–27). The initial description of Tanner stages was by visual inspection alone, but because obesity has been significantly increasing in pre-pubertal children, palpation for thelarche (breast development) and gonadarche (testicular growth) is now considered the best method of assessment. Nonetheless, this type of assessment is difficult to obtain on a large longitudinal cohort. Another measure of pubertal timing in girls is self-reported age of menarche, which is accurate when reported prospectively and in detail. However, there is no simple measure of pubertal development in boys analogous to menarche, which has led to males being studied less often than females. An additional hallmark of puberty is a period of rapid growth(28). As puberty approaches, the growth velocity slows, known as the preadolescent dip, before it suddenly accelerates during mid-puberty, resulting in each individual experiencing a peak height velocity (PHV). For females, the age at peak height velocity (APHV) usually occurs earlier than males, typically during Tanner stage 3; males usually reach peak height velocity during Tanner stage 4(29–31).

In this study, we aimed to examine the association of exposure to maternal diabetes during pregnancy with pubertal timing, estimated using APHV, and speed of pubertal growth, estimated using PHV, using data from an ethnically diverse observational cohort study in Colorado.

Methods

The Exploring Perinatal Outcomes among Children (EPOCH) study is an historical prospective multiethnic cohort study that recruited 604 mother/child dyads in Colorado. Eligible participants were children exposed to maternal diabetes and a random sample of children not exposed. Participants were offspring of singleton pregnancies, born at a single hospital in Denver between 1992 and 2002, whose biological mothers were members of the Kaiser Permanente of Colorado Health Plan (KPCO). Children and their mothers were invited to participate in two research visits, at average ages of 10.5 years and 16.5 years. Longitudinal heights were obtained from the child's medical records from birth to their current age. For this report, we used data from 417 youth who completed both research visits and had at least 4 or more longitudinal height measurements from 2 years of age to their current age. One participant was removed from the dataset due to an extremely unrealistic outcome measure. The study was approved by the appropriate Institutional Review Boards. All participants provided written informed consent and the youth provided written assent.

Exposure to maternal diabetes was defined as a physician diagnosis of diabetes during the index pregnancy, ascertained from the KPCO perinatal database (details about exposure definition and methodology were previously published(11)). Seventy mothers were diagnosed with gestational diabetes mellitus (GDM) and seven mothers were diagnosed with type 1 diabetes mellitus prior to pregnancy. Because our research question focused on the effects of hyperglycemia during pregnancy on puberty we included all diabetes types.

Youth height and weight measurements

Longitudinal heights and weights were obtained from the child's medical records from birth to their current age. For youth with a KPCO enrollment gap or who were no longer a KPCO member, standing heights and weights were obtained from their non-KPCO providers. Longitudinal heights from birth to 2 years of age were excluded from these analyses to focus on the pubertal growth spurt rather than the postnatal growth spurt that occurs the first two years of life. The median number of measurements per participant from 2 years of age to current age was 19.

In addition, standard anthropometric measures were recorded at each research visit. Height and weight were measured in light clothing and without shoes. Weight was measured to the nearest 0.1 kg, using a portable electronic SECA scale. Height was measured to the nearest 0.1 cm, using a portable SECA stadiometer. Height and weight were measured and recorded twice, and an average was taken. Scales and stadiometers were calibrated every 2 months using standard weights for scales and an aluminum measuring rod for the stadiometer. Height and weight taken at the first research visit were used to calculate early childhood BMI. Z-scores for height and BMI were calculated using data from the National Center for Health Statistics and Centers of Disease Control and Prevention (CDC).

Pubertal timing and speed of pubertal growth

Pubertal timing and speed of pubertal growth were defined by APHV and PHV, respectively, and were estimated using longitudinal height records and Superimposition by Translation and Rotation (SITAR) growth curve analysis(32). The SITAR method uses a shape invariant spline curve and a nonlinear random-effects model to estimate an average growth curve for the entire sample and each individual's deviation from this average curve. There are 3 subject-specific measures termed size (cm), velocity (PHV), and timing (APHV); conceptually, these subject-specific measures define how much bigger or smaller each child is compared with the population average (size), how much faster and slower the child's growth velocity is compared with the population average (velocity), and how much earlier or later the child experienced peak velocity compared with the population average (timing). The models were fitted with the SITAR package in R(33). Because SITAR estimates individual measures using the underlying growth curve of the population, four separate models were fitted by sex and exposure status (exposed females, unexposed females, exposed males, unexposed males). The SITAR model allows for the flexibility in fitting the spline curves by adjusting the degrees of freedom (*df*), in doing so, the optimal models are selected by evaluating the BIC score. The lowest BIC score indicates the best fit model. Due to the variability of the longitudinal height measurements, observations with a residual greater than 4 were excluded from the analysis (n=42), this accounted for <0.7% of the total observations used. SITAR models of height by age were fitted for each group with 4 *df* for exposed females, 5 *df* for unexposed females, 6 *df* for exposed males, 6 *df* for unexposed males. Subject-specific measures size, velocity, and timing were estimated as random effects.

Other measurements

Race/ethnicity was self-reported at each research visit using the 2000 US Census base questions and categorized as Hispanic (any race), non-Hispanic white, non-Hispanic black, or non-Hispanic other. Maternal pre-pregnancy BMI (kg/m²) was calculated from the KPCO measured maternal weight before the last menstrual cycle preceding pregnancy and measured maternal height that was collected at the first research visit. Maternal level of education and total household income at the time of birth were self-reported during the first research visit. Mother's insulin use during pregnancy, delivery method, and child's birth weight were collected from the KPCO perinatal database. Pubertal development at the time of each research visit, including Tanner stage and age of menarche (females only), was ascertained by child's self-report and are used only to describe the study participants. Using a diagrammatic representation of Tanner staging adapted from Marshall and Tanner(25,26), Tanner stage based on breast development was used for females and Tanner stage based on pubic hair was used for males. Age of menarche was reported to the half year for females.

Statistical analyses

Accelerated failure time (AFT) models with a log-logistic distribution were used to evaluate the association between exposure to maternal diabetes and pubertal timing (APHV) and speed of pubertal growth (PHV). A log-logistic distribution was selected to allow modeling for a non-monotonic hazard function that allows for increased hazards during the earlier time

periods and decreases at later times. There was no right censoring of any individuals because APHV and PHV were estimated and achieved by all participants. A time-ratio (TR) was estimated using the AFT beta coefficients, which can be interpreted as the ratio of the median time-to-event for a given level of a covariant to the referent level. The associations between exposure to maternal diabetes and APHV and PHV were examined separately using three main models: an initial base model including adjustment for child's sex and race/ethnicity; a second model including an additional adjustment for potential socio-economic confounders (mother's education and total household income at birth); and a final model with an additional adjustment for early childhood BMI z-score, explored as potential mediator of hypothesized main effects. A potential effect modification by child's sex was evaluated using an interaction term (exposure status*child's sex) in the base models. If the interaction term was not significant based on a p-value <0.05, it was removed from the model. As a secondary analysis in a subset of data, we explored the effect of the mother's pre-pregnancy BMI on the relationships of maternal diabetes with pubertal timing and speed of pubertal growth. SAS statistical software, version 9.4 (SAS Institute, Inc, Cary, North Carolina) was used for all AFT analyses.

Results

Longitudinal heights (range: 4-52 height measurements per participant; average: 19 height measurements; total: 6332 height measurements) were collected on 417 children (77 exposed and 340 unexposed to maternal diabetes) and included in the analytic cohort. There were no significant differences in exposure status, child's race or sex between the analytical cohort and the larger EPOCH cohort of 604 participants. Anthropometric and demographic characteristics are summarized in Table I. Mean ($\pm$ SD) current ages of exposed and unexposed youth were 15.8 ± 1.0 and 16.8 ± 1.2 years respectively ($p < 0.001$). Of the 77 exposed youth, 44.2% were female and of the 340 unexposed youth 51.2% were female. Youth exposed to maternal diabetes were heavier at birth than youth unexposed ($3,392 \pm 498$ vs. $3,191 \pm 613$ g, respectively, $p = 0.008$). A total of 29% of mothers with diabetes were treated with insulin during their pregnancy. Achieved height z-scores were not significantly different between youth exposed to maternal diabetes compared with those unexposed for both research visits (0.35 ± 0.88 and 0.34 ± 0.99 , $p = 0.93$; 0.26 ± 0.88 and 0.17 ± 1.06 , $p = 0.47$, respectively). The mean ages at menarche were 11.9 ± 1.4 and 12.3 ± 1.3 years for exposed and unexposed females, respectively ($p = 0.20$). Mean unadjusted APHV (pubertal timing) for exposed and unexposed children were 12.1 ± 1.4 and 12.2 ± 1.4 years, respectively ($p = 0.62$). Mean unadjusted PHV (speed of pubertal growth) for exposed and unexposed children were 9.3 ± 0.4 and 8.6 ± 0.6 cm/year, respectively ($p < 0.001$).

Pubertal timing and exposure to maternal diabetes

Table 2 shows the association between exposure to maternal diabetes and pubertal timing adjusting for child's sex and race/ethnicity (model 1), as well as additional covariates—socio-economic characteristics (model 2). The interaction between exposure status and child's sex on APHV was not significant (Figure, A). After adjustment for the child's sex and race/ethnicity (model 1), the median age of pubertal timing was 2% younger, or ~3 month age difference, in exposed youth compared with unexposed youth ($\beta = -0.021$, $p =$

0.028). Males reached APHV significantly later than females ($\beta = 0.151$, $p < 0.001$) and Hispanics and non-Hispanic Blacks reached APHV significantly earlier than non-Hispanic Whites ($\beta = -0.031$, $p < 0.001$; $\beta = -0.065$, $p < 0.001$, respectively). Further adjustment for socio-economic characteristics (model 2) did not influence our findings. In model 3, when adjusting for early childhood BMI, explored here as a potential mediator, the relationship between exposure to maternal diabetes and APHV was attenuated and became nominally non-significant ($\beta = -0.017$, $p = 0.059$).

Speed of pubertal growth and exposure to maternal diabetes.

Figure, B shows the effect modification by child's sex of the association between exposure to maternal diabetes and PHV, adjusting for race/ethnicity ($p < 0.001$). Exposed females had a 10.5% greater PHV than unexposed females ($\beta = 0.10$, $p < 0.001$), and exposed males had a 4.0% greater PHV compared with unexposed males ($\beta = 0.04$, $p < 0.001$). Further adjustment for socio-economic characteristics did not influence our findings, and further adjustment for early childhood BMI, the proposed mediator, had no effect on the relationship between exposure and PHV (data not shown). In a post-hoc analysis, we evaluated the independent effect of exposure on PHV, while controlling for APHV and found that APHV was significantly associated with PHV ($\beta = -0.007$, $p < 0.001$), but the effect of exposure to maternal diabetes on PHV remained significant and only changed the beta coefficients of exposure about 2% in girls and 6% in boys.

Sub-set analyses including maternal pre-pregnancy BMI

Mother/child dyads (N 298) had available pre-pregnancy BMI data (78.9% were unexposed; 46% were females). In the subset with available data, the strength of the association between exposure to maternal diabetes and APHV was reduced ($\beta = -0.018$), and the association was not significant ($p = 0.12$) (Table 3). Further adjustment for maternal pre-pregnancy BMI had no additional effect. For speed of pubertal growth, in the subset with available data, we observed similar associations as in the larger dataset and addition of maternal pre-pregnancy BMI had no effect on the relationship between exposure to maternal diabetes and PHV (data not shown). The interaction between exposure to maternal diabetes and child's sex was still significant, where exposed females had a 10.7% greater PHV compared with unexposed females ($p < 0.001$) and exposed males had a 3.9% greater PHV compared with unexposed males ($p < 0.001$), even after adjustment for pre-pregnancy BMI.

Discussion

In an ethnically diverse population, we found that youth exposed to maternal diabetes during the intrauterine life have an earlier pubertal timing and a faster speed of pubertal growth than youth who were not exposed, independent of child's sex, race/ethnicity, and socio-economic factors. Increased BMI in early childhood largely mediates the association with earlier pubertal timing but has no effect on the association with the speed of pubertal growth. The difference in the speed of pubertal growth between exposed and unexposed is greater in female than in male offspring and is independent of pubertal timing.

Our finding of a relationship between exposure to maternal diabetes and earlier pubertal timing is consistent with the limited number of other studies that have explored this relationship(22–24). Kubo et al found that daughters of mothers who were obese prior to pregnancy, based on pre-pregnancy BMI, and had GDM during their pregnancy, had an earlier pubertal onset compared with girls of mothers who had neither condition.²⁶ Additionally, Monteilh et al found that GDM was associated with a 2 month earlier pubertal onset of pubic hair among males. The results from these studies showed that the effects of GDM on pubertal timing were similar to our findings of about a 3-month time difference(22,23). This difference in pubertal timing shifts the distribution of the population by increasing the number of adolescents who experience earlier pubertal timing, which has been shown in previous studies to be linked to an increase in chronic disease risk later in life. Grunnet et al found that female offspring of mothers with GDM were 2 times more likely to reach puberty before female offspring of mothers without GDM.(24) However, similar to our results, when controlling for offspring's own BMI the results were attenuated and became non-significant. Childhood BMI has been associated with both exposure to maternal diabetes and pubertal timing(1,11,34), including in our cohort(11), and it is thus a likely mediator of this relationship. In our study early childhood BMI seemed to account for most of the association between exposure to maternal diabetes and earlier pubertal timing, but others(23) showed that the relationship still remains significant after adjustment, suggesting potential additional pathways by which exposure to maternal diabetes results in earlier pubertal timing. Further research exploring potential mechanisms by which *in utero* exposure to maternal diabetes may result in earlier pubertal timing is warranted.

Looking at a different measure of pubertal timing, age of menarche for females only, our study found a non-significant effect of exposure to maternal diabetes during pregnancy on pubertal timing (results not shown). However, this variable was collected to the half-year in our study (i.e. 12.0 years at age of menarche or 12.5 years at age of menarche) and, therefore, may have not been sensitive enough to detect an effect of exposure to maternal diabetes on menarcheal age.

During normal pubertal development, maturation of the hypothalamic-pituitary-gonadal (HPG) axis, along with increased adrenal androgen and growth hormone secretion, drive achievement of pubertal milestones and growth. Previous animal and human studies have shown that maternal diabetes and obesity are associated with offspring adiposity and metabolic dysfunction(11,13–17,35). In a previous report from our cohort, we found that adolescents exposed to diabetes *in utero* had significantly higher fasting insulin levels, compared with unexposed adolescents(35). Thus, it is plausible that hyperinsulinemia seen in exposed offspring may permanently alter the HPG axis and subsequent sex hormone secretion (36). Sex hormone alterations may explain the sex differences we observed in the association between exposure to maternal diabetes and speed of pubertal growth. Also, researchers have shown that pubertal timing is related to physical growth (37,38), so we explored whether our results of exposure to maternal diabetes being associated with PHV may be explained by the earlier APHV. However, we found that the effect of exposure to maternal diabetes on PHV remained significant even after controlling for APHV, suggesting that the observed effect on the speed of pubertal growth is not explained by an earlier

pubertal timing. Further study of these potential mechanisms is important, as linear growth is an important biomarker of a child's development and overall health.

The difference in the size of the association between maternal diabetes and pubertal timing vs. pubertal growth observed in our study may be related to the multiple hormonal pathways associated with pubertal onset and maturation. Pubertal timing is primarily dependent on the reemergence of gonadotropin-releasing hormone (GnRH) secretion, which is the initial step in the HPG axis, while pubertal growth is not only dependent on the HPG axis, but also the GH-IGF-1 axis and the interaction of these axes(39,40). It is likely that exposure to maternal diabetes influences these axes and their associated hormones differently, resulting in the somewhat larger effect of maternal diabetes exposure on pubertal speed vs pubertal timing.

We also explored the effect of additional adjustment for maternal pre-pregnancy BMI on the relationship between exposure to maternal diabetes and pubertal timing and speed of pubertal growth in the offspring, albeit only in a subset with available data. Previous studies looking at the effect of maternal diabetes on puberty measures have not explored the influence pre-pregnancy BMI. We were only able to explore this in a subset, in which, due to the smaller sample size, the relationship between exposure to maternal diabetes *in utero* and pubertal timing was not significant, even without controlling for maternal pre-pregnancy BMI. Additional adjustment for maternal pre-pregnancy BMI had no further effect. Because pre-pregnancy BMI is highly related to maternal diabetes, it may be part of the fetal overnutrition pathway(41), in which case this modeling may also result in over adjustment. Nevertheless, adjustment for maternal pre-pregnancy BMI had no effect on the association between exposure to maternal diabetes and speed of pubertal growth.

Although our findings represent an important addition to the existing evidence regarding risk factors for earlier puberty, further investigation into potentially responsible mediators, specifically hormones related to the GH-IGF-1 and HPG axes, is needed. Additionally, further study is required to understand the role that puberty may play on the relationship between exposure to diabetes *in utero* and subsequent disease risk in the offspring. It is possible that puberty may be another critical or sensitive period in the lifecycle when the effects of *in utero* exposure to maternal diabetes may be enhanced via biological programming.

This prospective study is one of the largest multi-ethnic cohorts to investigate the relationships between maternal diabetes and offspring pubertal outcomes. Estimating accurately the timing and intensity of puberty is challenging and the current methods, such as using height as a measure of pubertal timing or estimating the mean age at a given Tanner stage or the transition from one stage to another, have some variability in their estimates. However, in this study we focused on using height as a measure of pubertal timing and intensity because this is a valid measure of pubertal timing and we had longitudinal height measures on individuals from 6 to 19 years of age. Also, we wanted to be able to compare these pubertal measures between boys and girls, which is not possible using Tanner stages. The height measurements used in calculating APHV and PHV were not obtained in a standardized fashion, which may have introduced some variability and decreased the precision of the APHV and PHV estimates, attenuating the associations of interest.

Additionally, we have not yet followed the cohort long enough to have data on achieved adult height, and therefore we cannot address the question of whether the observed difference in the speed of pubertal growth according to exposure status will affect final achieved height. Because achieved height at 2 times points during childhood and adolescence was not affected, we speculate that exposure to diabetes *in utero* may not affect achieved adult height but rather the shape of childhood growth trajectory. Unfortunately, our study was unable to control for genetic factors associated with pubertal timing and growth. However, this would not bias our results unless the genes in question were also associated with maternal diabetes, which has not been shown to date. Further research is needed to examine whether the relationship between exposure to maternal diabetes and pubertal development mediates, at least in part, the increased risk of obesity and cardio-metabolic outcomes associated with this exposure. Such findings may then provide evidence for life-stage targeted interventions aimed at reducing or halting the transgenerational vicious cycle of diabetes and obesity.

In summary, these novel findings provide evidence that exposure to diabetes during the intrauterine life may affect pubertal development and growth in the offspring and support the hypothesis that perinatal exposures are among the multiple contributors to the general trend of earlier puberty seen over the last couple of decades.

Acknowledgements.

We thank the EPOCH participants, their families, and the study staff for their dedicated contribution to this work. We also thank Dr Tim Cole (UCL Institute of Child Health, London, UK) for his guidance in using the SITAR method.

Supported by the NIH, National Institute of Diabetes, Digestive, and Kidney Diseases (R01 DK068001). Its contents are the authors' sole responsibility and do not necessarily represent official NIH views. The funders had no role in the design, conduct, or reporting of this work. The authors declare no conflicts of interest.

Abbreviations:

APHV	age at peak height velocity
PHV	peak height velocity
SITAR	Superimposition by Translation and Rotation
KPCO	Kaiser Permanente of Colorado
EPOCH	Early Perinatal Outcomes in Children
DM	diabetes mellitus
GDM	gestational diabetes mellitus
HPG	hypothalamic-pituitary-gonadal

References

1. Biro FM, Khoury P, Morrison JA. Influence of obesity on timing of puberty. *Int J Androl* 2006 29:272–7; discussion 286–290. [PubMed: 16371114]

2. Euling SY, Herman-Giddens ME, Lee PA, Selevan SG, Juul A, Sørensen TIA, et al. Examination of US puberty-timing data from 1940 to 1994 for secular trends: panel findings. *Pediatrics*. 2008 2;121 Suppl 3:S172–191. [PubMed: 18245511]
3. Elks CE, Ong KK, Scott RA, van der Schouw YT, Brand JS, Wark PA, et al. Age at menarche and type 2 diabetes risk: the EPIC-InterAct study. *Diabetes Care*. 2013 11;36:3526–34. [PubMed: 24159179]
4. Prentice P, Viner RM. Pubertal timing and adult obesity and cardiometabolic risk in women and men: a systematic review and meta-analysis. *Int J Obes* 2005. 2013 8;37:1036–43.
5. Bodicoat DH, Schoemaker MJ, Jones ME, McFadden E, Griffin J, Ashworth A, et al. Timing of pubertal stages and breast cancer risk: the Breakthrough Generations Study. *Breast Cancer Res* 2014;16:R18. [PubMed: 24495528]
6. Charalampopoulos D, McLoughlin A, Elks CE, Ong KK. Age at Menarche and Risks of All-Cause and Cardiovascular Death: A Systematic Review and Meta-Analysis. *Am J Epidemiol*. 2014 7 1;180:29–40. [PubMed: 24920784]
7. Ong KK, Bann D, Wills AK, Ward K, Adams JE, Hardy R, et al. Timing of voice breaking in males associated with growth and weight gain across the life course. *J Clin Endocrinol Metab* 2012 8;97:2844–52. [PubMed: 22654120]
8. Hardy R, Kuh D, Whincup PH, Wadsworth ME. Age at puberty and adult blood pressure and body size in a British birth cohort study. *J Hypertens*. 2006 1;24:59–66. [PubMed: 16331102]
9. Day FR, Elks CE, Murray A, Ong KK, Perry JRB. Puberty timing associated with diabetes, cardiovascular disease and also diverse health outcomes in men and women: the UK Biobank study. *Sci Rep* 2015 6 18;5:11208. [PubMed: 26084728]
10. Dabelea D The Predisposition to Obesity and Diabetes in Offspring of Diabetic Mothers. *Diabetes Care*. 2007 7 1;30:S169–74. [PubMed: 17596467]
11. Crume TL, Ogden L, West NA, Vehik KS, Scherzinger A, Daniels S, et al. Association of exposure to diabetes in utero with adiposity and fat distribution in a multiethnic population of youth: the Exploring Perinatal Outcomes among Children (EPOCH) Study. *Diabetologia*. 2011 1;54:87–92. [PubMed: 20953862]
12. Pirc LK, Owens JA, Crowther CA, Willson K, Blasio MJD, Robinson JS. Mild gestational diabetes in pregnancy and the adipoinular axis in babies born to mothers in the ACHOIS randomised controlled trial. *BMC Pediatr*. 2007 4 12;7:18. [PubMed: 17430602]
13. Dabelea D, Hanson RL, Lindsay RS, Pettitt DJ, Imperatore G, Gabir MM, et al. Intrauterine exposure to diabetes conveys risks for type 2 diabetes and obesity: a study of discordant sibships. *Diabetes*. 2000 12;49:2208–11. [PubMed: 11118027]
14. West NA, Crume TL, Maligie MA, Dabelea D. Cardiovascular risk factors in children exposed to maternal diabetes in utero. *Diabetologia*. 2011 3;54:504–7. [PubMed: 21153896]
15. Garcia-Vargas L, Addison SS, Nistala R, Kurukulasuriya D, Sowers JR. Gestational Diabetes and the Offspring: Implications in the Development of the Cardiorenal Metabolic Syndrome in Offspring. *Cardiorenal Med*. 2012 5;2:134–42. [PubMed: 22851962]
16. Boney CM, Verma A, Tucker R, Vohr BR. Metabolic syndrome in childhood: association with birth weight, maternal obesity, and gestational diabetes mellitus. *Pediatrics*. 2005 3;115:e290–296. [PubMed: 15741354]
17. Lawlor DA, Fraser A, Lindsay RS, Ness A, Dabelea D, Catalano P, et al. Association of existing diabetes, gestational diabetes and glycosuria in pregnancy with macrosomia and offspring body mass index, waist and fat mass in later childhood: findings from a prospective pregnancy cohort. *Diabetologia*. 2010 1;53:89–97. [PubMed: 19841891]
18. DeSisto CL. Prevalence Estimates of Gestational Diabetes Mellitus in the United States, Pregnancy Risk Assessment Monitoring System (PRAMS), 2007–2010 *Prev Chronic Dis* [Internet]. 2014 [cited 2016 11 23];11 Available from: https://www.cdc.gov/pcd/issues/2014/13_0415.htm
19. Getahun D, Nath C, Ananth CV, Chavez MR, Smulian JC. Gestational diabetes in the United States: temporal trends 1989 through 2004. *Am J Obstet Gynecol* 2008 5;198:525.e1–5. [PubMed: 18279822]
20. Ferrara A Increasing Prevalence of Gestational Diabetes Mellitus: A public health perspective. *Diabetes Care*. 2007 7 1;30:S141–6. [PubMed: 17596462]

21. Biro FM, Huang B, Crawford PB, Lucky AW, Striegel-Moore R, Barton BA, et al. Pubertal correlates in black and white girls. *J Pediatr* 2006 2;148:234–40. [PubMed: 16492435]
22. Monteilh C, Kieszak S, Flanders WD, Maisonet M, Rubin C, Holmes AK, et al. Timing of maturation and predictors of Tanner stage transitions in boys enrolled in a contemporary British cohort. *Paediatr Perinat Epidemiol* 2011 1;25:75–87. [PubMed: 21133972]
23. Kubo A, Ferrara A, Laurent CA, Windham GC, Greenspan LC, Deardorff J, et al. Associations Between Maternal Pregravid Obesity and Gestational Diabetes and the Timing of Pubarche in Daughters. *Am J Epidemiol* 2016 7 1;184:7–14. [PubMed: 27268032]
24. Grunnet LG, Hansen S, Hjort L, Madsen CM, Kampmann FB, Thuesen ACB, et al. Adiposity, Dysmetabolic Traits, and Earlier Onset of Female Puberty in Adolescent Offspring of Women With Gestational Diabetes Mellitus: A Clinical Study Within the Danish National Birth Cohort. *Diabetes Care*. 2017 12;40:1746–55. [PubMed: 29038315]
25. Marshall WA, Tanner JM. Variations in the Pattern of Pubertal Changes in Boys. *Arch Dis Child* 1970 2 1;45:13–23. [PubMed: 5440182]
26. Marshall WA, Tanner JM. Variations in pattern of pubertal changes in girls. *Arch Dis Child* 1969 6 1;44:291–303. [PubMed: 5785179]
27. Dorn LD, Dahl RE, Woodward HR, Biro F. Defining the boundaries of early adolescence: A user's guide to assessing pubertal status and pubertal timing in research with adolescents. *Appl Dev Sci* 2006;10:30–56.
28. Gluckman PD, Hanson MA. Changing times: the evolution of puberty. *Mol Cell Endocrinol* 2006 7 25;254–255:26–31.
29. Rogol AD, Roemmich JN, Clark PA. Growth at puberty. *J Adolesc Health Off Publ Soc Adolesc Med* 2002 12;31:192–200.
30. De Leonibus C, Marcovecchio ML, Chiavaroli V, de Giorgis T, Chiarelli F, Mohn A. Timing of puberty and physical growth in obese children: a longitudinal study in boys and girls. *Pediatr Obes*. 2014 Aug;9:292–9.
31. Cole TJ, Pan H, Butler GE. A mixed effects model to estimate timing and intensity of pubertal growth from height and secondary sexual characteristics. *Ann Hum Biol* 2014 2;41:76–83. [PubMed: 24313626]
32. Cole TJ, Donaldson MDC, Ben-Shlomo Y. SITAR--a useful instrument for growth curve analysis. *Int J Epidemiol* 2010 12;39:1558–66. [PubMed: 20647267]
33. sitar v1.0.8 | RDocumentation [Internet]. [cited 2016 11 25]. Available from: <https://www.rdocumentation.org/packages/sitar/versions/1.0.8>
34. Kubo A, Ferrara A, Windham GC, Greenspan LC, Deardorff J, Hiatt RA, et al. Maternal Hyperglycemia During Pregnancy Predicts Adiposity of the Offspring. *Diabetes Care*. 2014 11 1;37:2996–3002. [PubMed: 25150158]
35. Sauder K, Hockett C, Ringham B, Glueck D, Dabelea D. Intrauterine exposure to maternal diabetes or obesity and offspring insulin resistance and β -cell function: the EPOCH study. In San Diego, CA; 2017.
36. Bordini B, Rosenfield RL. Normal Pubertal Development: Part I: The Endocrine Basis of Puberty. *Pediatr Rev*. 2011 6 1;32:223–9.
37. Huang B, Biro FM, Dorn L. Determination of Relative Timing of Pubertal Maturation through Ordinal Logistic Modeling: Evaluation of Growth and Timing Parameters. *J Adolesc Health Off Publ Soc Adolesc Med* 2009 10;45:383–8.
38. Llop-Viñolas D, Vizmanos B, Closa Monasterolo R, Escribano Subías J, Fernández-Ballart JD, Martí-Henneberg C. Onset of puberty at eight years of age in girls determines a specific tempo of puberty but does not affect adult height. *Acta Paediatr Oslo Nor* 1992 2004 7;93:874–9.
39. Casazza K, Hanks LJ, Alvarez JA. Role of Various Cytokines and Growth Factors in Pubertal Development [Internet]. Vol. 55 Karger Publishers; 2010 [cited 2016 11 23]. Available from: <http://www.karger.com/Article/Abstract/321969>
40. Laron Z Insulin-like growth factor 1 (IGF-1): a growth hormone. *Mol Pathol*. 2001 10;54:311–6. [PubMed: 11577173]
41. Gillman MW, Rifas-Shiman S, Berkey CS, Field AE, Colditz GA. Maternal gestational diabetes, birth weight, and adolescent obesity. *Pediatrics*. 2003 3;111:e221–226. [PubMed: 12612275]

Figure.

The sex difference in the association of exposure to diabetes *in utero* with (a) offspring age at peak height velocity and (b) offspring peak height velocity using a log-logistic AFT model, adjusting for child's race (model 1).

Table 1.

Characteristics of youth and their mothers by maternal diabetes exposure

Variable	Unexposed to DM (n=340)				Exposed to DM (n=77)				p-value
	N	%	Mean (SD)		N	%	Mean (SD)		
Current age (years)			16.8 (1.2)		77		15.8 (1.0)		<0.001
Sex (female)	174	51.2			34	44.2	34 (44.2)		0.27
Child's birthweight (g)			3191 (613)				3392 (498)		0.008
Child's height z-score at visit 1 (cm)			0.34 (0.99)				0.35 (0.88)		0.93
Child's height z-score at visit 2 (cm)			0.17 (1.06)				0.26 (0.88)		0.47
Race/Ethnicity									0.18
Non-Hispanic White	165	48.5			48	62.3			
Hispanic	126	37.1			22	28.6			
Non-Hispanic Black	29	8.5			4	5.2			
Non-Hispanic Other	20	5.9			3	3.9			
Self-reported Tanner stage at visit 2 ^a									0.71
Pre-pubertal (TTS-1)	0	0			0	0			
Pubertal (TTS-2)	3	0.9			0	0			
Pubertal (TTS-3)	17	5			4	5.2			
Pubertal (TTS-4)	133	39.4			32	41.6			
Pubertal (TTS-5)	185	54.7			41	53.3			
Child's age at menarche (female only)			12.3 (1.3)				11.9 (1.4)		0.2
Mother's pre-pregnancy BMI (kg/m ²)	235		25.6 (6.0)		63		26.9 (5.7)		0.12
Self-reported household income at birth									0.51
\$49,999	155	45.7	155 (45.7)		32	41.6			
\$50,000	184	54.3	184 (54.3)		45	58.4			
Mother's self-reported education at birth									0.83
High School	61	17.9			13	16.9			
> High School	279	82.1			64	82.1			
Mother's insulin use during pregnancy (yes)					22	29			
Age at peak height velocity (yrs)			12.21 (1.35)				12.12 (1.38)		0.62
Peak height velocity (cm/yr)			8.64 (0.63)				9.30 (0.39)		<0.001

SL
Females breast TS; Males pubic hair TS

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 2.

The association between exposure to maternal diabetes and age at peak height velocity.

Variables	Model 1 ^a			Model 2 ^b			Model 3 ^c		
	TR	β	95% CI	p-value	TR	β	95% CI	p-value	TR
Exposure status									
Unexposed to DM	1	Referent	Referent	Referent	1	Referent	Referent	Referent	Referent
Exposed to DM	0.98	-0.021	(-0.04, -0.002)	0.028	0.98	-0.022	(-0.041, -0.003)	0.025	0.98
Race/Ethnicity									
Non-Hispanic Whites	1	Referent	Referent	Referent	1	Referent	Referent	Referent	Referent
Non-Hispanic Black	0.94	-0.065	(-0.094, -0.036)	<0.001	0.94	-0.061	(-0.091, -0.032)	<0.001	0.96
Hispanics	0.97	-0.031	(-0.047, -0.015)	<0.001	0.97	-0.027	(-0.044, -0.01)	0.002	0.98
Non-Hispanic other	0.97	-0.03	(-0.064, 0.005)	0.09	0.97	-0.027	(-0.062, 0.007)	0.12	0.98
Sex									
Females	1	Referent	Referent	Referent	1	Referent	Referent	Referent	Referent
Males	1.16	0.151	(0.136, 0.165)	<0.001	1.16	0.15	(0.135, 0.165)	<0.001	1.17
Household income									
<\$50,000/year					1	Referent	Referent	Referent	Referent
\$50,000/year					1.01	0.01	(-0.006, 0.03)	0.23	1
Maternal education									
High school					1	Referent	Referent	Referent	Referent
Any college					1.01	0.010	(-0.01, 0.03)	0.57	1.01
BMI z-score								0.97	-0.026

^aModel 1 adjusted for child's race/ethnicity and sex

^bModel 2 adjusted for child's race/ethnicity, sex, household income and maternal education

^cModel 3 adjusted for child's race/ethnicity, sex, household income, maternal education and child's BMI z-score at visit 1

The effect of adjustment for pre-pregnancy BMI on the association between exposure to maternal diabetes *in utero* and age at peak height in a subset (n=298).

Table 3.

Variables	Model 1 ^a				Model 2 ^b			
	TR	β	95% CI	p-value	TR	β	95% CI	p-value
Exposure status								
Unexposed to DM	1	Referent	Referent	Referent	1	Referent	Referent	Referent
Exposed to DM	0.98	-0.018	(-0.041, 0.005)	0.12	0.99	-0.012	(-0.035, 0.01)	0.28
Race/Ethnicity								
Non-Hispanic Whites	1	Referent	Referent	Referent	1	Referent	Referent	Referent
Non-Hispanic Black	0.94	-0.059	(-0.094, -0.023)	0.001	0.96	-0.046	(-0.082, -0.011)	0.01
Hispanics	0.97	-0.029	(-0.05, -0.009)	0.005	0.98	-0.023	(-0.043, 0.002)	0.03
Non-Hispanic other	0.99	-0.012	(-0.054, 0.03)	0.58	0.99	-0.006	(-0.048, 0.035)	0.76
Sex								
Females	1	Referent	Referent	Referent	1	Referent	Referent	Referent
Males	1.17	0.154	(0.135, 0.173)	<0.001	1.17	0.156	(0.138, 0.174)	<0.001
Pre-pregnancy BMI					1	-0.003	(-0.004, -0.001)	<0.001

^a Adjusted for child's race/ethnicity, and sex

^b Adjusted for child's race/ethnicity, sex, and maternal pre-pregnancy BMI