Quantum chemical calculation and binding modes of H1R; a combined study of molecular docking and DFT for suggesting therapeutically potent H1R antagonist

Yasir Mohamed Riza¹ · Md. Rimon Parves¹ · Fahmida Alam Tithi¹ · Sanjida Alam¹

Received: 14 April 2018 / Accepted: 15 February 2019 / Published online: 25 February 2019
© Springer-Verlag GmbH Germany, part of Springer Nature 2019

Abstract

Histamine-1 receptor (H1R) belongs to the family of rhodopsin-like G-protein-coupled receptors expressed in cells that mediate allergies and other pathophysiological diseases. For alleviation of allergic symptoms, H1R antagonists are therapeutic drugs; of which the most frequently prescribed are second generation drugs, such as; Cetirizine, Loratadine, Hydroxyzine, Desloratadine, Bepotastine, Acrivastine and Rupatadine. To understand their potency, binding affinity and interaction; we have employed molecular docking and quantum chemical study such as; Induced-fit docking and calculation of quantum chemical descriptors. This study also introduces the binding site characterization of H1R, with its known antagonists and Curcumin (our proposed alternative H1R antagonist); useful for future drug target site. The interactive binding site residues of H1R are found to be; Lys-191, Tyr-108, Asp-107, Tyr-100, Lys-179, Lys-191, Thr-194, Trp-428, Phe-432, Tyr-458, Hie-450, with most of these shown to be inhibited by naturally-occurring compound curcumin. Amongst the FDA approved drugs, Hydroxyzine showed best ligand binding affinity, calculated as -141.491 kcal/mol and naturally occurring compound, Curcumin showed binding affinity of -87.046 kcal/mol. The known antagonists of H1R has been used for hypothesizing curcumin as naturally occurring lead compound for the target using accurate molecular docking simulation study. Though the pharmacological action of known inhibitor is already established, they could differ from their reactivity, which we have also focused in our study for predicting drug reactivity.

Keywords H1R antagonists · Binding site characterization · QM/MM · Density functional theory · Induced fit docking · MM-GBSA

Introduction

Histamine receptors are a class of G protein coupled receptors that have been classified as; H1R, H2R, H3R, and H4R (Hill et al. 1997) among which H1R is responsible for causing allergic inflammation, Immune modulation and regulation of T cell function and auto-immunity (Jutel et al.

2002; Ma et al. 2002; Musio et al. 2006; Pedotti et al. 2003; Schaefer et al. 1999). Histamine influences its inflammation reaction by regulating its own receptor expression on endothelial cells (Schaefer et al. 1999). The signal transduction of H1R is typical for, and convergent with, that of other $G\alpha_{q/11}$ protein-coupled receptors (Bakker et al. 2004; Brown et al. 2002). H1R activates phospholipase C, resulting in the catalysis of phosphatidylinositol-4,5-bisphosphate to inositol-1,4,5-trisphosphate and diacylglycerol (DAG) which induces calcium ion release which is a trigger for symptoms of allergic response, vascular permeability and smooth muscle contraction. H1R also activates NF- κ B in both a constitutive and agonist-dependent manner (Bakker et al. 2001) and induces granulocyte macrophage colony-stimulating factor (GM-CSF), interleukin (IL)-8, IL-6 and IL-16 release from airway epithelial cells (Arima et al. 1999; Jeannin et al. 1994; Kanda and Watanabe 2004). H1R is effectively associated with the expression of some chemokines, and cytokines

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s40203-019-0050-3>) contains supplementary material, which is available to authorized users.

✉ Yasir Mohamed Riza
yasirmohamedriza@gmail.com

¹ Department of Biochemistry and Biotechnology, Faculty of Basic Medical and Pharmaceutical Sciences, University of Science and Technology Chittagong (USTC), Foy's Lake, Khushi-4202, Chittagong, Bangladesh

that is involved in inflammation (Matsubara et al. 2005; Caron et al. 2001). The CC chemokines, monocyte chemoattractant protein (MCP) 1 and 3, RANTES and Eotaxin induced by Histamine H1 receptor suggests a prolonged inflammatory cycle, in the case of allergic rhinitis, between the cell that releases the histamine and their intensified migration to nasal mucosa (Fujikura et al. 2001). H1R plays a primary role in allergic rhinitis, urticaria, anaphylaxis and to a lesser degree, asthma (White 1990) and it also mediates inflammatory responses in human keratinocytes (Giustizieri et al. 2004). When the immune system overreacts to normally harmless substances in the environment, allergic reaction occurs as a result. This hypersensitivity occurs in various forms including atopic dermatitis, asthma, allergic rhinitis, and food allergies. The occurrence of allergy-related disorders has been increasing yearly. Especially in the case of food allergies where it has been estimated to have affected 4% of the adult population whilst 6% of children population (Venter et al. 2008). Many drugs have been developed such as immunosuppressants, antihistamines, and steroids to treat allergic disorders (Conen et al. 2013; Kim et al. 2013; Wong et al. 2013). Antihistamines effectively inhibit histamine-mediated symptoms, such as sneezing and nasal discharge, because they can block histamine (H1) receptors (Tarnasky and Van 1990; Wang et al. 2001). However, for alleviating H1 mediated disease symptoms, second generation H1R antagonist are more commonly used due to its selectivity and relatively lessened sedative effect. Thusly, we have selected the most popular and most commonly used second generation H1R antagonists to comparatively analyze and understand their binding pattern mechanisms and their quantum chemical attributes, to ascertain which of these commercial drugs is ideal for alleviating H1 mediated disease symptoms. However, although these drugs are effective, when taken for long periods, these drugs may have adverse effects such as cataracts, growth retardation, osteoporosis, diabetes, and hypertension (de Benedictis et al. 2012; Elphick and Southern 2012). Due to these potential problems, natural products may provide an alternative to effectively treat allergies with fewer adverse effects. *Curcuma longa* Linn (*Zingiberaceae*), has a rhizome called turmeric (given the name curry spice by the British), which has been widely consumed as a seasoning and used therapeutically to treat inflammatory disorders of respiratory and gastrointestinal systems in predominantly Asian countries such as India, Japan, Korea and China (Gilani et al. 2005; Srinivasan 1953). Turmeric has also been studied expansively for its pharmacological activities such as its anticancer, antioxidant, anti-microbial, neuroprotective and anti-inflammatory effects (Chandra and Gupta 1972; Lutomski et al. 1974; Rajakrishnan et al. 1999; Toda et al. 1985). Curcumin, the main active ingredient or compound, which is obtained from turmeric has been also claimed to be anti-inflammatory and antioxidant agent (Aggarwal et al.

2007; Balasubramanyam et al. 2003). Curcumin's immunoregulatory activities has been extensively studied ever since they were first reported in 1999 (Antony et al. 1999). For example, curcumin attenuated allergic airway inflammation by regulating the balance of CD4⁺CD25⁺ regulatory T cells (Tregs)/T-helper (Th) in ovalbumin (OVA)-sensitized mice (Ma et al. 2013). In allergic disorders, both turmeric and curcumin were found to ameliorate allergic contact dermatitis allergic rhinitis, food allergy, allergic conjunctivitis and asthma (Chung et al. 2012; Oh et al. 2011; Ringman et al. 2005; Thakare et al. 2013; Thompson and Tan 2006) through regulation of immune responses mediated by mast cells, IgE responses and Th2 responses, A cognate of T cell kinase ZAP70 in mast cells, Syk kinase, plays a crucial role in the IgE-dependent activation of mast cells. Syk kinase-dependent signaling events is inhibited by Curcumin in mast cells, which would result in broad beneficial effects, due to the suppression of multiple inflammatory mediators, which each play overlapping but distinct roles in the inflammatory response and thus might contribute to its antiallergic activity (Lee et al. 2008). Thus, we have selected the active main ingredient of Turmeric spice, Curcumin, as an alternative to the commercially produced second generation H1-receptor antagonists, for treatment of allergic symptoms.

Method and materials

Ligand preparations

Selected drugs were retrieved from Drug Bank (Wishart et al. 2018) and prepared using LigPrep wizard of Schrödinger suite (LigPrep, Schrödinger, LLC, New York, NY, 2017). Ionization state of these small molecules were generated at pH 7.0 ± 2.0 by using Epik (Shelley et al. 2007) for proper enumeration of tautomer and protonation states for better understanding in biological condition. Up to 32 possible stereoisomers per ligand were retained. By assigning bond order and bond angles, all the small molecules were minimized using OPLS3 (Harder et al. 2016) force field.

Protein preparation

The crystal structure of Histamine 1 receptor (PDB ID: 3RZE) was retrieved from Protein Data Bank (Berman et al. 2000) and prepared in protein preparation wizard (Madhavi Sastry et al. 2013) of Schrödinger maestro 11.2. Firstly, the protein was preprocessed by assigning bond orders, creating zero order bonds to metals and disulfide bonds. After removing water and adding Hydrogen bonds, the protein was preprocessed for review and modification, to be refined. By assigning H bond and keeping sample water orientation,

the protonation states were optimized at neutral pH. Finally, a restrained minimization of the protein structure was performed using the default constraint of 0.30 Å RMSD and OPLS3 force field. Afterwards, the protein quality was checked in Ramachandran plot (Lovell et al. 2003) in protein preparation wizard.

Receptor grid generation and induced fit docking

Receptor grid was generated by specifying co-crystallized ligand. The default parameters were used to generate the grid for ligand docking. Glide induced-fit docking protocol (Sherman et al. 2006) was used for extensive accuracy of ligand docking. Prepared optimized ligands were introduced to Induced fit docking using the default parameters.

MM-GBSA calculation

The docked pose view files were introduced to Prime program (Prime, Schrödinger, LLC, New York, NY, 2017) for calculating the binding free energy employing MM-GBSA approach which is a physics-based method that computes the force field energies in implicit solvent of the bound and unbound molecules involved in the binding process. The MM-GBSA binding energy is calculated by using the Eq. 1, where E denotes energy and includes terms such as protein–ligand van der Waals contacts, electrostatic interactions, ligand de solvation, and internal strain (ligand and protein) energies, using VSGB2.0 solvation model (Li et al. 2011) and OPLS3 force field. Residues within 5.0 Å of the ligand were allowed to relax throughout the minimization (minimization of all atoms in each residue) of the complex, keeping the rest of the structure fixed.

$$\Delta G_{\text{binding}} = E_{\text{complex}} - E_{\text{protein}} - E_{\text{ligand}} \quad (1)$$

Quantum chemical calculations

Quantum chemical calculations have been performed by employing density functional theory (DFT) with B3LYP (Lee et al. 1988) method using 6-31G [d, p] basis set (Eas-ton et al. 1996) in the Gaussian program (Frisch et al. 2016).

Ligand optimization, Electronic energies, Enthalpy, Gibbs free energy of each compound were also investigated. Besides this, the dipole moments and drug polarizability has also been calculated. For observing molecular reactivity and stability, the frontier HOMOs, LUMOs and gap energies were calculated (Hoque et al. 2015; Parr and Zhou 1993).

P450 site of metabolism

Only the naturally-occurring compound curcumin (ID: DB11672), was undertaken to predict Site of metabolism, Intrinsic reactivity and Fe accessibility. The P450 Site of Metabolism (SOM) of selected compound was determined by induced fit docking on CYP2D6 isoform using the P450 Site of Metabolism module of Schrodinger Suite. Only intrinsic reactivity was determined for CYP3A4 isoform, based on ligand-based models to predict how a druglike compound is metabolized for CYP 3A4. Haloperidol (ID: DB00502), an FDA approved small molecule cytochrome substrate, was used in the role of a control.

Results and discussion

Quantum chemical calculation

Nowadays, quantum mechanics has reliable application in the study of electronic structure, wave function, modelling-simulation, vibrational spectra, IR spectra, ground state studies of an individual atom, NMR prediction and so on. Therefore, the frontier molecular descriptors such as HOMO (Highest occupied molecular orbital) and LUMO (lowest unoccupied molecular orbital), Gap energy, Hardness, Softness, can also be studied by employing quantum chemistry. The electronic energy, enthalpy and Gibbs free energy, along with the stoichiometry of selected drugs are reported in Table 1. HOMO, LUMO, Gap energy, Dipole moment of those drugs are reported in Table 2. Optimized structure, HOMO and LUMO plots are given in Figs. 1 and 2, respectively. The lowest gap energy was found for DB09488

Table 1 Electronic energy, enthalpy, Gibbs free energy in hartree

Drug ID	Drug name	Stoichiometry	Electronic energy (eV)	Enthalpy	Gibbs free energy
DB00341	Cetirizine	$C_{21}H_{25}ClN_2O_3$	−1670.674	−1610.216	−1610.302
DB00455	Loratadine	$C_{22}H_{23}ClN_2O_2$	−1572.396	−1571.959	−1572.04
DB00557	Hydroxyzine	$C_{21}H_{27}ClN_2O_2$	−1536.634	−1536.158	−1536.243
DB00967	Desloratadine	$C_{19}H_{19}ClN_2$	−1305.18	−1304.821	−1304.885
DB04890	Bepotastine	$C_{21}H_{25}ClN_2O_3$	−1610.705	−1610.246	−1610.334
DB09488	Acrivastine	$C_{22}H_{24}N_2O_2$	−1112.668	−1112.225	1112.306
DB11614	Rupatadine	$C_{26}H_{26}ClN_3$	−1630.907	−1630.415	−1630.501
DB11672	Curcumin	$C_{21}H_{20}O_6$	−1263.596	−1263.199	−1263.287

Table 2 Energy in atomic unit of HOMO, LUMO, Gap energy and dipole moment in Debye

Drug ID	Drug name	HOMO	LUMO	Gap energy	Dipole moment
DB00341	Cetirizine	-0.21913	-0.02237	0.19676	3.921
DB00455	Loratadine	-0.22145	-0.03411	0.18735	2.285
DB00557	Hydroxyzine	-0.21655	-0.02201	0.19454	2.559
DB00967	Desloratadine	-0.21699	-0.02894	0.18805	3.688
DB04890	Bepotastine	-0.21782	-0.03635	0.18147	4.886
DB09488	Acrivastine	-0.12481	-0.06658	0.05823	2.809
DB11614	Rupatadine	-0.22021	-0.03209	0.18812	1.126
DB11672	Curcumin	-0.20535	-0.07206	0.13329	3.750

Fig. 1 Optimized structure of selected drug candidates

(Acrivastine) and DB11672 (Curcumin) which was calculated to be 0.05823 eV and 0.13329 eV, respectively.

Induced fit docking simulation and MM-GBSA calculation

In induced fit docking, both ligands and receptors are flexible. The glide induced fit docking has extra refinement

features for advanced refinement and prediction of conformational changes within receptor active-site and minimization that proceeds to redock. Conformational changes are very crucial for accuracy of docking, which is why prime refinement and prediction is very necessary to ascertain accurate docking and characterizing of the binding sites. Ligand poses are ranked by glide score in an empirical function, which approximates ligand binding free energy. In this

Fig. 2 Plots of HOMO and LUMO. The positive electron density is indicated by red color while green color indicates negative electron density

study; DB11672, DB04890, DB11614, DB09488, DB00557, DB00341, DB00341, DB00967 and DB00455 showed a glide score of -12.111 kcal/mol, -11.066 kcal/mol, -11.974 kcal/mol, -10.769 kcal/mol, -12.076 kcal/mol, -11.522 kcal/mol, -11.094 kcal/mol and -12.233 kcal/mol, respectively. The best ligand binding affinity was observed for DB00557 (Hydroxyzine), which was calculated to be -140.491 kcal/mol. The details of docking results of selected candidates are reported in Table 3. Among all docked complexes, the amino acid Tyr-431 was commonly found in H-bonding interaction except for in DB00341. However, interestingly the amino acid Trp-428 was found in π - π stacking interaction amongst all docked complexes. Therefore, Tyr-108 was also found to form π -cation bonding in the complexes except for in the cases of DB00455

Table 3 Induced fit docking and MM-GBSA ΔG bind results in kcal/mol

Entry name	Drug name	Glide gscore	Prime MM-GBSA ΔG (kcal/mol)
DB11672	Cetirizine	-12.111	-87.046
DB04890	Loratadine	-11.066	-39.379
DB11614	Hydroxyzine	-11.974	-48.656
DB09488	Desloratadine	-10.769	-89.457
DB00557	Bepotastine	-12.076	-140.491
DB00341	Acrivastine	-11.522	-47.834
DB00967	Rupatadine	-11.094	-57.412
DB00455	Curcumin	-12.233	-63.840

and DB11672. The most prominent interaction, however, was found for DB11672 (Curcumin), which interacts with Lys-191, Lys-179, Thr-194, Asp-107, Tyr-108, Tyr-431 by

H bonding formation. Detailed ligand–receptor interaction diagrams are given in Fig. 3. Please find their docked poses in Supplementary Fig. 5.

Fig. 3 Receptor–ligand interaction

P450 site of metabolism

The enzymes of cytochrome p450 brings about the detoxification of drugs and other foreign compounds by undergoing various reactions such as hydroxylation, dealkylation, and double-bond oxidation (Liu et al. 2013). A metabolic reaction can ensue at a highly reactive site of a molecule, on the condition that it is properly oriented within the CYP enzyme to interact with the reaction center. Therefore, to understand the crucial aspects of iron accessibility, intrinsic reactivity to form combined SOM (site of metabolism), we performed induced fit docking using CYP2D6 isoform, as well as, CYP3A4 isoform for the compound Curcumin (Fig. 4a) and the Control, Haloperidol (Fig. 4b). Choice of CYP2D6 isoform, which is the most drug-metabolizing enzyme isoform aside from CYP3A4, allows for greater recognition of substrates due to its higher selectivity (Rydberg and Olsen 2011). Detailed scoring parameters have been given in Fig. 4a, b for the label atoms, with the accessibility of the atoms to the iron. This is defined as the natural logarithm of the number of poses for an atom, in which the atom was within 5 Å of Fe. Larger values indicate greater accessibility. As seen below, Curcumin (Fig. 4a) shows larger values in comparison to Haloperidol (Fig. 4b), thusly indicating greater accessibility. CYP3A4 isoform, the most prevalent CYP enzyme, which catalyzes the metabolism of nearly half the drugs on the market (Koe et al. 2014), had only its intrinsic reactivity assessed because ligand-based models

have been found to provide satisfactory prediction, with the intrinsic reactivity playing a significant role (Rydberg and Olsen 2011). Label atoms with the intrinsic reactivity for 3A4, were calculated with Hammett and Taft methodology. Positive values are more reactive, whilst negative values are less reactive. As shown in the figures below, certain label atoms of Curcumin express greater positive values indicating that they're relatively more reactive in comparison to its control's (Haloperidol) label atoms, which is therefore relatively less reactive. In the case of Combined SOM, all three annotations are displayed at the same time, in graphical form. The Fe-accessibility is indicated by a set of green rays drawn out from the atom to a given maximum length. Each full-length ray represents one unit of accessibility, so the number of full-length rays is the integer part of the score. Thus, the remaining partial ray represents the decimal part of the score. The circles represent the overall score and the intrinsic reactivity. The radius of the circle is proportional to this score. The intensity of the red color of the circle is proportional to the intrinsic reactivity. The blue perimeter indicates whether the atom passed the filtering stage, which has cutoffs for the intrinsic reactivity and the number of poses. Thusly, as illustrated down below, Curcumin displays relatively greater Fe-accessibility (greater set of green rays), increased overall score and intrinsic reactivity (greater radius, intensity and number of red circles), and a greater number of label atoms that passed the filtering stage (greater number of blue perimeters around circles) compared to that

Fig. 4 a Metabolic maps of Curcumin. b Metabolic maps of Haloperidol (Control)

found with the control (Haloperidol). Overall, these findings indicate that Curcumin is potentially safer (relatively less toxic), compared to the FDA approved drug Haloperidol (ID: DB00502).

Conclusion

From our study, we have discovered the common binding sites residues of H1R, which are involved in interaction with FDA approved commercialized drugs which might be useful for future drug target site. Those active sites have also been shown to be inhibited by our proposed H1R antagonist, curcumin. Amongst the FDA approved drugs, DB00557 (Hydroxyzine), exhibited the best binding affinity. However, the best FDA approved drugs evaluated by in silico docking approach has been found to be less reactive than curcumin as reported in Table 2. Besides this, the dipole moment of DB11672 (Curcumin) has found to be greater than DB00557 (Hydroxyzine), as reported in Table 2. From ligand interaction analysis DB11672 (Curcumin) emphasized six H bonding network with receptor's active sites, whereas DB00557 (Hydroxyzine), interacts with active site residues by four hydrogen bonding networks. Based on all the evidences given above, this study concludes with the firm belief that use of H1R inhibitor from naturally occurring compound Curcumin, is seen to be a comparatively more effective and a therapeutically more significant cure for H1R mediated allergic symptoms based on In-Silico approach.

Acknowledgements We would like to express our deepest gratitude and thanks to Allah, our parents, our families, our teachers and all those who've helped and supported us along the way.

Funding The authors received no funding from an external source.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Ethics approval None.

References

- Aggarwal BB, Sundaram C, Malani N, Ichikawa H (2007) Curcumin: the Indian solid gold. *Adv Exp Med Biol* 595:1–75. https://doi.org/10.1007/978-0-387-46401-5_1
- Antony S, Kuttan R, Kuttan G (1999) Immunomodulatory activity of curcumin. *Immunol Invest* 28:291–303
- Arima M et al (1999) Expression of interleukin-16 by human epithelial cells: inhibition by dexamethasone. *Am J Respir Cell Mol Biol* 21:684–692
- Bakker RA, Schoonus SB, Smit MJ, Timmerman H, Leurs R (2001) Histamine H1-receptor activation of nuclear factor- κ B: roles for G β γ - and G α q/11-subunits in constitutive and agonist-mediated signaling. *Mol Pharmacol* 60:1133–1142
- Bakker RA, Casarosa P, Timmerman H, Smit MJ, Leurs R (2004) Constitutively active Gq/11-coupled receptors enable signaling by co-expressed Gi/o-coupled receptors. *J Biol Chem* 279:5152–5161
- Balasubramanyam M, Koteswari AA, Kumar RS, Monickaraj SF, Maheswari JU, Mohan V (2003) Curcumin-induced inhibition of cellular reactive oxygen species generation: novel therapeutic implications. *J Biosci* 28:715–721
- Berman HM et al (2000) The protein data bank. *Nucleic Acids Res* 28:235–242
- Brown RE, Sergeeva OA, Eriksson KS, Haas HL (2002) Convergent excitation of dorsal raphe serotonin neurons by multiple arousal systems (orexin/hypocretin, histamine and noradrenaline). *J Neurosci* 22:8850–8859
- Caron G, Delneste Y, Roelandts E, Duez C, Herbault N, Magistrelli G, Bonnefoy J-Y, Pestel J, Jeannin P (2001) Histamine induces CD86 expression and chemokine production by Human Immature Dendritic Cells. *J Immunol* 166(10):6000–6006
- Chandra D, Gupta SJ (1972) Anti-inflammatory and anti-arthritic activity of volatile oil of *Curcuma longa* (Haldi). *Indian J Med Res* 60:138
- Chung S-H, Choi SH, Choi JA, Chuck RS, Joo C-K (2012) Curcumin suppresses ovalbumin-induced allergic conjunctivitis. *Mol Vis* 18:1966–1972
- Conen S et al (2013) The role of P-glycoprotein in CNS antihistamine effects. *Psychopharmacology* 229:9–19
- de Benedictis FM, Bush AJA, Medicine CC (2012) Corticosteroids in respiratory diseases in children. *Am J Respir Crit Care Med* 185:12–23
- Easton RE, Giesen DJ, Welch A, Cramer CJ, Truhlar DG (1996) The MIDI! basis set for quantum mechanical calculations of molecular geometries and partial charges. *Theor Chim Acta* 93:281–301. <https://doi.org/10.1007/bf01127507>
- Elphick HE, Southern KW (2012) Antifungal therapies for allergic bronchopulmonary aspergillosis in people with cystic fibrosis. *Cochrane Database Syst Rev*. <https://doi.org/10.1002/14651858.cd002204.pub2>
- Frisch MJ, Trucks GW, Schlegel HB, Scuseria GE, Robb MA, Cheeseman JR, Scalmani G, Barone V, Petersson GA, Nakatsuji H, Li X, Caricato M, Marenich AV, Bloino J, Janesko BG, Gomperts R, Mennucci B, Hratchian HP, Ortiz JV, Izmaylov AF, Sonnenberg JL, Williams-Young D, Ding F, Lipparini F, Egidi F, Goings J, Peng B, Petrone A, Henderson T, Ranasinghe D, Zakrzewski VG, Gao J, Rega N, Zheng G, Liang W, Hada M, Ehara M, Toyota K, Fukuda R, Hasegawa J, Ishida M, Nakajima T, Honda Y, Kitao O, Nakai H, Vreven T, Throssell K, Montgomery Jr JA, Peralta JE, Ogliaro F, Bearpark MJ, Heyd JJ, Brothers EN, Kudin KN, Staroverov VN, Keith TA, Kobayashi R, Normand J, Raghavachari K, Rendell AP, Burant JC, Iyengar SS, Tomasi J, Cossi M, Millam JM, Klene M, Adamo C, Cammi R, Ochterski JW, Martin RL, Morokuma K, Farkas O, Foresman JB, Fox DJ (2016) Gaussian 16 Rev. B.01. Gaussian, Inc., Wallingford CT
- Fujikura T, Shimosawa T, Yakuo I (2001) Regulatory effect of histamine H1 receptor antagonist on the expression of messenger RNA encoding CC chemokines in the human nasal mucosa. *J Allergy Clin Immunol* 107:123–128
- Gilani AH, Shah AJ, Ghayur MN, Majeed KJLS (2005) Pharmacological basis for the use of turmeric in gastrointestinal and respiratory disorders. *Life Sci* 76:3089–3105
- Giustizieri ML, Albanesi C, Fluhr J, Gisondi P, Norgauer J, Girolomoni G (2004) H1 histamine receptor mediates inflammatory responses in human keratinocytes. *J Allergy Clin Immunol* 114:1176–1182. <https://doi.org/10.1016/j.jaci.2004.07.054>

- Harder E et al (2016) OPLS3: a force field providing broad coverage of drug-like small molecules and proteins. *J Chem Theory Comput* 12:281–296. <https://doi.org/10.1021/acs.jctc.5b00864>
- Hill S et al (1997) International Union of Pharmacology. XIII. Classification of histamine receptors. *Pharmacol Rev* 49:253–278
- Hoque MM, Halim MA, Sarwar MG, Khan MW (2015) Palladium-catalyzed cyclization of 2-alkynyl-*N*-ethanoyl anilines to indoles: synthesis, structural, spectroscopic, and mechanistic study. *J Phys Org Chem* 28:732–742. <https://doi.org/10.1002/poc.3477>
- Jeannin P et al (1994) Histamine induces interleukin-8 secretion by endothelial cells. *Blood* 84:2229–2233
- Jutel M, Watanabe T, Akdis M, Blaser K, Akdis CA (2002) Immune regulation by histamine. *Curr Opin Immunol* 14:735–740. [https://doi.org/10.1016/s0952-7915\(02\)00395-3](https://doi.org/10.1016/s0952-7915(02)00395-3)
- Kanda N, Watanabe S (2004) Histamine enhances the production of granulocyte-macrophage colony-stimulating factor via protein kinase C α and extracellular signal-regulated kinase in human keratinocytes. *J Investig* 122:863–872
- Kim SH, Kwon JE, Kim H-BJA (2013) Successful treatment of steroid-dependent eosinophilic cellulitis with cyclosporine. *Allergy Asthma Immunol Res* 5:62–64
- Koe XF, Tengku Muhammad TS, Chong AS-C, Wahab HA, Tan ML (2014) Cytochrome P450 induction properties of food and herbal-derived compounds using a novel multiplex RT-qPCR in vitro assay, a drug-food interaction prediction tool. *Food Sci Nutr* 2:500–520. <https://doi.org/10.1002/fsn3.122>
- Lee C, Yang W, Parr RG (1988) Development of the Colle–Salvetti correlation-energy formula into a functional of the electron density. *Phys Rev B Condens Matter* 37:785–789
- Lee JH et al (2008) Curcumin, a constituent of curry, suppresses IgE-mediated allergic response and mast cell activation at the level of Syk. *J Allergy Clin Immunol* 121:1225–1231. <https://doi.org/10.1016/j.jaci.2007.12.1160>
- Li J, Abel R, Zhu K, Cao Y, Zhao S, Friesner RA (2011) The VSGB 2.0 model: a next generation energy model for high resolution protein structure modelling. *Proteins* 79:2794–2812. <https://doi.org/10.1002/prot.23106>
- Liu X et al (2013) In silico prediction of cytochrome P450-mediated site of metabolism (SOM). *Protein Pept Lett* 20:279–289
- Lovell SC et al (2003) Structure validation by C α geometry: phi, psi and C β deviation. *Proteins* 50:437–450. <https://doi.org/10.1002/prot.10286>
- Lutomski J, Kedzia B, Debska W (1974) Effect of an alcohol extract and of active ingredients from *Curcuma longa* on bacteria and fungi (author's transl). *Planta Med* 26:9–19. <https://doi.org/10.1055/s-0028-1097963>
- Ma RZ et al (2002) Identification of Bphs, an autoimmune disease locus, as histamine receptor H1. *Science* 297:620–623
- Ma C, Ma Z, Fu Q, Ma S (2013) Curcumin attenuates allergic airway inflammation by regulation of CD4 + CD25 + regulatory T cells (Tregs)/Th17 balance in ovalbumin-sensitized mice. *Fitoterapia* 87:57–64. <https://doi.org/10.1016/j.fitote.2013.02.014>
- Madhavi Sastry G, Adzhigirey M, Day T, Annabhimoju R, Sherman W (2013) Protein and ligand preparation: parameters, protocols, and influence on virtual screening enrichments. *J Comput Mol Des* 27:221–234. <https://doi.org/10.1007/s10822-013-9644-8>
- Matsubara M, Tamura T, Ohmori K, Hasegawa K (2005) Histamine H1 receptor antagonist blocks histamine-induced proinflammatory cytokine production through inhibition of Ca $^{2+}$ -dependent protein kinase C, Raf/MEK/ERK and IKK/ κ B/NF- κ B signal cascades. *Biochem Pharmacol* 69(3):433–449
- Musio S et al (2006) A key regulatory role for histamine in experimental autoimmune encephalomyelitis: disease exacerbation in histidine decarboxylase-deficient mice. *J Immunol* 176:17–26
- Oh S-W, Cha J-Y, Jung J-E, Chang B-C, Kwon H-J, Lee B-R, Kim D-Y (2011) Curcumin attenuates allergic airway inflammation and hyper-responsiveness in mice through NF- κ B inhibition. *J Ethnopharmacol* 136:414–421. <https://doi.org/10.1016/j.jep.2010.07.026>
- Parr RG, Zhou Z (1993) Absolute hardness: unifying concept for identifying shells and subshells in nuclei, atoms, molecules, and metallic clusters. *Acc Chem Res* 26:256–258. <https://doi.org/10.1021/ar00029a005>
- Pedotti R, De Voss JJ, Steinman L, Galli SJ (2003) Involvement of both ‘allergic’ and ‘autoimmune’ mechanisms in EAE, MS and other autoimmune diseases. *Trends Immunol* 24:479–484
- Rajakrishnan V, Viswanathan P, Rajasekharan K, Menon VP (1999) Neuroprotective role of curcumin from *Curcuma longa* on ethanol-induced brain damage. *Phytother Res Int J Devot Pharmacol Toxicol Eval Nat Prod Deriv* 13:571–574
- Ringman JM, Frautschy SA, Cole GM, Masterman DL, Cummings JL (2005) A potential role of the curry spice curcumin in Alzheimer's disease. *Curr Alzheimer Res* 2:131–136
- Rydberg P, Olsen L (2011) Ligand-based site of metabolism prediction for cytochrome P450 2D6. *ACS Med Chem Lett* 3:69–73. <https://doi.org/10.1021/ml200246f>
- Schaefer U, Schmitz V, Schneider A, Neugebauer E (1999) Histamine induced homologous and heterologous regulation of histamine receptor subtype mRNA expression in cultured endothelial cells. *Shock* 12:309–315
- Shelley JC, Cholleti A, Frye LL, Greenwood JR, Timlin MR, Uchimaya M (2007) Epik: a software program for pK a prediction and protonation state generation for drug-like molecules. *J Comput Aided Mol* 21:681–691. <https://doi.org/10.1007/s10822-007-9133-z>
- Sherman W, Day T, Jacobson MP, Friesner RA, Farid R (2006) Novel procedure for modeling ligand/receptor induced fit effects. *J Med Chem* 49:534–553. <https://doi.org/10.1021/jm050540c>
- Srinivasan K (1953) A chromatographic study of the curcuminoids in *Curcuma longa* L. *J Pharm Pharmacol* 5:448–457
- Tarnasky P, Van JAP (1990) Antihistamine therapy in allergic rhinitis. *J Fam Pract* 30:71–80
- Thakare VN, Osama MM, Naik SR (2013) Therapeutic potential of curcumin in experimentally induced allergic rhinitis in guinea pigs. *Int Immunopharmacol* 17:18–25. <https://doi.org/10.1016/j.intimp.2013.04.025>
- Thompson DA, Tan BB (2006) Tetrahydracurcumin-related allergic contact dermatitis. *Contact Dermat* 55:254–255. <https://doi.org/10.1111/j.1600-0536.2006.00857.x>
- Toda S, Miyase T, Arichi H, Tanizawa H, Takino YJC, Bulletin P (1985) Natural antioxidants. III. Antioxidative components isolated from rhizome of *Curcuma longa* L. *Chem Pharm Bull* 33:1725–1728
- Venter C, Pereira B, Voigt K, Grundy J, Clayton CB, Higgins B, Arshad SH, Dean T (2008) Prevalence and cumulative incidence of food hypersensitivity in the first 3 years of life. *Allergy* 63:354–359
- Wang E-J, Casciano CN, Clement RP, Johnson WW (2001) Evaluation of the interaction of loratadine and desloratadine with P-glycoprotein. *Drug Metab Dispos* 29:1080–1083
- White MV (1990) The role of histamine in allergic diseases. *J Allergy Clin Immunol* 86:599–605. [https://doi.org/10.1016/S0091-6749\(05\)80223-4](https://doi.org/10.1016/S0091-6749(05)80223-4)
- Wishart DS et al (2018) DrugBank 5.0: a major update to the DrugBank database for 2018. *Nucleic Acids Res* 46:D1074–D1082. <https://doi.org/10.1093/nar/gkx1037>
- Wong R, Wong M, Robinson PD, Fitzgerald DA (2013) Omalizumab in the management of steroid dependent allergic bronchopulmonary aspergillosis (ABPA) complicating cystic fibrosis. *Paediatr Respir Rev* 14:22–24