Published in final edited form as:

Sci Bull (Beijing). 2018 December 15; 63(23): 1527–1528. doi:10.1016/j.scib.2018.09.020.

A new guideline for diagnosis and treatment of syncope in children and adolescents that stimulates further thought and discussion

Julian M. Stewart

Physiology and Medicine, Center for Hypotension, The New York Medical College, Valhalla, NY 10595, USA

A new guideline for diagnosis and treatment of syncope in children and adolescents was published by Chinese Pediatric Cardiology Society (CPCS) [1]. This guideline admirably outlines a differential diagnosis of syncope in the Chinese population, placing appropriate emphasis on vasovagal syncope (VVS, simple faint), the most common cause. The authors appropriately discuss other more ominous causes of syncope such as cardiogenic syncope and neurogenic orthostatic hypotension. In particular, cardiogenic syncope can have lethal consequences if not promptly treated. The paper provides a useful algorithm for diagnostic evaluation of syncope patients placing primary emphasis on history, physical exam, and ECG, and further discussing testing procedures such as standing tests used in screening patients, and upright tilt table testing which remains the reference standard for orthostatic testing in younger patients. Indeed, most vasovagal and other forms of reflex or situational syncopes can be best diagnosed on clinical history [2]. The authors further consider other forms of orthostatic intolerance such as orthostatic hypertension [3], postural tachycardia syndrome (POTS), and “initial orthostatic hypotension” experienced as transient lightheadedness upon rapid standing [4]. The authors regard POTS as a cause of vasovagal syncope. This supposition is controversial and deserves further comment.

POTS has been reported to be associated with vasovagal syncope to varying degree [5,6]. Postural Tachycardia Syndrome was a term devised by workers at Mayo clinic to describe patients with symptoms of chronic orthostatic intolerance synchronous with excessive upright tachycardia but who had no postural hypotension as characterizes neurogenic orthostatic hypotension and vasovagal syncope [7]. POTS is best considered “a heterogeneous and multifactorial disorder” [8] of diverse etiology rather than a disease or causal entity. POTS can be clinically distinguished from typical recurrent vasovagal syncope in that POTS is chronic with day-to day symptoms while VVS is episodic; between episodes VVS patients are generally well.

Tilt table studies may add confusion to this etiological scheme. In brief, tilt table results need not correspond to real world syncope. More than 40% of recurrent VVS patients have

julian_stewart@nymc.edu.

Conflict of interest

The author declare that he has no conflict of interest.

heart rates exceeding POTS thresholds during tilt tests [9], while prolonged orthostatic testing of POTS patients beyond the original standard 10-minute tilt produces “false positive tests” including increasing numbers of false positive POTS [10] and a relatively high incidence of vasovagal syncope in healthy young volunteers (and POTS patients) with no real world history of syncope [11]. Specificity is further reduced by nitroglycerin [12]. However, recent reconsideration of such false positive VVS suggests that they represent a subpopulation with a hypotensive susceptibility [13]. Moreover, the vasovagal syncope response can be induced in almost everyone at different thresholds of orthostatic stress [14,15] and may be an evolutionary reflex response to hemorrhage [16].

In summary, the guideline provides a useful review of syncope and orthostatic intolerance in general while stimulating further thought and discussion.

Acknowledgments

This work was supported by RO1 HL 134674 and RO1 HL 112736 from the National Heart Lung and Blood Institute.

Biography

Julian Stewart graduated from Cornell University with a degree in physics, receiving M.D. and Ph.D. (cardiac physiology) from the University of Chicago, and internship and residency at New York University. He completed a Pediatric Cardiology fellowship at Cornell University-New York Hospital. He is currently Professor of Pediatrics, Physiology, and Medicine, and Associate Chairman of Pediatrics at New York Medical College. He directs the Center for Hypotension, funded by the National Heart, Lung and Blood Institute of the National Institutes of Health, and has studied the circulatory bases of orthostatic intolerance in the young for more than 18 years with a focus on mechanisms of vasovagal syncope and postural tachycardia syndrome.

References

- [1]. Wang C, Li Y, Liao Y, et al. Chinese Pediatric Cardiology Society (CPCS) guideline for diagnosis and treatment of syncope in children and adolescents. *Sci Bull* 2018;63:1558–64.
- [2]. Wieling W, van Dijk N, de Lange FJ, et al. History taking as a diagnostic test in patients with syncope: developing expertise in syncope. *Eur Heart J* 2015;36:277–80. [PubMed: 25514928]
- [3]. Robertson D Orthostatic hypertension: the last hemodynamic frontier. *Hypertension* 2011;57:158–9. [PubMed: 21199993]
- [4]. Wieling W, Krediet CT, van Dijk N, et al. Initial orthostatic hypotension: review of a forgotten condition. *Clin Sci (Lond)* 2007;112:157–65. [PubMed: 17199559]
- [5]. Ojha A, McNeeley K, Heller E, et al. Orthostatic syndromes differ in syncope frequency. *Am J Med* 2010;123:245–9. [PubMed: 20193833]

- [6]. Raj SR. The Postural Tachycardia Syndrome (POTS): pathophysiology, diagnosis & management. *Indian Pacing Electrophysiol J* 2006;6:84–99. [PubMed: 16943900]
- [7]. Schondorf R, Low PA. Idiopathic postural orthostatic tachycardia syndrome: an attenuated form of acute pandysautonomia? *Neurology* 1993;43:132–7. [PubMed: 8423877]
- [8]. Benarroch EE. Postural tachycardia syndrome: a heterogeneous and multifactorial disorder. *Mayo Clin Proc* 2012;87:1214–25. [PubMed: 23122672]
- [9]. Medow MS, Merchant S, Suggs M, et al. Postural heart rate changes in young patients with vasovagal syncope. *Pediatrics* 2017;139:e13286.
- [10]. Plash WB, Diedrich A, Biaggioni I, et al. Diagnosing postural tachycardia syndrome: comparison of tilt testing compared with standing haemodynamics. *Clin Sci (Lond)* 2013;124:109–14. [PubMed: 22931296]
- [11]. Lewis DA, Zlotocha J, Henke L, et al. Specificity of head-up tilt testing in adolescents: effect of various degrees of tilt challenge in normal control subjects. *J Am Coll Cardiol* 1997;30:1057–60. [PubMed: 9316539]
- [12]. Vlahos AP, Tzoufi M, Katsouras CS, et al. Provocation of neurocardiogenic syncope during head-up tilt testing in children: comparison between isoproterenol and nitroglycerin. *Pediatrics* 2007;119:e419–25. [PubMed: 17224456]
- [13]. Sutton R, Brignole M. Twenty-eight years of research permit reinterpretation of tilt-testing: hypotensive susceptibility rather than diagnosis. *Eur Heart J* 2014;35:2211–2. [PubMed: 24980488]
- [14]. el-Bedawi KM, Hainsworth R. Combined head-up tilt and lower body suction: a test of orthostatic tolerance. *Clin Auton Res* 1994;4:41–7. [PubMed: 8054836]
- [15]. LeLorier P, Klein GJ, Krahn A, et al. Combined head-up tilt and lower body negative pressure as an experimental model of orthostatic syncope. *J Cardiovasc Electrophysiol* 2003;14:920–4. [PubMed: 12950533]
- [16]. Alboni P, Alboni M. Vasovagal syncope as a manifestation of an evolutionary selected trait. *J Atr Fibrillation* 2014;7:1035. [PubMed: 27957092]