Cr(VI) reduction by an extracellular polymeric substance (EPS) produced from a strain of *Pseudochrobactrum saccharolyticum*

Dongyan Long^{1,2} · Muhammad Zaffar Hashmi³ · Xiaomei Su⁴ · Siwatt Pongpiachan⁵

Received: 24 October 2018 / Accepted: 20 February 2019 / Published online: 28 February 2019
© King Abdulaziz City for Science and Technology 2019

Abstract

A better understanding of the Cr(VI) reduction position and mechanisms by a Cr(VI)-reducing strain is important for the bioremediation of Cr pollution in the environment. In the present study, we were interested in figuring out the role of extracellular polymeric substances (EPS) as the main area for Cr(VI) reduction in the newly reported strain of *Pseudochrobactrum saccharolyticum* LY10. We investigated the subcellular distribution and reduction capability of each cellular component as the main area of Cr(VI) reduction by scanning electron microscopy and soft X-ray spectromicroscopy. The results suggested that most of Cr was presented in the supernatants as Cr(III) after reduction. In the cells, Cr was mostly distributed in the EPS and cell wall, while the EPS had the maximum Cr(VI) reduction rate (81.5%) as compared with the cell wall (30.1%). Soft X-ray spectromicroscopy analysis indicated that Cr accumulated more in the EPS. Therefore, the results suggested that the EPS were the main area for Cr(VI) reduction in the bacteria of *P. saccharolyticum* LY10.

Keywords EPS · *Pseudochrobactrum saccharolyticum* LY10 · Cr(VI) reduction · Subcellular distribution · Soft X-ray spectromicroscopy

Introduction

Chromium (Cr) might pose a severe threat to public health and the environment if discharged without adequate pretreatments (Zhong et al. 2014; Lai and Lo 2008; Beukes et al. 2017). Cr(VI) is one of the prevalent states found in the environment and is more toxic because of its mobility compared

with the Cr(III). Therefore, the reduction of Cr(VI) has been widely studied and is of great concern to many researchers. Cr(VI) has also been identified as 1 of the 17 chemicals posing the greatest threat to humans by the United States Environmental Protection Agency (Lan et al. 2014). Compared with the traditional chemical reduction methods, microbial reduction of Cr(VI) has been considered as a promising process for its cost-effective and eco-friendly advantages (Cervantes et al. 2001; Chai et al. 2009; Morel et al. 2011; Ma et al. 2019; Su et al. 2019).

The microbial reduction of Cr(VI) has been reported for a variety of bacterial strains in aerobic or anaerobic conditions, such as *Paracoccus denitrificans* (Mazoch et al. 2004), *Bacillus subtilis* (Morokutti et al. 2005), *Vibrio fischeri* (Ful-ladosa et al. 2006), *Pseudomonas aeruginosa* (Kilic et al. 2010), and *Pseudomonas putida* (Dogan et al. 2011). A novel Cr(VI) reducing bacteria (*Pseudochrobactrum saccharolyticum* LY10) has also been isolated and characterized for its high Cr(VI) reducing ability, typical alkali tolerance, and halotolerance in our previous study (Long et al. 2013).

The direct enzymatic reaction or other indirect biotransformation to metabolites are the major processes for reduction of Cr(VI) (Lovley 1993; Lee et al. 2000; Valls et al. 2000). Basically, microbial reduction of Cr(VI) can

✉ Muhammad Zaffar Hashmi
hashmi_qau@yahoo.com

¹ College of Environmental and Natural Resource Sciences, Zhejiang University, Yuhangtang Road 388, Hangzhou 310058, Zhejiang, People's Republic of China

² Chaotianmen Sub-district Office, Yuzhong District, Chongqing 400011, People's Republic of China

³ Department of Meteorology, COMSATS University, Islamabad, Pakistan

⁴ College of Geography and Environmental Science, Zhejiang Normal University, Yingbin Road 688#, Jinhua 321004, People's Republic of China

⁵ School of Social and Environmental Development, National Institute of Development Administration (NIDA), 118 Moo3, Sereethai Road, Klong-Chan, Bangkok, Bangkok 10240, Thailand

be divided into two processes, intracellular and extracellular. Some microbes can directly use the related intracellular reductase to reduce Cr(VI), and some others use extracellular metabolites to reduce the highly toxic Cr(VI). Previous studies have shown that *P. putida* MK1 could use the soluble reductase ChrR to reduce Cr(VI) (Park et al. 2000), whereas *Bacillus megaterium* TKW3 uses a membrane-associated reductase to reduce Cr(VI) (Cheung and Gu 2005). Under anaerobic conditions, microorganisms can use extracellular metabolites with Cr(VI) reducibility to reduce Cr(VI) extracellularly. These studies found that sulfate-reducing bacteria can reduce the sulfate radical to the divalent negative sulfur. This reduction occurs by a series of enzymes, including sulfuric adenosyltransferase and adenylyl sulfate reductase (APS reductase), that generate the metabolite of H₂S as an effective Cr(VI) reducing agent (Tebo and Obraztsova 1998). In addition, some Fe(III) reduction microorganisms reduce Fe(III) to Fe(II) and use the resulting Fe(II) to reduce Cr(VI) (Wielinga et al. 2001). Therefore, there are different Cr(VI) reduction positions and mechanisms for different microorganisms.

Extracellular polymeric substances (EPS), mainly containing polysaccharides, proteins, extracellular DNA, and lipids, exhibit a dynamic double-layered structure with loosely bound EPS and tightly bound EPS (Flemming and Wingender 2010). EPS have an abundance of negatively charged functional groups that react with heavy metals as ligands for which the adsorption characteristics and kinetics have been reported previously (Wang et al. 2014a). The EPS produced by several bacterial species are recognized as biofloculants, bioabsorbents, encapsulating materials, heavy metal-removing agents, drug delivery agents, ion exchange resins, a natural immunomodulator and antioxidant and anti-biofilm agents (Kavita et al. 2013). In addition, previous studies have shown that the Cr(VI) sorption of loosely bound EPS and tightly bound EPS is followed by a charge transfer from the nitrogen and oxygen to the Cr(VI), and the transfer of charge can result in the reduction of Cr(VI) to Cr(III) (Wang et al. 2014b). However, whether the EPS are the main sites of Cr(VI) reduction have not been proven.

The soft X-ray spectromicroscopy has been proved to be a powerful technology for the spectral imaging and chemical speciation analysis of metals. It combined the advantages of scanning transmission X-ray microscopy with high spatial resolution and the near-edge absorption fine structure spectroscopy with high-energy resolution (Smith 2001; Chao et al. 2005). Compared with transmission electron microscopy and other technologies, the soft X-ray spectromicroscopy analysis causes relatively little damage to the samples and can be used in nano-scale research on a variety of forms such as solid and liquid (Jacobsen et al. 2000). For environment pollutants, the soft X-ray spectromicroscopy can also be used to study the chemical distribution of organic

compounds and inorganic metals by K-edge and L-edge, respectively.

Overall, we hypothesized that the EPS were the main sites for Cr(VI) reduction in the newly reported strain of *P. saccharolyticum* LY10. To prove this, the Cr(VI) subcellular distribution and the reduction capability of various components of *P. saccharolyticum* LY10 cells were investigated in the present study. Furthermore, scanning electron microscopy (SEM), and specially, the soft X-ray spectromicroscopy method were applied to observe the images of cells or single cell under Cr(VI) stress to confirm the main area of Cr(VI) reduction in the cell.

Methodology

Bacteria and media

The novel Cr(VI)-reducing strain of *P. saccharolyticum* LY10, isolated in our previous study (Long et al. 2013) and characterized for its high Cr(VI)-reducing ability, alkali tolerance and halotolerance, was selected for the present study. The common Luria–Bertani (LB) medium with tryptone (10 g), yeast extract (5 g), NaCl (5 g), and distilled water (1 L) in pH 8.0 was used during the incubation of bacteria (McLean and Beveridge 2001; Chen et al. 2012). Cr(VI) was added into the LB medium of K₂CrO₄ (200 mM) after the filtration by 0.22 µm membrane.

Subcellular components separation

Pseudochrobactrum saccharolyticum LY10 was inoculated in liquid LB medium containing 110 mg L⁻¹ Cr(VI) and 220 mg L⁻¹ Cr(VI), and then was cultured at 28 °C, 160 rpm for 96 h. The broth was centrifuged for 30 min at 6000g and then, the supernatant and cells was separated. The concentration of total Cr and Cr(VI) in the supernatant was measured. The obtained cells were further separated for the subcellular components (Chen 2011). The cells were first resuspended in 10 mL 0.9% NaCl solution and centrifuged at 10,000g, 4 °C for 30 min. The remaining supernatant was the extracellular polymeric substances (EPS). The broth was further resuspended with 10 mL 25 mM Tris–HCl buffer (pH 7.0), and the cells were broken by ultrasonic (200 w, 30:30 s, 20 times). The solution was centrifuged (2500g, 4 °C, 20 min) to remove the unbroken cells, which were redissolved in 10 mL 25 mM Tris–HCl buffer and broken completely by ultrasonic. The mixed broken solution was centrifuged (30,000g, 4 °C, 30 min) to obtain the precipitate (cell wall). While the remaining supernatant was centrifuged at 110,000g, 4 °C for 2 h, the precipitate was the cell membrane, and the rest of the supernatant was soluble fraction of cells.

The pellet obtained after centrifugation was digested with concentrated nitric acid at 200 °C for 2 h and diluted with 5% hydrochloric acid at constant volume. The soluble components were treated with concentrated nitric acid (5%) overnight. A Thermo Element MKII-M6 atomic absorption spectrophotometer was used to measure the total Cr content in different components. Colorimetric method was used to determine the concentration of Cr(VI) at 540 nm using the DPC method with a detection limit of 0.2 $\mu\text{g L}^{-1}$.

Cr(VI) reduction by different cell components

The *P. saccharolyticum* LY10 was incubated in LB liquid medium without Cr(VI) and cultured at 28 rpm for 24 h. The EPS, cell wall, cell membrane, and soluble fraction were obtained by the centrifugation method. Following purification, the ability of different cell components to reduce Cr(VI) was tested. Generally, 10 mL of phosphate buffer (0.1 M, pH 7.0) was added separately into the cell components, and a Cr(VI) stock solution was then added to a final Cr(VI) concentration of 5.2 mg L^{-1} in the system. The Cr(VI) was sampled in each system under 28 °C, 160 rpm for 16 h to measure the remaining Cr(VI) concentration and further analyze the Cr(VI) reduction capability of different cellular components.

SEM analysis

Pseudochrobactrum saccharolyticum LY10 was inoculated in liquid LB medium that contained 55 mg L^{-1} Cr(VI), 110 mg L^{-1} Cr(VI), or 220 mg L^{-1} Cr(VI). Cultures were maintained at 28 °C, 160 rpm for 96 h, and then the cells were collected after centrifugation (6000 rpm, 4 °C, 10 min). The cells were placed in 2.5% glutaraldehyde solution overnight at 4 °C. The samples were rinsed with 0.1 M, pH 7.0 phosphate buffer three times for 15 min. The samples were fixed for 1–2 h with 1% osmium tetroxide solution and rinsed with 0.1 M, pH 7.0 phosphate buffer three times for 15 min each. The samples were dehydrated for 15 min with different concentrations of ethanol solution: 50%, 70%, 80%, 90%, and 95%, followed by dehydration with 100% ethanol twice for 20 min. The samples were treated with a mixture of ethanol and isoamyl acetate (V/V = 1/1) for 30 min, then pure isoamyl acetate for 1–2 h. Then, the samples were dried at the critical point and plated. The processed samples were analyzed using an XL30ESEM environmental scanning electron microscope (Netherlands).

Analysis of soft X-ray spectromicroscopy

Sample preparation

Pseudochrobactrum saccharolyticum LY10 was inoculated in liquid LB medium that contained 220 mg L^{-1} Cr(VI) with

sampling after 24 h and 48 h. The cultures were centrifuged at 5000 rpm, 4 °C for 1 min, and the cells were washed twice with sterile water to remove any Cr interference from the culture medium. The clean bacterial cells were resuspended in sterile water. The cells were dripped to a silicon nitride window, allowed to air dry, and analyzed by the soft X-ray spectroscopy micro-beam line.

Sample measurement and data analysis

The sample detection by soft X-ray spectromicroscopy was completed at beam line 08U of the Shanghai Synchrotron Radiation Facility (SSRF). Data were analyzed using dual-energy ratio-contrast image method. For each specimen, two absorption images were scanned separately with energies E1 and E2, of which E1 was focused on the absorption edge of the chosen element and E2 was focused below the edge (Smith 2001; Chao et al. 2005). Because the change in photon energies was only a few percent or less, the absorption of X-rays by all the other elements in the specimen was negligible, and consequently any observed difference in the transmitted X-ray flux between the two images obtained at the two energies could be attributed to the presence of the element being studied (Smith 2001; Chao et al. 2005). The total electron yield (TEY) method was applied to detect the Cr(VI) reference sample of $\text{K}_2\text{Cr}_2\text{O}_7$ and the Cr(III) reference sample of CrPO_4 . The absorption edge (E1) of Cr(VI) and Cr(III) was confirmed to be 579.8 eV and 576.8 eV, respectively. The E2 which was below the edge focused at 573.8 eV. The three energies were used to scan the sample for image. The scanning range was determined according to the sample of which the scanning step was 0.03 μm and every pixel acquisition time was 20 ms. The distribution map of Cr(VI) was obtained by digital division of two absorption-contrast images at energies of 579.8 eV (at the peak) and 573.8 eV (pre-peak). Similarly, the Cr(III) maps was obtained by the digital ratios of 576.8 eV and 573.8 eV (Smith 2001; Chao et al. 2005). Microsoft Excel and SPSS software version 13.0 were used for the data processing and analysis. In addition, the probability value of 0.05 was used to determine significance in the analysis of variance.

Results

Subcellular distribution of Cr in cells

The Cr content was determined for each component as shown in Fig. 1a. It was found that the respective Cr contents were 102.3 and 208.8 mg L^{-1} in supernatants of the 110 and 220 mg L^{-1} cultures, which accounted for more than 92% of the total amount of Cr, while the immobilized, cellular Cr was only 5.1–7.1% of the total amount. However, most of

the Cr was in the form of Cr(III) in the supernatants, which accounted for 76.9 and 54.0% of the total Cr in supernatants for the two treatments, respectively. To further study the distribution of Cr in different subcellular components, the gradient centrifugation method was applied to divide the whole cells into four components including the EPS, cell wall, cell membrane, and intracellular soluble fraction. Figure 1b showed that the Cr contents were mostly distributed in the EPS and cell wall, accounting for more than 90% of the total Cr in cells. It was noted that the content of Cr was more in the EPS (47.1%) compared with the cell wall (43.3%) when the initial concentration of Cr(VI) was increased to 220 mg L⁻¹. There was only 6.4 and 7.5% of the intracellular soluble fraction at 110 and 220 mg L⁻¹ of Cr(VI) concentration, respectively, whereas the Cr distribution was lowest in the cell membrane (1.5 and 2.1%, respectively).

Cr(VI) reduction by different subcellular fractions

The four subcellular fractions were freshly obtained by the gradient centrifugation method and incubated for Cr(VI) reduction in medium containing 5.2 mg L⁻¹ Cr(VI) for 16 h. The ability of different subcellular fractions to reduce Cr(VI) was compared and presented in Fig. 2. The results indicated that the four fractions had obviously different abilities to reduce Cr(VI). Interestingly, the EPS had the maximum Cr(VI) reduction rate (81.5%) within 16 h, while the cell

wall had the lowest capacity of Cr(VI) reduction with only 30.1% of Cr(VI) reduced within the same amount of time. The reduction rate of Cr(VI) was 69.2 and 49.3% for intracellular soluble fraction and cell membrane, respectively.

SEM observation of cells under Cr(VI) stress

SEM was used to observe the micro-appearance of *P. saccharolyticum* LY10 under different Cr(VI) stressors. In the

Fig. 2 The Cr(VI)-reducing abilities of different subcellular fractions of *P. saccharolyticum* LY10 [experiments were conducted with initial 5.2 mg L⁻¹ Cr(VI)]

Fig. 1 Distribution profiles of Cr in different fractions of *P. saccharolyticum* LY10 culture (**a** distribution in supernatant; **b** the subcellular distribution profile of Cr in cells)

absence of Cr(VI), SEM results revealed that the cells were rod shaped with regular shapes, and they were dispersed throughout the system with no clear connections by extracellular materials (Fig. 3a). At the 55 mg L^{-1} Cr(VI) concentration, EPS was clearly outside of each cell (Fig. 3b). With the increase of Cr concentration, the shape of the cells changed (Fig. 3c). When the *P. saccharolyticum* LY10 was exposed to 220 mg L^{-1} of Cr(VI) for 96 h, a large number of EPS emerged surrounding the cells, and some cells were wrapped up by EPS.

Distribution of Cr in the single cell by soft X-ray spectromicroscopy analysis

The soft X-ray spectromicroscopy analysis was used to study the distribution of different forms of Cr in a single cell of *P. saccharolyticum* LY10. The results indicated that an obvious accumulation of EPS could be observed surrounding the single cell with exposure to 220 mg L^{-1} Cr(VI), even when the cell has been washed with sterilized

water twice prior to analysis (Fig. 4). This result confirmed the conclusion that large amounts of EPS appeared and closely connected with the cells under the toxic Cr(VI) stress, which was previously observed by SEM.

The distribution of Cr(VI) or Cr(III) in a single cell, determined by the soft X-ray spectromicroscopy analysis, was presented in the Fig. 5. The figures indicated that the distribution of both Cr(VI) and Cr(III) was greater in the EPS surrounding the cell, while there was rarely any Cr(VI) and Cr(III) inside the cells. After 24 h incubation, the surface density of Cr(VI) in the single cell ranged from 1.54×10^{-6} to $4.84 \times 10^{-6} \text{ g/cm}^2$, while the surface density of Cr(III) was 1.09×10^{-6} to $2.78 \times 10^{-6} \text{ g/cm}^2$. However, the surface density of Cr(VI) decreased to 1.29×10^{-6} to $2.91 \times 10^{-6} \text{ g/cm}^2$ after exposure for 48 h, while the surface density of Cr(III) increased to 1.76×10^{-6} to $5.22 \times 10^{-6} \text{ g/cm}^2$. The calculated surface density suggested that the Cr(VI) in the EPS was gradually reduced to Cr(III) with incubation time (Fig. 5, 48 h).

Fig. 3 SEM images of *P. saccharolyticum* LY10 cells [a cells were not treated with Cr(VI); b cells were incubated with 55 mg L^{-1} Cr(VI); c 110 mg L^{-1} Cr(VI); d 220 mg L^{-1} Cr(VI)]

Fig. 4 Morphology of a single *P. saccharolyticum* LY10 cell by soft X-ray spectromicroscopy analysis ($3.3\ \mu\text{m} \times 3.3\ \mu\text{m}$)

Discussion

After 96 h of incubation, the Cr(VI) concentration decreased from 110 to $220\ \text{mg L}^{-1}$ to 23.6 and $96.1\ \text{mg L}^{-1}$, respectively, which confirmed the high efficiencies of Cr(VI) removal by *P. saccharolyticum* LY10 (Long et al. 2013). The Cr(VI) reduction by the microorganism could be divided into intracellular and extracellular reduction (Pal and Paul 2008). The Cr(VI) compounds had a tetrahedral structure, a similar molecular structure with SO_4^{2-} . Thus, Cr(VI) could transfer into the cells using the sulfate channels on the cell membrane (Ramirez-Diaz et al. 2008). On the contrary, the Cr(III) compounds had an octahedral structure and were unable to pass the cell membrane (Peterson et al. 1997). Therefore, the extracellularly reduced Cr(III) could not enter the cell membrane, and it was also difficult to discharge the Cr(III) reduced from Cr(VI) intracellularly through the cell membrane. Figure 1 showed that a large percentage of total Cr (more than 92%) was present in the supernatants, most of which was in the form of Cr(III), and only 5.1–7.1% was

Fig. 5 The distribution of different Cr species in a single cell of *P. saccharolyticum* LY10 by soft X-ray spectromicroscopy analysis (24 h: $3.3\ \mu\text{m} \times 3.3\ \mu\text{m}$; 48 h: $3.3\ \mu\text{m} \times 5.4\ \mu\text{m}$)

immobilized by the cells, suggesting that Cr(VI) reduction occurred mainly extracellularly rather than intracellularly.

Previous studies on the subcellular distribution of heavy metals in the cell have already suggested that the intracellular soluble fractions were the main distribution area (Kong et al. 1992; Keim et al. 2001). However, the present study revealed that Cr contents were mostly distributed in the EPS and cell wall compared with other parts in the cell, as shown in Fig. 1b. Considering the Cr(VI) reduction capacity of different fractions, following the order: EPS > soluble fraction of cell > cell membrane > cell wall (Fig. 2), the cell wall had limited capacity of Cr(VI) reduction. Therefore, the removal of Cr(VI) by the strain *P. saccharolyticum* LY10 was mainly achieved extracellularly, and the EPS was the main area for the reduction process.

EPS produced by microorganisms are of particular relevance to the bioremediation process because of their involvement in flocculation and binding of metal ions from solutions (Salehizadeh and Shojaosadati 2003; Kachlany et al. 2001). Because EPS have many functional groups such as hydroxyl, amino, carboxyl, and phosphoric acid groups, the EPS can not only provide numerous binding sites for heavy metal ions but also effectively prevent the entry of toxic heavy metals into the microbial cells, and reduce the damage from toxic heavy metals to the cells (Valdman et al. 2001; Ueshima et al. 2008; Comte et al. 2006; Priester et al. 2006). Previous studies revealed that EPS of a biofilm has a much greater Pb^{2+} and Cd^{2+} fixation capacity than the microbial cells in the water environment (Kang et al. 2005). It was also found that more Cr was removed when large amounts of EPS were produced, suggesting that the bacterial EPS did contribute to the Cr(VI) removal (Jia et al. 2014). The removal of heavy metals by the bacterial EPS would not only increase the sensitivity of bacteria to heavy metals but also significantly reduce the cell immobilization and absorption ability of metal ions (Ueshima et al. 2008; Liu et al. 2002; Fang et al. 2011). Yang et al. (2007) studied the Cr(VI) resistance mechanism of *Enterobacter cloacae* using an atomic force microscope. Their results showed that the cell size of *E. cloacae* changed with the increase of Cr(VI) concentration, and the cell length increased from $2.3 \pm 0.6 \mu m$ without Cr(VI) stress to $3.2 \pm 0.7 \mu m$ under $400 mg L^{-1}$ Cr(VI). Specially, the EPS might be highly produced to hinder the entry of Cr(VI) into the cells, thereby protecting the cells.

The SEM results confirmed that there were no obvious EPS appearing between the cells without Cr(VI) exposure (Fig. 3a). However, with increasing Cr(VI) concentration, the EPS increased and tightly gathered near the cells, which might served as a protection mechanism for the cells of *P. saccharolyticum* LY10 (Fig. 3b–d). This was also confirmed by TEM-EDS analysis, which revealed that at high concentration, Cr(VI) exposure over time resulted in Cr accumulation in the extracellular of *P. saccharolyticum* LY10 in our

previous studies (Long et al. 2013). Further, when the strain *P. saccharolyticum* LY10 was exposed to $220 mg L^{-1}$ Cr(VI) for 96 h, black particles emerged surrounding the cells, and some cells were wrapped up by these extracellular particles. The black particles were further analyzed by energy dispersive X-ray spectrum (EDX), and the results showed that a large amount of Cr (15.1%) was contained in the black particles, whereas the Cr content in the cells was less than 10% (Long et al. 2013). These findings were consistent with results that EPS was the main area of Cr(VI) reduction in *P. saccharolyticum* LY10 which was obtained from the subcellular distribution of Cr in cells and the comparison of Cr(VI) reduction by different subcellular fractions.

The differences in Cr(VI) and Cr(III) distribution revealed that the majority of Cr(VI) was trapped outside the cells in EPS and was gradually reduced to Cr(III). Due to the octahedral structure of Cr(III), it could not pass through the cell membrane (Peterson et al. 1997), and, as a result, accumulated in the EPS and the cell wall (Fig. 5). This finding, again, suggested that the EPS secreted by *P. saccharolyticum* LY10 could most effectively trap Cr(VI) outside the cells and reduced it to Cr(III). Therefore, EPS was the main area of Cr(VI) reduction in the *P. saccharolyticum* LY10. Due to the high concentration ($220 mg L^{-1}$) of Cr(VI), EPS could not trap and reduce all of the Cr(VI) outside the cells. Consequently, a small amount of Cr(VI) with tetrahedral structure got into the cells using the sulfate channels on the cell membrane (Ramirez-Diaz et al. 2008). Subsequently, little Cr(VI) existed after 48 h exposure as shown in Fig. 5.

Microbial reduction of Cr(VI) has been identified as having two major enzymatic mechanisms (Cheung and Gu 2007). Aerobic reduction was thought to be a detoxification mechanism where cells used a soluble enzyme to reduce Cr(VI) to Cr(III) internally or externally to the plasma membrane, while reduction might also proceed through the use of CrO_4^{2-} as a terminal electron acceptor during anaerobic respiration (McLean and Beveridge 2001). Currently, many soluble enzymes have been found in different microorganisms with the ability to reduce chromate (Park et al. 2000; Cheung and Gu 2005, 2007). Cr(VI) served as a non-essential element to microorganisms (Ramirez-Diaz et al. 2008) and had a high cytotoxicity (Coetzee et al. 2018). To protect the cells from toxic damage due to the high concentration of Cr(VI) strain *P. saccharolyticum* LY10 secreted more EPS to wrap the cells. In the EPS Cr(VI) was reduced to Cr(III), and intercepted by adsorption. This combined mechanism could help the cells to survive under high Cr(VI) stress. With the increase in Cr(VI) concentrations, the protection mechanism of *P. saccharolyticum* LY10 became more apparent. The extracellular soluble reductase was the key enzyme found in the EPS and was potentially involved in Cr(VI) reduction (Viti et al. 2014). Compared with the intracellular reduction, extracellular reduction by EPS could efficiently decrease

the toxicity of Cr(VI) for key subcellular components and, provide better protection for the cells. Overall, the results suggested that EPS was the main Cr(VI) reduction site of *P. saccharolyticum* LY10, and an absorbed soluble reductase apparently catalyzed reduction.

Conclusion

The main area of Cr(VI) reduction in the strain of *P. saccharolyticum* LY10 was investigated in the present study. The results indicated that most of Cr was presented in the supernatants in the form of Cr(III) after reduction. In the cell, Cr was mostly distributed in the EPS and cell wall, accounting for more than 90% of the total Cr. However, EPS had the maximum Cr(VI) reduction rate (81.5%) compared with the intracellular fraction, cell wall and cell membrane, suggesting that the main pathway of Cr(VI) removal was extracellular reduction by the EPS rather than intracellular immobilization. The analysis of SEM and TEM confirmed that EPS appeared around the external walls with the Cr(VI) stress applied, and a large amount of Cr was contained in the EPS. Based on the soft X-ray spectromicroscopy analysis, the distribution of the Cr(VI) or Cr(III) in a single cell was much greater in the EPS surrounding the cell, while there was rarely chromium inside the cell. Therefore, we concluded that EPS were of great importance in the Cr(VI) resistance and reduction in the bacteria of *P. saccharolyticum* LY10 and were the main area for Cr(VI) reduction.

Acknowledgements This study was financially supported by the Hangzhou Science & Technology Planning Program (20120433B46) and National High-Tech Research and Development Program of China (2009AA063101). The authors acknowledge Zhi Guo, Xiangzhi Zhang, Lijuan Zhang and Haigang Li at soft X-ray beamline BL08U of Shanghai Synchrotron Radiation Facility (SSRF) for STXM imaging.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

References

- Beukes J, Du Preez S, Van Zyl P, Paktunc D, Fabritius T, Pääta M, Cramer M (2017) Review of Cr(VI) environmental practices in the chromite mining and smelting industry—relevance to development of the Ring of Fire, Canada. *J Clean Prod* 165:874–889
- Cervantes C, Campos-García J, Devars S, Gutiérrez-Corona F, Loza-Tavera H, Torres-Guzmán JC, Moreno-Sánchez R (2001) Interactions of chromium with microorganisms and plants. *FEMS Microbiol Rev* 25(3):335–347
- Chai L, Huang S, Yang Z, Peng B, Huang Y, Chen Y (2009) Cr(VI) remediation by indigenous bacteria in soils contaminated by chromium-containing slag. *J Hazard Mater* 167(1):516–522
- Chao W, Harteneck BD, Liddle JA, Anderson EH, Attwood DT (2005) Soft X-ray microscopy at a spatial resolution better than 15 nm. *Nature* 435(7046):1210–1213
- Chen G (2011) Molecular mechanisms of heavy metals tolerance and accumulation in unsaturated *Pseudomonas putida* CZ1 biofilm. Dissertation, Zhejiang University
- Chen Z, Huang Z, Cheng Y, Pan D, Pan X, Yu M, Pan Z, Lin Z, Guan X, Wu Z (2012) Cr(VI) uptake mechanism of *Bacillus cereus*. *Chemosphere* 87(3):211–216
- Cheung K, Gu J-D (2005) Chromate reduction by *Bacillus megaterium* TKW3 isolated from marine sediments. *World J Microbiol Biotechnol* 21(3):213–219
- Cheung K, Gu J-D (2007) Mechanism of hexavalent chromium detoxification by microorganisms and bioremediation application potential: a review. *Int Biodeterior Biodegrad* 59(1):8–15
- Coetzee JJ, Bansal N, Chirwa EM (2018) Chromium in environment, its toxic effect from chromite-mining and ferrochrome industries, and its possible bioremediation. *Expo Health*. <https://doi.org/10.1007/s12403-018-0284-z>
- Comte S, Guibaud G, Baudu M (2006) Relations between extraction protocols for activated sludge extracellular polymeric substances (EPS) and EPS complexation properties: Part I. Comparison of the efficiency of eight EPS extraction methods. *Enzyme Microb Technol* 38(1):237–245
- Dogan NM, Kantar C, Gulcan S, Dodge CJ, Yimaz BC, Mazmanci MA (2011) Chromium(VI) bioremoval by pseudomonas bacteria: role of microbial exudates for natural attenuation and biotreatment of Cr(VI) contamination. *Environ Sci Technol* 45(6):2278–2285. <https://doi.org/10.1021/Es1020951>
- Fang L, Wei X, Cai P, Huang Q, Chen H, Liang W, Rong X (2011) Role of extracellular polymeric substances in Cu(II) adsorption on *Bacillus subtilis* and *Pseudomonas putida*. *Bioresour Technol* 102(2):1137–1141
- Flemming H-C, Wingender J (2010) The biofilm matrix. *Nat Rev Microbiol* 8(9):623–633
- Fulladosa E, Desjardin V, Murat J-C, Gourdon R, Villaescusa I (2006) Cr(VI) reduction into Cr(III) as a mechanism to explain the low sensitivity of *Vibrio fischeri* bioassay to detect chromium pollution. *Chemosphere* 65(4):644–650
- Jacobsen C, Wirick S, Flynn G, Zimba C (2000) Soft X-ray spectroscopy from image sequences with sub-100 nm spatial resolution. *J Microsc Oxf* 197:173–184
- Jia Y, Huang H, Chen Z, Zhu Y-G (2014) Arsenic uptake by rice is influenced by microbe-mediated arsenic redox changes in the rhizosphere. *Environ Sci Technol* 48(2):1001–1007
- Kachlany SC, Levery SB, Kim JS, Reuhs BL, Lion LW, Ghiorse WC (2001) Structure and carbohydrate analysis of the exopolysaccharide capsule of *Pseudomonas putida* G7. *Environ Microbiol* 3(12):774–784
- Kang C, Guo J, Guo P, Zhao Y, Dong D (2005) Pb²⁺ adsorption characteristics by organic components of biofilms and influencing factors in natural water. *Chem J Chin Univ* 26(11):2043–2045
- Kavita K, Mishra A, Jha B (2013) Extracellular polymeric substances from two biofilm forming *Vibrio* species: characterization and applications. *Carbohydr Polym* 94(2):882–888
- Keim CN, Lins U, Farina M (2001) Elemental analysis of uncultured magnetotactic bacteria exposed to heavy metals. *Can J Microbiol* 47(12):1132–1136
- Kilic NK, Stensballe A, Otzen DE, Donmez G (2010) Proteomic changes in response to chromium(VI) toxicity in *Pseudomonas aeruginosa*. *Bioresour Technol* 101(7):2134–2140
- Kong S, Yonge DR, Johnstone DL, Petersen JN, Brouns TM (1992) Chromium distribution in subcellular components between fresh and starved subsurface bacterial consortium. *Biotechnol Lett* 14(6):521–524

- Lai KC, Lo IM (2008) Removal of chromium(VI) by acid-washed zero-valent iron under various groundwater geochemistry conditions. *Environ Sci Technol* 42(4):1238–1244
- Lan G, Hong X, Fan Q, Luo B, Shi P, Chen X (2014) Removal of hexavalent chromium in wastewater by polyacrylamide modified iron oxide nanoparticle. *J Appl Polym Sci* 131(20):440–450
- Lee D-C, Park C-J, Yang J-E, Jeong Y-H, Rhee H-I (2000) Screening of hexavalent chromium biosorbent from marine algae. *Appl Microbiol Biotechnol* 54(3):445–448
- Liu C, Gorby YA, Zachara JM, Fredrickson JK, Brown CF (2002) Reduction kinetics of Fe(III), Co(III), U(VI), Cr(VI), and Tc(VII) in cultures of dissimilatory metal-reducing bacteria. *Biotechnol Bioeng* 80(6):637–649
- Long D, Tang X, Cai K, Chen G, Chen L, Duan D, Zhu J, Chen Y (2013) Cr(VI) reduction by a potent novel alkaliphilic halotolerant strain *Pseudochrobactrum saccharolyticum* LY10. *J Hazard Mater* 256:24–32
- Lovley DR (1993) Dissimilatory metal reduction. *Ann Rev Microbiol* 47(1):263–290
- Ma L, Xu J, Chen N, Li M, Feng C (2019) Microbial reduction fate of chromium (Cr) in aqueous solution by mixed bacterial consortium. *Ecotoxicol Environ Saf* 170:763–770
- Mazoch J, Tesářík R, Sedláček V, Kučera I, Turánek J (2004) Isolation and biochemical characterization of two soluble iron (III) reductases from *Paracoccus denitrificans*. *Eur J Biochem* 271(3):553–562
- McLean J, Beveridge TJ (2001) Chromate reduction by a pseudomonad isolated from a site contaminated with chromated copper arsenate. *Appl Environ Microbiol* 67(3):1076–1084
- Morel MA, Ubalde MC, Braña V, Castro-Sowinski S (2011) Delftia sp. JD2: a potential Cr(VI)-reducing agent with plant growth-promoting activity. *Arch Microbiol* 193(1):63–68
- Morokutti A, Lyskowski A, Sollner S, Pointner E, Fitzpatrick TB, Kratky C, Gruber K, Macheroux P (2005) Structure and function of YcnD from *Bacillus subtilis*, a flavin-containing oxidoreductase. *Biochemistry* 44(42):13724–13733
- Pal A, Paul A (2008) Microbial extracellular polymeric substances: central elements in heavy metal bioremediation. *Indian J Microbiol* 48(1):49–64
- Park C, Keyhan M, Wielinga B, Fendorf S, Matin A (2000) Purification to homogeneity and characterization of a novel *Pseudomonas putida* chromate reductase. *Appl Environ Microbiol* 66(5):1788–1795
- Peterson ML, Brown GE, Parks GA, Stein CL (1997) Differential redox and sorption of Cr(III/VI) on natural silicate and oxide minerals: EXAFS and XANES results. *Geochim Cosmochim Acta* 61(16):3399–3412. [https://doi.org/10.1016/S0016-7037\(97\)00165-8](https://doi.org/10.1016/S0016-7037(97)00165-8)
- Priester JH, Olson SG, Webb SM, Neu MP, Hersman LE, Holden PA (2006) Enhanced exopolymer production and chromium stabilization in *Pseudomonas putida* unsaturated biofilms. *Appl Environ Microbiol* 72(3):1988–1996
- Ramirez-Diaz MI, Diaz-Perez C, Vargas E, Riveros-Rosas H, Campos-Garcia J, Cervantes C (2008) Mechanisms of bacterial resistance to chromium compounds. *Biomaterials* 21(3):321–332. <https://doi.org/10.1007/s10534-007-9121-8>
- Salehizadeh H, Shojasodati S (2003) Removal of metal ions from aqueous solution by polysaccharide produced from *Bacillus firmus*. *Water Res* 37 (17):4231–4235
- Smith N (2001) Science with soft X rays. *Phys Today* 54(1):29–54
- Su X, Wang Y, Xue B, Hashmi MZ, Lin H, Chen J, Wang Z, Mei R, Sun F (2019) Impact of resuscitation promoting factor (Rpf) in membrane bioreactor treating high-saline phenolic wastewater: performance robustness and Rpf-responsive bacterial populations. *Chem Eng J* 357:715–723
- Tebo BM, Obraztsova AY (1998) Sulfate-reducing bacterium grows with Cr(VI), U(VI), Mn(IV), and Fe(III) as electron acceptors. *FEMS Microbiol Lett* 162(1):193–198. <https://doi.org/10.1111/j.1574-6968.1998.tb12998.x>
- Ueshima M, Ginn BR, Haack EA, Szymanowski JE, Fein JB (2008) Cd adsorption onto *Pseudomonas putida* in the presence and absence of extracellular polymeric substances. *Geochim Cosmochim Acta* 72 (24):5885–5895
- Valdman E, Erijman L, Pessoa F, Leite S (2001) Continuous biosorption of Cu and Zn by immobilized waste biomass *Sargassum* sp. *Process Biochem* 36 (8):869–873
- Valls M, Atrian S, de Lorenzo V, Fernández LA (2000) Engineering a mouse metallothionein on the cell surface of *Ralstonia eutropha* CH34 for immobilization of heavy metals in soil. *Nat Biotechnol* 18(6):661–665
- Viti C, Marchi E, Decorosi F, Giovannetti L (2014) Molecular mechanisms of Cr(VI) resistance in bacteria and fungi. *FEMS Microbiol Rev* 38(4):633–659
- Wang J, Li Q, Li M-M, Chen T-H, Zhou Y-F, Yue Z-B (2014a) Competitive adsorption of heavy metal by extracellular polymeric substances (EPS) extracted from sulfate reducing bacteria. *Bioresour Technol* 163:374–376
- Wang Z, Gao M, Wang S, Xin Y, Ma D, She Z, Wang Z, Chang Q, Ren Y (2014b) Effect of hexavalent chromium on extracellular polymeric substances of granular sludge from an aerobic granular sequencing batch reactor. *Chem Eng J* 251:165–174
- Wielinga B, Mizuba MM, Hansel CM, Fendorf S (2001) Iron promoted reduction of chromate by dissimilatory iron-reducing bacteria. *Environ Sci Technol* 35(3):522–527. <https://doi.org/10.1021/Es001457b>
- Yang C, Cheng Y, Ma X, Zhu Y, Holman HY, Lin Z, Wang C (2007) Surface-mediated chromate-resistant mechanism of *Enterobacter cloacae* bacteria investigated by atomic force microscopy. *Langmuir* 23(8):4480–4485
- Zhong Q-Q, Yue Q-Y, Li Q, Gao B-Y, Xu X (2014) Removal of Cu(II) and Cr(VI) from wastewater by an amphoteric sorbent based on cellulose-rich biomass. *Carbohydr Polym* 111:788–796. <https://doi.org/10.1016/j.carbpol.2014.05.043>