Published in final edited form as:

Alcohol Clin Exp Res. 2019 March ; 43(3): 522–530. doi:10.1111/acer.13953.

Association between gabapentin receipt for any indication and AUDIT-C scores among clinical sub-populations with and without alcohol use disorder

Christopher T. Rentsch, PhD^{1,2,3}, David A. Fiellin, MD^{2,4}, Kendall J. Bryant, PhD⁵, Amy C. Justice, MD, PhD^{1,2,4}, and Janet P. Tate, ScD^{1,2}

¹Veterans Aging Cohort Study Coordinating Center, VA Connecticut Healthcare System, West Haven, CT, 06516, USA

²Department of Internal Medicine, Yale School of Medicine, New Haven, CT, 06511, USA

³Faculty of Epidemiology & Population Health, London School of Hygiene & Tropical Medicine, London, WC1E 7HT, UK

⁴Center for Interdisciplinary Research on AIDS, Yale School of Public Health, New Haven, CT, 06511, USA

⁵Director of HIV/AIDS Research, National Institute on Alcohol Abuse and Alcoholism, Bethesda, MD, 20892, USA

Abstract

Background: Current medications for alcohol use disorder (AUD) have limited efficacy and utilization. Some clinical trials have shown efficacy for gabapentin among treatment-seeking individuals. The impact of gabapentin on alcohol consumption in a more general sample remains unknown.

Methods: We identified patients prescribed gabapentin for 180 consecutive days for any clinical indication other than substance use treatment between 2009 and 2015 in the Veterans Aging Cohort Study. We propensity-score matched each gabapentin exposed patient with up to five unexposed patients. Multivariable difference-in-difference (DiD) linear regression models estimated the differential change in Alcohol Use Disorders Identification Test – Consumption (AUDIT-C) scores during follow-up between exposed and unexposed patients, by baseline level of alcohol consumption, and daily gabapentin dose. Analyses were stratified by AUD history. Clinically meaningful changes were *a priori* considered a DiD 1 point.

Results: Among patients with AUD, AUDIT-C scores decreased 0.39 points (95% CI 0.05, 0.73) more among exposed than unexposed patients ($p < 0.03$). Potentially clinically meaningful differences were observed among those with AUD and exposed to 1,500 milligrams/day (DiD 0.77, 95% CI 0.15, 1.38; $p < 0.02$). No statistically significant effects were found among patients with AUD at doses lower than 1,500 mg/day or baseline AUDIT-C 4. Among patients without

Corresponding author: Christopher T. Rentsch, PhD, VA Connecticut Healthcare System, Yale University School of Medicine, London School of Hygiene & Tropical Medicine, Keppel Street, WC1E 7HT, London, UK, Christopher.Rentsch@va.gov.

Conflicts of interest: The authors declare no conflict of interest.

AUD, we found no overall difference in changes in AUDIT-C scores, nor in analyses stratified by baseline level of alcohol consumption.

Conclusions: Patients exposed to doses of gabapentin consistent with those used in clinical trials, particularly those with AUD, experienced a greater decrease in AUDIT-C scores than matched unexposed patients.

Keywords

gabapentin; alcohol use disorder; electronic health records; propensity score

INTRODUCTION

Medications and counselling, although underused, are the most effective treatments for patients with alcohol use disorder (AUD) (Jonas et al., 2014, Magill et al., 2015). The efficacy of the three current medications approved by the U.S. Food and Drug Administration (FDA) for the treatment of AUD (i.e., naltrexone, acamprosate, and disulfiram) is limited (Lyon, 2017, Winslow et al., 2016, Litten et al., 2016a, Kranzler and Soyka, 2018), and novel strategies are actively being investigated (Koob and Mason, 2016, Koob and Volkow, 2016, Litten et al., 2016a). This has led researchers to assess the impact of a range of medications approved by the FDA for other indications on alcohol use, including topiramate, varenicline, baclofen, and gabapentin (Litten et al., 2016b, Soyka and Muller, 2017, Kranzler and Soyka, 2018).

Gabapentin, a structural analogue to gamma-aminobutyric acid (GABA), is FDA approved for treatment of partial seizure and postherpetic neuralgia and has shown some efficacy for treatment of AUD in treatment-seeking individuals (Falk et al., 2018, Mason et al., 2014). Anticonvulsants such as gabapentin are believed to decrease craving and alter the subjective effects of alcohol leading to decreased risk of relapse (Pani et al., 2014), although the mechanisms of action are not completely elucidated. A Cochrane Collaborative meta-analysis of three efficacy trials comparing gabapentin to placebo demonstrated that gabapentin use was associated with greater abstinence (decreased drinking days), and reduced heavy drinking, although it had no impact on percent days abstinent or craving (Pani et al., 2014). One efficacy trial comparing gabapentin to placebo demonstrated a dose-response effect on drinking outcomes among patients with Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV) alcohol dependence (American Psychiatric Association, 1994). In the 12-week trial, abstinence was 4% with placebo, 11% with 900 milligrams (mg) daily dose of gabapentin, and 17% with 1800 mg/day ($p=0.04$ for linear dose effect) (Mason et al., 2014). Similar benefits were seen for no heavy drinking and craving (Mason et al., 2014). These results support further evaluation of gabapentin among diverse patient populations.

FDA-approved medications for treatment of AUD are underused by specialists and generalist (e.g. primary care) providers (Finlay et al., 2017, Ford et al., 2017, Lee et al., 2015, Owens et al., 2018, Jonas et al., 2014, Mark et al., 2015, Mark et al., 2009, Mark et al., 2003a, Cohen et al., 2007, Williams et al., 2017, Harris et al., 2010, Harris et al., 2012). Limited efficacy, low patient demand, formulary restrictions, and lack of experience with

these medications for AUD treatment are known concerns among providers (Mark et al., 2009, Mark et al., 2003a, Mark et al., 2003b, Williams et al., 2018, Harris et al., 2013). However, a potential advantage of gabapentin for treatment of AUD is that it is commonly prescribed “off-label” for peripheral neuropathy, fibromyalgia, and other painful conditions (Shanthanna et al., 2017, Kesselheim et al., 2011), which suggests that clinicians have a level of comfort and familiarity with its use.

The impact of gabapentin on alcohol consumption when prescribed for indications other than treatment of AUD is unknown. Previous work has demonstrated that lower levels of alcohol exposure are associated with greater risk of physiologic injury and mortality among HIV-infected patients compared to uninfected patients (Justice et al., 2016) and low levels of alcohol negatively impacts prognosis in a range of medical conditions such as depression and liver disease (Sullivan et al., 2011, Sullivan et al., 2005, Lim et al., 2014). Therefore, we sought to determine the impact of gabapentin on changes in alcohol use among patients receiving gabapentin for common medical conditions, who reported any alcohol consumption and whether effects differed by AUD history, baseline level of alcohol consumption, and prescribed daily dose of gabapentin. We hypothesized that the effect of gabapentin on alcohol consumption would be greater among those with AUD, higher baseline levels of alcohol consumption, and prescribed higher doses.

MATERIALS AND METHODS

Study population

We used data from the Veterans Aging Cohort Study (VACS), which has been described in detail (Fultz et al., 2006, Justice et al., 2006). Briefly, VACS is a large observational cohort based on data from the national Veterans Health Administration (VA) electronic health records (EHR) that includes all HIV-infected patients in VA care (>50,000 HIV+ subjects) and uninfected patients (>100,000) 1:2 matched on region, age, race/ethnicity, and sex. VACS has been approved by the institutional review boards of the VA Connecticut Healthcare System and Yale School of Medicine, granted a waiver of informed consent, and deemed Health Insurance Portability and Accountability Act compliant.

For this analysis, we included HIV+ and HIV-uninfected patients who did (gabapentin exposed) and did not (gabapentin unexposed) receive gabapentin dispensed at VA pharmacies. For the gabapentin exposed group, we included all patients who received gabapentin for at least 180 continuous days for any indication between 1 January 2009 and 30 September 2015 from the following VA clinics: primary care, infectious disease, neurology, general internal medicine, physical medicine and rehabilitation services, pain, podiatry, orthopedics, women’s clinic, and rheumatology. These clinics were chosen because they were the source of most gabapentin prescriptions. To ensure that unexposed patients came from the same source population and had an equal opportunity to receive gabapentin, we randomly selected one outpatient visit date per calendar year to identify patients who attended one of the listed clinics but never received gabapentin. Importantly, we did not include patients with gabapentin prescriptions from substance use treatment programs; however, we did not exclude patients who subsequently visited a substance use treatment program during follow-up.

To allow us to follow exposed and unexposed patients over similar calendar time, we created an “index date” (also referred to as “baseline”) which was defined as the first fill date for gabapentin exposed patients and the random outpatient visit date for unexposed patients. We utilized a 12-month washout period to identify new episodes of gabapentin exposure. Therefore, patients who received gabapentin in 2008 were only eligible to be followed after a one-year period of no gabapentin exposure. We excluded patients who had no outpatient care in the VA in the year prior to their index date and those who had no measurement of alcohol consumption in the two years prior to their index date. We also excluded patients who reported no alcohol consumption based on the closest measurement to baseline.

Propensity score model and matching

To address concerns of confounding by indication, whereby patients with specific alcohol consumption patterns might be more likely to receive gabapentin, we generated propensity scores. Propensity scores are used to adjust for the conditional probability of being prescribed gabapentin given a set of covariates that are associated with both gabapentin receipt and alcohol consumption or associated with alcohol consumption only (Brookhart et al., 2006). Matching by propensity score provides a means to balanced exposure groups similar to random treatment allocation in a randomized controlled trial (Austin, 2011). We hypothesized that the effects of gabapentin on alcohol consumption may differ in patients with and without diagnosed AUD prior to their index date. Therefore, propensity scores (i.e. the predicted probability of gabapentin exposure) were estimated using separate multivariable logistic regression models for patients with and without AUD at baseline, as defined below. Estimating propensity scores separately has been shown to be unbiased, particularly in subgroup analyses with small sample sizes (Eeren et al., 2015, Green and Stuart, 2014, Rassen et al., 2012).

Variables used in the propensity score models were selected a priori based on clinical knowledge (Hernan et al., 2002) and included: year of index date, age at baseline, race, smoking status, body mass index at baseline, site prescribing pattern (the proportion of patients who initiated gabapentin stratified by year and HIV status), lab values closest to the index date (including haemoglobin, international normalized ratio, triglycerides, CD4 cell count), hepatitis C virus (HCV) status, HIV status, history of seizure prior to baseline, diabetes complications severity index (Young et al., 2008) at baseline, history of pain diagnoses prior to baseline (including neuropathy, osteoarthritis, or pain in the abdomen, back, chest, extremity, or neck, headache, or fracture), and history of medical and psychiatric conditions prior to baseline (including atrial fibrillation, myocardial infarction/coronary artery disease, peripheral vascular disease, diabetes, nephrolithiasis, glomerulonephritis, hyperlipidemia, pancreatitis, drug use disorders, post-traumatic stress disorder (PTSD), major or other depression, anxiety, bipolar disorder, schizophrenia and schizoaffective disorder). We also included variables that captured attendance to clinics (including primary care, dialysis, diabetic retinal screening, rheumatology, infectious disease, nephrology, neurology, pain, allergy, chiropractic, dental, diabetes, emergency department, electrocardiogram lab, ophthalmology, hematology, oncology, homeless program, nutrition, orthopedics, substance use, mental health, PTSD), frequency of all-cause hospitalizations, and the total number of unique clinics visited in the year prior to baseline.

Lastly, variables denoting receipt of other prescriptions (starting on or elapsing baseline) to treat pain (including non-steroidal anti-inflammatory drugs (NSAIDs), opioids, muscle relaxants, and antidepressants) and seizures were included in the model. Interaction terms were explored for significance, and six were kept in the final model (all $p < 0.05$). The model c-statistic was 0.83 for patients with AUD and 0.84 in patients without AUD, indicating adequate discrimination between gabapentin exposed and unexposed patients in both models (Hosmer and Lemeshow, 2000).

Since the distribution of propensity scores for exposed patients was different than that of unexposed patients, we used propensity score matching to exclude non-exchangeable unexposed patients (Figure 1) (Spendlin et al., 2016). We conducted propensity score matching within pre-specified subgroups of patients based on baseline Alcohol Use Disorders Identification Test – Consumption (AUDIT-C) scores and aggregated these subgroup strata to create the full matched cohort (Wang et al., 2018). Each exposed patient was matched to up to five unexposed patients with index dates in the same calendar year, using a greedy matching algorithm (Cormen, 2009).

Measures and follow-up

Baseline AUD was determined by one inpatient or two outpatient ICD-9 codes (303.X or 305–305.03) at any time prior to baseline. Alcohol consumption was assessed using the AUDIT-C, a three-question self-report alcohol screening questionnaire that detects heavy drinking and/or active AUD (Bush et al., 1998, Fiellin et al., 2000). AUDIT-C scores range from 0–12 with the likelihood of physiologic injury and mortality increasing as AUDIT-C scores increase (Justice et al., 2016). An AUDIT-C score of zero is defined as no current alcohol use, 1–3 suggests lower-risk drinking, 4–7 suggests at-risk drinking, and 8 suggest hazardous or heavy episodic alcohol consumption. Since 2007, the VA has required annual AUDIT-C screening on all patients in primary care (Bradley et al., 2006).

All patients were followed from their index date for a maximum of two years or until their last VA visit or death. Additionally, gabapentin exposed patients were censored at 30 days after the end of their last gabapentin prescription (allowing for a maximum 30-day gap between fills). To ensure equal follow-up time within matched sets, unexposed patients were censored at the total follow-up time of their matched exposed patient.

Statistical analyses

All statistical analyses were performed separately for patients with and without AUD at baseline. While evidence of alcohol consumption at baseline as measured by the AUDIT-C was a criterion for study inclusion, we did not restrict matching eligibility on the availability of a follow-up AUDIT-C (the outcome) as such a restriction would not be available at baseline in an analogous randomized clinical trial. Thus, 1,119 (44%) exposed patients in the propensity score matched sample did not have a follow-up AUDIT-C and were unable to be included in regression models. If an exposed patient did not have a follow-up AUDIT-C, we removed their entire matched set of unexposed patients to maintain a balanced sample. If an unexposed patient did not have a follow-up AUDIT-C, we kept the remaining patients in their matched set in the analytic sample as long as there was another unexposed patient in

the set. We used chi-square tests to examine balance between exposed and unexposed patients included in the full sample, propensity-score matched sample, and final analytic sample.

Among those in the final analytic sample, we calculated the average pre- and post-index AUDIT-C scores. Pre-index AUDIT-C scores were defined as the closest on or before the index date, within a maximum of two years. Post-index AUDIT-C scores were defined as the closest measure to the end of exposure or within 30 days of end of follow-up. We then used multivariable difference-in-difference (DiD) linear regression models (Donald and Lang, 2007, Lechner, 2011) to estimate the differential change between pre- and post-index AUDIT-C scores. We *a priori* considered a DiD estimate 1 point clinically meaningful (Rubinsky et al., 2013). To account for residual confounding not captured by propensity score matching, models were adjusted for any characteristic shown to be unbalanced between exposed and unexposed patients in addition to age, total number of medications prescribed during follow-up, and VACS Index. The VACS Index – a measure of physiologic injury incorporating age, CD4 count, HIV-1 RNA, hemoglobin, a marker of liver fibrosis (FIB-4), estimated glomerular filtration rate (eGFR), and HCV status – has been shown to predict acquired immunodeficiency syndrome (AIDS) and non-AIDS morbidity and mortality in multiple settings (Akgun et al., 2013, Akgun et al., 2014, Escota et al., 2015, Justice et al., 2013, Marquine et al., 2014, Tate et al., 2013, Womack et al., 2013).

We also performed subgroup analyses by self-reported level of alcohol consumption at baseline (as determined by AUDIT-C) and average daily gabapentin dose during follow-up. Daily dose was categorized to include roughly equal numbers of patients in each group: low (<600 milligrams [mg]), medium (600 mg-1,499 mg), and high (1,500 mg). Lastly, we conducted a sensitivity analysis excluding patients with a visit to a substance use treatment program during follow-up. All statistical analyses were performed using SAS version 9.4 (SAS Institute Inc., Cary, NC, USA).

RESULTS

Sample

We identified 5,721 gabapentin exposed patients and 52,243 gabapentin unexposed patients who reported any alcohol consumption in the two years prior to their index date. A total 2,520 exposed patients were matched (Supplemental Table 1); however, 1,119 (44%) did not have a follow-up AUDIT-C and were unable to be included in analysis. Among those with AUD in the final analytic sample, 29 (5.2%) were matched to five unexposed patients, 55 (9.8%) to four, 86 (15.3%) to three, 121 (21.5%) to two, and 271 (48.2%) to one unexposed patient. Among patients without AUD, 87 (10.4%) were matched to five unexposed patients, 125 (14.9%) to four, 124 (14.8%) to three, 167 (19.9%) to two, and 336 (40.0%) to one unexposed patient. Thus, the final analytic sample consisted of 562 exposed and 1,136 unexposed patients with AUD, and 839 exposed and 1,977 unexposed patients without AUD.

Prior to propensity score matching, the distribution of baseline characteristics significantly differed between gabapentin exposed and unexposed patients with or without AUD (Table

1). In the final analytic sample, gabapentin exposed and unexposed patients with or without AUD were well balanced (Table 1). There was a statistically significant difference in the proportion of antidepressant prescriptions at baseline among patients with AUD and the proportion of neuropathic pain diagnoses among patients without AUD. These covariates were included in adjusted models.

Gabapentin exposed patients who were not matched had higher propensity scores than those who were matched (median 0.23, interquartile range [IQR] 0.12–0.43 not matched; median 0.05, IQR 0.03–0.08 matched). Median follow-up time was 334 days (IQR 237–475 days) for patients with AUD and 385 days (IQR 266–574 days) for patients without AUD. Among exposed patients in the final sample, 31% were prescribed daily doses of gabapentin <600 mg, 44% were prescribed between 600–1,500 mg, and 25% were prescribed 1,500 mg.

Changes in AUDIT-C scores

There was no difference in the distribution of time between post-index AUDIT-C measures and end of follow-up between exposed and unexposed patients (Kruskal-Wallis $p=0.11$). Median difference between end of follow-up and post-index AUDIT-C was 106 days (IQR 30–195 days).

Overall, AUDIT-C scores decreased during the study period regardless of AUD history, baseline AUDIT-C, or gabapentin dose. Among patients with AUD, average AUDIT-C scores decreased from 4.16 (standard deviation [SD] 0.13) to 3.15 (SD 0.13) among exposed patients and 3.94 (SD 0.10) to 3.32 (SD 0.10) among unexposed patients (Table 2). The adjusted DiD estimate was statistically significant but the confidence interval did not include our *a priori* threshold for a clinically meaningful difference (DiD 0.39 points, 95% confidence interval [CI] 0.05, 0.73; $p=0.0264$, Figure 2). In analysis stratified by baseline AUDIT-C and among those with AUDIT-C of 1–3, average scores decreased 0.03 points among exposed and increased 0.57 points among unexposed patients (DiD 0.59, 95% CI 0.20, 0.99; $p=0.0032$). No significant differences were observed for higher baseline AUDIT-C. The largest DiD estimate was seen among patients with AUD at baseline and exposed to 1,500 mg/day of gabapentin (DiD 0.77 points, 95% CI 0.15, 1.38; $p=0.0149$), which was statistically significant and the confidence interval included our criteria for a clinically meaningful difference.

Patients without AUD had lower pre-index AUDIT-C scores and smaller decreases in AUDIT-C scores. Average AUDIT-C scores decreased from 2.61 (SD 0.07) to 2.02 (SD 0.07) among exposed patients and 2.49 (SD 0.05) to 2.05 (SD 0.05) among unexposed patients (Table 2). The adjusted DiD estimate was not statistically significant (DiD 0.14, 95% CI –0.01, 0.30; $p=0.0691$) (Table 2). The only statistically significant DiD estimate among patients without AUD was among those exposed to <600 mg/day of gabapentin (DiD 0.37 points, 95% CI 0.12, 0.61; $p=0.0034$), but this was not clinically meaningful. In sensitivity analyses excluding patients with a visit to a substance use treatment program during follow-up, DiD estimates were of a similar magnitude and direction although with wider confidence intervals due to smaller sample sizes (Supplemental Table 2).

DISCUSSION

This propensity-score matched analysis of the association between gabapentin use, prescribed for any indication, and patients' reported alcohol consumption found an overall statistically significant but not clinically meaningful difference in changes in AUDIT-C scores among patients with AUD. Analyses restricted to patients with AUD and exposed to 1,500 mg/day of gabapentin suggested a statistically significant and potentially clinically meaningful decrease in reported alcohol consumption. No other statistically significant effects were found among patients with AUD at doses lower than 1,500 mg/day. Interestingly, analyses among patients with AUD who reported low levels of alcohol consumption at baseline demonstrated no change in alcohol consumption among exposed and an increase in alcohol consumption among unexposed, resulting in a statistically significant DiD estimate. No other statistically significant effects were found among patients with AUD at doses lower than 1,500 mg/day or baseline AUDIT-C 4.

Among patients without AUD, we found no overall difference in changes in AUDIT-C scores, nor in analyses stratified by baseline level of alcohol consumption. While no effect was found among patients without AUD at doses 600 mg/day, analyses restricted to those prescribed <600 mg/day demonstrated a statistically significant but not clinically meaningful difference in changes in reported alcohol use. As shown in Table 2, this association was driven by an increase in reported alcohol consumption among unexposed patients and not a decrease among exposed patients.

One efficacy study evaluated the impact of gabapentin dose on alcohol-related outcomes. This randomized clinical trial, with a high dropout rate (43%, 65/150), noted a dose-response effect with improved outcomes at 1800 mg/day compared with placebo and 900 mg/day (Mason et al., 2014). Investigators chose these doses based on FDA-approved dose ranges for seizure and neuropathic pain. Our findings also demonstrate a greater impact of gabapentin at higher doses, which we defined as 1500 mg/day because we were not powered to limit to 1800 mg/day. Preliminary findings from another recent efficacy trial of a prodrug formulation of gabapentin, called gabapentin enacarbil, found no effect of any drinking measure among 346 patients with moderate or severe AUD (Falk et al., 2018). However, Falk et al suggest these results may be partially explained by the differential FDA-approved dosage of gabapentin and gabapentin enacarbil. The mechanism of action for gabapentin in the treatment of unhealthy alcohol use is not completely understood. Its activity is presumably related to the ability to increase or modulate GABA activity via voltage dependent calcium channels and direct synthesis (Leung et al., 2015). Its clinical anxiolytic and sedative effects may address withdrawal and craving in a dose-dependent manner among those with AUD. The impact of gabapentin on craving, however, is not clear (Pani et al., 2014).

This research differs from recent efficacy studies of the impact of gabapentin on alcohol consumption in a number of important ways. First, we evaluated the impact of gabapentin on alcohol consumption in a real-world setting among patients who did not receive their gabapentin prescription via a substance use treatment program. We addressed methodological challenges inherent to observational study designs by using uniform

exclusion criteria for exposed and unexposed patients, evaluating incident exposures, setting an index date for exposed and unexposed patients, and using propensity score matching to account for confounding by indication. Second, previous studies have shown that motivation to receive treatment for AUD can impact treatment outcomes (DiClemente et al., 2017, DiClemente et al., 2004, Field et al., 2009). Given that patients in our sample were prescribed gabapentin from non-substance use treatment providers, it is likely that patients were not receiving gabapentin to address their alcohol use. Nonetheless, more than one-third of exposed and unexposed patients with AUD were seen in a substance use treatment program during follow up. Findings from sensitivity analyses excluding patients with a visit to a substance use treatment program were largely consistent with our primary findings, though with less precision given the smaller sample sizes. Therefore, our findings may underestimate the impact of gabapentin in those who might be more motivated to treat their AUD. Third, we determined the association between gabapentin exposure and alcohol use in patients with any level of alcohol consumption. Decreasing alcohol consumption in patient populations who do not meet formal criteria for unhealthy alcohol use or AUD might result in improvement of other conditions such as HIV, depression and liver fibrosis (Lim et al., 2014, Sullivan et al., 2005, Sullivan et al., 2011, Justice et al., 2016). One of the advantages of using real-world, observational data to examine the impact of gabapentin exposure on alcohol consumption is the ability to determine whether an effect exists across a wide range of drinking behaviors. Some patients in our study who reported unhealthy alcohol use did not have diagnosed AUD, and notably gabapentin did not seem to have a clinically meaningful impact on their AUDIT-C scores.

There are limitations to our work. First, our sample was restricted to U.S. Veterans who were receiving care in the VA healthcare system, so our findings may not generalize to Veterans who did not receive care in the VA or to other patient populations. Second, due to the VACS sampling strategy and characteristics of the Veteran population, our sample was enriched with older men and patients with multiple medical comorbidities including HIV infection, which reflects a segment of patients aging with HIV disease but may not generalize to other clinical settings. Compared to estimates from the Veteran Population Projection Model 2016, our analytic sample accurately represented Veterans aged 60+, under represented younger Veterans, and somewhat over represented middle-aged Veterans. With respect to race/ethnicity, our sample over represented black Veterans. Third, AUDIT-C scores were collected as part of routine clinical care and may not reflect actual drinking patterns (Williams et al., 2015, McGinnis et al., 2016, Bradley et al., 2011). Finally, some of our analyses lacked adequate power due to small samples in certain patient subgroups. Nonetheless, we believe our findings from a large national integrated health care system provide novel information on the impact of gabapentin on alcohol use in individuals who may or may not have been receiving treatment for substance use.

This work has important implications for researchers and clinicians. We have used real-world data to demonstrate that gabapentin exposure of at least 180 consecutive days at doses <1,500 mg/day was not associated with a decrease in self-reported alcohol consumption among patients receiving gabapentin but not as treatment for their drinking. We did observe a potential threshold effect 1,500 mg/day among patients with diagnosed AUD, which is consistent with the dose response seen in a prior clinical trial (Mason et al., 2014). This

finding suggests the impact on drinking outcomes may not be present at lower doses and may be related to the medication's mechanism of action. Our selection of patients who had 180 consecutive days of gabapentin exposure reflects considerable stability and the impact of gabapentin on alcohol consumption may differ at shorter exposures. In addition to consideration of gabapentin dose and duration, it is important to recognize that since the gabapentin was provided by clinicians outside of substance use treatment programs, our findings were most commonly observed in the absence of counselling. Additional research that pairs information and/or motivational efforts targeted to address alcohol consumption among patients receiving gabapentin in general medical settings may be warranted.

In contrast to the limited use of FDA-approved medications to treat AUD, the widespread prescribing of gabapentin for other conditions indicates that many clinicians are familiar with it, which makes it a potentially useful addition to the array of medications available to treat AUD. However, emerging data indicates that gabapentin can be used non-medically for euphoria among certain patient subgroups (Peckham et al., 2017, Smith et al., 2016). Clinicians prescribing gabapentin will need to use caution to monitor patients for evidence of non-medical use or diversion of gabapentin. In addition, adverse effects known to be associated with gabapentin warrant evaluation in potentially vulnerable patient subgroups including those with HIV and HCV. Our findings indicate that future clinical trials should evaluate the impact of gabapentin on alcohol use in wider patient populations including non-treatment seeking patients with and without AUD.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Sources of support: This work is supported by National Institute on Alcohol Abuse and Alcoholism [R01-AA023733, U24-AA020794, U01-AA020790, U10-AA013566 (completed)]

REFERENCES

- AKGUN KM, GORDON K, PISANI M, FRIED T, MCGINNIS KA, TATE JP, BUTT AA, GIBERT CL, HUANG L, RODRIGUEZ-BARRADAS MC, RIMLAND D, JUSTICE AC & CROTHERS K 2013 Risk factors for hospitalization and medical intensive care unit (MICU) admission among HIV-infected Veterans. *J Acquir Immune Defic Syndr*, 62, 52–9. [PubMed: 23111572]
- AKGUN KM, TATE JP, CROTHERS K, CRYSTAL S, LEAF DA, WOMACK J, BROWN TT, JUSTICE AC & OURSLER KK 2014 An adapted frailty-related phenotype and the VACS index as predictors of hospitalization and mortality in HIV-infected and uninfected individuals. *J Acquir Immune Defic Syndr*, 67, 397–404. [PubMed: 25202921]
- AMERICAN PSYCHIATRIC ASSOCIATION 1994 Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition, Washington, DC, American Psychiatric Press.
- AUSTIN PC 2011 An Introduction to Propensity Score Methods for Reducing the Effects of Confounding in Observational Studies. *Multivariate Behav Res*, 46, 399–424. [PubMed: 21818162]
- BRADLEY KA, LAPHAM GT, HAWKINS EJ, ACHTMAYER CE, WILLIAMS EC, THOMAS RM & KIVLAHAN DR 2011 Quality concerns with routine alcohol screening in VA clinical settings. *J Gen Intern Med*, 26, 299–306. [PubMed: 20859699]

- BRADLEY KA, WILLIAMS EC, ACHTMAYER CE, VOLPP B, COLLINS BJ & KIVLAHAN DR 2006 Implementation of evidence-based alcohol screening in the Veterans Health Administration. *Am J Manag Care*, 12, 597–606. [PubMed: 17026414]
- BROOKHART MA, SCHNEEWEISS S, ROTHMAN KJ, GLYNN RJ, AVORN J & STURMER T 2006 Variable selection for propensity score models. *Am J Epidemiol*, 163, 1149–56. [PubMed: 16624967]
- BUSH K, KIVLAHAN DR, MCDONELL MB, FIHN SD & BRADLEY KA 1998 The AUDIT alcohol consumption questions (AUDIT-C): an effective brief screening test for problem drinking. Ambulatory Care Quality Improvement Project (ACQUIP). Alcohol Use Disorders Identification Test. *Arch Intern Med*, 158, 1789–95. [PubMed: 9738608]
- COHEN E, FEINN R, ARIAS A & KRANZLER HR 2007 Alcohol treatment utilization: findings from the National Epidemiologic Survey on Alcohol and Related Conditions. *Drug Alcohol Depend*, 86, 214–21. [PubMed: 16919401]
- CORMEN TH 2009 Introduction to algorithms, MIT press.
- DICLEMENTE CC, CORNO CM, GRAYDON MM, WIPROVNICK AE & KNOBLACH DJ 2017 Motivational interviewing, enhancement, and brief interventions over the last decade: A review of reviews of efficacy and effectiveness. *Psychol Addict Behav*, 31, 862–887. [PubMed: 29199843]
- DICLEMENTE CC, SCHLUNDT D & GEMMELL L 2004 Readiness and stages of change in addiction treatment. *Am J Addict*, 13, 103–19. [PubMed: 15204662]
- DONALD SG & LANG K 2007 Inference with difference-in-differences and other panel data. *The review of Economics and Statistics*, 89, 221–233.
- EEREN HV, SPREEUWENBERG MD, BARTAK A, DE ROOIJ M & BUSSCHBACH JJ 2015 Estimating subgroup effects using the propensity score method: a practical application in outcomes research. *Med Care*, 53, 366–73. [PubMed: 25738381]
- ESCOTA GV, PATEL P, BROOKS JT, BUSH T, CONLEY L, BAKER J, KOJIC EM, HAMMER J, ONEN NF & INVESTIGATORS SUNS 2015 Short communication: The Veterans Aging Cohort Study Index is an effective tool to assess baseline frailty status in a contemporary cohort of HIV-infected persons. *AIDS Res Hum Retroviruses*, 31, 313–7. [PubMed: 25495766]
- FALK DE, RYAN ML, FERTIG JB, DEVINE EG, CRUZ R, BROWN ES, BURNS H, SALLOUM IM, NEWPORT DJ, MENDELSON J, GALLOWAY G, KAMPMAN K, BROOKS C, GREEN AI, BRUNETTE MF, ROSENTHAL RN, DUNN KE, STRAIN EC, RAY L, SHOPTAW S, AIT-DAOUD TIOURIRINE N, GUNDERSON EW, RANSOM J, SCOTT C, LEGGIO L, CARAS S, MASON BJ, LITTEN RZ, NATIONAL INSTITUTE ON ALCOHOL, A. & ALCOHOLISM CLINICAL INVESTIGATIONS GROUP STUDY, G. 2018 Gabapentin Enacarbil Extended-Release for Alcohol Use Disorder: A Randomized, Double-Blind, Placebo-Controlled, Multisite Trial assessing Efficacy and Safety. *Alcohol Clin Exp Res*.
- FIELD CA, ADINOFF B, HARRIS TR, BALL SA & CARROLL KM 2009 Construct, concurrent and predictive validity of the URICA: data from two multi-site clinical trials. *Drug Alcohol Depend*, 101, 115–23. [PubMed: 19157723]
- FIELLIN DA, REID MC & O'CONNOR PG 2000 Screening for alcohol problems in primary care: a systematic review. *Archives of Internal Medicine*, 160, 1977–89. [PubMed: 10888972]
- FINLAY AK, ELLERBE LS, WONG JJ, TIMKO C, RUBINSKY AD, GUPTA S, BOWE TR, BURDEN JL & HARRIS AHS 2017 Barriers to and facilitators of pharmacotherapy for alcohol use disorder in VA residential treatment programs. *J Subst Abuse Treat*, 77, 38–43. [PubMed: 28476269]
- FORD JH, 2ND, ABRAHAM AJ, LUPULESCU-MANN N, CROFF R, HOFFMAN KA, ALANIS-HIRSCH K, CHALK M, SCHMIDT L & MCCARTY D 2017 Promoting Adoption of Medication for Opioid and Alcohol Use Disorders Through System Change. *J Stud Alcohol Drugs*, 78, 735–744. [PubMed: 28930061]
- FULTZ SL, SKANDERSON M, MOLE L, GANDHI N, BRYANT K, CRYSTAL S & JUSTICE AC 2006 Development and verification of a “virtual” cohort using the national VA Health Information System. *Medical Care*, 44, S25–S30. [PubMed: 16849965]

- GREEN KM & STUART EA 2014 Examining moderation analyses in propensity score methods: application to depression and substance use. *J Consult Clin Psychol*, 82, 773–83. [PubMed: 24731233]
- HARRIS AH, ELLERBE L, REEDER RN, BOWE T, GORDON AJ, HAGEDORN H, OLIVA E, LEMBKE A, KIVLAHAN D & TRAFTON JA 2013 Pharmacotherapy for alcohol dependence: perceived treatment barriers and action strategies among Veterans Health Administration service providers. *Psychol Serv*, 10, 410–9. [PubMed: 23356858]
- HARRIS AH, KIVLAHAN DR, BOWE T & HUMPHREYS KN 2010 Pharmacotherapy of alcohol use disorders in the Veterans Health Administration. *Psychiatr Serv*, 61, 392–8. [PubMed: 20360279]
- HARRIS AH, OLIVA E, BOWE T, HUMPHREYS KN, KIVLAHAN DR & TRAFTON JA 2012 Pharmacotherapy of alcohol use disorders by the Veterans Health Administration: patterns of receipt and persistence. *Psychiatr Serv*, 63, 679–85. [PubMed: 22549276]
- HERNAN MA, HERNANDEZ-DIAZ S, WERLER MM & MITCHELL AA 2002 Causal knowledge as a prerequisite for confounding evaluation: an application to birth defects epidemiology. *Am J Epidemiol*, 155, 176–84. [PubMed: 11790682]
- HOSMER D & LEMESHOW S 2000 *Applied Logistic Regression*, New York, John Wiley and Sons.
- JONAS DE, AMICK HR, FELTNER C, BOBASHEV G, THOMAS K, WINES R, KIM MM, SHANAHAN E, GASS CE, ROWE CJ & GARBUTT JC 2014 Pharmacotherapy for adults with alcohol use disorders in outpatient settings: a systematic review and meta-analysis. *JAMA*, 311, 1889–900. [PubMed: 24825644]
- JUSTICE AC, DOMBROWSKI E, CONIGLIARO J, FULTZ SL, GIBSON D, MADENWALD T, GOULET J, SIMBERKOFF M, BUTT A, RIMLAND D, RODRIGUEZ-BARRADAS M, GIBERT C, OURSLER KK, BROWN S, LEAF DA, GOETZ MB & BRYANT K 2006 Veterans Aging Cohort Study (VACS): Overview and description. *Med Care*, 44, S13–S24. [PubMed: 16849964]
- JUSTICE AC, MCGINNIS KA, TATE JP, BRAITHWAITE RS, BRYANT KJ, COOK RL, EDELMAN EJ, FIELLIN LE, FREIBERG MS, GORDON AJ, KRAEMER KL, MARSHALL BD, WILLIAMS EC & FIELLIN DA 2016 Risk of mortality and physiologic injury evident with lower alcohol exposure among HIV infected compared with uninfected men. *Drug Alcohol Depend*, 161, 95–103. [PubMed: 26861883]
- JUSTICE AC, MODUR S, TATE J, ALTHOFF K, JACOBSON LP, GEDBO KA, KITAHATA M, HORBERG M, BROOKS JT, BUCHACZ K, ROURKE SB, RACHLIS A, NAPRAVNIK S, ERON JJ, WILLIG JH, MOORE RD, KIRK GD, BOSCH RJ, RODRIGUEZ B, HOGG R, THORNE J, GOEDERT JJ, KLEIN MB, GILL J, DEEKS SG, STERLING TR, ANASTOS K & GANGE SJ 2013 Predictive Accuracy of the Veterans Aging Cohort Study Index for Mortality With HIV Infection: A North American Cross Cohort Analysis. *J Acquir Immune Defic Syndr*, 62, 149–163. [PubMed: 23187941]
- KESSELHEIM AS, DARBY D, STUDDERT DM, GLYNN R, LEVIN R & AVORN J 2011 False Claims Act prosecution did not deter off-label drug use in the case of neurontin. *Health Aff (Millwood)*, 30, 2318–27. [PubMed: 22147859]
- KOOB GF & MASON BJ 2016 Existing and Future Drugs for the Treatment of the Dark Side of Addiction. *Annu Rev Pharmacol Toxicol*, 56, 299–322. [PubMed: 26514207]
- KOOB GF & VOLKOW ND 2016 Neurobiology of addiction: a neurocircuitry analysis. *Lancet Psychiatry*, 3, 760–73. [PubMed: 27475769]
- KRANZLER HR & SOYKA M 2018 Diagnosis and Pharmacotherapy of Alcohol Use Disorder: A Review. *JAMA*, 320, 815–824. [PubMed: 30167705]
- LECHNER M 2011 The estimation of causal effects by difference-in-difference methods. *Foundations and Trends® in Econometrics*, 4, 165–224.
- LEE J, KRESINA TF, CAMPOPIANO M, LUBRAN R & CLARK HW 2015 Use of pharmacotherapies in the treatment of alcohol use disorders and opioid dependence in primary care. *Biomed Res Int*, 2015, 137020. [PubMed: 25629034]

- LEUNG JG, HALL-FLAVIN D, NELSON S, SCHMIDT KA & SCHAK KM 2015 The role of gabapentin in the management of alcohol withdrawal and dependence. *Ann Pharmacother*, 49, 897–906. [PubMed: 25969570]
- LIM JK, TATE JP, FULTZ SL, GOULET JL, CONIGLIARO J, BRYANT KJ, GORDON AJ, GIBERT C, RIMLAND D, GOETZ MB, KLEIN MB, FIELLIN DA, JUSTICE AC & LO RE V, 3RD 2014 Relationship between alcohol use categories and noninvasive markers of advanced hepatic fibrosis in HIV-infected, chronic hepatitis C virus-infected, and uninfected patients. *Clin Infect Dis*, 58, 1449–58. [PubMed: 24569533]
- LITTEN RZ, FALK DE, RYAN ML & FERTIG JB 2016a Discovery, Development, and Adoption of Medications to Treat Alcohol Use Disorder: Goals for the Phases of Medications Development. *Alcohol Clin Exp Res*, 40, 1368–79. [PubMed: 27184259]
- LITTEN RZ, WILFORD BB, FALK DE, RYAN ML & FERTIG JB 2016b Potential medications for the treatment of alcohol use disorder: An evaluation of clinical efficacy and safety. *Subst Abuse*, 37, 286–98. [PubMed: 26928397]
- LYON J 2017 More Treatments on Deck for Alcohol Use Disorder. *JAMA*.
- MAGILL M, KILUK BD, MCCRADY BS, TONIGAN JS & LONGABAUGH R 2015 Active Ingredients of Treatment and Client Mechanisms of Change in Behavioral Treatments for Alcohol Use Disorders: Progress 10 Years Later. *Alcohol Clin Exp Res*, 39, 1852–62. [PubMed: 26344200]
- MARK TL, KASSED CA, VANDIVORT-WARREN R, LEVIT KR & KRANZLER HR 2009 Alcohol and opioid dependence medications: prescription trends, overall and by physician specialty. *Drug Alcohol Depend*, 99, 345–9. [PubMed: 18819759]
- MARK TL, KRANZLER HR & SONG X 2003a Understanding US addiction physicians' low rate of naltrexone prescription. *Drug and Alcohol Dependence*, 71, 219–28. [PubMed: 12957340]
- MARK TL, KRANZLER HR, SONG X, BRANSBERGER P, POOLE VH & CROSSE S 2003b Physicians' opinions about medications to treat alcoholism. *Addiction*, 98, 617–26. [PubMed: 12751979]
- MARK TL, LUBRAN R, MCCANCE-KATZ EF, CHALK M & RICHARDSON J 2015 Medicaid coverage of medications to treat alcohol and opioid dependence. *J Subst Abuse Treat*, 55, 1–5. [PubMed: 25921475]
- MARQUINE MJ, UMLAUF A, ROONEY AS, FAZELI PL, GOUAUX BD, PAUL WOODS S, LETENDRE SL, ELLIS RJ, GRANT I, MOORE DJ & GROUP, H. I. V. N. R. P. 2014 The veterans aging cohort study index is associated with concurrent risk for neurocognitive impairment. *J Acquir Immune Defic Syndr*, 65, 190–7. [PubMed: 24442225]
- MASON BJ, QUELLO S, GOODELL V, SHADAN F, KYLE M & BEGOVIC A 2014 Gabapentin treatment for alcohol dependence: a randomized clinical trial. *JAMA Intern Med*, 174, 70–7. [PubMed: 24190578]
- MCGINNIS KA, TATE JP, WILLIAMS EC, SKANDERSON M, BRYANT KJ, GORDON AJ, KRAEMER KL, MAISTO SA, CRYSTAL S, FIELLIN DA & JUSTICE AC 2016 Comparison of AUDIT-C collected via electronic medical record and self-administered research survey in HIV infected and uninfected patients. *Drug Alcohol Depend*, 168, 196–202. [PubMed: 27694059]
- OWENS MD, IOANNOU GN, TSUI JL, EDELMAN EJ, GREENE PA & WILLIAMS EC 2018 Receipt of alcohol-related care among patients with HCV and unhealthy alcohol use. *Drug Alcohol Depend*, 188, 79–85. [PubMed: 29754030]
- PANI PP, TROGU E, PACINI M & MAREMMANI I 2014 Anticonvulsants for alcohol dependence. *Cochrane Database of Systematic Reviews*.
- PECKHAM AM, FAIRMAN KA & SCLAR DA 2017 Policies to mitigate nonmedical use of prescription medications: how should emerging evidence of gabapentin misuse be addressed? *Expert Opin Drug Saf*, 1–5.
- RASSEN JA, GLYNN RJ, ROTHMAN KJ, SETOGUCHI S & SCHNEEWEISS S 2012 Applying propensity scores estimated in a full cohort to adjust for confounding in subgroup analyses. *Pharmacoepidemiol Drug Saf*, 21, 697–709. [PubMed: 22162077]
- RUBINSKY AD, DAWSON DA, WILLIAMS EC, KIVLAHAN DR & BRADLEY KA 2013 AUDIT-C scores as a scaled marker of mean daily drinking, alcohol use disorder severity, and probability

of alcohol dependence in a U.S. general population sample of drinkers. *Alcohol Clin Exp Res*, 37, 1380–90. [PubMed: 23906469]

- SHANTHANNA H, GILRON I, RAJARATHINAM M, ALAMRI R, KAMATH S, THABANE L, DEVEREAUX PJ & BHANDARI M 2017 Benefits and safety of gabapentinoids in chronic low back pain: A systematic review and meta-analysis of randomized controlled trials. *PLoS Med*, 14, e1002369. [PubMed: 28809936]
- SMITH RV, HAVENS JR & WALSH SL 2016 Gabapentin misuse, abuse and diversion: a systematic review. *Addiction*, 111, 1160–74. [PubMed: 27265421]
- SOYKA M & MULLER CA 2017 Pharmacotherapy of alcoholism - an update on approved and off-label medications. *Expert Opin Pharmacother*, 18, 1187–1199. [PubMed: 28658981]
- SPOENDLIN J, LAYTON JB, MUNDKUR M, MEIER C, JICK SS & MEIER CR 2016 The Risk of Achilles or Biceps Tendon Rupture in New Statin Users: A Propensity Score-Matched Sequential Cohort Study. *Drug Saf*, 39, 1229–1237. [PubMed: 27677637]
- SULLIVAN LE, FIELLIN DA & O'CONNOR PG 2005 The prevalence and impact of alcohol problems in major depression: a systematic review. *Am J Med*, 118, 330–41. [PubMed: 15808128]
- SULLIVAN LE, GOULET JL, JUSTICE AC & FIELLIN DA 2011 Alcohol consumption and depressive symptoms over time: a longitudinal study of patients with and without HIV infection. *Drug Alcohol Depend*, 117, 158–63. [PubMed: 21345624]
- TATE JP, JUSTICE AC, HUGHES MD, BONNET F, REISS P, MOCROFT A, NATTERMANN J, LAMPE FC, BUCHER HC, STERLING TR, CRANE HM, KITAHATA MM, MAY M & STERNE JA 2013 An internationally generalizable risk index for mortality after one year of antiretroviral therapy. *AIDS*, 27, 563–72. [PubMed: 23095314]
- WANG SV, JIN Y, FIREMAN B, GRUBER S, HE M, WYSS R, SHIN H, MA Y, KEETON S, KARAMI S, MAJOR JM, SCHNEEWEISS S & GAGNE JJ 2018 Relative Performance of Propensity Score Matching Strategies for Subgroup Analyses. *Am J Epidemiol*.
- WILLIAMS EC, ACHTMEYER CE, YOUNG JP, BERGER D, CURRAN G, BRADLEY KA, RICHARDS J, SIEGEL MB, LUDMAN EJ, LAPHAM GT, FOREHAND M & HARRIS AHS 2018 Barriers to and Facilitators of Alcohol Use Disorder Pharmacotherapy in Primary Care: A Qualitative Study in Five VA Clinics. *J Gen Intern Med*, 33, 258–267. [PubMed: 29086341]
- WILLIAMS EC, ACHTMEYER CE, YOUNG JP, RITTMUELLER SE, LUDMAN EJ, LAPHAM GT, LEE AK, CHAVEZ LJ, BERGER D & BRADLEY KA 2015 Local implementation of alcohol screening and brief intervention at five Veterans Health Administration primary care clinics: Perspectives of clinical and administrative staff. *J Subst Abuse Treat*.
- WILLIAMS EC, GUPTA S, RUBINSKY AD, GLASS JE, JONES-WEBB R, BENSLEY KM & HARRIS AHS 2017 Variation in receipt of pharmacotherapy for alcohol use disorders across racial/ethnic groups: A national study in the U.S. Veterans Health Administration. *Drug Alcohol Depend*, 178, 527–533. [PubMed: 28728114]
- WINSLOW BT, ONYSKO M & HEBER M 2016 Medications for alcohol use disorder. *Am Fam Physician*, 93, 457–465. [PubMed: 26977830]
- WOMACK JA, GOULET JL, GIBERT C, BRANDT CA, SKANDERSON M, GULANSKI B, RIMLAND D, RODRIGUEZ-BARRADAS MC, TATE J, YIN MT, JUSTICE AC & VETERANS AGING COHORT STUDY PROJECT, T. 2013 Physiologic frailty and fragility fracture in HIV-infected male veterans. *Clin Infect Dis*, 56, 1498–504. [PubMed: 23378285]
- YOUNG BA, LIN E, VON KORFF M, SIMON G, CIECHANOWSKI P, LUDMAN EJ, EVERSON-STEWART S, KINDER L, OLIVER M, BOYKO EJ & KATON WJ 2008 Diabetes complications severity index and risk of mortality, hospitalization, and healthcare utilization. *Am J Manag Care*, 14, 15–23. [PubMed: 18197741]

Figure 1. Distribution of propensity scores in gabapentin exposed and unexposed patients in the full cohort before matching, by alcohol use disorder (AUD) history

Figure 2.

Difference-in-difference estimates and 95% confidence intervals of self-reported changes in AUDIT-C scores associated with gabapentin exposure among non-treatment seeking patients and their propensity-score matched controls, by AUD history, baseline AUDIT-C, and prescribed dose of gabapentin

Notes: ** for $p < 0.05$, * for $p < 0.10$; Difference-in-difference = reported AUDIT-C decrease among gabapentin exposed patients minus reported AUDIT-C decrease among propensity-score matched unexposed patients

Abbreviations: AUDIT-C - Alcohol Use Disorders Identification Test – Consumption; AUD – alcohol use disorder; mg - milligrams

Table 1.

Distribution of baseline characteristics in gabapentin exposed and unexposed patients before and after propensity score (PS) matching among those meeting model inclusion criteria; PS were derived separately by AUD history at baseline

Characteristic	Full cohort		PS-matched, AUD		PS-matched, No AUD	
	Exposed n=5,721	Unexposed n=52,243	Exposed n=562	Unexposed n=562*	Exposed n=839	Unexposed n=839*
Age						
20–44	616 (10.8)	10,988 (21.0)	62 (11.0)	59 (10.5)	108 (12.9)	139 (16.5)
45–54	691 (12.1)	6,443 (12.3)	61 (10.9)	55 (9.8)	108 (12.9)	106 (12.6)
55–59	1,166 (20.4)	9,022 (17.3)	104 (18.5)	120 (21.3)	135 (16.1)	123 (14.6)
60+	3,248 (56.8)	25,790 (49.4)	335 (59.6)	328 (58.4)	488 (58.2)	471 (56.2)
Race						
Black	2,524 (44.1)	24,780 (47.4)	293 (52.1)	304 (54.1)	301 (35.9)	333 (39.7)
White	2,612 (45.7)	20,817 (39.9)	216 (38.4)	197 (35.1)	442 (52.7)	414 (49.3)
Hispanic	425 (7.4)	4,295 (8.2)	38 (6.8)	49 (8.7)	63 (7.5)	68 (8.1)
Other	160 (2.8)	2,351 (4.5)	15 (2.7)	12 (2.1)	33 (3.9)	24 (2.9)
Sex						
Male	5,555 (97.1)	50,862 (97.4)	550 (97.9)	551 (98.1)	801 (95.5)	812 (96.8)
Any hospitalization	439 (7.7)	1,410 (2.7)	28 (5.0)	23 (4.0)	26 (3.1)	17 (2.1)
HIV/HCV infection						
Uninfected	3,570 (62.4)	33,032 (63.2)	339 (60.3)	349 (62.1)	550 (65.6)	572 (68.2)
HCV mono-infected	517 (9.0)	2,827 (5.4)	66 (11.7)	67 (11.9)	18 (2.2)	18 (2.1)
HIV mono-infected	1,253 (21.9)	14,002 (26.8)	106 (18.9)	103 (18.3)	249 (29.7)	227 (27.1)
HIV/HCV co-infected	381 (6.7)	2,382 (4.6)	51 (9.1)	43 (7.7)	22 (2.6)	22 (2.6)
Conditions						
Seizure	550 (9.6)	2,539 (4.9)	54 (9.6)	54 (9.6)	43 (5.1)	47 (5.6)
Diabetes	1,798 (31.4)	9,138 (17.5)	141 (25.1)	135 (24.0)	260 (31.0)	234 (27.9)
Neuropathic pain	2,078 (36.3)	4,037 (7.7)	109 (19.4)	91 (16.1)	205 (24.4)	159 (19.0)
Any chronic pain	5,618 (98.2)	46,933 (89.8)	555 (98.8)	554 (98.6)	801 (95.5)	807 (96.2)
Other prescription						
Opioid	2,357 (41.2)	5,120 (9.8)	140 (24.9)	130 (23.2)	193 (23.0)	182 (21.6)

Characteristic	Full cohort		PS-matched, AUD		PS-matched, No AUD	
	Exposed n=5,721	Unexposed n=52,243	Exposed n=562	Unexposed n=562*	Exposed n=839	Unexposed n=839*
<i>Antidepressant</i>	646 (11.3)	1,250 (2.4)	38 (6.8)	22 (4.0)	59 (7.0)	44 (5.3)
<i>NSAID</i>	3,286 (57.4)	11,395 (21.8)	247 (44.0)	260 (46.4)	321 (38.3)	332 (39.5)
<i>Muscle relaxant</i>	1,024 (17.9)	1,726 (3.3)	49 (8.7)	43 (7.6)	72 (8.6)	62 (7.3)
<i>Anticonvulsant</i>	275 (4.8)	1,055 (2.0)	35 (6.2)	27 (4.8)	28 (3.3)	26 (3.1)
Visit to substance use treatment program during follow-up	1,287 (22.5)	11,479 (22.0)	224 (39.9)	214 (38.2)	22 (2.6)	23 (2.7)

Notes: all statistics reported as n (%); up to five unexposed patients were matched to each exposed patient;

* unexposed matches were weighted according to number of matches; among the PS-matched sample, all $p > 0.05$ except for antidepressant prescription among AUD and neuropathic pain among no AUD

Abbreviations: PS - propensity score; HIV - human immunodeficiency virus; HCV - hepatitis C virus; AUD - alcohol use disorder; NSAID - non-steroidal anti-inflammatory drug

Table 2.

Estimated average pre- and post-index date AUDIT-C scores and difference-in-differences (DiD) by baseline alcohol use diagnosis (AUD), AUDIT-C, and average dose of gabapentin during follow-up

		AUD at baseline		No AUD at baseline	
		Exposed n=562	Unexposed n=1,136	Exposed n=839	Unexposed n=1,977
All patients	Pre	4.16 (0.13)	3.94 (0.10)	2.61 (0.07)	2.49 (0.05)
	Post	3.15 (0.13)	3.32 (0.10)	2.02 (0.07)	2.05 (0.05)
	D ^a	-1.01 (0.14)	-0.62 (0.10)	-0.59 (0.07)	-0.44 (0.04)
	DiD (95% CI)	0.39 (0.05, 0.73), p=0.0264		0.14 (-0.01, 0.30), p=0.0691	
By baseline AUDIT-C					
1-3	Pre	n=310 2.18 (0.13)	n=649 2.19 (0.09)	n=674 1.97 (0.06)	n=1,586 1.92 (0.04)
	Post	2.16 (0.13)	2.75 (0.09)	1.65 (0.06)	1.70 (0.04)
	D ^a	-0.03 (0.17)	0.57 (0.11)	-0.31 (0.07)	-0.21 (0.05)
	DiD (95% CI)	0.59 (0.20, 0.99), p=0.0032		0.10 (-0.07, 0.27), p=0.2341	
4-7	Pre	n=189 5.29 (0.16)	n=378 5.18 (0.12)	n=165 5.09 (0.11)	n=391 4.92 (0.08)
	Post	4.12 (0.16)	3.82 (0.12)	3.40 (0.11)	3.55 (0.08)
	D ^a	-1.17 (0.21)	-1.37 (0.15)	-1.69 (0.15)	-1.37 (0.09)
	DiD (95% CI)	-0.19 (-0.70, 0.32), p=0.4636		0.32 (-0.02, 0.66), p=0.0615	
8	Pre	n=63 10.03 (0.27)	n=109 9.89 (0.21)		
	Post	4.69 (0.27)	4.76 (0.21)		
	D ^a	-5.33 (0.37)	-5.12 (0.28)		
	DiD (95% CI)	0.21 (-0.70, 1.11), p=0.6500			
By average dose, mg					
<600	Pre	n=168 3.78 (0.22)	n=1,136 3.95 (0.10)	n=270 2.67 (0.11)	n=1,977 2.49 (0.05)

§

		AUD at baseline		No AUD at baseline	
		Exposed n=562	Unexposed n=1,136	Exposed n=839	Unexposed n=1,977
600-1,499	Post	3.11 (0.22)	3.32 (0.10)	1.87 (0.11)	2.05 (0.05)
	D ^a	-0.67 (0.26)	-0.62 (0.10)	-0.81 (0.12)	-0.44 (0.04)
	DiD (95% CI)	0.05 (-0.50, 0.59), p=0.8650		0.37 (0.12, 0.61), p=0.0034	
		n=267	n=1,136	n=352	n=1,977
1,500	Pre	4.25 (0.18)	3.95 (0.10)	2.59 (0.10)	2.49 (0.05)
	Post	3.21 (0.18)	3.32 (0.10)	2.16 (0.10)	2.05 (0.05)
	D ^a	-1.04 (0.21)	-0.62 (0.10)	-0.43 (0.10)	-0.44 (0.04)
	DiD (95% CI)	0.42 (-0.03, 0.87), p=0.0685		-0.01 (-0.23, 0.21), p=0.9255	
1,500	Pre	n=127	n=1,136	n=217	n=1,977
	Post	4.47 (0.26)	3.95 (0.10)	2.57 (0.12)	2.49 (0.05)
	D ^a	3.08 (0.26)	3.32 (0.10)	2.01 (0.12)	2.05 (0.05)
	DiD (95% CI)	-1.39 (0.30)	-0.62 (0.10)	-0.56 (0.13)	-0.44 (0.04)
		0.77 (0.15, 1.38), p=0.0149		0.12 (-0.15, 0.39), p=0.3873	

Abbreviations: AUDIT-C - Alcohol Use Disorders Identification Test - Consumption; Pre - pre-index AUDIT-C score; Post - post-index AUDIT-C score; D^a - change in AUDIT-C score; DiD - difference-in-difference estimate; CI - confidence interval

Notes: statistics reported as mean (standard error); all models adjusted for age, total medication count, and VACS Index; additional adjusting covariates were antidepressant prescriptions at baseline among patients with AUD and the proportion of neuropathic pain diagnoses among patients without AUD

^aToo few patients with no AUD at baseline and AUDIT-C 8, included those with AUDIT-C 8 with AUDIT-C 4-7