Motor Cortical Plasticity in Schizophrenia: A Meta-analysis of Transcranial Magnetic Stimulation – Electromyography Studies

Urvakhsh Meherwan Mehta^{1,*}, Milind Vijay Thanki¹, Jaya Padmanabhan², Alvaro Pascual-Leone², and Matcheri S Keshavan³

¹Department of Psychiatry, National Institute of Mental Health & Neurosciences (NIMHANS), Bangalore, India

²Division of Cognitive Neurology, Berenson-Allen Center for Noninvasive Brain Stimulation, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, Massachusetts, USA

³Department of Psychiatry, Beth Israel Deaconess Medical Center and Massachusetts Mental Health Center, Harvard Medical School, Boston, Massachusetts, USA

Abstract

Background—Several lines of investigations converge upon aberrant synaptic plasticity as a potential pathophysiological characteristic of schizophrenia. *In vivo* experiments using neuromodulatory perturbation techniques like Transcranial Magnetic and Direct Current Stimulation (TMS & tDCS) have been increasingly used to measure ‘motor cortical plasticity’ in schizophrenia. A systematic quantification of cortical plasticity and its moderators in schizophrenia is however lacking.

Method—The PubMed/MEDLINE database was searched for studies up to December 31st, 2017 that examined case-control experiments comparing neuromodulation following single-session of TMS or tDCS. The primary outcome was the standardized mean difference for differential changes in motor evoked potential (MEP) amplitudes measured with single-pulse TMS (MEP_{sp}) between patients and healthy subjects following TMS or tDCS. After examining heterogeneity, meta-analyses were performed using fixed effects models.

Results—A total of 16 datasets comparing cortical plasticity (MEP_{sp}) between 189 schizophrenia patients and 187 healthy controls were included in the meta-analysis. Patients demonstrated diminished MEP_{sp} with effect sizes (Cohen’s *d*) ranging from 0.66 (LTP-like plasticity) to 0.68 (LTD-like plasticity). Heterosynaptic plasticity studies demonstrated a greater effect size (0.79) compared to homosynaptic plasticity studies (0.62), though not significant ($P=0.43$). Clinical, perturbation protocol- and measurement-related factors, and study quality did not significantly moderate the aberrant plasticity demonstrated in schizophrenia.

Conclusions—Schizophrenia patients demonstrate diminished LTP- and LTD-like motor cortical plasticity, which is not influenced by the various clinical and experimental protocol related

*Corresponding Author: Urvakhsh Meherwan Mehta, Associate Professor of Psychiatry, Wellcome Trust / DBT India Alliance Early Career Fellow, National Institute of Mental Health & Neurosciences (NIMHANS), Bangalore 560029, India, urvakhsh@gmail.com, Phone: +91-80-26995805, +91-9448208167.

confounders. These consistent findings should encourage the use of perturbation-based biomarkers to characterize illness trajectories and treatment response.

Keywords

Cortical plasticity; Psychosis; Transcranial Magnetic Stimulation; Motor cortex; Biological marker

1 Introduction

Cortical plasticity is conceptualized as an intrinsic attribute of the nervous system that enables its ongoing, adaptive modification in response to a multitude of environmental stimuli (Pascual-Leone et al., 2005). This dynamic malleability of the nervous system (structural and functional) in response to neuronal activity triggered by internal or external stimuli is thought to be central to its processing of learning and adaptive behavior (Hart and Hobert, 2018). There have been incremental advances to our understanding about the precise synaptic morphological changes that govern neuronal plasticity and remodeling. These processes maintain the numerous overlapping and interconnected neural circuits that support brain function and are also implicated in brain disease states. For instance, the dynamic interplay between activity-regulated genes (e.g., *arc/arg3.1*) and presynaptic (e.g., neurexin) or postsynaptic (e.g., neuregulin) cell-adhesion molecules in shaping the excitatory and inhibitory circuit-level plasticity is critical in maintaining normal brain functions and is implicated in the genesis of severe neuropsychiatric disorders including schizophrenia (Leslie and Nedivi, 2011; Schizophrenia Working Group of the Psychiatric Genomics Consortium, 2014; Südhof, 2017).

Various lines of human neuroimaging and animal research suggest that the core manifestations of schizophrenia and its neurodevelopmental trajectories can be understood as resulting from aberrant neuronal plasticity or dysplasticity that encompasses hypoplastic cognitive and volitional neural systems, as well as, hyperplastic salience detection and emotion processing systems (Keshavan et al., 2015). In keeping with these hypotheses, the last couple of decades has witnessed a rise of non-invasive, *in vivo* brain stimulation experiments to induce and quantify cortical plasticity responses in patients with schizophrenia (Cohen et al., 1998; Pascual-Leone et al., 1999). For example, Transcranial Magnetic Stimulation (TMS), by means of its ability to induce neuronal action potentials can be employed as both, a neuromodulator – to induce plastic change, and, as a neurophysiological probe – to quantify the degree of plastic change, in combination with electromyography (EMG), electroencephalography (EEG) or functional magnetic resonance imaging (fMRI). In contrast, Transcranial Direct Current Stimulation (tDCS) can alter resting membrane potentials of underlying neurons and hence be used as a pure neuromodulator.

Such techniques potentially enable researchers to quantify the ‘dysplasticity’ in schizophrenia. As a novel investigational approach, they do provide an excellent opportunity to translate these results from experimental laboratories to clinical practice informing us about potential neural markers that identify at-risk states, support diagnoses and predict treatment outcomes or even select personalized treatments. The central concept (figure-1)

supporting these *in vivo* brain stimulation experiments is to use an external cortical perturbation (successive electric or magnetic fields) and measure the magnitude and duration of change in cortical reactivity induced by this perturbation by means of TMS-EEG or TMS-EMG protocols (Freitas et al., 2013). Such perturbation-based remodeling of brain reactivity provides us with unique insights into local and network-level dynamics that can complement traditional resting-state and task-induced brain reactivity measurements. Importantly, such approaches would be translatable and thus enable reliable bridging of animal to human model systems (Diester et al., 2015).

Over the years, several experiments have now attempted to quantify and compare cortical plasticity in schizophrenia and healthy individuals using such perturbation protocols and their variants (Bhandari et al., 2016; A. Hasan et al., 2013; Voineskos et al., 2013). The ability of these protocols to cause *in vivo* transitory cortical reactivity increases or decreases (Cooke and Bliss, 2006) parallels the commonly observed *in vitro* long-term potentiation/depression (LTP/LTD)-like experience-dependent synaptic plasticity accompanying Hebbian learning and memory (Artola et al., 1990; Toni et al., 1999). Despite having diverse mechanistic foundations, these perturbation protocols (TMS or tDCS), appear to have a final common individual-specific LTP/LTD-like effect that can be quantified by TMS-EMG (most common) or TMS-EEG measurements (figure-1) as changes in the amplitudes of motor evoked potentials (MEPs) or TMS-evoked EEG potentials (TEPs), respectively.

Several studies demonstrate impaired LTP-like and LTD-like cortical plasticity in schizophrenia (McClintock et al., 2011), however these perturbation protocols induce changes that tend to be fragile and variable – a function of not just the biological state or experimental protocol of perturbation and measurement, but also of the pre-stimulation neuronal state of the stimulated cortical area (Huang et al., 2017). A systematic quantification of cortical plasticity aberrations in schizophrenia is however lacking. Moreover, the experimental or patient-related factors that contribute to the variability in cortical plasticity can potentially refine our current understanding about the characteristics of cortical plasticity in schizophrenia. This exercise will also be a step forward in standardizing experimental protocols for employment in future perturbation-based biomarker studies.

In this context, we performed a meta-analysis of studies examining cortical plasticity in schizophrenia using perturbation-based strategies, with the aim to (a) quantify the magnitude of cortical plasticity abnormality in schizophrenia, and (b) study the moderating influence of patient-related (e.g., age, gender, duration of illness, medication dose, symptoms) and experiment-related (e.g., perturbation and recording techniques, methodological rigor) factors on cortical plasticity in schizophrenia.

2 Method

2.1 Study selection

Two authors (UMM and MVT) independently searched the PubMed/MEDLINE databases for studies up to December 31st 2017 using the following key phrases “(neuronal plasticity [Medical Subject Heading] OR cortical plasticity [in titles or abstracts] AND schizophrenia

[Medical Subject Heading]]". In addition, manual search of bibliographic cross-referencing from selected studies and reviews was performed to ensure no potential studies were excluded. Studies were included in the meta-analysis if they reported sufficient data (means and standard deviations, graphs, inferential statistics) to compute effect size measures from case-control experiments examining changes in cortical reactivity as measured using investigational Transcranial Magnetic Stimulation (TMS) protocols in patients with schizophrenia and healthy controls, following any perturbation intervention (e.g., TMS, tDCS). There were a total of 401 hits, of which, 13 studies providing 16 datasets were included in the meta-analysis. Data from three studies were not used in the meta-analysis as two of these studies (Strube et al., 2015b, 2015a) had subjects pooled from studies already included and one study did not provide sufficient data that could be used to compute effect size measures (Zhou et al., 2017). Figure-2 provides details of the study selection process.

2.2 Data analysis

Standardized Mean Differences (Cohen's d) were derived from sample size, means and standard deviations. If group mean $\pm$ SD were not reported, they were extracted from graphs using Plot Digitizer Software (Huwaldt and Steinhorst, 2014). If we could not extract these values from graphs, group*time interaction effect F values or t values for each cortical reactivity parameter were used along with the sample sizes of the two groups to derive Cohen's d and its variance using the *MAd* package (Del Re and Hoyt, 2014) in R statistical programming language version 3.4.3 (R Core Team, 2014). This derivation of standardized mean differences is based on existing guidelines (Cooper et al., 2009) and has been used in earlier meta-analyses (Wykes et al., 2011). Among the various motor cortical reactivity measurements, MEP change was the commonest, followed by resting motor threshold (RMT) and cortical silent period (CSP). An omnibus meta-analysis was performed to examine if perturbation methods induced differential changes in MEP amplitudes with single pulse TMS (MEP) between patients and healthy controls irrespective of the expected direction – LTD- or LTP-like plasticity using the *metafor* package in R (Viechtbauer, 2010). A positive effect size indicated greater induction of MEP by the neuromodulatory method in healthy subjects compared to schizophrenia patients. Subsequently, secondary meta-analyses were conducted on studies examining LTD- and LTP-like plasticity protocols to determine if the effects were consistent across different types of cortical plasticity being examined. Consistency of the computed mean weighted effect sizes were tested with the Cochran's Q test of heterogeneity. Based on the tests of heterogeneity, random or fixed effects models were implemented for the meta-analyses. Publication bias was examined by visualization of funnel plots and performing Egger's regression test and rank correlation test for funnel plot asymmetry. Leave-one-out sensitivity analyses were conducted for determining consistency and robustness of results. An alpha probability error significance level of < 0.05 (two-tailed) was used for the random/fixed effects models, test for heterogeneity, publication bias and moderator analyses. Meta-regression analyses were performed using mixed-effects modeling in *metafor* package in R (Viechtbauer, 2010) to independently control for (entered one by one) the effects of differences in perturbation protocol related parameters, clinical and socio-demographic parameters, cortical reactivity measurement related parameters, as well as, the classificatory system used to diagnose schizophrenia (ICD 10 or DSM IV).

2.3 Methodological rigor and reporting quality ratings

Each of the 13 studies that were included for the meta-analysis were rated for methodological rigor and reporting quality. A tailor-made checklist was designed with a scoring system. The checklist was designed with the items that are relevant to various nuances of assessing change in cortical reactivity (as measured using TMS) following a perturbation method (e.g., ensuring proper matching of cases and controls, recording and eliminating potential confounding variables, a-priori outcome definitions, etc. – see table-S2 in supplementary material for details). The checklist provided a maximum score of 21 and a proportion of the maximum score derived for each study / dataset was used in the meta-regression analysis.

3 Results

3.1 Summary of included studies (table-1)

Neuromodulatory perturbation protocols used in these 13 studies included low frequency repetitive TMS (LF-rTMS), continuous theta burst stimulation (cTBS), cathodal and anodal tDCS and paired associative stimulation (PAS). The site of delivery of these perturbation protocols was the left motor cortex hand area (M1) in all studies except Oxley et al., 2014 (left premotor cortex) and Hasan et al., 2013 (bilateral M1). The neurophysiological probe used in these studies to determine and quantify cortical reactivity was always single-pulse TMS-EMG, recorded predominantly from the right hand muscles (first dorsal interosseous or abductor pollicis brevis). The average sample size across the 16 datasets was 16 patients and 15 healthy subjects, who had a mean duration of illness of 5.78 ± 1.15 years, receiving antipsychotic medications amounting to a cumulative average of 396.56 ± 82.92 milligrams of chlorpromazine equivalents per day.

Most studies found a diminished or equivocal (Hasan et al., 2011; Strube et al., 2016) cortical plasticity response (MEP), and two found an opposite pattern of change in cortical reactivity as compared to healthy subjects – MEP facilitation with LF-rTMS (Oxley et al., 2004) and MEP suppression with 300 pulses of cTBS (Hasan et al., 2015). It is noteworthy that, none of the studies included in the meta-analysis demonstrated an exaggerated plasticity response. Five studies reported data on changes in RMT. While the two LF-rTMS protocols reported an increase in RMT in healthy subjects but not in schizophrenia patients (Fitzgerald, 2004; Oxley et al., 2004), the three cathodal tDCS protocols reported no group difference in RMT change (Alkomiet Hasan et al., 2013b; Hasan et al., 2012a, 2012b). CSP was reported to be unchanged in three (Alkomiet Hasan et al., 2013b; Hasan et al., 2012b, 2011) of four cathodal tDCS studies, while one LF-rTMS study reported diminished CSP in healthy subjects but not in schizophrenia patients (Fitzgerald, 2004). Meta-analyses were performed only for the results reported on MEP and not for changes in CSP or RMT because of the small number and heterogeneous nature of study designs. While the focus of most studies was on comparing the magnitude of plastic change induced, several studies also reported differences in time-courses of the plastic change between groups. Both these PAS studies demonstrated a group-difference (diminished MEP facilitation) at later time-points – 20 minutes (Ribolsi et al., 2017) and 30 minutes (Frantseva et al., 2008) following the PAS protocol. Other studies also recorded cortical reactivity across multiple time-points, but they

did not report any significant time-course difference and hence performed further analyses with average post-stimulation MEPs across all time-points [e.g., (Alkomiet Hasan et al., 2013b, 2013a; Hasan et al., 2015)]. The PAS method of Ribolsi et al. used a cortico-cortical pairing of TMS-pulses, as opposed to other PAS studies that used the traditional pairing of median nerve stimulation with motor cortex stimulation. See table-S1 in supplementary material for more details regarding experimental protocols, clinical data and quality measures.

3.2 Meta-analysis

A total of 13 studies providing 16 datasets comparing cortical plasticity measures (MEP) between an overall number of 189 schizophrenia patients (mean age 32.29 years; 68.1% males) and 187 healthy controls (mean age 31.78±3.4 years; 66.36% males) were included in the meta-analysis. One dataset did not measure MEP (Daskalakis et al., 2008), and hence the primary meta-analysis for change in MEP following a perturbation method was run on 15 datasets. There was good consistency in the effect sizes across all the 15 datasets [Q ($df = 14$) = 6.46, $p = 0.953$]. The meta-analysis, performed using the fixed-effects model, revealed significantly diminished motor cortical plasticity responses in the patient group, with an effect size of 0.673 (95% CI, 0.49 to 0.86; $P < .0001$). There were no potential outliers and influential cases as identified by the leave-one-out (estimates ranging from 0.64 to 0.7) and influence functions. There was little suggestion towards a publication bias as observed from the funnel plot and non-significant tests of plot symmetry (Egger's regression $z = 0.34$, $p = 0.73$; Kendall's tau rank correlation $\tau = 0.29$, $p = 0.14$). We then included the study by Daskalakis et al., which measured thumb angular displacement as a measure of plasticity following an exercise protocol, as this also is likely to indicate LTP-like plasticity properties. The plasticity impairment in the patient group persisted with an effect size of 0.68 (95% CI, 0.49 to 0.86; $P < .0001$) by applying a fixed-effects model [Q ($df = 15$) = 6.6720, $P = 0.9661$]. The forest plots depicting these results are included in figures-3 and 4; funnel plots for these analyses are given in the supplementary material.

3.3 Moderator effects

3.3.1 Examining differences based on plasticity categories—We categorized plasticity measurements based on two principles. The first principle depended on whether the perturbation protocol was likely to facilitate excitability (LTP-like; nine datasets) or inhibition (LTD-like; seven datasets) in the motor cortex. The second principle depended on whether plasticity was induced by activating one set of synapses (homosynaptic plasticity; 12 datasets) or simultaneous paired activation of two sets of synapses (heterosynaptic plasticity; four datasets). Secondary meta-analyses for LTD- / LTP-like plasticity, and homosynaptic / heterosynaptic plasticity studies were also performed using the fixed effects model, as tests of heterogeneity were not significant for inconsistent effect sizes. LTD-like plasticity deficits were seen in schizophrenia patients with an effect size of 0.68 (95% CI, 0.41 to 0.95; $P < .0001$), while LTP-like plasticity deficits were present with an effect size of 0.66 (95% CI, 0.41 to 0.92; $P < .0001$). Similarly, homosynaptic plasticity deficits were present with an effect size of 0.62 (95% CI, 0.41 to 0.84; $P < .0001$), and heterosynaptic plasticity deficits were present with an effect size of 0.79 (95% CI, 0.43 to 1.16; $P < .0001$). The type of plasticity abnormality in the patient group [LTD-like vs. LTP-like; $Q(1) = 0.01$,

$P = 0.92$ and homosynaptic vs. heterosynaptic; $Q(1) = 0.61$, $P = 0.43$] did not significantly moderate the overall abnormality. None of the datasets were identified as outliers that could have potentially influenced the results in a significant manner.

3.3.2 Examining effects of the perturbation protocols used—Two studies each examined cortical plasticity by using TMS and cTBS methods, three studies used anodal-tDCS and four studies each used PAS and cathodal-tDCS. Only one study examined use-dependent plasticity by implementing an exercise protocol to change angular displacement of thumb twitches induced by single pulse TMS. We examined the moderating effects of both, general (e.g., exercise, TMS, PAS, tDCS) and specific (e.g., exercise, TMS, PAS, TBS, cathodal-tDCS or anodal-tDCS) type of perturbation protocols on the results and found no significant effect of one particular method [16 datasets, $Q(1) = 2.05$, $P = 0.82$ and 15 datasets, $Q(1) = 0.62$, $P = 0.73$ respectively]. However, on examining the bubble-plot comparing weighted effect sizes against perturbation protocols, there was an indication that the results coming from LF-rTMS and PAS studies were closer to the true effect size (0.68) determined in the meta-analysis (figure-4).

3.3.3 Examining effects of clinical and socio-demographic variables—Mean age of the subjects in each study [16 datasets, $Q(1) = 0.31$, $P = 0.57$], proportion of males recruited [16 datasets, $Q(1) = 0.27$, $P = 0.59$], classificatory system followed to diagnose schizophrenia [16 datasets, $Q(1) = 0.13$, $P = 0.71$], symptom severity (13 datasets, PANSS total scores) [$Q(1) = 0.01$, $P = 0.96$], duration of illness [11 datasets, $Q(1) = 0.01$, $P = 0.93$], chlorpromazine equivalents [16 datasets, $Q(1) = 0.02$, $P = 0.86$] and global functioning of patients [11 datasets, $Q(1) = 0.07$, $P = 0.79$], did not have a moderating effect on the plasticity differences between the two study groups across the datasets.

3.3.4 Examining effects of TMS/tDCS experimental variables—Similarly, results were not significantly moderated by baseline resting motor threshold of patients [10 datasets, $Q(1) = 0.216$, $P = 0.64$] and controls [10 datasets, $Q(1) = 0.20$, $P = 0.65$], baseline amplitude of motor evoked potential of patients [13 datasets, $Q(1) = 0.17$, $P = 0.67$], stimulation intensity used to measure motor evoked potentials [14 datasets, $Q(1) = 0.01$, $P = 0.92$], and the number of trials [15 datasets, $Q(1) = 0.001$, $P = 0.97$] averaged to determine cortical excitability (motor evoked potentials). Stimulator device type did not moderate the results for cortical plasticity differences as measured using TMS [eight datasets, $Q(1) = 0.49$, $P = 0.48$] and tDCS [seven datasets, $Q(1) = 0.88$, $P = 0.34$] neuromodulatory perturbation protocols. Quality of the individual studies rated on a continuum also did not significantly alter the outcome [16 datasets, $Q(1) = 0.41$, $P = 0.52$].

4 Discussion

We found that schizophrenia patients had impairments in cortical plasticity outcomes measured using TMS-EMG experiments with effect sizes ranging from 0.66 (LTP-like plasticity) to 0.68 (LTD-like plasticity). Heterosynaptic plasticity studies demonstrated a greater effect size (0.79) compared to homosynaptic plasticity studies (0.62), though not statistically significant ($P = 0.43$). Clinical, perturbation protocol-related and cortical reactivity measurement-related factors did not significantly moderate the aberrant plasticity

demonstrated in schizophrenia patients. Quality of individual studies also did not have a moderating effect on plasticity outcomes between the two groups studied. None of the indices indicated potential publication bias. These findings suggest an overall moderate effect size of cortical plasticity impairment in schizophrenia. These findings can be interpreted within three contexts - clinical, study methodological and neurobiological.

Among clinical scenarios, earlier studies in patients with schizophrenia have reported that the illness course and medication status determined plasticity outcomes. One study that examined recent-onset and multiple-episode chronic schizophrenia patients reported diminished cortical plasticity only in the chronic patient group (Hasan et al., 2011). This could either suggest a neurotoxic effect of chronic schizophrenia or an effect of antipsychotic medications – dopamine receptor blockade with sulpiride diminished motor cortical plasticity responses in healthy subjects (Nitsche et al., 2006). However, in our pooled sample analysis, duration of illness did not moderate the cortical plasticity outcomes significantly. Nearly all the studies included in the meta-analysis recruited stable patients on antipsychotic medications with a mean duration of illness ~5.78 years (4.4 to 7.6 years range). There were two studies (Daskalakis et al., 2008; Fitzgerald, 2004) that recruited un-medicated patients; neither of them reported significant differences in cortical plasticity responses between medicated and un-medicated patients. Nevertheless, the sample sizes of un-medicated or recent-onset patients in these studies were 10. We believe this would have made the pooled sample underpowered to detect any moderating effect of these potential confounding variables. In addition, we did not observe any moderating effect of cumulative antipsychotic dose (in chlorpromazine equivalents) on the cortical plasticity group-differences. However, there is also a possibility that altered plasticity may in fact antedate – and perhaps be causally related to – the manifestation of clinical symptoms of schizophrenia. Aberrant LTD-like motor cortical plasticity was indeed reported in a cathodal-tDCS study in non-psychotic first-degree relatives of individuals with schizophrenia (Alkomiet Hasan et al., 2013b). Longitudinal studies will be better able to examine the translational potential of such neuromodulatory perturbation protocols in identifying individuals who are at risk for psychotic disorders.

Another clinical source of variation could be the smoking status of patients. A pooled analysis of earlier studies demonstrated impaired LTD-like cortical plasticity only in non-smoking schizophrenia patients, but normal LTD-like cortical plasticity responses in schizophrenia patients who were chronic smokers (Strube et al., 2015a). Interestingly, varenicline, a partial agonist at the $\alpha 4\beta 2$ nicotinic acetylcholine receptor, increased LTP-like cortical plasticity in schizophrenia patients who were non-smokers. These findings highlight not only the cholinergic regulation of N-methyl-D-aspartate (NMDA)-mediated glutamatergic neurotransmission (and thereby plasticity) in schizophrenia (Marchi et al., 2015), but also emphasize the need for meticulous patient selection during recruitment in such studies. This is crucial, especially given the large response variability observed across diverse neuromodulatory perturbation protocols (López-Alonso et al., 2015; van der Kamp et al., 1996). Unfortunately, smoking status was not routinely reported in most studies, and hence we could not perform a meta-regression to examine its moderating effects on cortical plasticity outcomes in this analysis. Yet another potential clinical confounding effect could be mediated by diabetes, which often presents with subtle neurocognitive deficits and is

known to be highly comorbid in individuals with schizophrenia (Annamalai et al., 2017). Early reports suggest impaired LTP-like cortical plasticity effects assessed using intermittent theta burst stimulation in patients with type-2 diabetes mellitus (Fried et al., 2016). None of the studies included in this analysis provided information on comorbid diabetes and hence we could not examine its effect on cortical plasticity outcomes. Lastly, brain atrophy, by means of a resulting greater scalp-to-brain distance and added current shunting in the cerebrospinal fluid spaces, could also reduce perturbation effects (Wagner et al., 2007). Since patients with schizophrenia also have varying degrees of brain atrophy (Padmanabhan et al., 2015; Walton et al., 2018), MRI brain imaging to correct for scalp-to-brain distance coupled with projective studies where intensity is accordingly adjusted would be desirable.

From a study methodological standpoint, we examined different perturbation-protocols and the cortical reactivity measurement using TMS-EMG set-up. In summary, none of the above moderated the between-group cortical plasticity outcome in a significant manner. This is despite the various perturbation methods having potentially different, yet inconclusive cellular and network-level mechanisms that have a final common output that resembles LTP/LTD-like experience dependent synaptic plasticity (Huang et al., 2017). However, bubble-plot observations suggested LF-rTMS and PAS perturbation protocols to have less fluctuating effect sizes that were closer to the true, overall effect-size (figure-5) as compared to other methods. Newer perturbation methods like theta burst stimulation (Huang et al., 2005), quadripulse stimulation (Ugawa, 2016), quadripulse theta burst stimulation (Jung et al., 2016) and high-definition tDCS (Kuo et al., 2013) that are likely to induce more robust and less variable cortical reactivity changes have not been sufficiently studied in the schizophrenia population. Given the rapid advancements in engineering newer perturbation protocols, it is important that the newer methods are applied in a clinical population after careful experiments that demonstrate their robustness in terms of validity with respect to putative neurobiological underpinnings, and test-retest reliability. We now know from healthy subject experiments that there is a minimum number of trials to record MEPs that is needed to achieve sufficient test-retest reliability of cortical inhibition and plasticity outcomes with neuromodulatory protocols (Chang et al., 2016). However, we did not find a significant moderation of differential cortical plasticity outcomes by the number of trials used in the included studies to measure MEPs (see supplementary material). The absence of sham stimulation protocols in any of these experiments should also merit a deliberation on possible placebo effects the perturbation protocols could have had on motor cortical reactivity changes. For example, it is known that a sense of expectation and anticipation result in motor cortical excitability changes (Lou et al., 2013). Such expectancy related placebo effects on motor cortical physiology, though not elaborately characterized, could be avoided by either having sham stimulations to account for the sensory effects or alternative control strategies like stimulating a non-motor brain region (Duecker and Sack, 2015). There were very few studies that examined time-course effects of the various neuromodulatory perturbation protocols on MEP between groups. Similarly, there was limited data about effects on other cortical reactivity measures (e.g., RMT and CSP) thus limiting conclusive inferences. These could be the focus of future studies.

Lastly, the overarching diminished cortical plasticity (LTP/LTD-like or homo/heterosynaptic) responses to a range of non-invasive brain stimulation perturbation protocols

in schizophrenia partly validates the ‘dysplastic’ brain model of schizophrenia (Keshavan et al., 2015), especially with respect to the hypo-plasticity observed in most studies. There was insufficient data to systematically examine the association between hypoplastic motor cortical changes and symptom dimensions in the patient group. This is understandable since most studies had recruited stable schizophrenia patients. Examining motor cortical plasticity in actively symptomatic or resistant patients with schizophrenia and patients with homogenous sets of symptoms will enable testing of the aberrant hyper-plasticity that is hypothesized as to be related to some symptom dimensions like affective instability and phasic exacerbations of positive symptomatology (Keshavan et al., 2015). Nevertheless, a diminished plastic brain response to perturbation protocols in schizophrenia lays emphasis on the role of an aberrant, hypofunctioning NMDA receptor system and its downstream effects on maintaining dendritic arborization, glial density, myelination and an optimal excitation/inhibition balance (Friston et al., 2016; Stephan et al., 2009) necessary for optimal network-level brain functioning. Incidentally, the activity-dependent trafficking of the NMDA receptors between synaptic and extra-synaptic sites is also central to optimal synaptic plasticity and experience-dependent learning (Lau and Zukin, 2007). While there is consistent evidence of a diminished motor cortical plasticity response in schizophrenia, very little is known about its specificity to schizophrenia. Indeed, cortical plasticity has been systematically evaluated and found to be abnormal in individuals with depression (Kuhn et al., 2016), autism (Oberman et al., 2016; Pedapati et al., 2016), Alzheimer’s dementia (Koch et al., 2017). However, there are no head-to-head comparisons of how cortical plasticity varies across these diverse psychiatric disorders.

Restricting our interpretation on cortical plasticity in schizophrenia to data emerging from TMS-EMG studies focusing on the motor cortex alone might reveal only a partial understanding of its complex pathophysiology. Hence, the importance of TMS-EEG/MRI studies that could provide information with better temporo-spatial and molecular resolution cannot be underestimated. Indeed, evidence from parallel investigations comparing electroencephalographic event related potentials evoked by tetanized auditory stimuli (Mears and Spencer, 2012) and visual high frequency stimulation (Cavus et al., 2012) suggest potentiation deficits in schizophrenia; these deficits also have an association with their neurocognitive performance (Jahshan et al., 2017) and functional outcomes (Kantrowitz et al., 2016). To the best of our knowledge, there are no studies that have combined the assessment of motor and sensory cortex plasticity in schizophrenia. This is an important research gap, which can be addressed by combining motor and non-motor plasticity experiments, especially because the developmental critical periods for cortical plasticity are likely to differ across specialized cortical processes supported by distinct cortical neural networks. As a next step, it needs to be evaluated whether cortical plasticity aberrations observed in the motor cortex parallel those in other cortical regions. The emergence of TMS-EEG cortical plasticity methods (Chung et al., 2017) has the potential to bridge this gap in the near future by enabling readouts from motor, as well as, non-motor cortical areas.

It is possible that each of the perturbation protocols used to elicit a neuroplastic response acts via different neurophysiological and neurochemical mechanisms that are relevant to the pathophysiology of schizophrenia (Wilson et al., 2018). The cortical-level in vivo plasticity elicited using such methods need not always directly correlate with LTP- and LTD-like

synaptic-level plasticity, as the cortical-level plasticity is potentially influenced by a range of complex mechanisms such as sliding thresholds, metaplasticity and an optimal cortical excitatory–inhibitory balance (Hensch, 2005). This provides us with the unique opportunity to examine (a) the behavioral correlates of diverse cortical plasticity properties in schizophrenia and (b) their longitudinal trajectories over distinct illness phases that parallel brain development and potentially represent a brain health index (Freitas et al., 2013). Meticulous homogenous subject selection, standardized neuromodulatory protocol delivery, recording and measuring potential confounding variables (e.g., smoking status, comorbid diabetes, genotyping of important plasticity-related polymorphisms) will go a long way in reducing the inter-individual response variability. Implementing these refined approaches can potentially improve the reliability of cortical plasticity responses, and the true inter-subject variability can then be utilized as potential markers of response to not only non-invasive brain stimulation therapies (Osoegawa et al., 2018), but also to antipsychotic medications and cognitive training interventions in schizophrenia.

In conclusion, this meta-analysis of TMS-EMG studies demonstrated hypoactive mechanisms of motor cortical plasticity in patients with schizophrenia as compared to healthy subjects. These deficits were reported in largely stable schizophrenia groups and were not moderated by clinical factors, methodological variability (perturbation-method or cortical reactivity recordings) or study quality. Future studies need to focus on examining the specificity, behavioral correlates and developmental trajectories of these cortical plasticity responses in schizophrenia. Potential clues towards heterogeneous responses (e.g., smoking status, illness stage and medication status, comorbid diabetes and specific genotypes) need to be systematically measured and accounted for. The robust findings in terms of moderate effect size deficits in cortical plasticity in schizophrenia should encourage future studies to examine this novel investigational protocol to supplement conventional neuroimaging and electrophysiological experiments as disease biomarkers.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

References

- Annamalai A, Kosir U, Tek C. Prevalence of obesity and diabetes in patients with schizophrenia. *World J Diabetes*. 2017; 8:390–396. DOI: 10.4239/wjd.v8.i8.390 [PubMed: 28861176]
- Artola A, Bröcher S, Singer W. Different voltage-dependent thresholds for inducing long-term depression and long-term potentiation in slices of rat visual cortex. *Nature*. 1990; 347:69–72. DOI: 10.1038/347069a0 [PubMed: 1975639]
- Bhandari A, Voineskos D, Daskalakis ZJ, Rajji TK, Blumberger DM. A Review of Impaired Neuroplasticity in Schizophrenia Investigated with Non-invasive Brain Stimulation. *Front Psychiatry*. 2016; 7doi: 10.3389/fpsy.2016.00045
- Cavus I, Reinhart RM, Roach BJ, Gueorgieva R, Teyler TJ, Clapp WC, Ford JM, Krystal JH, Mathalon DH. Impaired visual cortical plasticity in schizophrenia. *Biol Psychiatry*. 2012; 71:512–20. DOI: 10.1016/j.biopsych.2012.01.013 [PubMed: 22364738]
- Chang WH, Fried PJ, Saxena S, Jannati A, Gomes-Osman J, Kim Y-H, Pascual-Leone A. Optimal number of pulses as outcome measures of neuronavigated transcranial magnetic stimulation. *Clin Neurophysiol Off J Int Fed Clin Neurophysiol*. 2016; 127:2892–2897. DOI: 10.1016/j.clinph.2016.04.001

- Chung SW, Lewis BP, Rogasch NC, Saeki T, Thomson RH, Hoy KE, Bailey NW, Fitzgerald PB. Demonstration of short-term plasticity in the dorsolateral prefrontal cortex with theta burst stimulation: A TMS-EEG study. *Clin Neurophysiol Off J Int Fed Clin Neurophysiol*. 2017; 128:1117–1126. DOI: 10.1016/j.clinph.2017.04.005
- Cohen LG, Ziemann U, Chen R, Classen J, Hallett M, Gerloff C, Butefisch C. Studies of neuroplasticity with transcranial magnetic stimulation. *J Clin Neurophysiol Off Publ Am Electroencephalogr. Soc.* 1998; 15:305–324.
- Cooke SF, Bliss TV. Plasticity in the human central nervous system. *Brain*. 2006; 129:1659–1673. DOI: 10.1093/brain/awl082 [PubMed: 16672292]
- Cooper, HM, Hedges, LV, Valentine, JC, editors. *The handbook of research synthesis and meta-analysis*. 2nd ed. Russell Sage Foundation; New York: 2009.
- Daskalakis ZJ, Christensen BK, Fitzgerald PB, Chen R. Dysfunctional neural plasticity in patients with schizophrenia. *Arch Gen Psychiatry*. 2008; 65:378–85. DOI: 10.1001/archpsyc.65.4.378 [PubMed: 18391126]
- Del Re, AC; Hoyt, WT. MAD: Meta-Analysis with Mean Differences. R package version 0.8-2. 2014.
- Diester, I, Hefti, F, Mansuy, I, Pascual-Leone, A, Robbins, TW, Rubin, LL, Sawa, A, Wernig, M, Dölen, G, Mucke, L, Sommer, B. Bridging the Gap between Patients and Models Translational Neuroscience: Toward New Therapies. Hyman, SE, Nikolic, K, editors. MIT Press; Cambridge, MA: 2015. 209–244.
- Duecker F, Sack AT. Rethinking the role of sham TMS. *Front Psychol*. 2015; 6doi: 10.3389/fpsyg.2015.00210
- Fitzgerald P. Reduced plastic brain responses in schizophrenia: a transcranial magnetic stimulation study*1. *Schizophr Res*. 2004; 71:17–26. DOI: 10.1016/j.schres.2004.01.018 [PubMed: 15374568]
- Frantseva MV, Fitzgerald PB, Chen R, Moller B, Daigle M, Daskalakis ZJ. Evidence for impaired long-term potentiation in schizophrenia and its relationship to motor skill learning. *Cereb Cortex*. 2008; 18:990–6. DOI: 10.1093/cercor/bhm151 [PubMed: 17855721]
- Freitas C, Farzan F, Pascual-Leone A. Assessing brain plasticity across the lifespan with transcranial magnetic stimulation: why, how, and what is the ultimate goal? *Front Neurosci*. 2013; 7doi: 10.3389/fnins.2013.00042
- Fried PJ, Schilberg L, Brem A-K, Saxena S, Wong B, Cypess AM, Horton ES, Pascual-Leone A. Humans with Type-2 Diabetes Show Abnormal Long-Term Potentiation-Like Cortical Plasticity Associated with Verbal Learning Deficits. *J Alzheimers Dis*. 2016; 55:89–100. DOI: 10.3233/JAD-160505
- Friston K, Brown HR, Siemerkus J, Stephan KE. The dysconnection hypothesis (2016). *Schizophr Res*. 2016; 176:83–94. DOI: 10.1016/j.schres.2016.07.014 [PubMed: 27450778]
- Hart MP, Hobert O. Neurexin controls plasticity of a mature, sexually dimorphic neuron. *Nature*. 2018; 553:165–170. DOI: 10.1038/nature25192 [PubMed: 29323291]
- Hasan A, Aborowa R, Nitsche MA, Marshall L, Schmitt A, Gruber O, Falkai P, Wobrock T. Abnormal bihemispheric responses in schizophrenia patients following cathodal transcranial direct stimulation. *Eur Arch Psychiatry Clin Neurosci*. 2012a; 262:415–23. DOI: 10.1007/s00406-012-0298-7 [PubMed: 22318337]
- Hasan, Alkomiet; Bergener, T; Nitsche, MA; Strube, W; Bunse, T; Falkai, P; Wobrock, T. Impairments of Motor-Cortex Responses to Unilateral and Bilateral Direct Current Stimulation in Schizophrenia. *Front Psychiatry*. 2013a; 4doi: 10.3389/fpsyg.2013.00121
- Hasan A, Brinkmann C, Strube W, Palm U, Malchow B, Rothwell JC, Falkai P, Wobrock T. Investigations of motor-cortex cortical plasticity following facilitatory and inhibitory transcranial theta-burst stimulation in schizophrenia: A proof-of-concept study. *J Psychiatr Res*. 2015; 61:196–204. DOI: 10.1016/j.jpsychires.2014.12.006 [PubMed: 25555304]
- Hasan A, Falkai P, Wobrock T. Transcranial brain stimulation in schizophrenia: targeting cortical excitability, connectivity and plasticity. *Curr Med Chem*. 2013; 20:405–413. [PubMed: 23157633]
- Hasan, Alkomiet; Misewitsch, K; Nitsche, MA; Gruber, O; Padberg, F; Falkai, P; Wobrock, T. Impaired Motor Cortex Responses in Non-Psychotic First-Degree Relatives of Schizophrenia

- Patients: A Cathodal tDCS Pilot Study. *Brain Stimulat.* 2013b; 6:821–829. DOI: 10.1016/j.brs.2013.03.001
- Hasan A, Nitsche MA, Herrmann M, Schneider-Axmann T, Marshall L, Gruber O, Falkai P, Wobrock T. Impaired long-term depression in schizophrenia: a cathodal tDCS pilot study. *Brain Stimul.* 2012b; 5:475–83. DOI: 10.1016/j.brs.2011.08.004 [PubMed: 21945231]
- Hasan A, Nitsche MA, Rein B, Schneider-Axmann T, Guse B, Gruber O, Falkai P, Wobrock T. Dysfunctional long-term potentiation-like plasticity in schizophrenia revealed by transcranial direct current stimulation. *Behav Brain Res.* 2011; 224:15–22. DOI: 10.1016/j.bbr.2011.05.017 [PubMed: 21645555]
- Hensch TK. Critical period plasticity in local cortical circuits. *Nat Rev Neurosci.* 2005; 6:877–888. DOI: 10.1038/nrn1787 [PubMed: 16261181]
- Huang Y-Z, Edwards MJ, Rounis E, Bhatia KP, Rothwell JC. Theta burst stimulation of the human motor cortex. *Neuron.* 2005; 45:201–206. DOI: 10.1016/j.neuron.2004.12.033 [PubMed: 15664172]
- Huang Y-Z, Lu M-K, Antal A, Classen J, Nitsche M, Ziemann U, Ridding M, Hamada M, Ugawa Y, Jaberzadeh S, Suppa A, et al. Plasticity induced by non-invasive transcranial brain stimulation: A position paper. *Clin Neurophysiol.* 2017; 128:2318–2329. DOI: 10.1016/j.clinph.2017.09.007 [PubMed: 29040922]
- Huwaldt, JA; Steinhurst, S. Plot Digitizer 2.6.6. 2014. [WWW Document]. URL <http://plotdigitizer.sourceforge.net>
- Jahshan C, Wynn JK, Mathalon DH, Green MF. Cognitive correlates of visual neural plasticity in schizophrenia. *Schizophr Res.* 2017; 190:39–45. DOI: 10.1016/j.schres.2017.03.016 [PubMed: 28336195]
- Jung NH, Gleich B, Gattinger N, Hoess C, Haug C, Siebner HR, Mall V. Quadri-Pulse Theta Burst Stimulation using Ultra-High Frequency Bursts – A New Protocol to Induce Changes in Cortico-Spinal Excitability in Human Motor Cortex. *PLOS ONE.* 2016; 11:e0168410.doi: 10.1371/journal.pone.0168410 [PubMed: 27977758]
- Kantrowitz JT, Epstein ML, Beggel O, Rohrig S, Lehrfeld JM, Revheim N, Lehrfeld NP, Reep J, Parker E, Silipo G, Ahissar M, et al. Neurophysiological mechanisms of cortical plasticity impairments in schizophrenia and modulation by the NMDA receptor agonist D-serine. *Brain J Neurol.* 2016; 139:3281–3295. DOI: 10.1093/brain/aww262
- Keshavan MS, Mehta UM, Padmanabhan JL, Shah JL. Dysplasticity, metaplasticity, and schizophrenia: Implications for risk, illness, and novel interventions. *Dev Psychopathol.* 2015; 27:615–35. DOI: 10.1017/S095457941500019X [PubMed: 25997775]
- Koch G, Di Lorenzo F, Loizzo S, Motta C, Travaglione S, Baiula M, Rimondini R, Ponzo V, Bonni S, Toniolo S, Sallustio F, et al. CSF tau is associated with impaired cortical plasticity, cognitive decline and astrocyte survival only in APOE4-positive Alzheimer's disease. *Sci Rep.* 2017; 7doi: 10.1038/s41598-017-14204-3
- Kuhn M, Mainberger F, Feige B, Maier JG, Mall V, Jung NH, Reis J, Klöppel S, Normann C, Nissen C. State-Dependent Partial Occlusion of Cortical LTP-Like Plasticity in Major Depression. *Neuropsychopharmacology.* 2016; 41:1521–1529. DOI: 10.1038/npp.2015.310 [PubMed: 26442602]
- Kuo H-I, Bikson M, Datta A, Minhas P, Paulus W, Kuo M-F, Nitsche MA. Comparing Cortical Plasticity Induced by Conventional and High-Definition 4×1 Ring tDCS: A Neurophysiological Study. *Brain Stimulat.* 2013; 6:644–648. DOI: 10.1016/j.brs.2012.09.010
- Lau CG, Zukin RS. NMDA receptor trafficking in synaptic plasticity and neuropsychiatric disorders. *Nat Rev Neurosci.* 2007; 8:413–426. DOI: 10.1038/nrn2153 [PubMed: 17514195]
- Leslie JH, Nedivi E. Activity-regulated genes as mediators of neural circuit plasticity. *Prog Neurobiol.* 2011; 94:223–237. DOI: 10.1016/j.pneurobio.2011.05.002 [PubMed: 21601615]
- López-Alonso V, Fernández-Del-Olmo M, Costantini A, Gonzalez-Henriquez JJ, Cheeran B. Intra-individual variability in the response to anodal transcranial direct current stimulation. *Clin Neurophysiol Off J Int Fed Clin Neurophysiol.* 2015; 126:2342–2347. DOI: 10.1016/j.clinph.2015.03.022

- Lou J-S, Dimitrova DM, Hammerschlag R, Nutt J, Hunt EA, Eaton RW, Johnson SC, Davis MD, Arnold GC, Andrea SB, Oken BS. Effect of expectancy and personality on cortical excitability in Parkinson's disease. *Mov Disord Off J Mov Disord Soc.* 2013; 28:1257–1262. DOI: 10.1002/mds.25471
- Marchi M, Grilli M, Pittaluga AM. Nicotinic modulation of glutamate receptor function at nerve terminal level: a fine-tuning of synaptic signals. *Front Pharmacol.* 2015; 6doi: 10.3389/fphar.2015.00089
- McClintock SM, Freitas C, Oberman L, Lisanby SH, Pascual-Leone A. Transcranial Magnetic Stimulation: A Neuroscientific Probe of Cortical Function in Schizophrenia. *Biol Psychiatry.* 2011; 70:19–27. DOI: 10.1016/j.biopsych.2011.02.031 [PubMed: 21571254]
- Mears RP, Spencer KM. Electrophysiological assessment of auditory stimulus-specific plasticity in schizophrenia. *Biol Psychiatry.* 2012; 71:503–11. DOI: 10.1016/j.biopsych.2011.12.016 [PubMed: 22277333]
- Nitsche MA, Lampe C, Antal A, Liebetanz D, Lang N, Tergau F, Paulus W. Dopaminergic modulation of long-lasting direct current-induced cortical excitability changes in the human motor cortex: Dopamine in human neuroplasticity. *Eur J Neurosci.* 2006; 23:1651–1657. DOI: 10.1111/j.1460-9568.2006.04676.x [PubMed: 16553629]
- Oberman LM, Ifert-Miller F, Najib U, Bashir S, Heydrich JG, Picker J, Rotenberg A, Pascual-Leone A. Abnormal Mechanisms of Plasticity and Metaplasticity in Autism Spectrum Disorders and Fragile X Syndrome. *J Child Adolesc Psychopharmacol.* 2016; 26:617–624. DOI: 10.1089/cap.2015.0166 [PubMed: 27218148]
- Osoegawa C, Gomes JS, Grigolon RB, Brietzke E, Gadelha A, Lacerda ALT, Dias ÁM, Cordeiro Q, Laranjeira R, de Jesus D, Daskalakis ZJ, et al. Non-invasive brain stimulation for negative symptoms in schizophrenia: An updated systematic review and meta-analysis. *Schizophr Res.* 2018; doi: 10.1016/j.schres.2018.01.010
- Oxley T, Fitzgerald PB, Brown TL, de Castella A, Daskalakis ZJ, Kulkarni J. Repetitive transcranial magnetic stimulation reveals abnormal plastic response to premotor cortex stimulation in schizophrenia. *Biol Psychiatry.* 2004; 56:628–33. DOI: 10.1016/j.biopsych.2004.08.023 [PubMed: 15522245]
- Padmanabhan JL, Tandon N, Haller CS, Mathew IT, Eack SM, Clementz BA, Pearlson GD, Sweeney JA, Tamminga CA, Keshavan MS. Correlations Between Brain Structure and Symptom Dimensions of Psychosis in Schizophrenia, Schizoaffective, and Psychotic Bipolar I Disorders. *Schizophr Bull.* 2015; 41:154–162. DOI: 10.1093/schbul/sbu075 [PubMed: 24907239]
- Pascual-Leone A, Amedi A, Fregni F, Merabet LB. The plastic human brain cortex. *Annu Rev Neurosci.* 2005; 28:377–401. DOI: 10.1146/annurev.neuro.27.070203.144216 [PubMed: 16022601]
- Pascual-Leone A, Tarazona F, Keenan J, Tormos JM, Hamilton R, Catala MD. Transcranial magnetic stimulation and neuroplasticity. *Neuropsychologia.* 1999; 37:207–217. [PubMed: 10080378]
- Pedapati EV, Gilbert DL, Erickson CA, Horn PS, Shaffer RC, Wink LK, Laue CS, Wu SW. Abnormal Cortical Plasticity in Youth with Autism Spectrum Disorder: A Transcranial Magnetic Stimulation Case-Control Pilot Study. *J Child Adolesc Psychopharmacol.* 2016; 26:625–631. DOI: 10.1089/cap.2015.0183 [PubMed: 27007257]
- R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing; Vienna, Austria: 2014.
- Ribolsi M, Lisi G, Ponzo V, Siracusano A, Caltagirone C, Nioiu C, Koch G. Left hemispheric breakdown of LTP-like cortico-cortical plasticity in schizophrenic patients. *Clin Neurophysiol.* 2017; 128:2037–2042. DOI: 10.1016/j.clinph.2017.06.255 [PubMed: 28843131]
- Schizophrenia Working Group of the Psychiatric Genomics Consortium. Biological insights from 108 schizophrenia-associated genetic loci. *Nature.* 2014; 511:421–427. DOI: 10.1038/nature13595 [PubMed: 25056061]
- Stephan KE, Friston KJ, Frith CD. Dysconnection in schizophrenia: from abnormal synaptic plasticity to failures of self-monitoring. *Schizophr Bull.* 2009; 35:509–27. DOI: 10.1093/schbul/sbn176 [PubMed: 19155345]

- Strube W, Bunse T, Nitsche MA, Palm U, Falkai P, Hasan A. Differential response to anodal tDCS and PAS is indicative of impaired focal LTP-like plasticity in schizophrenia. *Behav Brain Res.* 2016; 311:46–53. DOI: 10.1016/j.bbr.2016.05.027 [PubMed: 27185738]
- Strube W, Bunse T, Nitsche MA, Wobrock T, Aborowa R, Misewitsch K, Herrmann M, Falkai P, Hasan A. Smoking restores impaired LTD-like plasticity in schizophrenia: a transcranial direct current stimulation study. *Neuropsychopharmacology.* 2015a; 40:822–830. [PubMed: 25308351]
- Strube W, Nitsche MA, Wobrock T, Bunse T, Rein B, Herrmann M, Schmitt A, Nieratschker V, Witt SH, Rietschel M, Falkai P, et al. BDNF-Val66Met-Polymorphism Impact on Cortical Plasticity in Schizophrenia Patients: A Proof-of-Concept Study. *Int J Neuropsychopharmacol.* 2015b; 18doi: 10.1093/ijnp/pyu040
- Südhof TC. Synaptic Neurexin Complexes: A Molecular Code for the Logic of Neural Circuits. *Cell.* 2017; 171:745–769. DOI: 10.1016/j.cell.2017.10.024 [PubMed: 29100073]
- Toni N, Buchs PA, Nikonenko I, Bron CR, Muller D. LTP promotes formation of multiple spine synapses between a single axon terminal and a dendrite. *Nature.* 1999; 402:421–425. DOI: 10.1038/46574 [PubMed: 10586883]
- Ugawa Y. Quadripulse stimulation: Less variable? Why? *Clin Neurophysiol.* 2016; 127:e29–e30. DOI: 10.1016/j.clinph.2015.11.088
- van der Kamp W, Zwinderman AH, Ferrari MD, van Dijk JG. Cortical excitability and response variability of transcranial magnetic stimulation. *J Clin Neurophysiol Off Publ Am Electroencephalogr Soc.* 1996; 13:164–171.
- Viechtbauer W. Conducting meta-analyses in R with the metafor package. *J Stat Softw.* 2010; 36:1–48.
- Voineskos D, Rogasch NC, Rajji TK, Fitzgerald PB, Daskalakis ZJ. A review of evidence linking disrupted neural plasticity to schizophrenia. *Can J Psychiatry.* 2013; 58:86–92. [PubMed: 23442895]
- Wagner T, Valero-Cabre A, Pascual-Leone A. Noninvasive human brain stimulation. *Annu Rev Biomed Eng.* 2007; 9:527–565. DOI: 10.1146/annurev.bioeng.9.061206.133100 [PubMed: 17444810]
- Walton E, Hibar DP, van Erp TGM, Potkin SG, Roiz-Santiañez R, Crespo-Facorro B, Suarez-Pinilla P, van Haren NEM, de Zwarte SMC, Kahn RS, Cahn W, et al. Prefrontal cortical thinning links to negative symptoms in schizophrenia via the ENIGMA consortium. *Psychol Med.* 2018; 48:82–94. DOI: 10.1017/S0033291717001283 [PubMed: 28545597]
- Wilson MT, Fulcher BD, Fung PK, Robinson PA, Fornito A, Rogasch NC. Biophysical modeling of neural plasticity induced by transcranial magnetic stimulation. *Clin Neurophysiol.* 2018; 129:1230–1241. DOI: 10.1016/j.clinph.2018.03.018 [PubMed: 29674089]
- Wykes T, Huddy V, Cellard C, McGurk SR, Czobor P. A meta-analysis of cognitive remediation for schizophrenia: methodology and effect sizes. *Am J Psychiatry.* 2011; 168:472–85. DOI: 10.1176/appi.ajp.2010.10060855 [PubMed: 21406461]
- Zhou D, Pang F, Liu S, Shen Y, Liu L, Fang Z, Wang C, Hu Z, Yuan T-F. Altered Motor-Striatal Plasticity and Cortical Functioning in Patients with Schizophrenia. *Neurosci Bull.* 2017; 33:307–311. DOI: 10.1007/s12264-016-0079-9 [PubMed: 27838828]

Figure-1. Illustration of TMS-EEG or TMS-EMG cortical plasticity experiments

Figure-2. Flowchart representing the study selection process

Figure-3. Cortical plasticity (MEP) meta-analysis forest plot for all datasets

Forest plot for cortical plasticity (MEP) meta-analyses between schizophrenia patients and healthy subjects; MEP = change in motor evoked potential amplitudes following a neuromodulatory perturbation protocol. See supplementary table-S1 for more details

4A LTD-like cortical plasticity

4B LTP-like cortical plasticity

Figure-4. Cortical plasticity (MEP Δ) meta-analysis forest plots for LTD-like and LTP-like cortical plasticity

4A and 4B are forest plots for LTD-like and LTP-like cortical plasticity (MEP Δ) meta-analyses respectively; MEP Δ = change in motor evoked potential amplitudes following a neuromodulatory perturbation protocol. See supplementary table-S1 for more details

Figure-5. Bubble plots of effect sizes representing cortical plasticity (MEP) impairments in schizophrenia patients compared to healthy subjects
 X-axes (5A: generic division & 5B: specific division) represent the different neuromodulatory perturbation protocols used in the selected studies and Y-axes represent the effect sizes (size of the circle is weighted for sample size)

Table-1
Summary of different studies examining plasticity differences between patients with schizophrenia and healthy subjects

Datasets	Year	Perturbation method	Cortical reactivity	Subjects	Summary	Plasticity (MEP) in SZ
Long Term Depression (LTD) – like plasticity						
Transcranial Magnetic Stimulation (TMS) perturbation protocols						
Oxley et al., 2004#	2004	LF-rTMS	MEP [§] , RMT, SICI	12 SZ (DSM IV), 12 HS	LF-rTMS produced MEP suppression in HS as expected, but there was MEP facilitation in SZ. RMT increased in HS but reduced in SZ; there was no significant difference between group difference in SICI change	↔
(Fitzgerald, 2004)	2004	LF-rTMS	MEP [§] , RMT	26 SZ (DSM IV), 18 HS	LF-rTMS produced significantly reduced MEP suppression in SZ, compared to HS. There was no significant difference between medicated and non-medicated SZ. LF-rTMS produced an increase in RMT and diminished CSP duration in HS, but not SZ	↓
(Hasan et al., 2015) (cTBS600)	2015	cTBS	MEP	10 SZ (ICD 10), 10 HS	c-TBS-600 produced significant MEP suppression in HS but not in SZ	↓
Transcranial Direct Current Stimulation (tDCS) perturbation protocols						
(Hasan et al., 2012b)	2012	c-tDCS	MEP, RMT, CSP	21 SZ (ICD 10), 21 HS	c-tDCS produced significantly reduced (a) MEP suppression and (b) CSP duration elongation in SZ compared to HS. The RMT change was not significant in either groups	↓
(Alkomiet Hasan et al., 2013a)	2013	Left c-tDCS	MEP	9 SZ (ICD 10), 9 HS	Left c-tDCS produced reduced MEP suppression in SZ compared to HS	↓
(Hasan et al., 2012a)	2012	c-tDCS	MEP, RMT, CSP	18 SZ (ICD 10), 18 HS	c-tDCS produced significantly reduced MEP suppression on the stimulated and contralateral hemispheres in SZ compared to HS. There were no group differences in effects of c-tDCS on RMT or CSP	↓
(Alkomiet Hasan et al., 2013b)	2013	c-tDCS	MEP, RMT, CSP	15 SZ (ICD 10), 15 FDR and 10 HS	c-tDCS produced significantly reduced MEP suppression on the stimulated and contralateral hemispheres in SZ compared to HS. RMT and CSP showed no group differences after c-tDCS	↓
Long Term Potentiation (LTP) – like plasticity						
Transcranial Magnetic Stimulation (TMS) perturbation protocols						
(Hasan et al., 2015) (cTBS300)	2015	cTBS	MEP	10 SZ (ICD 10), 10 HS	c-TBS-300 produced significant MEP suppression in SZ, but MEP facilitation in HS	↔
Transcranial Direct Current Stimulation (tDCS) perturbation protocols						

Datasets	Year	Perturbation method	Cortical reactivity	Subjects	Summary	Plasticity (MEP) in SZ
(Hasan et al., 2011)	2011	a-tDCS	MEP, CSP, SICI, ICF	22 SZ (ICD 10), 22 HS	a-tDCS produced significantly reduced MEP facilitation in multiple-episode SZ compared to recent-onset SZ and HS. There was no group difference for change in CSP, SICI and ICF	↓ / ≈
(Alkomiet Hasan et al., 2013a)	2013	Left c-tDCS + Right a-tDCS	MEP	9 SZ (ICD 10), 9 HS	Left c-tDCS + Right a-tDCS produced significantly reduced MEP facilitation in SZ compared to HS	↓
(Strube et al., 2016) (a-tDCS)	2016	a-tDCS	MEP, SICI, ICF	20 SZ (ICD 10), 20 HS	a-tDCS demonstrated MEP facilitation in SZ and HS. SICI and ICF did not show any group differences following a-tDCS	≈
Paired Associative Stimulation (PAS) perturbation protocols						
(Frantseva et al., 2008)	2008	PAS	MEP	15 SZ or SA (DSM IV), 15 HS	PAS produced significantly reduced MEP facilitation in SZ compared to HS	↓
(Ribolsi et al., 2017)	2017	PAS	MEP	12 SZ (DSM IV), 12 HS	Parieto-motor PAS produced reduced and slower MEP facilitation in SZ (more so in the left hemisphere), compared to HS	↓
(Strube et al., 2016) (PAS)	2016	PAS	MEP, SICI, ICF	20 SZ (ICD 10), 20 HS	PAS produced significantly reduced MEP facilitation in SZ as compared to HS. SICI and ICF did not show any group differences following PAS	↓
(Bridgman et al., 2016)	2016	PAS	MEP	9 SZ (DSM IV), 10 HS	*Reduced MEP facilitation following PAS in SZ as compared to HS	↓
Others						
(Daskalakis et al., 2008)	2008	Exercise	Angular displacement	20 SZ (6 antipsychotic free) (DSM IV), 20 HS	Exercise induced significantly reduced mean angular displacement in SZ compared to HS, irrespective of their medication status	↓

Note: SZ = schizophrenia patients; HS = healthy subjects; SA = schizoaffective disorder; FDR = first degree relatives; MEP = motor evoked potentials; RMT = resting motor threshold; CSP = cortical silent period; SICI = short interval intracortical inhibition; ICF = intracortical facilitation; c-tDCS = cathodal tDCS; a-tDCS = anodal tDCS; cTBS = continuous theta burst stimulation; all perturbation protocols were delivered to the motor cortex, except [#]Oxley et al., which was delivered to the premotor cortex; all MEP recordings were performed using SI_{1m}V as the stimulation dose, except ^{\$}Oxley et al., and ^{\$}Fitzgerald et al., where the stimulating doses ranged between 102% to 120% RMT; *results reported from the placebo arm of the study; MEP = change in motor evoked potential amplitudes following a neuromodulatory perturbation protocol; ↓ = lesser than healthy subjects; ≈ = same as healthy subjects; ↔ = opposite to healthy subjects