Published in final edited form as:

Acta Neuropathol. 2019 March ; 137(3): 527–530. doi:10.1007/s00401-018-01955-0.

Antisense RNA foci are associated with nucleoli and TDP-43 mislocalization in C9orf72-ALS/FTD: A quantitative study

Olubankole Aladesuyi Arogundade^{1,*}, Jennifer E. Stauffer^{1,2,*}, Shahram Saberi^{1,3,*}, Sandra Diaz Garcia¹, Sahana Malik^{1,4}, Hani Basilim^{1,5}, Maria J Rodriguez¹, Takuya Ohkubo¹, and John Ravits^{1,†}

¹Department of Neuroscience, University of California, San Diego, 9500 Gilman Drive, La Jolla, CA, 92093, MS-0624, USA

²Present address: The Jackson Laboratory, 600 Main St, Bar Harbor, ME 04609, USA

³Present address: Department of Pathology, Icahn School of Medicine at Mount Sinai, New York, 1 Gustave L. Levy Place, Box 1194, NY 10029, USA

⁴Present address: Cooper Medical School of Rowan University, 401 South Broadway, Camden, NJ 08103, USA

⁵Present address: King Abdullah Medical Complex Jeddah (KAMCJ), Ameer Naif Road, Al-Shera'a, Jeddah 23816, Saudi Arabia

Three main mechanisms are thought to contribute to neurodegeneration in C9ORF72 amyotrophic lateral sclerosis (ALS) and frontotemporal dementia (FTD) (C9-ALS/FTD): toxicity from transcribed expanded repeat RNAs, toxicity from RAN-translated dipeptide repeat proteins (DPRs), and loss of C9ORF72 protein function [4, 5, 9, 11, 12, 18]. Sense and antisense RNA foci have both been consistently observed in C9-ALS/FTD neuropathology [1, 3, 5, 8, 9, 18] and hypothesized to cause neurodegeneration by sequestering critical RNA binding proteins [2, 6]. Antisense but not sense RNA foci have been shown to correlate with mislocalization of TDP-43, a signature protein of ALS and frontal temporal lobar degeneration (FTLD) [1]. A unique circumferential studding of nucleoli by antisense RNA foci was observed in a case report of two C9-FTLD cases [17] and recently re-discussed with two additional cases [16]. Understanding the relative contributions from sense and antisense strands to pathogenesis is critical, since antisense oligonucleotides (ASOs) are expected to ameliorate toxicity whether stemming from RNAs or DPRs.

We evaluated RNA foci in five C9-ALS cases with and without FTD in our short-postmortem interval ALS repository [13]. C9-ALS/FTD was confirmed by repeat-primed PCR and Southern blotting. Consistent with the literature, both sense and antisense RNA foci were widely distributed throughout the nervous systems [1, 3, 5, 9, 18]. They were not

[†]Address correspondence: jravits@ucsd.edu.

*Equal contributions

Conflict of Interest

The authors declare that they have no conflict of interest.

concentrated in the motor or frontal cortex, the anatomical regions of direct relevance to C9-ALS/FTD (Supplemental Fig. 1a and b). Overall, the mean prevalence of sense and antisense RNA foci over multiple CNS regions were 16% and 20% in neurons and 1% and 1% in glia, respectively (Supplemental Fig. 1c). Antisense RNA foci were significantly denser than sense RNA foci within the RNA foci-positive neurons ($p < 0.0001$; Supplemental Fig. 1d). As reported in C9-FTD and one case of C9-ALS, we also observed that in large neurons, RNA foci were frequently localized to nucleoli, identified as a prominent immunofluorescent density within the nucleus, often directly abutting their circumference (Fig. 1a). We confirmed nucleolar localization using the nucleolar marker nucleolin (Fig. 1b). We quantified the association and overall on average, 27% of antisense RNA foci but only 9% of sense RNA foci were perinucleolar. We saw that perinucleolar antisense RNA foci were more frequent in disease-related areas than non-disease-related areas and were more likely than sense RNA foci to be perinucleolar in disease-related areas but this did not reach significance in disease-unrelated areas: 35% of antisense RNA foci but only 13% of sense foci in disease-related areas were perinucleolar, compared to 17% and 4%, respectively, in disease-unrelated areas (Supplemental Fig. 1e and f). The preponderance of nucleolar antisense RNA foci but not sense RNA foci tended to correlate with the burden of TDP-43 pathology in frontal cortex (Supplemental Table 1).

We evaluated the relationship between RNA foci and TDP-43 mislocalization using co-fluorescent *in situ* hybridization and immunofluorescence. We did this in neurons in the anterior horn of spinal cord, where identification of motor neurons by location in the anterior horn is readily done (Fig. 1c and d). Nucleolar localization of RNA foci was confirmed by imaging through the z-plane. As previously reported [1], antisense RNA foci were associated with TDP-43 pathology and sense foci were not: antisense foci were seen in 67% of neurons with cytoplasmic aggregation of TDP-43 and 16% of neurons with preserved nuclear TDP-43 ($p = 0.008$; data not shown), compared to sense foci which were seen in 20% of neurons with cytoplasmic aggregation of TDP-43 and 18% of neurons with preserved nuclear TDP-43 ($p = 0.5$, data not shown). Importantly, the difference between antisense and sense foci held true for the perinucleolar RNA foci: perinucleolar antisense RNA foci were seen in 67% of neurons with cytoplasmic aggregation of TDP-43 and 20% of neurons with preserved nuclear TDP-43 ($p = 0.008$; Fig. 1e), compared to perinucleolar sense RNA foci, which were seen in 9% of neurons with cytoplasmic TDP-43 and 11% of neurons with preserved nuclear TDP-43 ($p = 0.7$; Fig. 1f and Table 1). Overall, the co-occurrence of TDP-43 abnormalities and perinucleolar antisense RNA foci was significantly greater than expected ($p < 0.001$; Fig. 1g), while the co-occurrence of TDP-43 abnormalities and perinucleolar sense RNA foci was as expected ($p = 0.6$; Fig. 1g).

The association of antisense RNA foci and nucleoli is important in light of recent discussions about nucleolar stress contributing to the pathogenesis of C9-ALS/FTD [6, 7, 10, 15]. The nucleolus is the site of many localized functions such as ribosomal biogenesis and disruptions to nucleolar function can dramatically affect many functions such as rates of protein synthesis. Alterations in nucleolar morphology as well as altered rates of protein synthesis have been found in C9-FTLD/ALS frontal cortex [10] and cellular model [6, 15], although not confirmed by all [14]. Thus far, sense RNA foci [6] and the arginine-containing DPRs, poly-GR and poly-PR [7, 15] have been implicated to cause or at least to be

associated with nucleolar stress in C9-ALS/FTD pathogenesis. While antisense RNA foci were previously found to correlate with mislocalization of TDP-43 in C9-ALS [1] and antisense RNA foci studding of nucleoli have been observed in C9-FTLD [17], the co-occurrence of nucleolar antisense RNA foci with TDP-43 neuropathology has not been reported or quantitated. In our study, we found that nucleolar antisense RNA foci are significantly more associated with disease, especially with neuronal TDP-43 cytoplasmic aggregation and our findings bring the different aspects together. It has been suggested that nucleolar antisense RNA foci may form early in disease and precede TDP-43 aggregation [17].

In summary, antisense RNA foci have unique neuropathological features compared to sense RNA foci including nucleolar localization, anatomic distribution, and correlation with TDP-43 pathology. This re-enforces that antisense RNA may play an important role in pathogenesis and be an important target for therapy.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

This research was supported by grants from ALS Association (5356S3), Target ALS (20134792), National Institute of Neurological Diseases and Stroke (NIH R01NS088578 and NS047101), and Pam Golden. OAA is supported by National Science Foundation Graduate Research Fellowship (DGE-1650112). TO is supported by NINDS CReATe Consortium (U54NS092091). Len Petrucelli provided extra sense probe for FISH. We thank the patients and their families for their generous contribution to this research.

References

1. Cooper-Knock J, Higginbottom A, Stopford MJ, Highley JR, Ince PG, Wharton SB, et al. (2015) Antisense RNA foci in the motor neurons of C9ORF72-ALS patients are associated with TDP-43 proteinopathy. *Acta Neuropathol* 130:63–75. doi: 10.1007/s00401-015-1429-9. [PubMed: 25943887]
2. Cooper-Knock J, Walsh MJ, Higginbottom A, Highley JR, Dickman MJ, Edbauer D, et al. (2014) Sequestration of multiple RNA recognition motif-containing proteins by C9orf72 repeat expansions. *Brain* 137:2040–2051. doi: 10.1093/brain/awu120. [PubMed: 24866055]
3. DeJesus-Hernandez M, Finch NCA, Wang X, Gendron TF, Bieniek KF, Heckman MG, et al. (2017) In-depth clinico-pathological examination of RNA foci in a large cohort of C9ORF72 expansion carriers. *Acta Neuropathol* 134(2):255–269. doi: 10.1007/s00401-017-1725-7. [PubMed: 28508101]
4. DeJesus-Hernandez M, Mackenzie IR, Boeve BF, Boxer AL, Baker M, Rutherford NJ, et al. (2011) Expanded GGGGCC Hexanucleotide Repeat in Noncoding Region of C9ORF72 Causes Chromosome 9p-Linked FTD and ALS. *Neuron* 72(2):245–56. doi: 10.1016/j.neuron.2011.09.011. [PubMed: 21944778]
5. Gendron TF, Bieniek KF, Zhang YJ, Jansen-West K, Ash PEA, Caulfield T, et al. (2013) Antisense transcripts of the expanded C9ORF72 hexanucleotide repeat form nuclear RNA foci and undergo repeat-associated non-ATG translation in c9FTD/ALS. *Acta Neuropathol* 126:829–844. doi: 10.1007/s00401-013-1192-8. [PubMed: 24129584]
6. Haeusler AR, Donnelly CJ, Periz G, Simko EAJ, Shaw PG, Kim MS, et al. (2014) C9orf72 nucleotide repeat structures initiate molecular cascades of disease. *Nature* 507(7491):195–200. doi: 10.1038/nature13124. [PubMed: 24598541]

7. Kwon I, Xiang S, Kato M, Wu L, Theodoropoulos P, Wang T, et al. (2014) Poly-dipeptides encoded by the C9orf72 repeats bind nucleoli, impede RNA biogenesis, and kill cells. *Science* 345(6201): 1139–45. doi: 10.1126/science.1254917. [PubMed: 25081482]
8. Lagier-Tourenne C, Baughn M, Rigo F, Sun S, Liu P, Li H-R, et al. (2013) Targeted degradation of sense and antisense C9orf72 RNA foci as therapy for ALS and frontotemporal degeneration. *Proc Natl Acad Sci* 110(47):E4530–9. doi: 10.1073/pnas.1318835110. [PubMed: 24170860]
9. Mizielińska S, Lashley T, Norona FE, Clayton EL, Ridler CE, Fratta P, et al. (2013) C9orf72 frontotemporal lobar degeneration is characterised by frequent neuronal sense and antisense RNA foci. *Acta Neuropathol* 126:845–857. doi: 10.1007/s00401-013-1200-z. [PubMed: 24170096]
10. Mizielińska S, Ridler CE, Balendra R, Thoeng A, Woodling NS, Grässer FA, et al. (2017) Bidirectional nucleolar dysfunction in C9orf72 frontotemporal lobar degeneration. *Acta Neuropathol Commun* 5(1):29. doi: 10.1186/s40478-017-0432-x. [PubMed: 28420437]
11. Mori K, Arzberger T, Grässer FA, Gijssels I, May S, Rentzsch K, et al. (2013) Bidirectional transcripts of the expanded C9orf72 hexanucleotide repeat are translated into aggregating dipeptide repeat proteins. *Acta Neuropathol* 126:881–893. doi: 10.1007/s00401-013-1189-3. [PubMed: 24132570]
12. Renton AE, Majounie E, Waite A, Simón-Sánchez J, Rollinson S, Gibbs JR, et al. (2011) A hexanucleotide repeat expansion in C9ORF72 is the cause of chromosome 9p21-linked ALS-FTD. *Neuron* 72(2):257–68. doi: 10.1016/j.neuron.2011.09.010. [PubMed: 21944779]
13. Saberi S, Stauffer JE, Jiang J, Garcia SD, Taylor AE, Schulte D, et al. (2018) Sense-encoded poly-GR dipeptide repeat proteins correlate to neurodegeneration and uniquely co-localize with TDP-43 in dendrites of repeat-expanded C9orf72 amyotrophic lateral sclerosis. *Acta Neuropathol* 135(3): 459–474. doi: 10.1007/s00401-017-1793-8. [PubMed: 29196813]
14. Schludi MH, May S, Grässer FA, Rentzsch K, Kremmer E, Küpper C, et al. (2015) Distribution of dipeptide repeat proteins in cellular models and C9orf72 mutation cases suggests link to transcriptional silencing. *Acta Neuropathol*. 130(4):537–55. doi: 10.1007/s00401-015-1450-z. [PubMed: 26085200]
15. Tao Z, Wang H, Xia Q, Li K, Li K, Jiang X, et al. (2015) Nucleolar stress and impaired stress granule formation contribute to C9orf72 RAN translation-induced cytotoxicity. *Hum Mol Genet* 24:2426–2441. doi: 10.1093/hmg/ddv005. [PubMed: 25575510]
16. Vatsavayi SC, Nana AL, Yokoyama JS, Seeley WW (2018) C9orf72-FTD/ALS pathogenesis: evidence from human neuropathological studies. *Acta Neuropathol* doi: 10.1007/s00401-018-1921-0 [Epub ahead of print].
17. Vatsavayi SC, Yoon SJ, Gardner RC, Gendron TF, Vargas JNS, Trujillo A, et al. (2016) Timing and significance of pathological features in C9orf72 expansion-associated frontotemporal dementia. *Brain* 139(Pt 12):3202–3216. [PubMed: 27797809]
18. Zu T, Liu Y, Banez-Coronel M, Reid T, Pletnikova O, Lewis J, et al. (2013) RAN proteins and RNA foci from antisense transcripts in C9ORF72 ALS and frontotemporal dementia. *Proc Natl Acad Sci* 110(51):E4968–77. doi: 10.1073/pnas.1315438110. [PubMed: 24248382]

Fig. 1. Antisense RNA foci have unique neuropathological features compared to sense RNA foci.

(a): Sense and antisense RNA foci (arrowheads) are observed in the nucleoplasm or on the periphery of nucleoli of neurons (outer, middle and innermost dashed lines mark the cytoplasm, nucleoplasm and nucleolus, respectively). (b): Nucleolar localization is confirmed by nucleolar marker nucleolin. (c and d): Neurons containing antisense RNA foci often have cytoplasmic aggregation of TDP-43 and neurons with sense RNA foci often have preserved nuclear TDP-43 (inset scale, 500 nm). (e): The percentage of neurons with nucleolar antisense RNA foci was significantly increased in neurons with cytoplasmic aggregation of TDP-43 compared to preserved nuclear TDP-43 ($p = 0.008$). (f): The percentage of neurons with nucleolar sense RNA foci was not significantly different in neurons with cytoplasmic aggregation of TDP-43 compared to preserved nuclear TDP-43 ($p = 0.7$, MannWhitney test). (g): The co-occurrence of nucleolar antisense RNA foci and cytoplasmic aggregation of TDP-43 is significantly greater than expected while the co-occurrence of nucleolar sense RNA foci and cytoplasmic

aggregation of TDP-43 is not. Area proportional Venn-diagrams represent Fisher's exact test. Data is from motor neurons within the anterior horn of spinal cord.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1:

Cytoplasmic aggregation of TDP-43 and nucleolar RNA foci in motor neurons within the anterior horn of spinal cord by co-FISH-IF

		Nucleolar RNAfoci				
		Cytoplasmic TDP-43		Nuclear TDP-43		
		Observed	Expected	Observed	Expected	<i>p</i> -value
Antisense	Foci-positive	43	34	32	41	< 0.001
	Foci-negative	2	11	21	12	
Sense	Foci-positive	8	9	24	23	0.6
	Foci-negative	9	8	19	20	