CORRESPONDENCE

Treatment Options for Abdominal Obesity in Adults

A Meta-Analysis and Systematic Review of Randomized Controlled Trials

by Dr. biol. hum. Dorothea Kesztyüs, Julia Erhardt, Dr. med. Dorothee Schönsteiner, and Prof. Dr. med. Tibor Kesztyüs in issue 29–30/2018

Confusing Terminology

In the key statement of their important review article, Kesztyüs and colleagues express the following demand: “Further research regarding the treatment of abdominal obesity is urgently needed.” I hope they mean studies with clinically relevant endpoints (mortality, morbidity) and not merely studies with surrogate outcomes—for example, “reduction of abdominal obesity,” as one might deduce from the conclusions of their article.

This uncertainty may also be related to the fact that the authors seem to be confused about their terminology. Erroneously, they mention “pharmaceutical trials,” “pharmaceutical interventions,” and “pharmacological interventions.”

Pharmaceutical trials investigate the quality (identity, ingredients, purity, and so on) of a medical drug. Efficacy and harmlessness are, however, determined in clinical-pharmacological (phase I) and clinical (phase II–IV) trials.

Terminology problems are not merely a formality, as is shown by a further comment relating to a study reported by Tuomilehto et al. (2), in which the waist circumference of obese participants with prediabetes decreased by 4.4 cm and their relative risk of developing type 2 diabetes fell by 58%.

Kesztyüs et al., however, mentioned “a 58% reduction in the incidence of type 2 diabetes mellitus.” Since the incidence relates to the onset of an event—that is, the absolute risk—58% shrinks to 12% (control group 23%, intervention group 11%).

Unclear terminology is not trivial but can contribute to misunderstandings.

DOI: 10.3238/arztebl.2019.0039a

References

1. Kesztyüs D, Erhardt J, Schönsteiner D, Kesztyüs T: Treatment options for abdominal obesity in adults—a meta-analysis and systematic review of randomized controlled trials. *Dtsch Arztebl Int* 2018; 115: 487–93.
2. Tuomilehto J, Lindström J, Eriksson JG, et al.: Prevention of type 2 diabetes mellitus by changes in lifestyle among subjects with impaired glucose tolerance. *N Engl J Med* 2001; 344: 1343–50.

Prof. Dr. med. Frank P. Meyer
Wanzleben-Börde
U_F_Meyer@gmx.de

Creating Associations With Positive Experiences Is Key

Kesztyüs et al. in their excellent review article confirm that for abdominal obesity, among others, a disciplined, long-term adaptation of dietary habits and lifestyle (365

days/year) is the method of choice (1). In terms of the organismic required muscle activity, allow me to add a recent French meta-analysis to the authors’ reference list. The working group of Maillard et al. confirmed high intensity interval training (HIIT) as a time efficient strategy for reducing abdominal-visceral fat especially in overweight or obese persons, independent of sex (2). In view of the difficulties that many people experience with committing to exercise, we should not underestimate the degree to which overweight/obese persons associate HIIT with positive experiences and perceive it as a feasible and enjoyable form of physical activity (3, 4). For this reason, combined lifestyle interventions in abdominal obesity should—as far as the individual training/health status of the patient permits—integrate into everyday life frequently repeated, short intensive exercise elements with realistic implementation steps.

DOI: 10.3238/arztebl.2019.0039b

References

1. Kesztyüs D, Erhardt J, Schönsteiner D, Kesztyüs T: Treatment options for abdominal obesity in adults—a meta-analysis and systematic review of randomized controlled trials. *Dtsch Arztebl Int* 2018; 115: 487–93.
2. Maillard F, Pereira B, Boisseau N: Effect of high-intensity interval training on total, abdominal and visceral fat mass: a meta-analysis. *Sports Med* 2018; 48: 269–88.
3. Kong Z, Fan X, Sun S, Song L, Shi Q, Nie J: Comparison of high-intensity interval training and moderate-to-vigorous continuous training for cardiometabolic health and exercise enjoyment in obese young women: a randomized controlled trial. *PLoS One* 2016; 11: e0158589.
4. Smith-Ryan AE: Enjoyment of high-intensity interval training in an overweight/obese cohort: a short report. *Clin Physiol Funct Imaging* 2017; 37: 89–93.

Dr. oec. troph. Martin Hofmeister
Verbraucherzentrale Bayern e. V.
Referat Lebensmittel und Ernährung
München
hofmeister@vzbayern.de

More Effective Therapeutic Approaches Should Be Deployed

Restricting the literature analysis to non-surgical treatment options (1) rules out the most effective therapy—that is, obesity surgery and metabolic surgery. This narrow perspective is not justified in scientific terms, since surgery for obesity and type 2 diabetes has long been accepted by the S3 guideline (2).

The authors included in their meta-analysis 15 randomized trials of modifications of behavior, nutrition, and physical activity/exercise as well as pharmaceutical approaches. The mean waist circumference—a

measure of visceral obesity—was reduced by merely 2.65 cm, to 100.05 cm; even after combined lifestyle interventions, the loss was a mere 4.11 cm. To call such a treatment result moderate is simply a euphemism, in view of the defined thresholds. The achieved result is in most cases clinically irrelevant and, furthermore, is not maintained in the long term, and therefore it does not justify the enormous resource consumption.

Admittedly, prevention would constitute the only solution for this chronic disorder. Non-operative treatment modalities have been proved to fail in most patients in its current form. In routine clinical practice, the therapeutic goals—defined in the S3 guideline as long-term weight loss >15% (20%) kg/m² in persons with a BMI >35 (40) kg/m²—are achieved only in exceptional situations.

Consequently, a new, different, more effective therapeutic approach needs to be deployed—and according to what is currently known this is the combination with surgical methods. In the meantime, the guideline has conceded the pointlessness of conservative methods alone in patients with a BMI >50 kg/m². This is the threshold for the guideline to recommend the primary indication for surgery.

Optimizing obesity treatments requires a paradigm shift: early surgery and intensified conservative treatment afterwards. The responsibility for structured long-term follow-up must return to general practitioners and family physicians—this is relevant for reimbursement purposes.

DOI: 10.3238/arztebl.2019.0039c

References

1. Kesztyüs D, Erhardt J, Schönsteiner D, Kesztyüs T: Treatment options for abdominal obesity in adults—a meta-analysis and systematic review of randomized controlled trials. *Dtsch Arztebl Int* 2018; 115: 487–93.
2. S3-Leitlinie: Chirurgie der Adipositas und metabolischer Erkrankungen. www.awmf.org/uploads/tx_szleitlinien/088-001_S3_Chirurgie-Adipositas-metabolische-Erkrankungen_2018-02.pdf (last accessed on 10 October 2018).

Prof. Dr. med. Norbert Runkel

Klinik für Adipositaschirurgie und metabolische Chirurgie
Exzellenzzentrum, Sana Klinikum Offenbach
Norbert.Runkel@sana.de

Dr. med. Sonja Chiappetta

Prof. Dr. med. Rudolf Weiner

In Reply:

We thank the correspondents for their contributions in response to our systematic review of therapeutic approaches in abdominal obesity. Confusing terminology does not change the sobering result of our review—namely, that a blatant lack of effective measures and research exists in this area. In all this, the notable exception is obviously in particular bariatric surgery, which we excluded from our literature search as it is an invasive method that requires a separate review—and, of course, appreciation. This perspective is also represented in the review by Peirson et al. of treatments for general overweight and obesity, since surgical interventions concern an extent of obesity and comorbid states that is seen less often in primary care (1).

We hope that our review (2) contributed to putting abdominal obesity as a high-risk form of obesity into greater focus for doctors' attention and to clarify that it is not merely a problem of esthetics. The comment about high intensity interval training (HIIT) was most welcome, as this is a method that would probably find wide acceptance by those affected because of the way in which it is conceived. In conclusion we wish to emphasize that a fundamental lifestyle change in each phase of the pathogenesis and development of abdominal obesity and its comorbidities is to be recommended—whether that be in prevention, therapy, or rehabilitation.

DOI: 10.3238/arztebl.2019.0040

References

1. Peirson L, Douketis J, Ciliska D, Fitzpatrick-Lewis D, Ali MU, Raina P: Treatment for overweight and obesity in adult populations: a systematic review and meta-analysis. *CMAJ Open* 2014; 2: E306–17.
2. Kesztyüs D, Erhardt J, Schönsteiner D, Kesztyüs T: Treatment options for abdominal obesity in adults—a meta-analysis and systematic review of randomized controlled trials. *Dtsch Arztebl Int* 2018; 115: 487–93.

Dr. biol. hum. Dorothea Kesztyüs, MPH

Institut für Allgemeinmedizin
Universität Ulm
dorothea.kesztues@uni-ulm.de

Julia Erhardt

Dr. med. Dorothee Schönsteiner

Prof. Dr. med. Tibor Kesztyüs

Conflict of interest statement

All correspondents declare that no conflict of interest exists.