Published in final edited form as:

J Am Chem Soc. 2019 February 20; 141(7): 2825–2831. doi:10.1021/jacs.8b13192.

Aminoxyl-Catalyzed Electrochemical Diazidation of Alkenes Mediated by a Metastable Charge-Transfer Complex

Juno C. Siu[‡], Joseph B. Parry[‡], and Song Lin^{*}

Department of Chemistry and Chemical Biology, Cornell University, Ithaca, New York 14853, United States

Abstract

We report the development of a new aminoxyl radical catalyst, CHAMPO, for the electrochemical diazidation of alkenes. Mediated by an anodically generated charge-transfer complex in the form of CHAMPO–N₃, radical diazidation was achieved across a broad scope of alkenes without the need for a transition metal catalyst or a chemical oxidant. Mechanistic data support a dual catalytic role for the aminoxyl serving as both a single-electron oxidant and a radical group transfer agent.

The discovery of reactions mediated by organic radicals continues to provide solutions to challenging synthetic problems in traditional two-electron chemistry.¹ In this context, design and implementation of new catalytic strategies have both expanded the toolbox available for accessing new synthetic targets and transformed the fundamental understanding of reactions involving open-shell pathways.² For example, persistent aminoxyl radicals [e.g., (2,2,6,6-tetramethylpiperidin-1-yl)oxyl (TEMPO)] have been extensively explored in catalytic oxidation reactions with both conventional chemical³ and electrochemical techniques,⁴ which has given rise to synthetically useful processes for small-molecule and polymer syntheses.

Despite significant advances, we contend that the scope of TEMPO chemistry remains to be fully explored. TEMPO and related *N*-oxyl radicals can undergo one-electron redox processes, granting access to three discrete oxidation states.⁴ This feature distinguishes these radicals from common organic compounds and likens them to many transition metal complexes. In this fashion, TEMPO has been shown to enable single-electron oxidation events in an inner-sphere manner via the formation of metastable closed-shell intermediates.^{5,6} Nevertheless, the systematic use of the “metallic” character of *N*-oxyls in catalyst development remains meager.⁷ To date, reactions catalyzed by *N*-oxyls are largely confined to oxidations of alcohols,⁸ aldehydes,⁹ amines,¹⁰ (thio)amides,¹¹ and peroxy radicals.¹² In

^{*}Corresponding Author songlin@cornell.edu.

[‡]These authors contributed equally.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/jacs.8b13192.

Experimental procedures and characterization data (PDF)

Crystallographic data for CHAMPO (CCDC1881714) (CIF)

Notes

The authors declare no competing financial interest.

these transformations, the persistent radical is used primarily in two capacities: as a single/two-electron oxidant or an H atom abstractor. In this report, we expand the scope of *N*-oxyl catalysis in the development of metal-free electrochemical diazidation of alkenes.

We previously reported a first-generation electrochemical protocol for the diazidation of alkenes catalyzed by Mn (Scheme 1A, eq 1).¹³ This work was built on foundational contributions by Minisci,¹⁴ Magnus,¹⁵ Snider,¹⁶ Xu,¹⁷ and others¹⁸ in alkene diazidation using conventional chemical methods. These reactions provide 1,2-diazides in a single step from common alkenes, giving access to vicinally dinitrogenated structures that are highly prevalent in biomedically and synthetically relevant molecules.¹⁹ Nevertheless, all methods reported to date require the use of a transition metal or a chemical oxidant, or both. These reactive species often lead to environmentally deleterious wastes and present an explosion hazard when used alongside N_3^- . Although our electro-chemical protocol¹³ circumvents the use of strong chemical oxidants, the need for a metal catalyst and a Brønsted acid (HOAc) as the sacrificial oxidant remains undesirable from a safety perspective.

While studying the electrochemical alkene diazidation, we discovered a new electrochemical method for the azidoxylation of alkenes mediated by TEMPO (Scheme 1A, eq 2).²⁰ Detailed mechanistic investigation showed that TEMPO plays two roles in this reaction. Aside from being a radical trap, TEMPO also promotes the single-electron oxidation of N_3^- to $N_3^{\bullet}$.²¹ This process is mediated by a metastable charge-transfer complex (CTC) formed between anodically generated TEMPO⁺ and N_3^- . Upon azidyl addition to the alkene, the nascent carbon-centered radical I rapidly combines with TEMPO to deliver the masked vicinal aminoalcohol (Scheme 1B, red pathway).

We envisioned that the TEMPO-consuming azidoxylation reaction could be engineered into a TEMPO-catalyzed alkene diazidation if intermediate I was terminated by an $N_3^{\bullet}$ equivalent instead of directly by TEMPO (Scheme 1B, blue pathway). Specifically, we hypothesize that the TEMPO- N_3 adduct may behave as an azidyl group transfer agent akin to the function of a redox-active transition metal complex (e.g., $[Mn^{III}]-N_3$). This operation will furnish the desired vicinal diazide while returning the aminoxyl from an oxoammonium ion to a neutral radical oxidation state.

We investigated the plausibility of *N*-oxyl-catalyzed electrochemical diazidation (Table 1). In our alkene azidoxylation, diazidation products were occasionally observed in small quantities when sterically encumbered alkenes were used. The diazide formation was more pronounced at a higher applied voltage and using less TEMPO. Under these conditions, the $[TEMPO^+]/[TEMPO]$ ratio is maximized to favor the formation of the CTC and mitigate the undesired azidoxylation. For example, the electrolysis of alkene **1a**, NaN_3 , and 1 equiv TEMPO in MeCN/H₂O at a cell voltage of 2.7 V²² yielded 36% 1,2-diazide **2a** in addition to 45% **3a** (entry 1). Decreasing TEMPO loading to a catalytic amount still provided encouragingly 11% **2a** alongside 7% **3a** (entry 2). However, the catalyst was unproductively consumed in the azidoxylation pathway, which was also indicated by the color change of the reaction medium. Upon starting electrolysis, the solution turned dark red, which is characteristic of the TEMPO- N_3 CTC. This color disappeared after a mere 10 min as a

result of TEMPO depletion. This visual indicator thus provided a convenient readout for reaction optimization.

We reasoned that heating the reaction would promote homolysis of the C–O bond in byproduct **3a**, thus returning both TEMPO and the carbon-centered radical to the catalytic cycle.²³ Counterintuitively, this operation led to rapid decomposition of the CTC and suppressed diazidation. Instead, cooling the reaction improved the yield to 46% at 0 °C and 61% at –10 °C (entries 3 and 4). However, the color of the CTC decayed after 2 h with concomitant formation of **3a** (ca. 5%).

Studies by Bobbitt,^{8k} Minter/Sigman,²⁴ and Stahl²⁵ showed that the distal C4 substituent of the aminoxyl has a profound influence on its reactivity in alcohol oxidation. Accordingly, we surveyed various catalysts²⁶ and found that 4-acetamido-TEMPO (ACT) proved the most promising, furnishing 71% **2a** and <5% **3a** (entry 5). However, the CTC decayed after ca. 3 h. We hypothesized that catalyst degradation could also occur via β -H⁺ elimination of the corresponding oxoammonium ion²⁷ in a manner akin to the Hofmann elimination.

We reasoned that increasing the steric encumbrance about the *N*-oxyl center should disfavor catalyst consumption via azidooxygenation²⁸ and β -elimination.²⁹ Accordingly, we synthesized cyclohexane-substituted (4-acetamidopiperidin-1-yl)oxyl (CHAMPO).^{27,30} With CHAMPO as the electrocatalyst, the lifetime of the CTC significantly improved and the color of the solution persisted upon full conversion of the alkene (4 h).³¹ Diazide **2a** was obtained in 97% isolated yield with complete elimination of the *N*-oxyl trapping product (entry 6). Importantly, exclusion of catalyst led to olefin consumption but negligible diazide formation (entry 7).³² In addition, common single-electron oxidants in lieu of electricity did not provide appreciable diazide.³³ We observed the same catalyst trend using trisubstituted alkene **1b**: from TEMPO to ACT to CHAMPO, improvement in product yield and catalyst stability was evident (entries 8–10).

We subsequently explored the scope of this new electrochemical diazidation (Table 2). A broad range of alkenes with diverse structures and electronic properties underwent the desired transformation with high efficiency (**2a**, **c–p**). Unactivated monosubstituted alkenes were less reactive toward N₃[•] addition and more prone to azidooxygenation due to reduced steric hindrance.^{20a,34} Therefore, 15 mol % CHAMPO was used to ensure synthetically useful yields (**2c**, **d**).

Alkenes with a catalog of labile functional groups underwent electrochemical diazidation smoothly (**2b**, **q–y**). For example, electrophiles prone to nucleophilic displacement by N₃[–], such as alkyl bromide and epoxide groups, remained intact. Oxidatively sensitive functionalities, such as enolizable ketone, aldehyde, sulfide, and ferrocene groups, also were tolerated. Intriguingly, a primary alcohol-derived alkene was converted to the corresponding diazide in 60% yield with traces of aldehyde byproducts. Chemoselectivity between various C=C bonds was observed with carvone or *N*-allyl-*N*-prenyltoluenesulfonamide as substrate. Owing to the electrophilic nature of N₃[•], the more electron-releasing alkene was preferentially diazidated.³⁵

The 1,2-diazide products can be readily reduced to the corresponding vicinal diamines using Staudinger reduction or catalytic hydrogenation. Notably, the different steric environments of the two azido groups in **2k** enabled chemoselective reduction to form β -azidoamide **5**. The diazides could also be readily converted to bistriazoles or 1,2-dinitro compounds,³⁶ which are common structures in energetic materials.

Compared with our first-generation Mn-catalyzed alkene diazidation, the new protocol presents several practical advances. For example, replacing the transition metal catalyst with an organocatalyst and the protic acid (HOAc) with H₂O eliminates the generation of hazardous metal azides and hydrazoic acid. These features make the diazide synthesis safer and more amenable to practical applications. We conducted further experiments to exemplify the user friendliness of the protocol (Table 3). Using inexpensive Ni foam as the cathode instead of Pt only marginally decreased product yield (entry 2). Using carbon as both cathode and anode did not provide **2k** in appreciable amounts. To make the reaction system metal-free, we carried out electrolysis in a divided cell that completely separates the diazidation from the cathodic proton reduction on Pt (entry 3). The electrolysis was also successfully performed using commercial ElectraSyn 2.0³⁷ or AA batteries as the power source (entries 4 and 5). Electrolyte choice proved unimportant (entry 6). In fact, given the ionic nature of NaN₃, electrolysis in the absence of an additional electrolyte yielded **2k** quantitatively (entry 7). Finally, diazidation on a synthetically relevant scale resulted in a promising isolated yield of 69% (entry 8).

Aside from the radical pathway in our initial working mechanistic hypothesis (see Scheme 1B), two additional reaction pathways may be plausible to account for diazidation reactivity. Carbon-centered radical **I** may undergo single-electron oxidation, either directly on the anode or by CHAMPO⁺, to generate the corresponding carbocation (**II**; Scheme 2A). The cation would subsequently react with N₃⁻ to form the diazide. Alternatively, azidooxygenated product **3** could be an intermediate en route to the diazide via oxidation and mesolytic cleavage³⁸ to form **II** (Scheme 2B).

Experimental evidence (Scheme 3) is consistent with our original mechanistic proposal in which the aminoxyl–N₃ adduct mediates the radical diazidation, whereas for certain types of substrates, carbocation pathway I also might operate. Reactions with radical clocks showed that cyclopropane **7** and bisallylamine **9/10** underwent radical-induced ring rupture or 5-*exo*-atrig cyclization, respectively. In all cases, no direct alkene diazidation was observed. In principle, upon radical ring opening or cyclization, the resultant carbon-centered radical could be further oxidized to the cation before nucleophilic azidation; however, these carbocations, either highly electron-deficient or at a primary carbon, would be rather unstable. We also carried out diazidation of amide **13**, which—if operating via a carbocation mechanism—would cyclize to furnish oxazoline **15**.³⁹ However, electrolysis produced diazide **14** cleanly as the only observable product. Together, these data strongly support the radical mechanism.

We further studied alkene substrates that would lead to resonance-stabilized carbocation intermediates (Scheme 3C) and observed radical-to-carbocation oxidation. For example, camphor-derived alkene **16** was converted to 11% diazide **17** in addition to 42% 1,3-

azidoalcohol **18** via carbocation rearrangement. We reasoned that the norbornyl structure selectively stabilizes carbocation (**II**) but not its radical precursor (**I**), thus favoring the oxidation of **I** to **II**. Notably, neither rearrangement-azidation (19) or direct 1,2-azidohydroxylation products were observed. This result strongly supports the hypothesis that the alkene diazidation undergoes predominantly a radical pathway,^{17b,40} whereas the azidohydroxylation adopts a cationic pathway that triggers the rearrangement.⁴¹ Furthermore, 4-methoxy- α -methylstyrene (**1g**) was converted to diazide **2g** (major) and azidoalcohol **20** (minor). A key finding from these experiments is that if a carbocation is formed, its capture by H₂O is inevitable. Importantly, the majority of the substrates we investigated (see Table 2) produced no appreciable amounts of azidoalcohol products. This piece of information lends further support for the radical mechanism involving *N*-oxyl-mediated azidyl transfer.

Finally, we conducted a control experiment using azidoxy-generated products 3g or 3k instead of the alkene as the substrate for electrolysis and observed no diazide formation. This result excluded the possibility that *N*-oxyl adducts are key intermediates in the alkene diazidation (see Scheme 2B).

Accordingly, we propose a CHAMPO–N₃-mediated electrocatalytic cycle (Scheme 2C). We previously showed that *N*-oxyl radicals trigger an inner-sphere oxidation pathway that substantially lowers the potential required for accessing N₃[•] from N₃⁻²¹. In light of our new mechanistic information, we conclude that the CHAMPO–N₃ CTC plays two key roles: promoting the generation of free N[•] and mediating the second azidation through a putative radical group transfer event. Preliminary data suggest that the CTC is the resting state of the catalyst under a sufficient anodic potential. The first C–N₃ bond formation is slower than the second owing to the relatively slow, endergonic fragmentation of the CTC to CHAMPO and N₃[•]. This rate difference allows the two distinct radical addition events to take place successively with high chemoselectivity toward diazidation over side pathways that unproductively consumes **I**. In this mechanistic hypothesis, the metal-like behavior of *N*-oxyl radicals in the alkene diazidation is particularly intriguing. Our ongoing efforts will be devoted to further evaluating the mechanistic proposal and using the knowledge to guide the discovery of other synthetically useful transformations.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENTS

Financial support was provided by Cornell University and NIGMS (R01GM130928). S.L. thanks the ONR for a Young Investigator Award, which supports the synthesis of energetic materials. This study made use of the NMR facility supported by the NSF (CHE-1531632) and the ESR facility supported by NIGMS (P41GM103521). We thank IKA for the donation of ElectraSyn 2.0, Dr. Samantha MacMillan for help with X-ray crystallography analysis, Drs. Ivan Keresztes and Frank Schroeder for help with NMR spectral analysis, Dr. Timoth Chauvire for help with ESP data analysis, Yifan Shen for reproducing experimental results, and Dean Kim for experimental assistance.

REFERENCES

- (1). (a) Renaud P; Sibi MP Radicals in Organic Synthesis; Wiley-VCH: Weinheim, 2001. (b) Yan M; Lo JC; Edwards JT; Baran PS Radicals: Reactive Intermediates with Translational Potential. *J. Am. Chem. Soc* 2016, 138, 12692. [PubMed: 27631602] (c) Ischay MA; Yoon TP Accessing the Synthetic Chemistry of Radical Ions. *Eur. J. Org. Chem* 2012, 2012, 3359. (d) Moeller KM Using Physical Organic Chemistry To Shape the Course of Electrochemical Reactions. *Chem. Rev* 2018, 118, 4817. [PubMed: 29498518]
- (2). (a) Studer A; Curran DP Catalysis of Radical Reactions: A Radical Chemistry Perspective. *Angew. Chem., Int. Ed* 2016, 55, 58. (b) Skubi KL; Blum TR; Yoon TP Dual Catalysis Strategies in Photochemical Synthesis. *Chem. Rev* 2016, 116, 10035. [PubMed: 27109441] (c) Narayanam JMR; Stephenson CR Visible light photoredox catalysis: applications in organic synthesis. *Chem. Soc. Rev* 2011, 40, 102. [PubMed: 20532341] (d) Francke R; Little RD Redox catalysis in organic electrosynthesis: basic principles and recent developments. *Chem. Soc. Rev* 2014, 43, 2492. [PubMed: 24500279] (e) Yan M; Kawamata Y; Baran PS Synthetic Organic Electrochemical Methods Since 2000: On the Verge of a Renaissance. *Chem. Rev* 2017, 117, 13230. [PubMed: 28991454] (f) Crossley SWM; Martinez RM; Obradors C; Shenvi RA Mn-, Fe-, and Co-Catalyzed Radical Hydrofunctionalizations of Olefins. *Chem. Rev* 2016, 116, 8912. [PubMed: 27461578]
- (3). (a) Bobbitt JM; Brückner C; Merbouh N Oxoammonium and Nitroxide-Catalyzed Oxidation of Alcohols. *Org. React* 2009, 74, 103. (b) Wertz S; Studer A Nitroxide-catalyzed transition-metal-free aerobic oxidation processes. *Green Chem* 2013, 15, 3116. (c) Leadbeater NE; Bobbitt JM TEMPO-Derived Oxoammonium Salts as Versatile Oxidizing Agents. *Aldrichimica Acta* 2014, 47, 65.
- (4). Nutting JE; Rafiee MR; Stahl SS Tetramethylpiperidine *N*-Oxyl (TEMPO), Phthalimide *N*-Oxyl (PINO), and Related *N*-Oxyl Species: Electrochemical Properties and Their Use in Electrocatalytic Reactions. *Chem. Rev* 2018, 118, 4834. [PubMed: 29707945]
- (5). (a) Qian X-Y; Li S-Q; Song J; Xu H-C TEMPO-Catalyzed Electrochemical C–H Thiolation: Synthesis of Benzothiazoles and Thiazolopyridines from Thioamides. *ACS Catal* 2017, 7, 2730. (b) Peterson BM; Lin S; Fors BP Electrochemically Controlled Cationic Polymerization of Vinyl Ethers. *J. Am. Chem. Soc* 2018, 140, 2076. [PubMed: 29385348] (c) Furukawa K; Shibuya M; Yamamoto Y Chemoselective Catalytic Oxidation of 1,2-Diols to α -Hydroxy Acids Controlled by TEMPO–ClO₂ Charge-Transfer Complex. *Org. Lett* 2015, 17, 2282. [PubMed: 25886211]
- (6). For examples of stable TEMPO–X-type adducts formed via single-electron transfer, see: (a) Li Y.; Pouliot M; Vogler T; Renaud P; Studer A α -Aminooxylation of Ketones and β -Chloro- α -aminooxylation of Enones with TEMPO and Chlorocatecholborane. *Org. Lett* 2012, 14, 4474. [PubMed: 22900473] (b) Wang X; Ye Y; Zhang S; Feng J; Xu Y; Zhang Y; Wang J Copper-Catalyzed C(sp³)–C(sp³) Bond Formation Using a Hypervalent Iodine Reagent: An Efficient Allylic Trifluoromethylation. *J. Am. Chem. Soc* 2011, 133, 16410. [PubMed: 21936560]
- (7). TEMPO has been used (stoichiometrically or catalytically) in lieu of transition metals for formal C–C cross coupling. See: (a) Murarka S; Studer A Transition Metal-Free TEMPO-Catalyzed Oxidative Cross-Coupling of Nitrones with Alkynyl-Grignard Reagents. *Adv. Synth. Catal* 2011, 353, 2708. (b) Maji MS; Murarka S; Studer A Transition-Metal-Free Sonogashira-Type Coupling of ortho-Substituted Aryl and Alkynyl Grignard Reagents by Using 2,2,6,6-Tetramethylpiperidine-*N*-oxyl Radical as an Oxidant. *Org. Lett* 2010, 12, 3878. [PubMed: 20704186]
- (8). (a) For examples, see: Semmelhack MF; Chou CS; Cortes DA Nitroxyl-mediated electrooxidation of alcohols to aldehydes and ketones. *J. Am. Chem. Soc* 1983, 105, 4492. (b) Hoover J; Stahl S Highly Practical Copper(I)/TEMPO Catalyst System for Chemo-selective Aerobic Oxidation of Primary Alcohols. *J. Am. Chem. Soc* 2011, 133, 16901. [PubMed: 21861488] (c) Badalyan A; Stahl SS Cooperative electrocatalytic alcohol oxidation with electron-proton-transfer mediators. *Nature* 2016, 535, 406–410. [PubMed: 27350245] (d) Liu R; Liang X; Dong C; Hu X Transition-Metal-Free: A Highly Efficient Catalytic Aerobic Alcohol Oxidation Process. *J. Am. Chem. Soc* 2004, 126, 4112. [PubMed: 15053593] (e) Shibuya M; Osada Y; Sasano Y; Tomizawa M; Iwabuchi Y Highly Efficient, Organocatalytic Aerobic Alcohol Oxidation. *J. Am. Chem. Soc* 2011, 133, 6497. [PubMed: 21473575] (f) Shibuya M; Doi R; Shibuta T; Uesugi S; Iwabuchi Y

Organocatalytic One-Pot Oxidative Cleavage of Terminal Diols to Dehomologated Carboxylic Acids. *Org. Lett* 2012, 14, 5006. [PubMed: 22991924] (g)Hickey D; McCammant M; Giroud F; Sigman M; Minter S Hybrid Enzymatic and Organic Electrocatalytic Cascade for the Complete Oxidation of Glycerol. *J. Am. Chem. Soc* 2014, 136, 15917. [PubMed: 25350383] (h)Zhao M; Li J; Mano E; Song Z; Tschaen D; Grabowski E; Reider P Oxidation of Primary Alcohols to Carboxylic Acids with Sodium Chlorite Catalyzed by TEMPO and Bleach. *J. Org. Chem* 1999, 64, 2564. See also ref 5c. Stoichiometric oxidation: (i) Bailey WF; Bobbitt JM; Wiberg KB Mechanism of the Oxidation of Alcohols by Oxoammonium Cations. *J. Org. Chem* 2007, 72, 4504. [PubMed: 17488040] (j)Ganem B Biological spin labels as organic reagents. Oxidation of alcohols to carbonyl compounds using nitroxyls. *J. Org. Chem* 1975, 40, 1998. (k)Bobbitt JM Oxoammonium Salts. 4-Acetylamino-2,2,6,6-tetramethylpiperidine-1-oxoammonium Perchlorate: A Stable and Convenient Reagent for the Oxidation of Alcohols. *Silica Gel Catalysis. J. Org. Chem* 1998, 63, 9367.

- (9). For examples, see: (a) Miles K; Abrams M; Landis C; Stahl S KetoABNO/NO_x Cocatalytic Aerobic Oxidation of Aldehydes to Carboxylic Acids and Access to α -Chiral Carboxylic Acids via Sequential Asymmetric Hydroformylation/Oxidation. *Org. Lett* 2016, 18, 3590. [PubMed: 27410397] (b)Rafiee M; Konz Z; Graaf M; Koolman H; Stahl S Electrochemical Oxidation of Alcohols and Aldehydes to Carboxylic Acids Catalyzed by 4-Acetamido-TEMPO: An Alternative to “Anelli” and “Pinnick” Oxidations. *ACS Catal* 2018, 8, 6738.
- (10). For a review, see: (a) Bartelson A; Lambert K; Bobbitt J; Bailey W Recent Developments in the Nitroxide-Catalyzed Oxidation of Amines: Preparation of Imines and Nitriles. *ChemCatChem* 2016, 8, 3421–3430. For examples, see: (b) Lennox A; Goes S; Webster M; Koolman H; Djuric S; Stahl S Electrochemical Aminoxyl-Mediated α -Cyanation of Secondary Piperidines for Pharmaceutical Building Block Diversification. *J. Am. Chem. Soc* 2018, 140, 11227. [PubMed: 30141925] (c)Sammelhack MF; Schmid CR Nitroxyl-mediated electro-oxidation of amines to nitriles and carbonyl compounds. *J. Am. Chem. Soc* 1983, 105, 6732. (d)Kim J; Stahl S Cu/ Nitroxyl-Catalyzed Aerobic Oxidation of Primary Amines into Nitriles at Room Temperature. *ACS Catal* 2013, 3, 1652. [PubMed: 24015373] (e)Sonobe T; Oisaki K; Kanai M Catalytic aerobic production of imines en route to mild, green, and concise derivatizations of amines. *Chem. Sci* 2012, 3, 3249. (f)Wu Y; Yi H; Lei A Electrochemical Acceptorless Dehydrogenation of N-Heterocycles Utilizing TEMPO as Organo-Electrocatalyst. *ACS Catal* 2018, 8, 1192. (g)Romero NA; Margrey KA; Tay NE; Nicewicz DA Site-selective arene C-H amination via photoredox catalysis. *Science* 2015, 349, 1326. [PubMed: 26383949] (h)Somfai P; Yu L Synthesis of α -Keto Amides by a Pyrrolidine/TEMPO-Mediated Oxidation of α -Keto Amines. *Synlett* 2016, 27, 2587. (i)Wertz S; Studer A Metal-Free 2,2,6,6-Tetramethylpiperidin-1-yloxy Radical (TEMPO) Catalyzed Aerobic Oxidation of Hydroxylamines and Alkoxyamines to Oximes and Oxime Ethers. *Helv. Chim. Acta* 2012, 95, 1758.
- (11). (a)Tarantino KT; Miller DC; Callon TA; Knowles RR Bond-Weakening Catalysis: Conjugate Aminations Enabled by the Soft Homolysis of Strong N–H Bonds. *J. Am. Chem. Soc* 2015, 137, 6440. [PubMed: 25945955] See also refs 5a, b. Stoichiometric reaction: (b) Xu F; Zhu L; Zhu S; Yan X; Xu H-C Electrochemical Intramolecular Amino-oxygenation of Unactivated Alkenes. *Chem. - Eur. J* 2014, 20, 12740. [PubMed: 25145684]
- (12). Griesser M; Shah R; Van Kessel TA; Zilka O; Haidasz AE; Pratt AD The Catalytic Reaction of Nitroxides with Peroxyl Radicals and Its Relevance to Their Cytoprotective Properties. *J. Am. Chem. Soc* 2018, 140, 3798. [PubMed: 29451786]
- (13). Fu N; Sauer G; Saha A; Loo A; Lin S Metal-catalyzed electrochemical diazidation of alkenes. *Science* 2017, 357, 575. [PubMed: 28798126]
- (14). Minisci F Free-radical additions to olefins in the presence of redox systems. *Acc. Chem. Res* 1975, 8, 165.
- (15). (a)Magnus P; Roe MB; Hulme C New trialkylsilyl enol ether chemistry: direct 1,2-bis-azidation of triisopropylsilyl enol ethers: an azido-radical addition process promoted by TEMPO. *J. Chem. Soc., Chem. Commun* 1995, 263. (b)Magnus P; Lacour J; Evans PA; Roe MB; Hulme C Hypervalent Iodine Chemistry: New Oxidation Reactions Using the Iodosylbenzene–Trimethylsilyl Azide Reagent Combination. Direct α - and β -Azido Functionalization of Triisopropylsilyl Enol Ethers. *J. Am. Chem. Soc* 1996, 118, 3406.

- (16). Snider BB; Lin H An Improved Procedure for the Conversion of Alkenes and Glycols to 1,2-Diazides Using $\text{Mn}(\text{OAc})_3 \cdot 2\text{H}_2\text{O}$ in Acetonitrile Containing Trifluoroacetic Acid. *Synth. Commun* 1998, 28, 1913.
- (17). (a)Yuan YA; Lu DF; Chen YR; Xu H Iron-Catalyzed Direct Diazidation for a Broad Range of Olefins. *Angew. Chem., Int. Ed* 2016, 55, 534.(b)Shen S-J; Zhu C-L; Lu D-F; Xu H Iron-Catalyzed Direct Olefin Diazidation via Peroxyester Activation Promoted by Nitrogen-Based Ligands. *ACS Catal* 2018, 8, 4473. [PubMed: 29785320]
- (18). For examples, see: (a) Fumagalli G; Rabet PTG; Boyd S; Greaney MF Three-Component Azidation of Styrene-Type Double Bonds: Light-Switchable Behavior of a Copper Photoredox Catalyst. *Angew. Chem., Int. Ed* 2015, 54, 11481.(b)Lu M-Z; Wang C-Q; Loh T-P Copper-Catalyzed Vicinal Oxyazidation and Diazidation of Styrenes under Mild Conditions: Access to Alkyl Azides. *Org. Lett* 2015, 17, 6110. [PubMed: 26651361] (c)Peng H; Yuan Z; Chen P; Liu G Palladium-Catalyzed Intermolecular Oxidative Diazidation of Alkenes. *Chin. J. Chem* 2017, 35, 876.(d)Fristad WE; Brandvold TA; Peterson JR; Thompson SR Conversion of alkenes to 1,2-diazides and 1,2-diamines. *J. Org. Chem* 1985, 50, 3647.(e)Moriarty RM; Khosrowshahi JS A versatile synthesis of vicinal diazides using hypervalent iodine. *Tetrahedron Lett* 1986, 27, 2809.Related azidoamination: (f) Shen K; Wang Q Copper-Catalyzed Alkene Aminoazidation as a Rapid Entry to 1,2-Diamines and Installation of an Azide Reporter onto Azaheterocycles. *J. Am. Chem. Soc* 2017, 139, 13110. [PubMed: 28825822]
- (19). (a)Lucet D; Le Gall T; Mioskowski C The Chemistry of Vicinal Diamines. *Angew. Chem., Int. Ed* 1998, 37, 2580.(b)Chemler SR; Fuller PH Heterocycle synthesis by copper facilitated addition of heteroatoms to alkenes, alkynes and arenes. *Chem. Soc. Rev* 2007, 36, 1153. [PubMed: 17576482]
- (20). (a)Siu JC; Sauer GS; Saha A; Macey RL; Fu N; Chauvire T; Lancaster KL; Lin S Electrochemical Azidooxygenation of Alkenes Mediated by a TEMPO–N₃ Charge-Transfer Complex. *J. Am. Chem. Soc* 2018, 140, 12511. [PubMed: 30160949] For related seminal work on alkene azidooxygenation using a nonelectrochemical method, see: (b) Zhang B; Studer A Stereoselective Radical Azidooxygenation of Alkenes. *Org. Lett* 2013, 15, 4548. [PubMed: 23930944]
- (21). The potential for azidyl formation is lowered by ca. 200 mV. See ref 20a.
- (22). This corresponds to ca. 0.7 V initial anodic potential ($E_{a,i}$) vs Fc/Fc^+ (ferrocene/ferrocenium).
- (23). Studer A; Harms K; Knoop C; Müller C; Schulte T New Sterically Hindered Nitroxides for the Living Free Radical Polymerization: X-ray Structure of an α -H-Bearing Nitroxide. *Macromolecules* 2004, 37, 27.
- (24). Hickey D; Schiedler D; Matanovic I; Doan P; Atanasov P; Minter S; Sigman M Predicting Electrocatalytic Properties: Modeling Structure–Activity Relationships of Nitroxyl Radicals. *J. Am. Chem. Soc* 2015, 137, 16179. [PubMed: 26635089]
- (25). (a)Gerken J; Pang Y; Lauber M; Stahl S Structural Effects on the pH-Dependent Redox Properties of Organic Nitroxyls: Pourbaix Diagrams for TEMPO, ABNO, and Three TEMPO Analogs. *J. Org. Chem* 2018, 83, 7323. [PubMed: 29182282] (b)Rafiee M; Miles KC; Stahl SS Electrocatalytic Alcohol Oxidation with TEMPO and Bicyclic Nitroxyl Derivatives: Driving Force Trumps Steric Effects. *J. Am. Chem. Soc* 2015, 137, 14751. [PubMed: 26505317]
- (26). For a full list of catalysts surveyed, see SI.
- (27). (a)Ma Z; Huang Q; Bobbitt JM Oxoammonium salts. 5. A new synthesis of hindered piperidines leading to unsymmetrical TEMPO-type nitroxides. Synthesis and enantioselective oxidations with chiral nitroxides and chiral oxoammonium salts. *J. Org. Chem* 1993, 58, 4837.(b)Moad G; Rizzardo E; Solomon DH The reaction of acyl peroxides with 2,2,6,6-tetramethylpiperidinyl-1-oxy. *Tetrahedron Lett* 1981, 22, 1165.
- (28). This strategy has been employed in nitroxide-mediated polymerization (NMP) to reverse *N*-oxyl trapping and thus decrease the temperature needed to maintain chain growth. See ref 23.
- (29). In addition to increased steric encumbrance, CHAMPO also has fewer β -H's. The cyclohexane motifs could also stabilize the CHAMPO^+ ion through hyperconjugation. See ref 27a.
- (30). Haugland M; El-Sagheer A; Porter R; Peña J; Brown T; Anderson E; Lovett J 2'-Alkynyl nucleotides: A Sequence- and Spin Label-Flexible Strategy for EPR Spectroscopy in

DNA. J. Am. Chem. Soc 2016, 138, 9069 See also ref 26a. CHAMPO's $E_{1/2} = +0.29$ V. [PubMed: 27409454]

- (31). When the electrolysis was left continued for 12 h instead, the color remained persisted in the end of the reaction.
- (32). Schafer H Oxidative Addition of the Azide Ion to Olefins. A Simple Route to Diamines. Angew. Chem., Int. Ed. Engl 1970, 9, 158.
- (33). For instance, CAN gave < 10% diazide in addition to 42% azidoalcohol among other byproducts; K₂S₂O₇ or PhIO led to minimal conversion of the alkene.
- (34). Azidooxygenation byproducts were observed with TEMPO or ACT as the catalyst.
- (35). Workentin MS; Wagner BD; Luszyk J; Wayner DDM A Molecular Lock. J. Am. Chem. Soc 1995, 117, 119.
- (36). For bistriazole synthesis via dipolar cycloaddition, see SI; for dinitroalkane synthesis, see: Carmeli M; Rozen S Oxidation of Azides by the HOF·CH₃CN: A Novel Synthesis of Nitro Compounds. J. Org. Chem 2006, 71, 4585. [PubMed: 16749792]
- (37). (a)Yan M; Kawamata Y; Baran PS Synthetic Organic Electrochemistry: Calling All Engineers. Angew. Chem., Int. Ed 2018, 57, 4149.(b)Fu N; Sauer GS; Lin S A general, electrocatalytic approach to the synthesis of vicinal diamines. Nat. Protoc 2018, 13, 1725. [PubMed: 30072721]
- (38). (a)Zhu Q; Gentry EC; Knowles RR Catalytic Carbocation Generation Enabled by the Mesolytic Cleavage of Alkoxyamine Radical Cations. Angew. Chem., Int. Ed 2016, 55, 9969.(b)Gentry EC; Rono LJ; Hale ME; Matsuura R; Knowles RR Enantioselective Synthesis of Pyrroloindolines via Noncovalent Stabilization of Indole Radical Cations and Applications to the Synthesis of Alkaloid Natural Products. J. Am. Chem. Soc 2018, 140, 3394. [PubMed: 29432006] (c)O'Brya G; Braslau R Terminal Functionalization of Polymers via Single Electron Oxidation of N-Alkoxyamines. Macromolecules 2006, 39, 9010.
- (39). Xiong P; Long H; Song J; Wang Y; Li J-F; Xu H-CJ *Am. Chem. Soc.* Electrochemically Enabled Carbohydroxylation of Alkenes with H₂O and Organotrifluoroborates. J. Am. Chem. Soc 2018, 140, 16387.
- (40). Legnani L; Prina-Cerai G; Delcaillau T; Willems S; Morandi B Efficient access to unprotected primary amines by iron-catalyzed aminochlorination of alkenes. Science 2018, 362, 434. [PubMed: 30361368]
- (41). Legnani L; Morandi B Direct Catalytic Synthesis of Unprotected 2-Amino-1-Phenylethanols from Alkenes by Using Iron(II) Phthalocyanine. Angew. Chem., Int. Ed 2016, 55, 2248.

(A) Background: Electrochemical diazidation and azidoxygenation

For seminal developments in chemical diazidation, see refs. 14–18.

For seminal developments in chemical azidoxygenation, see ref. 20b.

(B) Working mechanistic hypothesis for aminoxyl ($NO\cdot$)-catalyzed alkene diazidation **Scheme 1.**

Mechanistic Hypothesis for *N*-Oxyl-Catalyzed Alkene Diazidation

(A) Cationic pathway I: Direct oxidation of carbon-centered radical I

(B) Cationic pathway II: Oxidation of the azidoxygenation product 3

(C) Radical pathway

Scheme 2.
Plausible Mechanistic Pathways

(A) Radical clocks: **(B) Carbocation trap:** **(C) Stabilized carbocations:** **(D) Mesolytic cleavage control:**

Scheme 3.
 Mechanistic Probes^a

Table 1.

Reaction Optimization^a

$\text{R}-\text{CH}=\text{CH}_2 \xrightarrow[\text{C}(+)/\text{Pt}(-), E_{\text{cell}} = 2.7 \text{ V } (E_{\text{a,i}} \approx 0.7 \text{ V}), \text{ T } (^\circ\text{C})]{\text{catalyst (10 mol\%), NaN}_3 \text{ (3.0 equiv)}} \text{R}-\text{CH}(\text{N}_3)-\text{CH}_2\text{N}_3$			
$\text{R}-\text{CH}(\text{N}_3)-\text{CH}_2\text{N}_3 + \text{N}(\text{R}')_2 \rightarrow \text{R}-\text{CH}(\text{O}-\text{N}(\text{R}')_2)-\text{CH}_2\text{N}_3$			
N-oxyl trapping byproduct 3			
Entry	Catalyst	T (°C)	Yield (%)
<div> 1a </div>			
1 ^b	TEMPO	22	36
2	TEMPO	22	11
3	TEMPO	0	46
4	TEMPO	-10	61
5	ACT	-10	71
6	CHAMPO	-10	97 ^c
7	none	-10	0
<div> 1b </div>			
8	TEMPO	-10	47
9	ACT	-10	68
10	CHAMPO	-10	87 ^c

^aConditions: **1** (0.2 mmol, 1 equiv), NaN₃ (3 equiv; 1.5 equiv per C–N bond), LiClO₄ (1.8 equiv), MeCN/H₂O (3.8 mL, 12:1); cell voltage (E_{cell}) = 2.7 V; yields determined by ¹H NMR. ^bWith 1 equiv TEMPO. ^cIsolated yield.

Table 2.

Substrate Scope^a

^aConditions: **1** (0.2 mmol, 1 equiv), CHAMPO (10 mol %), NaN₃ (3 equiv), LiClO₄ (1.8 equiv), MeCN/H₂O (3.8 mL, 12:1). ^b15 mol % CHAMPO. ^cLimonene-1,2-oxide (**1r**) was purchased as a pair of diastereomers at C1,2. ^dDetermined by ¹H NMR owing to the volatility of the product. ^eConditions: **2v**, PPh₃, H₂O/THF, reflux; then *p*TSA. ^fConditions: **2k**, PPh₃, H₂O/THF, reflux; then Ac₂O, NEt₃, DCM, 0 °C. ^gConditions: **2a**, H₂, Pd/C, EtOAc, Boc₂O.

entry	variation from “standard conditions”	yield (%)
1	none	>99
2	Ni(–) instead of Pt(–)	92
3	divided cell, constant current of 8 mA	81
4	ElectraSyn 2.0 as power source, 0.3 mmol 1k	82
5	AA batteries (×2) as power source	89
6	using TBAPF ₆ instead of LiClO ₄	95
7	no LiClO ₄ , $E_{\text{cell}} = 3.4 \text{ V}^b$	>99
8	1 mmol scale	69 ^c

^b A higher cell voltage is applied to compensate for the increased solution resistance between cathode and anode.

^c Isolated yield; reaction was carried out at a higher substrate concentration (see SI).