Phylogenetic diversity of sulfate-reducing bacteria of sediments of Chilika Lake, India, determined through analysis of the dissimilatory sulfite reductase (*dsrAB*) gene

Sri Sasi Jyothsna Tadinada^{1,3} · Rahul Kamidi^{1,4} · Saikat Dutta³ · Sasikala Chintalapati¹ · Venkata Ramana Chintalapati²

Received: 24 October 2018 / Accepted: 26 February 2019 / Published online: 7 March 2019
© King Abdulaziz City for Science and Technology 2019

Abstract

In this study, the sulfate-reducing bacteria, (SRB) were identified and reported for the first time through analysis of functional gene *dsrAB*, from the DNA of sediment samples collected from 10 sites of the Chilika lake. The finding illustrates Forty six Operational Taxonomic Units (OTUs), identified from the DGGE which were obtained from the 10 sediment samples. Of these, 34 OTUs exhibited around 78–96% sequence similarity and 12 OTUs showed 97 to 100% sequence similarity to the *dsrAB* gene of reported type strains of SRB. The sequence information obtained revealed the presence and distribution of diverse types of SRB which include phylotypes related to *Desulfovibrio*, *Desulfonatronovibrio*, *Desulfomicrobium*, *Desulfobotulos* and *Desulfobacca*. Upon comparison of *dsrAB* gene sequences of SRB obtained through this study with those collected from the GenBank, and through the dendrogram constructed, it was observed that except 13 OTUs that clustered closely with the reported type strains, all other 36 OTUs clustered distantly and had no representative member of SRB. This indicated the presence of phylogenetically diverse groups of SRB inhabiting the lake Chilika.

Keywords Sulfate-reducing bacteria · Sediments · *Desulfovibrio* · Dissimilatory sulfite reductase · *DsrAB* gene · Chilika Lagoon · Operational taxonomic units (OTU's)

Introduction

Approximately 70% of the Earth's environment is marine, which includes substantial sediment deposits (Fowler 1990). Though these sediments contain the largest global organic carbon reservoir, they have been considered relatively inactive (Jørgensen 1983), until the most recent studies conducted on direct measurement of microbial populations

using comprehensive microbial ecological techniques, which revealed the presence of extensive prokaryotic biosphere existing in marine sediments (Coolen et al. 2002; D'Hondt et al. 2004; Fröhlich-Nowoisky et al. 2016). Anoxic sediments are important sites of aquatic ecosystems as they play a crucial role in the decomposition of organic matter in these ecosystems. Due to the depleted oxygen levels, the decomposition of organic matter is channeled through fermentative, methanogenic or respiratory pathways involving reduced nitrate, ferric iron, manganese or sulfate as terminal electron acceptors (Henrichs and Reeburgh 1987; Throckmorton et al. 2015). The dissimilatory sulfate reduction accounts for the bulk of carbon mineralization in marine anoxic sediments (Crill and Martens 1987). A greater part in the formation of sulfidogenic sediments is played by the sulfate-reducing bacteria (SRB) in such ecosystems. These bacteria reduce sulfate to sulfide using a variety of electron donors like H₂, fatty acids, alcohols and aromatic compounds (Widdel and Bak 1992; Schink and Stams 2013). According to Bussmann and Reichardt (1991), the diversity of sulfate-reducing bacteria of surface waters is influenced

✉ Sri Sasi Jyothsna Tadinada
srisasijyothsna.t@ramky.com

¹ Bacterial Discovery Laboratory, Centre for Environment, Institute of Science and Technology, Jawaharlal Nehru Technological University, Hyderabad, India
² Department of Plant Sciences, School of Life Science, University of Hyderabad, Hyderabad, India
³ Environmental Consultancy, Ramky Enviro Services Private Limited, 12th Floor, Ramky Grandiose, Ramky Towers Complex, Gachibowli, Hyderabad, India
⁴ Central Sericultural Research and Training Institute, Central Silk Board, Berhampore, West Bengal, India

by the seasonal variations and hence fluctuates accordingly while that of sediments is more constant. Till today out of 188 validly published species of SRB, 31 species belonging to 19 different genera have been isolated from sediments (<http://www.bacterio.cict.fr/>) and uncultured sulfate-reducing bacteria have also been reported from sediments of various other regions of the world (Manabu and Susumu 1996). Hence sediments are known to harbor a variety of sulfate-reducing bacteria and thus an attempt was made to study the diversity of SRB from sediment samples collected from diverse regions of Chilika Lagoon, India.

Chilika is the largest lagoon along the coast of India situated between the latitude of 19°28' and 19°54'N and longitude of 85°05' and 85°38'E. The water spread area of the lagoon varies between 1165 and 906 Sq Km during the monsoon and summer, respectively (Mohanty and Panda 2009). There is an assemblage of freshwater, brackish and marine eco-system in the lagoon. It has very high productivity along with the presence of a variety of habitats allowing an amazing number of species. Reports on floral and faunal diversity of the lake Chilika, as reported by various researchers are available, but data pertaining to the microbial diversity of the same lagoon is scant. The lake also harbors rich black sulfide smelling sediments at various regions which drew our attention in studying the SRB responsible for the same.

As the classical methods of cultivation and identification of SRB upon purification are more time consuming and may result in the identification of cultured diversity alone, the present study focused on identification of uncultured SRB from sediment samples of Chilika Lake. The identification of SRB was based on the presence of *dsrAB* genes, encoding the α and β subunits of the enzyme dissimilatory sulfite reductase (*DSR*), which is an essential enzyme coded by almost all known SRB, as it catalyzes the final steps of sulfate reduction (Wanger et al. 1998). Using a combined process of PCR, nested PCR and denaturing gradient gel electrophoresis with SRB specific *dsrAB* gene targeting primers, the diversity of SRB in the sediment samples was identified, which is the first report of genetic diversity of SRB from Chilika Lake in general and from India, in particular. The results of the present study indicate the presence of diverse groups of SRB inhabiting the Lake Chilika, which are distantly related to the known and reported species of this group.

Materials and methods

Sample collection and processing for DNA extraction

Sediment samples were collected from various regions of the Chilika lake during December 2009 at global positioning

system coordinates 19°28' and 19°54'N and longitude of 85°05' and 85°38'E, respectively. Samples were collected in 25 mL sterile glass bottles with screw caps and stored at 4 °C until analysis. Van Veen grab sampler (Norinco make) was used for collection of sediments samples. All sediments were collected from a depth of 3–4 m. The pH and salinity of the sediment samples were analyzed before DNA extraction using pH electrode and conductivity meter (ADWA make). For this, 10 g. of each air dried sediment sample was suspended in 100 mL distilled water, and the pH, conductivity were read. Salinity was calculated from the electrical conductivity as described by (Hill et al. 1989; Eaton et al. 2005). The total genomic DNA was extracted from 2 g. of each sediment sample by bead beating method, using Soil DNA Isolation Kit Hi PurA MB542 (Hi Media) following the manufacturer's instructions.

PCR and nested PCR

The SRB specific functional gene *dsrAB* was amplified using the primer set *dsr1F* 5'-AC[C/G]CACTGGAAGCAC G-3' and *dsr4R* 5'-GTGTAGCAGTTACCGCA-3'. PCR components consisted of 15 μ L 2X red dye PCR master mix (150 mM Tris-HCl pH 8.5, 40 mM (NH₄)₂SO₄, 3.0 or 4.0 mM MgCl₂, 0.2% Tween 20, 0.4 mM dNTPs, 0.05 units/ μ L Ampliqon Taq DNA polymerase and Inert red dye and a stabilizer), 2 μ L bovine serum albumin (0.25 mg/mL), 2 μ L each of *dsr1F* and *dsr4R* primers (10 pmol/ μ L), 5 μ L 10X buffer and 4 μ L DNA (30–300 ng) making the final volume to 50 μ L using water (deoxyribonuclease free). Cycling parameters included an initial denaturation for 1 min at 94 °C followed by 34 cycles of 1 min at 94 °C, 1 min at 54 °C and 3 min at 72 °C concluding with a final extension for 10 min at 72 °C. The amplification resulted in a 1.9 kb product which was analyzed on a 1% agarose gel to ensure correct size. To obtain the *dsr* gene product suitable for running DGGE, a nested PCR was undertaken using a 1:10 diluted DNA template obtained from the initial *dsr* gene amplification. The primer set used for the nested PCR included *dsr1F* tagged with GC clamp (5'-CGCCCG CCG CGCCCGCGCCCGCCCGCCCGCCCGCCCG CCCC-3') in the 5' end (Muyzer et al. 1993) and *dsr5R* (5'-TGC CGA GGAGAACGATGT C-3') and the products were analyzed on a 2% agarose gel to ensure the correct size of the amplicon (243 bp) (Karr et al. 2005).

DGGE and sequencing

The 243 bp *dsrAB* amplicons were subjected to the denaturing gradient gel electrophoresis (DGGE) using the DCode universal mutation detection system (Bio-Rad,) according to the method described by Karr et al. (2005). The DGGE gels were stained using ethidium bromide (0.5 μ g/mL) and

visualized on a UV transilluminator. Bands were excised and eluted into sterile milliQ water and incubated for 30 min at 50 °C and finally kept on a shaker at room temperature overnight followed by storing at 4 °C before re-amplification. The eluted DNA was re-amplified using *dsrAB* primer set (*dsr1F* and *dsr5R*). The correct size of the final product of *dsrAB* gene amplicon (243 bp) was ensured on a 2% agarose gel. The gene sequences were obtained by cycle sequencing with the SequiTherm sequencing kit (Biozym) and the chain termination reaction (Sanger et al. 1977) using an automated laser fluorescence sequencer (Pharmacia) which was outsourced to Eurofins Bangalore, India.

Sequence analysis

Similarity to *dsrAB* sequences from other cultured and uncultured SRB was determined using BlastX function of BLAST (Altschul et al. 1990). The *dsrAB* gene sequences of Chilika lagoon sediments were manually aligned to the *dsrAB* sequences of known SRB obtained from GenBank. The CLUSTAL_W algorithm of MEGA 6 was used for sequence alignments and MEGA 6 (Tamura et al. 2013) software was used for phylogenetic analyses of the individual sequences. Distances were calculated using the Jukes and Cantor correction in a pair-wise deletion manner (Tamura et al. 2013). Neighbor-joining (NJ), minimum evolution (ME) and maximum parsimony (MP) methods in the MEGA6 software were used to construct phylogenetic trees. Percentage support values were obtained using a bootstrap procedure.

Nucleotide sequence accession numbers

The *dsrAB* gene sequences obtained from Chilika lake sediment samples were deposited in EMBL.

Results

Sample collection

Ten sediment samples were collected from Chilika lake which included nine ecologically important sites namely Barkul, Birds Island, Ghantasila Island, Heaven Island, Nalaban, Rambha and Satpada. Apart from these, sediments from the middle and interior sectors of the lake were also collected. The GPS coordinates of the samples collected are presented in Fig. 1. The pH and salinity particulars of the sample are given in Table 2. The salinity ranged from 1.8% – 5% and pH of the samples was observed to be neutral to moderately alkaline (7.0–8.0) Table 1. Also, each sample showed at least two morphologically different bacterial cell types under phase contrast microscope (Olympus B).

DNA Extraction, PCR, nested PCR and DGGE of *dsrAB* gene

DNA was extracted successfully from the sediment samples. Forty-six OTUs were generated from the DNA of 10 sediment samples through PCR, nested PCR and DGGE of the targeted functional gene *dsrAB* performed using *dsrAB* gene-specific primers. A minimum of 3 and a maximum of 8 OTUs were identified from each sample. The DNA obtained from sediment collected from the interior of Chilika lake yielded a maximum of 8 OTUs followed by Barkul and birds Island which yielded 6 OTUs each Table 2.

Sequence analysis and comparison

A phylogenetic tree constructed using MEGA 6 software to compare and understand the diversity of SRB based on the similarity of *dsrAB* sequences of Chilika lake with the other reported type strains of SRB obtained from the GenBank

Table 1 pH and salinity values of the sediment samples

No.	Place in Chilika (total no. of clones)	Sample pH	Sample salinity (%)	GPS of the sample
1	Interior of the Lake (8)	7.0	1.8	N 19°49' E 85°25'
2	Barkul (6)	8.0	4.9	N 19°37' E 85°11'
3	Birds Island (6)	8.0	1.7	N 19°27' E 85°08'
4	Satpada (5)	8.0	5.0	N 19°46' E 85°22'
5	Nalaban (5)	7.5	1.8	N 19°41' E 85°18'
6	Middle of Chilika Lake (4)	7.5	2.4	N 19°45' E 85°19'
7	Ghantasila Island (3)	8.0	1.8	N 19°31' E 85°06'
8	Rambha (3)	7.5	2.9	N 19°29' E 85°08'
9	Green pond at Satpada (3)	7.5	4.9	N 19°46' E 85°22'
10	Heaven Island (3)	8.0	3.7	N 19°40' E 85°12'

Fig. 1 Details of SRB obtained from each sampling site of Chilika lake with the GPS coordinates of the sampling sites

showed that majority of species belonged to non-spore forming Gram-negative SRB, except the one JSdsr 13.

Upon comparison among SRB sequences generated from the present study with those reported in the GenBank, twenty-six distinct clades were observed (Fig. 2). Thirteen deeply branching and distinct clades formed by a group of 1–4 OTUs each were observed within *Desulfovibrio* members, which contained no cultured representative species. Twelve other clades formed unclassified groups of SRB. Twenty-three deeply branching OTUs were observed which were quite distinct and had no representative genus with which the sequences were closely related.

Twenty-eight out of 46 OTUs clustered with the genus *Desulfovibrio* while OTUs close to other genera like *Desulfatiferula*, *Desulfobacca*, *Desulfobacter*, *Desulfobotulus*, *Desulfomicrobium*, *Desulfonatronovibrio*, *Desulfonatronum*, and *Desulfonema* were also obtained. A majority (34) of the sequences exhibited around 78–95% sequence similarity and a few (12) up to 100% sequence similarity to the *dsrAB* gene sequence of reported type strains of SRB.

Desulfovibrio was one such genus which was identified in all the sampled sediments. Only 18 out of 46 OTUs were close to the members other than the genus *Desulfovibrio*. Among these 18 OTUs, diverse phylotypes of SRB were identified belonging to 9 different genera without repeats which made the interpretation of results more complex. All the Chilika lake OTUs from the present study represented the SRB of δ -proteobacteria alone which included non-spore forming Gram-negative type of bacteria except

JSdsr13 which was close to the spore-forming sulfate-reducing bacterial genus *Desulfosporosinus* of the phylum Firmicutes. JSdsr 18 and JSdsr 19 were two such OTUs which did not have any representative strain of SRB reported in the GenBank. Each sediment sample was unique in the type of SRB inhabiting the sample except the common *Desulfovibrio* members as the dominant SRB. The 46 OTUs have been deposited in the EMBL. Accession numbers: FN811933–FN811935, FN811939–FN811941, FR694655–FR694674 and FR694697–FR694716.

Discussion

The sulfate reduction was prominent in sediments of Chilika lake, which were black and typically sulfide smelling. According to Barton (1995), the presence of sulfate reducers with high metabolic activity is easily revealed by the blackening of water and sediment due to precipitation of iron sulfide, and by the smell of hydrogen sulfide. However, no studies were performed in determining the sulfate-reducing bacteria responsible for the biological sulfate reduction of the lake, till today. Our previous studies on the cultured diversity of SRB of Chilika lake through 16S rRNA gene analysis have indicated the presence of strains of SRB belonging to the genus *Desulfovibrio* (Unpublished data). This led us to study the diversity of SRB of the Chilika lagoon. As most of the cultured diversity studies of SRB carried out by us resulted in the

Table 2 Phylogenetic analysis (based on the *dsrAB* gene) of enrichment cultures of SRB obtained from sediment samples of the lake Chilika, India

Place of sampling in Chilika Lake	<i>dsrAB</i> gene OTU obtained	EMBL Accession no. of <i>dsrAB</i> gene sequence	Nearest species of SRB, its place of isolation and % similarity based on <i>dsrAB</i> gene	Accession no of <i>dsrAB</i> gene sequence of the nearest SRB	Nearest type strain of SRB & (% similarity)	Accession no of <i>dsrAB</i> gene sequence of the type strain of SRB
Interior of lake	JSdsr 10	FN811933	<i>Desulfomicrobium baculum</i> DSM4028 Japan (94)	AB061530	<i>Desulfomicrobium baculum</i> DSM4028 Japan (94)	AB061530
	JSdsr 11	FN811934	Uncultured SRB E-P21 USA (100)	DQ404316	<i>Desulfovibrio desulfuricans</i> SRDQC China (98)	DQ450464
	JSdsr 31	FR694698	Uncultured SRB clone:C6 Japan (98)	AB263677	<i>Desulfobacterium cetonicum</i> DSM7267 Germany (83)	AF420282
	JSdsr 32	FR694699	—do—	—do—	<i>Desulfohalobium retbaense</i> DSM 5692 USA (78)	CP001734
	JSdsr 33	FR694700	Uncultured SRB clone LFsdsrC20 Antarctica (94)	AY273284	<i>Desulfovibrio burkinensis</i> DSM 6830 Germany (87)	AF418186
	JSdsr 46	FR694713	Uncultured SRB clone JSdsr7 India (100)	FN811939	<i>Desulfovibrio portus</i> MSL79 Japan. (86)	AB373125
	JSdsr 47	FR694714	Uncultured SRB clone RH.dsra-07-46 Israel (90)	FJ231254	<i>Desulfovibrio intestinalis</i> DSM 11,275 Germany (87)	AF418183
	JSdsr 49	FR694717	Uncultured SRB clone PIMO3C12 England (96)	AY741572	<i>Desulfobacterium oleovorans</i> DSM6200 Germany. (81)	AF418201

Table 2 (continued)

Place of sampling in Chikilika Lake	<i>dsrAB</i> gene OTU obtained	EMBL Accession no. of <i>dsrAB</i> gene sequence	Nearest species of SRB, its place of isolation and % similarity based on <i>dsrAB</i> gene	Accession no of <i>dsrAB</i> gene sequence of the nearest SRB	Nearest type strain of SRB & (% similarity)	Accession no of <i>dsrAB</i> gene sequence of the type strain of SRB
Barkul	JSdsr 3	FN811933	Uncultured SRB clone LGWK05 China (97)	EF065079	<i>Desulfovibrio burkinensis</i> DSM6830 Germany (88)	AF269147
	JSdsr 9	FN811941	Uncultured SRB clone RH.dsrA-07-5 Israel (96)	F1231220	<i>Desulfovibrio portus</i> MSL79 Japan (88)	AB373125
	JSdsr 34	FR694701	Uncultured SRB clone LFsdsrC20 Antarctica (95)	AY273284	<i>Desulfovibrio magneticus</i> RS-1 Japan. (84)	AP010904
	JSdsr 35	FR694702	—do—	—do—	<i>Desulfovibrio desulfuricans</i> G20 USA (83)	CP000112
	JSdsr 36	FR694703	Uncultured SRB clone LGWK05 China (81)	EF065079	<i>Desulfovibrio piger</i> ATCC29098 Germany (85)	AF482462
Birds island	JSdsr 48	FR694715	Uncultured SRB clone RH.dsrA-07-42 Israel (92)	F1231251	<i>Desulfovibrio portus</i> MSL79 China (91)	AB373125
	JSdsr 12	FR694657	<i>Desulfovibrio aminophilus</i> strain DSM 12,254, Austria (100)	AY626029	<i>Desulfovibrio aminophilus</i> strain DSM 12,254, Austria (100)	AY626029
	JSdsr 13	FR694658	<i>Desulfosporosinus orientis</i> DSM765 Germany (96)	AF271767	<i>Desulfosporosinus orientis</i> DSM765 Germany (96)	AF271767
	JSdsr 22	FR694667	<i>Desulfovibrio desulfuricans</i> ATCC27774 USA (93)	CP001358	<i>Desulfovibrio desulfuricans</i> ATCC27774 USA (93)	CP001358
	JSdsr 23	FR694668	<i>Desulfomicrobium</i> sp. MSL98 Japan (100)	AB373131	<i>Desulfovibrio aminophilus</i> DSM12254 Austria (88)	AY626029
	JSdsr 42	FR694709	Uncultured SRB clone 3 UK. (93)	GU181331	<i>Desulfovibrio vulgaris</i> DP4 USA (90)	CP000527
	JSdsr 43	FR694710	Uncultured SRB clone Mont23F (92)	EU189177	<i>Desulfovibrio intestinalis</i> DSM 11,275 Germany (89)	AF418183

Table 2 (continued)

Place of sampling in Chikilika Lake	<i>dsrAB</i> gene OTU obtained	EMBL Accession no. of <i>dsrAB</i> gene sequence	Nearest species of SRB, its place of isolation and % similarity based on <i>dsrAB</i> gene	Accession no of <i>dsrAB</i> gene sequence of the nearest SRB	Nearest type strain of SRB & (% similarity)	Accession no of <i>dsrAB</i> gene sequence of the type strain of SRB
Satpada	JSdsr 1	FN811933	Uncultured SRB clone LGWK05 China (90)	EF065079	<i>Bilophila wadsworthia</i> ATCC9260 Germany (87)	AF269147
	JSdsr 2	FN811934	—do—	—do—	—do—	—do—
	JSdsr 37	FR694704	<i>Desulfovibrio</i> sp. LVS-21 Hungary (92)	AM494497	<i>Bilophila wadsworthia</i> ATCC9260 Germany (91)	AF269147
	JSdsr 38	FR694705	Uncultured SRB clone JSdsr8 India (79)	FN811940	<i>Desulfovibrio aerotolerans</i> Dv05 Denmark (78)	AY749039
	JSdsr 39	FR694706	<i>Desulfovibrio</i> sp. LVS-15 Hungary (95)	AM494496	<i>Desulfovibrio desulfuricans</i> DSM642 Germany (94)	AF273034
Nalaban	JSdsr 8	FN811940	Uncultured SRB clone JSdsr8 India (100)	FN811940	<i>Desulfovibrio desulfuricans</i> DSM642 Germany (95)	AF273034
	JSdsr 26	FR694671	<i>Desulfatiferula olefinivorans</i> LM2801 France (100)	DQ826725	<i>Desulfatiferula olefinivorans</i> LM2801 France (100)	DQ826725
	JSdsr 27	FR694672	<i>Desulfonatronum lacustre</i> DSM10312 Germany (100)	AF418189	<i>Desulfonatronum lacustre</i> DSM10312 Germany (100)	AF418189
	JSdsr 28	FR694673	Uncultured SRB clone BV16F Israel (98)	EU552485	<i>Desulfovibrio carbinolicus</i> strain DSM3852 Austria (90)	AY626026
	JSdsr 45	FR694712	Uncultured SRB JSdsr9 India (91)	FN811941	<i>Desulfovibrio portus</i> MSL79 Japan (86)	AB373125

Table 2 (continued)

Place of sampling in Chilika Lake	<i>dsrAB</i> gene OTU obtained	EMBL Accession no. of <i>dsrAB</i> gene sequence	Nearest species of SRB, its place of isolation and % similarity based on <i>dsrAB</i> gene	Accession no of <i>dsrAB</i> gene sequence of the nearest SRB	Nearest type strain of SRB & (% similarity)	Accession no of <i>dsrAB</i> gene sequence of the type strain of SRB
Middle of Chilika lake	JSdsr 18	FR694663	Uncultured SRB clone OTU-67 China (90)	EU258889	Data unavailable	–
	JSdsr 19	FR694664	Uncultured SRB clone OTU-63 China (100)	EU258879	Data unavailable	–
	JSdsr 20	FR694665	<i>Desulfobacca acetoxidans</i> DSM1109 Germany (98)	AY167463	<i>Desulfobacca acetoxidans</i> DSM1109 Germany (98)	AY167463
	JSdsr 21	FR694666	Uncultured SRB clone LFsdsrC4 Antarctica (93)	AY273268	Data unavailable	–
Ghantasila Island	JSdsr 7	FN811938	Uncultured SRB clone JSdsr6 India (100)	FN811938	<i>Desulfovibrio salexigens</i> DSM2638 USA (96)	CP001649
	JSdsr 29	FR694674	<i>Desulfovibrio longus</i> DSM6739 Japan (95)	AB061540	<i>Desulfovibrio longus</i> DSM6739 Japan (95)	AB061540
	JSdsr 30	FR694675	Uncultured SRB clone:O12 C6 UK (98)	AB263701	<i>Desulfococcus oleovorans</i> Hxd3 USA (82)	CP000859
	JSdsr 24	FR694669	<i>Desulfonatronovibrio hydrogenvorans</i> DSM9292 Germany (96)	AF418197	<i>Desulfonatronovibrio hydrogenvorans</i> DSM9292 Germany (96)	AF418197
Rambha	JSdsr 25	FR694670	<i>Desulfovibrio</i> sp. TBP-1U USA (94)	AF327307	<i>Desulfovibrio acrylicus</i> DSM10141 Denmark (91)	FJ655912
	JSdsr 44	FR694711	<i>Desulfovibrio</i> sp. LVS-15 Hungary (97)	AM494496	<i>Desulfovibrio desulfuricans</i> DSM642 Germany (91)	AF273034
	JSdsr 16	FR694661	<i>Desulfonauticus submarinus</i> DSM15269 France (100)	AY187624	<i>Desulfonauticus submarinus</i> DSM15269 France (100)	AY187624
	JSdsr 17	FR694662	<i>Desulfospira joergensenii</i> DSM10085 Germany (95)	AF551760	<i>Desulfospira joergensenii</i> DSM10085 Germany (95)	AF551760
Green pond at satpada	JSdsr 41	FR694707	<i>Desulfovibrio</i> sp. LVS-15 Hungary (96)	AM494496	<i>Desulfovibrio desulfuricans</i> DSM642 Germany (95)	AF273034

Table 2 (continued)

Place of sampling in Chilika Lake	<i>dsrAB</i> gene OTU obtained	EMBL Accession no. of <i>dsrAB</i> gene sequence	Nearest species of SRB, its place of isolation and % similarity based on <i>dsrAB</i> gene	Accession no of <i>dsrAB</i> gene sequence of the nearest SRB	Nearest type strain of SRB & (% similarity)	Accession no of <i>dsrAB</i> gene sequence of the type strain of SRB
Heaven Island	JSdsr 14	FR694659	<i>Desulfobacca acetoxidans</i> DSM11109 Germany (97)	AY167463	<i>Desulfobacca acetoxidans</i> DSM11109 Germany (97)	AY167463
	JSdsr 15	FR694660	<i>Desulforhabdus amnigena</i> DSM10338 USA (100)	AF337901	<i>Desulforhabdus amnigena</i> DSM10338 USA (100)	AF337901
	JSdsr 40	FR694707	<i>Desulfovibrio aminophilus</i> DSM12254 Austria (100)	AY626029	<i>Desulfovibrio aminophilus</i> strain DSM12254 Austria	AY626029

identification of strains of *Desulfovibrionaceae* alone; the present study was targeted to study the genetic diversity of SRB of Chilika lake sediments. Hence the functional gene *dsrAB* which codes for the enzyme dissimilatory sulfite reductase (DSR) was chosen as the marker gene of SRB in the sediment samples.

So far only a limited number of sulfate reducers from subsurface sediments have been reported. Most of these belonged to the deltaproteobacterial genera *Desulfovibrio* and *Desulfomicrobium* or the spore-forming genera *Desulfosporosinus* or *Desulfotomaculum* within *Firmicutes* (Barnes et al. 1998; Köpke et al. 2005). However, whereas these investigators could identify SRB which were related exclusively to the *Desulfovibrio*, the sulfate reducers identified in our study were more diverse. The phylogenetic analysis revealed the presence of diverse groups of SRB inhabiting the sediments of the lake Chilika. However, some SRB OTUs showed similarities to deeply branching clades containing no reported representatives, indicating the presence of novel lineages of SRB.

Presence of the members of *Desulfovibrio* as majority in the Chilika Lake indicates the presence of oxic conditions or sulfur-oxidizing chemotrophs in or near the sediments as the members of *Desulfovibrio* have a higher degree of oxygen tolerance conditions than other SRB (Stahl et al. 2002). The members of *Desulfococcus*, *Desulfonema*, *Desulfosarcina* and *Desulfobacter* have also been identified, which were also reported in the deep sediments of methane hydrate containing marine sediments of Cascadia Margin (Bidle et al. 1999) and in the microbial mats located at the oxic-anoxic interphase (Teske et al. 1998). This study detected the presence of members of SRB reported to incompletely oxidize the organic compounds (lactate, propionate, ethanol etc.) producing acetate like *Desulfovibrio* and *Desulfomicrobium* out-competing the members which can completely oxidize lactate and other organic compounds (short chain fatty acids and alcohols) to CO₂ like *Desulfococcus*, *Desulfobacterium*, *Desulfosarcina* and *Desulfonema* (Hansen 1993). The dominance of members of *Desulfovibrionaceae* complies with the fact that their members are more dominant within marine sediments, mud flats and profundal lake sediments (Llobet-Brossa et al. 2002) rather than rhizosphere sediments of salt marsh. On the other hand, presence of members of *Desulfobacteraceae* is in agreement with the dominance of these members in the salt marsh sediments (Edgcomb et al. 1999; Hines et al. 1999) and coastal marine sediments (Sahm et al. 1999). SRB sequences related to the Gram-positive and thermophilic SRB were not detected in the sediment samples of Chilika indicating the absence of moderate thermophilic temperatures that could probably support the growth of moderate thermophiles like *Desulfotomaculum* with growth temperatures of 50–65 °C (Tanimoto and Bak 1994) which is too high for the Chilika lake sediments.

Fig. 2 Phylogenetic tree showing the relationships between selected strains of sulfate-reducing bacteria obtained from the GenBank and those obtained from Chilika Lake sediments based on the *dsrAB* gene. The dendrogram is constructed using the CLUSTAL_W algorithm of MEGA 6 and distance calculated using Jukes-Cantor, and

Neighbor-joining method. The *dsr AB* gene sequence accession numbers for the type strains and for the SRB clones obtained from Chilika Lake are given in parentheses. Bootstraps values (>95%) based on thousand replications are shown at the nodes of the tree. Bar 5 substitutions per 100 nucleotides

In conclusion, the present study on identification of SRB from sediment samples of Chilika Lake based on molecular approaches reveals the presence and distribution of diverse types of SRB in the lake sediments. The prevalence of

marine, brackish and freshwater types of habitats in a single ecosystem of Chilika Lake would have contributed to the physiologically and phylogenetically diverse groups of SRB inhabiting the lake. Further investigations of bacterial

communities including the phototrophic anoxygenic bacteria (Sucharita et al. 2010) and sulfur-oxidizing chemotrophs that go hand in hand in maintaining the sulfur cycle of the ecosystem is also important and would give a better understanding of the lake ecosystem. Apart from the sediment samples, diversity of SRB from various depths would also help to know the oxygen penetration and sulfide production rates that can indirectly present a picture of sulfur cycling of the lake.

Acknowledgements TS Sasi Jyothsna thanks CSIR for the award of research fellowship. Financial assistance received from DBT and DST (FIST) is acknowledged. We thank Ms. Azmatunnisa for the technical support. Also, I would like to extend my gratitude to Dr. B. Chakradhar, Director, RESPL, for critically reviewing the manuscript and serving as scientific advisor.

Compliance with ethical standards

Conflict of interest The authors state no conflict of interest for this work.

References

- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local alignment search tool. *J Mol Biol* 215:403–410. [https://doi.org/10.1016/S0022-2836\(05\)80360-2](https://doi.org/10.1016/S0022-2836(05)80360-2)
- Barnes SP, Bradbrook SD, Cragg BA (1998) Isolation of sulfate reducing bacteria from deep sea sediment layers of Pacific Ocean. *Geomicrobiol J* 15:67–83. <https://doi.org/10.1080/01490459809378066>
- Barton LL (1995) Sulfate reducing bacteria. In: Atkinson T, Sherwood RF Biotechnology handbooks-8. Series Kluwer Academic/ Plenum Publishers, New York
- Bidle KA, Kastner M, Bartlett DH (1999) A phylogenetic analysis of microbial communities associated with the methane hydrate containing marine fluids and sediments in the Cascadia Margin (ODP site 892B). *FEMS Microbiol Lett* 177:101–108. <https://doi.org/10.1111/j.1574-6968.1999.tb13719.x>
- Bussmann I, Reichardt W (1991) Sulfate-reducing bacteria in temporarily oxic sediments with bivalves. *Mar Ecol Prog Ser* 78:97–102
- Coolen MJL, Cypionka H, Sass AM, Sass, Overmann J (2002) Ongoing modification of Mediterranean Pleistocene sapropels mediated by prokaryotes. *Science* 296:2407–2410. <https://doi.org/10.1126/science.1071893>
- Crill PM, Martens CS (1987) Biogeochemical cycling in an organic rich coastal marine basin. Temporal and spatial variations in sulfate reduction rates. *Geochim Cosmochim Acta* 51:1175–1186. [https://doi.org/10.1016/0016-7037\(87\)90210-9](https://doi.org/10.1016/0016-7037(87)90210-9)
- D'Hondt S, Jørgensen BB, Miller DJ (2004) Distribution of microbial activity in deep sub seafloor sediments. *Science* 306:2216–2221. <https://doi.org/10.1126/science.1101155>
- Eaton AD, Clesceri LS, Greenberg AE, Franson MAH (2005) Standard methods for the examination of water and wastewater. *Am Public Health Assoc* 21:1600
- Edgcomb VP, McDonald JH, Devereux R, Smith DW (1999) Estimation of bacterial cell numbers in humic-rich salt marsh sediments using probes to 16S rDNA. *Appl Environ Microbiol* 164:271–279
- Fowler CMR (1990) The solid earth, an introduction to global geophysics. Cambridge University Press, Cambridge
- Fröhlich-Nowoisky J, Kampf CJ, Weber B, Huffman JA, Pöhlker C, Andreae MO, Su H (2016) Bioaerosols in the Earth system: Climate, health, and ecosystem interactions. *Atmos Res* 182:346–376. <https://doi.org/10.1038/ismej.2016.45>
- Hansen TA (1993) Carbon metabolism of sulfate reducing bacteria. In: Odum JM, Singleton R (eds) The sulfate reducing bacteria. Springer Verlag, New York, pp 21–40
- Henrichs SM, Reeburgh WS (1987) Anaerobic mineralization of marine sediment organic matter: rates and the role of anaerobic processes in the oceanic carbon economy. *Geo Microbiol J* 5:191–237. <https://doi.org/10.1080/01490458709385971>
- Hill KD, Dauphinee TM, Woods DJ (1989) The uniqueness of the practical salinity scale (1978): testing the scale with natural seawaters. *IEEE J Oceanic Eng* 14:265–267. <https://doi.org/10.1109/48.29606>
- Hines ME, Evans RS, Genthner BRS, Willis SG, Friedman S, Rooney-Verga JN, Devereux R (1999) Molecular phylogenetic and biogeochemical studies of Sulfate reducing bacteria in the rhizosphere of *Spartina alterniflora*. *Appl Environ Microbiol* 65:2209–2216
- Jørgensen BB (1983) Processes at the sediment water interface. In: Bolin B, Cook RB (eds) The major biochemical cycles and their interactions. Wiley, Chichester, pp 447–515
- Karr EA, Sattley WM, Rice MR, Jung DO, Madigan MT, Achenbach LA (2005) Diversity and distribution of sulfate-reducing bacteria in permanently frozen Lake Fryxell McMurdo Dry Valleys Antarctica. *Appl Environ Microbiol* 71:6353–6359. <https://doi.org/10.1128/AEM.71.10.6353-6359.2005>
- Köpke B, Wilms R, Engelen B, Cypionka H, Sass H (2005) Microbial diversity in coastal subsurface sediments: a cultivation approach using various electron acceptors and substrate gradients. *Appl Environ Microbiol* 71:7819–7830. <https://doi.org/10.1128/AEM.71.12.7819-7830.2005>
- Llobet-Brossa E, Rabus R, Bötcher ME, Könneke M, Finke N, Schramm A (2002) Community structure and activity of sulfate reducing bacteria in an inter tidal surface sediment: a multi-method approach. *Aquat Microbiol Ecol* 29:221–226. <https://doi.org/10.3354/ame029211>
- Manabu F, Susumu T (1996) Microdistribution of sulfate-reducing bacteria in sediments of a hypertrophic lake and their response to the addition of organic matter. *Ecol Res* 11:257–267
- Mohanty PK, Panda B (2009) Circulation & mixing processes in Chilika Lagoon. *Indian J Mar Sci* 38:205–214. <http://nopr.niscair.res.in/handle/123456789/4671>
- Muyzer G, De Waal EC, Uitterlinden AG (1993) Profiling of complex microbial populations by denaturing gradient gel electrophoresis analysis of polymerase chain reaction-amplified genes coding for 16S rRNA. *Appl Environ Microbiol* 59:695–700
- Sahm K, MacGregor BJ, Jørgensen B, Stahl DA (1999) Sulfate reduction and vertical distribution of sulfate reducing bacteria quantified by rRNA slot-blot hybridization in a coastal marine sediment. *Environ Microbiol* 1:56–74. <https://doi.org/10.1046/j.1462-2920.1999.00007.x>
- Sanger F, Nicklen S, Coulson AR (1977) DNA sequencing with chain terminating inhibitors. *Proc Natl Acad Sci USA* 74:5463–5467. <https://doi.org/10.1073/pnas.74.12.5463>
- Schink B, Stams AJ (2013) Syntrophism among prokaryotes. In: Rosenberg E, DeLong EF, Lory S, Stackebrandt E, Thompson T (eds) The Prokaryotes-Prokaryotic Communities and Ecophysiology. Springer, Berlin, pp 471–493
- Stahl DA, Fishbain S, Klein M, Baker BJ, Wanger W (2002) Origins and diversification of Sulfate-respiring microorganisms. *Antonie Leeuwenhoek* 81:189–195
- Sucharita K, Sasikala C, Ramana CV (2010) *Thiorhodococcus modestalkaliphilus* sp. nov. a phototrophic gamma proteobacterium from Chilika salt water lagoon, India. *J Gen Appl Microbiol* 56:93–99. <https://doi.org/10.2323/jgam.56.93>

- Tamura K, Stecher G, Peterson D, Filipski A, Kumar S (2013) MEGA6: molecular evolutionary genetics analysis version 6.0. *Mol Biol Evol* 30:2725–2729. <https://doi.org/10.1093/molbev/mst197>
- Tanimoto Y, Bak F (1994) Anaerobic degradation of methylmercaptan and dimethyl sulfide by newly isolated thermophilic sulfate reducing bacteria. *Appl Environ Microbiol* 60:2450–2455
- Teske A, Ramsingh NB, Habicht K, Fukui M, Kuver J, Jorgensen BB, Cohen Y (1998) Sulfate reducing bacteria and their activities in cyanobacterial mats of Solar lake (Sinai, Egypt). *Appl Environ Microbiol* 64:2943–2951
- Throckmorton HM, Heikoop JM, Newman BD, Altmann GL, Conrad MS, Muss JD, Wilson CJ (2015) Pathways and transformations of dissolved methane and dissolved inorganic carbon in Arctic tundra watersheds: Evidence from analysis of stable isotopes. *Glob Biogeochem Cycles* 29:1893–1910. <https://doi.org/10.1002/2014GB005044>
- Wanger M, Roger AJ, Flax JL, Brusseau GA, Stahl DA (1998) Phylogeny of dissimilatory sulfite reductases supports an early origin of sulfate respiration. *J Bacteriol* 180:2975–2982
- Widdel F, Bak F (1992) Gram-negative mesophilic sulfate-reducing bacteria. In: Balows A, TrIn Balows AHG, Trüper HG, Dworkin M, Harder W, Schleifer KH (eds) *The prokaryotes*, 2nd edn. Springer, New York, pp 3352–3378. https://doi.org/10.1007/978-1-4757-2191-1_21