Published in final edited form as:

Int J Hyg Environ Health. 2019 March ; 222(2): 168–176. doi:10.1016/j.ijheh.2018.09.001.

Five-year Exposure to PM_{2.5} and Ozone and Subclinical Atherosclerosis in Late Midlife Women: the Study of Women's Health Across the Nation

Chunzhe Duan^{1,†}, Evelyn Talbott¹, Maria Brooks¹, Sung Kyun Park², Rachel Broadwin³, Karen Matthews^{1,4}, and Emma Barinas-Mitchell^{1,*}

¹Department of Epidemiology, University of Pittsburgh, Graduate School of Public Health

²California Office of Environmental Health Hazard Assessment

³Department of Epidemiology and Department of Environmental Health Science, University of Michigan, School of Public Health

⁴Department of Psychiatry, University of Pittsburgh, School of Medicine

Abstract

Introduction: Effects of more than one-year exposure to air pollution on atherosclerosis is seldom studied. This paper aims to examine the association between five-year exposure to particulate matter 2.5 μm (PM_{2.5}), ozone (O₃) and atherosclerosis observed about seven years later in late midlife women.

Material and Methods: This study was conducted among 1188 women of the Study of Women's Health Across the Nation (SWAN) from five sites, Detroit, MI; Oakland, CA; Pittsburgh, PA; Chicago, IL; and Newark, NJ, with available data on both air pollutant exposure and carotid ultrasound scans. Five-year mean annualized exposure levels of two air pollutants, PM_{2.5} and ozone (O₃), were collected during 5 SWAN visits (1999-2005) from monitors 20 km within the participant's residential address. Linear regression models were used to estimate the association of prior five-year mean annualized exposure to PM_{2.5} and O₃ with common carotid intima-media thickness (cIMT) and inter-adventitial diameter (IAD) examined approximately seven years later (2009-2013). Logistic and multinomial logistic regressions were applied to assess the associations of air pollutants with plaque presence and plaque index, respectively.

*Corresponding author: Emma Barinas-Mitchell, 130 N. Bellefield Avenue, Suite 531, University of Pittsburgh, Pittsburgh, PA 15213, Phone: 412-624-3478, ejb4@pitt.edu.

[†]Present Address: H. Lee Moffitt Cancer Center, 12902 USF Magnolia Drive Tampa, FL 33612

Data access:

SWAN provides access to public use datasets that extend through the tenth annual follow-up visit. Some, but not all, of the data used for this manuscript are contained in the public use data sets. Links to each of the public use data sets are located on the SWAN web site: <http://www.swanstudy.org/swan-research/data-access/>. Investigators who require assistance accessing the public use data set may contact the SWAN Coordinating Center at the following email address: swanaccess@edc.pitt.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Conflict of Interest: All authors declared no actual or potential conflict of interest.

Results: At time of carotid ultrasound scan, women were on average 59.6 (± 2.7) years old and a majority was postmenopausal (88.4%). The women were White (48.4%), Black (31.2%), Chinese (13.3%) and Hispanic (7.1%). A 1 $\mu\text{g}/\text{m}^3$ higher 5-year mean annualized exposure to $\text{PM}_{2.5}$ was associated with an 8.0 μm (95% CI: 1.0 - 15.1) greater maximum cIMT at a later midlife, adjusting for cardiovascular disease risk factors; but was only related to IAD after adjusting for site. No association was found between either pollutant and plaque presence or plaque index.

Conclusions: Long-term exposure to $\text{PM}_{2.5}$ may contribute to elevated risk of atherosclerosis in the post-menopausal period.

Keywords

Long-term exposure; $\text{PM}_{2.5}$; Ozone (O_3); Subclinical Atherosclerosis; Women's Health

Introduction

Air pollution is ubiquitous and has deleterious effects on health. The most significant health outcomes related to air pollution are pulmonary and cardiovascular disease (CVD) (Brook, Rajagopalan et al. 2010, Franchini and Mannucci 2012). Since air pollution exists in our everyday environment, its cumulative effect may play a role in heart health (Atkinson, Carey et al. 2013, Hoek, Krishnan et al. 2013). Accumulating evidence from the last decade suggests that more than one-year exposure to air pollution may be particularly deleterious for CVD health. In analyses of several European cohorts, approximately 3-year exposure to particulate matter (PM) contributed to future all-cause mortality (Beelen, Raaschou-Nielsen et al. 2014) and coronary events (Cesaroni, Forastiere et al. 2014). Several cohort studies in the U.S. have also found a harmful effect of $\text{PM}_{2.5}$ on CVD mortality and morbidity in women (Miller, Siscovick et al. 2007, Puett, Hart et al. 2009, Hart, Puett et al. 2015).

Atherosclerosis, responsible for a majority of CVD events, is an inflammatory and lipids driven process, in which plaque builds up in the artery (Lu and Daugherty 2015). Biomarkers of subclinical atherosclerosis as outcomes have been widely adopted in research to study preclinical CVD. Carotid artery intima-media thickness (cIMT) measured via B-mode ultrasound, a surrogate biomarker of subclinical atherosclerosis (Bots and Grobbee 2002, Bauer, Caviezel et al. 2012), reflects early vascular changes and predicts CVD and stroke events independent of traditional CVD risk factors (Chambless, Folsom et al. 2000, Bots, Evans et al. 2003, Polak, Pencina et al. 2011, Bauer, Caviezel et al. 2012, van den Oord, Sijbrands et al. 2013). Ultrasound assessment of carotid plaque is a direct measure of atherosclerotic lesions. Plaque in conjunction with cIMT may be a better biomarker for predicting CVD risk compared to cIMT alone (Inaba, Chen et al. 2012), and utilizing quantitative measures of plaque may further improve prediction (Naqvi and Lee 2014).

The cross-sectional relationship between exposure to one-year air pollution and atherosclerosis has been examined in different populations (Künzli, Jerrett et al. 2005, Bauer, Moebus et al. 2010, Lenters, Uiterwaal et al. 2010, Breton, Wang et al. 2012, Rivera, Basagaña et al. 2012, Tonne, Yanosky et al. 2012, Kälisch, Hennig et al. 2013, Armijos, Weigel et al. 2015, Perez, Wolf et al. 2015). Most of these studies found that living close to major roads and exposure to higher air pollutant levels were related to atherosclerosis. There

are very few studies that have examined the association between more than 1-year exposure to air pollution and atherosclerosis; two of these studies were limited to young populations, whose atherosclerotic changes were not significant (Breton, Wang et al. 2012, Armijos, Weigel et al. 2015). Diez-Roux et al. observed an association between 20-years exposure to PM₁₀ using the EPA monitors and cIMT after adjusting for CVD risk factors (Diez Roux, Auchincloss et al. 2008). Rivera et al. related the 10-year exposure to NO₂, using land use regression and traffic proximity, to cIMT among a community population aged 32 - 86 years old in the Girona region of Spain (the REGICOR study) (Rivera, Basagaña et al. 2012). NO₂ and traffic proximity were related to cIMT in the unadjusted model, but only traffic proximity remained significantly associated with cIMT in the model adjusted for confounders. To our knowledge, there are no studies evaluating the effect of more than one-year cumulative exposure to air pollution on atherosclerosis in midlife women, a time of increased CVD susceptibility and accelerated progression of atherosclerosis in women (El Khoudary, Wildman et al. 2013). Thus, we investigated, in a cohort of women transitioning through the menopause, the association between 5-year mean annualized exposure to PM_{2.5} and O₃ and atherosclerosis burden assessed by cIMT and plaque, approximately 6.6 (range: 5.4 - 9.6) years later.

Material and Methods

Study Population

The Study of Women's Health Across the Nation (SWAN) is a community-based multi-center multi-ethnic cohort study of women's health as they transition the menopause. The SWAN study, conducted at seven sites across the U.S. began in 1996. This analysis involves the five sites that collected both air pollution and subclinical atherosclerosis outcome data, Detroit, MI; Oakland, CA; Pittsburgh, PA; Chicago, IL; and Newark, NJ. Each site recruited Whites and another minority racial group. Detroit, MI; Pittsburgh, PA and Chicago, IL recruited Black, Oakland, CA recruited Chinese; and Newark, NJ recruited Hispanics. Of the 1311 women who had a carotid ultrasound exam and readable images, 1188 (90.6%) had residential air pollutant estimates at visits 3 - 7 and were included. A detailed illustration of population inclusion is presented in supplementary figure S1. SWAN was approved by the Institutional Review Board at each site, and written informed consent was obtained from all participants. The study protocol conforms to the ethical guidelines of the 1975 Declaration of Helsinki.

Exposure to PM_{2.5} and O₃

Air pollution exposure was assessed by the Air Pollution Study, an ancillary study of SWAN. Detailed methods have been published elsewhere (Ostro, Malig et al. 2014, Green, Broadwin et al. 2016). Daily PM_{2.5} and O₃ values were retrieved from US EPA Air Quality System DataMart (US EPA 2015). Exposure to PM_{2.5} and O₃ was determined by monitors within 20 km from each participant's address. Monitor readings were used to calculate average exposure over 360 days before each of the five SWAN visits 3 - 7, corresponding to the years 1999-2005 (figure 1). The mean annualized level of air pollutants over this 5-year period, from this point forward referred to as *5-year mean annualized exposure*, was then calculated using the trapezoidal rule for the area under the curve (Yeh and Collegeville

1991) based on the yearly exposure from SWAN visits 3 - 7. Exposure data were limited to these SWAN visits, as visit 3 occurred in approximately 2000, when U.S. EPA first started to monitor PM_{2.5} level at the national level, and data for inflammatory/hemostatic biomarkers, lipids and glucose were available up to SWAN visit 7 (Green, Broadwin et al. 2016). In our assessment, we included all of the monitor information related to the participants' residential addresses. The number of PM_{2.5} monitors ranged from 6 - 19 and the number of ozone monitors ranged from 4 - 13 for each site. For residences with multiple monitors within 20 km, one monitor was selected to represent exposure. Selection was based on criteria balancing the number of women's clinic visits from Visit 3 to 7 with available exposure data for a monitor versus the distance to the monitor. Preference was typically given to the closer monitor. However, for example, if a monitor was located twice as far away as the closest monitor, it would be chosen if it provided more than doubled the number of clinic visits included in the study. If a participant moved (~13% of all women) during the year prior to her visit, exposure data from multiple addresses were weighted based on the time of move when assigning exposure, or evenly weighted if the move date was not available (Wu, Basu et al. 2017). The Detroit, MI site was excluded from O₃ exposure analyses since only four participants had available data

Assessment of cIMT and plaque

Ultrasound scans were implemented once, either at SWAN visit 12 or 13 (years 2009 to 2013). At each site, centrally trained and certified sonographers obtained carotid ultrasound images using a Terason t3000 Ultrasound System (Teratech Corp, Burlington, MA). Two images were obtained of the left and right distal common carotid artery (CCA) for later reading at the Ultrasound Reading Center (University of Pittsburgh Ultrasound Research Lab) (Wendelhag, Gustavsson et al. 1991). The inter-sonographer reliability for repeat carotid ultrasound scans was good to excellent, with intraclass correlation coefficient (ICC) for mean cIMT 0.72 - 0.95 and for inter-adventitial diameters (IAD) 0.80 - 0.98. AMS semi-automated edge detection software was used to measure the near and far wall cIMT by tracing the lumen-intima interface and the media-adventitia interface across the 1-cm CCA segment proximal to the carotid bulb. IAD were measured directly as the distance between the adventitial-medial interfaces on the near and the far wall. The reading reproducibility was excellent between and within readers, with ICC greater than 0.87. The mean of the average and maximal readings of both sides comprised mean and maximal cIMT, respectively. Sonographers at each site evaluated the presence and extent of plaque (grade) in each of 5 segments of the left and right carotid artery (distal and proximal CCA, bulb, and proximal internal and external carotid arteries). The grades from all segments of the combined left and right carotid artery were summed to create the plaque index (Sutton-Tyrrell, Kuller et al. 2002). Plaque index was categorized as: 0 (no observable plaque), 1 - 2, and greater than 2 (Sutton-Tyrrell, Lassila et al. 1998, Gepner, Young et al. 2017).

Assessment of other CVD risk factors

Covariate data were obtained at the SWAN baseline or follow-up visits. A detailed illustration of when the data were collected and extracted for this study is presented in figure 1. Self-reported race/ethnicity and education were collected at baseline. The smoking status was reported as never, past and current use at baseline, and the smoking status of the past-

year was asked again at each follow-up visit. All of the information was taken into account to determine the smoking status as past or current users at visit 12/13. We determined the medication use of antihypertensive, antidiabetic and hormone replacement therapy in the same fashion. Financial strain was assessed based on the question of how hard it was to pay for basics, which includes living expenses and medical treatment. A longitudinal category for financial strain was created and categorized into three groups, (a) consistently not hard; (b) mixed (more than 50% of time had no difficulties); (c) consistently hard (more than 50% of time had any difficulties) (Thurston, El Khoudary et al. 2014). The status of hypertension was determined by either systolic blood pressure (SBP) ≥ 130 mmHg, or a diastolic blood pressure (DBP) ≥ 85 mmHg (American Heart Association 2016, Sokol, Wirth et al. 2016), or use of antihypertensive medication. Diabetes status was determined by a fasting glucose level ≥ 126 mg/dL while not on steroid for two consecutive visits and/or reporting diabetes or use of antidiabetic medications (Mayo Clinic 2018). Physical measures of body mass index (BMI), and SBP and DBP were collected at all the visits using standard methodology. Blood samples were drawn after a 12-hour fast and were sent to the Medical Research Laboratories for analyses of high density lipoprotein cholesterol (HDL-c), low density lipoprotein cholesterol (LDL-c), total cholesterol, triglyceride, as well as high sensitivity C-reactive protein (hsCRP), tissue-type plasminogen activator antigen (tPA), and plasma plasminogen activator inhibitor 1 (PAI-1). Detailed methods have been published elsewhere (El Khoudary, Wang et al. 2016). These biomarker measures were available for visits baseline, 1, 3 - 7, 12 and 13. Corresponding to available air pollution data, 5-year mean levels of these biomarkers, as well as BMI and SBP, were calculated using SWAN visit 3 - 7 data. We observed that the continuous variables did not change very much over time (data not published yet).

Statistical analysis

Fasting glucose, insulin, triglyceride, hsCRP, tPA and PAI-1 were log transformed due to positive skewness. The distribution of the covariates and outcomes by PM_{2.5} quartiles are presented in table 1. We used linear or logistic regression to test the association between the participants' characteristics and PM_{2.5} level in quartiles (treated as continuous). Linear regression was used to estimate the association between visits 3 - 7 annualized mean exposure level of per 1 $\mu\text{g}/\text{m}^3$ PM_{2.5} and per 1 ppb ozone and visit 12/13 mean and maximum cIMT, and mean IAD. Logistic regression and multinomial logistic regression was used to estimate the relationship between the annualized mean exposure to air pollutants and plaque presence and plaque index, respectively. Multinomial logistic regression was used, since the assumption of proportional odds was violated (score test $p = 0.035$). A series of nested models were constructed to evaluate the association between air pollutant exposures and subclinical atherosclerosis adjusting for established confounders: (1) model controlling only for site, (2a) model additionally adjusting for SES/demographic factors, age, race, education, and financial strain, (2b) fully adjusted model (model 2a plus BMI, smoking, triglyceride, cholesterol, HDL-c, menopause status, hormone replacement therapy, fasting glucose, and antidiabetic and antihypertensive medication), and (3) extended model, additionally adjusting for potential intermediates, SBP, hsCRP, tPA and PAI-1. The main analyses are single pollutant models, which used PM_{2.5} or O₃ as the exposure. An exploratory analysis to identify high risk population was conducted by stratifying

participants' characteristics related to CVD risk. These strata have been used in several studies in the same domain (Diez Roux, Auchincloss et al. 2008, Bauer, Moebus et al. 2010, Kaufman, Adar et al. 2016). Analyses were performed with SAS (V9.3, SAS Institute, Cary, NC). All tests were two-sided with $\alpha = 0.05$.

Results

Among the 1188 women with air pollutant estimates at visits 3 - 7 and a carotid ultrasound scan at visit 12/13, 573 were White, 373 Black, 159 Chinese, and 83 Hispanic. At the time of the carotid ultrasound scan, women were a mean age of 59.6 years (SD = 2.7, range: 54 - 67), and approximately 90% were post-menopausal (table S1). In contrast, at visit 5, 5.0% were pre-, 43.0% were early peri-, 11.6% were late peri, and 40.3% were post menopause. Overall, the 5-year mean annualized PM_{2.5} exposure level was 14.9 (SD = 1.9) $\mu\text{g}/\text{m}^3$ (IQR: 13.6 - 16.1); and the annualized mean O₃ exposure level was 30.8 (SD = 6.2) ppb (IQR: 29.2 - 35.9). The correlation between the two pollutants PM_{2.5} and O₃ was $r=0.56$. Five-year mean annualized exposure to PM_{2.5} and ozone are shown in figure 2. Oakland, CA had the lowest PM_{2.5} exposure over time, with a mean of 12.0 $\mu\text{g}/\text{m}^3$ (SD=0.6), and relatively low level of O₃ exposure (mean=24.6 ppb and SD=5.9), but the range was very wide. All other four sites had comparable PM_{2.5} exposure levels. The mean cIMT was 0.79 mm (SD=0.12), maximum cIMT was 0.93 mm (SD=0.14) and the mean IAD was 7.20 mm (SD=0.67).

Baseline characteristics and 5-year mean levels of CVD risk factors by 5-year mean annualized exposure to PM_{2.5} quartiles are presented in table 1. Most of these factors were positively associated with PM_{2.5}. Those in lower PM_{2.5} exposure areas were more frequently never smokers and women who reported no financial strains. On average, women who lived in more polluted areas had higher BMI, blood pressure, LDL-c, insulin, hsCRP, tPA and PAI, and lower HDL-c. We also observed that women in the higher PM_{2.5} exposure quartiles had higher mean and maximum cIMT but were less likely to have plaque. In SWAN, it should be noted that Chinese women were more likely to have plaque.

In the main analyses of cIMT measures, PM_{2.5} was associated with mean cIMT in the model adjusted for site and SES (Model 2a, table 2). Women had a 7.57 μm (95% CI: 1.31 - 13.82) thicker cIMT for each 1 $\mu\text{g}/\text{m}^3$ 5-year mean annualized exposure to PM_{2.5}. However, this association was only marginally significant ($p = 0.076$) after adjusting for standard CVD risk factors (model 2b). Women with an average of 1 $\mu\text{g}/\text{m}^3$ 5-year mean annualized exposure to PM_{2.5}, had an 8.03 μm (95% CI: 1.01 - 15.05) thicker maximum cIMT on average (model 2b). In the models adjusted for SES plus diabetes (model 2b) and lipid (model 2c) biomarkers (supplementary tables S2 and S3), we noted there were small effect size reductions from model 2a to models 2b and 2c, but not as large as compared to model 3. After accounting for intermediates in the model, this association became null, which was mainly influenced by SBP (model 3). The association between PM_{2.5} and IAD was significant in the model adjusting for site, but not significant in the models adjusting for standard CVD risk factors. No clear association was observed between O₃ and any carotid measures (table 2). In the two-pollutant model, the estimates were slightly attenuated after adjusting for each other (supplementary tables S4 - S6). Neither of the two pollutants was significantly associated with plaque presence or plaque index in any model (table 3).

We also explored the association between air pollution and cIMT by participants' characteristics and CVD risk factors. We observed that 5-year mean annualized exposure to PM_{2.5} had a stronger effect on maximum cIMT in some of the subgroups (Figure 3), which was observed among diabetic, obese and more than college educated women. However, due to the disproportionate sample sizes across strata, no significant interactions were observed.

Discussion

In this 11-year observational study, we found that women who had a higher 5-year mean annualized PM_{2.5} over approximately 5 years during early midlife had a higher maximum cIMT in late midlife, controlling for traditional CVD risk factors. Significant associations were observed between PM_{2.5} and mean cIMT, independent of SES factors, and IAD after adjusting for site, but both these associations could be explained by standard CVD risk factors. No meaningful associations were found between PM_{2.5} and plaque and with O₃ and subclinical atherosclerosis.

This is the first study focused on the effects of 5-year mean annualized exposure to air pollution on atherosclerosis among women transitioning from peri-menopause to post-menopause with more than 10-year effects of observation. Our study observed women during the period of life when CVD risk accelerates rapidly. An earlier SWAN analysis indicated that late perimenopausal women have increased risk of CVD and the fastest progression of cIMT compared to women at an earlier stage of menopause (El Khoudary, Wildman et al. 2013). PM_{2.5} exposure was assessed in our study when most women were transitioning through the menopause. Thus, the exposure data was collected at the stage of life when it may have the greatest impact on atherosclerotic progression, vascular remodeling and CVD risk.

In this study, we revealed that women had an 8.03 μm (95% CI: 1.01 - 15.05) thicker maximum cIMT for each 1 $\mu\text{g}/\text{m}^3$ greater 5-year mean annualized exposure to PM_{2.5}, adjusting for traditional CVD risk factors. Consistent with our findings, Diez-Roux et al. found that 20-years exposure to observed PM₁₀ and imputed PM_{2.5} were associated with cIMT independently of CVD risk factors. As PM_{2.5} exposure levels were not available before 1999, they imputed the value using the spatial temporal model. Our study, using observed data, reached the same conclusion. Rivera and colleagues found that traffic intensity was associated with cIMT (Rivera, Basagaña et al. 2012). Traffic is one of the major sources of PM_{2.5} in urban areas (U.S. Environmental Protection Agency 2014). Although studies of more than one-year exposure to air pollutants and sub-clinical atherosclerosis are limited, the relationship between one-year PM_{2.5} exposure and cIMT has been observed in several cross-sectional and longitudinal studies (Bauer, Moebus et al. 2010, Künzli, Jerrett et al. 2010, Adar, Sheppard et al. 2013, Kaufman, Adar et al. 2016).

In many of the air pollution studies cited in the literature, one-year air pollution exposure is classified as long-term exposure. However, atherosclerosis is a life-long process likely resulting from long-term cumulative risk factor exposure, and its progression leading to clinical CVD events (Künzli, Perez et al. 2011). Thus, our primary aim was to assess the effects of greater than one-year exposure to PM_{2.5} and O₃ on atherosclerosis. However, we

also evaluated the effect of a shorter-term exposure of one year closer to the outcome measures and found that the associations between one-year exposure to PM_{2.5} at visit 7 and mean and maximum cIMT were slightly attenuated compared to 5-year mean annualized exposure levels (supplementary tables S7 - S9). The associations with PM_{2.5} and other subclinical atherosclerosis biomarkers, and ozone and all of the biomarkers were not changed very much. These findings confirm our hypothesis that more than one-year exposure to air pollution and longer follow-up time can contribute to atherosclerosis more meaningfully. Evidence from several large cohort studies reporting an association between more than one-year exposure to air pollution and future CVD morbidity and mortality also support our hypothesis. (Miller, Siscovick et al. 2007, Puett, Hart et al. 2009, Beelen, Raaschou-Nielsen et al. 2014, Cesaroni, Forastiere et al. 2014, Hart, Puett et al. 2015). All of these studies evaluated air pollution for one-year or across several years, and then followed up the participants for decades to clinical CVD events. From the Women's Health Initiative, Miller et al found that PM_{2.5} exposure in year 2000 was associated with cardiovascular events with a median follow-up of 6 years (Miller, Siscovick et al. 2007). In the Nurse's Health Study, the authors noted that up to 18-year exposure to PM was related to elevated CVD risk among diabetic women (Hart, Puett et al. 2015). These findings align with our hypothesis and results that several years of cumulative air pollution exposure may lead to elevated atherosclerosis risk as measured by maximum cIMT several years later.

The effect of air pollution on CVD risk has been extensively reviewed in several papers (Brook, Rajagopalan et al. 2010, Franklin, Brook et al. 2015). PM can trigger systemic oxidative stress and inflammation, increase adipokines and cytokines expression level, oxidize lipids, inflamed fat cells and thus, lead to atherosclerosis (Franklin, Brook et al. 2015). However, in this population-based study, we did not see an effect size change after adding hsCRP, tPA and PAI-1 in the fully adjusted model, which is consistent with findings from MESA Air study (Adar, Sheppard et al. 2013, Kaufman, Adar et al. 2016). We found that elevated systolic blood pressure and hypertension may be a factor mediating the effect of PM_{2.5} on cIMT, reducing the effect size by about 37%. There is ample evidence for the effects of PM_{2.5} on hypertension and elevation of blood pressure (Franklin, Brook et al. 2015), which are risk factors for atherosclerosis and CVD events (Allen, Siddique et al. 2014). Thus, it is plausible that elevated blood pressure may be a mediator of the effect of PM_{2.5} on atherosclerosis. Alternatively, the PM_{2.5} effect on carotid wall thickening, especially in the CCA and in the absence of plaque, may reflect adaptive vascular remodeling driven by hemodynamic stressors such as increased arterial pressure (Heusch, Libby et al. 2014).

In this study, we did not find evidence of an association between O₃ exposure and any biomarkers of subclinical atherosclerosis. There are very few studies examining the long-term exposure effect of O₃, since ambient O₃ level is largely dependent on ultra-violet intensity (U.S. Environmental Protection Agency 2015). Moreover, the acute effect of O₃ can be reversible in healthy individuals (Allen 2002) making the long-term effect of O₃ difficult to observe in areas with higher levels during the summer, and lower levels during the winter. In our sample, 5-year mean annualized exposure level of O₃ was about 30.8 (range: 17.8 - 46.1) ppb, much lower than the national ambient air quality standards (U.S. Environmental Protection Agency 2014). The only study examining O₃ exposure was among

college students in Los Angeles, CA (Breton, Wang et al. 2012). They noted that early childhood exposure, when students were 6 - 12-years old, impacted cIMT at age 19. Most of the participants in this study were originally from southern California, which may indicate that they were exposed to higher levels of O₃ all year round.

Our stratification analyses suggest that obese and diabetic women may be more susceptible to PM_{2.5}, as each 1 µg/m³ higher 5-year mean annualized exposure to PM_{2.5}, predicted a thicker maximum cIMT in these groups compared to the reference group. In contrast to the study by Rivera et al. (Rivera, Basagaña et al. 2012), we did not treat these factors as potential intermediates, but adjusting for these factors decreased the effect size about 20% compared to the minimally adjusted model. We did not observe a substantial effect size change from the models adjusting for diabetes and lipid biomarkers (supplementary tables S2 and S3). These biomarkers may not present strong mediation effects, however this would need to be tested formally in a large study population. In line with increased susceptibility in obese populations, several animal studies using APOE or LDLR knock-out mice, that are more susceptible to fat, showed an increase in plaque volume by 50% after exposing them to PM_{2.5} (Sun, Wang et al. 2005, Soares, Carvalho-Oliveira et al. 2009), although in our study PM_{2.5} was not associated with plaque measures. Diabetes may also interplay with oxidative stress and increase the risk in developing atherosclerosis (Bullon, Newman et al. 2014). In the Nurses' Health Study, they observed that women with diabetes exposed to higher levels of PM_{2.5} had an increased risk of incidence CVD, coronary heart disease and stroke (Hart, Puett et al. 2015).

We did not observe any association between either pollutant with plaque presence or plaque index. So far, there has been only one published study using plaque as an outcome (Gan, Allen et al. 2014). Consistent with our findings, in the M-CHAT study, the authors did not find significant associations between any of the pollutants, e.g. traffic, PM_{2.5}, NO_x, and noise, with plaque in the overall population. There is very weak evidence between PM_{2.5} and plaque in the current literature. Our study population may be still young to observe significant plaque burden. Although there were about 45% women who had plaque, most of them had only one or a few small plaques with plaque index of 1 - 2. The subclinical phase of atherosclerosis is very long (Künzli, Perez et al. 2011, Juhola, Magnussen et al. 2013). Thus, at this stage, plaque assessment may not yet reflect extensive subclinical atherosclerosis among these women.

Limitations

One of the limitations in our study is the gap between exposure assessment and outcomes collection, which was about 5 years. However, the data we have during year 2000 - 2005 corresponds to the higher exposure levels according to U.S. EPA (U.S. Environmental Protection Agency 2016). Moreover, several studies demonstrated that earlier long-term exposure to air pollution was related to CVD outcomes in a later time (Miller, Siscovick et al. 2007, Beelen, Raaschou-Nielsen et al. 2014, Cesaroni, Forastiere et al. 2014, Hart, Puett et al. 2015). Additionally, there likely is some overlap and gaps in the five years of exposure data since the study visits were not exactly 360 days apart. Another potential limitation is that the monitors used measured exposure estimation without any modeling. The modeled

data can overcome limitations due to missing values, and may add exposure variation if the spatial variance was well modelled. However, most of the modeling methods have limitations and uncertainties (Rao, Galmarini et al. 2011, Venkatram 2015). Moreover, if the extrapolation of the exposure modeling is to a period with extensive missing data from the monitors, the estimation may not be valid. Our outcome of cIMT only measured the CCA segment, which may not fully capture carotid arterial segments more prone to atherosclerosis. Rivera et al. reported a stronger association between 10-year NO₂ exposure and cIMT measured as the mean of 6 segments rather than the CCA segment only (Rivera, Basagaña et al. 2012). Lastly, prior cIMT information was not available to address pre-existing risk or conditions related to subclinical atherosclerosis.

Strengths

We have the advantage of studying air pollution exposure during the menopausal transition in women as a predictor of their post-menopausal subclinical atherosclerosis. SWAN collected extensive data of CVD risk factors of these women during the menopausal transition, allowing us to assess the independent association of five-year PM_{2.5} exposure with cIMT thickening in post-menopause. We uniquely established the association between 5-year mean annualized PM_{2.5} exposure and subclinical atherosclerosis about 7 years later. As atherosclerosis is a lifelong process (Künzli, Perez et al. 2011), longer exposure is more meaningful in establishing the relationship with atherosclerosis.

Conclusions

In conclusion, 5-year mean annualized exposure to PM_{2.5} in early mid-life independently contributes to atherosclerosis as measured by maximum CCA IMT at later mid-life in a multiethnic population-based cohort of women. The findings from this study extend the evidence in the current literature that long-term exposure to PM_{2.5} is harmful to heart health.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments:

Clinical Centers: *University of Michigan, Ann Arbor - Siobán Harlow, PI 2011 - present, MaryFran Sowers, PI 1994-2011; Massachusetts General Hospital, Boston, MA - Joel Finkelstein, PI 1999 - present, Robert Neer, PI 1994 - 1999; Rush University, Rush University Medical Center, Chicago, IL - Howard Kravitz, PI 2009 - present, Lynda Powell, PI 1994 - 2009; University of California, Davis/Kaiser - Ellen Gold, PI, University of California, Los Angeles - Gail Greendale, PI, Albert Einstein College of Medicine, Bronx, NY - Carol Derby, PI 2011 - present, Rachel Wildman, PI 2010 - 2011; Nanette Santoro, PI 2004 - 2010; University of Medicine and Dentistry - New Jersey Medical School, Newark - Gerson Weiss, PI 1994 - 2004; and the University of Pittsburgh, Pittsburgh, PA - Karen Matthews, PI.*

NIH Program Office: *National Institute on Aging, Bethesda, MD - Chhanda Dutta 2016- present; Winifred Rossi 2012-2016; Sherry Sherman 1994 - 2012; Marcia Ory 1994 - 2001; National Institute of Nursing Research, Bethesda, MD - Program Officers.*

Central Laboratory: *University of Michigan, Ann Arbor - Daniel McConnell (Central Ligand Assay Satellite Services).*

SWAN Repository: *University of Michigan, Ann Arbor - Siobán Harlow 2013 - Present; Dan McConnell 2011 - 2013; MaryFran Sowers 2000 - 2011.*

Coordinating Center: University of Pittsburgh, Pittsburgh, PA - Maria Mori Brooks, PI 2012 - present; Kim Sutton-Tyrrell, PI 2001 - 2012; New England Research Institutes, Watertown, MA - Sonja McKinlay, PI 1995 - 2001.

We thank the study staff at each site and all the women who participated in SWAN.

Financial Support: The Study of Women's Health Across the Nation (SWAN) has grant support from the National Institutes of Health (NIH), DHHS, through the National Institute on Aging (NIA), the National Institute of Nursing Research (NINR) and the NIH Office of Research on Women's Health (ORWH) (Grants U01NR004061; U01AG012505, U01AG012535, U01AG012531, U01AG012539, U01AG012546, U01AG012553, U01AG012554, U01AG012495, U01AG017719). This publication was supported in part by the National Center for Research Resources and the National Center for Advancing Translational Sciences, National Institutes of Health, through UCSF-CTSI Grant Number UL1 RR024131. The content of this manuscript is solely the responsibility of the authors and does not necessarily represent the official views of the NIA, NINR, ORWH or the NIH.

References

- Adar SD, Sheppard L, Vedal S, Polak JF, Sampson PD, Roux AVD, Budoff M, Jacobs DR, Jr, Barr RG and Watson K (2013). "Fine particulate air pollution and the progression of carotid intima-medial thickness: a prospective cohort study from the multi-ethnic study of atherosclerosis and air pollution." *PLoS medicine* 10(4): e1001430. [PubMed: 23637576]
- Allen J (2002). "The Ozone We Breathe" NASA http://earthobservatory.nasa.gov/Features/OzoneWeBreathe/ozone_we_breathe2.php (Accessed on January 24 2017).
- Allen NB, Siddique J, Wilkins JT, Shay C, Lewis CE, Goff DC, Jacobs DR, Liu K and Lloyd Jones D (2014). "Blood pressure trajectories in early adulthood and subclinical atherosclerosis in middle age." *Jama* 311(5): 490–497. [PubMed: 24496536]
- American Heart Association (2016). "About Metabolic Syndrome" http://www.heart.org/HEARTORG/Conditions/More/MetabolicSyndrome/About-MetabolicSyndrome_UCM_301920_Article.jsp#.Wws1i-4vzIU Access on May 27th 2018.
- Armijos RX, Weigel MM, Myers OB, Li WW, Racines M and Berwick M (2015). "Residential exposure to urban traffic is associated with increased carotid intima-media thickness in children." *J Environ Public Health* 2015: 713540. [PubMed: 25685160]
- Atkinson RW, Carey IM, Kent AJ, van Staa TP, Anderson HR and Cook DG (2013). "Long-term exposure to outdoor air pollution and incidence of cardiovascular diseases." *Epidemiology* 24(1): 44–53. [PubMed: 23222514]
- Bauer M, Caviezel S, Teynor A, Erbel R, Mahabadi AA and Schmidt-Trucksäss A (2012). "Carotid intima-media thickness as a biomarker of subclinical atherosclerosis." *Swiss Med Wkly* 142(10): 13705.
- Bauer M, Moebus S, Mohlenkamp S, Dragano N, Nonnemacher M, Fuchsluger M, Kessler C, Jakobs H, Memmesheimer M, Erbel R, Jockel KH and Hoffmann B (2010). "Urban particulate matter air pollution is associated with subclinical atherosclerosis: results from the HNR (Heinz Nixdorf Recall) study." *J Am Coll Cardiol* 56(22): 1803–1808. [PubMed: 21087707]
- Beelen R, Raaschou-Nielsen O, Stafoggia M, Andersen ZJ, Weinmayr G, Hoffmann B, Wolf K, Samoli E, Fischer P and Nieuwenhuijsen M (2014). "Effects of long-term exposure to air pollution on natural-cause mortality: an analysis of 22 European cohorts within the multicentre ESCAPE project." *The Lancet* 383(9919): 785–795.
- Bots ML, Evans GW, Riley WA and Grobbee DE (2003). "Carotid intima-media thickness measurements in intervention studies design options, progression rates, and sample size considerations: a point of view." *Stroke* 34(12): 2985–2994. [PubMed: 14615619]
- Bots ML and Grobbee DE (2002). "Intima media thickness as a surrogate marker for generalised atherosclerosis." *Cardiovascular drugs and therapy* 16(4): 341–351. [PubMed: 12652104]
- Breton CV, Wang X, Mack WJ, Berhane K, Lopez M, Islam TS, Feng M, Lurmann F, McConnell R, Hodis HN, Kunzli N and Avol E (2012). "Childhood air pollutant exposure and carotid artery intima-media thickness in young adults." *Circulation* 126(13): 1614–1620. [PubMed: 22896588]
- Brook RD, Rajagopalan S, Pope CA, Brook JR, Bhatnagar A, Diez-Roux AV, Holguin F, Hong Y, Luepker RV and Mittleman MA (2010). "Particulate matter air pollution and cardiovascular disease an update to the scientific statement from the American Heart Association." *Circulation* 121(21): 23312378.

- Bullon P, Newman HN and Battino M (2014). "Obesity, diabetes mellitus, atherosclerosis and chronic periodontitis: a shared pathology via oxidative stress and mitochondrial dysfunction?" *Periodontology* 2000 64(1): 139–153. [PubMed: 24320961]
- Cesaroni G, Forastiere F, Stafoggia M, Andersen ZJ, Badaloni C, Beelen R, Caracciolo B, de Faire U, Erbel R and Eriksen KT (2014). "Long term exposure to ambient air pollution and incidence of acute coronary events: prospective cohort study and meta-analysis in 11 European cohorts from the ESCAPE Project." *Bmj* 348: f7412. [PubMed: 24452269]
- Chambless LE, Folsom AR, Clegg LX, Sharrett AR, Shahar E, Nieto FJ, Rosamond WD and Evans G (2000). "Carotid wall thickness is predictive of incident clinical stroke: the Atherosclerosis Risk in Communities (ARIC) study." *American journal of epidemiology* 151(5): 478–487. [PubMed: 10707916]
- Diez Roux AV, Auchincloss AH, Franklin TG, Raghunathan T, Barr RG, Kaufman J, Astor B and Keeler J (2008). "Long-term exposure to ambient particulate matter and prevalence of subclinical atherosclerosis in the Multi-Ethnic Study of Atherosclerosis." *American journal of epidemiology* 167(6): 667–675. [PubMed: 18227099]
- El Khoudary SR, Wang L, Brooks MM, Thurston RC, Derby CA and Matthews KA (2016). "Increase HDL-C level over the menopausal transition is associated with greater atherosclerotic progression." *J Clin Lipidol* 10(4): 962–969. [PubMed: 27578129]
- El Khoudary SR, Wildman RP, Matthews K, Thurston RC, Bromberger JT and Sutton-Tyrrell K (2013). "Progression Rates of Carotid Intima-media Thickness and Adventitial Diameter during the Menopausal Transition." *Menopause* 20(1): 8–14. [PubMed: 22990755]
- Franchini M and Mannucci PM (2012). "Air pollution and cardiovascular disease." *Thrombosis research* 129(3): 230–234. [PubMed: 22113148]
- Franklin BA, Brook R and Pope CA (2015). "Air pollution and cardiovascular disease." *Current problems in cardiology* 40(5): 207–238. [PubMed: 25882781]
- Gan WQ, Allen RW, Brauer M, Davies HW, Mancini GJ and Lear SA (2014). "Long-term exposure to traffic-related air pollution and progression of carotid artery atherosclerosis: a prospective cohort study." *BMJ open* 4(4): e004743.
- Gepner AD, Young R, Delaney JA, Budoff MJ, Polak JF, Blaha MJ, Post WS, Michos ED, Kaufman J and Stein JH (2017). "Comparison of Carotid Plaque Score and Coronary Artery Calcium Score for Predicting Cardiovascular Disease Events: The Multi-Ethnic Study of Atherosclerosis." *Journal of the American Heart Association* 6(2): e005179. [PubMed: 28196817]
- Green R, Broadwin R, Malig B, Basu R, Gold EB, Qi L, Sternfeld B, Bromberger JT, Greendale GA, Kravitz HM, Tomey K, Matthews K, Derby CA, Jackson EA, Green R and Ostro B (2016). "Long- and Short-term Exposure to Air Pollution and Inflammatory/Hemostatic Markers in Midlife Women." *Epidemiology* 27(2): 211–220. [PubMed: 26600256]
- Hart JE, Puett RC, Rexrode KM, Albert CM and Laden F (2015). "Effect Modification of Long-Term Air Pollution Exposures and the Risk of Incident Cardiovascular Disease in US Women." *J Am Heart Assoc* 4(12).
- Hart JE, Puett RC, Rexrode KM, Albert CM and Laden F (2015). "Effect Modification of Long-Term Air Pollution Exposures and the Risk of Incident Cardiovascular Disease in US Women." *Journal of the American Heart Association* 4(12): e002301. [PubMed: 26607712]
- Heusch G, Libby P, Gersh B, Yellon D, Böhm M, Lopaschuk G and Opie L (2014). "Cardiovascular remodelling in coronary artery disease and heart failure." *The Lancet* 383(9932): 1933–1943.
- Hoek G, Krishnan RM, Beelen R, Peters A, Ostro B, Brunekreef B and Kaufman JD (2013). "Long-term air pollution exposure and cardio-respiratory mortality: a review." *Environ Health* 12(1): 43. [PubMed: 23714370]
- Inaba Y, Chen JA and Bergmann SR (2012). "Carotid plaque, compared with carotid intima-media thickness, more accurately predicts coronary artery disease events: a meta-analysis." *Atherosclerosis* 220(1): 128–133. [PubMed: 21764060]
- Juhola J, Magnussen CG, Berenson GS, Venn A, Burns TL, Sabin MA, Srinivasan SR, Daniels SR, Davis PH and Chen W (2013). "Combined effects of child and adult elevated blood pressure on subclinical atherosclerosis: the International Childhood Cardiovascular Cohort Consortium." *Circulation: CIRCULATIONAHA*. 113001614.

- Kälsch H, Hennig F, Moebus S, Möhlenkamp S, Dragano N, Jakobs H, Memmesheimer M, Erbel R, Jöckel K-H and Hoffmann B (2013). "Are air pollution and traffic noise independently associated with atherosclerosis: the Heinz Nixdorf Recall Study." *European heart journal* 35(13): 853–860. [PubMed: 24194529]
- Kaufman JD, Adar SD, Barr RG, Budoff M, Burke GL, Curl CL, Daviglius ML, Diez Roux AV, Gassett AJ, Jacobs DR, Jr., Kronmal R, Larson TV, Navas-Acien A, Olives C, Sampson PD, Sheppard L, Siscovick DS, Stein JH, Szpiro AA and Watson KE (2016). "Association between air pollution and coronary artery calcification within six metropolitan areas in the USA (the Multi-Ethnic Study of Atherosclerosis and Air Pollution): a longitudinal cohort study." *Lancet* 388(10045): 696–704. [PubMed: 27233746]
- Künzli N, Jerrett M, Garcia-Esteban R, Basagaña X, Beckermann B, Gilliland F, Medina M, Peters J, Hodis HN and Mack WJ (2010). "Ambient air pollution and the progression of atherosclerosis in adults." *PloS one* 5(2): e9096. [PubMed: 20161713]
- Künzli N, Jerrett M, Mack WJ, Beckerman B, LaBree L, Gilliland F, Thomas D, Peters J and Hodis HN (2005). "Ambient air pollution and atherosclerosis in Los Angeles." *Environmental health perspectives*: 201–206. [PubMed: 15687058]
- Künzli N, Perez L, von Klot S, Baldassarre D, Bauer M, Basagana X, Breton C, Dratva J, Elosua R and de Faire U (2011). "Investigating air pollution and atherosclerosis in humans: concepts and outlook." *Progress in cardiovascular diseases* 53(5): 334–343. [PubMed: 21414468]
- Lenters V, Uiterwaal CS, Beelen R, Bots ML, Fischer P, Brunekreef B and Hoek G (2010). "Longterm exposure to air pollution and vascular damage in young adults." *Epidemiology* 21(4): 512–520. [PubMed: 20407379]
- Lu H and Daugherty A (2015). "Recent Highlights of ATVB Atherosclerosis." *Arterioscler Thromb Vasc Biol* 35(3): 485–491. [PubMed: 25717174]
- Mayo Clinic (2018). "Diabetes - Diagnosis & Treatment." <https://www.mayoclinic.org/diseasesconditions/diabetes/diagnosis-treatment/drc-20371451> Accessed on May 20, 2018.
- Miller KA, Siscovick DS, Sheppard L, Shepherd K, Sullivan JH, Anderson GL and Kaufman JD (2007). "Long-term exposure to air pollution and incidence of cardiovascular events in women." *New England Journal of Medicine* 356(5): 447–458. [PubMed: 17267905]
- Naqvi TZ and Lee M-S (2014). "Carotid intima-media thickness and plaque in cardiovascular risk assessment." *JACC: Cardiovascular Imaging* 7(10): 1025–1038. [PubMed: 25051948]
- Ostro B, Malig B, Broadwin R, Basu R, Gold EB, Bromberger JT, Derby C, Feinstein S, Greendale GA and Jackson EA (2014). "Chronic PM2.5 exposure and inflammation: determining sensitive subgroups in mid-life women." *Environmental research* 132: 168–175. [PubMed: 24792413]
- Perez L, Wolf K, Hennig F, Penell J, Basagaña X, Aguilera I, Agis D, Beelen R, Brunekreef B and Cyrus J (2015). "Air Pollution and Atherosclerosis: A Cross-Sectional Analysis of Four European Cohort Studies in the ESCAPE Study." *Environmental health perspectives*.
- Polak JF, Pencina MJ, O'Leary DH and D'Agostino RB (2011). "Common carotid artery intima-media thickness progression as a predictor of stroke in multi-ethnic study of atherosclerosis." *Stroke* 42(11): 3017–3021. [PubMed: 21885840]
- Puett RC, Hart JE, Yanosky JD, Paciorek C, Schwartz J, Suh H, Speizer FE and Laden F (2009). "Chronic fine and coarse particulate exposure, mortality, and coronary heart disease in the Nurses' Health Study." *Environmental health perspectives* 117(11): 1697. [PubMed: 20049120]
- Rao ST, Galmarini S and Puckett K (2011). "Air Quality Model Evaluation International Initiative (AQMEII): advancing the state of the science in regional photochemical modeling and its applications." *Bulletin of the American Meteorological Society* 92(1): 23–30.
- Rivera M, Basagaña X, Aguilera I, Foraster M, Agis D, E. d. Pérez Groot, L., Mendez MA, Bouso L and Targa J (2012). "Association between long-term exposure to traffic-related air pollution and subclinical atherosclerosis: the REGICOR study." © *Environmental Health Perspectives*, 2012, vol. 121, p. 223–230. [PubMed: 23384708]
- Soares SR, Carvalho-Oliveira R, Ramos-Sanchez E, Catanozi S, da Silva LF, Mauad T, Gidlund M, Goto H and Garcia ML (2009). "Air pollution and antibodies against modified lipoproteins are associated with atherosclerosis and vascular remodeling in hyperlipemic mice." *Atherosclerosis* 207(2): 368–373. [PubMed: 19486979]

- Sokol A, Wirth MD, Manczuk M, Shivappa N, Zatonska K, Hurley TG and Hébert JR (2016). "Association between the dietary inflammatory index, waist-to-hip ratio and metabolic syndrome." *Nutrition research* 36(11): 1298–1303. [PubMed: 27865615]
- Sun Q, Wang A, Jin X, Natanzon A, Duquaine D, Brook RD, Aguinaldo JG, Fayad ZA, Fuster V, Lippmann M, Chen LC and Rajagopalan S (2005). "Long-term air pollution exposure and acceleration of atherosclerosis and vascular inflammation in an animal model." *Jama* 294(23): 3003–3010. [PubMed: 16414948]
- Sutton-Tyrrell K, Kuller LH, Matthews KA, Holubkov R, Patel A, Edmundowicz D and Newman A (2002). "Subclinical atherosclerosis in multiple vascular beds: an index of atherosclerotic burden evaluated in postmenopausal women." *Atherosclerosis* 160(2): 407–416. [PubMed: 11849665]
- Sutton-Tyrrell K, Lassila HC, Meilahn E, Bunker C, Matthews KA and Kuller LH (1998). "Carotid atherosclerosis in premenopausal and postmenopausal women and its association with risk factors measured after menopause." *Stroke* 29(6): 1116–1121. [PubMed: 9626281]
- Thurston RC, El Khoudary SR, Derby CA, Barinas-Mitchell E, Lewis TT, McClure CK and Matthews KA (2014). "Low socioeconomic status over 12 years and subclinical cardiovascular disease: the study of women's health across the nation." *Stroke* 45(4): 954–960. [PubMed: 24578209]
- Tonne C, Yanosky JD, Beevers S, Wilkinson P and Kelly FJ (2012). "PM mass concentration and PM oxidative potential in relation to carotid intima-media thickness." *Epidemiology* 23(3): 486–494. [PubMed: 22450694]
- U.S. Environmental Protection Agency (2014). "National Ambient Air Quality Standards (NAAQS)" <http://www.epa.gov/air/criteria.html> Accessed on May 23rd 2015.
- U.S. Environmental Protection Agency (2014). "Particulate Matter (PM)." <http://www.epa.gov/pmdesignations/faq.htm> Accessed on May 17th 2015.
- U.S. Environmental Protection Agency (2015). "Ground Level Ozone" <http://www.epa.gov/air/ozonepollution/> Accessed on May 17th 2015.
- U.S. Environmental Protection Agency (2016). "Air Quality Trends" <https://www3.epa.gov/airtrends/aqtrends.html> Accessed on March 23rd 2016.
- US EPA (2015). "AQS Data Mart" https://aqs.epa.gov/aqsweb/documents/data_mart_welcome.html (Accessed on Oct. 25th 2016).
- van den Oord SC, Sijbrands EJ, Gerrit L, van Klaveren D, van Domburg RT, van der Steen AF and Schinkel AF (2013). "Carotid intima-media thickness for cardiovascular risk assessment: systematic review and meta-analysis." *Atherosclerosis* 228(1): 1–11. [PubMed: 23395523]
- Venkatram A (2015). *Lectures on air pollution modeling*, Springer
- Wendelhag I, Gustavsson T, Suurkula M, Berglund G and Wikstrand J (1991). "Ultrasound measurement of wall thickness in the carotid artery: fundamental principles and description of a computerized analysing system." *Clinical Physiology and Functional Imaging* 11(6): 565–577.
- Yeh S-T and Collegeville G (1991). Using trapezoidal rule for the area under a curve calculation.

Highlights:

- Five-year effects of air pollution on atherosclerosis are examined in midlife women.
- Higher 5-year mean annualized exposure to PM_{2.5} contributed to greater maximum carotid intima-media thickness.
- PM_{2.5} exposure predicted inter-adventitial diameter when adjusting for site, but not after adjusting for covariates.
- No association was found between PM_{2.5} or ozone and plaque presence or severity.

Visit	00	01	02	03	04	05	06	07	08	09	10	11	12	13
Start year	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	interim	2009	2011
End year	1997	1999	2000	2001	2002	2003	2004	2005	2006	2009*	2008		2011	2013
Variables														
Age													X	X
Race/Ethnicity	X													
Education	X													
Carotid Ultrasound Scan													X	X
Air Pollution: PM _{2.5} and Ozone				X	X	X	X	X						
Menopause Status													X	X
Continuous Variables#				X	X	X	X	X						
Categorical Variables†													X	X

* Newark, NJ site skipped visit 10, and visit 9 were collecting data overlapped with visit 10's time frame

BMI, SBP, cholesterol, triglyceride, LDL-c, HDL-c, fasting blood glucose, insulin, hsCRP, tPA, and PAI-1

† Other categorical variables: smoking, diabetes, HTN, financial strain, and medications

Figure 1.

Timeline of the data collection and extraction for this analysis

Figure 2.
5-year mean annualized exposure levels of PM_{2.5} and O₃ by site

Figure 3.

Association between 5-year mean annualized exposure to $PM_{2.5}$ and mean of the maximum common carotid artery intima-media thickness (cIMT), by participant characteristics Note: The solid line shows the point estimate of $PM_{2.5}$ and maximum cIMT in the fully adjusted model (2b), with the shaded area indicates 95% CI. The dash line indicates 0, which is the null effect.

Table 1Characteristics by 5-year mean annualized exposure to PM_{2.5} exposure quartiles (N=1188)

	PM _{2.5} (µg/m ³)				p-trend
	9.8 – 13.58 (n=297)	13.58 – 15.34 (n=297)	15.34 – 16.14 (n=297)	16.14 – 20.84 (n=297)	
Ozone (ppb)	25.0 (6.2)	34.6 (3.3)	33.2 (3.3)	33.7 (4.2)	<.001
Age * (years)	60.1 (2.6)	59.3 (2.7)	59.3 (2.6)	60.0 (2.7)	.638
Race ** (%)					<.001
White	43.1	57.6	47.1	45.1	
Black	3.4	38.1	48.5	35.7	
Chinese	53.6	0	0	0	
Hispanic	0	4.4	4.4	19.2	
Ever Smoking * (%)	24.0	41.9	39.0	49.5	<.001
Education ** (%)					0.919
High School	20.9	24.8	23.7	22.3	
Some College/College	51.2	55.2	49.5	53.3	
> College	28.0	20.0	26.8	24.4	
Pay for Basis * (%)					<.001
Consistently Hard	17.5	38.1	36.7	38.4	
Mixed	29.6	27.6	24.6	25.9	
Consistently not hard	52.9	34.3	38.7	35.7	
Menopause Status * (%)					.389
Pre/Peri-menopause	1.7	2.7	2.4	2.4	
Post-menopause	91.6	86.9	86.5	87.9	
Other	6.7	10.4	11.1	9.8	
Ever hormone users (%)	42.1	43.4	43.8	43.8	.675
BMI	25.6 (5.8)	31.5 (7.7)	30.2 (6.7)	30.1 (6.4)	<.001
SBP (mmHg)	111.2 (11.4)	118.7 (15.5)	122.0 (15.7)	122.6 (13.5)	<.001
DBP (mmHg)	71.0 (7.5)	72.1 (8.7)	75.6 (9.1)	77.1 (7.9)	<.001
Hypertension * (%)	48.5	70.7	76.1	76.4	<.001
Hypertension Medication * (%)	33.3	55.6	55.6	60.6	<.001
Cholesterol (mg/dL)	203.2 (31.8)	201.7 (33.8)	197.9 (33.7)	204.2 (32.7)	.933
LDL-c (mg/dL)	115.8 (26.3)	118.8 (29.7)	116.5 (29.8)	122.5 (30.9)	.022
HDL-c (mg/dL)	62.3 (15.0)	58.1 (14.1)	56.9 (14.2)	56.9 (14.0)	<.001
Triglyceride (mg/dL) #	103.9 (76.7, 151.6)	101.4 (77.9, 151.5)	104.4 (80.4, 143.5)	108.0 (85.3, 146.0)	.616
Lipids Lowering Medication (%)	21.9	36.0	38.1	37.7	<.001
Glucose (mg/dL) #	88.9 (83.8, 94.3)	87.6 (83.3, 95.8)	88.6 (84.1, 95.8)	88.7 (83.8, 97.0)	.281

	PM _{2.5} (µg/m ³)				p-trend
	9.8 – 13.58 (n=297)	13.58 – 15.34 (n=297)	15.34 – 16.14 (n=297)	16.14 – 20.84 (n=297)	
Insulin (uIU/mL) [#]	9.3 (7.5, 12.4)	11.2 (8.3, 16.9)	11.4 (8.6, 15.9)	11.5 (8.5, 16.9)	<.001
Diabetes [*] (%)	7.4	15.8	16.5	19.2	<.001
Diabetic medication (%)	7.1	15.8	14.8	18.2	<.001
hsCRP(mg/L) [#]	1.2 (0.6, 2.9)	2.9 (1.0, 7.7)	2.9 (1.2, 5.9)	2.9 (1.3, 6.3)	<.001
tPA (ng/dL) [#]	6.5 (4.8, 8.4)	7.6 (5.5, 9.4)	7.4 (5.8, 9.6)	7.8 (6.0, 10.1)	<.001
PAI (ng/dL) [#]	14.4 (7.3, 26.1)	19.8 (10.9, 38.2)	19.1 (10.9, 35.1)	22.2 (11.9, 35.5)	<.001
Avg. cIMT [*] (mm)	0.76 (0.11)	0.80 (0.11)	0.80 (0.12)	0.81 (0.13)	<.001
Max. cIMT [*] (mm)	0.89 (0.13)	0.94 (0.13)	0.93 (0.14)	0.95 (0.15)	<.001
IAD [*] (mm)	7.12 (0.72)	7.24 (0.72)	7.24 (0.66)	7.20 (0.67)	.175
Plaque [*] (%)	47.8	41.9	40.4	40.4	.065
Plaque index [*] (%)					.025
0	52.2	58.1	59.6	59.6	
1–2	29.3	28.7	27.6	29.6	
3+	18.5	13.2	13.8	10.8	

* Data corresponds to carotid visits (SWAN visit 12/13);

** Data corresponds to baseline; All the other information was analyzed mean level of visit 3 – 7 corresponding to the available air pollution data;

[#] presented median (25th and 75th percentile), all the other continuous variables presented mean (std), and categorical variables presented percentage.

Table 2

Association between 5-year mean annualized exposure to PM_{2.5} and ozone and carotid artery intima-media thickness (cIMT) and inter-adventitial-diameter (IAD)

Models	Mean of average cIMT (μm)	Mean of maximum cIMT (μm)	Mean IAD (μm)
PM_{2.5} (μg/m³)			
Model 1	9.59 (3.19 – 15.98) ***	12.57 (5.21 – 19.93) ****	42.48 (6.77 – 78.19) **
Model 2a	7.57 (1.31 – 13.82) **	10.46 (3.29 – 17.63) ***	35.28 (–0.01 – 70.57) *
Model 2b	5.59 (–0.58 – 11.75) *	8.03 (1.01 – 15.05) **	21.18 (–12.60 – 54.96)
Model 3	3.22 (–2.83 – 9.26)	5.02 (–1.86 – 11.90)	8.99 (–24.26 – 42.26)
Model 2b + SBP	3.48 (–2.56 – 9.53)	5.40 (–1.48 – 12.28)	
Model 2b + HTN [†]	5.44 (–0.67 – 11.56) *	7.90 (0.94 – 14.86) **	
Model 2b + hsCRP	5.45 (–0.71 – 11.61) *	7.86 (0.84 – 14.87) **	
Model 2b + tPA	5.64 (–0.54 – 11.82) *	8.01 (0.97 – 15.04) **	
Model 2b + PAI-1	5.67 (–0.50 – 11.84) *	8.11 (1.08 – 15.14) **	
Ozone (ppb)[#]			
Model 1	–0.86 (–2.95 – 1.23)	–1.38 (–3.81 – 1.05)	–2.36 (–14.11 – 9.39)
Model 2a	–0.20 (–2.27 – 1.87)	–0.51 (–2.90 – 1.88)	0.16 (–11.60 – 11.91)
Model 2b	–0.49 (–2.51 – 1.53)	–0.92 (–3.23 – 1.38)	–0.47 (–11.67 – 10.72)
Model 3	–0.65 (–2.63 – 1.33)	–1.09 (–3.34 – 1.15)	–2.11 (–13.16 – 8.93)

Note: Model 1 adjust for site; model 2a further adjust for age, race, education, and longitudinal financial strain; model 2b additionally adjust for BMI, smoking, cholesterol, HDL-c, triglyceride, menopause status, hormone use, fasting glucose, antidiabetic medication, and antihypertensive medication; model 3 adjusting for intermediates: systolic blood pressure (SBP), C-reactive protein, tPA and PAI-1.

Data presented as effect (95% confidence interval)

* Marginal significant, 0.05 < p < 0.1

** p < 0.05

*** p < 0.01

[#] Michigan site was excluded, since there were extensive missing values with ozone exposure, N = 826.

Table 3

Association between 5-year mean annualized exposure to PM_{2.5} and ozone and plaque presence and plaque index

Models	Odd Ratio of Plaque Presence	Odds Ratio of Plaque index		
		0	1–2	> 2
<u>PM_{2.5} (µg/m³)</u>				
Model 1	0.98 (0.88 – 1.09)	Reference	0.99 (0.88 – 1.12)	0.96 (0.82 – 1.12)
Model 2a	1.00 (0.89 – 1.11)	Reference	1.01 (0.89 – 1.14)	0.99 (0.85 – 1.16)
Model 2b	0.98 (0.87 – 1.10)	Reference	1.01 (0.89 – 1.15)	0.93 (0.79 – 1.10)
Model 3	0.97 (0.86 – 1.09)	Reference	1.01 (0.88 – 1.15)	0.91 (0.76 – 1.08)
<u>Ozone (ppb)[#]</u>				
Model 1	1.01 (0.97 – 1.05)	Reference	1.03 (0.99 – 1.07)	0.98 (0.93 – 1.03)
Model 2a	1.01 (0.97 – 1.05)	Reference	1.03 (0.98 – 1.07)	0.98 (0.93 – 1.03)
Model 2b	1.01 (0.97 – 1.05)	Reference	1.02 (0.98 – 1.07)	0.98 (0.93 – 1.04)
Model 3	1.01 (0.97 – 1.05)	Reference	1.03 (0.98 – 1.07)	0.98 (0.92 – 1.03)

Note: Model 1 adjust for site; model 2a further adjust for age, race, education, and longitudinal financial strain; model 2b additionally adjust for BMI, smoking, menopause status, hormone use, cholesterol, HDL-c, fasting glucose, antidiabetic medication use, and antihypertensive medication use; model 3 adjusting for intermediates: systolic blood pressure (SBP), C-reactive protein, tPA and PAI-1.

Data presented as odds ratio (95% confidence interval)

[#] Michigan site was excluded, since there were extensive missing values with ozone exposure, N = 826.