Published in final edited form as:

Chem Commun (Camb). 2018 November 01; 54(88): 12487–12490. doi:10.1039/c8cc05809a.

Ru(II) Complexes with Diazine Ligands: Electronic Modulation of Coordinating Group is Key to the Design of “Dual Action” Photoactivated Agents

Dmytro Havrylyuk^a, Megha Deshpande^a, Sean Parkin^a, and Edith C. Glazer^a

^aDepartment of Chemistry, 505 Rose St., University of Kentucky, Lexington, KY 40506

Abstract

Coordination complexes can be used to photocage biologically active ligands, providing control over the location, time, and dose of a delivered drug. Dual action agents can be created if both the ligand released and the ligand-deficient metal center effect biological processes. Ruthenium (II) complexes coordinated to pyridyl ligands generally are only capable of releasing one ligand in H₂O, wasting equivalents of drug molecules, and producing a Ru(II) center that is not cytotoxic. In contrast, Ru(II) polypyridyl complexes containing diazine ligands eject both monodentate ligands, with the quantum yield (Φ_{ps}) of the second phase varying as a function of ligand pK_a and the pH of the medium. This effect is general, as it is effective with different Ru(II) structures, and demonstrates that diazine-based drugs are the preferred choice for the development of light-activated dual action Ru(II) agents.

Graphical Abstract

Light-triggered Ru(II) molecules that produce active species capable of forming covalent adducts with DNA are being actively explored for photoactivated chemotherapy (PACT).¹ A complementary and potentially compatible approach is “photocaging,” where a biologically active, monodentate ligand is masked by coordination to a metal center.^{2–7} Photo-release of this ligand allows for temporal and spatial control over activity. However, a persistent issue that has limited the utility of Ru(II) photocages is the sluggish photochemistry associated

Conflicts of interest

There are no conflicts to declare.

with ejecting the monodentate ligand. Strain-inducing bidentate ligands are known to activate dissociative photochemical pathways within cells,^{8–11} and have been used to increase the photo-lability of the monodentate ligand.^{12–14}

“Dual action” light-activated Ru(II) compounds, where both the metal center and the liberated ligands induce different, potentially synergistic biological effects, are also of interest.^{15–16} The same issue hampers development of such dual action agents, however, as photocages; the photoreactivity of the second ligand is often orders of magnitude lower than the first.¹⁷ This sub-optimal photochemistry results in the waste of one equivalent of the drug ligand, but more importantly, the opening of two binding sites on the metal appears important for the creation of the Ru(II) cytotoxic species.^{18,19–20} This is in notable contrast to platinum species such as phenanthrinplatin, a highly potent cytotoxin with only one reactive site.²¹

The limitation of photoejection of only one ligand has been demonstrated for Ru(II) photocages containing pyridine, imidazole, aliphatic amine derivatives,²² and phosphine ligands.²³ These light-induced ligand dissociation phenomena have been explored by ultrafast spectroscopy techniques and computational approaches.^{24–25} In contrast, release of two monodentate ligands has been shown for complex with 5-cyanouracil,²⁶ demonstrating that Ru(II) bound nitriles undergo light-activated ligand exchange more efficiently than other monodentate ligands.⁷ However, the second ligand ejection is slow, and the majority of nitrile-containing drugs, including anticancer agents, are derivatives of nitrogen-containing heterocycles.²⁷ Therefore, coordination with a Ru scaffold by a direct synthetic pathway could be complicated by the presence of coordination isomers in the prodrug molecule.

Considering the difference between acid-base properties and photochemical behaviors for Ru(II) complexes with aliphatic amines or pyridine (hard and borderline bases, respectively) and nitriles (soft bases),^{28,29} we hypothesized that *cis*-[Ru(bpy)₂(L₂)]²⁺ (bpy = 2,2'-bipyridyl) complexes with monodentate diazines (which are softer bases than pyridine) would efficiently release two ligands upon light irradiation in aqueous media. This hypothesis is partially supported by the fact that incorporation of the bidentate diazine ligand bipyrimidine in Ru(II) complexes results in a photochemically active species without any strain-inducing groups, in marked contrast to bipyridine complexes.^{29–30} Diazines are privileged heterocycles in medicinal chemistry,⁹ and are bioisosteres for pyridine and phenyl rings, so this study was further motivated by the presence of these moieties in several known drugs and the potential for incorporation in many more. The results of this report may be applied as a started platform to develop improved light-activated photocages and dual-action ruthenium complexes with diazine-based antitumor agents.

Three complexes with isomeric diazine coligands, *cis*-[Ru(bpy)₂(pyd)₂]²⁺ (pyd = pyridazine; compound **2**), *cis*-[Ru(bpy)₂(pym)₂]²⁺ (pym = pyrimidine; **3**), and *cis*-[Ru(bpy)₂(pyz)₂]²⁺ (pyz = pyrazine; **4**) were synthesized and their photochemical properties compared to *cis*-[Ru(bpy)₂(py)₂]²⁺ (**1**; py = pyridine; Scheme 1). The complexes were prepared under low light conditions by refluxing *cis*-[Ru(bpy)₂Cl₂] with a 10-fold excess of the desired diazine in ethanol:water (1:1). Complexes **1**, **2** and **4** have been reported previously,^{31–32} including the photophysical properties³³ and some photosubstitution reactions³⁴ for complexes **1** and

2. NMR spectroscopy and X-ray crystal structure have been reported for **1**,³² though noteworthy differences were found in the NMR of compound **1**.³⁵

Structural analysis by x-ray crystallography revealed distorted octahedral geometries for complexes **2** and **3**. The complexes exhibited altered [Ru–N] bond lengths in comparison to pyridine-containing **1** (Table S4);³² the Ru–N(pyd) and Ru–N(pym) bonds (**2**, **3**) are equal, in contrast to Ru–N(py) bonds (**1**) where different length were found (2.063 and 2.13 Å). The N–Ru–N bond angles between diazine ligands are also distorted from ideal 90° and 180°, with the deviations of N2*–Ru–N2 (angle between trans-nitrogens in the bpy coligands) for complexes **2** (8.04°) and **3** (6.74°) larger than for **1** (5°) (Fig. S1A, C). The pyridazine ligands are bent from N2 (the top bpy nitrogen, Fig. S1B) with different bond angles (88.44° and 97.13°), and the bond angle between pyridazines is 92.53° in contrast to 90° between pyridine ligands in complex **1**.³² This distortion was anticipated to affect the photochemical reactivity of the complexes.

All four Ru(II) complexes (**1–4**) were relatively stable in the dark for 72 h at 37 °C (Fig. S21), and exhibited selective photoejection of the first monodentate ligand in water (monitored by absorption spectroscopy; Scheme 1; Fig. S2–5) when irradiated with 470 nm light. The presence of an isosbestic point was interpreted as indicating the direct conversion to a single product. Compound **1** formed *cis*-[Ru(bpy)₂(py)(H₂O)]²⁺ (see absorption profile in Fig. 2B) and no further ligand loss was observed.

In marked contrast, a second photoreaction was observed for compounds **2–4** (Fig. 1A, C, S3–5). The quantum yields of photosubstitution by water (ϕ_{PS}) are shown in Table 1 and half-lives ($t_{1/2}$) are provided in Table S13. Only the diazine complexes ejected two ligands. The first ligand photoejection is facile, with $t_{1/2}(1)$ of less than 1 min for complexes **1–4** and ϕ_{PS} of 0.031–0.11. In contrast, the ϕ_{PS} (**2**) for **2–4** in water were inverted relative to ϕ_{PS} (**1**), indicating both the sensitivity of the photochemistry to the identity of diazine ligand, and the presence of a specific chemical feature that drives the disparity in the yields of these sequential processes. The ϕ_{PS} values for the second ligand ejection ranged from 0.0005–0.0033, with **4** exhibiting the highest quantum yield. The product was identified as *cis*-[Ru(bpy)₂(H₂O)₂]²⁺ by both HPLC and absorption spectroscopy (Fig. 1A, S3–5). The retention time (RT) and absorption profile was compared with [Ru(bpy)₂]-based products after irradiation of strained compound **5** [Ru(bpy)₂(dmbpy)]²⁺ (dmbpy – 6,6'-dimethyl-2,2'-bipyridyl).⁸ Further exposure of *cis*-[Ru(bpy)₂(H₂O)₂]²⁺ to light after photoejection resulted in a decrease in the intensity of the MLCT (metal-to-ligand charge-transfer) absorption band at 490 nm following irradiation after 90 (**4**), 120 (**3**) and 150 (**2**) minutes, likely due to oxidation of the Ru(II) center.¹³

An inverse relationship was found between $\phi_{PS}(2)$ for complexes **2–4** and the pK_a values for protonated parent ligands, as shown in Fig. 2A. The weaker basic diazine ligands are more photolabile, and there was no ejection of the pyridine, which is the strongest base in this series. The quantum yields also correlate with chemical shifts of the α protons of the diazine ligands (Fig. 2B). In addition, ϕ_{PS} and $t_{1/2}(2)$ were found to be sensitive to the environment (pH), as compounds **2–4** demonstrated 1.7–2-fold faster ligand ejection in HCl–KCl buffer, pH = 2 than in sodium phosphate buffer, pH=7.4 (Table S4). A similar effect was observed

with 1.8–3.5-fold faster $t_{1/2}$ values in D₂O vs. H₂O. A possible explanation is that engagement of the non-bonding electrons of the uncoordinated aza nitrogen, either through protonation or hydrogen bonding, accelerates the photochemistry either by forming a pre-encounter complex with the incoming ligand or by polarizing the electrons on the diazine ligand.

DNA damage was assessed by gel electrophoresis (Fig. 3). Incubation of each Ru(II) complex with plasmid DNA in the dark showed no interactions (Fig. S8), in contrast to the two types of DNA damage observed upon irradiation with 470 nm light. The diazine complexes **2–4** undergo ligands loss and covalent attachment to DNA, as observed by the reduction of DNA mobility and loss of ethidium bromide (EtBr) staining. Complex **1** created a combination of covalent damage and single strand breaks to form relaxed circular DNA, possibly through sensitization of singlet oxygen (¹O₂). There is some indication of single strand breaks for compound **2** as well, which has the slowest $t_{1/2}$ (2) of the diazine-containing complexes. Systems that both photoeject and create reactive oxygen species have previously been identified as useful dual mechanism agents;^{8, 36} if a diazine-containing drug were ligated, these would become triple action agents.

In order to determine if incorporation of diazines for improvement of Ru(II) complexes photo-lability is generalizable to other Ru(II) structures, the photochemistry of two *trans*-Ru(II) complexes containing pyridine (**6**) and pyrazine (**7**) ligands was investigated. Previously, *trans* Ru(II) complex containing thermally exchangeable ligands exhibited higher potency than an analogous *cis* compound,³⁷ making this an appealing scaffold for the creation of dual action agents. Accordingly, *trans*-[Ru(qpy)(pyz)]²⁺ (**7**; qpy = 2,2':6',2'':6'',2'''-quaterpyridine) was synthesized and compared to the pyridine analogue **6**, which is photo-stable in aqueous media. In contrast, the *trans*-Ru(II) complex with pyrazine ligands ejected the pyrazine ligand upon irradiation for nine hours (Fig. S9).

Finally, to demonstrate the useful application for a pure “photocaging” approach,^{38–39} two Ru(II) complexes (**8**, **9**) were synthesized using a [Ru(tpy)(bpy)]²⁺ scaffold (tpy=2,2':6'-2''-terpyridine). The photochemical reaction of the complex containing pyridine, [Ru(tpy)(bpy)(py)]²⁺ (**8**), did not reach completion after nine hours irradiation (Fig. 4A, B), which is consistent with previous reports of $\phi_{PS} < 10^{-5}$ in MeCN.^{40,41} In contrast [Ru(tpy)(bpy)(pyz)]²⁺ (**9**) exhibited significantly enhanced photolability of pyrazine upon irradiation in aqueous media, with $t_{1/2}$ =10 min and complete ligand exchange in two hours (Fig. 4C, D, $\phi_{PS} = 0.007$ in water). This significantly improved photochemistry makes [Ru(tpy)(bpy)]²⁺ a useful photocage in water if a diazine ligand is used.

In conclusion, rather than incorporating strain-inducing bidentate ligands in Ru(II) scaffolds^{8–10, 42} or substituting monodentate ligands with electron withdrawing groups,⁴³ these results demonstrate that simply using diazine ligands radically improves photochemical features. The approach works for a variety of Ru(II) scaffolds, and facilitates the ejection of two ligands from the *cis*-[Ru(bpy)₂]²⁺ cage. We posit that simple electronic tuning by switching from pyridine to diazine systems is a far more efficient and flexible approach for the creation of functional light-activated metal complexes. These results

suggest that photochemistry can be tuned by judicious use of ligands based on pK_a values and HSAB theory.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We gratefully acknowledge the National Institutes of Health (Grant GM107586) for the support of this research. Crystallographic work was made possible by the National Science Foundation (NSF) MRI program, grants CHE-0319176 and CHE-1625732.

Notes and references

1. Farrer NJ; Salassa L; Sadler PJ, Dalton Trans 2009, (48), 10690–701. [PubMed: 20023896]
2. Zayat L; Calero C; Albores P; Baraldo L; Etchenique R, J. Am. Chem. Soc 2003, 125 (4), 882–3. [PubMed: 12537482]
3. Zayat L; Salierno M; Etchenique R, Inorg. Chem 2006, 45 (4), 1728–31. [PubMed: 16471986]
4. Respondek T; Garner RN; Herroon MK; Podgorski I; Turro C; Kodanko JJ, J. Am. Chem. Soc 2011, 133 (43), 17164–7. [PubMed: 21973207]
5. Li A; Turro C; Kodanko JJ, Chem. Commun 2018, 54 (11), 1280–1290.
6. Renfrew AK, Metallomics 2014, 6 (8), 1324–35. [PubMed: 24850462]
7. Mosquera J; Sanchez MI; Mascarenas JL; Eugenio Vazquez M, Chem Commun (Camb) 2015, 51 (25), 5501–4. [PubMed: 25582525]
8. Howerton BS; Heidary DK; Glazer EC, J. Am. Chem. Soc 2012, 134 (20), 8324–7. [PubMed: 22553960]
9. Wachter E; Heidary DK; Howerton BS; Parkin S; Glazer EC, ChemComm 2012, 48 (77), 9649–51.
10. Hidayatullah AN; Wachter E; Heidary DK; Parkin S; Glazer EC, Inorg. Chem 2014, 53 (19), 10030–2. [PubMed: 25198057]
11. Havrylyuk D; Heidary DK; Nease L; Parkin S; Glazer EC, Eur. J. Inorg. Chem 2017, 2017 (12), 1687–1694. [PubMed: 29200939]
12. Arora K; White JK; Sharma R; Mazumder S; Martin PD; Schlegel HB; Turro C; Kodanko JJ, Inorg Chem 2016, 55 (14), 6968–79. [PubMed: 27355786]
13. Huisman M; White JK; Lewalski VG; Podgorski I; Turro C; Kodanko JJ, Chem Commun (Camb) 2016, 52 (85), 12590–12593. [PubMed: 27711349]
14. Wei J; Renfrew AK, J. Inorg. Biochem 2018, 179, 146–153. [PubMed: 29180165]
15. Zamora A; Denning CA; Heidary DK; Wachter E; Nease LA; Ruiz J; Glazer EC, Dalton Trans 2017, 46, 2165–2173. [PubMed: 28121322]
16. Chan H; Ghrayche JB; Wei J; Renfrew AK, Eur. J. Inorg. Chem 2017, 2017 (12), 1679–1686.
17. Smith NA; Zhang P; Greenough SE; Horbury MD; Clarkson GJ; McFeely D; Habtemariam A; Salassa L; Stavros VG; Dowson CG; Sadler PJ, Chem Sci 2017, 8 (1), 395–404. [PubMed: 28451184]
18. The cytotoxicity appears to depend on the coligands used.
19. Cuello-Garibo JA; Meijer MS; Bonnet S, Chem Commun (Camb) 2017, 53 (50), 6768–6771. [PubMed: 28597879]
20. Sgambellone MA; David A; Garner RN; Dunbar KR; Turro C, J Am Chem Soc 2013, 135 (30), 11274–82. [PubMed: 23819591]
21. Park GY; Wilson JJ; Song Y; Lippard SJ, Proc Natl Acad Sci U S A 2012, 109 (30), 11987–92. [PubMed: 22773807]
22. White JK; Schmehl RH; Turro C, Inorganica Chim Acta 2017, 454, 7–20. [PubMed: 28042171]
23. Franco S; Manuel SR; Samira NG; Antonio R, Eur. J. Inorg. Chem 2016, 2016 (10), 1528–1540.

24. Salassa L; Garino C; Salassa G; Gobetto R; Nervi C, *J Am Chem Soc* 2008, 130 (29), 9590–7. [PubMed: 18588292]
25. Greenough SE; Roberts GM; Smith NA; Horbury MD; McKinlay RG; Zurek JM; Paterson MJ; Sadler PJ; Stavros VG, *Phys. Chem. Chem. Phys* 2014, 16 (36), 19141–55. [PubMed: 25060066]
26. Garner RN; Gallucci JC; Dunbar KR; Turro C, *Inorg. Chem* 2011, 50 (19), 9213–5. [PubMed: 21879748]
27. Fleming FF; Yao L; Ravikumar PC; Funk L; Shook BC, *J. Med. Chem* 2010, 53 (22), 7902–17. [PubMed: 20804202]
28. House JE, *Inorganic Chemistry*, p. 297–98.
29. Crutchley RJ; Lever ABP, *Inorg. Chem* 1982, 21 (6), 2276–2282.
30. Vicendo P; Mouysset S; Paillous N, *Photochem. Photobiol* 1997, 65 (4), 647–55. [PubMed: 9114740]
31. Adeyemi SA; Johnson EC; Miller FJ; Meyer TJ, *Inorg. Chem* 1973, 12 (10), 2371–2374.
32. Hitchcock PB; Seddon KR; Turp JE; Yousif YZ; Zora JA; Constable EC; Wernberg O, *J. Chem. Soc., Dalton Trans* 1988, (7), 1837–1842.
33. Caspar JV; Meyer TJ, *Inorg. Chem* 1983, 22 (17), 2444–2453.
34. Pinnick DV; Durham B, *Inorg. Chem* 1984, 23 (10), 1440–1445.
35. In contrast to the described ¹H NMR spectrum (360 MHz, (CD₃)₂SO) for **1**, some resonances for H₅ and H_{5'} of bpy ligands (**1**–**4**) are present as doublet of doublets of doublets (ddd) (Figure S10). Detailed NMR spectra and interpretations are presented in Supporting Information.
36. Loftus LM; White JK; Albani BA; Kohler L; Kodanko JJ; Thummel RP; Dunbar KR; Turro C, *Chemistry – A European Journal* 2016, 22 (11), 3704–3708.
37. Wachter E; Zamora A; Heidary DK; Ruiz J; Glazer EC, *Chem. Commun* 2016, 52 (66), 10121–4.
38. Lameijer LN; Ernst D; Hopkins SL; Meijer MS; Askes SHC; Le Devedec SE; Bonnet S, *Angew. Chem. Int. Ed. Engl* 2017, 56 (38), 11549–11553. [PubMed: 28666065]
39. Li A; Yadav R; White JK; Herroon MK; Callahan BP; Podgorski I; Turro C; Scott EE; Kodanko JJ, *Chem Commun (Camb)* 2017, 53 (26), 3673–3676. [PubMed: 28304025]
40. Ru(II) complexes have higher quantum yields in MeCN than in water.
41. Knoll JD; Albani BA; Durr CB; Turro C, *J. Phys. Chem. A* 2014, 118 (45), 10603–10. [PubMed: 25027458]
42. Knoll JD; Albani BA; Turro C, *Acc Chem Res* 2015, 48 (8), 2280–7. [PubMed: 26186416]
43. Vu AT; Santos DA; Hale JG; Garner RN, *Inorg. Chim. Acta* 2016, 450, 23–29.

Fig. 1.

Determination of photoejection products by HPLC. A) Chromatogram of **1** (blue line, after 60 min irradiation), **2** (red line), **3** (green line) and **4** (violet line) after 120 min irradiation in comparison to **5** (black dash line, after 15 min irradiation). B) Absorption profile of Ru(II) photoproducts of **1** (blue line, RT = 8.35 min, cis -[Ru(bpy)₂(py)(H₂O)]²⁺) and **5** (black dash line, RT = 7.87 min, cis -[Ru(bpy)₂(H₂O)₂]²⁺). C) Absorption profile of Ru(II) photoproducts of **2** after ejection of first (cyan line, 5 min irradiation, RT = 8.36 min) and second pyridazine ligand (red line, 120 min irradiation, RT = 7.99 min) overlaid with cis -[Ru(bpy)₂(H₂O)₂]²⁺ (black dash line, RT = 7.87 min).

Fig. 2. Correlation between quantum yields for the second ligand ejection for **2** (green ○), **3** (red □), **4** (blue △) and pKa of protonated free diazines (A) or chemical shifts of α protons of diazine ligands (B).

Fig. 3.

Agarose gel electrophoresis showing the dose response of compounds A) **1**, B) **2**, C) **3** and D) **4** incubated with 40 μg/mL pUC19 DNA with irradiation (470 nm light). Lanes 1 and 12, DNA ladder; lane 2, EcoRI; lane 3, Cu(OP)₂; lane 4–11, 0–500 μM. EcoRI and Cu(OP)₂ are controls for linear and relaxed circle DNA. See SI for full gels.

Fig. 4. Photochemistry of [Ru(tpy)(bpy)(L)]²⁺ complexes. A) Absorption spectra of **8** with irradiation (blue line, t=0, red line, t= 540 min). B) HPLC of **8** as a function of irradiation time. C) Absorption spectra of **9** with irradiation (blue line, t=0, red line, t= 100 min). D) Comparison of % conversion for **8** (red open □) and **9** (green open ○). The integrated area under the peaks from B) was used for % conversion of **8**.

Scheme 1.

Photochemical reactions of complexes included in this study in H₂O.

Table 1.Photophysical and photochemical^a properties of compounds **1–9** in H₂O

Compound	$\lambda_{\text{max abs}}$ (nm) ^b		ϕ_{PS} ^d	
	A	B	(1)	(2)
1	455	-	0.031	-
2	420	450	0.11	0.0005
3	415	460	0.070	0.0011
			0.059 ^e	0.0013 ^e
4	405	445	0.11	0.0033
5			0.022	-
6			nd	nd
7			nd	nd
8			nd	-
9			0.007	-

^a Measured using a 13 mW/cm² 470 nm LED.^b For the MLCT.^c From the fit to a single exponential.^d See SI for a detailed description.^e Determined by HPLC.