Published in final edited form as:

J Am Chem Soc. 2018 November 14; 140(45): 15181–15185. doi:10.1021/jacs.8b09909.

Diastereoselective and Enantioselective Conjunctive Cross-Coupling Enabled by Boron Ligand Design

Jesse A. Myhill, Christopher A. Wilhelmsen, Liang Zhang, and James P. Morken*

Department of Chemistry, Merkert Chemistry Center, Boston College, Chestnut Hill, Massachusetts 02467, United States

Abstract

Enantio- and diastereoselective conjunctive cross-coupling of β -substituted alkenylboron “ate” complexes is studied. While β substitution shifts the chemoselectivity of the catalytic reaction in favor of the Suzuki-Miyaura product, use of a boronic ester ligand derived from acenaphthoquinone allows the process to favor the conjunctive product, even with substituted substrates.

Graphical Abstract

Configurationally-defined benzhydryl stereocenters are important structural motifs that appear in a broad array of natural products and therapeutic agents.¹ Accordingly, a variety of catalytic methods have been developed to target their construction.² While recent advances in benzylic cross-coupling have provided important tools to target these features, an added synthetic challenge arises when benzylic stereocenters are sited adjacent to additional stereogenic centers. In these situations, multistep organic synthesis is often required for construction of the stereochemical dyad of interest.³ Our group has been developing a catalytic conjunctive cross-coupling reaction that converts vinylboron “ate” complexes and electrophiles to enantiomerically-enriched secondary or tertiary alkylboronic esters bearing a single stereocenter (eq. 1, Scheme 1).⁴ To address the problem of benzhydryl construction as outlined above, we questioned whether β -substituted alkenylboronic esters might engage in conjunctive cross-coupling and deliver compounds that bear vicinal stereogenic centers (eq. 2). In this report, we describe the development of this process and provide insight about how

*Corresponding Author: morken@bc.edu.

Supporting Information. Procedures, characterization and spectral data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

The authors declare no competing financial interest.

the structure of boron ligands can tip the reaction outcome in favor of the conjunctive coupling product or the classic Suzuki-Miyaura product.

Preliminary efforts to employ β -substituted alkenyl boronates in conjunctive cross-coupling reactions revealed a process that is dominated by direct Suzuki-Miyaura cross-coupling. For instance, in contrast to the conjunctive cross-coupling with the unsubstituted vinylB(neo)-derived “ate” complex **1a** which occurs in 84% yield and 97:3 er (Scheme 1, eq. 3), when styrenylB(neo) was converted to the derived “ate” complex **1b** and subjected to coupling with *p*-anisyltriflate, 1 mol% Pd(OAc)₂ and 1.2 mol% Mandyphos (**L**), only 13% of the conjunctive coupling product **2b** was obtained, with stilbene derivative **3b** being the predominant product. The disparate reaction outcomes with **1a** and **1b** as substrate may be understood by consideration of competing catalytic reaction pathways. Mechanistic studies intimate that the metal-induced metallate rearrangement which underlies the conjunctive coupling occurs by metal-alkene binding (**A**, equation 1, box),^{4f} followed by simultaneous R_M migration and C-Pd bond formation at C_β. In contrast, recent studies of transmetalation from organoboronic acid derivatives to bis(phosphine)Pd complexes suggests that the direct Suzuki-Miyaura reaction could arise by either (a) an open transition state originating from a palladium alkene complex where C-Pd bond formation occurs at C_α with concomitant rupture of the C-B bond (**B**, shaded box, Scheme 1)⁵, or (b) association of Pd with a boronic ester oxygen (**C**), which is then followed by transmetalation reaction through a closed four-centered transition state involving a five-coordinate Pd complex.⁶ For the reaction of substrate **1b**, the added substitution at the β carbon of the alkene (R=H→Ph) would serve to inhibit conjunctive coupling as it requires Pd-C bond formation at a more hindered site. However, transmetalation through either **B** or **C**, ultimately leading to C-Pd bond formation at C_α, would be relatively unaltered by the addition of β -substituents on the alkenyl group. Thus, the net effect of alkene β -substitution is to disfavor the metallate shift relative to the direct transmetalation reaction and thereby alter the course of the reaction. Following this analysis, it was considered that alternate ligand sets on the boron atom that selectively interfere with either Pd(II)-oxygen binding or that hinder access to C_α might reestablish the metallate shift as a major reaction pathway.

To investigate the effect of boron ligands on the chemoselectivity of the coupling reaction in equation 2, we converted a series of styrenyl boronic esters to the derived “ate” complexes and subjected them to coupling with Pd(OAc)₂ and MandyPhos ligand (*S_p*,*S_{p'}*-**L**). As depicted in Table 1, with relatively unencumbered ligands on boron such as **L1** (neopentyl glycol) or **L2** (*cis*-1,2-cyclopentane diol) the Suzuki-Miyaura product remains the predominant reaction product. However, when the boron center is more encumbered such as with **L3** (pinacol) and **L4**, the proportion of conjunctive cross-coupling (**C3**) product was increased, consistent with the notion that hindered access to either the oxygen atom or C_α may retard the rate of the direct transmetalation reaction relative to metallate shift. It can also be noted that chemoselectivity with boron centers bearing even more highly encumbered ligands (**L5**, **L6**) begins to suffer, an outcome we consider might be due to hindered olefin binding. Lastly, support for the notion that the oxygen atom may be involved in the direct transmetalation process is provided by the reaction with hydrocarbon-based

ligand **L7** (9-BBN) wherein the metallate shift-based path operates to the near exclusion of the Suzuki-Miyaura reaction.

Considering the observations in entries 1–6 of Table 1, we sought ligands that are both readily accessible and also effectively shield the oxygen atom and/or C $_{\alpha}$ without blocking access to the alkene C $_{\beta}$. As shown in entry 8, ligand **L8** (termed “mac”), readily prepared by methylation of acenaphthoquinone (*vide infra*), appears to satisfy these requirements, delivering the conjunctive product in good yield and enantioselectivity. Of note, the reaction chemoselectivity could be enhanced by conducting the reaction at 40 °C and in the presence of CsF; under these conditions the conjunctive coupling product is formed in 76% isolated yield, >20:1 dr, and with 99:1 enantioselectivity.

Diol **4** is readily available from acenaphthoquinone by a single-step diastereoselective (4.3:1 dr) carbonyl addition reaction employing trimethylaluminum in toluene,⁷ followed by crystallization of the diol from ethyl acetate solvent. Conversion of boronic acids to derived B(mac) esters is readily accomplished by esterification in the presence of catalytic FeCl₃.⁸ In contrast to common boronic ester ligands (i.e. pinacol, neopentylglycol), when the B(mac) derivatives are converted to the four coordinate “ate” complexes, the issue of stereoisomerism arises and it was considered that this feature might pose a complication for selective and efficient catalytic coupling reactions. To study the properties of B(mac) “ate” complexes, the addition of phenyllithium to *n*-butylB(mac) in THF-d₈ at room temperature was analyzed by ¹H NMR analysis. As shown in Scheme 2b, the kinetic addition product appeared to arise from addition of the nucleophile *cis* relative to the vicinal methyl groups forming *trans*-**6** in a 5:1 ratio.⁹ Of note, stirring the reaction for 30 minutes at 60 °C resulted in a 2:1 diastereomer ratio, and after 12 hours an equilibrium 1:1 ratio of the isomers was observed. Importantly, when reaction partners are reversed such that of *n*-BuLi was added to PhB(mac), an inverted initial kinetic ratio (ca. 1:12.5) results, but the same thermodynamic ratio is achieved upon heating (see Supporting Information for details). These experiments suggest that diastereomeric “ate” complexes should be interconverting during the course of reaction and that the configuration of these species will likely be inconsequential to conjunctive couplings.

With ready access to B(mac)-derived substrates and an understanding of their properties, these compounds were examined in conjunctive coupling reactions. As depicted in Table 2, the reaction can operate with both electron-rich and electron-poor electrophiles and, after oxidative work-up, provides the derived secondary alcohols with high diastereoselectivity (>20:1 dr) and enantioselectivity (generally above 98:2 er.). While substrates with heteroatom-containing functional groups do not pose an inherent problem, diminished yield was observed with electron-deficient electrophiles, an outcome that arises from competitive Suzuki-Miyaura coupling reactions (in general for Table 2, the remaining mass balance is comprised of Suzuki-Miyaura products). Fortunately, even with these substrates the stereoinduction remains high. It should also be noted that a substrate with a hindered migrating alkenyl group, reacted with anomalously low enantioselectivity, an observation that has been made in other systems.^{4a,c} Examination of both aryl triflate and aryl bromide electrophiles showed that both electrophiles could be employed to furnish the conjunctive product; however, bromide electrophiles required the addition of KOTf to ameliorate the

inhibitory activity of bromide salts^{4b} and provided diminished product yields relative to their triflate counterparts. The reaction was also found to operate with several different migrating groups, with electron-rich migrating groups giving higher yields than those that are electron-poor. Surprisingly, under the current conditions, “ate” complexes containing alkyl migrating groups did not engage in the conjunctive coupling pathway and provided direct Suzuki-Miyaura coupling products instead. With respect to the substituent at the β -carbon of the alkenyl boronate, substrates containing electron-rich and electron-deficient arenes, as well as those containing alkyl groups were found to participate in the reaction. While an electron-rich β -substituent appeared to increase the conversion of the reaction, it also resulted in more competition from the Suzuki-Miyaura pathway (**23**; 1:1 conjunctive:Suzuki). Substrates with electron-deficient β -substituents furnished less Suzuki-Miyaura reaction, but required higher temperatures for the conjunctive process to proceed (**24**, 4:1 conjunctive:Suzuki). Lastly, while substrates with β -alkyl substitution (**16-22**) gave only modest yields, they reacted with outstanding selectivity and furnished products that would be otherwise difficult to access with single-step catalytic processes.

Although B(mac)-derived products are stable to column chromatography, they are poorly soluble in many organic solvents such that for the purposes of Table 2, it was most convenient to oxidize the coupling products and isolate the corresponding alcohols. To learn about other transformations of the B(mac) derivatives, a larger scale coupling reaction was undertaken and the reaction product examined. As shown in Scheme 3, a gram-scale coupling provided secondary boronic ester **26** with reaction efficiency and selectivity comparable to smaller scale reactions in Table 2. Consistent with the observations in Table 2, purified intermediate **26** underwent oxidation efficiently, providing alcohol **2b** in outstanding yield and selectivity. Also noteworthy, is that direct amination¹⁰/Boc protection of **26** furnished carbamate **27** in excellent yield and as a single diastereomer. Lastly, it was found that B(mac)-derivate **26** underwent efficient modified Matteson homologation¹¹ to furnish primary B(mac) derivative **28** as a single diastereomer.

The diastereoselective conjunctive coupling establishes strategically-useful functional group arrays in ways that are not straightforward to access by contemporary asymmetric synthesis. This feature is illustrated by the application of the conjunctive coupling to the synthesis of obtusafuran¹² (Scheme 4). While this target was previously prepared by a convenient enantioselective ketone reduction and cyclization, construction of the requisite ketone starting material required five steps of chemical synthesis.¹³ Employing *trans*-2-propenylB(mac) in a conjunctive coupling with PhLi and aryl bromide **29**, furnished **30** in excellent yield and stereoselection. Subsequent oxidative cyclization¹⁴ and deprotection provided the target in just three synthesis steps (two reaction vessels) from simple starting materials.

In summary, we have established a catalytic, diastereo-, and enantioselective conjunctive coupling of β -substituted alkenylboronic esters. This process employs an encumbered diolato ligand to control the reaction of alkenylboron “ate” complexes, tipping the reaction in favor of a metallate shift-based pathway rather than direct transmetalation. Further studies on the mechanistic origin of chemoselectivity with B(mac) derived “ate” complexes will be reported in due course.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGEMENTS

This work was supported by the NIH (R35-GM1217140). We thanks Solvias for a generous donation of MandyPhos ligands.

REFERENCES

- (1). Ameen D; Snape TJ, Chiral 1,1-Diaryl Compounds as Important Pharmacophores. *Med. Chem. Commun* 2013, 4, 893.
- (2). For a review: see:Mondal S; Roy D; Panda G, Overview on the Recent Strategies for the Enantioselective Synthesis of 1, 1-Diarylalkanes, Triarylmethanes and Related Molecules Containing the Diarylmethine Stereocenter. *ChemCatChem* 2018, 10, 1941. For selected catalytic reactions, see:Imao D; Glasspoole BW; Laberge VS; Crudden CM, Cross Coupling Reactions of Chiral Secondary Organoboron Esters With Retention of Configuration. *J. Am. Chem. Soc* 2009, 131, 5024. [PubMed: 19301820] Zhou Q; Srinivas HD; Dasgupta S; Watson MP, Nickel-Catalyzed Cross-Couplings of Benzylic Pivalates with Arylboroxines: Stereospecific Formation of Diarylalkanes and Triarylmethanes. *J. Am. Chem. Soc* 2013, 135, 3307. [PubMed: 23425080] Harris MR; Hanna LE; Greene MA; Moore CE; Jarvo ER, Retention or Inversion in Stereospecific Nickel-Catalyzed Cross-Coupling of Benzylic Carbamates with Arylboronic Esters: Control of Absolute Stereochemistry with an Achiral Catalyst. *J. Am. Chem. Soc* 2013, 135, 3303. [PubMed: 23414579] Lewis CA; Gustafson JL; Chiu A; Balsells J; Pollard D; Murry J; Reamer RA; Hansen KB; Miller SJ, A Case of Remote Asymmetric Induction in the Peptide-Catalyzed Desymmetrization of a Bis(phenol). *J. Am. Chem. Soc* 2008, 130, 16358. [PubMed: 19006301] Gao F; Lee Y; Mandai K; Hoveyda AH, Quaternary Carbon Stereogenic Centers through Copper-Catalyzed Enantioselective Allylic Substitutions with Readily Accessible Aryl- or Heteroarylithium Reagents and Aluminum Chlorides. *Angew. Chem. Int. Ed* 2010, 49, 8370.
- (3). Examples: Tapentadol:Zhang Q; Li J-F; Tian G-H; Zhang R-X; Sun J; Suo J; Feng X; Fang D; Jiang X-R; Shen J-S, A Practical and Enantioselective Synthesis of Tapentadol. *Tetrahedron Asymm.* 2012, 23, 577. Taranabant:Wallace DJ; Campos KR; Shultz CS; Klapars A; Zewge D; Crump BR; Phenix BD; McWilliams JC; Krska S; Sun Y; Chen CY; Spindler F, New Efficient Asymmetric Synthesis of Taranabant, a CB1R Inverse Agonist for the Treatment of Obesity. *Org. Process Res. Dev* 2009, 13, 84. Mixanpril:Turcaud S; Gonzalez W; Michel J; Roques BP; Fournie-Zaluski C, Diastereoselective Synthesis of Mixanpril, an Orally Active Dual Inhibitor of Neutral Endopeptidase and Angiotensin Converting Enzyme. *Bioorg. Med. Chem. Lett* 1995, 5, 1893. Paroxetine:Zhang Y; Liao YT; Liu XH; Yao Q; Zhou YH; Lin LL; Feng XM, Catalytic Michael/Ring-Closure Reaction of α,β -Unsaturated Pyrazoleamides with Amidomalonates: Asymmetric Synthesis of (-)-Paroxetine. *Chem. Eur. J* 2016, 22, 15119. [PubMed: 27576747]
- (4). Zhang L; Lovinger GJ; Edelstein EK; Szymaniak AA; Chierchia MP; Morken JP, Catalytic Conjunctive Cross-Coupling Enabled by Metal-Induced Metallate Rearrangement. *Science*, 2016, 351, 70. [PubMed: 26721996] Lovinger GJ; Aparece MD; Morken JP, Pd-Catalyzed Conjunctive Cross-Coupling between Grignard-Derived Boron "Ate" Complexes and C(sp²) Halides or Triflates: NaOTf as a Grignard Activator and Halide Scavenger. *J. Am. Chem. Soc* 2017, 139, 3153. [PubMed: 28161943] Edelstein EK; Namirembe S; Morken JP, Enantioselective Conjunctive Cross-Coupling of Bis(alkenyl)borates: A General Synthesis of Chiral Allylboron Reagents. *J. Am. Chem. Soc* 2017, 139, 5027. [PubMed: 28335597] Chierchia MP; Law C; Morken JP, Nickel-Catalyzed Enantioselective Conjunctive Cross-Coupling of 9-BBN Borates. *Angew. Chem. Int. Ed* 2017, 56, 11870. Lovinger GJ; Morken JP, Ni-Catalyzed Enantioselective Conjunctive Coupling with C(sp³) Electrophiles: A Radical-Ionic Mechanistic Dichotomy. *J. Am. Chem. Soc* 2017, 139, 17293. [PubMed: 29116784] Myhill JA; Zhang L; Lovinger GA; Morken JP, Enantioselective Construction of Tertiary Boronic Esters by Conjunctive Cross-Coupling. *Angew. Chem. Int. Ed* 2018, 57, 12799. For recent aligned studies on electrophile-induced metallate rearrangement, see the following citations and references cited therein: Wilson

- CM; Ganesh V; Noble A; Aggarwal VK, Enantiospecific sp^2 – sp^3 Coupling of *ortho*- and *para*-Phenols with Secondary and Tertiary Boronic Esters. *Angew. Chem. Int. Ed* 2017, 56, 16318. Ganesh V; Odachowski M; Aggarwal VK, Alkynyl Moiety for Triggering 1,2-Metallate Shifts: Enantiospecific sp^2 – sp^3 Coupling of Boronic Esters with *p*-Arylacetylenes. *Angew. Chem. Int. Ed*, 2017, 56, 9752. Wang Y; Noble A; Sandford C; Aggarwal VK, Enantiospecific Trifluoromethyl-Radical-Induced Three-Component Coupling of Boronic Esters with Furans. *Angew. Chem. Int. Ed*, 2017, 56, 1810. Panda S; Ready JM, Tandem Allylation/1,2-Boronate Rearrangement for the Asymmetric Synthesis of Indolines with Adjacent Quaternary Stereocenters. *J. Am. Chem. Soc.*, 2018, 140, 13242. [PubMed: 30240211]
- (5). Ortuño MA; Lledós A; Maseras F; Ujaque G, The Transmetalation Process in Suzuki–Miyaura Reactions: Calculations Indicate Lower Barrier via Boronate Intermediate. *ChemCatChem* 2014, 6, 3132.
- (6). Huang Y-L, Weng C-M, and Hong F-E, Density Functional Studies on Palladium-Catalyzed Suzuki–Miyaura Cross-Coupling Reactions Assisted by N- or P-Chelating Ligands. *Chem. Eur. J* 2008, 14, 4426. [PubMed: 18381730] (b) Note that studies by Denmark suggest transmetalation of diphosphine complexes related to B may occur by dissociation of a phosphine in order to access a transition state involving a four-coordinate Pd-complex, see: Thomas AA; Wang H; Zahrt AF; Denmark SE, Structural, Kinetic, and Computational Characterization of the Elusive Arylpalladium(II)boronate Complexes in the Suzuki–Miyaura Reaction. *J. Am. Chem. Soc* 2017, 139, 3805. [PubMed: 28266847] With monophosphine Pd complexes, transmetalation through B–O–Pd bonded four-coordinate structures appears to predominate, see: Thomas AA; Zahrt AF; Delaney CP; Denmark SE, Elucidating the Role of the Boronic Esters in the Suzuki–Miyaura Reaction: Structural, Kinetic, and Computational Investigations. *J. Am. Chem. Soc* 2018, 140, 4401. [PubMed: 29543441] Thomas AA; Denmark SE, Pre-transmetalation Intermediates in the Suzuki–Miyaura Reaction Revealed: The Missing Link. *Science* 2016, 352, 329. [PubMed: 27081068] Carrow BP; Hartwig JF, Distinguishing Between Pathways for Transmetalation in Suzuki–Miyaura Reactions. *J. Am. Chem. Soc* 2011, 133, 2116. [PubMed: 21280669] Amatore C; Jutand A; Le Duc G, Kinetic Data for the Transmetalation/Reductive Elimination in Palladium-Catalyzed Suzuki–Miyaura Reactions: Unexpected Triple Role of Hydroxide Ions Used as Base. *Chem. Eur. J* 2011, 17, 2492. [PubMed: 21319240] Matos K; Soderquist JA, Alkylboranes in the Suzuki–Miyaura Coupling: Stereochemical and Mechanistic Studies. *J. Org. Chem* 1998, 63, 461. [PubMed: 11672034]
- (7). Baidossi W; Rosenfeld A; Wassermann BC; Schutte S; Schumann H; Blum J, [(3-Dimethylamino)propyl]dimethylaluminum: A Convenient Reagent for Methylation and Ethynylation of Carbonyl Compounds. *Synthesis* 1996, 9, 1127.
- (8). Wood JL; Marciasini LD; Vaultier M; Pucheault M, Iron Catalysis and Water: A Synergy for Refunctionalization of Boron. *Synlett* 2014, 25, 0551.
- (9). The configuration of the “ate” complexes was secured by both NOE and chemical shift analysis. See Supporting Information for details.
- (10). Mlynarski SN; Karns AS; Morken JP, Direct Stereospecific Amination of Alkyl and Aryl Pinacol Boronates. *J. Am. Chem. Soc* 2012, 134, 16449. [PubMed: 23002712]
- (11). Sonawane RP; Jheengut V; Rabalakos C; Larouche-Gauthier R; Scott HK; Aggarwal VK, Enantioselective Construction of Quaternary Stereogenic Centers from Tertiary Boronic Esters: Methodology and Applications. *Angew. Chem. Int. Ed* 2011, 50, 3760.
- (12). Gregson M; Ollis WD; Redman BT; Sutherland IO; Dietrichs HH; Gottlieb OR, Obtusastylene and Obtustylene, Cinnamylphenols from *Dalbergia Retusa*. *Phytochemistry* 1978, 17, 1395.
- (13). Chen C; Weisel M, Concise Asymmetric Synthesis of (+)-Conocarpan and Obtusafuran. *Synlett* 2013, 24, 189.
- (14). (a) Wang X; Lu Y; Dai H-X; Yu J-Q, Pd(II)-Catalyzed Hydroxyl-Directed C–H Activation/C–O Cyclization: Expedient Construction of Dihydrobenzofurans. *J. Am. Chem. Soc.*, 2010, 132, 12203. [PubMed: 20715820] (b) Wang H; Li G; Engle KM; Yu J-Q; Davies HML, Sequential C–H Functionalization Reactions for the Enantioselective Synthesis of Highly Functionalized 2,3-Dihydrobenzofurans. *J. Am. Chem. Soc.*, 2013, 135, 6774. [PubMed: 23600441]

Prior work: **via:** **This work:** **Preliminary observation:** **prospective complexes leading to transmetalation:** **Scheme 1.**Conjunctive Cross-Coupling of β -Substituted Alkenylboronates

Scheme 2.

Methylated Acenaphthoquinone (mac) as a Ligand for Boronic Esters.

Scheme 3.

Synthetic Transformations of AlkylB(mac) Derivatives.

Scheme 4.

Construction of (+)-Obtusafuran by Consecutive Cross-Coupling.

Table 1.

Effect of Boron Ligand on Chemoselectivity in Consecutive Coupling Reactions.^a

entry	BL ₂	C3:SM	C3 yield	dr	er
1	L1 (neo)	1:5.8	13 (10)	>20:1	99:1
2	L2	1:>20	<5	nd	nd
3	L3 (pin)	1:2	35(30)	>20:1	98:2
4	L4	1.7:1	56(46)	>20:1	99:1
5	L5	1:2	30	>20:1	nd
6	L6	1:3	20	>20:1	nd
7	L7 (9-BBN)	>20:1	92(75)	>20:1	67:33
8	L8 (mac)	2.5:1	75(70)	>20:1	99:1
9 ^b	L8 (mac)	4.2:1	83(76)	>20:1	99:1

(a) Yields are by ¹H NMR versus an internal standard, yield in parentheses is after isolation. For entry 7, yield is of the derived alcohol; for the others, yield is of the boronic ester. Enantiomer ratio of derived alcohol was determined by SFC analysis on a chiral stationary phase and in comparison to authentic enantiomer mixture. Diastereomer ratio determined by analysis of the ¹H NMR spectrum.

(b) Reaction at 40 °C and with 1 equiv CsF.

Table 2.

Pd/MandyPhos-Catalyzed Conjugative Coupling of β -Substituted B(mac)-Derived “Ate” Complexes.^a

(^a) Yields are isolated yields of purified material and represent average yield of two different experiments. Enantiomer ratios were determined by SFC analysis on a chiral stationary phase and are in comparison to authentic racemic materials. Diastereoselectivity determined by analysis of the ¹H NMR spectrum. For reactions of organobromide electrophiles, an equivalent of KOTf was added.

(^b) Reaction at 60 °C.

(^c) Reaction conducted with 2% Pd and 2.2% L1.