Cochrane
Library

Cochrane Database of Systematic Reviews

Reduced salt intake for heart failure (Protocol)

Mahtani KR, Heneghan CJ, Nunan D, Onakpoya IJ, Roberts NW, Hobbs FDR

Mahtani KR, Heneghan CJ, Nunan D, Onakpoya IJ, Roberts NW, Hobbs FDR.

Reduced salt intake for heart failure.

Cochrane Database of Systematic Reviews 2018, Issue 2. Art. No.: CD011214.

DOI: 10.1002/14651858.CD011214.pub2.

www.cochranelibrary.com

TABLE OF CONTENTS

HEADER	1
REASON FOR WITHDRAWAL	1
WHAT'S NEW	1
SOURCES OF SUPPORT	1

[Intervention Protocol]

Reduced salt intake for heart failure

Kamal R Mahtani¹, Carl J Heneghan¹, David Nunan¹, Igho J Onakpoya¹, Nia W Roberts², FD Richard Hobbs¹

¹Nuffield Department of Primary Care Health Sciences, University of Oxford, Oxford, UK. ²Bodleian Health Care Libraries, University of Oxford, Oxford, UK

Contact address: Kamal R Mahtani, Nuffield Department of Primary Care Health Sciences, University of Oxford, New Radcliffe House, Radcliffe Observatory Quarter, Oxford, Oxfordshire, OX2 6GG, UK. kamal.mahtani@phc.ox.ac.uk.

Editorial group: Cochrane Heart Group.

Publication status and date: Withdrawn from publication for reasons stated in the review, published in Issue 2, 2018.

Citation: Mahtani KR, Heneghan CJ, Nunan D, Onakpoya IJ, Roberts NW, Hobbs FDR. Reduced salt intake for heart failure. *Cochrane Database of Systematic Reviews* 2018, Issue 2. Art. No.: CD011214. DOI: 10.1002/14651858.CD011214.pub2.

Copyright © 2018 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

The editorial group responsible for this previously published document have withdrawn it from publication.

REASON FOR WITHDRAWAL

This protocol has been withdrawn as the author team is unable to progress to the final review stage.

WHAT'S NEW

Date	Event	Description
22 February 2018	Amended	This protocol has been withdrawn as the author team is unable to progress to the final review stage

SOURCES OF SUPPORT

Internal sources

- National Institute for Health Research (NIHR) School for Primary Care Research (SPCR), UK.

A proportion of this project is funded by the National Institute for Health Research (NIHR) School for Primary Care Research (SPCR).

Disclaimer:

This article presents independent research part-funded by the National Institute for Health Research (NIHR). The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health.

External sources

- Royal College of General Practitioners, UK.

This research was part-funded by the Scientific Foundation board of the Royal College of General Practitioners (Grant number SFB 2013-14)