Published in final edited form as:

Exerc Sport Sci Rev. 2019 April ; 47(2): 66–74. doi:10.1249/JES.0000000000000182.

Exercise and Vascular Insulin Sensitivity in Skeletal Muscle and Brain

T. Dylan Olver¹, M. Harold Laughlin^{2,3}, and Jaume Padilla^{3,4,5}

¹Dept. Veterinary Biomedical Sciences, University of Saskatchewan, SK. Canada

²Dept. Biomedical Sciences, University of Missouri, Columbia, MO. USA

³Dalton Cardiovascular Research Center, University of Missouri, Columbia, MO, USA

⁴Dept. Nutrition and Exercise Physiology, University of Missouri, Columbia, MO, USA

⁵Dept. Child Health, University of Missouri, Columbia, MO, USA

Abstract

We present the hypothesis that exercise-induced hyperemia, perhaps through vascular shear stress, represents an important factor responsible for the effects of physical activity (PA) on vascular insulin sensitivity. Specifically, we postulate PA involving the greatest amount of skeletal muscle mass and the greatest central neural recruitment maximizes perfusion and consequently enhances vascular insulin sensitivity in skeletal muscle and brain.

Summary for TOC:

Physical activity-induced improvements in vascular insulin sensitivity in the skeletal muscle and brain are likely related to the exercise hyperemic response.

Keywords

Vascular insulin signaling; Skeletal muscle; brain; Physical Activity; exercise; blood flow; shear stress

INTRODUCTION

In 1939 Abramson and colleagues examined the vascular actions of insulin shock therapy and reported that insulin-induced hypoglycemia resulted in increased peripheral limb blood flow in patients with Schizophrenia (1). Shortly thereafter, Ferris and co-workers (2) and later Porta and co-workers (3) demonstrated insulin-induced hypoglycemia either had no effect or a small vasodilatory effect on intracranial/cerebral blood flow in patients with Schizophrenia. Granted these initial observations do not reflect non-pathological conditions,

Corresponding Author: M. Harold Laughlin, Ph.D, Curator's Professor, Department of Biomedical Sciences, Rm. W117, Veterinary Medicine Building, University of Missouri, Columbia, MO. 65211, Phone: 573-884-4391, Fax: 573-884-6890, laughlinm@missouri.edu.

Conflict of Interest: None

these findings highlight the earliest direct evidence of insulin-stimulated vasodilation in the peripheral vasculature.

Since then, researchers have validated and extended upon earlier findings by demonstrating that physiological and pharmacological doses of exogenous insulin stimulate peripheral vasodilation during euglycemia (4–8) and by providing evidence that endogenous insulin also stimulates vasodilation (9–12). Notably, the magnitude of insulin-stimulated peripheral vasodilation appears blunted in the setting of various insulin resistant states in humans (i.e., obesity and type 2 diabetes; T2D) (9,11,13–15) and in experimental animal models (diet-induced and genetic models of T2D as well as insulin-treated T1D) (16–25). There is growing consensus that insulin-stimulated vasodilation is a physiologically relevant phenomenon and depressed insulin-stimulated vasodilation serves as a hallmark of vascular dysfunction (21,26–29). Being physically active or participating in structured exercise training can improve insulin-stimulated vasodilation in the setting of health and disease, with such improvements occurring most readily in tissue regions that undergo a relative increase in activity during the transition from rest to exercise (21,29). Herein, we address the hypothesis that exercise-induced improvements in vascular insulin sensitivity occur in a region-specific manner within skeletal muscle and cerebral vasculatures and in relation to local increases in blood flow during exercise (Figure 1).

Brief overview of endothelial insulin signaling

Available evidence indicates the peripheral hemodynamic effects of insulin are mediated by a combination of neurohumoral (6,8,30–34) and endothelial vasodilator and vasoconstrictor signals (19,21,26,28,29,35–37). This review will focus primarily on the effects of insulin on the vascular endothelium and endothelial mechanisms through which exercise training exerts therapeutic benefits on vascular insulin sensitivity.

At the molecular level, insulin binds to endothelial insulin receptor substrate-1 (IRS-1), which activates two primary signaling cascades, the phosphatidylinositol-4,5-bisphosphate 3-kinase (PI3K)/protein kinase B (Akt)/nitric oxide (NO) pathway, and the Ras/mitogen activated protein kinase (MAPK)/endothelin-1 (ET-1) pathway (26,37) (Figure 2). Activation of the former pathway induces anti-atherogenic and vasodilator signaling, whereas activation of latter pathway induces pro-atherogenic and vasoconstrictor signaling. Under healthy circumstances (in many vascular beds/arteries) the net effect of insulin administration is a NO-dependent vasodilation suggesting activation of the PI3K/Akt/NO pathway predominates (26,28,37).

Selective vascular insulin resistance refers to a shift in endothelial insulin signaling reflected by decreased IRS-1-induced activation of PI3K/Akt/NO with no change or increased activation of MAPK/ET-1 (26,37). Thus, the imbalance promotes pro-atherogenic and ET-1 dependent vasoconstrictor signaling (16,17). Current evidence suggests selective vascular insulin resistance may occur early in the development of metabolic derangement (11,18) and is implicated in decreased elasticity and progression of atherosclerosis in conduit arteries, impaired blood flow control, reduced capillary perfusion and nutrient delivery, and limited trans-endothelial nutrient/insulin transport within specified tissues/organs (21,26–29). The underlying causes of vascular insulin resistance are likely multifactorial. In the setting of

insulin deficiency or insulin resistance, lipotoxicity, glucotoxicity, inflammation and oxidative stress may all serve a role in directly and indirectly impairing insulin signaling (26,37,38). For example, the aforementioned metabolic perturbations are implicated in elevating endothelial cell diacylglycerol which activates protein kinase C isoforms and subsequently reduces tyrosine phosphorylation of IRS-1, decreasing the downstream activation of the Akt and NO pathway. Further, hyperinsulinemia may also enhance MAPK activation directly which can inhibit subsequent IRS-1 activation and downstream signaling (26,37). Indeed, published animal work shows obesity/T2D or palmitic acid-induced deficits in insulin-stimulated vasodilation in isolated arterioles can be reversed acutely with inhibition of protein kinase C beta (18) or theta (39) as well as ET-1A receptor inhibition (40).

Routine exercise consistently enhances vascular insulin sensitivity (for review see (21,29)). Initial cross sectional data reveal insulin-stimulated vasodilation in the lower limb is greater in endurance trained athletes compared to otherwise healthy sedentary controls (41). In healthy control rodents and rodents with insulin-treated type 1 diabetes, improvements in insulin-stimulated vasodilation following endurance training are associated with increases in endothelial NO synthase protein (20). Of note, in the latter study, insulin treatment was adjusted bi-weekly to equalize hyperglycemia/blood glucose concentrations for all rats with type 1 diabetes, highlighting the therapeutic effect of endurance training occurred independent of chronic improvements in hyperglycemia. Further, when comparing wheel-running with metformin or food restriction in obese/T2D rodents, although all treatments exerted similar beneficial effects on body composition and HbA1c, only wheel-running improved insulin-induced dilation in isolated skeletal muscle arterioles (22,42,43). Collectively, these data suggest the beneficial effect of regular exercise is conferred independent of fat loss and enhanced glycemic control and may be related to local or systemic signals subjected to the vasculature during exercise (Figure 1). In order to better understand the mechanisms responsible for the therapeutic benefits of acute and chronic exercise, we now discuss some of the signals experienced by the vasculature in response to exercise as well as some key chronic vascular adaptations to exercise training.

EXERCISE AND VASCULAR SHEAR STRESS – SKELETAL MUSCLE

Generally, skeletal muscle is grouped into three fiber types classified according to both their contractile and metabolic properties (44): slow-twitch oxidative (SO), fast-twitch, glycolytic (FG), and fast-twitch, oxidative, glycolytic (FOG). Skeletal muscle blood flow increases with exercise intensity and is distributed heterogeneously within and among skeletal muscles in relation to fiber type composition and recruitment during exercise (45–54). During the transition from rest to maximal exercise, skeletal muscle blood flow may increase up to a 10-fold and can reach up to 300–400 mL·100 g⁻¹·min⁻¹ (47,55,56). Of interest, the vasculature supplying skeletal muscle fibers that undergo a relative increase in activity during exercise may experience increased vascular shear stress (i.e., the frictional force of blood acting against the lumen wall) during exercise (47,55).

Vascular shear stress causes the deformation of a population of mechanoreceptors, that appear to be coupled to glycocalyx, located on the apical surface of endothelial cells. This in

turn activates the mechanosensitive PI3K/Akt/NO signaling pathway (55,57) (Figure 2). It is currently believed that acute or repeated exposure to exercise-induced increases in vascular shear stress, and by extension acute or repeated exposure to enhanced endothelial PI3K/Akt/NO signaling, is involved in changes in vasomotor control (i.e., improved endothelial-dependent vasodilation) (23,58,59) and vascular remodeling (reflected by an increase in the number, diameter and density of arterioles and capillaries) in active skeletal muscle (47,51,55,60–73). It is important to note that available evidence, predominantly from animals, suggests exercise training modulates vascular function and structure of arteries/arterioles in spatially-distinct (47,63,73–86), intensity-dependent manner (23,47,60–64,71) and adaptations tend to be concentrated in the muscle tissue that experiences the greatest relative increase in activity during training sessions (60–62,72,87–93). For example, both moderate intensity continuous endurance and high intensity interval training increase capillary density in animals (71) and humans (21,65,66); however, work in animals suggests such increases in capillary density following training occur predominantly in SO vs. FOG/FG muscle fibers, respectively (21,51,71,91,92). This phenomenon is perhaps not as clear in humans, as recent work from Cocks and colleagues (65) shows that while the increase in capillary density in the vastus lateralis of obese men tends to be greater following moderate intensity continuous (40–60 min·day⁻¹ at 65% $\dot{V}O_{2peak}$, 5 days·week⁻¹, for four weeks) than high intensity interval sprint (4–6 X repeat Wingate tests, interspersed with 4.5 min rest, 5 days·week⁻¹, for four weeks) cycling training (19 vs. 6%, respectively), training-induced increases in capillary-to-fiber ratio or capillary contacts per fiber were similar between training regimens and not fiber type specific. Of note, these results may be influenced by the skeletal muscle examined and muscle fibers sampled between groups as well as pre-/post-training, as the vastus lateralis contains SO, FOG and FG fibers and the depth and fiber composition of the muscle biopsy may impact the results (94,95).

Acute and chronic exercise – vascular insulin signaling (see Table)

In hallmark studies, Dela and colleagues (96,97) examined the effect of either one-legged cycling (30 min·day⁻¹ at 70% $\dot{V}O_{2peak}$, 6 days·week⁻¹, for ten weeks) (96) or resistance exercise training (leg press, knee extension and hamstring curls; 3–4 sets of 8–12 repetitions at ~70–80% 1 repetition-max, performed 3 days·week⁻¹, for 6 weeks) (97) on lower limb blood flow responses during a euglycemic hyperinsulinemic clamp in patients with T2D. They found that insulin-mediated vasodilation and glucose clearance were greater in the trained limb, but not the untrained limb, from pre- to post-training. They reported similar findings in young and old men following the same one-legged cycling program (98) and more recently, similar findings were documented following a single bout of one-legged cycling exercise (~50% peak work load for 60 minutes, with 3 X 5 minute intervals at 100% peak workload) (99). Eskelinen and coworkers (100) examined insulin-stimulated glucose uptake in the upper and lower limbs following either moderate intensity endurance (40–60 min·day⁻¹ at 60% $\dot{V}O_{2peak}$, 3 days·week⁻¹, for two weeks) or high intensity interval sprint (4–6 X repeat Wingate tests, 3 days·week⁻¹, for two weeks) cycling training and reported that training-induced improvements in skeletal muscle glucose uptake were restricted to the lower limbs (i.e. active skeletal muscle). Further, while both training programs improved glucose uptake in the vastus lateralis, intermedius and medialis, only interval sprint training

increased glucose uptake in the rectus femoris (100). The authors speculated that the latter observation related to different muscle activation patterns during exercise and the inability of moderate intensity endurance exercise to activate the rectus femoris (see the Table). The relationship between skeletal muscle fiber recruitment during exercise and vascular insulin sensitivity is further exemplified by work in obese rats with T2D that shows that training-induced improvements in insulin-mediated vasodilation of resistance arteries following moderate intensity continuous and high intensity interval training are most robust in red and white portions of the gastrocnemius muscle, respectively (23,43). Taken together, these findings highlight the following important concepts: 1) the beneficial effects of exercise training on insulin sensitivity are related to improved insulin-stimulated vasodilation; 2) improvements in insulin-mediated vasodilation occur in relation to skeletal muscle activation and skeletal muscle fiber recruitment within each muscle during exercise (i.e., in relation to the exercise hyperemic response); and 3) such improvements can be conferred with acute and chronic exercise as well as independently of long term improvements in body composition or metabolic status.

Shear stress and insulin signaling - crossing paths

Vascular shear stress and insulin-stimulated vasodilation are both reliant on activation of the PI3K/Akt/NO pathway (37,57) (Figure 2). Of note, the vascular insulin sensitizing effect of exercise appears to be restricted to portions of the vasculature that undergo a relative increase in blood flow during exercise (23,43,96,99,101). Further, recent human work reveals that improvements in insulin-stimulated lower limb microvascular perfusion and leg glucose uptake 4 hours following one legged cycling exercise are blunted (to non-exercised levels) with NO inhibition administered during the insulin clamp (99). Given exercise-related improvements in insulin signaling can occur independent of longstanding changes in metabolic status, are localized to active skeletal muscle and blunted by NO inhibition, this raises the possibility that vascular shear stress primes endothelial cells to become more insulin sensitive and enhance insulin-stimulated NO signaling. In support of this view, recently published work established that the insulin sensitizing effects of vascular shear stress are independent of skeletal muscle contraction (102). First, it was demonstrated in cultured human aortic endothelial cells that compared with cells exposed to low shear conditions (3 dynes/cm²), those exposed to 1 hour of increased shear stress (20 dynes/cm²) subsequently exhibited a shift in insulin signaling characterized by an increased activation of endothelial NO synthase relative to MAPK. Second, it was demonstrated in isolated porcine skeletal muscle resistance arteries that compared with arteries kept under no-flow conditions, those exposed to 1 hour of increased shear stress (20 dynes/cm²) subsequently displayed enhanced insulin-stimulated vasodilation. Prior shear stress only augmented insulin-stimulated vasodilation and not endothelium-independent vasodilation, suggesting the effect is conferred through an endothelial-dependent mechanism. Lastly, the translational relevance of these *in vitro* findings was validated by demonstrating that compared with the control leg, single leg heating resulting in increased blood flow, and likely attendant increase in shear stress, caused a subsequent augment in popliteal artery blood flow and calf microvascular perfusion in response to a systemic infusion of insulin. Collectively, these findings support the hypothesis that vascular shear stress may be a primary mechanism through which exercise enhances vascular insulin sensitivity.

Other vascular adaptations likely responsible for the effects of exercise training on insulin-stimulated glucose/insulin delivery and trans-endothelial insulin transport are increases in microvascular perfusion/capillary recruitment (99,103) and increased capillary volume or density (65,104–106). In this regard, an increased capillary density following moderate continuous aerobic training ($45 \text{ min} \cdot \text{day}^{-1}$ at $75\% \dot{V}O_{2\text{max}}$, $3 \text{ days} \cdot \text{week}^{-1}$, for six months) independently increases insulin sensitivity in older adults (104). Indeed, enhanced vascular density and volume following aerobic training (21) provides greater potential for delivery of nutrients and insulin to the target organ (Figure 1).

FUTURE PERSPECTIVES

Exercise and endothelial insulin signaling in the brain

In contrast to skeletal muscle, global brain blood flow increases only modestly ($\sim 0\text{--}20\%$) during incremental exercise until $\sim 60\% \dot{V}O_{2\text{max}}$ and may remain unchanged or decrease at higher intensities (likely the result of decreasing PaCO_2 values). However, evidence from miniature swine, dogs, ponies and humans reveals that, exercise increases regional brain blood flow in a structurally-specific and intensity-dependent manner (up to $\sim 70\%$ above basal values within the most active structures (47,55,107)). Further, recent human data indicates cerebral blood flow may increase in the recovery period immediately following high intensity exercise (108). Brain regions recruited at the onset and during sustained exercise include those involved in central command, motor execution, equilibrium, cardiorespiration, auditory, olfactory, and visual regions (47,55). Though much less established, work in humans and animals indicate cerebrovascular adaptations to exercise training may be both structurally- and intensity-dependent (55,109,110). With regards to the effects of exercise on vascular insulin signaling in the cerebrovasculature, recent work in obese rats with T2D reveals increased physical activity (wheel-running) improves insulin-mediated cranial/cerebral vasodilation and maintains cerebellum blood flow during insulin stimulation *in vivo*. Further, improvements in insulin-stimulated posterior cerebral artery vasodilation following wheel running were associated with an increased contribution of insulin-induced NO and decreased contribution of insulin-induced MAPK/ET-1 signaling (40). Of note, branches of the posterior cerebral artery supply the cerebellum and the cerebellum is believed to be active during exercise, highlighting (similar to skeletal muscle) exercise training-induced improvements in insulin-stimulated vasodilation may occur in relation to regional brain activation and corresponding blood flow responses during exercise (111–113).

Recent work in obese/T2D swine reveals impaired insulin-induced pial artery vasodilation coincides with depressed insulin-stimulated Akt signaling in the prefrontal cortex (18). The significance of insulin modulation of brain blood flow control and the role of insulin in central nervous system (CNS) function remains to be elucidated fully. However, it has become clearer that insulin signaling is critical to normal CNS function (e.g., insulin action in the brain is involved in improving memory & mood, inhibiting food intake, reducing bodyweight, increasing peripheral insulin sensitivity, reducing gluconeogenesis & lipolysis in the fasting state and increasing postprandial thermogenesis) and CNS insulin resistance is implicated in the abnormal regulation of behavior and neurocognitive impairment/

pathologies (114). The brain relies predominantly on insulin-independent glucose uptake and insulin stimulation has little effect on brain glucose metabolism in healthy individuals. However, insulin stimulation increases brain glucose uptake in individuals with impaired glucose tolerance (i.e., insulin-stimulated glucose uptake is maximal in the fasting state under healthy but not insulin resistance conditions) (115). In stark contrast to skeletal muscle, reduced insulin-stimulated brain glucose uptake may reflect improved metabolic/insulin sensitivity. Improvements in insulin sensitivity in the brain may be attributed to enhanced insulin transport across the blood brain barrier. Briefly, one mechanism through which trans-endothelial insulin transport is achieved is by binding to insulin receptors located on the luminal membrane of brain endothelial cells, becoming internalized through an endocytotic process and being shuttled across the endothelial cell. Insulin is then released and can interact with vascular smooth muscle cells, pericytes and astrocytes or enter into the brain interstitial fluid and interact directly with neurons (116). Recent work in mice highlights receptor-mediated trans-endothelial insulin transport in the microvasculature is NO-dependent (117) and represents a primary mechanism through which insulin uptake is achieved in the brain. Further, diet-induced obesity in mice impairs trans-endothelial insulin transport in the brain (118). Whether exercise improves trans-endothelial insulin transport in the brain and whether such improvements are related to exercise-induced vascular shear stress has not been examined.

Recently, it was demonstrated that high intensity interval sprint cycling training (4–6 X repeat Wingate tests, 3 days·week⁻¹, for two weeks), but not moderate intensity endurance cycling training (40–60 min·day⁻¹ at 60% $\dot{V}O_{2peak}$, 3 days·week⁻¹, for two weeks), reduced glucose uptake during insulin stimulation in the cerebellum, superior frontal gyrus, medial frontal gyrus, temporal cortex, thalamus, cingulate gyrus, and occipital cortex (119). The author's speculated the superior effect of high intensity interval sprint training may be the result of greater exercise-related neural recruitment, exercise-related changes in brain metabolism and possibly improvements in insulin transport across the blood brain barrier. Sprint exercise lasting 30 seconds (performed on a cycle ergometer similar to the repeat Wingate protocol above) is associated with a transient increase in the index of cerebral blood flow during the sprint, followed by a gradual decline to near or below baseline values by the end of the sprint and then a significant hyperemic response during the post-sprint recovery (lasting at least 60 seconds after the sprint) (108). Thus, it is conceivable the hemodynamic effects of high intensity interval exercise contribute to cerebrovascular adaptations to training that improve cerebrovascular insulin sensitivity (i.e., improved insulin-stimulated NO signaling, enhanced vascular density/volume following training and possibly improvements in trans-endothelial insulin transport). Indeed, mounting evidence from humans and animal models indicates regular exercise enhances cerebrovascular NO signaling, vascular volume and capillary density in a region specific and intensity-dependent manner within the brain (55,109,110). Further, it is speculated these adaptations are related to regional brain activation and vascular shear stress experienced during the exercise bout (55,109,110). Whether these exercise-induced signals or training-induced adaptations confer a benefit to CNS insulin signaling is an area for further study.

Vascular training

Aerobic and resistance exercise training improve vascular insulin sensitivity with improvements occurring in the vascular supply of skeletal muscles and possibly brain regions that undergo a relative increase in activity/blood flow during exercise (see the Table). We contend that the added benefit of including different exercise modalities and incorporating low, moderate and high intensity exercise is not simply ‘more is better’. Consider that different exercise modes and intensities require activation of different skeletal muscles, different fiber recruitment patterns and activation of different brain regions or populations of neurons within the same brain region. Therefore, employing different modes and intensities of exercise will modulate the spatial distribution of functional and structural vascular adaptations to exercise training in both the skeletal muscle and the brain. As such, we postulate that providing the duration of the stimulus is sufficient (i.e., one that exceeds a minimum threshold to induce adaptation), exercise programs designed to activate the most skeletal muscle and the most muscle fibers within each skeletal muscle (i.e., greatest relative increase in fiber recruitment during exercise) and require global, diverse and integrated neural output (i.e., learning and employing complex and coordinated motor/sensorimotor activation, employs elements of strategy, consists of repetitive and novel stimuli and engages multiple senses) will promote the most direct and widespread vascular adaptations. For this reason, we recommend modulating mode and intensity of exercise to maximize skeletal muscle fiber and CNS neural recruitment with the goal of generating the greatest enhancement in vascular insulin signaling in the most amount of vascular tissue (Figure 1).

Participation in recreational physical activity or structured exercise training are widely accepted to be beneficial for metabolic and neurocognitive function. Likewise, they are essential in maintaining or improving vascular insulin sensitivity; however, the underlying mechanisms remain unresolved. We speculate that exercise-induced improvements in vascular insulin signaling are conferred, at least in part, through exercise-induced vascular shear stress. Accordingly, going forward it is important that future research experimentally tests if indeed exercise-induced shear stress enhances vascular insulin sensitivity, as well as determine the molecular mechanisms by which this occurs. Additional unanswered questions are:

- to what extent the skeletal muscle and brain microvasculature is subjected to increased shear stress during exercise?
- are the molecular mechanisms by which exercise enhances micro- and macrovascular insulin sensitivity the same?
- do exercise-induced improvements in vascular insulin signaling contribute to improvements in metabolic and neurocognitive function?
- do the effects of exercise-induced vascular shear stress extend beyond the vasculature (i.e., does vascular NO production impart benefits in surrounding tissues?)?

These and other questions are critical in elucidating the underlying mechanisms responsible for the insulin sensitizing effects of exercise on the vasculature and essential for the development of prescriptive-based personalized exercise medicine.

Acknowledgments

Disclosure of Funding: J.P. is supported by National Institutes of Health (NIH) Grants K01-HL-125503 and R01-HL-137769. T.D.O. is supported by the Saskatchewan Health Research Foundation Establishment Grant.

REFERENCES

1. Abramson DI, Schkloven N, Margolis MN, Mirsky IA. Influence of massive doses of insulin on peripheral blood flow in man. *Am J Physiol.* 1939;128:124–32.
2. Ferris EB, Rosebaum M, Aring CD, Ryder HW, Roseman E, Hawkins JR. Intracranial blood flow in insulin coma. *Arch Neur Psych.* 1941;46(3):509–12.
3. Porta DP, Maiolo AT, Negri VU, Rossella E. Cerebral blood flow and metabolism in therapeutic insulin coma. *Metabolism.* 1964;13(2):131–40. [PubMed: 14127087]
4. Richter EA, Kiens B, Mizuno M, Strange S. Insulin action in human thighs after one-legged immobilization. *J Appl Physiol.* 1989;67(1):19–23. [PubMed: 2668254]
5. Richter EA, Mikines KJ, Galbo H, Kiens B. Effect of exercise on insulin action in human skeletal muscle. *J Appl Physiol.* 1989;66(2):876–85. [PubMed: 2496078]
6. Steinberg HO. Effects of insulin on the vascular system In: Johnstone MT, Veves A, editors. *Diabetes and cardiovascular disease.* Springer, Humana Press; 2001 p. 265–84.
7. Creager MA, Liang C-S, Coffman JD. Beta adrenergic-mediated vasodilator response to insulin in the human forearm. *J Pharmacol Exp Ther.* 1985;235(3):709–14. [PubMed: 3001273]
8. Liang C-S, Doherty JU, Faillace R, et al. Insulin infusion in conscious dogs. *J Clin Invest.* 1982;69:1321–36. [PubMed: 6123523]
9. Baron AD, Laakso M, Brechtel G, Hoit B, Watt C, Edelman S V. Reduced postprandial skeletal muscle blood flow contributes to glucose intolerance in human obesity. *J Clin Endocrinol Metab.* 1990;70(6):1525–33. [PubMed: 2189883]
10. Olver TD, Mattar L, Grisé KN, et al. Glucose-stimulated insulin secretion causes an insulin-dependent nitric oxide-mediated vasodilation in the blood supply of the rat sciatic nerve. *Am J Physiol Regul Integr Comp Physiol.* 2013;305(2):R157–63. [PubMed: 23616106]
11. Olver TD, Hazell TJ, Hamilton CD, Shoemaker JK, Lemon PWR. Impaired superficial femoral artery vasodilation and leg blood flow in young obese women following an oral glucose tolerance test. *Appl Physiol Nutr Metab.* 2012;37(1):176–83. [PubMed: 22300358]
12. Mikus CR, Fairfax ST, Libla JL, et al. Seven days of aerobic exercise training improves conduit artery blood flow following glucose ingestion in patients with type 2 diabetes. *J Appl Physiol.* 2011;111(3):657–64. [PubMed: 21737826]
13. Laakso M, Edelman S V, Brechtel G, Baron AD. Impaired insulin-mediated skeletal muscle blood flow in patients with NIDDM. *Diabetes.* 1992;41(9):1076–83. [PubMed: 1499861]
14. Laakso M, Edelman S V, Brechtel G, Baron AD. Decreased effect of insulin to stimulate skeletal muscle blood flow in obese man. A novel mechanism for insulin resistance. *J Clin Invest.* 1990;85(6):1844–52. [PubMed: 2189893]
15. Reynolds LJ, Credeur DP, Manrique C, Padilla J, Fadel PJ, Thyfault JP. Obesity, type 2 diabetes, and impaired insulin stimulated blood flow: role of skeletal muscle NO synthase and endothelin-1. *J Appl Physiol.* 2016;122(1):38–47. [PubMed: 27789766]
16. Eringa EC, Stehouwer CDA, Merlijn T, Westerhof N, Sipkema P. Physiological concentrations of insulin induce endothelin-mediated vasoconstriction during inhibition of NOS or PI3-kinase in skeletal muscle arterioles. *Cardiovasc Res.* 2002;56(3):464–71. [PubMed: 12445887]
17. Eringa EC, Stehouwer CDA, Roos MH, Westerhof N, Sipkema P. Selective resistance to vasoactive effects of insulin in muscle resistance arteries of obese Zucker (fa/fa) rats. *Am J Physiol Endocrinol Metab.* 2007;293(5):E1134–9. [PubMed: 17623751]
18. Olver TD, Grunewald ZI, Jurrissen TJ, et al. Microvascular insulin resistance in skeletal muscle and brain occurs early in the development of juvenile obesity in pigs. *Am J Physiol Regul Integr Comp Physiol.* 2017;314(2):R252–64. [PubMed: 29141949]

19. Olver TD, McDonald MW, Klakotskaia D, et al. A chronic physical activity treatment in obese rats normalizes the contributions of ET-1 and NO to insulin-mediated posterior cerebral artery vasodilation. *J Appl Physiol.* 2017;122(4):1040–50. [PubMed: 28183819]
20. Olver TD, McDonald MW, Grisé KN, et al. Exercise training enhances insulin-stimulated nerve arterial vasodilation in rats with insulin-treated experimental diabetes. *Am J Physiol Regul Integr Comp Physiol.* 2014;306(12):R941–50. [PubMed: 24740656]
21. Olver TD, Laughlin HM. Endurance, interval sprint training and resistance training; impact on microvascular dysfunction in type 2 diabetes. *Am J Physiol Heart Circ Physiol.* 2016;310:H337–50. [PubMed: 26408541]
22. Crissey JM, Padilla J, Jenkins NT, et al. Metformin does not enhance insulin-stimulated vasodilation in skeletal muscle resistance arteries of the OLETF rat. *Microcirculation.* 2013;20(8):764–75. [PubMed: 23879830]
23. Martin JS, Padilla J, Jenkins NT, et al. Functional adaptations in the skeletal muscle microvasculature to endurance and interval sprint training in the type 2 diabetic OLETF rat. *J Appl Physiol.* 2012;113(8):1223–32. [PubMed: 22923508]
24. Katakam PVG, Snipes JA, Steed MM, Busija DW. Insulin-induced generation of reactive oxygen species and uncoupling of nitric oxide synthase underlie the cerebrovascular insulin resistance in obese rats. *J Cereb Blood Flow Metab.* 2012;32(5):792–804. [PubMed: 22234336]
25. Meijer RI, Gray S, Aylor K, Barrett EJ. Pathways for insulin access to the brain: the role of the microvascular endothelial cell. *Am J Physiol Hear Circ Physiol.* 2016;311:H132–8.
26. King GL, Park K, Li Q. Selective insulin resistance and the development of cardiovascular diseases in diabetes: the 2015 Edwin Bierman Award lecture. *Diabetes.* 2016;65(6):1462–71. [PubMed: 27222390]
27. Kullmann S, Heni M, Hallschmid M, Fritsche A, Preissl H, Häring H-U. Brain insulin resistance at the crossroads of metabolic and cognitive disorders in humans. *Physiol Rev.* 2016;96:1169–209. [PubMed: 27489306]
28. Barrett EJ, Wang H, Upchurch CT, Liu Z. Insulin regulates its own delivery to skeletal muscle by feed-forward actions on the vasculature. *Am J Physiol Metab.* 2011;301(2):E252–63.
29. Padilla J, Olver TD, Thyfault JP, Fadel PJ. Role of habitual physical activity in modulating vascular actions of insulin. *Exp Physiol.* 2015;100(7):759–71. [PubMed: 26130183]
30. Vollenweider P, Randin D, Tappy L, Jequier E, Nicod P, Scherrer U. Impaired insulin-induced sympathetic neural activation and vasodilation in skeletal muscle in obese humans. *J Clin Invest.* 1994;93:2365–71. [PubMed: 8200969]
31. Scherrer U, Vollenweider P, Randin D, Jequier E, Nicod P, Tappy L. Suppression of insulin-induced sympathetic activation and vasodilation by dexamethasone in humans. *Circulation.* 1993;88:388–94. [PubMed: 8339402]
32. Allwood MJ, Ginsburg J, Paton A. The effect of insulin hypoglycemia on blood flow in intact and sympathectomized extremities in man. *J Physiol.* 1957;139:97–107.
33. Dela F, Stallknecht B, Biering-Sørensen F. An intact central nervous system is not necessary for insulin-mediated increases in leg blood flow in humans. *Eur J Physiol.* 2000;441(2–3):241–50.
34. Sartori C, Trueb L, Nicod P, Scherrer U. Effects of sympathectomy and nitric oxide synthase inhibition on vascular actions of insulin in humans. *Hypertension.* 1999;34(4):586–9. [PubMed: 10523331]
35. Cardillo C, Nambi SS, Kilcoyne CM, et al. Insulin stimulates both endothelin and nitric oxide activity in the human forearm. *Circulation.* 1999;100(8):820–5. [PubMed: 10458717]
36. Mahmoud AM, Szczurek MR, Blackburn BK, et al. Hyperinsulinemia augments endothelin-1 protein expression and impairs vasodilation of human skeletal muscle arterioles. *Physiol Rep.* 2016;4(16):1–15.
37. Muniyappa R, Montagnani M, Koh KK, Quon MJ. Cardiovascular actions of insulin. *Endocr Rev.* 2007;28(5):463–91. [PubMed: 17525361]
38. Kim F, Gallis B, Corson MA. TNF-alpha inhibits flow and insulin signaling leading to NO production in aortic endothelial cells. *Am J Physiol Cell Physiol.* 2001;280:C1057–65. [PubMed: 11287317]

39. Bakker W, Sipkema P, Stehouwer CDA, et al. Protein kinase C theta activation induces insulin-mediated constriction of muscle resistance arteries. *Diabetes*. 2008;57(3):706–13. [PubMed: 18086904]
40. Olver TD, McDonald MW, Klakotskaia D, et al. A chronic physical activity treatment in obese rats normalizes the contributions of ET-1 and NO to insulin-mediated posterior cerebral artery vasodilation. *J Appl Physiol*. 2017;122(4):1040–50. [PubMed: 28183819]
41. Hardin D, Azzaelli B, Edwads J, et al. Mechanisms of enhanced insulin sensitivity in endurance trained athletes: effects on blood flow and differential expression of GLUT 4 in skeletal muscle. *J Endocrinology Metab*. 1995;80:2437–46.
42. Mikus CR, Rector RS, Arce-Esquivel AA, et al. Daily physical activity enhances reactivity to insulin in skeletal muscle arterioles of hyperphagic Otsuka Long-Evans Tokushima Fatty rats. *J Appl Physiol*. 2010;109(4):1203–10. [PubMed: 20634354]
43. Mikus CR, Roseguini BT, Uptergrove GM, et al. Voluntary wheel running selectively augments insulin-stimulated vasodilation in arterioles from white skeletal muscle of insulin-resistant rats. *Microcirculation*. 2012;19(8):729–38. [PubMed: 22804760]
44. Saltin B, Gollnick PD. Skeletal muscle adaptability: significance for metabolism and performance. In: *Handbook of Physiology: Skeletal Muscle Section*. 1983 p. 10: 555–631.
45. Delp MD, Laughlin MH. Regulation of skeletal muscle perfusion during exercise. *Acta Physiol Scand*. 1998;162:411–9. [PubMed: 9578387]
46. Laughlin MH, Armstrong RB. Muscular blood flow distribution patterns as a function of running speed in rats. *Am J Physiol Hear Circ Physiol*. 1982;12:296–306.
47. Laughlin MH, Davis MJ, Secher NH, et al. Peripheral circulation. *Compr Physiol*. 2012;2(1):321–447. [PubMed: 23728977]
48. Jasperse JJ, Laughlin MH. Exercise and skeletal muscle microcirculation In: *Microvascular Research: Biology and Pathology* San Diego, CA: Elsevier; 2005 p. 527–35.
49. Laughlin MH, Woodman CR, Schrage WG, Gute D, Price EM. Interval sprint training enhances endothelial function and eNOS content in some arteries that perfuse white gastrocnemius muscle. *J Appl Physiol*. 2004;96(1):233–44. [PubMed: 12923113]
50. McAllister RM, Jasperse JL, Laughlin MH. Nonuniform effects of endurance exercise training on vasodilation in rat skeletal muscle. *J Appl Physiol*. 2005;98(2):753–61. [PubMed: 15448126]
51. Laughlin MH, Cook JD, Tremble R, Ingram D, Collieran PN, Turk JR. Exercise training produces non-uniform increases in arteriolar density of rat soleus and gastrocnemius muscle. *Microcirculation*. 2006;13(3):175–86. [PubMed: 16627360]
52. Binder KW, Murfee WL, Song JI, Laughlin MH, Price RJ. Computational network model prediction of hemodynamic alterations due to arteriolar remodeling in interval sprint trained skeletal muscle. *Microcirculation*. 2007;14(3):181–92. [PubMed: 17454671]
53. Armstrong RB, Laughlin MH. Blood flows within and among rat muscles as a function of time during high speed treadmill exercise. *J Physiol*. 1983;344:189–208. [PubMed: 6655579]
54. Armstrong RB, Laughlin MH. Rat muscle blood flows during high-speed locomotion. *J App Physiol*. 1985;59(4):1322–8.
55. Olver TD, Ferguson BS, Laughlin MH. Molecular mechanisms for exercise training-induced changes in vascular structure and function: skeletal muscle, cardiac muscle, and the brain In: Bouchard C, editor. *Progress in Molecular Biology and Translational Science*. Elsevier Inc.; 2015 p. 227–57.
56. Andersen P, Saltin B. Maximal perfusion of skeletal muscle in man. *J Physiol*. 1985;366:233–49. [PubMed: 4057091]
57. Dimmeler S, Assmus B, Hermann C, Haendeler J, Zeiher AM. Fluid shear stress stimulates phosphorylation of Akt in human endothelial cells: Involvement in suppression of apoptosis. *Circ Res*. 1998;83:334–41. [PubMed: 9710127]
58. Jenkins NT, Padilla J, Martin JS, et al. Differential vasomotor effects of insulin on gastrocnemius and soleus feed arteries in the OLETF rat model: role of endothelin-1. *Exp Physiol*. 2014;99(1):262–71. [PubMed: 23995100]

59. Bender SB, Newcomer SC, Harold Laughlin M. Differential vulnerability of skeletal muscle feed arteries to dysfunction in insulin resistance: impact of fiber type and daily activity. *Am J Physiol Heart Circ Physiol*. 2011;300(4):H1434–41. [PubMed: 21317309]
60. Laughlin MH, Korthuis RJ, Sexton WL, Armstrong RB. Regional muscle blood flow capacity and exercise hyperemia in high-intensity trained rats. *J Appl Physiol*. 1988;64(6):2420–7. [PubMed: 3403424]
61. Armstrong RB, Laughlin MH. Exercise blood flow patterns within and among rat muscles after training. *Am J Physiol Heart Circ Physiol*. 1984;246(1 Pt 2):H59–68.
62. Sexton WL, Laughlin MH. Influence of endurance exercise training on distribution of vascular adaptations in rat skeletal muscle. *Am J Physiol Heart Circ Physiol*. 1994;266(2 Pt 2):H483–HH490.
63. Laughlin MH, Korthuis RJ, Dunker DJ, J BR. Control of blood flow to cardiac and skeletal muscle during exercise New York: American Physiological Society and Oxford University Press; 1996 Chapter 16.
64. Musch TI, Haidet GC, Ordway GA, Longhurst JC, Mitchell JH. Training effects on regional blood flow response to maximal exercise in foxhounds. *J Appl Physiol*. 1987;62(4):1724–32. [PubMed: 3597244]
65. Cocks M, Shaw CS, Shepherd SO, et al. Sprint interval and continuous training have equal benefits on aerobic capacity, insulin sensitivity, muscle capillarisation and endothelial eNOS/NAD(P)H oxidase protein ratio in obese men. *J Physiol*. 2016;594(8):2307–21. [PubMed: 25645978]
66. Cocks M, Shaw CS, Shepherd SO, et al. Sprint interval and endurance training are equally effective in increasing muscle microvascular density and eNOS content in sedentary males. *J Physiol*. 2013;591:641–56. [PubMed: 22946099]
67. Kim HJ, Lee JS, Kim CK. Effect of exercise training on muscle glucose transporter 4 protein and intramuscular lipid content in elderly men with impaired glucose tolerance. *Eur J Appl Physiol*. 2004 12;93(3):353–8. [PubMed: 15480742]
68. Prior SJ, Blumenthal JB, Katzel LI, Goldberg AP, Ryan AS. Increased skeletal muscle capillarization after aerobic exercise training and weight loss improves insulin sensitivity in adults with IGT. *Diabetes Care*. 2014;37(5):1469–75. [PubMed: 24595633]
69. Egginton S, Hudlick O. Selective long-term electrical stimulation of fast glycolytic fibres increases capillary supply but not oxidative enzyme activity in rat skeletal muscles. *Exp Physiol*. 2000;85(5):567–74. [PubMed: 11038408]
70. Egginton S. Invited review: activity-induced angiogenesis. *Eur J Physiol*. 2009;457:963–77.
71. Laughlin MH, Roseguini B. Mechanisms for exercise training-induced increases in skeletal muscle blood flow capacity: differences with interval sprint training versus aerobic endurance training. *J Physiol Pharmacol*. 2008;59(Suppl 7):71–88. [PubMed: 19075325]
72. Laughlin MH, Newcomer SC, Bender SB. Importance of hemodynamic forces as signals for exercise-induced changes in endothelial cell phenotype. *J Appl Physiol*. 2008;104(3):588–600. [PubMed: 18063803]
73. Lash JM, Bohlen HG. Time- and order-dependent changes in functional and NO-mediated dilation during exercise training. *J Appl Physiol*. 1997;82(2):460–8. [PubMed: 9049725]
74. Mcallister M, Kimani JK, Webster JL, Parker JL, Laughlin MH. Effects of exercise training on responses of peripheral and visceral arteries in swine. *J Appl Physiol*. 1996;80(1):216–25. [PubMed: 8847306]
75. Parker JL, Mattox ML, Laughlin MH. Contractile responsiveness of coronary arteries from exercise-trained rats. *J Appl Physiol*. 1997;83(2):434–43. [PubMed: 9262438]
76. Sessa WC, Pritchard K, Seyedi N, Wang J, Hintze TH. Chronic exercise in dogs increases coronary vascular nitric oxide production and endothelial cell nitric oxide synthase gene expression. *Circ Res*. 1994;74(2):349–53. [PubMed: 7507417]
77. Wang J, Wolin MS, Hintze TH. Chronic exercise enhances endothelium-mediated dilation of epicardial coronary artery in conscious dogs. *Circ Res*. 1993;73(5):829–38. [PubMed: 8403254]
78. Sun D, Huang A, Koller A, Kaley G. Adaptation of flow-induced dilation of arterioles to daily exercise. *Microvasc Res*. 1998;56(1):54–61. [PubMed: 9683563]

79. Olver TD, Reid SM, Smith AR, et al. Effects of acute and chronic interval sprint exercise performed on a manually propelled treadmill on upper limb vascular mechanics in healthy young men. *Physiol Rep*. 2016;4(13).
80. Delp MD, Laughlin MH. Time course of enhanced endothelium-mediated dilation in aorta of trained rats. *Med Sci Sport Exerc*. 1997;1454–61.
81. Delp D, McAllister M, Harold M. Exercise training alters aortic vascular reactivity in hypothyroid rats. *Am J Physiol Heart Circ Physiol*. 1995;(19):1428–35.
82. Delp MD, McAllister RM, Laughlin MH. Exercise training alters endothelium-dependent vasoreactivity of rat abdominal aorta. *J Appl Physiol*. 1993;75(3):1354–63. [PubMed: 8226551]
83. Koller A, Huang A, Sun D, Kaley G. Exercise training augments flow-dependent dilation in rat skeletal muscle arterioles. Role of endothelial nitric oxide and prostaglandins. *Circ Res*. 1995;76:544–50. [PubMed: 7534658]
84. Laughlin MH, Pollock JS, Amann JF, Hollis ML, Woodman CR, Price EM. Training induces nonuniform increases in eNOS content along the coronary arterial tree. *J Appl Physiol*. 2001;90(2):501–10. [PubMed: 11160048]
85. McAllister RM, Laughlin MH. Short-term exercise training alters responses of porcine femoral and brachial arteries. *J Appl Physiol*. 1997;82(5):1438–44. [PubMed: 9134890]
86. Muller JM, Myers PR, Laughlin MH. Vasodilator responses of coronary resistance arteries of exercise-trained pigs. *Circulation*. 1994;89(5):2308–14. [PubMed: 8181157]
87. Sexton WL, Korthuis RJ, Laughlin MH. High-intensity exercise training increases vascular transport capacity of rat hindquarters. *Am J Physiol Heart Circ Physiol*. 1988;254(2 Pt 2):H274–8.
88. Bevegård BS, Shepherd JT. Reaction in man of resistance and capacity vessels in forearm and hand to leg exercise. *J Appl Physiol*. 1966;21(1):123–32. [PubMed: 5903898]
89. Clausen JP, Trap-Jensen J. Effects of training on the distribution of cardiac output in patients with coronary artery disease. *Circulation*. 1970;42(4):611–24. [PubMed: 11993304]
90. Rowell LB. *Human Cardiovascular Control*. Oxford University Press; 1993.
91. Gute D, Fraga C, Laughlin MH, Amann JF. Regional changes in capillary supply in skeletal muscle of high-intensity endurance-trained rats a. *J Appl Physiol*. 1996;81(2):619–26. [PubMed: 8872626]
92. Gute D, Laughlin MH, Amann JF. Regional changes in capillary supply in skeletal muscle of interval-sprint and low intensity, endurance trained rats. *Microcirculation*. 1994;1:183–93. [PubMed: 8790589]
93. Mackie BG, Terjung RL. Blood flow to different skeletal muscle fiber types during contraction. *Am J Physiol Heart Circ Physiol*. 1983;245(2):H265–75.
94. Edgerton VR, Smith JL, Simpson DR. Muscle fibre type populations of human leg muscles. *Histochem J*. 1975;7(3):259–66. [PubMed: 123895]
95. Dwyer D, Browning J, Weinstein S. The reliability of muscle biopsies taken from vastus lateralis. *J Sci Med Sport* /. 1999;2(4):333–40. [PubMed: 10710011]
96. Dela F, Larsen JJ, Mikines KJ, Ploug T, Petersen LN. Insulin-stimulated muscle glucose clearance in patients with NIDDM: effects of one-legged physical training. *Diabetes*. 1995;44:1010–20. [PubMed: 7657022]
97. Holten MK, Zacho M, Gaster M, Juel C, Wojtaszewski JFP, Dela F. Strength training increases insulin-mediated glucose uptake, GLUT4 content, and insulin signaling in skeletal muscle in patients with type 2 diabetes. *Diabetes*. 2004;53:294–305. [PubMed: 14747278]
98. Dela F, Mikines KJ, Larsen JJ, Galbo H. Training-induced enhancement of insulin action in human skeletal muscle: the influence of aging. *J Gerontol*. 1996;51(4):247–52.
99. Sjøberg KA, Frøsig C, Kjøbsted R, et al. Exercise increases human skeletal muscle insulin sensitivity via coordinated increases in microvascular perfusion and molecular signaling. *Diabetes*. 2017;66(6):1501–10. [PubMed: 28292969]
100. Eskelinen J-J, Heinonen I, Löytyniemi E, et al. Muscle-specific glucose and free fatty acid uptake after sprint interval and moderate intensity training in healthy middle-aged men. *J Appl Physiol*. 2015;118:1172–80. [PubMed: 25767035]

101. McDonald MW, Olver TD, Dotzert MS, et al. Aerobic exercise training improves insulin-induced vasorelaxation in a vessel-specific manner in insulin-treated experimental diabetes. *Diabetes Vasc Dis Res.* 2018;accepted(aheadofprint):Jan-2018–00002.
102. Walsh L, Ghiarone T, Olver TD, et al. Increased endothelial shear stress improves insulin-stimulated vasodilation in skeletal muscle. *J Physiol.* 2018;accepted(aheadofprint):doi: 10.1113/JP277050.
103. Wagenmakers AJM, Strauss JA, Shepherd SO, Keske MA, Cocks M. Increased muscle blood supply and transendothelial nutrient and insulin transport induced by food intake and exercise: Effect of obesity and ageing. *J Physiol.* 2016;594(8):2207–22. [PubMed: 25627798]
104. Prior SJ, Goldberg AP, Ortmeier HK, et al. Increased skeletal muscle capillarization independently enhances insulin sensitivity in older adults after exercise training and detraining. *Diabetes.* 2015;64(10):3386–95. [PubMed: 26068543]
105. Lillioja S, Young AA, Culter CL, et al. Skeletal muscle capillary density and fiber type are possible determinants of in vivo insulin resistance in man. *J Clin Invest.* 1987;80:415–24. [PubMed: 3301899]
106. Akerstrom T, Laub L, Vedel K, et al. Increased skeletal muscle capillarization enhances insulin sensitivity. *Am J Physiol Endocrinol Metab.* 2014;307(12):E1105–16. [PubMed: 25352432]
107. Hiura M, Nariai T, Ishii K, et al. Changes in cerebral blood flow during steady-state cycling exercise: a study using oxygen-15-labeled water with PET. *J Cereb Blood Flow Metab.* 2014 3;34(3):389–96. [PubMed: 24301294]
108. Curtelin D, Morales-Alamo D, Torres-Peralta R, et al. Cerebral blood flow, frontal lobe oxygenation and intra-arterial blood pressure during sprint exercise in normoxia and severe acute hypoxia in humans. *J Cereb Blood Flow Metab.* 2018;38(1):136–50. [PubMed: 28186430]
109. Schmidt W, Endres M, Dimeo F, Jungehulsing GJ. Train the vessel, gain the brain: physical activity and vessel function and the impact on stroke prevention and outcome in cerebrovascular disease. *Cerebrovasc Dis.* 2013;35(4):303–12. [PubMed: 23594423]
110. Lucas SJE, Cotter JD, Brassard P, Bailey DM. High-intensity interval exercise and cerebrovascular health: curiosity, cause, and consequence. *J Cereb Blood Flow Metab.* 2015;35(6):902–11. [PubMed: 25833341]
111. Delp MD, Armstrong RB, Godfrey DA, Laughlin MH, Ross CD, Wilkerson MK. Exercise increases blood flow to locomotor, vestibular, cardiorespiratory and visual regions of the brain in miniature swine. *J Physiol.* 2001 6 15;533(3):849–59. [PubMed: 11410640]
112. Isaacs KR, Anderson BJ, Alcantara AA, Black JE, Greenough WT. Exercise and the brain: angiogenesis in the adult rat cerebellum after vigorous physical activity and motor skill learning. *J Cereb Blood Flow Metab.* 1992 1;12(1):110–9. [PubMed: 1370068]
113. Vissing J, Andersen M, Diemer NH. Exercise-induced changes in local cerebral glucose utilization in the rat. *J Cereb Blood Flow Metab.* 1996 7;16(4):729–36. [PubMed: 8964814]
114. Kullmann S, Heni M, Hallschmid M, Fritsche A, Preissl H, Häring H-U. Brain Insulin Resistance at the Crossroads of Metabolic and Cognitive Disorders in Humans. *Physiol Rev.* 2016;96(4):1169–209. [PubMed: 27489306]
115. Hirvonen J, Virtanen KA, Nummenmaa L, et al. Effects of insulin on brain glucose metabolism in impaired glucose tolerance. *Diabetes.* 2011;60(2):443–7. [PubMed: 21270256]
116. Martin JS, Padilla J, Jenkins NT, et al. Functional adaptations in the skeletal muscle microvasculature to endurance and interval sprint training in the type 2 diabetic OLETF rat. *J Appl Physiol.* 2012;113(8):1223–32. [PubMed: 22923508]
117. Wang H, Wang AX, Aylor K, Barrett EJ. Nitric oxide directly promotes vascular endothelial insulin transport. *Diabetes.* 2013;62(12):4030–42. [PubMed: 23863813]
118. Meijer RI, Gray S, Aylor K, Barrett EJ. Pathways for insulin access to the brain: the role of the microvascular endothelial cell. *Am J Physiol Heart Circ Physiol.* 2016;311(5):H1132–8. [PubMed: 27591216]
119. Honkala SM, Johansson J, Motiani KK, et al. Short-term interval training alters brain glucose metabolism in subjects with insulin resistance. *J Cereb Blood Flow Metab.* 2017;0(00):1–11.
120. Ebeling P, Bourey R, Koranyi L, Tet al. Mechanism of enhanced insulin sensitivity in athletes. *J Clin Invest.* 1993;92:1623–31. [PubMed: 8408617]

121. Bisquolo VAF, Cardoso JCG, Ortega KC, et al. Previous exercise attenuates muscle sympathetic activity and increases blood flow during acute euglycemic hyperinsulinemia. *J Appl Physiol.* 2005;98:866–71. [PubMed: 15542577]
122. Vinet A, Obert P, Dutheil F, et al. Impact of a Lifestyle Program on Vascular Insulin Resistance in Metabolic Syndrome Subjects: The RESOLVE Study. *J Clin Endocrinol Metab.* 2015;100(2): 442–50. [PubMed: 25353072]
123. Russell RD, Hu D, Greenaway T, et al. Skeletal muscle microvascular-linked improvements in glycemic control from resistance training in individuals with type 2 diabetes. *Diabetes Care.* 2017;40(9):1–16.

Keypoints:

- Insulin resistance in the skeletal muscle and brain vasculatures represents a unifying pathology linking metabolic, neuro- and vascular diseases.
- Routine physical activity (PA) is an effective therapeutic approach to optimize vascular insulin signaling in the skeletal muscle and brain.
- PA that involves a large skeletal muscle mass and modulates intensity to recruit more muscle fibers within each muscle, as well as, requires a significant central neural recruitment will maximize vascular adaptations that promote vascular insulin sensitivity in the skeletal muscle and brain.
- Future research should focus on whether vaso-metabolic and vaso-neural treatment outcomes in preclinical and clinical populations with insulin resistance are improved when such mechanistic concepts are integrated into prescriptive-based PA interventions.

Figure 1.

Skeletal muscle and brain vascular adaptations to physical activity include enhanced endothelial insulin sensitivity and capillary density. Adaptations occur primarily in arteries that supply tissue regions that undergo a relative increase in activity during exercise. Whereas low-intensity walking involves recruitment of a small amount of total muscle mass, few muscle fibers within each muscle and a small population of neurons in a small number of brain regions; combined low, moderate, and high-intensity cross country skiing involves recruitment of a large amount of total muscle mass, many muscle fibers within each muscle and requires activation of a larger population of neurons from a greater number of brain regions. Therefore, we speculate that compared to low-intensity walking (or any similar exercise that only involves a small amount of skeletal muscle and neural recruitment), cross country skiing (or any similar whole-body exercise where intensity and duration can be manipulated to modulate the total number of skeletal muscle fibers and brain regions involved) confers more widespread improvements in vascular insulin sensitivity and capillary density. Enhanced endothelial insulin sensitivity corresponds with greater production of nitric oxide (NO) relative to endothelin-1 (ET-1). This shift is associated with reduced vascular inflammation and attenuated progression of atherosclerosis in conduit arteries, improved insulin-induced dilation in resistance arteries, increased insulin-stimulated capillary recruitment, enhanced nutrient delivery and potentially augmented trans-endothelial insulin transport in skeletal muscle and the brain. Improvements in nutrient delivery are further enhanced by physical activity-induced increases in microvascular volume/capillary density. The cumulative effect of enhanced vascular insulin sensitivity and

capillary density may include improved metabolic and neurocognitive function in the skeletal muscle and brain, respectively.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 2.

Schematic depicting the dependence of both vascular shear stress and endothelial insulin signaling on the phosphatidylinositol-4,5-bisphosphate 3-kinase (PI3K)/protein kinase B (Akt)/nitric oxide (NO) pathway signaling cascade. Briefly, vascular shear stress causes the deformation of luminal endothelial mechanoreceptors/glycocalyx which stimulate intracellular activation of the PI3K/Akt/NO pathway and insulin stimulates activation of both the endothelial PI3K/Akt/NO and the mitogen activated protein kinase (MAPK)/endothelin-1 (ET-1) pathway. The vascular actions of insulin are influenced by the balance between the two pathways. The dependence of both vascular shear stress and endothelial insulin signaling on the PI3K/Akt/NO pathway raises the possibility that vascular shear stress enhances vascular insulin sensitivity by promoting activation of the PI3K/Akt/NO pathway. VASC=vascular, IRS1/2=insulin receptor substrate 1/2, ET_{A/B}=endothelin receptor, CaM=calcium calmodulin complex, cyt=cytosolic, PIP=phosphatidylinositol 4,5-phosphate, GC=guanylate cyclase, PDE5=phosphodiesterase 5, eNOS=endothelial NO synthase, O₂=oxygen, BH₄=tetrahydrobiopterin, P=phosphorylated.

Table.

Human studies examining the effect of exercise on vascular insulin signaling

Study type/authors	Population	Physical Activity	Main Findings
Cross sectional observations			
Ebeling et al. (1993)(120)	Healthy men (28±2 yr)	Bandy players vs sedentary control	Insulin did not induce significant vasodilation, but Brandy players displayed increased blood flow and insulin-stimulated glucose uptake in the forearm
Hardin et al (1995)(41)	Healthy men (25±1 yr)	Endurance athletes vs sedentary control	Runners displayed greater insulin-stimulated vasodilation and glucose uptake in the lower limb
Acute exercise intervention			
Richter et al. (1989)(5)	Healthy men (21–24 yr)	Single leg cycle ergometer exercise (75% of max work capacity for 60 min)	Exercise improved insulin-stimulated glucose uptake but not lower limb vasodilation
Bisquolo et al (2005)(121)	Healthy men (32±2 yr)	Cycle ergometer exercise (50% peak O ₂ uptake for 45 min)	Exercise improved insulin-stimulated vasodilation in the forearm
Sjøberg et al (2017) (99)	Healthy men (25±1 yr)	Single leg cycle ergometer exercise (50% peak work load for 60 min, with 3 × 5 min intervals at 100% peak workload)	Exercise improved insulin-stimulated glucose uptake and vasodilation in the lower limb
Exercise training intervention			
Dela et al. (1995)(96)	Healthy men and men with T2D (58±3 yr)	Single leg cycle ergometer exercise (30 min per day at 70% $\dot{V}O_{2peak}$, 6 d per wk, for 10 wk)	Training improved insulin-stimulated glucose uptake and insulin-stimulated vasodilation in the trained limb
Dela et al. (1996)(98)	Healthy young (23±1 yr) and old men (59±1 yr)	Single leg cycle ergometer exercise (30 min per day at 70% $\dot{V}O_{2peak}$, 6 d per wk, for 10 wk)	Training improved insulin-stimulated glucose uptake and insulin-stimulated vasodilation in the trained limb
Holten et al (2004)(97)	Healthy men and men with T2D (62±2 yr)	Single leg strength training exercise (leg press, knee extensions, ham string curls, 3–4 sets for 8–12 reps at 70–80% 1 RM, 3 d per wk, for 6 wk)	Training improved insulin-stimulated glucose uptake and insulin-stimulated vasodilation in the trained limb
Vinet et al. (2015)(122)	Healthy men and women and men and women with metabolic syndrome (59±1 yr)	Lifestyle intervention: diet education program with monitored walking or aqua gym and resistance training (8 exercises, 3 sets of 10 repetitions, 4–5 d per wk for 6 months)	Lifestyle intervention improved insulin-stimulated cutaneous perfusion response
Eskelinen et al (2015)(100)	Healthy men (40–55 yr)	Moderate continuous endurance cycling (60% $\dot{V}O_{2max}$ for 40–60 min, 3 d per wk for 2 wk) or high intensity interval sprint cycling (4–6 repeated Wingate tests, interspersed with 4.5 min recovery, 3 d per wk for 2 wk)	Training improved insulin-stimulated glucose uptake in lower vs upper limbs, with high intensity training affecting a greater amount of quadriceps muscle tissue
Honkala et al. (2017)(119)	Insulin resistant men and women (44–54 yr)	Moderate continuous endurance cycling (60% $\dot{V}O_{2max}$ for 40–60 min, 3 d per wk for 2 wk) or high intensity interval sprint cycling (4–6 repeated Wingate tests, interspersed with 4.5 min recovery, 3 d per wk for 2 wk)	Only high intensity sprint training increased aerobic capacity and reduced insulin-stimulated glucose uptake in the brain
Russel et al. (2017)(123)	Men and women with T2D (52±2 yr)	Whole body resistance training (leg press, pull down, chest press, lunges, seated row, back fly, bicep curl, incline chest press, dumbbell shoulder press, leg extension, leg curl, dips, lateral raise, triceps extensions, dumbbell	Training improved forearm skeletal muscle, but not skin, microvascular blood flow response to an oral glucose challenge

Study type/authors	Population	Physical Activity	Main Findings
		deadlift and push-ups, 1 set of 6–15 reps at 65–85% 1 RM, 3 d per wk for 6 wk)	

T2D=type 2 diabetes, RM= repetition maximum

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript