CORRESPONDENCE

The Relevance of Irreversible Loss of Brain Function as a Reliable Sign of Death

by Prof. Dr. med. Stephan A. Brandt, and Prof. Dr. med. Heinz Angstwurm (ret.)

in issue 41/2018

Spiritual Issues Should Be Considered

My comment on the article by Brandt and Angstwurm (1) is that much greater value should be placed on spiritual or religious issues than on sober scientific approaches. The question is that of life and death. The fact that an ethics committee has to be summoned in order to decide whether brain death means a person's actual death or not, shows the degree to which human intellectuality has come to overlay our hidden spiritual insights.

An article by Walter van Laack (2) explains how research results of near-death experiences imply that a consciousness exists that is independent of the brain and possibly survives a person's death. We can certainly assume that the entire complexity of near-death experiences is cannot be explained in terms of neurophysiology. Today's medical scientists are aware that consciousness exists independently of brain function. We can therefore conclude that a brain dead human body is not a corpse.

An article by Cathrin Marshall (3) makes apparent the enormous emotional stress that a transplant team is under when a brain-dead body is opened up from the jugular vein to the pubal symphysis, and the organs are explanted from the hollow basin thus created.

Intuition is a result of psychological-spiritual experiences and cannot be taught. Human beings require time to reach spiritual decisions, especially when crucial decisions—such as organ donation—are concerned.

The opt-out system for organ donation interferes massively with the spiritual decision-finding process. It is a psychological assault and for this reason should be strictly rejected on moral grounds.

DOI: 10.3238/arztebl.2019.0097a

References

1. Brandt SA, Angstwurm H: The relevance of irreversible loss of brain function as a reliable sign of death. *Dtsch Arztebl Int* 2018; 115: 675–81.
2. Van Laack: Nahtoderfahrungen. *Der Allgemeinarzt* 2017; 39: 80–3.
3. Marshall C: Grenzsituationen im Arbeitsbereich des OP-Pflegepersonals – Die Explantation. www.transplantation-information.de/...datei/explantation_facharbeit_2004.pdf (last accessed on 8 January 2019).

Dr. rer. nat. Dr. med. Alois Bahemann
Arzt für Allgemeinmedizin – Homöopathie
info@praxis-bahemann.de

Metaphysical and Cultural Aspects

The review article (1) explicitly excludes metaphysical and cultural questions. For this reason it is not capable of producing a consensus—neither as a profound explanation for potential organ donors nor as the basis of a universally binding law, not even within the medical profession. The authors overlooked all human beings for whom consciousness (to use the words of Max Planck) is fundamental. From this perspective, “brain dead” people are terminal patients in whom life-prolonging measures are not indicated, but who are still neither dead nor without sensation, but dying, suffering, and defenseless, not able to communicate their inner lives because the tool for this—the brain—is missing.

From a medical perspective the article is not immune to challenge:

- A complete loss of function of the cerebrum, cerebellum, and brainstem cannot be confirmed by applying the currently used diagnostic methods (2). The criteria of brain death do not rule out that parts of the brain perceive external stimuli and can react to such stimuli (3).

- The fact that the organs of the organism have their own coordinating function independently of the brain is not something that can be disproved in one sentence. Also, a great difference exists between total loss of function and isolation of one organ.

- Historically it can be clearly documented that the Harvard Committee with its definition of brain death primarily considered the needs of transplantation medicine (4).

Experts in the German Medical Association have attempted for many years to ignore the international consensus on brain death (brain dead patients are dying patients but are still alive) and maintain their own concept. How much longer is this going to work?

DOI: 10.3238/arztebl.2019.0097b

References

1. Brandt SA, Angstwurm H: The relevance of irreversible loss of brain function as a reliable sign of death. *Dtsch Arztebl Int* 2018; 115: 675–81.
2. Rehder S: Grauzone Hirntod – Organspende verantworten. Augsburg: Sankt Ulrich Verlag GmbH 2010; p. 58.
3. Owen A: Zwischenwelten – Ein Wissenschaftler erforscht die Grauzone zwischen Leben und Tod. München: Droemer 2017.
4. Henderson DS: Death and donation—Rethinking brain death as a means for procuring transplantable Eugene, Oregon (USA): Pickwick Publications 2011; p. 6–10.

Dr. med. Hans-Joachim Ritz
Hannover, hajoritz@arcor.de

Is Irreversible Loss of Brain Function a Reliable Sign of Death?

Describing irreversible loss of brain function (ILBF) as “the biological end of the life of a human being” (1) does not tally with the thanatological definition of biological death as the death of the last cell (2). The two are not equivalent. The review article itself makes this clear with its description of the function of the pituitary gland.

One collaborator listed in Box 3 in the article (1) doubts the admissibility of equating death with ILBF (3). The cited position of the German Ethics Council (4) is equivocal in this context, but this was not mentioned in the article.

From a scientific-medical perspective, several definitions of death exist (for example, clinical death; ILBF as individual death but, by contrast, ILBF as an “extreme minimal living state”[4]; biological death). Even the legislator differentiates in § 3 I 1 No 2 of the German Transplant Act (TPG) between death as an admissibility criterion and, in § 3 II No 2 TPG, brain death (=final, irreversible loss of all function of the cerebrum, cerebellum, and brainstem) as an inadmissibility criterion, and names in § 5 I 2 TPG the final irreversible cardiac arrest.

In spite of the explanation and the erroneous description of the fourth revision of the German Medical Association’s guideline, no complete and acceptable rules for establishing death according to §§ 3 I 1 No 2, 16 I 1 No 1 TPG exist. This is obvious—among other reasons—from the fact that the guideline is lacking information on putrefaction as a reliable sign of death, as well as well-founded guidance on establishing death (apart from ILBF).

For whom or what the German Medical Association as a non-registered association deserved a special role in the sense of a guarantor function (1) was not explained.

DOI: 10.3238/arztebl.2019.0098a

References

1. Brandt SA, Angstwurm H: The relevance of irreversible loss of brain function as a reliable sign of death. *Dtsch Arztebl Int* 2018; 115: 675–81.
2. Bajanowski T, Brinkmann B: Todesbegriffe und Todesfeststellung. In: Brinkmann B, Madea B (eds.): *Handbuch gerichtliche Medizin* 2004; p. 13–18.
3. Birnbacher D: Der Hirntod – der Tod? *DZPhil* 2012; 60: 422–3.
4. Deutscher Ethikrat (eds.): *Hirntod und Entscheidungen zur Organspende. Stellungnahme*. www.ethikrat.org/fileadmin/Publikationen/Stellungnahmen/deutsch/stellungnahme-hirntod-und-entscheidung-zur-organspende.pdf (last accessed on 17 October 2018): p. 102; p.159.

Assessor Prof. Dr. med. Dr. med. habil. Markus Parzeller

Institut für Rechtsmedizin, Universitätsklinikum Frankfurt, Goethe-Universität Frankfurt am Main
parzeller@em.uni-frankfurt.de

Prof. Dr. med. Marcel A. Verhoff

Institut für Rechtsmedizin, Universitätsklinikum Frankfurt, Goethe-Universität Frankfurt am Main

Dr. med. Barbara Zedler

Institut für Rechtsmedizin der Universitätskliniken Gießen und Marburg

Pertinent Points Need to Be More Convincingly Stated

The excellent, clear, and persuasive explanation of the importance of brain death will serve as a great orientation aid to colleagues (1). Two points, however, could have been stated more convincingly:

- The insistence that even in primary infratentorial lesions the loss of cerebral function should always be documented is justified in the article as being “in the interest of internal consistency.” In these patients with primary loss of brainstem function and sustained function of the large hemispheres, as well as possibly reproducible visually evoked potentials, the current state of science does not allow for ascertaining the degree to which perception—that is, consciousness—is still a possibility. A more precise way of reasoning would be: In primary infratentorial lesions the irreversible loss of supratentorial brain function must be confirmed, because all brain functions are to be excluded—including any consciousness.

- In terms of the popular but incorrect notion that brain death was defined in order to enable organ explantation, the authors rightly say that the German Society for Surgery published brain death criteria before the Harvard Ad Hoc Committee in 1968. This argument is likely to convince but few because the public debate about how to define the death of an organ donor received the attention of the public earlier than that, in December 1967, after the first heart transplantation. The authors of the first scientific accounts, neurosurgeons Wertheimer in 1959 and Tönnis and Frohwein in Germany in 1963 (2) regarded brain death merely as a justification to stop treatment (3). The option of organ donation was not known and was therefore not discussed. The diagnosis of brain death was used for organ donations only after these scientific contributions. The German Society for Surgery and the Harvard Ad Hoc Committee based their recommendations on these scientific findings.

DOI: 10.3238/arztebl.2019.0098b

References

1. Brandt SA, Angstwurm H: The relevance of irreversible loss of brain function as a reliable sign of death. *Dtsch Arztebl Int* 2018; 115: 675–81.
2. Tönnies W, Frohwein RA: Wie lange ist Wiederbelebung bei schweren Hirnverletzungen möglich? *Mschr Unfallheilkunde* 1963; 66: 169–90.
3. Frohwein RA, Firsching R: Hirntod-Diagnose in Deutschland. In: Deutsche Gesellschaft für Neurochirurgie (eds.): *Neurochirurgie in Deutschland: Geschichte und Gegenwart. 50 Jahre Deutsche Gesellschaft für Neurochirurgie*. Berlin, Wien: Blackwell Wissenschafts-Verlag 2001; 203–18.

Prof. Dr. med. Raimund Firsching

Direktor der Universitätsklinik für Neurochirurgie
raimund.firsching@med.ovgu.de

In Reply:

We thank our colleagues for their correspondence. They show the importance of differentiating the scientific-medical reasoning for the relevance of irreversible loss of brain function (ILBF) as a reliable sign of death from other questions associated with death.

On the one hand, the externally visible reliable signs of death mentioned by Parzeller et al. and the comments relating to § 3 TPG were not the subject of our article on the relevance of ILBF as a reliable sign of death. The objection regarding the fourth revision of the guideline was not justified, and neither were Ritz's misgivings about determination of ILBF—we therefore are unable to comment. In our opinion, the criticism of death as the biological end of human life cannot be deduced from the cited handbook article. The fact that the German Medical Association functioned as a guarantor is based on the multitude of responsibility transfers within the transplant law. In view of our pluralistic society, it is not surprising that the German Ethics Council expressed its position, as is well-known, in a majority and minority vote. By contrast we wrote our article (1) in an interdisciplinary fashion and reached uniform consent among all involved.

On the other hand, it is obvious that everyone's individual approach to death always includes—in addition to the medical aspects—the religious-metaphysical, cultural, and everything but medical aspects, as emphasized by Bahemann and Ritz. But let us clarify: with ILBF, the “brain tool” is lacking not only for communicating one's “inner life,” but also for perception and consciousness. We are not familiar with any “international consensus,” according to which people with

ILBF as established by using the appropriate protocol are “dying but still alive.” Rather, the relevance of ILBF as a reliable sign of death is accepted by the responsible medical professional societies and has become anchored in the law in several countries. The relevant declarations from the churches are included in the reference list with our article.

We wish to thank Firsching for clarifying once again two aspects, which also touches on the comment by Ritz, that the “definition of brain death” had primarily satisfied a requirement of transplantation medicine. In any case, considerations on the motives of scientific statements need to be clearly distinguished from what the statements actually say.

In the same spirit, let us refer once more to the final paragraph of our article. It is especially the medical profession's responsibility to distinguish cautiously and carefully between questions relating to subject matter and those relating to meaning. The acceptance of ILBF is also a question of trust in the medical profession and its responsible application.

DOI: 10.3238/arztebl.2019.0099

References

1. Brandt SA, Angstwurm H: The relevance of irreversible loss of brain function as a reliable sign of death. *Dtsch Arztebl Int* 2018; 115: 675–81

On behalf of the authors

Prof. Dr. med. Stephan A. Brandt

Klinik für Neurologie mit Experimenteller Neurologie,
Campus Charité Mitte, Charité – Universitätsmedizin, Berlin
Bundesärztekammer, Berlin
dezernat6@baek.de

Prof. Dr. med. Heinz Angstwurm (ret.)

Neurologische Klinik der
Ludwig-Maximilians-Universität München

Conflict of interest statement

All correspondents declare that no conflict of interest exists.