Published in final edited form as:

Biomol NMR Assign. 2019 April ; 13(1): 233–237. doi:10.1007/s12104-019-09883-0.

Chemical Shift Assignments of a Calmodulin Intermediate with Two Ca^{2+} Bound in Complex with the IQ-motif of Voltage-gated Ca^{2+} Channels ($\text{Ca}_v1.2$)

Ian Salvesson¹, David E. Anderson¹, Johannes W. Hell², and James B. Ames^{1,*}

¹Department of Chemistry, University of California, Davis, CA 95616

²Department of Pharmacology, University of California, Davis, CA 95616

Abstract

Calcium-dependent inactivation (CDI) of neuronal voltage-gated Ca^{2+} channels ($\text{Ca}_v1.2$) is important for synaptic plasticity, which is associated with learning and memory. The Ca^{2+} -dependent binding of calmodulin (CaM) to $\text{Ca}_v1.2$ is essential for CDI. Here we report NMR assignments for a CaM mutant (D21A/D23A/D25A/E32Q/D57A/D59A/N61A/E68Q, called CaM^{EF12}) that contains two Ca^{2+} bound at the third and fourth EF-hands (EF3 and EF4) and is bound to the IQ-motif (residues 1644–1665) from $\text{Ca}_v1.2$ (BMRB accession no. 27692).

Keywords

calmodulin; EF-hand; $\text{Ca}_v1.2$; IQ-motif; synaptic plasticity; CDI

Biological Context

The neuronal L-type voltage-gated Ca^{2+} channel ($\text{Ca}_v1.2$) controls the excitability of the postsynaptic membrane in hippocampal neurons, which plays an important role in learning and memory (Hell et al, 1993; Moosmang et al, 2005; Vogl et al, 2015). The cytosolic C-terminal region of $\text{Ca}_v1.2$ (residues 1644–1665, called IQ-motif) is important for promoting Ca^{2+} -dependent inactivation (CDI) of $\text{Ca}_v1.2$ (Erickson et al, 2001; Zuhlke et al, 1999). Ca^{2+} -saturated CaM (with 4 Ca^{2+} bound) binds tightly to the IQ-motif (Findeisen et al, 2013), and the $\text{Ca}_v1.2$ mutation I1654A that disables Ca^{2+} /CaM binding also prevents CDI (Ben Johny et al, 2013). The Ca^{2+} -free form of calmodulin (apoCaM) has been suggested to bind to $\text{Ca}_v1.2$ under basal conditions, which may explain how $\text{Ca}_v1.2$ is pre-associated with CaM (Adams et al, 2014). Atomic resolution structures are currently known for the IQ-motif bound to both apoCaM (Gabelli et al, 2014) and Ca^{2+} -saturated CaM (Van Petegem et al, 2005), but no structural information is known for the IQ-motif bound to the half-saturated intermediate state of CaM (with Ca^{2+} bound at EF3 and EF4 but not bound at EF1 and EF2). This CaM intermediate state is functionally present under basal conditions (Wang et al, 2018) and may play a role in CDI (Simms & Zamponi, 2014). To study the structure of the half-saturated CaM intermediate state, we first constructed a mutant protein (D21A/D23A/

*To whom correspondence should be addressed: jbames@ucdavis.edu.

D25A/E32Q/D57A/D59A/N61A/E68Q, called CaM^{EF12}) that disables Ca²⁺ binding to EF1 and EF2 but retains Ca²⁺-binding solely at EF3 and EF4. This CaM intermediate state is also important for regulation of ryanodine receptors (Sondergaard et al, 2015) and nitric oxide synthase (Piazza et al, 2016). We report detailed NMR resonance assignments for the CaM^{EF12} mutant with two Ca²⁺ bound (called Ca²⁺₂/CaM^{EF12}) that is bound to the IQ-motif of Ca_v1.2 (the complex is called Ca²⁺₂/CaM^{EF12}-IQ) as a first step toward solving the structure of this complex.

Methods and Experiments

Preparation of Ca²⁺₂/CaM^{EF12} bound to the Ca_v1.2 IQ-motif.

A cDNA of *Homo sapiens* calmodulin that contained the mutations (D21A/D23A/D25A/E32Q/D57A/D59A/N61A/E68Q) was subcloned into pET-11b vector (Novagen) that produced recombinant CaM^{EF12} without any affinity tag or extra residues. Uniformly ¹⁵N-labeled and ¹³C,¹⁵N-labeled CaM^{EF12} were expressed in bacterial cells grown on M9 minimal media supplemented with ¹⁵N-labeled NH₄Cl (0.5 g per liter of cell culture) and ¹³C-labeled glucose (3 g per liter). The recombinant CaM^{EF12} protein was purified using Ca²⁺-dependent hydrophobic interaction chromatography (Gopalakrishna & Anderson, 1982). A peptide fragment of the Ca_v1.2 IQ-motif (residues 1642–1665) was purchased from GenScript, dissolved in DMSO-d₆ and quantified using UV-Vis absorption. A 1.5-fold excess of Ca_v1.2 IQ-motif was added to Ca²⁺-bound CaM^{EF12} and the complex (Ca²⁺₂/CaM^{EF12}-IQ) was subsequently concentrated to 500 μM in the presence of 2 mM CaCl₂ using a 3K Amicon concentrator.

NMR spectroscopy.

Samples of Ca²⁺₂/CaM^{EF12}-IQ complex for NMR analysis were prepared by exchanging the protein complex into a buffer containing 20 mM Tris-d₁₁ (pH 7.5) with 1 mM CaCl₂, and 92% H₂O/8% D₂O. All NMR experiments were performed at 303K on a Bruker Avance 800 MHz spectrometer equipped with a four channel interface and triple resonance cryogenic (TCI) probe. The ¹⁵N-¹H HSQC spectrum (Fig. 1) was recorded with 256 × 2048 complex points for ¹⁵N(F1) and ¹H(F2), respectively. Assignment of backbone resonances was obtained by analyzing the following spectra: HNCA, HNCACB, CBCA(CO)NH, HNCO (Ikura et al, 1990). Side chain resonances were assigned by analyzing HCCH-TOCSY (Ikura et al, 1991). The NMR data were processed using NMRPipe (Delaglio et al, 1995) and analyzed using Sparky NMRFAM (Lee et al, 2015).

Assignments and Data Deposition

A two-dimensional ¹⁵N-¹H HSQC NMR spectrum of Ca²⁺₂/CaM^{EF12} bound to unlabeled Ca_v1.2 IQ peptide illustrates representative NMR assignments (Fig. 1). NMR assignments were based on 3D heteronuclear NMR spectra recorded from ¹³C/¹⁵N-labeled Ca²⁺₂/CaM^{EF12} bound to unlabeled IQ peptide (residues 1642–1665). The NMR spectra of Ca²⁺₂/CaM^{EF12}-IQ exhibited well-dispersed peaks with uniform intensities indicative of a stably folded protein complex. A few amide resonances exhibit noteworthy downfield shifts, including T29, I101, and N138 that are each located in β-strands (see arrows in Fig. 2) and

are expected to form antiparallel β -sheets with strong backbone amide hydrogen bonds. Two downfield peaks assigned to G99 (EF3) and G135 (EF4) are likely caused by a strong hydrogen bond between the backbone NH of G99 and G135 with the side chain carboxyl group of D94 and D130, respectively caused by Ca^{2+} binding to EF3 and EF4. The absence of corresponding downfield shifts for G26 (EF1) and G62 (EF2) is consistent with a lack of Ca^{2+} binding at EF1 and EF2. More than 92% of the main chain ^{13}C resonances ($^{13}\text{C}\alpha$, $^{13}\text{C}\beta$, and ^{13}CO), 90% of backbone amide resonances (^1HN , ^{15}N), and 74% of methyl side chain resonances were assigned. The unassigned residues (14–15, 30–32, 43, 59–61, 75, 78, 94) had overlapped backbone amide resonances and/or weak NMR intensities that obscured their assignment. In particular, the EF-hand loop residues in EF1 (residues 30–32) and EF2 (residues 58–61) could not be assigned due to weak NMR intensities, perhaps caused by conformational disorder in these Ca^{2+} -free binding loops. A complete listing of the chemical shift assignments (^1H , ^{15}N , ^{13}C) of $\text{Ca}^{2+}_2/\text{CaM}^{\text{EF12}}$ bound to unlabeled $\text{Ca}_v1.2$ IQ peptide have been deposited in the BioMagResBank (<http://www.bmrb.wisc.edu>) under accession number 27692.

The secondary structure of $\text{Ca}^{2+}_2/\text{CaM}^{\text{EF12}}$ in the complex was calculated based on the chemical shift index (Wishart et al, 1992) of each assigned amino acid residue and ANN-secondary structure prediction using TALOS (Shen et al, 2009) (Fig. 2). As expected, $\text{Ca}^{2+}_2/\text{CaM}^{\text{EF12}}$ contains four EF-hands comprised of eight α -helices. The helices are named $\alpha 1$ (residues 8–20), $\alpha 2$ (residues 30–39), $\alpha 3$ (residues 46–56), $\alpha 4$ (residues 66–76), $\alpha 5$ (residues 83–93), $\alpha 6$ (residues 103–113), $\alpha 7$ (residues 119–129) and $\alpha 8$ (residues 139–146) depicted by blue rectangles in Fig. 2. In addition, a short and conserved β -strand was observed in EF1 (residues 27–29), EF2 (residues 63–65), EF3 (residues 100–102) and EF4 (residues 136–138) shown by orange arrows in Fig. 2. The overall secondary structure of $\text{Ca}^{2+}_2/\text{CaM}^{\text{EF12}}$ is similar to that observed in the crystal structure of Ca^{2+} -saturated CaM bound to IQ peptide (Van Petegem et al, 2005). A plot of the amide chemical shift perturbation caused by the binding of the IQ peptide reveals that CaM^{EF12} residues in the N-terminal domain (residues 1–80 in EF1 and EF2) exhibit relatively small chemical shift differences (CSDs) upon binding to the IQ peptide (Fig. 3). In stark contrast, the CaM residues in the C-terminal domain (residues 81–149) exhibit much larger CSDs induced by IQ binding (Fig. 3). The larger CSD values for the C-terminal domain residues suggest that the IQ-motif is bound to the C-terminal domain (EF3 and EF4). The relatively smaller CSD values for the N-terminal domain residues suggest that the IQ peptide may not make contact with the CaM N-terminal domain (EF1 and EF2), in contrast to what is seen in the crystal structure of Ca^{2+} -saturated CaM bound to IQ. The NMR assignments of $\text{Ca}^{2+}_2/\text{CaM}^{\text{EF12}}$ bound to the $\text{Ca}_v1.2$ IQ-motif presented here are an important first step toward determining the full three-dimensional structure of $\text{Ca}^{2+}_2/\text{CaM}^{\text{EF12}}$ bound to $\text{Ca}_v1.2$.

Acknowledgements

We thank Jeff Walton for technical support and help with NMR experiments. Work supported by NIH grants (EY012347) to J.B.A and (AG055357) to J.W.H.

References

- Adams PJ, Ben-Johny M, Dick IE, Inoue T, Yue DT (2014) Apocalmodulin itself promotes ion channel opening and Ca(2+) regulation. *Cell* 159: 608–622 [PubMed: 25417111]
- Ben Johny M, Yang PS, Bazzazi H, Yue DT (2013) Dynamic switching of calmodulin interactions underlies Ca2+ regulation of CaV1.3 channels. *Nature communications* 4: 1717
- Delaglio F, Grzesiek S, Vuister GW, Zhu G, Pfeiffer J, Bax A (1995) NMRPipe: a multidimensional spectral processing system based on UNIX pipes. *J Biomol NMR* 6: 277–293 [PubMed: 8520220]
- Erickson M, Alseikhan B, Peterson B, Yue D (2001) Preassociation of calmodulin with voltage-gated Ca(2+) channels revealed by FRET in single living cells. *Neuron* 31: 973–985 [PubMed: 11580897]
- Findeisen F, Rumpf CH, Minor DL, Jr. (2013) Apo states of calmodulin and CaBP1 control CaV1 voltage-gated calcium channel function through direct competition for the IQ domain. *Journal of molecular biology* 425: 3217–3234 [PubMed: 23811053]
- Gabelli SB, Boto A, Kuhns VH, Bianchet MA, Farinelli F, Aripirala S, Yoder J, Jakoncic J, Tomaselli GF, Amzel LM (2014) Regulation of the NaV1.5 cytoplasmic domain by calmodulin. *Nature communications* 5: 5126
- Gopalakrishna R, Anderson WB (1982) Ca2+-induced hydrophobic site on calmodulin: application for purification of calmodulin by phenyl-Sepharose affinity chromatography. *Biochem Biophys Res Commun* 104: 830–836 [PubMed: 6803791]
- Hell JW, Westenbroek RE, Warner C, Ahljianian MK, Prystay W, Gilbert MM, Snutch TP, Catterall WA (1993) Identification and differential subcellular localization of the neuronal class C and class D L-type calcium channel α 1 subunits. *J Cell Biol* 123: 949–962 [PubMed: 8227151]
- Ikura M, Kay LE, Bax A (1990) A novel approach for sequential assignment of ^1H , ^{13}C , and ^{15}N spectra of proteins: heteronuclear triple-resonance three-dimensional NMR spectroscopy. Application to calmodulin. *Biochemistry* 29: 4659–4667 [PubMed: 2372549]
- Ikura M, Spera S, Barbato G, Kay LE, Krinks M, Bax A (1991) Secondary structure and side-chain ^1H and ^{13}C resonance assignments of calmodulin in solution by heteronuclear multidimensional NMR spectroscopy. *Biochemistry* 30: 9216–9228 [PubMed: 1909892]
- Lee W, Tonelli M, Markley JL (2015) NMRFAM-SPARKY: enhanced software for biomolecular NMR spectroscopy. *Bioinformatics* 31: 1325–1327 [PubMed: 25505092]
- Moosmang S, Haider N, Klugbauer N, Adelsberger H, Langwieser N, Muller J, Stiess M, Marais E, Schulla V, Lacinova L, Goebbels S, Nave K, Storm D, Hofmann F, Kleppisch T (2005) Role of hippocampal Cav1.2 Ca2+ channels in NMDA receptor-independent synaptic plasticity and spatial memory. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 25: 9883–9892 [PubMed: 16251435]
- Piazza M, Dieckmann T, Guillemette JG (2016) Structural Studies of a Complex Between Endothelial Nitric Oxide Synthase and Calmodulin at Physiological Calcium Concentration. *Biochemistry*
- Shen Y, Delaglio F, Cornilescu G, Bax A (2009) TALOS+: a hybrid method for predicting protein backbone torsion angles from NMR chemical shifts. *J Biomol NMR* 44: 213–223 [PubMed: 19548092]
- Simms BA, Zamponi GW (2014) Neuronal voltage-gated calcium channels: structure, function, and dysfunction. *Neuron* 82: 24–45 [PubMed: 24698266]
- Sondergaard MT, Tian X, Liu Y, Wang R, Chazin WJ, Chen SR, Overgaard MT (2015) Arrhythmogenic Calmodulin Mutations Affect the Activation and Termination of Cardiac Ryanodine Receptor-mediated Ca2+ Release. *J Biol Chem* 290: 26151–26162 [PubMed: 26309258]
- Van Petegem F, Chatelain FC, Minor DL, Jr. (2005) Insights into voltage-gated calcium channel regulation from the structure of the CaV1.2 IQ domain-Ca2+/calmodulin complex. *Nature structural & molecular biology* 12: 1108–1115
- Vogl AM, Brockmann MM, Giusti SA, Maccarrone G, Vercelli CA, Bauder CA, Richter JS, Roselli F, Hafner AS, Dedic N, Wotjak CT, Vogt-Weisenhorn DM, Choquet D, Turck CW, Stein V, Deussing JM, Refojo D (2015) Neddylation inhibition impairs spine development, destabilizes synapses and deteriorates cognition. *Nature neuroscience* 18: 239–251 [PubMed: 25581363]

- Wang K, Holt C, Lu J, Brohus M, Larsen K, Overgaard M, Wimmer R, Van Petegem F (2018) Arrhythmia mutations in calmodulin cause conformational changes that affect interactions with the cardiac voltage-gated calcium channel. *Proceedings of the National Academy of Sciences of the United States of America* 115: E10556–E10565 [PubMed: 30348784]
- Wishart DS, Sykes BD, Richards FM (1992) The chemical shift index: a fast and simple method for the assignment of protein secondary structure through NMR spectroscopy. *Biochemistry* 31: 1647–1651. [PubMed: 1737021]
- Zuhlke RD, Pitt GS, Deisseroth K, Tsien RW, Reuter H (1999) Calmodulin supports both inactivation and facilitation of L-type calcium channels. *Nature* 399: 159–162 [PubMed: 10335846]

Fig. 1: Two-dimensional ^{15}N - ^1H HSQC NMR spectrum of ^{15}N -labeled $\text{Ca}^{2+}_2/\text{CaM}^{\text{EF12}}$ bound to unlabeled $\text{Ca}_v1.2$ IQ peptide at pH 7.5 recorded at 800-MHz ^1H frequency (A). Expanded view of the crowded spectral region in the center of the spectrum (B). Constant-time ^{13}C - ^1H HSQC spectrum of ^{13}C -labeled $\text{Ca}^{2+}_2/\text{CaM}^{\text{EF12}}$ bound to unlabeled $\text{Ca}_v1.2$ IQ peptide (C). Representative assignments are indicated; complete assignments are available as BMRB accession no. 27692.

Fig. 2: Secondary structure of $\text{Ca}^{2+}_2/\text{CaM}^{\text{EF12}}$ bound to $\text{Ca}_v1.2$ IQ. Normalized order parameter (upper panel) and ANN-secondary structure probability (lower panel) plotted as a function of residue number. Secondary structure elements (rectangle for helix and arrow for β -strand) were calculated on the basis of chemical shift index and sequential NOE patterns.

Fig. 3:

Amide chemical shift perturbation for $\text{Ca}^{2+}_2/\text{CaM}^{\text{EF12}}$ in the presence versus absence of bound IQ peptide. The chemical shift difference (CSD) was calculated as $\text{CSD} = \{(\text{HN}_\text{A} - \text{HN}_\text{B})^2 + ({}^{15}\text{N}_\text{A} - {}^{15}\text{N}_\text{B})^2\}^{1/2}$. HN_A and HN_B are amide proton chemical shift of $\text{Ca}^{2+}_2/\text{CaM}^{\text{EF12}}$ in the presence and absence of IQ peptide respectively, and ${}^{15}\text{N}_\text{A}$ and ${}^{15}\text{N}_\text{B}$ are amide ${}^{15}\text{N}$ chemical shift of $\text{Ca}^{2+}_2/\text{CaM}^{\text{EF12}}$ in the presence and absence of IQ peptide respectively. Chemical shift assignments for $\text{Ca}^{2+}_2/\text{CaM}^{\text{EF12}}$ (without IQ) were derived from wildtype apo-CaM (BMRB 27190) for N-terminal domain residues (1–79) and from wild type Ca^{2+} -saturated CaM (BMRB 547) for the C-terminal domain residues (80–149). The CSD values of the mutated residues in EF1 and EF2 (21, 23, 25, 57, 59, and 61) are not meaningful (due to the mutation) and are not shown.