Published in final edited form as:

Chem Asian J. 2019 February 15; 14(4): 532–536. doi:10.1002/asia.201801732.

Photoredox-Catalyzed Alkenylation of Benzylsulfoniums

Shinya Otsuka^{a,b}, Dr. Keisuke Nogi^a, Prof. Dr. Tomislav Rovis^b, and Prof. Dr. Hideki Yorimitsu^a

^aDepartment of Chemistry, Graduate School of Science, Kyoto University, Sakyo-ku, Kyoto 606-8502 (Japan)

^bDepartment of Chemistry, Columbia University, 3000 Broadway, New York, NY 10027 (USA)

Abstract

Visible light-mediated radical alkenylation of benzylsulfoniums was achieved by means of *fac*-Ir(ppy)₃ as a photocatalyst, giving allylbenzenes as products. A variety of functional groups such as halogen, ester, and cyano groups were well tolerated in this transformation. Starting benzylsulfoniums could be readily prepared from benzyl alcohols via an acid-mediated substitution, increasing the synthetic utility of this transformation.

Graphical Abstract:

Visible light-mediated radical alkenylation of benzylsulfoniums was achieved by means of *fac*-Ir(ppy)₃ as a photocatalyst, giving allylbenzenes as products. A variety of functional groups such as halogen, ester, and cyano groups were well tolerated in this transformation. Starting benzylsulfoniums could be readily prepared from benzyl alcohols via an acid-mediated substitution, increasing the synthetic utility of this transformation.

Keywords

photoredox catalysis; alkenylation; sulfonium; radical reaction

Organosulfur compounds widely exist in natural compounds, living bodies, pharmaceuticals, and functional materials.^[1] From the viewpoint of organic synthesis, organosulfur compounds can be regarded as important synthetic intermediates, due to their unique reactivities that organohalogen compounds do not have. In order to utilize such characteristics for organic synthesis, recently, transition metal-catalyzed transformations of organosulfur compounds with cleavage of their C–S bonds have been actively explored.^[2]

Publisher's Disclaimer: This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

Among various transition metal-catalyzed transformations, the Mizoroki–Heck alkenylation is one of the most important reactions in organic synthesis.^[3,4] However, Mizoroki–Heck-type alkenylation of organosulfur compounds has been left unexplored due to the strongly coordinating and potentially catalyst-poisoning nature of leaving anionic sulfur moieties. In order to overcome these problems, utilization of arylsulfoniums instead of aryl sulfides is beneficial because 1) oxidative addition of arylsulfoniums is easier than that of aryl sulfides due to their electron-deficient nature; and 2) leaving neutral sulfides would be less debilitating to the catalyst. Taking advantage of these benefits, Zhang's group and our group recently succeeded in developing palladium-catalyzed Mizoroki–Heck-type reactions of arylsulfoniums.^[5] Unfortunately, however, these methods were applicable only to arylsulfoniums, and alkylsulfoniums did not react at all. In order to convert alkylsulfoniums to the corresponding alkene products, we then turned our strategy to photoredox-catalyzed radical-mediated alkenylation.^[6] Recently Lei and co-workers reported photoredox-catalyzed^[7] radical-mediated alkenylation of benzyl bromides under very mild reaction conditions.^[8] We envisioned that Lei's alkenylation would be applicable to benzylsulfoniums. Our working hypothesis is shown in Scheme 1: **A**) photoexcited catalyst reduces benzylsulfonium to generate benzyl radical and dialkyl sulfide, **B**) addition of the benzyl radical to alkene takes place, **C**) single electron transfer from the radical adduct to the oxidized photocatalyst generates cation **I**, and **D**) deprotonation affords alkenylation product. Indeed, various benzylsulfoniums and alkenes could be coupled under visible light irradiation. It is worth mentioning that electrochemical,^[9] chemical,^[10] and photosensitized^[11,12] formations of aryl or alkyl radicals from sulfonium salts have been well studied, while its application to organic synthesis has been limited to α -carbonyl or α -cyanosulfoniums,^[10,12] triarylsulfoniums,^[13] and 1-trifluoromethyldibenzothiophenium (Umemoto reagent).^[14]

In the presence of 2 mol% of *fac*-Ir(ppy)₃ as a photocatalyst, the reaction of 1-benzyltetrahydrothiophenium bromide (**1a**) with 1,1-diphenylethylene (**2a**) was tested under blue LED irradiation. By using Na₂CO₃ as a base and DMF as a solvent, a 39% yield of the alkenylated product **3aa** was obtained (Table 1, entry 1). The use of K₂CO₃ instead of Na₂CO₃ gave the product **3aa** in a slightly lower yield (entry 2). NaOAc was found to be inferior, and **3aa** was not obtained when Cs₂CO₃ was used^[15] (entries 3 and 4). Organic bases such as Et₃N, *i*-Pr₂EtN, and DABCO resulted in no or negligible formations of the desired product (See Table S1 in the Supporting Information). By increasing the amount of **2a** to 2 equivalents, the yield of **3aa** increased to 52% (entry 5). Since the reduction potential of the sulfonium salt **1a** is –1.48 V (vs SCE, see Figure S1 in the supporting information), strongly reducing *fac*-Ir(ppy)₃ (Ir^{III}*/Ir^{IV} = –1.73 V)^[16] was crucial. As another photocatalyst, [Ir(ppy)₂(dtbpy)](PF₆) having a higher reduction potential (Ir^{III}*/Ir^{IV} = –0.96 V)^[17] did not give **3aa** at all (entry 6). The result of solvent screening revealed that NMP was the best solvent (entry 7 and Table S1). Eventually, the reaction in higher concentration (0.5 M) furnished **3aa** in 71% yield (entry 8). In the absence of either photocatalyst or light, **3aa** was not obtained at all (entries 9 and 10).^[18]

With the optimized reaction conditions in hand, we then tried this alkenylation with various benzylsulfoniums (Scheme 2). The efficiency of the alkenylation depends on the electronic

nature of benzylsulfoniums: the yields of the products from electron-deficient sulfoniums were higher than that from **1a**. Functional groups such as ester and cyano groups were well tolerated in this reaction, giving **3ba**, **3ca**, and **3da**, respectively. A methyl substituent on the *ortho* position in **1e** did not retard the reaction. Electron-deficient sulfoniums **1f** and **1g** afforded **3fa** and **3ga** in high yields, respectively. Bromo-substituted benzylsulfoniums **1h** and **1i** were successfully converted to **3ha** and **3ia** without touching their aryl–Br bonds. Substituents on the *meta* positions did not hamper the reaction; products **3ja** and **3ka** were obtained in good yields. π -Extended naphthylsulfoniums **1l** and **1m** also gave alkenylation products **3la** and **3ma**.

The scope of alkenes was also investigated (Scheme 3). 1,1-Diphenylethylenes bearing electron-withdrawing fluoro or electron-donating methoxy groups participated in the reaction, giving **3ib** or **3bc** in good yields. The methylsulfanyl groups in **2e** left untouched in the reaction and **3be** was obtained in 77% yield. The alkenylation of **1b** with unsymmetrical 1,1-diarylethylenes **2f** and **2g** afforded *E/Z* mixtures of **3bf** and **3bg** in 87% and 84% yields, respectively. Styrene derivatives also underwent the present alkenylation and the reaction of **1i** with 4-methoxystyrene (**2h**) afforded 1,3-diarylpropene **3ih** in 66% yield as a mixture of *E* and *Z* isomers.^[19] Under the standard conditions, less electronrich 4-acetoxystyrene (**2i**) gave the desired product **3bi** in 20% yield along with hydrated alcohol **3bi'** in 20% yield. We speculated that benzylic cation **I** in Scheme 1 would be trapped by a trace amount of water in the reaction mixture to generate **3bi'**.^[14a,l] Therefore we ran the reaction in the presence of tetrabutylammonium bromide (TBAB) in order to enhance the deprotonation from **I** by increasing the solubility of the base. As a result, the formation of **3bi'** was suppressed and **3bi** was obtained in 37% yield. The alkenylation with simple styrene in the presence of TBAB gave **3bj** while the yield was lower than that of electron-rich styrenes. The reaction with α -methylstyrene (**2k**) also proceeded while a mixture of three isomers, **3ik'**, (*E*)-**3ik**, and (*Z*)-**3ik**, was obtained.

We then checked the effect of the counteranion of benzylsulfoniums. Instead of benzylsulfonium bromide **1i**, benzylsulfonium triflate **4i** was involved in the reaction (Scheme 4a). As a result, not only the desired product **3ia** but also tertiary alcohol **3ia'** was observed in the crude mixture. We inferred that the difference would be derived from the efficiency of the step **D** in Scheme 1. When **1i** is employed, counteranion of the benzylic cation **I** should be Br[−] or NaCO₃[−] (by generating NaBr) and the anion would deprotonate **I** to afford **3ia**. On the other hand, in the case of **4i**, far less basic TfO[−] would be reluctant to deprotonate **I**. As a result, the reaction of **I** with water afforded **3ia'**. An addition of TBAB was again effective to accelerate the deprotonation and suppress the formation of **3ia'**, as is the case with **3bi** and **3bj**, affording **3ia** in 72% yield (Scheme 4b, upper). Additionally, we tried to obtain **3ia'** selectively by running the reaction in the presence of water. As expected, **3ia'** was obtained in 65% yield (Scheme 4b, lower). The reaction in the presence of MeOH gave ether **3ia''** in 58% yield. Interestingly, formation of **3ia'** is less efficient when 2-bromobenzyl bromide was used instead of **4i**: the yield of **3ia'** was only 27% and **3ia** was the major product. Therefore, it can be emphasized that the synthesis of **3ia'** or **3ia''** is one feature of using benzylsulfoniums.

Since 1-benzyltetrahydrothiophenium triflates **4** can be synthesized from benzyl alcohols and tetrahydrothiophene,^[20] the scope of 1-benzyltetrahydrothiophenium triflates was also examined (Scheme 5). The sulfonium salt prepared from 2-chlorobenzyl alcohol could be alkenylated to afford **3na** in 69% yield. Electron-rich benzylsulfonium **4o** was also converted to **3oa** in 54% yield although the yield was lower than those from electron-deficient ones. It is noteworthy that this method allows one to avoid the use of unstable electron-rich benzyl bromides. Alkenylation of 2-arylbenzylsulfonium **4q** proceeded in spite of the steric hindrance, giving **3qa** in 70% yield. This example shows usefulness of this reaction (Scheme 6); while palladium-catalyzed cross-coupling of 2-bromobenzyl bromide with arylboronic acids was reported to take place at the benzylic C–Br bond (Scheme 6a),^[21] that of 2-bromobenzyl alcohols took place at the aromatic C–Br bonds without any protection of the hydroxy moiety.^[22] After the arylation, the hydroxy group was “activated” to the corresponding sulfonium triflate and then sulfonium **4g** was alkenylated under the photoredox catalysis. This arylation–activation–alkenylation sequence would provide a new synthetic strategy for allylbenzene derivatives (Scheme 6b).

Finally, we conducted the present alkenylation in a 1.5 mmol scale; sulfonium bromide **1b** and sulfonium triflate **4i** reacted with **2a**, respectively. In both cases, the corresponding alkenylation products **3ba** and **3ia** were obtained without loss of the yields (Scheme 7).

In conclusion, we have developed alkenylation of benzylsulfoniums by means of photoredox catalysis. The starting benzylsulfonium triflates could be prepared from the corresponding benzyl alcohols, allowing this transformation valuable. Further development on photoredox-mediated transformations of organosulfur compounds is currently underway in our laboratories.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

This work was supported by JSPS KAKENHI Grant Numbers JP16H04109, JP18H04254, JP18H04409, and JP18K14212. S.O. acknowledges JSPS Predoctoral Fellowship and JSPS Overseas Challenge Program for Young Researchers. T.R. thanks NIGMS (GM125206).

References

- [1]. a) An Introduction to Organosulfur Chemistry (Ed.: Cremllyn RJ), Wiley, Hoboken, 1996b) Organosulfur Chemistry I (Ed.: Page PCB), Springer, Heidelberg, 1999;c) Organosulfur Chemistry II, (Ed.: Page PCB), Springer, Heidelberg, 1999;d) Advances in Sulfur Chemistry, Vol. 2 (Ed.: Rayner CM), JAI Press, Greenwich, 2000;e) Sulfur Compounds: Advances in Research and Application (Ed.: Acton AQ), Scholarly Eds., Atlanta, 2012.
- [2]. a)Reviews on catalytic C–S bond transformations of organosulfur compounds including aryl sulfides: Dubbaka SR, Vogel P, Angew. Chem. Int. Ed 2005, 44, 7674; Angew. Chem. 2005, 117, 7848;b) Wang L, He W, Yu Z, Chem. Soc. Rev 2013, 42, 599; [PubMed: 23079733] c) Modha SG, Mehta VP, Van der Eycken EV, Chem. Soc. Rev 2013, 42, 5042; [PubMed: 23467811] d) Pan F, Shi Z-J, ACS Catal. 2014, 4, 280;e) Gao K, Otsuka S, Baralle A, Nogi K, Yorimitsu H, Otsuka A, J. Synth. Org. Chem. Jpn 2016, 74, 1119;f) Ortgies DH, Hassanpour A, Chen F, Woo S,

- Forgione P, Eur. J. Org. Chem 2016, 408;g) Otsuka S, Nogi K, Yorimitsu H, Top. Curr. Chem 2018, 376, 13.
- [3]. a) Mizoroki T, Mori K, Ozaki, Bull. Chem. Soc. Jpn 1971, 44, 581;b) Heck RF, Nolley JP, Jr., J. Org. Chem 1972, 37, 2320.
- [4]. a) Bräse S, de Meijere A in Metal-Catalyzed Cross-Coupling Reactions, 2nd ed. (Eds.: de Meijere A, Diederich F), Wiley-VCH, Weinheim, 2004, Chap. 5;b) Beller M, Zapf A, Riermeier TH in Transition Metals for Organic Synthesis: Building Blocks and Fine Chemicals, 2nd ed. (Eds.: Beller M, Bolm C), Wiley-VCH, Weinheim, 2004, Chap. 2.13;c) The Mizoroki-Heck Reaction (Ed.: Oestreich M), Wiley, Hoboken, 2009;d) Beletskaya IP, Cheprakov AV, Chem. Rev 2000, 100, 3009; [PubMed: 11749313] e) Dounay AB, Overman LE, Chem. Rev 2003, 103, 2945. [PubMed: 12914487]
- [5]. a) Wang S-M, Song H-X, Wang X-Y, Liu N, Qin H-L, Zhang C-P, Chem. Commun 2016, 52, 11893;b) Uno D, Minami H, Otsuka S, Nogi K, Yorimitsu H, Chem. Asian. J 2018, 13, 2397. [PubMed: 29729085]
- [6]. a) Selected examples of transition metal-catalyzed radical-mediated alkenylation: Ikeda Y, Nakamura T, Yorimitsu H, Oshima K, J. Am. Chem. Soc 2002, 124, 6514; [PubMed: 12047154] b) Terao J, Watabe H, Miyamoto M, Kambe N, Bull. Chem. Soc. Jpn 2003, 76, 2209;c) Liu C, Tang S, Liu D, Yuan J, Zheng L, Meng L, Lei A, Angew. Chem. Int. Ed 2012, 51, 3638; Angew. Chem. 2012, 124, 3698;d) Zhang X, Yi H, Liao Z, Zhang G, Fan C, Qin C, Liu J, Lei A, Org. Biomol. Chem 2014, 12, 6790; [PubMed: 25065706] e) Zhu K, Dunne J, Shaver MP, Thomas SP, ACS Catal. 2017, 7, 2353.
- [7]. a) Reviews: Visible Light Photocatalysis in Organic Chemistry (Eds.: Stephenson CRJ, Yoon TP, MacMillan DWC), Wiley, Hoboken, 2018;b) Zeitler K, Angew. Chem. Int. Ed 2009, 48, 9785; Angew. Chem. 2009, 121, 9969;c) Yoon TP, Ischay MA, Du J, Nat. Chem 2010, 2, 527; [PubMed: 20571569] d) Narayanam JMR, Stephenson CRJ, Chem. Soc. Rev 2011, 40, 102; [PubMed: 20532341] e) Xuan, Xiao W-J, Angew. Chem. Int. Ed 2012, 124, 6934; Angew. Chem. 2012, 51, 6828;f) Tucker JW, Stephenson CRJ, J. Org. Chem 2012, 77, 1617; [PubMed: 22283525] g) Pier CK, Rankic DA, MacMillan DWC, Chem. Rev 2013, 113, 5322. [PubMed: 23509883]
- [8]. Liu Q, Yi H, Liu J, Yang Y, Zhang X, Zeng Z, Lei A, Chem. Eur. J 2013, 19, 5120. [PubMed: 23426910]
- [9]. a) Bonner WA, J. Am. Chem. Soc 1952, 74, 5078;b) Colichman EL, Love DL, J. Org. Chem 1953, 18, 40;c) Finkelstein M, Petersen RC, Ross SD, J. Electrochem. Soc 1963, 110, 422;d) Anderson JD, Baizer MM, Petrovich JP, J. Org. Chem 1966, 31, 3890;e) Wagenknecht JH, Baizer MM, J. Electrochem Soc 1967, 114, 1095;f) Moses PR, Chambers JQ, J. Electroanal. Chem 1974, 49, 105;g) Chambers JQ, Maupin PH, II, Liao C-S, J. Electroanal. Chem 1978, 87, 239;h) Beak P, Sullivan TA, J. Am. Chem. Soc 1982, 104, 4450;i) Saeva FD, Morgan BP, J. Am. Chem. Soc 1984, 106, 4121.
- [10]. a) Kataoka T, Tsutsumi K, Kano K, Mori K, Miyake M, Yokota M, Shimizu H, Hori M, J. Chem. Soc., Perkin Trans 1 1990, 3017;b) Kataoka T, Iwama T, Shimizu H, Hori M, Phosphorus, Sulfur, and Silicon, 1992, 67, 169.
- [11]. Wang X, Saeva FD, Kampmeier JA, J. Am. Chem. Soc 1999, 121, 4364.
- [12]. a) Hedstrand DM, Kruizinga WH, Kellogg RM, Tetrahedron Lett. 1978, 14, 1255;b) Van Bergen TJ, Hedstrand DM, Kruizinga WH, Kellogg RM, J. Org. Chem 1979, 44, 4953.
- [13]. Donck S, Baroudi A, Fensterbank L, Goddard J-P, Ollivier C, Adv. Synth. Catal 2013, 355, 1477.
- [14]. a) Yasu Y, Koike T, Akita M, Angew. Chem. Int. Ed 2012, 51, 9567; Angew. Chem. 2012, 124, 9705;b) Mizuta S, Verhoog S, Engle KM, Khotavivattana T, O'Duill M, Wheelhouse K, Rassias G, Médebielle M, Gouverneur V, J. Am. Chem. Soc 2013, 135, 2505; [PubMed: 23373772] c) Yasu Y, Koike T, Akita M, Org. Lett 2013, 15, 2136; [PubMed: 23600821] d) Ji Y-L, Lin J-H, Xiao J-C, Gu Y-C, Org. Chem. Front 2014, 1280;e) Ge G-C, Huang X-J, Ding C-H, Wan S-L, Dai L-X, Hou X-L, Chem. Commun 2014, 50, 3048;f) Yu J, Wang Y-L, Gong T-J, Xiao B, Fu Y, Chem. Commun 2014, 50, 12915;g) Yasu Y, Arai Y, Tomita R, Koike T, Akita M, Org. Lett 2014, 16, 780; [PubMed: 24422891] h) Tomita R, Yasu Y, Koike T, Akita M, Beilstein J. Org. Chem 2014, 10, 1099; [PubMed: 24991259] i) Noto N, Miyazawa, Koike T, Akita M, Org. Lett 2015, 17, 3710; [PubMed: 26181341] j) Asano M, Tomita R, Koike T, Akita M, J. Fluorine Chem,

2015, 179, 83;k) Noto N, Koike T, Akita M, J. Org. Chem 2016, 81, 7064; [PubMed: 27304729]
l) Tomita R, Koike T, Akita M, Chem. Commun 2017, 53, 4681.

- [15]. When Cs₂CO₃ was used as a base, the crude mixture was messy. Strongly basic Cs₂CO₃ would have generated the corresponding sulfonium ylide. For generation of sulfonium ylides from benzylsulfoniums, see: Zhou R, Deng X, Zheng J, Sheng Q, Sun X, Tang Y, Chin. J. Chem 2011, 29, 995.
- [16]. Flamigni L, Barbieri A, Sabatini C, Ventura B, Barigelletti F, Top. Curr. Chem 2007, 281, 143.
- [17]. Lowry MS, Goldsmith JI, Slinker JD, Rohl R, Pascal RA, Malliaras GG, Bernhard S, Chem. Mater 2005, 17, 5712.
- [18]. A light on-off experiment was conducted to understand the reaction mechanism. As shown in Figure S2 in the Supporting Information, the alkenylation proceeded only when blue LED was irradiated. This result suggests that the reaction would not proceed through radical chain mechanism.
- [19]. The E/Z ratio of 3ih would depend on the reaction time. With a longer reaction time, the ratio of the E isomer of 3ih was increased probably due to E/Z isomerization.
- [20]. See the Supporting Information and Scheme 6.
- [21]. Langle S, Abarbri M, Duchêne A, Tetrahedron Lett. 2003, 44, 9255.
- [22]. Wolfe JP, Singer RA, Yang BH, Buchwald SL, J. Am. Chem. Soc 1999, 121, 9550.

Scheme 1.

Working hypothesis of this work.

Scheme 2.
Scope of benzylsulfonium bromides.

Scheme 3. Scope of alkenes. [a] With 1.0 equiv of *n*Bu₄NBr.

(a) Reaction of benzylsulfonium triflate **4i** in the standard conditions

(b) Divergent synthesis from benzylsulfonium triflate **4i**

Scheme 4.
Reaction of benzylsulfonium triflate **4i**.

Scheme 5.
Scope of benzylsulfonium triflates.

a) Arylation of 2-bromobenzyl bromide (ref. 21)

b) Arylation–activation–alkenylation sequence

Scheme 6.

Synthesis of **3qa** starting from 2-bromobenzyl alcohol.

Scheme 7.
Scale-up reactions.

Table 1.

Optimization of reaction conditions.^[a]

entry	catalyst	base	solvent	NMR yield
1	<i>fac</i> -Ir(ppy) ₃	Na ₂ CO ₃	DMF	39%
2	<i>fac</i> -Ir(ppy) ₃	K ₂ CO ₃	DMF	34%
3	<i>fac</i> -Ir(ppy) ₃	NaOAc	DMF	16%
4	<i>fac</i> -Ir(ppy) ₃	CS ₂ CO ₃	DMF	0%
5	<i>fac</i> -Ir(ppy) ₃	Na ₂ CO ₃	DMF	52% ^[b]
6	[Ir(ppy) ₂ (dtbpy)](PF ₆)	Na ₂ CO ₃	DMF	0% ^[b]
7	<i>fac</i> -Ir(ppy) ₃	Na ₂ CO ₃	NMP	64% ^[b]
8	<i>fac</i> -Ir(ppy) ₃	Na ₂ CO ₃	NMP	71% ^[b,c]
9	—	Na ₂ CO ₃	NMP	0% ^[b,c]
10	<i>fac</i> -Ir(ppy) ₃	Na ₂ CO ₃	NMP	0% ^[b,c,d]

^[a] **1a** (0.50 mmol), **2a** (0.50 mmol), *fac*-Ir(ppy)₃ (0.010 mmol), base (1.0 mmol), solvent (2.5 mL).^[b] **2a** (1.0 mmol).^[c] NMP (1.0 mL).^[d] In the dark.