Published in final edited form as:

Bone. 2019 May ; 122: 31–37. doi:10.1016/j.bone.2019.01.023.

“Effects of disease-afflicted and aging neurons on the musculoskeletal system”

Gregorio Valdez^{1,2}

¹Fralin Biomedical Research Institute, Virginia Tech Carilion, Roanoke, Virginia, USA

²Department of Biological Sciences, Virginia Tech, Blacksburg, Virginia, USA

Abstract

The musculoskeletal system includes skeletal muscles, bones and innervating axons from neurons in the central and peripheral nervous system. Together, they form the largest structure in the body. They also initiate and coordinate locomotion, provide structural stability, and contribute to metabolism and homeostasis. Because of these functions, much effort has been devoted to ascertaining the impact of acute and chronic stress, such as disease, injury and aging, on the musculoskeletal system. This review will examine the role of the nervous system in the deleterious changes that accrue in skeletal muscles and bones during the progression of neurologic diseases and with advancing age.

Keywords

Synapse; Neuromuscular junction; Motor Neurons; Sensory Neurons; Sympathetic Neurons; Aging; Diseases

1. Introduction:

The musculoskeletal system requires innervating axons to form, mature and function. It is therefore not surprising that skeletal muscles and bones undergo deleterious morphological and functional changes when innervating axons become dysfunctional or degenerate due to diseases, old age and injuries. Exemplifying this point, the function and size of skeletal muscle diminish in diseases that result in the degeneration of α -motor neuron axons, such as amyotrophic lateral sclerosis (ALS) and spinal muscular atrophy (SMA) [1,2]. Skeletal muscles also atrophy following injuries that either compromise the function or result in the destruction of α -motor neuron axons [3]. A similar relationship has been established between bones and innervating axons. Bones lose mass and become weak and brittle when sensory and sympathetic neurons dysfunction [4]. Data also continue to accrue indicating that age-related deleterious changes in innervating axons contribute to the reduction of

Gregorio Valdez, Fralin Biomedical Research Institute, Virginia Tech Carilion, Virginia Tech, 2 Riverside Circle, Roanoke, VA 24016, gvaldez1@vtc.vt.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

skeletal muscle and bone mass, called sarcopenia and osteoporosis respectively, as well as loss of function of the musculoskeletal system in old age [5,6]. This review will summarize published studies on the effect of diseases and aging on skeletal muscles and bones and their innervating axons. It will also propose future lines of inquiries to further understand the role of the nervous system in the myriad of pathogenic features associated with aged and disease-afflicted skeletal muscles and bones.

2. Impact of disease-affected and aging neurons on skeletal muscles:

A skeletal muscle is a complex organ that requires the exquisite collaboration of various resident cells and innervating axons to form, function and repair damages to remain viable. In this section, we will provide a general summary of the type, location and function of innervating axons in skeletal muscles. We will also highlight the importance of these innervating axons in the maintenance and repair of skeletal muscles. Additionally, we will review the contribution of innervating axons to pathophysiological changes that accrue in skeletal muscles during the progression of diseases and normal aging.

2.1. Innervation of skeletal muscles:

Skeletal muscles are innervated by three neuronal types (Table 1) with distinctive roles: (1) motor neurons, which relay information from the central nervous system (CNS) to muscle to induce muscle contraction and are subdivided into α and γ subtypes; (2) sensory neurons, which relay proprioceptive (i.e. the rate and extent of contraction of a given muscle), temperature and pain signals (nociceptors), from muscles to the CNS; and (3) sympathetic neurons, which modulate a variety of adaptive cellular processes, particularly those involved in flight or fight responses. Among these neurons, α -motor neurons are essential for skeletal muscle to form, mature, contract and survive. These neurons exert their influence on skeletal muscles primarily at the neuromuscular junction (NMJ), a synapse formed between α -motor neuron axon nerve endings (also referred as the presynaptic region) and extrafusal muscle fibers [7]. At NMJs, α -motor neurons release acetylcholine to induce muscle contraction. It also releases molecules, such as neuronal isoforms of agrin, called z-agrin, to promote the formation and stability of the postsynaptic region (the region within an extrafusal muscle fiber specialized for decoding cholinergic transmission) [8,9]. Through these actions, α -motor neuron neurons play central roles in the function and viability of skeletal muscles, and thus in the initiation of all voluntary movements.

The coordination and precision of muscle movement requires the action of γ -motor neurons and type Ia/II proprioceptive sensory neurons. These neurons converge on intrafusal muscle fibers, to form a structure called the muscle spindle [10]. γ -motor neurons innervate the polar region of intrafusal muscle fibers where they release ACh to cause contraction of this muscle fiber type. The nerve endings of Ia/II proprioceptive sensory neurons, on the other hand, terminate throughout the equatorial region of intrafusal muscle fibers and function to relay information to the CNS, and directly to α -motor neurons, about the tone and contraction rate of skeletal muscles. A third proprioceptive sensory neuron subtype, type Ib, innervates Golgi tendon organs and also relays information regarding the contractile status of skeletal muscles to various components of the CNS, including α -motor neurons [11].

Through these actions, either through direct or indirect connections with α -motor neurons in the spinal cord, these neurons influence the function and health of skeletal muscles.

Sensory neurons that sense pain and temperature, commonly referred to as nociceptors, have been found in skeletal muscles [12–18]. The axons of these nociceptors are either thinly myelinated (A δ -fibers) or unmyelinated (C fibers), and are positive for the neuropeptide substance P (SP) and calcitonin gene-related peptide (CGRP) [19]. Unlike proprioceptive sensory neurons, nociceptors do not contact muscle fibers nor any other cell within skeletal muscles. These neurons depolarize (are activated) only when levels of potassium and ATP abruptly and significantly increase [20]. Hence, these nociceptors relay pain information to the CNS following the tearing of many muscle fibers at once, often caused by injuries. This explains the lack of pain sensation during the progressive and asynchronous atrophy of muscles, in which only a few muscle fibers die at any given time and therefore do not significantly alter potassium and ATP levels in the interstitial space, with advancing age and progression of diseases such as ALS, SMA and the spectrum of muscular dystrophies.

In addition to motor and sensory neurons, skeletal muscles are innervated by sympathetic postganglionic nerve endings positive for the neuropeptide Y (NPY) and tyrosine hydroxylase (TH) [18,21–24]. Using light microscopy, these nerve endings have been found in various locations within skeletal muscles of rabbits, rodents and humans [13,22,24,25]. They are present in muscle spindles and adjacent to extrafusal muscle fibers in addition to surrounding blood vessels [24,26]. More recently, sympathetic nerve endings were found adjacent to and directly abutting NMJs in the soleus and extensor digitorum longus muscles in developing and adult mice [22]. Their presence at NMJs and in muscle spindles suggests that they play important functions beyond modulating blood vessels. This has been supported by recent findings showing that sympathetic nerve endings activate the α -1a and β -2 adrenergic receptors (ADRA1A and ADRB2, respectively) in skeletal muscles, leading to a marked increase in the production of 3',5'-cyclic adenosine monophosphate (cAMP) [27] and the peroxisome proliferator-activated receptor gamma coactivator 1- α (PGC-1 α) [22,23,28]. These two signaling molecules have been shown to mediate a variety of cellular processes that influence mitochondrial function, satellite cell proliferation and NMJ health in skeletal muscles [29–31]. Through these and potentially other molecules, sympathetic nerve endings alter the release of ACh at NMJs, regulate the secretion and reuptake of calcium from the sarcoplasmic reticulum, and modulate glucose uptake and metabolic activity. These molecular changes are necessary for skeletal muscles to adapt to changing physiological and functional demands not only for flight-or-flight responses, but also associated with development, exercise, diseases, injuries and aging.

2.2. Impact of disease-afflicted neurons on skeletal muscles:

Skeletal muscles must be innervated by α -motor neurons in order to function and survive. Hence, it is not surprising that diseases that affect α -motor neurons and their axons invariably result in the malfunction and atrophy of skeletal muscles. This is the case in ALS [32–36], SMA and myasthenia gravis (MG) [1,37–41], which directly target α -motor neurons and particularly their nerve endings in skeletal muscles. In ALS and SMA, α -motor neurons and their axons completely degenerate, resulting in the loss of muscle function and

mass. MG is a disease that causes muscle weakness and fatigue resulting from mutations or autoimmune-related loss of function of various genes critical for the function of the presynaptic or postsynaptic region of the NMJ. Among presynaptic genes are the P/Q-type voltage-gated calcium channel (VGCC) and choline acetyl transferase [42]. The nerve endings of α -motor neurons are also affected in muscular dystrophy [43,44], diabetes, and vascular exclusion [45–48]. While the effect of these diseases on α -motor neuron nerve endings is likely indirect, their dysfunction may nevertheless further compromise the function and viability of skeletal muscles.

There is evidence that γ -motor neurons and proprioceptive sensory neurons undergo deleterious changes in ALS and SMA. In a mouse model for ALS, γ -motor neurons were found to increase the contraction rate of intrafusal muscles fibers, and thus augment the activity of Ia proprioceptive sensory neurons [49]. This heightened firing frequency of Ia proprioceptive sensory neurons has been suggested to contribute to calcium-induced cytotoxicity of α -motor neurons by augmenting levels of glutamate at synaptic sites in the spinal cord [50]. Additionally, recent findings indicate that proprioceptive sensory neurons may be directly affected and contribute to α -motor neuron degeneration in ALS and SMA. In a mouse model for SMA, the loss of proprioceptive sensory synaptic inputs correlates temporally and spatially with degeneration of α -motor neurons [51]. More recently, our lab demonstrated that Ia and II proprioceptive sensory nerve endings in the spinal cord and in skeletal muscles also degenerate in two mouse models for ALS, either harboring SOD1^{G93A} or TDP43^{A315T} [52]. Importantly, we showed that these neurons are intrinsically compromised in ALS [53], indicating that they may contribute to disease pathology by altering the function of α -motor neurons. Together, these published findings support roles for γ -motor and proprioceptive sensory neurons in ALS and SMA, two diseases that cause atrophy of skeletal muscles. In contrast to proprioceptive sensory neurons, the impact of diseases on nociceptive sensory neurons, and thus their contribution to skeletal muscle dysfunction and atrophy, remains unknown.

Sympathetic neurons have also been implicated in the function and health of skeletal muscles in various diseases. In this regard, drugs that augment the actions of the sympathetic system, such as ephedrine or salbutamol, provide therapeutic benefits presumably by restoring the normal function of NMJs [42]. Supporting this possibility, the inverse experiment, which relied on other chemicals to induce degeneration of sympathetic nerves, was shown to reduce the size of NMJs and density of nAChRs [23]. These changes, however, were reversed following the systemic administration of the ADRB2 agonist clenbuterol [54]. Sympathetic neurons have also been implicated in the progression of SMA and ALS, two diseases that invariably result in degeneration of the NMJ and atrophy of skeletal muscles [55–57]. Together, these published findings demonstrate important roles for sympathetic neurons in skeletal muscle pathogenesis. They have also revealed that a close anatomical and functional relationship exists between sympathetic and motor nerve endings at NMJs [54].

2.3. Effect of aging neurons on skeletal muscles:

The loss of skeletal muscle function and mass is a hallmark of old age in humans and animals. In humans, there is a significant and constant decline in muscle mass and strength [58], called sarcopenia, beginning in the fifth decade of life. Similar deleterious changes have been shown to occur on skeletal muscles of aging rodents. Over the last few decades data has accrued supporting an important role for the nervous system in the erosion of skeletal muscle function and mass with advancing age [59,60]. Not surprisingly, the majority of studies have examined the role of α -motor neurons. To date, we know that aging results in significant molecular, morphological and functional changes in α -motor neurons [61–65]. Aging also impairs the propagation of action potentials down α -motor neuron axons [66–68], alters the distribution of synaptic vesicles [69] and reduces the density of active zones at the presynaptic region of α -motor neuron nerve endings [70,71]. These destructive cellular changes with advancing age ultimately cause the nerve endings of α -motor neurons to retract from NMJs following a similar pattern as observed in ALS [32,33,72–78]. Because these deleterious changes occur progressively, it has been hypothesized that degeneration of α -motor neuron nerve endings contribute to aging of skeletal muscles, including sarcopenia. In support of this possibility, the two lifestyles best known to slow aging of skeletal muscles, a calorically restricted diet and exercise [33], attenuate the deleterious effects of aging on innervating α -motor axons. The caloric-restriction mimetic, resveratrol, was also found to delay aging of skeletal muscles partly by preventing retraction of α -motor axons from NMJs [79].

Because coordinated movements, such as gait and balance, become increasingly more difficult to properly carry out with aging [80,81], our lab recently examined the impact of aging on Ia/II proprioceptive sensory neurons in mice [82]. We found that the majority of Ia proprioceptive afferents fail to maintain their annulospirals structures and type II proprioceptive afferents degenerate in muscle spindles of the extensor digitorum longus (EDL) and soleus muscles of 11-and 15-month old mice compared to 2-month-old mice. Accompanying these changes, we found fewer proprioceptive sensory neurons populating dorsal root ganglia in old mice. Despite these findings, it remains unknown if changes in proprioceptive sensory neuron nerve endings contribute to the loss of skeletal muscle function and mass with aging. If they do, they likely act by altering the function of α -motor neurons. There is also a paucity of knowledge regarding the effect of aging on γ -motor, nociceptive, Ib proprioceptive, and sympathetic neurons innervating skeletal muscles.

3. Impact of disease-affected and aging neurons on bones:

While bone is a hard tissue, a characteristic necessary to act as a scaffold, it is also highly adaptable to changes. Exemplifying this, bones undergo significant remodeling as they grow during development and repair damages following injuries. These features of bones, however, are compromised in diseases and in old age. Here, we will review the potential contribution of innervating axons on the deleterious morphological changes associated with disease-afflicted and aged bones.

3.1. Innervation of bones:

Sensory and sympathetic neurons innervate various regions of bones [83,84] (Table 2). Their nerve endings are found on the periosteum, a fibrous membrane covering the entire surface of bones that contains blood vessels and lymphatic vessels. Nerve endings are also present in two distinct anatomical regions of bones, the diaphysis and epiphysis. The diaphysis is the tubular region that runs from the distal to the proximal end of bones. This area has an inner region, called the medullary canal where yellow bone marrow resides. The medullary canal is covered by the endosteum, a delicate fibrous membrane. The epiphysis is the wider and polar region of a bone. It is made up of spongy bone filled with red marrow. Axons enter deeper regions of bones together with blood vessels and through the foramen, located near the equatorial region of bones.

Sensory axons are most prevalent in the periosteum, and surround this area in a fishnet-like pattern presumably to better detect mechanical distortion of the underlying cortical bone that can result from injuries, diseases and even aging [85]. Sensory nerves are also present in cortical bone as well as bone marrow but at a lower density compared to the periosteum. To date, the preponderance of evidence indicates that bones are innervated largely by nociceptive sensory neurons, which is not surprising given the sharp and intense pain sensation following an injury to this tissue [86]. As expected, these nociceptive axons are small in diameter, unmyelinated or thinly myelinated, and exhibit electrical conduction velocities of less than 2 m/s or between 2 to 30 m/s. In contrast to skeletal muscles, nociceptors innervating bones are heterogeneous, and can be subdivided based on expression of various combinations of SP, CGRP, the tyrosine receptor kinase A (TrkA), and isolectin B4 (IB4). Suggesting that sensory innervation affects bone development in addition to relaying painful sensation, there is a strong correlation between sprouting of developing sensory nerves and formation of new bone [87]. Moreover, the destruction of sensory nerves slows the formation of new bone and increases fragility of bones [88]. More recently, Chen H et al [89] established that sensory neurons play an active and direct role in the maintenance and repair of bones. The authors found that osteoblasts release prostaglandin E2 (PGE2) to activate the PGE2 receptor 4 (EP4) located in sensory nerves. Importantly, they demonstrated that inhibiting this communication, between osteoblast and sensory nerves, has deleterious effects on the integrity and mass of bones.

Sympathetic postganglionic neurons positive for TH, NPY and dopamine-beta-hydroxylase also innervate bones [21,88,90,91]. These nerve endings are found sparsely on the periosteum compared to the inner regions of the bone, the epiphysis and diaphysis. In addition to regulating vasomotor function, sympathetic nerves play important roles in developing and adult bones. They directly affect bones by releasing norepinephrine to activate ADRB2 in osteoblasts and osteoclasts. Highlighting the functional role of this sympathetic signaling axis in bone, deletion of ADRB2 increases the formation of cancellous bone [92], the spongy bone located in the epiphysis region of long bones. Similarly, blocking ADRB2 with propranolol, a non-selective antagonist, increases cancellous bone [93,94]. Conversely, constitutively stimulating the ADRB2 using agonists, such as isoproterenol, clenbuterol, and salbutamol, results in the loss of bone mass in mice and rats [93]. Furthermore, Chen H et al [89] recently demonstrated that sympathetic

hyperactivation promotes bone loss. There is also evidence indicating that sympathetic nerves regulate bone through the beta-1 adrenergic receptor (ADRB1). Specifically, Khosla S et al [95] found a strong correlation between selective blockers of the ADRB1 with better structural integrity and higher metabolic activity in bones of postmenopausal women. The role of sympathetic nerves in bones, however, varies depending on their location. For example, stimulation of the sympathetic nervous system in bone marrow promotes the differentiation of resident stem cells and the subsequent egress of hematopoietic stem and progenitor cells [92] in contrast to inhibiting the actions of osteoblasts in other bone regions. Nevertheless, these and other published evidence have established a central role for sympathetic nerve endings in the formation, maintenance and repair of bones.

3.2. Impact of disease-afflicted neurons on bones:

Sensory and sympathetic neurons are affected in various diseases known to compromise the structural integrity and to reduce the mass of bones. In the reflex sympathetic dystrophy syndrome (RSDS; also called complex regional pain syndrome), damaged sympathetic nerves extensively sprout within bones leading to osteoporosis and osteopenia [96]. In sensory and autonomic neuropathies (HSAN), a broad spectrum disease with different etiologies, the degeneration of sensory and sympathetic nerves cause deleterious changes on bones residing in the distal portions of the lower extremities [97]. Sensory and sympathetic neurons are also affected in a number of genetic bone disorders such as osteogenesis imperfecta, Paget's disease and juvenile arthritis [83]. In these genetic bone diseases, their nerve endings also extensively sprout, partly explaining the severe pain and loss of bone mass resulting from heightened sympathetic innervation that accompanies such disorders. In addition, the high incidence of osteoporosis and bone fractures in neurological disorders with a CNS etiology, including Alzheimer's disease, Parkinson's disease, ALS and SMA, suggest roles for sympathetic and sensory nerve endings in the deterioration of bones [98]. However, sensory and sympathetic nerve endings innervating bones have yet to be examined prior to and during the progression of such neurological diseases.

3.3. Effect of aging neurons on bones:

There is a paucity of knowledge regarding the effect of aging on sensory and sympathetic nerve endings innervating bones. In one of the few studies published to date, Jimenez-Andrade JM et al [99] visualized sensory axons in the femur periosteum of young, middle-aged and old rats using antibodies against CGRP and neurofilament (NF). This study found no difference in the density of sensory axons across ages. More recently, Chartier SR et al [83] examined sensory axons in the femur of young and aged mice. In the periosteum, bone marrow, and cortical bone, the density of sensory nerve endings was unchanged. These findings partly corroborate observations in humans indicating that the number of sensory nerve endings increases in aged and degenerating vertebrae [100]. Together, these published observations indicate that sensory nerve endings remain unchanged or potentially sprout to occupy more territory even though bones clearly undergo deleterious structural changes in old age. Interestingly, sensory neurons also extensively sprout in fractured adult bones, a morphological change associated with increased pain sensation. Thus, the exuberant sprouting of sensory nerve endings may underlie the increased sensitivity to bone pain in the elderly population [20,101]. While the aforementioned studies provide insights into the

effect of aging on sensory neurons innervating bones, there is still much to learn about the effect of aging on these nociceptors. For example, the impact of aging on the functional and molecular characteristics of sensory nerve endings remains unknown. This information is essential to further understand the role of these neurons in aging bones.

Chartier et al [83] also examined sympathetic nerve endings in different regions of the femur of aging mice using an antibody against TH. Similar to sensory nerve endings, the density of sympathetic nerve endings was found unchanged in the periosteum and bone marrow of aged femurs. This was not the case in cortical bone where the density of sympathetic nerve endings was found to be decreased by over 75% in the femur of aged mice. While the contribution of sympathetic nerves to age-related changes in bones remains unknown, severing or blocking the function of these nerve endings promotes bone growth in adulthood. Thus, the loss of sympathetic nerve endings in cortical bone, which constitutes over 80% of long bones such as the femur, may contribute to the progressive enlargement of bones with advancing age. In support of this possibility, deletion of adenylyl cyclase 5, a downstream target of beta-2AR, was shown to increase bone mass with advancing age [102].

4. Concluding remarks:

The nervous system clearly plays an important and at times central role in the pathogenic changes that accumulate in bones and skeletal muscles due to diseases and with advancing age. Because of this, it is critical to answer a number of outstanding questions to target neurons that innervate bones and skeletal muscles. These include: 1) What are the initial molecular, subcellular and functional changes that precipitate dysfunction and degeneration of motor, sensory and sympathetic neuron nerve endings in bones and skeletal muscles?; 2) Does aging of innervating neurons precede, cause or contribute to pathophysiological changes in skeletal muscles and bones?; 3) Does aging simultaneously affect innervating neurons, skeletal muscles and bones?; 4) Do agents that preserves the function of sympathetic and sensory neurons delay or prevent musculoskeletal aging?; and 5) What underlies the disparate effect of altering the function of sympathetic neurons on bones compared to skeletal muscles? Additionally, it will be important to understand if and how different neuronal populations innervating skeletal muscles and bones cross-talk in the periphery and in the CNS. This question is particularly relevant given the evidence that sympathetic neurons potentially interact with sensory neurons in bones [89,103] and motor neurons and proprioceptive sensory neurons in skeletal muscles [22– 25]. The fact that connected bones and skeletal muscles adapt in concert to different functional demands and pathological insults raises the possibility that neuronal circuits in the CNS coordinate changes between these two components of the musculoskeletal system. Answering these questions is essential to understand the contribution of the nervous system to pathophysiological changes in the musculoskeletal system. This knowledge may also reveal broader relationships between different neurons that innervate bones and skeletal muscles that were not appreciated in the past, and pave the path to better mitigate disease and age-related deleterious changes in the musculoskeletal system.

Acknowledgement

The author thanks Thomas Taetzsch for providing critical comments and editing the manuscript.

Funding

This work was supported by the National Institute on Aging (R01AG055545) and the National Institute of Neurological Disorders and Stroke (R21NS106313).

References:

- [1]. Arnold WD, Kassar D, Kissel JT, Spinal muscular atrophy: diagnosis and management in a new therapeutic era., *Muscle Nerve* 51 (2015) 157–67. doi:10.1002/mus.24497. [PubMed: 25346245]
- [2]. Kiernan MC, Vucic S, Cheah BC, Turner MR, Eisen A, Hardiman O, Burrell JR, Zoing MC, Amyotrophic lateral sclerosis., *Lancet* (London, England) 377 (2011) 942–55. doi:10.1016/S0140-6736(10)61156-7.
- [3]. Carlson BM, The Biology of Long-Term Denervated Skeletal Muscle., *Eur. J. Transl. Myol* 24 (2014) 3293. doi:10.4081/ejtm.2014.3293. [PubMed: 26913125]
- [4]. Grässel S, The role of peripheral nerve fibers and their neurotransmitters in cartilage and bone physiology and pathophysiology, 2014. doi:10.1186/s13075-014-0485-1.
- [5]. Taetzsch T, Valdez G, NMJ maintenance and repair in aging, *Curr. Opin. Physiol* 4 (2018) 57–64. doi:10.1016/j.cophys.2018.05.007. [PubMed: 30560223]
- [6]. Roberts S, Colombier P, Sowman A, Mennan C, Rölfing JHD, Guicheux J, Edwards JR, Ageing in the musculoskeletal system., *Acta Orthop* 87 (2016) 15–25. doi:10.1080/17453674.2016.1244750. [PubMed: 27748151]
- [7]. Sanes JR, Lichtman JW, Development of the vertebrate neuromuscular junction., *Annu. Rev. Neurosci* 22 (1999) 389–442. doi:10.1146/annurev.neuro.22.1.389. [PubMed: 10202544]
- [8]. Gautam M, Noakes PG, Moscoso L, Rupp F, Scheller RH, Merlie JP, Sanes JR, Defective neuromuscular synaptogenesis in agrin-deficient mutant mice., *Cell* 85 (1996) 525–35. [PubMed: 8653788]
- [9]. Samuel MA, Valdez G, Tapia JC, Lichtman JW, Sanes JR, Agrin and synaptic laminin are required to maintain adult neuromuscular junctions., *PLoS One* 7 (2012) e46663. doi:10.1371/journal.pone.0046663. [PubMed: 23056392]
- [10]. Chen H-H, Hippenmeyer S, Arber S, Frank E, Development of the monosynaptic stretch reflex circuit., *Curr. Opin. Neurobiol* 13 (2003) 96–102. [PubMed: 12593987]
- [11]. Proske U, The golgi tendon organ, *Trends Neurosci* 2 (1979) 7–8. doi:10.1016/0166-2236(79)90004-3.
- [12]. Jankowski MP, Rau KK, Ekmann KM, Anderson CE, Koerber HR, Comprehensive phenotyping of group III and IV muscle afferents in mouse., *J. Neurophysiol* 109 (2013) 2374–81. doi:10.1152/jn.01067.2012. [PubMed: 23427306]
- [13]. Ljung BO, Forsgren S, Fridén J, Sympathetic and sensory innervations are heterogeneously distributed in relation to the blood vessels at the extensor carpi radialis brevis muscle origin of man., *Cells. Tissues. Organs* 165 (1999) 45–54. doi:10.1159/000016673. [PubMed: 10460973]
- [14]. Mense S, Muscle pain: mechanisms and clinical significance., *Dtsch. Arztebl. Int* 105 (2008) 214–9. doi:10.3238/artzebl.2008.0214. [PubMed: 19629211]
- [15]. Leighton B, Foot EA, The role of the sensory peptide calcitonin-gene-related peptide(s) in skeletal muscle carbohydrate metabolism: effects of capsaicin and resiniferatoxin., *Biochem. J* 307 (Pt 3) (1995) 707–12. [PubMed: 7741700]
- [16]. Mense S, Chapter 5. Group III and IV receptors in skeletal muscle: are they specific or polymodal?, *Prog. Brain Res* 113 (1996) 83–100. doi:10.1016/S0079-6123(08)61082-1. [PubMed: 9009729]
- [17]. Barry CM, Kestell G, Gillan M, Haberberger RV, Gibbins IL, Sensory nerve fibers containing calcitonin gene-related peptide in gastrocnemius, latissimus dorsi and erector spinae muscles and

- thoracolumbar fascia in mice, *Neuroscience* 291 (2015) 106–117. doi:10.1016/j.neuroscience.2015.01.062. [PubMed: 25681518]
- [18]. Peyronnard JM, Charron LF, Lavoie J, Messier JP, Motor, sympathetic and sensory innervation of rat skeletal muscles., *Brain Res* 373 (1986) 288–302. <http://www.ncbi.nlm.nih.gov/pubmed/3719313> (accessed November 26, 2018). [PubMed: 3719313]
- [19]. Woolf CJ, Ma Q, Nociceptors—Noxious Stimulus Detectors, *Neuron* 55 (2007) 353–364. doi:10.1016/J.NEURON.2007.07.016. [PubMed: 17678850]
- [20]. Mantyh PW, The neurobiology of skeletal pain, (n.d.). doi:10.1111/ejn.12462.
- [21]. Houweling P, Kulkarni RN, Baldock PA, Neuronal control of bone and muscle, *Bone* 80 (2015) 95–100. doi:10.1016/J.BONE.2015.05.006. [PubMed: 26453499]
- [22]. Straka T, Vita V, Prokshi K, Hörner S, Khan M, Pirazzini M, Williams M, Hafner M, Zaglia T, Rudolf R, Postnatal Development and Distribution of Sympathetic Innervation in Mouse Skeletal Muscle, *Int. J. Mol. Sci* 19 (2018) 1935. doi:10.3390/ijms19071935.
- [23]. Rodrigues ACZ, Messi ML, Wang Z-M, Abba MC, Pereyra A, Birbrair A, Zhang T, O'Meara M, Kwan P, Lopez EIS, Willis MS, Mintz A, Files DC, Furdul C, Oppenheim RW, Delbono O, The sympathetic nervous system regulates skeletal muscle motor innervation and acetylcholine receptor stability, *Acta Physiol* (2018) e13195. doi:10.1111/apha.13195.
- [24]. Radovanovic D, Peikert K, Lindström M, Domellöf FP, Sympathetic innervation of human muscle spindles., *J. Anat* 226 (2015) 542–8. doi:10.1111/joa.12309. [PubMed: 25994126]
- [25]. Bombardi C, Grandis A, Chiocchetti R, Bortolami R, Johansson H, Lucchi ML, Immunohistochemical localization of alpha1a-adrenoreceptors in muscle spindles of rabbit masseter muscle, *Tissue Cell* 38 (2006) 121–125. doi:10.1016/j.tice.2005.12.003. [PubMed: 16510160]
- [26]. Barker D, Saito M, Autonomic Innervation of Receptors and Muscle Fibres in Cat Skeletal Muscle, *Proc. R. Soc. B Biol. Sci* 212 (1981) 317–332. doi:10.1098/rspb.1981.0042. [PubMed: 6115396]
- [27]. Silveira WA, Gonçalves DA, Graça FA, Andrade-Lopes AL, Bergantin LB, Zanon NM, Godinho RO, Kettelhut IC, Navegantes LCC, Activating cAMP/PKA signaling in skeletal muscle suppresses the ubiquitin-proteasome-dependent proteolysis: implications for sympathetic regulation, *J. Appl. Physiol* 117 (2014) 11–19. doi:10.1152/jappphysiol.01055.2013. [PubMed: 24833777]
- [28]. Snyder-Warwick AK, Satoh A, Santosa KB, Imai S-I, Jablonka-Shariff A, Hypothalamic Sirt1 protects terminal Schwann cells and neuromuscular junctions from age-related morphological changes., *Aging Cell* 17 (2018) e12776. doi:10.1111/accel.12776.
- [29]. Chakkalakal JV, Jones KM, Basson MA, Brack AS, The aged niche disrupts muscle stem cell quiescence., *Nature* 490 (2012) 355–60. doi:10.1038/nature11438. [PubMed: 23023126]
- [30]. Dupuis L, Gonzalez de Aguilar J-L, Echaniz-Laguna A, Eschbach J, Rene F, Oudart H, Halter B, Huze C, Schaeffer L, Bouillaud F, Loeffler J-P, Muscle mitochondrial uncoupling dismantles neuromuscular junction and triggers distal degeneration of motor neurons., *PLoS One* 4 (2009) e5390. doi:10.1371/journal.pone.0005390. [PubMed: 19404401]
- [31]. Gonzalez-Freire M, de Cabo R, Studenski SA, Ferrucci L, The Neuromuscular Junction: Aging at the Crossroad between Nerves and Muscle., *Front. Aging Neurosci* 6 (2014) 208. doi:10.3389/fnagi.2014.00208. [PubMed: 25157231]
- [32]. Moloney EB, de Winter F, Verhaagen J, ALS as a distal axonopathy: molecular mechanisms affecting neuromuscular junction stability in the presymptomatic stages of the disease., *Front. Neurosci* 8 (2014) 252. doi:10.3389/fnins.2014.00252. [PubMed: 25177267]
- [33]. Valdez G, Tapia JC, Kang H, Clemenson GD, Jr., Gage FH, Lichtman JW, Sanes JR, Attenuation of age-related changes in mouse neuromuscular synapses by caloric restriction and exercise, *Proc Natl Acad Sci U S A* 107 (2010) 14863–14868. doi:1002220107 [pii]10.1073/pnas.1002220107. [PubMed: 20679195]
- [34]. Etang AFDL, Maharjan N, Braña MC, Rueggsegger C, Rehmann R, Goswami A, Roos A, Troost D, Schneider BL, Weis J, Saxena S, Marinesco-Sjögren syndrome protein SIL1 regulates motor neuron subtype-selective ER stress in ALS, *18* (2015). doi:10.1038/nn.3903.

- [35]. Valsecchi V, Boido M, De Amicis E, Piras A, Vercelli A, Expression of Muscle-Specific MiRNA 206 in the Progression of Disease in a Murine SMA Model., PLoS One 10 (2015) e0128560. doi: 10.1371/journal.pone.0128560. [PubMed: 26030275]
- [36]. Bayat V, Jaiswal M, Bellen HJ, The BMP signaling pathway at the Drosophila neuromuscular junction and its links to neurodegenerative diseases., Curr. Opin. Neurobiol 21 (2011) 182–8. doi: 10.1016/j.conb.2010.08.014. [PubMed: 20832291]
- [37]. Titulaer MJ, Lang B, Verschuuren JJ, Lambert-Eaton myasthenic syndrome: from clinical characteristics to therapeutic strategies., Lancet. Neurol 10 (2011) 1098–107. doi:10.1016/S1474-4422(11)70245-9. [PubMed: 22094130]
- [38]. Drummond PD, Sensory–autonomic interactions in health and disease, in: Handb. Clin. Neurol, 2013: pp. 309–319. doi:10.1016/B978-0-444-53491-0.00024-9.
- [39]. Poliachik SL, Bain SD, Threet D, Huber P, Gross TS, Transient muscle paralysis disrupts bone homeostasis by rapid degradation of bone morphology, Bone 46 (2010) 18–23. doi:10.1016/J.BONE.2009.10.025. [PubMed: 19857614]
- [40]. Vincent A, Lang B, Kleopa KA, Autoimmune Channelopathies and Related Neurological Disorders, Neuron 52 (2006) 123–138. doi:10.1016/J.NEURON.2006.09.024. [PubMed: 17015231]
- [41]. Boido M, Vercelli A, Neuromuscular Junctions as Key Contributors and Therapeutic Targets in Spinal Muscular Atrophy., Front. Neuroanat 10 (2016) 6. doi:10.3389/fnana.2016.00006. [PubMed: 26869891]
- [42]. Cruz PMR, Palace J, Beeson D, Congenital myasthenic syndromes and the neuromuscular junction, Curr. Opin. Neurol 27 (2014) 566–575. doi:10.1097/WCO.0000000000000134. [PubMed: 25159927]
- [43]. Li L, Xiong W-C, Mei L, Neuromuscular Junction Formation, Aging, and Disorders, Annu. Rev. Physiol 80 (2018) 159–188. doi:10.1146/annurev-physiol-022516-034255. [PubMed: 29195055]
- [44]. Park KHJ, Mechanisms of Muscle Denervation in Aging: Insights from a Mouse Model of Amyotrophic Lateral Sclerosis., Aging Dis 6 (2015) 380–9. doi:10.14336/AD.2015.0506. [PubMed: 26425392]
- [45]. Tu H, Zhang D, Corrick RM, Muellemann RL, Wadman MC, Li Y-L, Morphological Regeneration and Functional Recovery of Neuromuscular Junctions after Tourniquet-Induced Injuries in Mouse Hindlimb, Front. Physiol 8 (2017) 207. doi:10.3389/fphys.2017.00207. [PubMed: 28428759]
- [46]. Jin KL, Mao XO, Greenberg DA, Shvartsman D, Cezar C, Lichtman JW, Vandenburgh HH, Mooney DJ, Vascular endothelial growth factor: Direct neuroprotective effect in in vitro ischemia, Proc. Natl. Acad. Sci 97 (2000) 10242–10247. doi:10.1073/pnas.97.18.10242. [PubMed: 10963684]
- [47]. Fahim MA, Hasan MY, Alshuaib WB, Early morphological remodeling of neuromuscular junction in a murine model of diabetes., J. Appl. Physiol 89 (2000) 2235–40. doi:10.1152/jappl.2000.89.6.2235. [PubMed: 11090573]
- [48]. Dalkin W, Taetzsch T, Valdez G, The Fibular Nerve Injury Method: A Reliable Assay to Identify and Test Factors That Repair Neuromuscular Junctions, J. Vis. Exp (2016). doi:10.3791/54186.
- [49]. Lalancette-Hebert M, Sharma A, Lyashchenko AK, Shneider NA, Γ motor neurons survive and exacerbate α motor neuron degeneration in ALS., Proc. Natl. Acad. Sci. U. S. A 113 (2016) E8316–E8325. doi:10.1073/pnas.1605210113. [PubMed: 27930290]
- [50]. Foran E, Trotti D, Glutamate transporters and the excitotoxic path to motor neuron degeneration in amyotrophic lateral sclerosis., Antioxid. Redox Signal 11 (2009) 1587–602. doi:10.1089/ars.2009.2444. [PubMed: 19413484]
- [51]. Mentis GZ, Blivis D, Liu W, Drobac E, Crowder ME, Kong L, Alvarez FJ, Sumner CJ, O'Donovan MJ, Early functional impairment of sensory-motor connectivity in a mouse model of spinal muscular atrophy., Neuron 69 (2011) 453–67. doi:10.1016/j.neuron.2010.12.032. [PubMed: 21315257]
- [52]. Vaughan SK, Kemp Z, Hatzipetros T, Vieira F, Valdez G, Degeneration of proprioceptive sensory nerve endings in mice harboring amyotrophic lateral sclerosis-causing mutations., J. Comp. Neurol 523 (2015) 2477–94. doi:10.1002/cne.23848. [PubMed: 26136049]

- [53]. Vaughan SK, Sutherland NM, Zhang S, Hatzipetros T, Vieira F, Valdez G, The ALS-inducing factors, TDP43A315T and SOD1G93A, directly affect and sensitize sensory neurons to stress., *Sci. Rep* 8 (2018) 16582. doi:10.1038/s41598-018-34510-8. [PubMed: 30410094]
- [54]. Khan MM, Lustrino D, Silveira WA, Wild F, Straka T, Issop Y, O'Connor E, Cox D, Reischl M, Marquardt T, Labeit D, Labeit S, Benoit E, Molgó J, Lochmüller H, Witzemann V, Kettelhut IC, Navegantes LCC, Pozzan T, Rudolf R, Sympathetic innervation controls homeostasis of neuromuscular junctions in health and disease., *Proc. Natl. Acad. Sci. U. S. A* 113 (2016) 746–50. doi:10.1073/pnas.1524272113. [PubMed: 26733679]
- [55]. Oey PL, Vos PE, Wieneke GH, Wokke JHJ, Blankestijn PJ, Karemaker JM, Subtle involvement of the sympathetic nervous system in amyotrophic lateral sclerosis., *Muscle Nerve* 25 (2002) 402–8. [PubMed: 11870718]
- [56]. Schara U, Barisic N, Deschauer M, Lindberg C, Straub V, Strigl-Pill N, Wendt M, Abicht A, Müller JS, Lochmüller H, Ephedrine therapy in eight patients with congenital myasthenic syndrome due to DOK7 mutations, *Neuromuscul. Disord* 19 (2009) 828–832. doi:10.1016/J.NMD.2009.09.008. [PubMed: 19837590]
- [57]. Pane M, Staccioli S, Messina S, D'Amico A, Pelliccioni M, Mazzone ES, Cuttini M, Alfieri P, Battini R, Main M, Muntoni F, Bertini E, Villanova M, Mercuri E, Daily salbutamol in young patients with SMA type II, *Neuromuscul. Disord* 18 (2008) 536–540. doi:10.1016/J.NMD.2008.05.004. [PubMed: 18579379]
- [58]. Hepple RT, Sarcopenia--a critical perspective., *Sci. Aging Knowledge Environ* 2003 (2003) pe31. doi:10.1126/sageke.2003.46.pe31. [PubMed: 14627844]
- [59]. Delbono O, Neural control of aging skeletal muscle., *Aging Cell* 2 (2003) 21–9. [PubMed: 12882331]
- [60]. Larsson L, Ansved T, Effects of ageing on the motor unit., *Prog. Neurobiol* 45 (1995) 397–458. [PubMed: 7617890]
- [61]. Das MM, Avalos P, Suezaki P, Godoy M, Garcia L, Chang CD, Vit J-P, Shelley B, Gowing G, Svendsen CN, Human neural progenitors differentiate into astrocytes and protect motor neurons in aging rats., *Exp. Neurol* 280 (2016) 41–9. [PubMed: 27032721]
- [62]. Rygiel KA, Grady JP, Turnbull DM, Respiratory chain deficiency in aged spinal motor neurons., *Neurobiol. Aging* 35 (2014) 2230–8. [PubMed: 24684792]
- [63]. Tomlinson BE, Irving D, The numbers of limb motor neurons in the human lumbosacral cord throughout life., *J. Neurol. Sci* 34 (1977) 213–9. [PubMed: 925710]
- [64]. Lexell J, Evidence for nervous system degeneration with advancing age., *J. Nutr* 127 (1997) 1011S–1013S. [PubMed: 9164286]
- [65]. Kawamura Y, O'Brien P, Okazaki H, Dyck PJ, Lumbar motoneurons of man II: the number and diameter distribution of large-and intermediate-diameter cytons in “motoneuron columns” of spinal cord of man., *J. Neuropathol. Exp. Neurol* 36 861–70. [PubMed: 894329]
- [66]. Fahim MA, Endurance exercise modulates neuromuscular junction of C57BL/6NNia aging mice, *J Appl Physiol* 83 (1997) 59–66. [PubMed: 9216945]
- [67]. Arasaki K, Iwamoto H, Tamaki M, Influence of aging and the motor unit type on the inverse relationship between the refractory period and the conduction velocity of the alpha motor nerve fibers., *Muscle Nerve* 18 (1995) 1019–20. doi:10.1002/mus.880180915. [PubMed: 7643864]
- [68]. Pousinha PA, Correia AM, Sebastião AM, Ribeiro JA, The giant miniature endplate potentials frequency is increased in aged rats., *Neurosci. Lett* 584 (2015) 224–9. doi:10.1016/j.neulet.2014.10.035. [PubMed: 25449868]
- [69]. Prakash YS, Sieck GC, Age-related remodeling of neuromuscular junctions on type-identified diaphragm fibers., *Muscle Nerve* 21 (1998) 887–95. [PubMed: 9626248]
- [70]. Nishimune H, Numata T, Chen J, Aoki Y, Wang Y, Starr MP, Mori Y, Stanford JA, Active zone protein Bassoon co-localizes with presynaptic calcium channel, modifies channel function, and recovers from aging related loss by exercise., *PLoS One* 7 (2012) e38029. doi:10.1371/journal.pone.0038029. [PubMed: 22701595]
- [71]. Lee KM, Chand KK, Hammond LA, Lavidis NA, Noakes PG, Functional decline at the aging neuromuscular junction is associated with altered laminin- α 4 expression, *Aging (Albany. NY)* 9 (2017) 880–899. doi:10.18632/aging.101198. [PubMed: 28301326]

- [72]. Valdez G, Tapia JC, Lichtman JW, Fox MA, Sanes JR, Shared resistance to aging and ALS in neuromuscular junctions of specific muscles., *PLoS One* 7 (2012) e34640. doi:10.1371/journal.pone.0034640. [PubMed: 22485182]
- [73]. Punga AR, Ruegg MA, Signaling and aging at the neuromuscular synapse: lessons learnt from neuromuscular diseases., *Curr. Opin. Pharmacol* 12 (2012) 340–6. doi:10.1016/j.coph.2012.02.002. [PubMed: 22365504]
- [74]. Rudolf R, Khan MM, Labeit S, Deschenes MR, Degeneration of neuromuscular junction in age and dystrophy., *Front. Aging Neurosci* 6 (2014) 99. doi:10.3389/fnagi.2014.00099. [PubMed: 24904412]
- [75]. Tintignac LA, Brenner H-R, Rüegg MA, Mechanisms Regulating Neuromuscular Junction Development and Function and Causes of Muscle Wasting., *Physiol. Rev* 95 (2015) 809–52. doi: 10.1152/physrev.00033.2014. [PubMed: 26109340]
- [76]. Vandervoort AA, Aging of the human neuromuscular system., *Muscle Nerve* 25 (2002) 17–25. [PubMed: 11754180]
- [77]. Fahim MA, Robbins N, Ultrastructural studies of young and old mouse neuromuscular junctions., *J. Neurocytol* 11 (1982) 641–56. [PubMed: 7131048]
- [78]. Rocha MC, Pousinha PA, Correia AM, Sebastião AM, Ribeiro JA, Early changes of neuromuscular transmission in the SOD1(G93A) mice model of ALS start long before motor symptoms onset., *PLoS One* 8 (2013) e73846. doi:10.1371/journal.pone.0073846. [PubMed: 24040091]
- [79]. Stockinger J, Maxwell N, Shapiro D, deCabo R, Valdez G, Caloric Restriction Mimetics Slow Aging of Neuromuscular Synapses and Muscle Fibers, *Journals Gerontol. Ser. A* (2017). doi: 10.1093/gerona/glx023.
- [80]. Fried LP, Ferrucci L, Darer J, Williamson JD, Anderson G, Untangling the concepts of disability, frailty, and comorbidity: implications for improved targeting and care., *J. Gerontol. A. Biol. Sci. Med. Sci* 59 (2004) 255–63. [PubMed: 15031310]
- [81]. Sorond FA, Cruz-Almeida Y, Clark DJ, Viswanathan A, Scherzer CR, De Jager P, Csizsar A, Laurienti PJ, Hausdorff JM, Chen WG, Ferrucci L, Rosano C, Studenski SA, Black SE, Lipsitz LA, Aging, the Central Nervous System, and Mobility in Older Adults: Neural Mechanisms of Mobility Impairment., *J. Gerontol. A. Biol. Sci. Med. Sci* 70 (2015) 1526–32. [PubMed: 26386013]
- [82]. Vaughan SK, Stanley OL, Valdez G, Impact of Aging on Proprioceptive Sensory Neurons and Intrafusal Muscle Fibers in Mice., *J. Gerontol. A. Biol. Sci. Med. Sci* (2016) glw175. doi: 10.1093/gerona/glw175.
- [83]. Chartier SR, Mitchell SAT, Majuta LA, Mantyh PW, The Changing Sensory and Sympathetic Innervation of the Young, Adult and Aging Mouse Femur, *Neuroscience* 387 (2018) 178–190. doi:10.1016/J.NEUROSCIENCE.2018.01.047. [PubMed: 29432884]
- [84]. Mach DB, Rogers SD, Sabino MC, Luger NM, Schwei MJ, Pomonis JD, Keyser CP, Clohisy DR, Adams DJ, O’Leary P, Mantyh PW, Origins of skeletal pain: sensory and sympathetic innervation of the mouse femur., *Neuroscience* 113 (2002) 155–66. [PubMed: 12123694]
- [85]. Matsuo K, Ji S, Miya A, Yoda M, Hamada Y, Tanaka T, Takao-Kawabata R, Kawaai K, Kuroda Y, Shibata S, Innervation of the tibial epiphysis through the intercondylar foramen, *Bone* 120 (2019) 297–304. doi:10.1016/j.bone.2018.11.007. [PubMed: 30439572]
- [86]. Castañeda-Corral G, Jimenez-Andrade JM, Bloom AP, Taylor RN, Mantyh WG, Kaczmarek MJ, Ghilardi JR, Mantyh PW, The majority of myelinated and unmyelinated sensory nerve fibers that innervate bone express the tropomyosin receptor kinase A, *Neuroscience* 178 (2011) 196–207. doi:10.1016/j.neuroscience.2011.01.039. [PubMed: 21277945]
- [87]. Gray C, Hukkanen M, Kontinen YT, Terenghi G, Arnett TR, Jones SJ, Burnstock G, Polak JM, Rapid neural growth: Calcitonin gene-related peptide and substance P-containing nerves attain exceptional growth rates in regenerating deer antler, *Neuroscience* 50 (1992) 953–963. doi: 10.1016/0306-4522(92)90218-Q. [PubMed: 1280352]
- [88]. Eleftheriou F, Campbell P, Ma Y, Control of Bone Remodeling by the Peripheral Sympathetic Nervous System, *Calcif. Tissue Int* 94 (2014) 140–151. doi:10.1007/s00223-013-9752-4. [PubMed: 23765388]

- [89]. Chen H, Hu B, Lv X, Zhu S, Zhen G, Wan M, Jain A, Gao B, Chai Y, Yang M, Wang X, Deng R, Wang L, Cao Y, Ni S, Liu S, Yuan W, Chen H, Dong X, Guan Y, Yang H, Cao X, Prostaglandin E2 mediates sensory nerve regulation of bone homeostasis, *Nat. Commun* 10 (2019) 181. doi: 10.1038/s41467-018-08097-7. [PubMed: 30643142]
- [90]. Togari A, Arai M, Kondo A, The role of the sympathetic nervous system in controlling bone metabolism, *Expert Opin. Ther. Targets* 9 (2005) 931–940. doi:10.1517/14728222.9.5.931. [PubMed: 16185149]
- [91]. Takeda S, Karsenty G, Molecular bases of the sympathetic regulation of bone mass, *Bone* 42 (2008) 837–840. doi:10.1016/J.BONE.2008.01.005. [PubMed: 18295563]
- [92]. Pierroz DD, Bonnet N, Bianchi EN, Bouxsein ML, Baldock PA, Rizzoli R, Ferrari SL, Deletion of β -adrenergic receptor 1, 2, or both leads to different bone phenotypes and response to mechanical stimulation, *J. Bone Miner. Res* 27 (2012) 1252–1262. doi:10.1002/jbmr.1594. [PubMed: 22407956]
- [93]. Bonnet N, Pierroz DD, Ferrari SL, Adrenergic control of bone remodeling and its implications for the treatment of osteoporosis., *J. Musculoskelet. Neuronal Interact* 8 (n.d.) 94–104. [PubMed: 18622078]
- [94]. Bonnet N, Laroche N, Vico L, Dolleans E, Benhamou CL, Courteix D, Dose Effects of Propranolol on Cancellous and Cortical Bone in Ovariectomized Adult Rats, *J. Pharmacol. Exp. Ther* 318 (2006) 1118–1127. doi:10.1124/JPET.106.105437. [PubMed: 16740619]
- [95]. Khosla S, Drake MT, Volkman TL, Thicke BS, Achenbach SJ, Atkinson EJ, Joyner MJ, Rosen CJ, Monroe DG, Farr JN, Sympathetic β 1-adrenergic signaling contributes to regulation of human bone metabolism., *J. Clin. Invest* 128 (2018) 4832–4842. doi:10.1172/JCI122151. [PubMed: 30153111]
- [96]. Goh EL, Chidambaram S, Ma D, Complex regional pain syndrome: a recent update., *Burn. Trauma* 5 (2017) 2. doi:10.1186/s41038-016-0066-4.
- [97]. Axelrod FB, Gold-von Simson G, Hereditary sensory and autonomic neuropathies: types II, III, and IV., *Orphanet J. Rare Dis* 2 (2007) 39. doi:10.1186/1750-1172-2-39. [PubMed: 17915006]
- [98]. Poncelet AN, Rose-Innes AP, Neurologic Disorders Associated with Bone and Joint Disease, *Amin. Neurol. Gen. Med* (2014) 433–457. doi:10.1016/B978-0-12-407710-2.00022-9.
- [99]. Jimenez-Andrade JM, Mantyh PW, Sensory and sympathetic nerve fibers undergo sprouting and neuroma formation in the painful arthritic joint of geriatric mice, *Arthritis Res. Ther* 14 (2012) R101. doi:10.1186/ar3826. [PubMed: 22548760]
- [100]. Brown MF, Hukkanen MV, McCarthy ID, Redfern DR, Batten JJ, Crock HV, Hughes SP, Polak JM, Sensory and sympathetic innervation of the vertebral endplate in patients with degenerative disc disease., *J. Bone Joint Surg. Br* 79 (1997) 147–53. [PubMed: 9020464]
- [101]. Yezierski RP, The effects of age on pain sensitivity: preclinical studies., *Pain Med* 13 Suppl 2 (2012) S27–36. doi:10.1111/j.1526-4637.2011.01311.x. [PubMed: 22497745]
- [102]. Yan L, Vatner DE, O'Connor JP, Ivessa A, Ge H, Chen W, Hirotani S, Ishikawa Y, Sadoshima J, Vatner SF, Type 5 adenylyl cyclase disruption increases longevity and protects against stress., *Cell* 130 (2007) 247–58. doi:10.1016/j.cell.2007.05.038. [PubMed: 17662940]
- [103]. Ryu V, Watts AG, Xue B, Bartness TJ, Bidirectional crosstalk between the sensory and sympathetic motor systems innervating brown and white adipose tissue in male Siberian hamsters, *Am. J. Physiol. Integr. Comp. Physiol* 312 (2017) R324–R337. doi:10.1152/ajpregu.00456.2015.

Highlights

- Role of α -motor nerve endings in skeletal muscles health
- Innervation of NMJs by sympathetic neurons
- Cross-talk between sympathetic and sensory neurons in bones

Table 1.

Neuronal types present in skeletal muscle.

Neuron	Connection with	Function
α -motor	Extrafusal muscle fiber	Muscle contraction
γ -motor	Intrafusal muscle fiber	Muscle contraction
Type Ia proprioceptive sensory	Muscle spindle	Proprioception
Type Ib proprioceptive sensory	Muscle spindle	Proprioception
Type II proprioceptive sensory	Golgi organ	Proprioception
Nociceptive sensory	None - Free nerve endings	Pain sensation
Sympathetic	Blood vessels, NMJs, muscle spindles	Flight or fight response

Table 2.

Neuronal types present in bone.

Neuron	Bone region innervated	Function
Nociceptive sensory	Periosteum, diaphysis, epiphysis, cortical bone, bone marrow	Detection of pain, bone remodeling
Sympathetic	Periosteum, diaphysis, epiphysis	Vasoregulation, bone remodeling