Published in final edited form as:

Prog Cardiovasc Dis. 2019 ; 62(2): 163–171. doi:10.1016/j.pcad.2018.08.006.

Meta-analysis of Exercise Training on Left Ventricular Ejection Fraction in Heart Failure with Reduced Ejection Fraction: A 10-year Update

Wesley J. Tucker^{a,1}, Rhys I. Beaudry^{a,1}, Yuanyuan Liang^b, Alexander M. Clark^c, Corey R. Tomczak^d, Michael D. Nelson^a, Oyvind Ellingsen^{e,f}, and Mark J. Haykowsky^{a,*}

^aCollege of Nursing and Health Innovation, University of Texas at Arlington, Arlington, TX, USA

^bUniversity of Maryland School of Medicine, Baltimore, MD, USA

^cFaculty of Nursing, University of Alberta, Edmonton, Alberta, Canada

^dCollege of Kinesiology, University of Saskatchewan, Saskatoon, Saskatchewan, Canada

^eK. G. Jebsen Center of Exercise in Medicine, Department of Circulation and Medical Imaging, Norwegian University of Science and Technology, Trondheim, Norway

^fDepartment of Cardiology, St. Olav's Hospital, Trondheim, Norway

Abstract

Background: The role of exercise training modality to attenuate left ventricular (LV) remodeling in heart failure patients with reduced ejection fraction (HFrEF) remains uncertain. The authors performed a systematic review and meta-analysis of published reports on exercise training (moderate-intensity continuous aerobic, high-intensity interval aerobic, and resistance exercise) and LV remodeling in clinically stable HFrEF patients.

Methods: We searched MEDLINE, Cochrane Central Registry of Controlled Trials, CINAHL, and PubMed (2007 to 2017) for randomized controlled trials of exercise training on resting LV ejection fraction (EF) and end-diastolic and end-systolic volumes in HFrEF patients.

Results: 18 trials reported LV ejection fraction (LVEF) data, while 8 and 7 trials reported LV end-diastolic and LV end-systolic volumes, respectively. Overall, moderate-intensity continuous training (MICT) significantly increased LVEF (weighted mean difference, WMD = 3.79%; 95% confidence interval, CI, 2.08 to 5.50%) with no change in LV volumes versus control. In trials ≥ 6 months duration, MICT significantly improved LVEF (WMD = 6.26%; 95% CI 4.39 to 8.13%) while shorter duration (<6 months) trials modestly increased LVEF (WMD = 2.33%; 95% CI 0.84 to 3.82%). High-intensity interval training (HIIT) significantly increased LVEF compared to control (WMD = 3.70%; 95% CI 1.63 to 5.77%) but was not different than MICT (WMD =

*Address reprint requests to: Mark J. Haykowsky, PhD, University of Texas at Arlington, College of Nursing and Health Innovation, 411 S Nedderman Drive, Arlington, TX 76010., mark.haykowsky@uta.edu (M.J. Haykowsky).

¹These authors contributed equally to this work.

Statement of conflict of interest

None of the authors have any conflicts of interests with regard to this publication.

3.17%; 95% CI -0.87 to 7.22%). Resistance training performed alone or combined with aerobic training (MICT or HIIT) did not significantly change LVEF.

Conclusions: In clinically stable HFrEF patients, MICT is an effective therapy to attenuate LV remodeling with the greatest benefits occurring with long-term (≥ 6 months) training. HIIT performed for 2 to 3 months is superior to control, but not MICT, for improvement of LVEF.

Keywords

Exercise therapy; Moderate-intensity continuous training; High-intensity interval training; Resistance training; Left ventricular ejection fraction

Heart failure (HF) is a major health care problem associated with a high individual and societal health care burden.¹ Approximately 50% of HF patients have reduced ejection fraction (HFrEF) with structural cardiac remodeling characterized by left ventricular (LV) dilation and reduced function.² Given that LV remodeling is associated with increased mortality, therapies that attenuate this process are important therapeutic targets.^{3,4}

Exercise training (ET) is an effective intervention to attenuate LV remodeling in clinically stable HFrEF patients.⁵ However, the magnitude of this change may depend on the type of exercise undertaken.⁶ In 2007, a meta-analysis by Haykowsky et al. found a significant increase in resting LV ejection fraction (LVEF) and reduction in LV volumes and improved peak oxygen uptake (peak VO_2) following aerobic moderate-intensity continuous training (MICT) but these changes were not replicated when aerobic and strength training were combined.⁶ Wisloff et al. extended these findings by demonstrating significantly greater anti-remodeling benefits with high-intensity interval training (HIIT) (e.g. brief, intermittent bursts of vigorous aerobic exercise interspersed with short periods of low-intensity active recovery⁷) compared to MICT in older (mean age: 76 years) post-infarction HFrEF patients.⁸ Yet in contrast, other investigators^{9–11} have not found additional anti-remodeling benefits with HIIT versus MICT in HFrEF patients (mean age: 60 years). Given this continued uncertainty and importance of understanding optimal ET regimen for the large and vulnerable population with HFrEF, we performed a 10-year update of our prior systematic review and meta-analysis of randomized controlled trials that examined the effect of ET on LV remodeling and peak oxygen consumption (peak VO_2) in clinically stable HFrEF patients.

Methods

Data sources

The authors searched (Jan 2007 to Jan 2017) MEDLINE, Cochrane Central Register of Controlled Trials, CINAHL, and PubMed using the following MESH terms and text words used in our original meta-analysis⁶: *heart failure, exercise, exercise therapy, exercise test, therapeutic exercise, cardiac rehabilitation, and kinesiotherapy*. We also hand-searched reference lists of all identified studies, and previous systematic reviews. We excluded non-English articles.

Study selection

After removal of all duplicate citations, two investigators (W.J.T. and R.I.B.) independently reviewed the titles and abstracts of all potentially eligible citations reporting the effect of ET on LVEF (primary outcome) and/or volumes in HFrEF patients. Both investigators obtained the full text of potentially relevant articles and independently reviewed those using pre-standardized data abstraction forms and eligibility criteria that were defined a priori. Studies were excluded that were: non-randomized, non-HFrEF, did not contain an ET intervention, did not report EF, LVEF $\geq 45\%$, did not have a usual care control group, had an ET intervention that also received a drug intervention, or had HF patients who were not clinically stable for at least 1 month prior to starting the ET intervention. In addition, studies that did not meet the Weston et al.¹² criteria of HIIT (e.g. 85%–95% peak heart rate or 80–100% peak work rate/peak VO_2) were excluded.

Data extraction and quality assessment

Two investigators (W.J.T. and R.I.B.) extracted all outcome data independently. When necessary, original investigators were contacted to clarify data or provide data in the required format (mean $\pm$ SD). Authors for 3 studies provided additional data.^{11,13,14} Quality was assessed using the previously validated Jadad scale¹⁵ (a 5-point scale used to assess the methodological quality of clinical trials by evaluating the adequacy of reporting for study randomization, double-blinding, and disclosure of withdrawals and dropouts) and adequacy of allocation concealment.

Data synthesis and analysis

Data were analyzed using the change score (post–pre) from baseline for both the exercise and control groups and the corresponding standard deviation was estimated using standard statistical methods assuming a correlation of 0.5 between pre and post measures.¹⁶ To pool the effect sizes, two models were used: 1) a fixed effect model using the inverse variance (IV) method; and 2) a random effects model using the method of DerSimonian and Laird (D + L), with the estimate of heterogeneity being taken from the IV fixed-effect model.^{17,18} Results were reported as weighted mean differences (WMD) with 95% confidence intervals (CIs) using random effects model. Heterogeneity was quantified using the I^2 statistic and a value of $I^2 \geq 50\%$ is considered substantial heterogeneity.¹⁸ Subgroup analyses by the length of exercise program (≥ 6 months vs. <6 months) were conducted to investigate possible sources of heterogeneity. Publication bias was tested visually using funnel plots¹⁹ and quantitatively using the Begg adjusted-rank correlation test²⁰ and Egger regression asymmetry test.²¹ There was no evidence of publication bias identified. All analyses were performed using Stata/SE 15 (StataCorp, College Station, Texas).

Results

Study screening, selection and evaluation

After removing duplicates, our search yielded 3714 citations from electronic databases. In addition, five citations were identified through hand-searches of reference lists. After initial screening, 51 full manuscripts were reviewed of which 33 were excluded for reasons

highlighted in (Fig 1). Screening consistency between independent evaluators was $\kappa = 0.88$ for interrater reliability.

Studies included in the systematic review

Eighteen unique randomized controlled ET trials were identified (Table 1).^{8,9,11,13,14,22–34} The trials included clinically stable older (mean age: 63 years, 77% male, predominantly ischemic etiology) HFrEF patients. Fourteen trials incorporated MICT,^{8,9,11,13,23–31,33} four trials compared HIIT with MICT or control,^{8,9,11,26} and four trials examined the effects of resistance training performed alone or in combination with aerobic training (resistance training alone, $n = 1$; combined aerobic and resistance training, $n = 3$).^{14,22,32,34} The training duration varied between 1 month and 10 years, with most of the trials lasting between 3 and 6 months.

Given the nature of the intervention, no trial was double-blind. Although all trials were randomized, only 6 (32%) described the randomization procedures or allocation concealment. Thirteen trials (68%) adequately detailed subject withdrawals/dropouts and 14 trials (74%) blinded ascertainment of LV remodeling outcomes. Accordingly, trials scored relatively poorly on the Jadad Scale: 5 received a score of 1 out of 5, 9 received a score of 2 out of 5, and 5 received a score of 3 out of 5.

Quantitative data synthesis

ET and LVEF—Compared to controls, MICT significantly improved LVEF (14 trials; 810 patients; WMD = 3.79%, 95% CI 2.08 to 5.50, Fig 2), however substantial heterogeneity across trials was found ($I^2 = 57.0\%$). Accordingly, further analysis based on ET program length (<6 months versus ≥6 months) revealed that MICT lasting <6 months modestly increased LVEF (9 trials; 463 patients; WMD = 2.33%, 95% CI 0.84 to 3.82%; $I^2 = 3.8\%$) while ET performed for 6 months or longer resulted in large significant increases in LVEF (5 trials; 347 patients; WMD = 6.26%; 95% CI 4.39 to 8.13%; $I^2 = 32.3\%$).

HIIT significantly increased LVEF compared to control (4 trials; 267 patients; WMD = 3.70%; 95% CI 1.63 to 5.77%; $I^2 = 8.5\%$, Fig 3) but no significant difference occurred with resistance training performed alone or in combination with aerobic training (4 trials; 95 patients; WMD = 1.94%; 95% CI –2.04 to 5.92%; $I^2 = 0.0\%$, Fig 4). Finally, in the 4 trials comparing HIIT and MICT, the increase in LVEF was not significantly different between groups (256 patients; WMD = 3.17%; 95% CI –0.87 to 7.22%; $I^2 = 66.7\%$, Fig 5).

ET and LV volumes—Compared to controls, ET was not associated with a significant change in LV end-diastolic volume (MICT, 6 trials,^{8,9,11,28,30,31} $n = 317$; WMD: 1.85 ml; 95% CI –5.03 to 8.73 ml, $I^2 = 0.0\%$; HIIT, 3 trials,^{8,9,11} $n = 235$; WMD: –2.13 ml, CI –9.57 to 5.32 ml, $I^2 = 0.0\%$; Resistance training performed alone or in combination with aerobic training, 2 trials,^{14,32} $n = 39$, WMD: –1.24 ml; 95% CI –31.64 to 29.16, $I^2 = 0.0\%$). No significant difference was found between HIIT and MICT for end-diastolic volume (3 trials,^{8,9,11} $n = 227$; WMD: –4.17 ml, CI –11.08 to 2.75 ml, $I^2 = 0.0\%$).

LV end-systolic volume did not significantly differ between MICT or resistance training performed alone or combined with aerobic training versus controls (MICT, 5 trials,

^{8,9,11,28,31} $n = 324$; WMD: -6.14 ml; 95% CI -12.83 to 0.54 ml, $I^2 = 0.0\%$; Resistance training performed alone or in combination with aerobic training, 2 trials^{14,32}; $n = 39$, WMD: -6.34 ml; 95% CI -34.78 to 22.09 ml, $I^2 = 0.0\%$). However, a significant decrease in LV end-systolic volume was found between HIIT and control (3 trials,^{8,9,11} $n = 235$; WMD: -10.66 ml, 95% CI -17.73 to -3.58 ml, $I^2 = 0.0\%$) but not versus MICT (3 trials,^{8,9,11} $n = 235$; WMD: -4.77 ml, 95% CI -10.90 to 1.37 ml, $I^2 = 0.0\%$).

ET and peak VO_2 —MICT significantly increased peak VO_2 compared to controls (13 trials^{8,9,11,13,23-26,28-31,33}; 749 patients, WMD = 2.67 ml·kg·min⁻¹; 95% CI 1.81 to 3.53 ml·kg·min⁻¹; $I^2 = 70.6\%$). Given heterogeneity across trials, further analysis by training length revealed that MICT performed for <6 months was associated with a modest increase in peak VO_2 (9 trials,^{8,9,11,24-26,30,31,33} 463 patients; WMD = 2.01 ml·kg·min⁻¹; 95% CI 1.06 to 2.96 ml·kg·min⁻¹; $I^2 = 54.0\%$) while a larger increase occurred with ET 6 months (4 trials,^{13,23,28,29} 286 patients, WMD = 3.88 ml·kg·min⁻¹; 95% CI 2.79 to 4.98 ml·kg·min⁻¹; $I^2 = 48.4\%$).

Compared to control, peak VO_2 was significantly higher with HIIT (4 trials,^{8,9,11,26} $n = 267$ patients; WMD = 3.63 ml·kg·min⁻¹; 95% CI 1.99 to 5.28 ml·kg·min⁻¹; $I^2 = 71.6\%$) and resistance training performed alone or in combination with aerobic training (3 trials^{14,22,34}; 79 patients; WMD = 3.19 ml·kg·min⁻¹; 95% CI 1.22 to 5.17 ml·kg·min⁻¹; $I^2 = 0.0\%$). Finally, the improvement in peak VO_2 was not different between HIIT and MICT (4 trials^{8,9,11,26}; 256 patients; WMD = 1.89 ml·kg·min⁻¹; 95% CI -0.39 to 4.16 ml·kg·min⁻¹; $I^2 = 86.7\%$).

Discussion

This 10-year review of ET trials in patients with HFrEF identified three major new findings: 1) In clinically stable HFrEF patients, MICT significantly improves LVEF compared to control with the greatest improvement occurring with long-term (> 6 months) training; 2) The increase in LVEF is significantly greater with HIIT compared to control but not compared to MICT; and 3) Resistance training performed alone or with aerobic training (MICT or HIIT) does not improve LVEF.

A 2007 meta-analysis by Haykowsky et al. found that MICT significantly increased LVEF (WMD: $+2.6\%$) and decreased LV volumes in clinically stable HFrEF patients.⁶ We extend these findings here to clinically stable HFrEF patients who are on optimal HF therapy, and show that the anti-remodeling benefit with MICT is dependent on the length of the ET program. The mechanisms underpinning this favorable adaptation could be attributed to a reduction in LV afterload which is associated with improved vascular endothelial function.^{35,36} Specifically, Hambrecht et al. reported a significant reduction in resting and peak exercise total peripheral resistance after 6-months of MICT compared to control.³⁶ The change in resting total peripheral resistance was also positively related to the change in LV end-diastolic diameter while changes in resting and peak exercise total peripheral resistance were inversely related to the changes in stroke volume.³⁶ In a subgroup of patients studied, the improvement in leg blood flow in response to intra-arterial acetylcholine infusion (endothelial-dependent vasodilator) after MICT was inversely related to changes in peak

exercise total peripheral resistance and positively related to the increase in peak VO_2 .^{35,36} Finally, our finding that peak VO_2 was significantly greater after long versus short-duration MICT is likely due to “central” adaptations as the change in maximal cardiac output is 3.5-fold higher after 6 months compared to 2 to 3 months of training in clinically stable HFrEF patients.^{26,36–39}

Four trials in this review compared the effects of HIIT versus MICT (2 to 3 months in duration, Table 1) or control on LV remodeling and peak VO_2 in clinically stable HFrEF patients.^{8,9,11,26} Our finding that HIIT was not associated with a greater improvement in LVEF and peak VO_2 compared to MICT may be related to the underlying intensity of the interval training exercise stimulus. For example, Wisloff et al. reported a superior increase in resting systolic mitral annulus excursion and velocity, LV outflow tract peak ejection velocity, isovolumic relaxation time, peak annular velocity during early filling, brachial artery endothelial function, mitochondrial function and peak VO_2 after 12 weeks of HIIT ($n = 9$) compared to MICT ($n = 9$) and controls ($n = 9$).⁸ Importantly, subjects in the HIIT group exercised at 92% of their peak heart rate while the MICT group exercised at 74% of their peak heart rate. In contrast, in the SMARTEX trial, no significant difference was found between HIIT and MICT for resting LVEF and peak VO_2 , however 51% of the HIIT participants exercised below their prescribed ET intensity while 80% of the MICT participants exercised above their prescribed intensity level.¹¹ Therefore, the interval training intensity may be an important determinant driving physiological benefit to LV function and peak VO_2 .

Our finding that resistance training performed alone or combined with aerobic training (MICT or HIIT) significantly increased peak VO_2 compared to control but was not associated with a significant change in LVEF is consistent with a recent meta-analysis in HFrEF patients.⁴⁰ The failure of this form of exercise to attenuate LV remodeling may be due to the increased pressure load associated with resistance exercise.^{41,42}

Study limitations

The trials in this review compared HIIT to MICT over 12 weeks in duration, consequently the long-term effects of HIIT on LV remodeling and peak VO_2 and its determinants remain unknown. Secondly, few studies compared the effects on LV remodeling of resistance training alone or resistance combined with aerobic training (MICT or HIIT). However, our results confirm our prior finding that this form of ET does not attenuate LV remodeling in clinically stable HFrEF patients.⁶ Not all studies included LV volume measures, however LVEF has been used extensively as a remodeling index⁴³, and a decrease in this outcome can occur via a reduction in end-diastolic and/or end-systolic volume. A final limitation common to cardiac rehabilitation and ET trials was that a majority of HFrEF patients in this analysis were males (77%).^{44,45} Thus, it is unclear if these findings extend to older female HFrEF patients. Despite these limitations, and unlike our 2007 meta-analysis using the same search strategy,⁶ the majority of the subjects in the trials included in this updated analysis were on evidence-based HF therapy, and most of the investigators performing the outcome measure analysis were blinded to group allocation.

Conclusions

In clinically stable HFrEF patients, MICT is an effective therapy to attenuate LV remodeling and improve peak VO_2 compared to control with the greatest benefits occurring with long-term (6 months) training. HIIT performed for 2 to 3 months is superior to control, but not MICT, for the improvement in LVEF and peak VO_2 . Finally, resistance training performed alone or combined with aerobic training (MICT or HIIT) significantly improved peak VO_2 without a change in LVEF.

Acknowledgments

The authors thank the following investigators^{11,13,14} for providing additional data for inclusion in this meta-analysis. Dr. Tucker is supported by American Heart Association (AHA) Grant (AHA Award Number: 18POST33990210). Dr. Haykowsky is supported by the Moritz Chair in Geriatrics at the University of Texas at Arlington, and National Institutes of Health (NIH) R15NR016826-01 grant.

Funding

None to declare.

Abbreviations and acronyms:

EF	ejection fraction
ET	exercise training
HF	heart failure
HFrEF	heart failure with reduction ejection fraction
HIIT	high-intensity interval training
LV	left ventricle
MICT	moderate-intensity continuous training
Peak VO_2	peak oxygen consumption
WMD	weighted mean difference

References

1. Benjamin EJ, Blaha MJ, Chiuve SE, et al. Heart disease and stroke statistics-2017 update: a report from the American Heart Association. *Circulation* 2017;135:e146–e603. [PubMed: 28122885]
2. Konstam MA, Kramer DG, Patel AR, Maron MS, Udelson JE. Left ventricular remodeling in heart failure: current concepts in clinical significance and assessment. *JACC Cardiovasc Imaging* 2011;4:98–108. [PubMed: 21232712]
3. Kim GH, Uriel N, Burkhoff D. Reverse remodelling and myocardial recovery in heart failure. *Nat Rev Cardiol* 2018;15:83–96. [PubMed: 28933783]
4. Nijst P, Martens P, Mullens W. Heart failure with myocardial recovery-the patient whose heart failure has improved: what next? *Prog Cardiovasc Dis* 2017;60:226–236. [PubMed: 28551473]
5. Kondamudi N, Haykowsky M, Forman DE, Berry JD, Pandey A. Exercise training for prevention and treatment of heart failure. *Prog Cardiovasc Dis* 2017;60:115–120. [PubMed: 28684221]

6. Haykowsky MJ, Liang Y, Pechter D, Jones LW, McAlister FA, Clark AM. A meta-analysis of the effect of exercise training on left ventricular remodeling in heart failure patients: the benefit depends on the type of training performed. *J Am Coll Cardiol* 2007;49:2329–2336. [PubMed: 17572248]
7. Karlsen T, Aamot IL, Haykowsky M, Rognmo O. High intensity interval training for maximizing health outcomes. *Prog Cardiovasc Dis* 2017;60:67–77. [PubMed: 28385556]
8. Wisloff U, Stoylen A, Loennechen JP, et al. Superior cardiovascular effect of aerobic interval training versus moderate continuous training in heart failure patients: a randomized study. *Circulation* 2007;115:3086–3094. [PubMed: 17548726]
9. Belardinelli R, Lacalaprice F, Ventrella C, Volpe L, Faccenda E. Waltz dancing in patients with chronic heart failure: new form of exercise training. *Circ Heart Fail* 2008;1:107–114. [PubMed: 19808280]
10. Haykowsky MJ, Timmons MP, Kruger C, McNeely M, Taylor DA, Clark AM. Meta-analysis of aerobic interval training on exercise capacity and systolic function in patients with heart failure and reduced ejection fractions. *Am J Cardiol* 2013;111:1466–1469. [PubMed: 23433767]
11. Ellingsen O, Halle M, Conraads V, et al. High-intensity interval training in patients with heart failure with reduced ejection fraction. *Circulation* 2017;135:839–849. [PubMed: 28082387]
12. Weston KS, Wisloff U, Coombes JS. High-intensity interval training in patients with lifestyle-induced cardiometabolic disease: a systematic review and meta-analysis. *Br J Sports Med* 2014;48:1227–1234. [PubMed: 24144531]
13. Passino C, Del Ry S, Severino S, et al. C-type natriuretic peptide expression in patients with chronic heart failure: effects of aerobic training. *Eur J Cardiovasc Prev Rehabil* 2008;15:168–172. [PubMed: 18391643]
14. Santos JM, Kowatsch I, Tsutsui JM, et al. Effects of exercise training on myocardial blood flow reserve in patients with heart failure and left ventricular systolic dysfunction. *Am J Cardiol* 2010;105:243–248. [PubMed: 20102926]
15. Jadad AR, Moore RA, Carroll D, et al. Assessing the quality of reports of randomized clinical trials: is blinding necessary? *Control Clin Trials* 1996;17:1–12. [PubMed: 8721797]
16. Follmann D, Elliott P, Suh I, Cutler J. Variance imputation for overviews of clinical trials with continuous response. *J Clin Epidemiol* 1992;45:769–773. [PubMed: 1619456]
17. Dersimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986;7: 177–188. [PubMed: 3802833]
18. Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ* 2003;327:557–560. [PubMed: 12958120]
19. Light R, Pillemer DB. *Summing Up. The Science of Reviewing Research* Cambridge, MA: Harvard University Press 1984.
20. Begg CB, Mazumdar M. Operating characteristics of a rank correlation test for publication bias. *Biometrics* 1994;50:1088–1101. [PubMed: 7786990]
21. Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997;315:629–634. [PubMed: 9310563]
22. Antunes-Correa LM, Nobre TS, Groehs RV, et al. Molecular basis for the improvement in muscle metaboreflex and mechanoreflex control in exercise-trained humans with chronic heart failure. *Am J Physiol Heart Circ Physiol* 2014;307:H1655–H1666. [PubMed: 25305179]
23. Belardinelli R, Georgiou D, Cianci G, Purcaro A. 10-year exercise training in chronic heart failure: a randomized controlled trial. *J Am Coll Cardiol* 2012;60:1521–1528. [PubMed: 22999730]
24. Brubaker PH, Moore JB, Stewart KP, Wesley DJ, Kitzman DW. Endurance exercise training in older patients with heart failure: results from a randomized, controlled, single-blind trial. *J Am Geriatr Soc* 2009;57:1982–1989. [PubMed: 20121952]
25. Eleuteri E, Mezzani A, Di Stefano A, et al. Aerobic training and angiogenesis activation in patients with stable chronic heart failure: a preliminary report. *Biomarkers* 2013;18:418–424. [PubMed: 23805979]
26. Fu TC, Wang CH, Lin PS, et al. Aerobic interval training improves oxygen uptake efficiency by enhancing cerebral and muscular hemodynamics in patients with heart failure. *Int J Cardiol* 2013;167:41–50. [PubMed: 22197120]

27. Hassanpour Dehkordi A, Khaledi Far A. Effect of exercise training on the quality of life and echocardiography parameter of systolic function in patients with chronic heart failure: a randomized trial. *Asian J Sports Med* 2015;6, e22643. [PubMed: 25883771]
28. Holtriegel R, Winzer EB, Linke A, et al. Long-term exercise training in patients with advanced chronic heart failure: sustained benefits on left ventricular performance and exercise capacity. *J Cardiopulm Rehabil Prev* 2016;36:117–124. [PubMed: 26906148]
29. Klecha A, Kawecka-Jaszcz K, Bacior B, et al. Physical training in patients with chronic heart failure of ischemic origin: effect on exercise capacity and left ventricular remodeling. *Eur J Cardiovasc Prev Rehabil* 2007;14:85–91. [PubMed: 17301632]
30. Malfatto G, Branzi G, Osculati G, et al. Improvement in left ventricular diastolic stiffness induced by physical training in patients with dilated cardiomyopathy. *J Card Fail* 2009;15:327–333. [PubMed: 19398081]
31. Muller L, Myers J, Kottman W, Luchinger R, Dubach P. Long-term myocardial adaptations after cardiac rehabilitation in heart failure: a randomized six-year evaluation using magnetic resonance imaging. *Clin Rehabil* 2009;23:986–994. [PubMed: 19880673]
32. Palevo G, Keteyian SJ, Kang M, Caputo JL. Resistance exercise training improves heart function and physical fitness in stable patients with heart failure. *J Cardiopulm Rehabil Prev* 2009;29:294–298. [PubMed: 19561521]
33. Sandri M, Kozarez I, Adams V, et al. Age-related effects of exercise training on diastolic function in heart failure with reduced ejection fraction: the Leipzig Exercise Intervention in Chronic Heart Failure and Aging (LEICA) Diastolic Dysfunction Study. *Eur Heart J* 2012;33:1758–1768. [PubMed: 22267243]
34. Stevens AL, Hansen D, Herbots L, et al. Exercise training improves insulin release during glucose tolerance testing in stable chronic heart failure patients. *J Cardiopulm Rehabil Prev* 2015;35:37–46. [PubMed: 25533903]
35. Hambrecht R, Fiehn E, Weigl C, et al. Regular physical exercise corrects endothelial dysfunction and improves exercise capacity in patients with chronic heart failure. *Circulation* 1998;98:2709–2715. [PubMed: 9851957]
36. Hambrecht R, Gielen S, Linke A, et al. Effects of exercise training on left ventricular function and peripheral resistance in patients with chronic heart failure: a randomized trial. *JAMA* 2000;283:3095–3101. [PubMed: 10865304]
37. Hambrecht R, Niebauer J, Fiehn E, et al. Physical training in patients with stable chronic heart failure: effects on cardiorespiratory fitness and ultrastructural abnormalities of leg muscles. *J Am Coll Cardiol* 1995;25:1239–1249. [PubMed: 7722116]
38. Coats AJ, Adamopoulos S, Radaelli A, et al. Controlled trial of physical training in chronic heart failure. Exercise performance, hemodynamics, ventilation, and autonomic function. *Circulation* 1992;85:2119–2131. [PubMed: 1591831]
39. Dubach P, Myers J, Dziekan G, et al. Effect of high intensity exercise training on central hemodynamic responses to exercise in men with reduced left ventricular function. *J Am Coll Cardiol* 1997;29:1591–1598. [PubMed: 9180124]
40. Jewiss D, Ostman C, Smart NA. The effect of resistance training on clinical outcomes in heart failure: a systematic review and meta-analysis. *Int J Cardiol* 2016;221:674–681. [PubMed: 27423089]
41. Cheetham C, Green D, Collis J, Dembo L, O'Driscoll G. Effect of aerobic and resistance exercise on central hemodynamic responses in severe chronic heart failure. *J Appl Physiol* (1985) 2002;93:175–180. [PubMed: 12070202]
42. McKelvie RS, McCartney N, Tomlinson C, Bauer R, MacDougall JD. Comparison of hemodynamic responses to cycling and resistance exercise in congestive heart failure secondary to ischemic cardiomyopathy. *Am J Cardiol* 1995;76:977–979. [PubMed: 7484846]
43. Bristow MR, Kao DP, Breathett KK, et al. Structural and functional phenotyping of the failing heart: is the left ventricular ejection fraction obsolete? *JACC Heart Fail* 2017;5: 772–781. [PubMed: 29096787]
44. Niederseer D, Thaler CW, Niederseer M, Niebauer J. Mismatch between heart failure patients in clinical trials and the real world. *Int J Cardiol* 2013;168:1859–1865. [PubMed: 23352093]

45. O'Connor CM, Whellan DJ, Lee KL, et al. Efficacy and safety of exercise training in patients with chronic heart failure: HF-ACTION randomized controlled trial. JAMA 2009;301:1439–1450. [PubMed: 19351941]

Fig 1.

Flow of trials through the selection process. LVEF: left ventricle ejection fraction.

Fig 2. Moderate-intensity continuous training (MICT) and left ventricle ejection fraction (LVEF) in all trials and split by intervention duration (<6 months or ≥ 6 months).

Fig 3.
High-intensity interval training (HIIT) and left ventricle ejection fraction (LVEF).

Fig 4.
Resistance training performed alone or combined with aerobic training and left ventricle ejection fraction (LVEF).

Fig 5.
High-intensity interval training (HIIT) versus moderate-intensity continuous training (MICT) and left ventricle ejection fraction (LVEF).

Table 1

Description of randomized controlled exercise training intervention studies included.

Study, year (Ref.)	Study sample	Sample size, n	Men, %	Mean age, years	HF+EF etiology	Exercise training intervention	Intervention duration, months	EF measure	Drug therapy, %
Antunes-Correa 2014 ²²	NYHA class II–III HF; LVEF <40%	ET, 17	76	56	Ischemic 35% Idiopathic 35%	RT + MICT 3 days/week, MICT 30–40 min at 60–72% VO _{2peak} + 10 min RT strength training exercises	3	Echo, SBP	ACE: 100; BB: 100; DIU: 71; SPIRO: 82
Belardinelli 2008 ⁹	NYHA class II–III HF; LVEF <40%	ET, 44	82	60	Prior MI 55% Prior CABG 77%	HIIT 3 days/week, 21 min/session, Dance 5 min slow, 3 min fast (88% HR _{peak} reported)	2	Echo, modified SSP	ACE: 80; BB: 80; DIGI: 14; DIU: 73; NIT: 18
Belardinelli 2012 ²³	NYHA class II–III HF; LVEF <40%	ET, 63	78	60	Ischemic 80% Non-ischemic 20	MICT 3 days/week, 40 min/session, 60–70% VO _{2peak}	120	Echo, not Specified	ACE: 71; AIIA: 24; BB: 46; DIGI: 19; DIU: 52
Brubaker 2009 ²⁴	LVEF <45%	ET, 30	63	70	Not reported	MICT 3 days/week, 30–40 min/session, 40% HRR × 2 weeks followed by 60–70% HRR.	4	Echo, not Specified	ACE: 80; BB: 20; CCB: 30; DIGI: 70; DIU: 87; NIT: 23
Eleuteni 2013 ²⁵	NYHA class II HF; LVEF <40%	ET, 11	100	66	Not reported	MICT 5 days/week, 30 min/session, 59–62% VO _{2peak}	3	Echo, BP	ACE: 86; BB: 14; CCB: 21; DIGI: 72; DIU: 86; NIT: 41
Ellingsen 2017 ¹¹	NYHA class II–III HF; LVEF <35%	ET, 77	81	65	Ischemic 60% Prior MI 57%, prior CABG 26%, prior PCI 46%	HIIT 3 days/week, 38 min/session, Intervals, 4 × 4 min at median 90% HR _{max} reported (88–92%, 95% CI) 3 min active recovery between	3	Echo, Modified SBP	ACE: 92; BB: 95; DIGI: 22; DIU: 75
Fu 2013 ²⁶	NYHA class II–III HF; LVEF <40%, or LVEF >40% with episodes of acute pulmonary edema	ET, 14	64	68	Ischemic 67% Non-ischemic 33%	MICT 3 days/week, 47 min/session, median 77% HR _{max} reported (74–82%, 95% CI)	3	Echo, not Specified	ACE: 79; BB: 93; CCB: 64; DIGI: 21; DIU: 50

Study, year (Ref.)	Study sample	Sample size, n	Men, %	Mean age, years	HFREF etiology	Exercise training intervention	Intervention duration, months	EF measure	Drug therapy, %
Hassanpour Dehkordi 2015 ²⁷	NYHA class II–III HF; LVEF <40%	ET, 13	62	66	Ischemic 60% Non-ischemic 40%	MICT 3 days/week, 30 min/session, 60% VO _{2peak}	3		ACE: 77; BB: 92; CCB: 62; DIGI: 31; DIU: 46
		CNT, 13	69	68	Ischemic 67% Non-ischemic 33%		3		ACE: 77; BB: 92; CCB: 69; DIGI: 23; DIU: 46
		ET, 30	60	60	Ischemic 70% Non-ischemic 30%	MICT 3 days/week, 25–35 min/session, 60–70% HRR	6	Echo, not Specified	ACE: 80; DIGI: 70; DIU: 93
		CNT, 31	74	58	Ischemic 65% Non-ischemic 35%		6		ACE: 87;DIGI: 71; DIU: 90
Holtriegel 2016 ²⁸	NYHA class IIIB HF; LVEF <30%	ET, 18	100	60	Ischemic 56% DCM 44%	MICT 5 days/week, 30 min/session, 50–60% VO _{2peak}	12	Echo, not Specified	ACE: 100; BB: 94; DIGI: 56; DIU: 94
Klecha 2007 ²⁹	NYHA class II–III HF; LVEF <35%	CNT, 19	100	62	Ischemic 53% DCM 47%		12		ACE: 100; BB: 100; DIGI: 21; DIU: 100
		ET, 25	80	60	Ischemic 100%	MICT 3 days/week, 25 min/session, 80% HR _{max}	6	MRI, not specified	ACE: 100; BB: 100; DIGI: 36; DIU: 64; NIT: 36
Malfatto 2009 ³⁰	DCM	CNT, 25	72	61	Ischemic 100%		6		ACE: 100; BB: 100; DIGI: 32; DIU: 68 NIT: 68
		ET, 27	70	65	Ischemic 52% Non-ischemic 48%	MICT 3 days/week, 40 min/session, 60% VO _{2peak}	3	Echo, not specified	ACE: 89; BB: 81; DIGI: 4; DIU: 67; SPIRO: 52
		CNT, 27	74	67	Ischemic 59% Non-ischemic 41%		3		ACE: 81; BB: 78; DIGI: 7; DIU: 70; SPIRO: 48
Muller 2009 ³¹	LVEF <40%	ET, 8	100	47	Not reported	MICT 5 days/week, 30 min/session, +45 min walk twice daily, 60–80% HRR	1	MRI, Stack	ACE: 100; BB: 75; DIGI: 38; DIU: 88
Palevo 2009 ³²	NYHA class II–III HF; LVEF <40%	CNT, 8	100	56			1		ACE: 88; BB: 50; DIGI: 25; DIU: 75
		ET, 10	94 ^d	70	Ischemic 100%	RT 3 days/week, 12 strength exercises, 2 sets, 12–15 reps, 60% 1RM	2	Echo, not specific	ACE: 100; BB: 100
Passino 2008 ¹³	NYHA class <IV HF; LVEF <45%	CNT, 6		65	Ischemic 100%		2		ACE: 100; BB: 100
		ET, 71	87	61	Ischemic 55% Idiopathic 45%	MICT 3 days/week, 30 min/session, 65% VO _{2peak}	9	Echo, not specific	ACE: 75; BB: 73; DIU: 79; SPIRO: 38
		CNT, 19	74	63	Ischemic 42% Idiopathic 58%		9		ACE: 74; BB: 74; DIU: 84; SPIRO: 37
Sandri 2012 ³³	NYHA class II–III HF; LVEF <40%	ET, 30	80	61	Ischemic 60% DCM 40%	MICT 4 days/week, 4 sessions/day, 20 min/session, 70% VO _{2peak}	1	Echo, 4C Simpson Disk	ACE: 83 BB: 100; DIU: 83; SPIRO: 43
Santos 2010 ¹⁴	LVEF <45%	CNT, 30	83	61	Ischemic 67% DCM 33%		1		ACE: 83; BB: 100; DIU: 83; SPIRO: 50
		ET, 13	69	53	Not reported	RT + MICT 3 days/week, MICT 25–40 min per session at HR of AT + RT 10 min of strength exercises	4	Echo, modified SBP	ACE: 100; BB: 100; DIGI: 38; DIU: 85
		CNT, 10	40	59.4			4		ACE: 70; BB: 100; DIGI: 0; DIU: 40

Study, year (Ref.)	Study sample	Sample size, <i>n</i>	Men, %	Mean age, years	HF/EF etiology	Exercise training intervention	Intervention duration, months	EF measure	Drug therapy, %
Stevens 2015 ³⁴	NYHA class III HF	ET, 15	67	67	Ischemic 40%	RT + HIIT 5 sessions every 2 weeks, 24–60 min/session. HIIT 4 bouts of 6–10 min cycle/walk at critical power (~80% VO _{2peak}) with 2 min rest between bouts RT 2–3 sets, 10–15 reps 50–70% 1RM	3	Echo, modified	ACE: 87, BB: 93, DIU: 80
Wisloff 2007 ⁸	>12 months post-MI, LVEF <40%	CNT, 7	86	64	Ischemic 57%	HIIT 3 days/week, 38 min/session, Intervals, 4 × 4 min at mean 92 ± 2% HR _{max} reported, 3 min active recovery between.	3	SBP	ACE: 100, BB: 100, DIU: 86
		ET, 9	78	77	Post-MI 100%		3	Echo, modified SBP	ACE: 100, BB: 100, DIU: 56 NIT: 44
		ET, 9	78	74	Post-MI 100%	MICT 3 days/week, 47 min/session, mean 74 ± 2% HR _{max} reported	3		ACE: 100, BB: 100, DIU: 44, NIT: 56
		CNT, 9	67	76	Post-MI 100%	MICT 1 day ever 3 weeks, 47 min/session, mean 71 ± 2% HR _{max} reported	3		ACE: 100, BB: 100, DIU: 56, NIT: 44

All data are mean values unless otherwise indicated. *Abbreviations:* ACE: angiotensin converting enzyme inhibitor, AIA: angiotensin 1 antagonist, AIIA: angiotensin 2 antagonist, AT: anaerobic threshold, BB: beta-blocker, CABG: coronary artery bypass graft, CCB: calcium channel blocker, CI: confidence interval, CNT: control, DCM: dilated cardiomyopathy, DIGI: digoxin, DIU: diuretic, EF: ejection fraction, ET: exercise training, HF: heart failure, HIIT: high-intensity interval training, HR_{max}: heart rate max, HRR: heart rate reserve, LVEF: left ventricular ejection fraction, MI: myocardial infarction, MICT: moderate-intensity continuous training, NIT: nitrates, NYHA: New York Heart Association, PCI: percutaneous coronary intervention, RT: resistance training, SBP: Simpson’s Biplane, SPIRO: spirinolactone, SSP: Simpson’s Single Plane, VO_{2peak}: peak oxygen consumption, 1RM: 1-repetition maximum, 4C: 4-chamber.

^aMedian values.