Published in final edited form as:

Chem Commun (Camb). 2019 March 19; 55(24): 3469–3472. doi:10.1039/c9cc01068h.

Red-emitting pyrene-benzothiazolium: Unexpected selectivity to lysosomes for real-time cell imaging without alkalinizing effect

Chathura S. Abeywickrama^a, Kaveesha J. Wijesinghe^c, Robert V. Stahelin^d, and Yi Pang^{a,b,*}

^a. Department of Chemistry, University of Akron, Akron, Ohio 44325, USA.

^b. Maurice Morton Institute of Polymer Science, University of Akron, Akron, Ohio 44325, USA.

^c. Department of Chemistry and Biochemistry, University of Notre Dame, Notre Dame, Indiana, USA

^d. Department of Medicinal Chemistry and Molecular Pharmacology, Purdue University, West Lafayette, Indiana, 47907, USA.

Abstract

A series of pyrene-benzothiazolium probes were synthesized. By replacing the pyridinium with a benzothiazolium unit, the selectivity of pyrene-derivatives is found to switch from nuclear to cellular lysosomes. New probes do not require proton participation and exhibit high biocompatibility and long-term imaging ability.

Graphical Abstract

Lysosomes are membrane-encapsulated organelles that contain degradative enzymes to digest macromolecules within the cell.^{1,2} Since the discovery of lysosomes by Duve in 1955,³ the organelles are known to contain a variety of digestive enzymes called “hydrolases”, which exhibit optimal activity in a narrow acidic pH environment within the lysosomal lumen (pH 4.5 – 6).⁴ Abnormal lysosome activities or lysosome malfunction has been attributed to many disease conditions including neurodegenerative disorders, lysosome storage diseases, cancer and inflammations.^{4–13}

* Corresponding Author: yp5@uakron.edu.

†Footnotes relating to the title and/or authors should appear here.

Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here].
See DOI: 10.1039/x0xx00000x

Imaging lysosomes in live or fixed cell samples has recently received much attention due to their involvement in cell processing activities and cancer.^{12,14–17} Most probe design incorporates a basic amino group for lysosome selectivity, which include commercial LysoTracker® Green DND-26 ($\lambda_{\text{ex}} \sim 504$ nm, $\lambda_{\text{em}} \sim 510$ nm) and LysoTracker® Red DND-99 ($\lambda_{\text{ex}} \sim 577$ nm, $\lambda_{\text{em}} \sim 590$ nm).¹⁸ Since the existing fluorescent probes require an acidic media to operate, their accumulation in lysosomes will cause the pH increase (the “alkalinizing effect”) that perturbs the normal cell activity.^{19–22} Although new lysosome probes have been developed recently with spirolactams²³ and peptide²⁴ based molecules, protonation of an amine group remains an essential step, in order to guide the probe molecules into the acidic lysosome lumen. There are also interests in developing new probes that exhibit emission in the near infrared (NIR) region.^{25,24,26} However, no lysosome probes are found to operate without resorting to protonation in the lumen, which is known to alter the intracellular pH to affect normal cell activities.

As a planar molecular skeleton with extensive conjugation, pyrene has been used for developing fluorescent sensors,^{27–32} including those for intracellular membranes³⁰ and mitochondria.³¹ Our recent study shows that probe **1** is a bright red-emitting dye for cell nucleus staining.³² The nucleus selectivity of **1** is likely achieved via binding to DNA minor groove (Figure 1a). During the interaction, the ammonium cation is required to fit into the shallow space of the DNA minor groove for effective polar interaction with the DNA backbone. In an effort to examine the substituent impact on the cell staining, herein we report the synthesis of **2**, where the substituent R on the benzothiazolium could provide sufficient steric hindrance to prevent the ammonium cation to fully enter the DNA minor groove, thereby decreasing its interaction with the cell nucleus. Surprisingly, compound **2** exhibited remarkable selectivity for intracellular lysosomes, in sharp contrast to **1**. Since **2** has no group for protonation, its ability to generate bright red fluorescence, upon binding to lysosomes, provides a promising strategy to achieve long term tracking of lysosomes by eliminating harmful “alkalinizing effects”.

Synthesis.

The probe **2** with different substituents was synthesized in good yields (ESI Figure S1).^{26,32} Chemical structures of probe **2** were characterized by ¹H NMR and ¹³C NMR spectroscopy and high-resolution mass spectrometry (ESI, Figure S2 – S3).

Optical properties.

In dichloromethane (DCM), probes **2** exhibited $\lambda_{\text{abs}} \approx 530$ nm - 540 nm (Table 1 ESI Figure S4), which, is significantly red-shifted from the monomeric pyrene ($\lambda_{\text{abs}} \approx 352$ nm). The observed large bathochromic shift could be attributed to the strong intra-molecular charge transfer (ICT) from pyrene to benzothiazolium moiety in **2**, which was verified by spectroscopic study at low temperature (in liquid nitrogen at -188 °C) (ESI Figure S5). The probe **2** also showed bright red emission with $\lambda_{\text{em}} \sim 630$ nm – 640 nm ($\Phi_{\text{fl}} \approx 0.22$ – 0.48 in DMSO), along with a large Stokes' shift ($\lambda \approx 140$ nm, 4170 cm⁻¹). Interestingly, fluorescent quantum yield of **2** was significantly lower in an aqueous environment ($\Phi_{\text{fl}} \approx$

0.05) than that in a non-aqueous media (Table 1), making it possible for cell staining under “wash-free” conditions.

Biological cell studies.

Bright red emission was observed when using **2** (500 nM) to stain different cell lines (COS-7, HEK293, A549 and Huh 7.5) (ESI Figure S7). Clear cell imaging was observable after 30 minutes of staining without any washing post-staining. The non-uniform pattern observed from the cell imaging (Figure 2) suggested that the dyes might be selectively bound to certain intracellular organelles. Possibility of binding to mitochondria was ruled out by co-staining **2** with commercial MitoTracker® Green FM on COS-7 cells (ESI Figure S8). The intracellular location of **2** was eventually confirmed by co-staining with LysoTracker® Green DND-26, which showed excellent co-localization (Figure 2, ESI Figure S16). The calculated Mander's correlation coefficients for probes **2a** - **2d** were > 0.9 in all cell lines, indicating the exceptional lysosome specificity (ESI figure S9). Consistent results were also observed in the co-localization with LysoTracker® Green DND-26 in HEK 293 cells (ESI Figure S10) and in cancer cell lines (A549 and Huh 7.5), showing the probe's ability to visualize lysosome in live cells. The observed lysosome selectivity from **2** was in sharp contrast to the nucleus selectivity reported from **1**,³² showing a large impact of the associated cyanine segment on the organelle selectivity.

The LC₅₀ values for probe **2** were calculated to be in the 25 – 30 µM range by using CellTiter-Glow® Luminescent cell viability assay in COS-7 cells (ESI figure S22). The result further confirmed that these probes are suitable for biological cell studies since the working concentration of the probes are far below their calculated LC₅₀ value. Bright fluorescent confocal microscopy images could be obtained when probe **2** concentration was as low as 100 nM (ESI Figure S15).

Long Term Imaging.

Lack of cellular pH perturbation, in addition to its low toxicity, raises possibility of using **2** for long term lysosome imaging. Thus, A549 cells were treated with **2d** (500 nM) over a period of 6 hours, which gave stable fluorescent confocal microscopy images (Figure 3a, ESI Figure S18). In sharp contrast, cells stained with commercial LysoTracker® Red DND-99 showed significant morphological changes and notable cell shrinkage after 1.5 hours (Figure 4a, ESI Figure S19). The fluorescence signals of **2d** and LysoTracker® Red was further analyzed by plotting their average fluorescence intensity vs. staining time (Figure 3b). No apparent intensity change were observed from **2b** in A549 cells over a period of 6 hours. However, LysoTracker® exhibited a significant decrease in relative intensity with time, which could be attributed to the fluorescence quenching due to pH elevation within lysosomes (i.e. alkalinizing effect). In another experiment, COS-7 cells were incubated with probe **2d** for 30 min and 24 hrs, and clear fluorescence signal was still observable from **2d** even after 24 hrs in comparison with initial confocal microscopy images (ESI Figure S21). The ability of **2** in giving stable fluorescence signals in intracellular environment (e.g. on lysosomes, Figure 3b) thus opened the possibility for long term imaging, due to elimination of pH sensitivity.

Photostability.

Photostability was evaluated by studying the samples of A549 cells that were stained with **2b**, **2d** or LysoTracker® Red for 30 minutes separately. The resulting samples were continuously irradiated with a 561 nm laser on the microscope (ESI Figure S23 – 24). The fluorescence intensity, averaged by measurement on samples ($n > 30$), was plotted as a function of irradiation time (Figure 3c). Probe **2d** exhibited excellent stability in comparison with the commercial LysoTracker® Red, without showing significant photobleaching during the experiment period (Figure 3c). The result also suggested that the substituent on the benzothiazolium segment had a large impact on the photostability of **2**. A possible reason for poor photostability of **2b** is attributed to possible photodegradation of the excited probe in the presence of molecular oxygen where singlet oxygen ($^1\text{O}_2$) or superoxide anion (O_2^-) upon interactions of the excited probe **2** with molecular oxygen (See ESI Figure S28).^{33,34} Therefore, relative electron density on cyanine N atom can affect their photostability (**2d** exhibited a greater photostability as a result of having electron rich benzyl substituent).³⁴

Possible probe Interaction.

Since the probe structure does not include a basic amino group, the intriguing question is how the probe **2** was accumulated in cellular lysosomes. Since both pyrene and benzothiazolium segments are quite inert to protonation, the acidic environment in the organelle might not promote accumulation of **2** in the lysosome lumen. The hydrophobic nature of pyrene segment thus led us to assume that probe **2** might enter hydrophobic regions (i.e., lysosomal membrane or lysosomal proteins) rather than the aqueous lysosomal lumen. One possibility is that **2** could be associated with the lysosomal membrane (hydrophobic region). In order to evaluate the hypothesis, we carried out a study involving *in vitro* generated lipid vesicles that can mimic an artificial phospholipid bilayer structure similar to cellular membranes. In the control experiment, probe **2** did not show any appreciable fluorescent signal in aqueous HEPES buffer (pH=7.4). Interestingly, the pyrene-based molecules (**2b** and **2d**) exhibited a large fluorescent turn on in the presence of liposome vesicles (Figure 4a). This observation indicated that the possibility to associate with hydrophobic membrane components of the lysosome, generating fluorescence turn on signal. The assumption is consistent with the literature report where pyrene-based probes are able to stain biological membranes due to their hydrophobic nature.³⁰

Fluorescence of **2** in different pH aqueous solutions revealed that emission was only slightly affected (about $\pm 10\%$) in a wide pH range (pH=3.2 – 11.4) (ESI Fig. S6). Since the fluorescence of **2** did not change significantly from neutral (pH $\approx$ 7.2) to acidic (pH $\approx$ 4.5 within lysosome lumen), in intracellular it is possible that the dye is binding to a hydrophobic region of lysosomes. In another experiment, an acidic aqueous solution of (pH = 4.5) probe **2d** was titrated with 10% bovine serum albumin (BSA). Interestingly, addition of BSA led to a significant fluorescent enhancement ($\approx 600\%$) (Figure 4b). It should be pointed out that BSA binding did not cause fluorescence enhancement in commercial LysoTracker® probes, since the PET effect from an amine group would be stronger in BSA than in acidic aqueous.³⁵ Therefore, interaction mechanism of the probe **2** in cellular

lysosomes should be significantly different from that of acidotropic commercial LysoTracker® probes.

In conclusion, new fluorescent probes series have been successfully developed by coupling pyrene and benzothiazolium segments. In addition to bright red emission with large Stokes' shift ($\lambda > 130$ nm), the pyrene-based probes **2** exhibited the following attractive properties that are distinctive from the existing lysosome probes: (1) low cytotoxicity ($LC_{50} > 25 \mu M$); (2) large fluorescence turn on in hydrophobic environments, such as hydrophobic lysosomal lipid membrane, to enable wash-free staining; and (3) stable emission throughout a wide pH range (1.0 – 12.0), as the probe structure is insensitive to acidity. Although their structures do not contain an amino group, the new probes (**2a – 2d**) exhibit remarkable selectivity for visualizing lysosomes in both healthy and tumor cells. By eliminating the “alkalinizing effect”, the new probes have significant advantages over the existing commercial LysoTracker® probes and can be used for long term tracking of lysosome activity. This study points to the possibility of developing lysosome probes by targeting the lysosome membranes, rather than the acidic lysosome lumens where the enzymatic activities occur. Since probes **2** do not exhibit “alkalinizing effect” and exert minimal perturbation on the digestive activities of enzymes, they could be valuable tools for monitoring lysosome activities in biological research.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENT

Y.P. acknowledges support from the University of Akron via Coleman endowment, and partial support from NIH (Grant no. 1R15GM126438-01A1). K.J.W. was supported by a NIH CBBI fellowship (T32GM075762). The imaging studies were supported by the Indiana University School of Medicine-South Bend Imaging and Flow Cytometry core (to R.V.S.).

References

1. Yapici NB, Bi Y, Li P, Chen X, Yan X, Mandalapu SR, Faucett M, Jockusch S, Ju J and Gibson KM, Sci. Rep.
2. Chen X, Bi Y, Wang T, Li P, Yan X, Hou S, Bammert CE, Ju J, Gibson KM, Pavan WJ and Bi L, Sci. Rep., DOI:10.1038/srep09004.
3. de Duve C and Wattiaux R, Annu. Rev. Physiol, 1966, 28, 435–492. [PubMed: 5322983]
4. Kirkegaard T and Jäättelä M, Biochim. Biophys. Acta - Mol. Cell Res, 2009, 1793, 746–754.
5. Maes H and Agostinis P, Mitochondrion, 2014, 58–68.
6. Honscheid P, Datta K and Muders MH, Int J Radiat Biol, 2014, 90, 628–635. [PubMed: 24678799]
7. Jiang P and Mizushima N, Cell Res., 2014, 24, 69. [PubMed: 24323045]
8. He X, Li J, An S and Jiang C, Ther. Deliv, 2013, 4, 1499–1510. [PubMed: 24304248]
9. Reyjal J, Cormier K and Turcotte S, in Advances in Experimental Medicine and Biology, 2014, vol. 772, pp. 167–188. [PubMed: 24272359]
10. Platt FM, Nat. Rev. Drug Discov, DOI:10.1038/nrd.2017.214.
11. Ashoor R, Yafawi R, Jessen B and Lu S, PLoS One., DOI:10.1371/journal.pone.0082481.
12. Jaattela M, Oncogene, 2004, 23, 2746–2756. [PubMed: 15077138]
13. Boya P and Kroemer G, Oncogene, 2008, 27, 6434–6451. [PubMed: 18955971]

14. Reinheckel T, Deussing J, Roth W and Peters C, Biol. Chem, 2001, 382, 735–741. [PubMed: 11517926]
15. Balaji KN, Schaschke N, Machleidt W, Catalfamo M and Henkart PA, J. Exp. Med, 2002, 196, 493–503. [PubMed: 12186841]
16. Felbor U, Kessler B, Mothes W, Goebel HH, Ploegh HL, Bronson RT and Olsen BR, Proc Natl Acad Sci U S A, 2002, 99, 7883–7888. [PubMed: 12048238]
17. Guicciardi ME, Leist M and Gores GJ, Oncogene, 2004, 23, 2881–2890. [PubMed: 15077151]
18. Haugland RP, The handbook: a guide to fluorescent probes and labeling technologies, Molecular probes, 2005.
19. Wiederschain GY, Biochem, 2011, 76, 1276–1276.
20. Zhang X, Wang C, Han Z and Xiao Y, ACS Appl. Mater. Interfaces, 2014, 6, 21669–21676. [PubMed: 25382852]
21. Zhu H, Fan J, Xu Q, Li H, Wang J, Gao P and Peng X, Chem. Commun, 2012, 48, 11766–11768.
22. Kim HM, An MJ, Hong JH, Jeong BH, Kwon O, Hyon JY, Hong SC, Lee KJ and Cho BR, Angew. Chemie - Int. Ed, 2008, 47, 2231–2234.
23. Wang B, Yu S, Chai X, Li T, Wu Q and Wang T, Chem. - A Eur. J, 2016, 22, 5649–5656.
24. Han Y, Li M, Qiu F, Zhang M and Zhang Y-H, Nat. Commun, 2017, 8, 1307. [PubMed: 29101340]
25. Zhang X, Wang C, Han Z and Xiao Y, ACS Appl. Mater. Interfaces, 2014, 6, 21669–21676. [PubMed: 25382852]
26. Dahal D, McDonald L, Bi X, Abeywickrama C, Gombiedza F, Konopka M, Paruchuri S and Pang Y, Chem. Commun, 2017, 53, 3697–3700.
27. Niko Y, Cho Y, Kawauchi S and Konishi G, RSC Adv, 2014, 4, 36480.
28. Niko Y, Kawauchi S and Konishi GI, Chem. - A Eur. J, 2013, 19, 9760–9765.
29. Niko Y, Kawauchi S, Otsu S, Tokumaru K and Konishi GI, J. Org. Chem, 2013, 78, 3196–3207. [PubMed: 23425392]
30. Niko Y, Didier P, Mely Y, Konishi GI and Klymchenko AS, Sci. Rep., DOI:10.1038/srep18870.
31. Niko Y, Moritomo H, Sugihara H, Suzuki Y, Kawamata J and Konishi G, J. Mater. Chem. B, 2015, 3, 184–190.
32. Abeywickrama CS, Wijesinghe KJ, Stahelin RV and Pang Y, Chem. Commun, 2017, 53, 5886–5889.
33. Chen C, Zhou B, Lu D and Xu G, J. Photochem. Photobiol. A Chem, 1995, 89, 25–29.
34. Chen X, Peng X, Cui A, Wang B, Wang L and Zhang R, J. Photochem. Photobiol. A Chem, 2006, 181, 79–85.
35. Bertman KA, Abeywickrama CS, Baumann HJ, Alexander N, McDonald L, Shriver LP, Konopka M and Pang Y, J. Mater. Chem. B, 2018, 6, 5050–5058.

Figure 1.

Structure of pyrene-based probes **1** and **2** and their selectivity switch from nucleus to lysosomes. And schematic illustration of lysosome structure (c).

Figure 2.

Fluorescence confocal microscopy images of COS-7, A549 and Huh 7.5 cells incubated for 30 minutes with probe **2d** (500 nM) with LysoTracker® Green DND-26 (70 nM). Images a, d, and g shows fluorescence microscope images of LysoTracker® Green DND-26 (70 nM) images b, e, and h represent images of probe **2d**. Images c, f, and i represent overlapped figures for **2d** with LysoTracker® Green. Probe **2d** was excited with a 561 nm laser and LysoTracker® Green was excited with a 488 nm laser line. All images were obtained at 63x oil magnification. Mander's overlap coefficient calculated for **2d** was found to be 0.93, 0.94 and 0.90, respectively, in COS-7, A549 and Huh 7.5 cells. (Number of replicates = 3, n = 40 cells).

Figure 3.

(a) - Time based photo-stability experiment for A549 cells incubated with **2d** (500 nM) and LysoTracker® Red DND-99 (100 nM) for 30 minutes. Images were obtained at 30 minute intervals under consistent parameters of the microscope. Probe **2d** and LysoTracker® Red were excited with 561 nm laser. (b) - Relative average fluorescence intensity obtained from A549 cells stained with probe **2d** (500 nM) and LysoTracker Red (70 nM) for a period of 3 hours. Probe **2d** and LysoTracker® Red were excited with a 561 nm laser. (c) - The plot of recovered fluorescence intensity versus irradiation time calculated for LysoTracker® Red (100 nM), **2b** (100 nM) and **2d** (100 nM). All probes were stained in A549 cells, and excited with a 561 nm laser.

Figure 4.

(a) Fluorescence emission obtained for **2b** and **2d** (1×10^{-5} M) upon introduction in to 1 mM lipid vesicles (PC:PE 80:20) in HEPES buffer (pH = 7.4). Probes were excited at 530 nm. (b) Represents the fluorescence emission enhancement observed in **2d** (1×10^{-5} M) upon addition of 10% bovine serum albumin (BSA) in to an aqueous solution (pH 4.5) of probe **2d** (1×10^{-6} M). Probe **2d** was excited at 500 nm for emission collection.

Table 1.Optical properties of probes **2a – 2d**

Solvent	Property	Probe			
		2a	2b	2c	2d
DCM	λ_{abs}	530 nm	531 nm	532 nm	542 nm
	λ_{em}	633 nm	632 nm	634 nm	641 nm
	(QY)	(0.15)	(0.16)	(0.12)	(0.17)
DMSO	λ_{abs}	479 nm	483 nm	481 nm	492 nm
	λ_{em}	638 nm	636 nm	638 nm	645 nm
	(QY)	(0.22)	(0.27)	(0.31)	(0.48)
EtOH	λ_{abs}	491 nm	495 nm	494 nm	507 nm
	λ_{em}	627 nm	628 nm	628 nm	637 nm
	(QY)	(0.13)	(0.15)	(0.19)	(0.21)
Water	λ_{abs}	468 nm	470 nm	471 nm	481 nm
	λ_{em}	632 nm	631 nm	630 nm	640 nm
	(QY)	(0.054)	(0.051)	(0.063)	(0.061)