Published in final edited form as:

Ann Epidemiol. 2019 April ; 32: 20–27. doi:10.1016/j.annepidem.2019.01.010.

Perceived stress and incident sexually transmitted infections in a prospective cohort

Rodman Turpin¹, Rebecca M. Brotman², Ryan S. Miller³, Mark A. Klebanoff⁴, Xin He¹, and Natalie Slopen¹

¹Department of Epidemiology and Biostatistics, University of Maryland School of Public Health

²Institute for Genome Sciences, University of Maryland School of Medicine

³Division of Pediatric Endocrinology, University of Maryland School of Medicine

⁴Center for Perinatal Research, The Research Institute at Nationwide Children's Hospital

Abstract

Purpose: Psychosocial stress has been associated with susceptibility to many infectious pathogens. We evaluated the association between perceived stress and incident sexually transmitted infections (STIs, *Chlamydia trachomatis*, *Neisseria gonorrhoeae*, and *Trichomonas vaginalis* genital infections) in a prospective study of women. Stress may increase vulnerability to STIs by suppressing immune function and altering the protective vaginal microbiota.

Methods: Utilizing the 1999 Longitudinal Study of Vaginal Flora (n=2,439), a primarily African-American cohort of women, we fitted Cox proportional hazards models to examine the association between perceived stress and incident STIs. We tested bacterial vaginosis (measured by Nugent Score) and sexual behaviors (condom use, number of partners, partner concurrence) as mediators using Vanderweele's difference method with bootstrapping.

Results: Baseline perceived stress was associated with incident STIs both in the unadjusted model (HR=1.019; 95% CI 1.009–1.029) and after adjusting for confounders (aHR=1.015; 95% CI 1.005–1.026). Nugent score and sexual behaviors significantly mediated 21 % and 65% of this adjusted association respectively, and 78% when included together in the adjusted model.

Conclusions: This study advances understanding of the relationship between perceived stress and STIs and identifies high-risk sexual behaviors and development of BV—both known risk factors for STIs—as potential mechanisms underlying this association.

Address correspondence to: Rodman Turpin, PhD, MS, University of Maryland, College Park, School of Public Health, Department of Epidemiology and Biostatistics, 4200 Valley Dr #2242, College Park, MD 20742, RTurpin@umd.edu, Phone: 301-405-3575, Fax: 301-314-9366.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Obtaining Data: LSVF data are available by submitting a request to the NICHD/DIPHR Biospecimen Repository Access and Data Sharing (BRADS) program.

Conflict of Interest: The authors have no conflicts of interest to disclose.

Keywords

Psychological Stress; Sexually Transmitted Diseases; Gonorrhea; Chlamydia; Trichomonas; Bacterial Vaginosis; Sexual Behavior; Prospective Studies

Introduction

According to 2016 Centers for Disease Control (CDC) estimates, there are approximately 20 million new sexually transmitted infections (STIs) in the U.S. each year.¹ In 2008, the annual national burden of STIs was in excess of \$16 billion in direct medical costs to the healthcare system.^{2, 3} In addition to their own morbidities, STIs such as genital *Neisseria gonorrhoeae* (GC), *Chlamydia trachomatis* (CT), and *Trichomonas vaginalis* (TV), are risk factors for infertility⁴ and acquisition of human immunodeficiency virus (HIV).⁵ Population-based surveys show marked and persistent racial/ethnic disparities in the STI and HIV prevalence.^{6–8} Individual risk behaviors, such as number of sexual partners and a lack of condom use, do not fully account for the high rate of infection among African-Americans in the U.S.^{9–15} Perceived stress is a well-known psychosocial factor that increases risk for many health outcomes¹⁶, as stress can impair multiple physiological systems^{17, 18} and increase the likelihood of high risk behaviors.^{19, 20} Limited research has examined psychosocial stress in relation to vulnerability to STIs, although extensive research indicates that high psychosocial stress is associated immune suppression, leading to heightened infection susceptibility, severity and persistence of infections.^{16, 21–27}

Perceived stress refers to thoughts or feelings one has about their level of experienced stress. It is associated with STI risk due to multiple biological and behavioral mechanisms. On the biological front, several studies—including one with participants from the Longitudinal Study of Vaginal Flora (LSVF) (the same cohort used in the current study)—have documented an independent association between psychosocial stress and greater prevalence and incidence of bacterial vaginosis (BV), a dysbiosis of the vaginal microbiota.^{25–27} Amabebe and Anumba also recently reviewed how cortisol, a biologic measure of stress, can potentially affect vaginal health.²⁸ BV has been linked to an approximate 2-fold increased risk for acquisition of STIs, including HIV, GC, CT, TV, herpes simplex virus and human papillomaviruses.^{29–35} The vaginal microbiome offers protection in part through the influential action of *Lactobacillus* spp. Species of vaginal *Lactobacillus* spp. can provide broad-spectrum protection through production of copious amounts of lactic acid³⁶, bacteriocins^{29, 37–40}, antagonistic bacteriocin-like substances⁴¹, and biosurfactants⁴², and through their ability to adhere to mucus, forming a physical barrier against incoming pathogens⁴³ and disrupting biofilms⁴⁴.

Considering behavioral pathways, high perceived stress has been associated with a 2.4-fold greater odds of reporting problems using condoms, and depression was associated with 50% lower odds of using condoms consistently.⁴⁵ A longitudinal study of 155 adolescents reported that perceived stress was significantly associated with emotional distress, which in turn, was associated with a higher number of sexual partners and high risk sexual partners.⁴⁶ Perceived stress may also influence STI risk through changes in partners: a study found that

among 1,002 women in South Africa, baseline depression was marginally associated ($p=.073$) with increased sexual partner concurrence in the following 12 months.⁴⁷

There are limited longitudinal data examining links between psychosocial stressors and STI.^{20, 48–50} In one prospective study of 627 African-American female adolescents (14–20 years), chronic socioeconomic strain was associated with STI acquisition and reinfection over 36 months.²⁰ In a 14 year study of nearly one million women in the Swedish Multi-Generation Register, death of a close relative was associated with incident STI.⁵⁰ In a study of 130 heterosexual dyads (ages 14–24 years) with discordant chlamydia status, depressive symptoms were associated with increased STI risk in the non-infected partner.⁵¹

The objective of this study was to evaluate the associations between psychosocial stress and incident STIs (genital CT, GC TV), testing BV and sexual behaviors as mediators. To our knowledge, no prior studies have examined potential behavioral and microbial mediators of the association between perceived stress and incident STIs.

Materials and Methods

Sample

From August 1999 to February 2002, 3,620 non-pregnant women (ages 15 to 44 years) were recruited to the LSVF through family planning and wellness clinics in the Birmingham, Alabama area.⁵² Exclusion criteria were immunocompromised status, hysterectomy, menopause, pelvic radiotherapy, antibiotic therapy greater than 30 days, non-fluency in English, plans to move within the next 12 months, participation in a clinical trial using antibiotics or genital microbicides, and mental/intellectual limitations preventing informed consent. Participants were followed for approximately 12 months with clinical exams and surveys. Participants who had a positive test for genital GC, CT or TV at baseline (screening described below) were excluded from our analytic sample (752 excluded). Next, we excluded participants with no visits after baseline (413 excluded). Of the remaining sample there was less than 1% missing data for all baseline measures other than income, which was imputed. All observations with incomplete baseline measures other than income were excluded (16 excluded). The final sample consisted of 2,439 participants (32.3% excluded).

All participants gave written informed consent, and the study was approved by the Institutional Review Boards of the National Institute of Child Health and Human Development, the Jefferson County Department of Health, the University of Alabama at Birmingham, as well as secondary data analysis at the University of Maryland School of Medicine.

Measures

Perceived Stress.—Perceived stress was measured at baseline using Cohen's 10-item Perceived Stress Scale (PSS), a well-validated measure of an individual's appraisal of stress in the past 30 days.^{53–55} Each of the items have a 5-point Likert response format, and the scale covers dimensions of overload, uncontrollability, and ability to cope, including questions such as "In the last month, how often have you been upset because of something

that happened unexpectedly?”. This scale demonstrated good internal consistency (Cronbach’s $\alpha = 0.85$). We used this measure as a continuous score in our main analyses.

Incident STI.—Participants were tested for GC, CT, and TV at each study visit. An endocervical swab was used to inoculate Thayer-Martin agar plates for culture of GC, while CT testing was performed by ligase chain reaction (Abbott Laboratories, Abbott Park, Illinois). TV was determined by means of a positive finding upon either culture (In-Pouch) or microscopic evaluation for trichomonads. Participants were removed from the risk set after their first observed incident STI, as there were very few cases with multiple distinct STIs during the study. For participants without an observed incident STI, their last study visit was used as the right-censoring time.

Bacterial Vaginosis.—BV was measured at each visit using Nugent’s scoring of a vaginal Gram stain (0 to 10).⁵⁶ Nugent score was categorized as low, intermediate, or high (0–3, 4–6, 7–10 respectively) based on standard scoring criteria. Time-varying measures for Nugent category were used up to the time point of first incident STI or right censoring. For each time point, the Nugent category at the previous wave was used in order to assess BV as a mediator of incident STI. This ensured that Nugent scores reflect the period after the end of the baseline stress period and before incident STI or right censoring. This also makes the Nugent Score concurrent with the beginning of the reference period for each sexual behavior, which extends three months prior to the time of measurement.

Sexual Risk Behaviors—Sexual risk behaviors were measured at each visit. Respondents reported on their number of sexual partners in the past 3 months (0 to 1, 2 or more), frequency of condom use (never, seldom, half the time, most of the time, always), and sexual concurrence of their partners (none, unlikely concurrence, possible concurrence, definite concurrence). Partner concurrence was assessed with the question “Do you think any of these sexual partners had sex with anyone else?” We used time-varying measures for all sexual behaviors up to first incident STI or right censoring.

Other Risk Factors for STIs.—We used potential confounders measured at the baseline interview, including self-identified race (Black, White, other), education (high school and under, above high school), marital status (married, not married), monthly income (less than \$500, \$500 to \$800, \$800 to \$3000, greater than \$3000), and age (continuous years). A literature-based approach was used to select confounders that have been shown in previous research to confound the association between stress and STI or have been independently associated with both stress and incident STI.^{19, 45, 46, 50}

Statistical Analysis

First, we compared baseline demographics, BV, and sexual behaviors between participants with and without incident STI using chi-square tests for categorical variables, and Satterthwaite t-tests for continuous variables. We used Cochran-Armitage tests of trend to test differences in proportions of ordinal variables. We also compared the theorized mediators across perceived stress. We also tested differences in survival rates between perceived stress quartiles (11, 12–17, 18–23, 24) using log-rank tests for trend. To assess

linearity of the PSS, we also analyzed the quartile-categorized PSS, using both adjusted and unadjusted Cox proportional hazards models. Categories of 0 and 1 for number of sexual partners were collapsed to account for potential unreported partner overlap for participants reporting 0 and 1 partners at adjacent visits. Adjacent categories of 0 and 1 was a commonly observed phenomenon in our data. To assess the validity of combining these categories, we also tested for differences between these categories in a post-hoc analysis.

Cox proportional hazards models were used to test the association between perceived stress and incident STIs. We used sequential model-building, where the first model only included the perceived stress term (Model 1). We then added terms for potential demographic confounders (Model 2). We then added time-varying terms for Nugent score and sexual behaviors (Model 3). The Kolmogorov-type supremum test was used to assess the proportional hazards assumption for all models.

Next, we tested sexual risk behaviors and Nugent score as mediators by calculating indirect association estimates for these variables both separately and combined. The indirect association estimates, adjusted for the same covariates included in Model 2, describe the proportion of the association between perceived stress and incident STI that is explained by the set of mediators. Vanderweele's difference method was used to estimate indirect associations of Nugent score, sexual behaviors, and both mediators combined by calculating the change in the association estimate when including each potential mediator in the model.⁵⁷ To calculate each indirect association, the beta estimate for the PSS term in each adjusted model containing mediators was subtracted from the beta estimate for the PSS term in the total adjusted model. Each percentage of mediated association was calculated from this. We used bootstrapping to generate 95% confidence intervals for each indirect association. All analyses were conducted using SAS Version 9.4 (SAS Institute, Cary, NC).

Missing data.—There was less than 5% missingness for all time-varying measures, and less than 1% missing data for all baseline measures except income, which had 8% missingness. To address this, monotone multiple imputation was used to generate 10 imputed datasets that were used for all model analyses. We imputed income using non-missing data from race, age, education level, and marital status. These variables were selected as they demonstrated significant associations with income in our dataset.

Results

Sample

The 2,439 participants contributed a total 1855.55 person-years. The person-time contributed by each participant ranged from 48 to 405 days. The PSS score ranged from 0 to 40 and was normally distributed (Supplement 1). In both unadjusted and adjusted models using the quartile-stratified PSS, we found clear evidence of dose-response, consistent with linearity (Supplement 2). Perceived stress was relatively stable over the follow-up period: 85% of participants remained within one stress quartile of their baseline stress quartile across all visits, with strong positive correlation between perceived stress measures across visits (Median $r=.62$; see Appendix 1). Women in our study were predominantly Black/African-American (78%) and unmarried (82%); most women had more than a high school

education (71%) and 72% had monthly income between \$500 and \$3000. There were 675 (27% of the sample) incident STI cases in the cohort across 1855.55 person-years, equating to an incidence rate of 36.4 cases per 100 person-years. Among incident STI cases, 46.4% were Chlamydia only, 35.7% were Trichomonas only, 10.2% were GC only, 3.0% were GC and Chlamydia, 1.3% were GC and Trichomonas, 3.1% were Chlamydia and Trichomonas, and 0.3% were all three infections.

Bivariate

Greater perceived stress was associated with incident STI ($p < .001$) (Table 1). Those with incident STI were more likely to be younger, Black, unmarried, and have elevated Nugent score, lower income, education of high school or less, sexual partner concurrence, multiple sexual partners, and frequent condom use. Additionally, greater perceived stress was associated with elevated Nugent score, multiple sexual partners, and sexual partner concurrence (Table 2). A Kaplan-Meier plot (Figure 1) demonstrates differences in survival between the four perceived stress quartiles. As shown, the lowest quartile of stress has lower risk of STI than any of the other quartiles. A log-rank test for trend indicated a significantly increasing risk of STI with increasing stress quartiles ($p < .001$).

Cox Proportional Hazards Models

Perceived stress was associated with incident STI (Table 3), with an approximate 2% increase in risk of incident STI with each 1 unit increase in perceived stress. At the highest values of perceived stress (40), this equates to nearly twice the risk of incident STI compared to those at the lowest values of perceived stress (0). When adjusting for confounders in Model 2, this was attenuated to approximately a 1.5% increase in risk of incident STI with each 1 unit increase in perceived stress. The association between perceived stress and incident STIs was further attenuated after incorporating potential mediators and was no longer statistically significant. Among mediators, intermediate Nugent score, high Nugent score, and partner sexual concurrence were associated with greater risk of STI. Significant demographic factors included race and education of high school or less. The Kolmogorov-type supremum test did not indicate a significant deviation from proportional hazards for any of the terms used ($p > .05$).

Mediation Analyses

Mediation analysis revealed significant indirect effects of Nugent score, sexual behaviors, and both combined (Table 4). Of the total association of stress with incident STI, 21% was mediated through Nugent score, 65% was mediated through sexual behaviors, and combined, Nugent score and sexual behaviors mediated 78% of the observed association (Figure 2).

Post-Hoc Analyses

Using the two-category number of sexual partners variable (0–1, 2+) did not result in any substantial change (<1% difference in indirect associations) compared to models using the three-category number of sexual partners variable (0, 1, 2+). Sexual behaviors were a

significant mediator using both categorizations for number of sexual partners, both in conjunction with and independent of Nugent score.

Discussion

In this longitudinal study of 2,450 reproductive-age women, baseline perceived stress was associated with incident STI over approximately a year, independent of major confounders. This is consistent with cross-sectional studies indicating that stress is associated with higher prevalence of STIs.^{20, 45, 50} Additionally, approximately one-fifth of the association of stress with incident STI was mediated through Nugent score. This finding unites two previous studies from the LSVF study team documenting an association between perceived stress and both development and prevalence of BV, and an association between BV and incident STI.^{27, 30} One-third of the mediating effects of Nugent score was independent of sexual behaviors; this may suggest direct impacts of stress on the dysbiosis of vaginal microbiota. The analysis of mediators allows for a more complete understanding of mechanisms of our observed association. While the analysis excluding mediators is useful for assessing the total effect of stress on incident STI, the model including mediators more clearly parses how this association manifests. This is an especially important approach to risk assessment, as it identifies specific factors to target for intervention. While the confidence intervals for the association between stress and incident STI in model 3 overlapped the null, this only reflects the direct effect of stress, not its total effect. Model 2 reflects the total effect of stress on STI. The substantial decrease in stress estimates in model 3 indicates that sexual risk behaviors and Nugent score are large mediators of this association (i.e., baseline stress affects STI through changes in sexual behaviors and Nugent score).

Chronic psychological stress can be accompanied by dysregulation in the immune system that may directly affect colonization by a STI and indirectly increase STI susceptibility due to changes in the vaginal microenvironment. Chronic psychological stress has been shown to affect the immune system by activating type-2 cytokine-driven and pro-inflammatory responses.²⁴ Chronic stress has been associated with a decrease in natural killer cell cytotoxicity and suppressed lymphocyte proliferative responses.^{58–60} Sympathetic fibers that innervate lymphoid tissues can act directly on lymphocytes via beta adrenergic receptors. Psychological stress may also influence release of corticotropin releasing hormone (CRH) from the hypothalamus. CRH causes release of adrenal corticotropic hormone from the pituitary gland, which induces cortisol secretion from the adrenal cortex. Cortisol can profoundly impair the immune response by blocking T cell proliferation.⁶¹ These changes may result in a decrease in Th1 cytokines, which are required for active cellular immunity to defend against a wide range of infections, and an increase in production of pro-inflammatory Th2 cytokines.²⁴ Inflammation could then cause destruction of infected epithelial layers, allowing bacterial STIs to access deeper tissues.^{62, 63} BV has been associated with genital pro-inflammatory cytokine upregulation, although some studies have found that downregulation of some cytokines also occurs.^{64–66} Shifts in immune function due to psychological stress may directly impact susceptibility to STIs through changes in host immune responses and the microbiota.

Stress may also influence additional behavioral risk factors for STI, including sex while using alcohol or other substances. In a prospective study by Seth et al. of 605 African-American female adolescents, depressive symptoms were associated with greater likelihood of not using condoms, having multiple sexual partners, having a main partner with concurrent sexual partners, a higher fear of discussing condoms with partners, and sex while influenced by alcohol or drugs.⁶⁷ Race demonstrated the strongest association with incident STI, independent of sexual risk behaviors and socioeconomic factors. There are many social and biological factors that converge to cause disproportionately high STI rates among African-Americans.⁶⁷ This association may reflect differences in sexual network dynamics, network STI prevalence, BV rates, and unmeasured individual, psychosocial and structural factors affected by racial discrimination.^{68, 69}

There are limitations to the present study. First, the study was limited to clinics in the Birmingham, Alabama area, and was comprised primarily of Black women, so we could not study modification by race. However, Black women living in rural areas are often neglected in STI literature and face increased risk of STI. A potential limitation is that the data was collected approximately 20 years ago. While specific stressors (e.g. types of discrimination experienced) may vary over time, we are not aware of evidence that the association between perceived stress and STI is likely to differ over time. Our results are consistent with recent studies on the association between stress and prevalent STI.^{19, 26, 45, 50} Culture-based screening methods for GC and TV are not as sensitive as modern methods such as PCR, so there may have been some false negatives. Finally, perceived stress is a subjective proxy for physiologic stress, and it is possible unobserved variables (e.g. recent major adverse events) may lead some individuals to over- or under-report their stress. While reporting of stress may differ between Black and White women, as Black women are likely to face additional stressors related to racism (e.g. additional experiences of discrimination, microaggressions), the PSS is highly validated and has been used in many studies across several populations and disciplines.⁵⁵ It also allows for the measurement of stress over the last 30 days, as opposed to a narrower measurement such as cortisol. Social desirability bias is also likely to be present for self-reported sexual behaviors, potentially resulting in misclassification.^{70, 71} Additionally, while our study covered multiple measures of sexual behaviors, these variables are not fully comprehensive. There are additional sexual risk factors, such as partner STI status, that were unmeasured and could not be included.

There are a number of strengths to this study. The use of Nugent scoring for BV assessment and laboratory-based screening for STIs, as well as the longitudinal study design, allowed us to assess the association between perceived stress and STIs, as well as estimate relative risk. The use of baseline perceived stress and subsequent sexual risk behaviors, Nugent score, and STI is especially important, as there are several potential bidirectional associations among these variables. For instance, bacterial vaginosis may lead to increases in stress, or changes in sexual partners may also lead to later changes in stress. This could result in a mediating pathway where stress mediates the association between these factors and STI. The use of temporally separated measures for baseline stress, intermediate mediators, and incident STI allows for assessment of a single theorized mediating pathway that can be interpreted with observed temporality. Much of the existing research on this topic has utilized small clinic-based samples, and often with predominantly White participants. The use of time-varying

mediators and covariates is a strength, as it allowed the most complete incorporation of the data available between the time of the baseline stress exposure and the time of STI or right censoring. Lastly, Vanderweele's method is particularly robust for assessing mediation, as it allows for testing multiple mediators and can parse their associations independent of each other.

Conclusions

In this large, 12-month prospective cohort study, perceived stress was associated with incident STIs, and sexual risk behaviors and BV appear to be key mechanisms of this association. To our knowledge, this is the first longitudinal study to assess the association between stress and incident STI and demonstrate how changes in Nugent score and sexual risk behaviors mediate the association between stress and incident STIs. This study highlights perceived stress and the intermediary risk factors as important focal points to consider in the development of STI prevention programs. Future research into social and biological mechanisms will be useful in further understanding how stress influences risk for genital infections and developing strategies to intervene.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements:

The authors would like to acknowledge the University of Maryland, College Park, and the University of Maryland, Baltimore, where this research was conducted.

Sources of Funding: This work was supported by grant NIAID R01-AI116799 from the National Institutes of Health, grant NO-1-HD-8-3293 from the National Institute of Child Health and Human Development and intramural fund Z01-HD002535 from the National Institute of Child Health and Human Development.

Appendix 1.

Pearson's r correlations^{***} of Perceived Stress Scale across waves (n=2,439).

	Wave 1	Wave 2	Wave 3	Wave 4	Wave 5
Wave 1	-	0.65	0.58	0.56	0.58
Wave 2	-	-	0.63	0.60	0.59
Wave 3	-	-	-	0.65	0.66
Wave 4	-	-	-	-	0.70
Wave 5	-	-	-	-	-

P-values for all correlations <.001.

List of Abbreviations:

CDC	Centers for Disease Control
GC	Neisseria gonorrhoeae

CT	Chlamydia trachomatis
TV	Trichomonas vaginalis
HIV	Human immunodeficiency virus
BV	Bacterial vaginosis
LSVF	Longitudinal Study of Vaginal Flora

REFERENCES

1. CDC. Sexually Transmitted Disease Surveillance, 2016. Atlanta, GA: U.S. Centers for Disease Control and Prevention, 2017.
2. Satterwhite CL, Torrone E, Meites E, et al. Sexually transmitted infections among US women and men: prevalence and incidence estimates, 2008. *Sex Transm Dis* 2013;40(3):187–93. [PubMed: 23403598]
3. Owusu-Edusei K, Chesson HW, Gift TL, et al. The estimated direct medical cost of selected sexually transmitted infections in the United States, 2008. *Sex Transm Dis* 2013;40(3):197–201. [PubMed: 23403600]
4. Mania-Pramanik J, Kerkar S, Sonawane S, et al. Current Chlamydia trachomatis Infection, A Major Cause of Infertility. *J Reprod Infertil* 2012;13(4):204–10. [PubMed: 23926547]
5. UNAIDS/WHO. Sexually transmitted diseases: policies and principles for prevention care. Geneva: UNAIDS Best Practice Collection Key Material, 1999.
6. Datta SD, Sternberg M, Johnson RE, et al. Gonorrhea and chlamydia in the United States among persons 14 to 39 years of age, 1999 to 2002. *AnnInternMed* 2007;147(2):89–96.
7. Miller WC, Ford CA, Morris M, et al. Prevalence of chlamydial and gonococcal infections among young adults in the United States. *JAMA* 2004;291 (18):2229–36. [PubMed: 15138245]
8. Centers for Disease Control and Prevention. Sexually Transmitted Disease Surveillance 2014. Atlanta: U.S. : Department of Health and Human Services., 2015.
9. Tanfer K, Cubbins LA, Billy JO. Gender, race, class and self-reported sexually transmitted disease incidence. *Fam Plann Perspect* 1995;27(5):196–202. [PubMed: 9104606]
10. Miller HG, Cain VS, Rogers SM, et al. Correlates of sexually transmitted bacterial infections among U.S. women in 1995. *Fam Plann Perspect* 1999;31 (1):4–9–23. [PubMed: 10029926]
11. Harawa NT, Greenland S, Cochran SD, et al. Do differences in relationship and partner attributes explain disparities in sexually transmitted disease among young white and black women? *J Adolesc Health* 2003;32(3):187–91. [PubMed: 12606112]
12. Ellen JM, Aral SO, Madger LS. Do differences in sexual behaviors account for the racial/ethnic differences in adolescents' self-reported history of a sexually transmitted disease? *Sex Transm Dis* 1998;25(3):125–9. [PubMed: 9524987]
13. Harawa NT, Greenland S, Bingham TA, et al. Associations of race/ethnicity with HIV prevalence and HIV-related behaviors among young men who have sex with men in 7 urban centers in the United States. *J Acquir Immune Defic Syndr* 2004;35(5):526–36. [PubMed: 15021318]
14. Rice RJ, Roberts PL, Handsfield HH, et al. Sociodemographic distribution of gonorrhea incidence: implications for prevention and behavioral research. *Am J Public Health* 1991;81(10):1252–8. [PubMed: 1928521]
15. Ellen JM, Kohn RP, Bolan GA, et al. Socioeconomic differences in sexually transmitted disease rates among black and white adolescents, San Francisco, 1990 to 1992. *Am J Public Health* 1995;85(11):1546–8. [PubMed: 7485669]
16. Cohen S, Janicki-Deverts D, Miller GE. Psychological stress and disease. *Jama-Journal of the American Medical Association* 2007;298(14):1685–7.
17. McEwen B, Stellar E. Stress and the individual: Mechanisms leading to disease. *Archives of Internal Medicine* 1993; 153(18):2093–101. [PubMed: 8379800]

18. McEwen BS. Physiology and neurobiology of stress and adaptation: Central role of the brain. *Physiological Reviews* 2007;87(3):873–904. [PubMed: 17615391]
19. Hulland EN, Brown JL, Swartzendruber AL, et al. The association between stress, coping, and sexual risk behaviors over 24-months among African American female adolescents. *Psychology, health & medicine* 2015;20(4):443–56.
20. Sales JM, Smearman EL, Swartzendruber A, et al. Socioeconomic-related Risk and STI Infection among African-American Adolescent Females. *J Adolesc Health* 2014;55(5):698–704. [PubMed: 24974317]
21. Cohen S, Alper CM, Doyle WJ, et al. Positive Emotional Style Predicts Resistance to Illness After Experimental Exposure to Rhinovirus or Influenza A Virus. *Psychosomatic Medicine* 2006;68(6): 809–15. [PubMed: 17101814]
22. Cohen S, Doyle WJ, Turner R, et al. Sociability and susceptibility to the common cold. *Psychological Science (Wiley-Blackwell)* 2003; 14(5):389.
23. Cohen S, Doyle WJ, Turner RB, et al. Childhood Socioeconomic Status and Host Resistance to Infectious Illness in Adulthood. *Psychosomatic Medicine* 2004;66(4):553–8. [PubMed: 15272102]
24. Cohen S, Janicki-Deverts D, Doyle WJ, et al. Chronic stress, glucocorticoid receptor resistance, inflammation, and disease risk. *Proceedings of the National Academy of Sciences* 2012;109(16): 5995–9.
25. Culhane JF, Rauh V, McCollum KF, et al. Maternal stress is associated with bacterial vaginosis in human pregnancy. *Maternal and child health journal* 2001;5(2):127–34. [PubMed: 11573838]
26. Culhane JF, Rauh VA, Goldenberg RL. Stress, bacterial vaginosis, and the role of immune processes. *Current infectious disease reports* 2006;8(6):459–64. [PubMed: 17064639]
27. Nansel TR, Riggs MA, Yu K-F, et al. The association of psychosocial stress and bacterial vaginosis in a longitudinal cohort. *American Journal of Obstetrics and Gynecology* 2006;194(2):381–6. [PubMed: 16458633]
28. Amabebe E, Anumba DOC. Psychosocial Stress, Cortisol Levels, and Maintenance of Vaginal Health. *Front Endocrinol (Lausanne)* 2018;9:568. [PubMed: 30319548]
29. Peipert JF, Lapane KL, Allsworth JE, et al. Bacterial vaginosis, race, and sexually transmitted infections: does race modify the association? *Sex TransmDis* 2008;35(4):363–7.
30. Brotman RM, Klebanoff MA, Nansel TR, et al. Bacterial vaginosis assessed by gram stain and diminished colonization resistance to incident gonococcal, chlamydial, and trichomonal genital infection. *J Infect Dis* 2010;202(12):1907–15. [PubMed: 21067371]
31. Cherpes TL, Meyn LA, Krohn MA, et al. Association between acquisition of herpes simplex virus type 2 in women and bacterial vaginosis. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America* 2003;37(3):319–25. [PubMed: 12884154]
32. Martin HL, Richardson BA, Nyange PM, et al. Vaginal lactobacilli, microbial flora, and risk of human immunodeficiency virus type 1 and sexually transmitted disease acquisition. *J Infect Dis* 1999;180(6):1863–8.
33. King CC, Jamieson DJ, Wiener J, et al. Bacterial vaginosis and the natural history of human papillomavirus. *Infect Dis Obstet Gynecol* 2011;2011:319460. [PubMed: 21869857]
34. Myer L, Denny L, Telerant R, et al. Bacterial vaginosis and susceptibility to HIV infection in South African women: a nested case-control study. *The Journal of infectious diseases* 2005;192(8):1372–80. [PubMed: 16170754]
35. Rea McClelland. Evaluation of the association between the concentrations of key vaginal bacteria and the increased risk of HIV acquisition in African women from five cohorts: a nested case-control study. *Lancet Infect Dis* 2018.
36. Boskey ER, Cone RA, Whaley KJ, et al. Origins of vaginal acidity: high D/L lactate ratio is consistent with bacteria being the primary source. *Human reproduction (Oxford, England)* 2001;16(9):1809–13.
37. Aroutcheva A, Gariti D, Simon M, et al. Defense factors of vaginal lactobacilli. *Am J Obstet Gynecol* 2001;185(2):375–9. [PubMed: 11518895]
38. Martin H, Richardson B, Nyange P, et al. Vaginal lactobacilli, microbial flora, and risk of human immunodeficiency virus type 1 and sexually transmitted disease acquisition. *J Infect Dis* 1999;180:1863–8. [PubMed: 10558942]

39. Myer L, Kuhn L, Stein ZA, et al. Intravaginal practices, bacterial vaginosis, and women's susceptibility to HIV infection: epidemiological evidence and biological mechanisms. *Lancet InfectDis* 2005;5(12):786–94.
40. Ness RB, Kip KE, Soper dE, et al. Bacterial vaginosis (BV) and the risk of incident gonococcal or chlamydial genital infection in a predominantly black population. *Sex TransmDis* 2005;32(7):413–7.
41. Ocana VS, Pesce De Ruiz Holgado AA, Nader-Macias ME. Characterization of a bacteriocin-like substance produced by a vaginal *Lactobacillus salivarius* strain. *Applied and Environmental Microbiology* 1999;65(12):5631–5. [PubMed: 10584033]
42. Reid G, Heinemann C, Velraeds M, et al. Biosurfactants produced by *Lactobacillus*. *Methods in enzymology* 1999;310:426–33. [PubMed: 10547809]
43. Boris S, Barbés C. Role played by lactobacilli in controlling the population of vaginal pathogens. *Microbes Infect* 2000;2(5):543–6. [PubMed: 10865199]
44. McMillan A, Dell M, Zellar MP, et al. Disruption of urogenital biofilms by lactobacilli. *Colloids Surf B Biointerfaces* 2011;86(1):58–64. [PubMed: 21497071]
45. Mazzaferro K, Murray P, Ness R, et al. Depression, Stress, and Social Support as Predictors of High-Risk Sexual Behaviors and STIs in Young Women. *J Adolesc Health* 2006;39(4):601–3. [PubMed: 16982400]
46. Ethier K, Kershaw T, Lewis J, et al. Self-esteem, emotional distress and sexual behavior among adolescent females: Inter-relationships and temporal effects. *J Adolesc Health* 2006;38(3):268–74. [PubMed: 16488825]
47. Nduna M, Jewkes R, Dunkle K, et al. Associations between depressive symptoms, sexual behaviour and relationship characteristics: a prospective cohort study of young women and men in the Eastern Cape, South Africa. *J Int AIDS Society* 2010;13(44).
48. Leserman J Role of depression, stress, and trauma in HIV disease progression. *Psychosom Med* 2008;70(5):539–45. [PubMed: 18519880]
49. Wickrama T, Merten MJ, Wickrama KAS. Early Socioeconomic Disadvantage and Young Adult Sexual Health. *American Journal of Health Behavior* 2012;36(6):834–48. [PubMed: 23026041]
50. Bond E, Lu D, Herweijer E, et al. Sexually transmitted infections after bereavement - a population-based cohort study. *BMC Infect Dis* 2016;16(1):419. [PubMed: 27528204]
51. Shrier LA, Schillinger JA, Aneja P, et al. Depressive symptoms and sexual risk behavior in young, chlamydia-infected, heterosexual dyads. *J Adolesc Health* 2009;45(1):63–9. [PubMed: 19541251]
52. Klebanoff MA, Schwabke JR, Zhang J, et al. Vulvovaginal symptoms in women with bacterial vaginosis. *Obstet Gynecol* 2004;104(2):267–72. [PubMed: 15291998]
53. Sharp LK, Kimmel LG, Kee R, et al. Assessing the Perceived Stress Scale for African American adults with asthma and low literacy. *J Asthma* 2007;44(4):311–6. [PubMed: 17530531]
54. Cole SR. Assessment of differential item functioning in the Perceived Stress Scale-10. *J Epidemiol Community Health* 1999;53(5):319–20. [PubMed: 10396541]
55. Taylor JM. Psychometric analysis of the Ten-Item Perceived Stress Scale. *Psychol Assess* 2015;27(1):90–101. [PubMed: 25346996]
56. Nugent RP, Krohn MA, Hillier SL. Reliability of diagnosing bacterial vaginosis is improved by a standardized method of gram stain interpretation. *J Clin Microbiol* 1991;29(2):297–301. [PubMed: 1706728]
57. VanderWeele TJ. Explanation in causal inference: developments in mediation and interaction. *Int J Epidemiol* 2016;45(6):1904–8. [PubMed: 27864406]
58. Cohen S, Miller GE, Rabin BS. Psychological stress and antibody response to immunization: a critical review of the human literature. *Psychosomatic medicine* 2001;63(1):7–18. [PubMed: 11211068]
59. Herbert TB, Cohen S. Stress and immunity in humans: a meta-analytic review. *Psychosom Med* 1993;55(4):364–79. [PubMed: 8416086]
60. Kiecolt-Glaser JK, the RGPo. Chronic stress alters the immune response to influenza virus vaccine in older adults. *National Acad Sciences*.

61. Rook GA. Glucocorticoids and immune function. *Baillieres Best Pract Res Clin Endocrinol Metab* 1999; 13(4):567–81. [PubMed: 10903815]
62. Svanborg C, Godaly G, Hedlund M. Cytokine responses during mucosal infections: role in disease pathogenesis and host defence. *Current opinion in microbiology* 1999;2(1):99–105. [PubMed: 10047563]
63. McGee ZA, Jensen RL, Clemens CM, et al. Gonococcal infection of human fallopian tube mucosa in organ culture: relationship of mucosal tissue TNF-alpha concentration to sloughing of ciliated cells. *Sex Transm Dis* 1999;26(3): 160–5. [PubMed: 10100774]
64. Yudin MH, Landers DV, Meyn L, et al. Clinical and cervical cytokine response to treatment with oral or vaginal metronidazole for bacterial vaginosis during pregnancy: a randomized trial. *Obstetrics and gynecology* 2003;102(3):527–34. [PubMed: 12962937]
65. Sturm-Ramirez K, Gaye-Diallo A, Eisen G, et al. High levels of tumor necrosis factor- alpha and interleukin-1 beta in bacterial vaginosis may increase susceptibility to human immunodeficiency virus. *The Journal of infectious diseases* 2000;182(2):467–73. [PubMed: 10915077]
66. Ryckman KK, Williams SM, Krohn MA, et al. Racial differences in cervical cytokine concentrations between pregnant women with and without bacterial vaginosis. *Journal of reproductive immunology* 2008;78(2):166–71. [PubMed: 18336917]
67. Seth P, Patel S, Sales J, et al. The impact of depressive symptomatology on risky sexual behavior and sexual communication among African American female adolescents. *Psych Hlth Med* 2011;16(3):346–56.
68. Kaplan KC, Holmes JM, Wallace M, et al. Racial Discrimination and HIV-related Risk Behaviors in Southeast Louisiana. *Am J Health Behav* 2016;40(1): 132–43. [PubMed: 26685822]
69. Prather C, Fuller TR, Marshall KJ, et al. The Impact of Racism on the Sexual and Reproductive Health of African American Women. *J Womens Health (Larchmt)* 2016;25(7):664–71. [PubMed: 27227533]
70. Zenilman JM, Weisman CS, Rompalo AM, et al. Condom use to prevent incident STDs: the validity of self-reported condom use. *Sex Transm Dis* 1995;22(1):15–21. [PubMed: 7709320]
71. Liu H, Morisky DE, Lin X, et al. Bias in Self-Reported Condom Use: Association Between Over-Reported Condom Use and Syphilis in a Three-Site Study in China. *AIDS Behav* 2016;20(6): 1343–52. [PubMed: 26696262]

Highlights

- Base perceived stress was associated with incident sexually transmitted infections.
- This association persisted independent of demographic and socioeconomic factors.
- Bacterial vaginosis partially mediated this association.
- Condom use, number of partners, and partner concurrence were also mediators.
- Mediation modeling method was used for causal relationship analysis.

Figure 1: Kaplan-Meier plot of survival probability (of no incident STI) over time in days stratified by baseline stress quartile (n=2,439, *** $P<.001$).

Figure 2:
Directed acyclic graph of the association of perceived stress with incident STI and mediating paths, adjusted for age, race, marital status, education, and income (n=2,439).

Table 1.

Baseline demographic and behavioral characteristics and Nugent score prior to event, stratified by incident STI status (n=2,439).

	Total		No Observed Incident STI		Observed Incident STI		P-value
	Mean	SD	Mean	SD	Mean	SD	
Perceived Stress Scale, past 3 months¹	17.12	7.5	16.77	7.6	18.02	7.0	<.001
Age (years)¹	25.70	7.1	26.05	7.1	24.76	7.0	<.001
	n	%	n	%	n	%	
Marital Status²							<.001
Not Married	1998	81.9	1395	79.1	603	89.3	
Married	441	18.1	369	20.9	72	10.7	
Race²							<.001
Black	1908	78.2	1290	73.1	618	91.6	
White	495	20.3	443	25.1	52	7.7	
Other	36	1.5	31	1.8	5	0.7	
Education Level²							<.001
High School	1739	71.3	1203	68.2	536	79.4	
>High School	700	28.7	561	31.8	139	20.6	
Monthly Income³							
< \$500	447	19.9	301	18.4	146	23.9	<.001
\$500 to \$800	659	29.3	440	26.9	219	35.8	
\$800 to \$3000	962	42.8	746	45.7	216	35.3	
> \$3000	178	7.9	147	9.0	31	5.1	
Nugent Score^{3,4}							
Low (0 to 3)	982	40.3	792	44.9	190	28.3	<.001
Medium (4 to 6)	515	21.4	362	20.6	153	22.8	
High (7 to 10)	939	38.6	610	34.5	329	49.0	
Condom Use, past 3 months^{3,4}							<.001
Never	1531	63.0	1165	66.3	366	54.5	
Seldom	112	4.6	77	4.4	35	5.2	
Half the time	85	3.5	56	3.2	29	4.3	
Most of the time	201	8.3	141	8.0	60	8.9	
Always	501	20.6	319	18.2	182	27.1	
Number of Sex Partners, past 3 months^{2,4}							<.001
0–1	2309	94.7	1690	95.6	619	91.7	
2	130	5.3	74	4.2	56	8.3	
Partner Concurrence^{3,4}							<.001

	Total		No Observed Incident STI		Observed Incident STI		P-value
	Mean	SD	Mean	SD	Mean	SD	
No concurrence	725	33.4	574	36.6	153	25.8	
Unlikely concurrence	1121	51.6	805	51.3	307	51.7	
Possibly concurrent	171	7.9	104	6.6	64	10.8	
Definitely concurrent	154	7.1	85	5.4	70	11.8	

¹ P-value calculated using Satterwaithe t-test;

² P-value calculated using Chi-square test;

³ P-value calculated using Cochran-Armitage trend test.

⁴ Nugent score is used from wave prior to observed incident STI or right censoring. Condom use, number of sexual partners, and sexual partner concurrence are used at wave of incident STI or right censoring. All other variables are used from baseline.

Table 2.

Mean baseline perceived stress scale across behavioral characteristics and Nugent score prior to event (n=2,439).

	Perceived Stress Scale, past 3 months		
	Mean	SD	P-value
Nugent Score ^{1,3}			.031
Low (0 to 3)	16.78	7.7	
Intermediate (4 to 6)	17.17	7.3	
High (7 to 10)	17.46	7.3	
Condom Use ^{1,3}			.165
Never	18.89	7.6	
Seldom	18.61	7.3	
Half the time	18.12	7.7	
Most of the time	17.49	7.3	
Always	17.13	7.3	
Number of Sex Partners, past 3 months ^{2,3}			<.001
0–1	16.93	7.4	
2	20.31	7.5	
Partner Concurrence ^{1,3}			<.001
No concurrence	16.49	7.4	
Unlikely concurrence	17.17	7.3	
Possibly concurrent	18.73	7.3	
Definitely concurrent	18.99	7.9	

¹ P-value calculated using Spearman's rank-sum correlation;

² P-value calculated using Satterwaithe t-test

³ Nugent score is used from wave prior to observed incident STI or right censoring. Condom use, number of sexual partners, and sexual partner concurrence are used at wave of incident STI or right censoring.

Table 3.

Hazard ratios (and 95% confidence intervals) for the association of perceived stress and incident STI (n=2,439).

	Model 1: Unadjusted Total Stress Association	Model 2: Adjusted Total Stress Association	Model 3: Adjusted Direct Stress Association
Perceived Stress Scale, past 3 months			
Continuous (0 to 40)	1.019 (1.009, 1.029)	1.015 (1.005, 1.026)	1.003 (0.990, 1.016)
Education			
> high school degree		1.00	1.00
≤High school degree		1.456 (1.195, 1.773)	1.460 (1.153, 1.848)
Monthly Income			
< \$500		1.00	1.00
\$500 to \$800		1.021 (0.844, 1.235)	0.920 (0.724, 1.169)
\$800 to \$3000		0.795 (0.635, 0.968)	0.730 (0.572, 0.931)
> \$3000		0.637 (0.435, 0.934)	0.778 (0.507, 1.195)
Race *			
Black		1.00	1.00
White		0.354 (0.266, 0.473)	0.416 (0.295, 0.585)
Age (years)		0.990 (0.979, 1.002)	0.997 (0.982, 1.012)
Marital status			
Not Married		1.00	1.00
Married		0.745 (0.576, 0.964)	0.795 (0.584, 1.073)
Nugent Score			
Low (0 to 3)			1.00
Intermediate (4 to 6)			1.595 (1.230, 2.068)
High (7 to 10)			1.612 (1.280, 2.029)
Condom Use, past 3 months			
Always			1.00
Most of the time			0.996 (0.712, 1.393)
Half the time			1.443 (0.969, 2.148)
Seldom			1.138 (0.747, 1.735)
Never			0.949 (0.748, 1.202)
Number of Sex Partners, past 3 months			
0–1			1.00
2			1.295 (0.900, 1.865)
Partner Concurrence			
No concurrence			1.00
Unlikely concurrence			1.490 (1.186, 1.872)
Possibly concurrent			1.709 (1.176, 2.483)

	Model 1: Unadjusted Total Stress Association	Model 2: Adjusted Total Stress Association	Model 3: Adjusted Direct Stress Association
Definitely concurrent			1.911 (1.325, 2.756)

* Terms for other racial groups not reported due to insufficient sample size in each group (n<20).

Note: Bold indicates statistically significant hazard ratio at $P<.05$.

Table 4.

Estimates (and bootstrap-generated two-sided 95% confidence intervals) for mediation effects of the association between perceived stress and incident STI: Longitudinal Study of Vaginal Flora, 1999–2002, among n=2,439 women STI-free at baseline.

	Model: Stress and only Confounders ¹	Model: Stress, Confounders ¹ , Nugent Score, and Sexual Behaviors ²	Model: Stress, Confounders ¹ , and Nugent Score	Model: Stress, Confounders ¹ , and Sexual Behaviors ²
Stress β	0.0150	0.0033	0.0121	0.0053
Indirect Effect	-	0.0117 (0.0091, 0.0156)	0.0029 (0.0014, 0.0048)	0.0097 (0.0065, 0.0134)
Percent Mediated	-	78%	21%	65%
p-value	-	<0.01	0.03	<0.01

Note: Indirect effects are the difference in estimates compared to the model with Stress and only confounders. Bold indicates statistically significant estimate at $P < .05$.

¹Confounders include age, race, income, education level, and marital status.

²Sexual behaviors include condom use, number of sexual partners, and sexual partner concurrence.