Published in final edited form as:

Am J Cardiol. 2019 May 01; 123(9): 1532–1537. doi:10.1016/j.amjcard.2019.02.006.

Cardiovascular Outcomes of Patients with Pulmonary Hypertension Undergoing Non-Cardiac Surgery

Nathaniel R. Smilowitz, MD^{a,b}, Andrew Armanious, MD^a, Sripal Bangalore, MD, MHA^a, Harish Ramakrishna, MD^{c,*}, and Jeffrey S. Berger, MD, MS^{a,d,*}

^aLeon H. Charney Division of Cardiology, Department of Medicine, New York University School of Medicine, New York, NY

^bDivision of Cardiology, Department of Medicine, Veterans Affairs New York Harbor Health Care System, New York, NY

^cDepartment of Anesthesiology, Mayo Clinic Arizona, Phoenix, AZ

^dDepartment of Surgery, New York University School of Medicine, New York, NY

Abstract

Pulmonary hypertension (PH), defined by a mean pulmonary artery pressure of >25mm Hg at rest, is strongly associated with morbidity and mortality in the perioperative period. The prevalence and outcomes of PH among patients referred for major non-cardiac surgery in the United States are unknown. Patients 18 years of age hospitalized for non-cardiac surgery were identified from Healthcare Cost and Utilization Project's (HCUP) National Inpatient Sample (NIS) data from 2004 to 2014. Pulmonary hypertension was defined by ICD-9 diagnosis codes. The primary outcome was perioperative major adverse cardiovascular events (MACCE), defined as in-hospital death, myocardial infarction (MI), or ischemic stroke. Among 17,853,194 hospitalizations for major non-cardiac surgery, 143,846 (0.81%) had PH. MACCE occurred in 8.3% of hospitalizations with any diagnosis of PH in comparison to 2.0% of those without PH ($p<0.001$), driven by an increased frequency of death (4.4% vs. 1.1%, $p<0.001$) and non-fatal MI (3.2% vs. 0.6%, $p<0.001$). After adjusting for demographics, clinical covariates, and surgery type, PH remained independently associated with MACCE (aOR 1.43, 95% CI 1.40 – 1.46). In conclusion, PH is associated with perioperative major adverse cardiovascular events. Careful patient selection, recognition of perioperative risks, and appropriate intraoperative hemodynamic monitoring may improve perioperative cardiovascular outcomes.

Corresponding Authors: Nathaniel R. Smilowitz, MD, Leon H. Charney Division of Cardiology, Department of Medicine, New York University School of Medicine, 530 First Avenue, HCC-14, New York, NY 10016, nathaniel.smilowitz@nyumc.org, Ph. 212-263-5656, Fax. 212-263-8534.

*Contributed equally as senior authors.

Disclosures: The investigators have no conflicts of interest to disclose.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Keywords

Acute Coronary Syndrome; Myocardial Infarction; Non-cardiac Surgery; Perioperative; Pulmonary Hypertension; Surgery

Introduction

Pulmonary hypertension (PH), defined as a mean pulmonary artery pressure ≥ 25 mmHg at rest,¹ can be identified by echocardiography in $\sim 0.7\%$ of patients referred for non-cardiac surgery.² Patients with PH are at an increased risk of adverse events in the perioperative period,^{3–5} with cardiovascular risks that may vary based on the severity of the hemodynamic abnormality, underlying mechanism of disease, the presence of cardiovascular risk factors and comorbidities, and the type of non-cardiac surgery to be performed. Unfortunately, few studies have identified large cohorts of patients with PH undergoing non-cardiac surgery in the contemporary era. In the present study, we sought to evaluate the incidence of perioperative adverse cardiovascular events after non-cardiac surgery in patients with and without PH using a large administrative database of hospital admissions from the United States.

Methods

Patients ≥ 18 years of age undergoing non-cardiac surgery from 2004 to 2014 were identified from the Healthcare Cost and Utilization Project's (HCUP) National Inpatient Sample (NIS).⁶ The NIS is a national administrative database that contains discharge-level data for a 20% stratified sample of patients admitted to United States hospitals.⁷ Patients were eligible for inclusion if the principal *International Classification of Diseases, Ninth Revision* (ICD-9) procedure code during hospital admission was a major therapeutic operating room procedure (HCUP Procedure Class 4).⁶ Patients who underwent cardiac transplantation, bone marrow transplantation, cardiac surgery, minimally invasive cardiac procedures, ophthalmologic surgery, dental surgery, radiation therapy, and non-operating room procedures were excluded from all analyses. Pulmonary hypertension was identified by the ICD-9 diagnosis codes for primary pulmonary arterial hypertension (416.0) and all other non-primary PH (416.8, 416.9), as previously defined.^{8,9}

The primary outcome was MACCE, defined as in-hospital death, myocardial infarction (MI), or ischemic stroke. Perioperative MI was identified using ICD-9 diagnosis codes for acute ST-segment elevation myocardial infarction (STEMI) (410.01 to 410.61, 410.81, and 410.91) and non-ST-segment elevation myocardial infarction (NSTEMI) (410.71).¹⁰ Acute ischemic stroke was identified using ICD-9 diagnosis codes 433.x1, 434.x1, 436, and 437.1.13. Other cardiovascular outcomes evaluated were acute pulmonary embolism (415.1x), cardiogenic shock (ICD-9 diagnosis code 785.51) and cardiac arrest (ICD-9 diagnosis code 427.5).

Categorical variables were reported as percentages and compared by Chi-square tests. Continuous variables were reported as mean $\pm$ standard deviation (SD) and compared using the Student's *t* test. Multivariable logistic regression models were generated to estimate odds

ratios for MACCE and its individual endpoints, adjusted for age, sex, race/ethnicity, obesity, tobacco use, hypertension, hyperlipidemia, diabetes mellitus, end stage renal disease, coronary artery disease, prior revascularization with either PCI or CABG, peripheral vascular disease, valvular heart disease, prior venous thromboembolism, chronic lung disease, malignancy, anemia, alcohol abuse, elective/urgent hospitalization, surgery type, and year of hospitalization. A sensitivity analysis was performed evaluating patients with primary pulmonary arterial hypertension. A second sensitivity analysis was performed excluding patients with a diagnosis of venous thromboembolism, in whom PH may have developed acutely in the post-operative state. Venous thromboembolism diagnoses during the surgical hospitalization were identified using ICD-9 diagnosis codes for pulmonary embolism (415.1x) and lower extremity deep vein thrombosis (451.1x, 451.81, 453.2, 453.4x).¹¹ A sensitivity analysis was also performed excluding younger individuals, restricting the analysis to adults age ≥ 45 years old. Sampling weights were applied to determine national incidence estimates when specified. Unweighted data were used in all other analyses. Statistical analyses were performed using SPSS 25 (IBM SPSS Statistics, Armonk, NY). Statistical tests are two-sided and p-values <0.05 were considered to be statistically significant. The NIS is a publicly available, de-identified dataset, and the study was exempt from institutional board review.

Results

A total of 17,853,194 hospitalizations for major non-cardiac surgery were identified. Pulmonary hypertension was present in 143,846 hospitalizations (0.81%) for non-cardiac surgery, corresponding to an estimated 693,819 surgical hospitalizations in the United States during this time period. Characteristics of patients with and without PH are shown in Table 1. Patients hospitalized for surgery with PH were older and less likely to be women than patients without PH. Smoking, obstructive sleep apnea, chronic pulmonary disease, and valvular heart disease were more common in patients with PH than those without PH. In both sexes, the frequency of PH diagnoses increased with age (Figure 1). The proportion of patients undergoing non-cardiac surgery with a diagnosis of PH increased over time, from 0.4% in 2004 to 1.2% in 2014 (Figure 2).

Primary pulmonary arterial hypertension (WHO Group 1 PH) was identified in 3,500 cases, representing 2.4% of surgical hospitalizations with any diagnosis of PH and 0.02% of hospitalizations undergoing non-cardiac surgery. This corresponds to an estimated 16,691 hospitalizations for surgery in the United States during this time period. Patients with WHO Group 1 PH tended to be younger (68.24 vs. 71.66, $p<0.001$) and were more likely to be women (67.7% vs. 63.2%, $p<0.001$). Characteristics of patients with WHO Group 1 PH in comparison to other types of PH are shown in Supplemental Table 2.

Among the 143,846 surgical hospitalizations with a diagnosis of PH, right sided heart catheterization was performed during 2,961 (2.1%) hospitalizations. The proportion of patients who underwent right heart catheterization during the surgical hospitalization declined with increasing age, from 4.3% of patients age 18–39 years to 0.9% of patients older than 80 years of age. For all subtypes of non-cardiac surgery, right sided heart catheterization was performed in fewer than 5% of surgical hospitalizations with an

associated diagnosis of PH. Among those with a diagnosis of pulmonary arterial hypertension (WHO Group 1 PH), right heart catheterization was performed in 4.3% (n=149) of surgical hospital admissions.

Major adverse cardiovascular events were more common in surgical hospitalizations with PH in comparison those without PH (8.3% vs. 2.0, $p<0.001$), driven by an increased frequency of death (4.4% vs. 1.1%, $p<0.001$), non-fatal myocardial infarction (3.2% vs. 0.6%, $p<0.001$) and stroke (1.5% vs. 0.5%, $p<0.001$; Table 2). The rate of perioperative MACCE did not decline over time (Supplemental Figure 1). After adjusting for demographics, clinical covariates, and surgery type, a diagnosis of PH was independently associated with MACCE (aOR 1.43, 95% CI 1.40 – 1.46), in-hospital death (aOR 1.51, 95% CI 1.47 – 1.55) and MI (aOR 1.49, 95% CI 1.44 – 1.54; Table 2). Pulmonary hypertension was associated with a decreased risk of stroke after multivariable adjustment. Patients with PH were more likely to have pulmonary embolism, cardiogenic shock and cardiac arrest in unadjusted and adjusted analyses.

Among the 3500 patients with WHO Group 1 PH, MACCE occurred in 451 (12.9%) of patients undergoing non-cardiac surgery. A total of 289 (8.3%) of patients died, 141 (4.3%) patients developed MI and 66 (1.9%) had an ischemic stroke. Pulmonary embolism occurred in 183 (5.3%) cases, cardiogenic shock occurred in 1.2% of patients and cardiac arrest occurred in 1.6% of cases. WHO Group 1 PH was also independently associated with MACCE after multivariable adjustment for demographics and clinical covariates (aOR 2.51, 95% CI 2.26 – 2.79; Supplemental Table 3).

The prevalence of PH varied by subgroup of non-cardiac surgery. Among patients undergoing transplant, vascular, thoracic and orthopedic surgery, the proportion of patients with PH was 3.92%, 2.33%, 2.00%, and 1.01%, respectively. Among patients with PH who were hospitalized for surgery, the most common surgical subtypes were orthopedic (38.3%, n=55105), vascular (21.4%, n=30771), and general surgery (19.7%, n=28263).

Patients with PH had a higher risk of perioperative MACCE than those without PH, with the highest surgery-specific risks of MACCE for patients undergoing neurosurgery (13.0% versus 4.1, $p<0.001$), thoracic surgery (12.2% vs. 5.6%, $p<0.001$), vascular surgery (11.2% versus 7.3%, $p<0.001$) and general surgery (9.3% versus 2.7%, $p<0.001$). Adjusted odds of MACCE associated with PH by surgical subtype are shown in Supplemental Figure 2. Surgical subtypes with low overall perioperative cardiovascular risk had the greatest increased odds of MACCE associated with PH.

In a sensitivity analysis excluding 12,211 PH patients with a diagnosis of pulmonary embolism or deep vein thrombosis, in whom PH may have developed acutely in the perioperative period, MACCE occurred in 7.9% (n=10,458), perioperative mortality was 4.2% (n=5,522), and MI occurred in 3.0% (n=3989). After the exclusion of patients with acute VTE, PH remained independently associated with MACCE after multivariable adjustment (aOR of 1.47, 95% CI 1.44 – 1.50). Similar findings were observed in a sensitivity analysis of surgical patients older than age 45 years (n=11,403,707), in whom PH

was present in 1.2% of hospitalizations and was independently associated with an increased risk of MACCE (aOR 1.41 [95% CI 1.39 – 1.44]).

Discussion

In a large cohort of 17,853,194 patients undergoing major non-cardiac surgery, PH was identified in 1 of every 125 cases overall, and in 1 of every 84 patients who were age 45 years. Based on these data, more than 60,000 patients with PH are estimated to undergo non-cardiac surgery in the United States each year. Pulmonary hypertension was most common in patients undergoing transplant, vascular, thoracic and orthopedic surgeries. Perioperative MACCE occurred more frequently among patients with PH in comparison to those without, driven by excess mortality, perioperative MI, and ischemic stroke. Among patients with PH, MACCE was diagnosed in 8.3% of cases and 4.4% of patients died in-hospital after surgery. The increased risks of MACCE associated with PH were observed across multiple major surgical subtypes. Pulmonary arterial hypertension, identified in 0.02% of patients undergoing non-cardiac surgery, was a particularly high-risk subgroup with a significant burden of cardiovascular complications.

These findings are consistent with prior reports of patients with PH undergoing noncardiac surgery from small single center retrospective studies. In prior series, perioperative mortality ranged from 4% to 26%, with morbidity 6% to 42% associated with PH.^{2,4,12–16} In a small case-control series of 173 patients, PH was associated with a higher incidence of decompensated heart failure, respiratory failure, hemodynamic instability, prolonged ventilatory support and longer ICU stays.¹⁴ Prior pulmonary embolism, right axis deviation on ECG, right ventricular hypertrophy, elevated RV systolic pressures, vasopressor use, anesthesia without nitrous oxide have been independently associated with perioperative mortality in patients with PH.⁴ The most recent guidelines recognize pulmonary arterial hypertension (WHO Group 1), high pulmonary artery systolic pressures (>70 mmHg), right ventricular dilation, pulmonary vascular resistance >3 Wood units, and New York Heart Association class III or IV symptoms as high risk features of PH.

Patients with PH are particularly vulnerable during the perioperative period, since they tolerate hemodynamic and fluid shifts poorly. General anesthesia, unavoidable for most intermediate and high-risk surgeries, may lead to systemic vasodilatation and perioperative hypotension. Extubation may also be challenging with right ventricular dysfunction and may lead to prolonged intubation. All of these factors may underlie the increased frequency of MACCE observed in the present analysis.

Despite the impact of PH on clinical outcomes, perioperative care guidelines provide limited recommendations for the evaluation and management of PH patients planned for noncardiac surgery. Patients with PH should undergo a comprehensive preoperative evaluation, with assessment of volume status, 12-lead electrocardiography, echocardiography, and appropriate chest imaging based on anticipated surgical risk. Right sided heart catheterization may be useful for invasive confirmation of pulmonary artery pressures and to establish the etiology of PH. However, few patients with PH underwent perioperative right heart catheterization in the present analysis.¹⁷ In addition, recent ACC/AHA guidelines

provide a Class IIa recommendation for pre-operative evaluation by a PH specialist.¹² Whenever possible, non-cardiac surgery should be performed in a center with PH expertise, and cardiac anesthesia involvement is indicated in patients with severe disease.¹⁷ Finally, the ACC/AHA guidelines provide a Class I recommendation to continue home pulmonary vasodilators in patients with PH who are already treated with these agents and are planned for non-cardiac surgery.¹²

There are a few notable limitations of the present analysis. First, diagnoses recorded in the NIS reflect administrative coding data are subject to reporting bias and/or errors. Pulmonary hypertension and perioperative major adverse cardiovascular events may be under-diagnosed. In particular, a recent study raises questions about the accuracy of administrative coding data to diagnose pulmonary arterial hypertension.¹⁸ Second, the results of echocardiography were not recorded in the NIS, and left ventricular function and pulmonary artery pressures are unknown. World Health Organization (WHO) Group classifications were not available. Consequently, the severity and mechanism of PH could not be ascertained. The prognostic significance of PH independent of left ventricular function requires confirmation. Third, in-hospital use of diuretics, pulmonary vasodilators, and anticoagulants could not be determined. Fourth, the sequence of non-cardiac surgery preceding non-fatal cardiovascular complications cannot be definitively established. However, high-risk non-cardiac surgery is contraindicated early after stroke or AMI, and patients with PH presenting with MACCE would be unlikely to undergo non-cardiac surgery during the same hospital admission. Fifth, data were analyzed as a simple random sample unless otherwise specified. Unweighted analyses may introduce empirical weighting that may not be reflective of the national population of patients undergoing surgery. Finally, long-term postoperative outcomes were not available from this in-hospital dataset.

In conclusion, this study is the largest to date to assess the outcomes of PH in non-cardiac surgery. We found that PH is present in 1 of every 125 patients undergoing major non-cardiac surgery and is an important risk factor for perioperative adverse cardiovascular events. Additional research is necessary to evaluate associations between the mechanism and severity of PH and perioperative outcomes. Meticulous preoperative assessment and planning, hemodynamic monitoring during the perioperative period, and appropriate post-operative care remain the cornerstones of care for patients with PH undergoing non-cardiac surgery.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Sponsor / Funding: This research is supported in part by an NYU CTSA grant, UL1 TR001445 and KL2 TR001446, from the National Center for Advancing Translational Sciences, National Institutes of Health.”

REFERENCES:

1. McLaughlin VV, Archer SL, Badesch DB, Barst RJ, Farber HW, Lindner JR, Mathier MA, McGoon MD, Park MH, Rosenson RS, Rubin LJ, Tapson VF, Varga J, Harrington RA, Anderson JL, Bates

- ER, Bridges CR, Eisenberg MJ, Ferrari VA, Grines CL, Hlatky MA, Jacobs AK, Kaul S, Lichtenberg RC, Lindner JR, Moliterno DJ, Mukherjee D, Pohost GM, Rosenson RS, Schofield RS, Shubrooks SJ, Stein JH, Tracy CM, Weitz HH, Wesley DJ. ACCF/AHA 2009 expert consensus document on pulmonary hypertension. *Circulation* 2009;119:2250–2294. [PubMed: 19332472]
2. Lai HC, Lai HC, Wang KY, Lee WL, Ting CT, Liu TJ. Severe pulmonary hypertension complicates postoperative outcome of non-cardiac surgery. *Br J Anaesth* 2007;99:184–190. [PubMed: 17576968]
3. Memtsoudis SG, Ma Y, Chiu YL, Walz JM, Voswinckel R, Mazumdar M. Perioperative mortality in patients with pulmonary hypertension undergoing major joint replacement. *Anesth Analg* 2010;111:1110–1116. [PubMed: 20841415]
4. Ramakrishna G, Sprung J, Ravi BS, Chandrasekaran K, McGoon MD. Impact of pulmonary hypertension on the outcomes of noncardiac surgery: predictors of perioperative morbidity and mortality. *J Am Coll Cardiol* 2005;45:1691–1699. [PubMed: 15893189]
5. Draughn D, Morris DM. Sustained pulmonary hypertension in surgical patients. *Am Surg* 1993;59:346–349. [PubMed: 8507056]
6. Smilowitz NR, Gupta N, Ramakrishna H, Guo Y, Berger JS, Bangalore S. Perioperative Major Adverse Cardiovascular and Cerebrovascular Events Associated With Noncardiac Surgery. *JAMA Cardiol* 2017;2:181–187. [PubMed: 28030663]
7. Steiner C, Elixhauser A, Schnaier J. The healthcare cost and utilization project: an overview. *Eff Clin Pract* 2002;5:143–151. [PubMed: 12088294]
8. Sikachi RR, Sahni S, Mehta D, Agarwal A, Agrawal A. Nationwide trends in inpatient admissions of pulmonary hypertension in the United States from 2000 to 2013. *Adv Respir Med* 2017;85:77–86. [PubMed: 28440533]
9. George MG, Schieb LJ, Ayala C, Talwalkar A, Levant S. Pulmonary hypertension surveillance: United States, 2001 to 2010. *Chest* 2014;146:476–495. [PubMed: 24700091]
10. Smilowitz NR, Gupta N, Guo Y, Berger JS, Bangalore S. Perioperative acute myocardial infarction associated with non-cardiac surgery. *Eur Heart J* 2017;38:2409–2417. [PubMed: 28821166]
11. Trinh VQ, Karakiewicz PI, Sammon J, Sun M, Sukumar S, Gervais MK, Shariat SF, Tian Z, Kim SP, Kowalczyk KJ, Hu JC, Menon M, Trinh QD. Venous thromboembolism after major cancer surgery: temporal trends and patterns of care. *JAMA Surgery* 2014;149:43–49. [PubMed: 24197279]
12. Fleisher LA, Fleischmann KE, Auerbach AD, Barnason SA, Beckman JA, Bozkurt B, Davila-Roman VG, Gerhard-Herman MD, Holly TA, Kane GC, Marine JE, Nelson MT, Spencer CC, Thompson A, Ting HH, Uretsky BF, Wijeyesundera DN. 2014 ACC/AHA Guideline on Perioperative Cardiovascular Evaluation and Management of Patients Undergoing Noncardiac Surgery. *J Am Coll Cardiol* 2014;64:e77–e137. [PubMed: 25091544]
13. Minai OA, Venkateshiah SB, Arroliga AC. Surgical intervention in patients with moderate to severe pulmonary arterial hypertension. *Conn Med* 2006;70:239–243. [PubMed: 16768070]
14. Kaw R, Pasupuleti V, Deshpande A, Hamieh T, Walker E, Minai OA. Pulmonary hypertension: an important predictor of outcomes in patients undergoing non-cardiac surgery. *Respir Med* 2011;105:619–624. [PubMed: 21195595]
15. Price LC, Montani D, Jais X, Dick JR, Simonneau G, Sitbon O, Mercier FJ, Humbert M. Noncardiothoracic nonobstetric surgery in mild-to-moderate pulmonary hypertension. *Eur Respir J* 2010;35:1294–1302. [PubMed: 19897552]
16. Meyer S, McLaughlin VV, Seyfarth HJ, Bull TM, Vizza CD, Gomberg-Maitland M, Preston IR, Barbera JA, Hassoun PM, Halank M, Jais X, Nickel N, Hoeper MM, Humbert M. Outcomes of noncardiac, nonobstetric surgery in patients with PAH: an international prospective survey. *Eur Respir J* 2013;41:1302–1307. [PubMed: 23143546]
17. Minai OA, Yared JP, Kaw R, Subramaniam K, Hill NS. Perioperative risk and management in patients with pulmonary hypertension. *Chest* 2013;144:329–340. [PubMed: 23880683]
18. Papani R, Sharma G, Agarwal A, Callahan SJ, Chan WJ, Kuo YF, Shim YM, Mihalek AD, Duarte AG. Validation of claims-based algorithms for pulmonary arterial hypertension. *Pulm Circ* 2018;8:1–8.

Figure 1.
Frequency of pulmonary hypertension among patients undergoing for non-cardiac surgery, by age and sex.

Figure 2.
The proportion of patients undergoing non-cardiac surgery with a diagnosis of pulmonary hypertension over time

Table 1.

Characteristics of patients age 18 years old hospitalized for non-cardiac surgery, with and without a pulmonary hypertension diagnosis.

Variables	Pulmonary Hypertension		p value
	Yes (n = 143,846)	No (n = 17,709,348)	
Age (years)	71.6 ± 14.6	53.5 ± 19.5	<0.001
Female Sex	91097 (63.3%)	11447976 (64.8%)	<0.001
Non-Hispanic White	96685 (67.2%)	10490933 (59.2%)	
Non-Hispanic Black	17352 (12.1%)	1701974 (9.6%)	
Hispanic	6915 (4.8%)	1646805 (9.3%)	<0.001
Unknown Race	5207 (3.6%)	886398 (5%)	
Other Race	17687 (12.3%)	2983238 (16.8%)	
Smoker	31034 (21.6%)	3208943 (18.1%)	<0.001
Obstructive Sleep Apnea	18007 (12.5%)	522743 (3%)	<0.001
Hypertension	98377 (68.4%)	7203953 (40.7%)	<0.001
Hyperlipidemia	50002 (34.8%)	3487250 (19.7%)	<0.001
Diabetes Mellitus	46724 (32.5%)	2930582 (16.5%)	<0.001
Coronary Artery Disease	51335 (35.7%)	2037382 (11.5%)	<0.001
Prior Percutaneous Coronary Intervention	7728 (5.4%)	431943 (2.4%)	<0.001
Prior Coronary Bypass Grafting	10592 (7.4%)	530568 (3%)	<0.001
Peripheral Vascular Disease	21267 (14.8%)	847146 (4.8%)	<0.001
Valvular Heart Disease	45043 (31.3%)	493374 (2.8%)	<0.001
Malignancy	8038 (5.6%)	677157 (3.8%)	<0.001
Anemia	46003 (32%)	2459097 (13.9%)	<0.001
History of Heart Failure	52791 (36.7%)	734558 (4.1%)	<0.001
Prior Alcohol abuse	3461 (2.4%)	331264 (1.9%)	<0.001
Prior Drug abuse	1972 (1.4%)	253540 (1.4%)	0.05
Collagen Vascular Disease	7623 (5.3%)	367138 (2.1%)	<0.001
HIV/AIDS	227 (0.2%)	20088 (0.1%)	<0.001
Hypothyroidism	23772 (16.5%)	1498825 (8.5%)	<0.001
Obesity	24937 (17.3%)	1730327 (9.8%)	<0.001
Chronic Kidney Disease	36649 (25.5%)	980978 (5.5%)	<0.001
Chronic Pulmonary Disease	51285 (35.7%)	2293988 (13%)	<0.001
Prior Venous Thromboembolism	7354 (5.1%)	335226 (1.9%)	<0.001
Atrial Fibrillation	49597 (34.5%)	957789 (5.4%)	<0.001
Elective Admission for Surgery	50725 (35.4%)	10154622 (57.5%)	<0.001
Surgery Type			<0.001
General	28263 (19.6%)	3520784 (19.9%)	
Endocrine	979 (0.7%)	181206 (1%)	
Genitourinary	5883 (4.1%)	1038193 (5.9%)	
Neurosurgery	4130 (2.9%)	874500 (4.9%)	

Variables	Pulmonary Hypertension		p value
	Yes (n = 143,846)	No (n = 17,709,348)	
Obstetrics/Gynecology	3552 (2.5%)	4254034 (24%)	
Orthopedics	55105 (38.3%)	5391446 (30.4%)	
Otolaryngology	865 (0.6%)	164638 (0.9%)	
Skin/Breast	6361 (4.4%)	654218 (3.7%)	
Thoracic	5988 (4.2%)	293127 (1.7%)	
Transplant	1949 (1.4%)	47788 (0.3%)	
Vascular	30771 (21.4%)	1289414 (7.3%)	

Hypertension was defined by Clinical Classification Software (CCS) diagnosis codes 98 and 99. Hyperlipidemia was defined by CCS diagnosis code 53. Anemia was based on AHRQ comorbidity measure definitions for chronic blood loss and deficiency anemias. Obesity was based on the AHRQ comorbidity measure for obesity.

Table 2.

Perioperative outcomes of non-cardiac surgery, with and without a diagnosis of pulmonary hypertension.

	Pulmonary Hypertension		Unadjusted p value	Adjusted* OR (95% CI)
	Yes (n = 143,846)	No (n = 17,709,348)		
Major Averse Cardiovascular Events	11985 (8.3%)	346110 (2%)	<0.001	1.43 (1.40 – 1.46)
Death	6354 (4.4%)	199234 (1.1%)	<0.001	1.51 (1.47 – 1.55)
Acute Myocardial Infarction	4535 (3.2%)	98026 (0.6%)	<0.001	1.49 (1.44 – 1.54)
Stroke	2192 (1.5%)	82046 (0.5%)	<0.001	0.93 (0.89 – 0.98)
Other Cardiovascular Complications				
Pulmonary Embolism	8824 (6.1%)	124266 (0.7%)	<0.001	3.35 (3.27 – 3.44)
Cardiogenic Shock	916 (0.6%)	12100 (0.1%)	<0.001	2.37 (2.20 – 2.55)
Cardiac Arrest	1651 (1.1%)	41834 (0.2%)	<0.001	1.72 (1.63 – 1.81)

* Adjusted for demographics, clinical comorbidities, urgent/emergent hospitalization, and non-cardiac surgery type as described in the methods.