Published in final edited form as:

J Phys Chem Lett. 2019 March 07; 10(5): 1146–1151. doi:10.1021/acs.jpclett.9b00227.

Folding Landscape of a Parallel G-quadruplex

Robert D. Gray, John O. Trent, Sengodagounder Arumugam, and Jonathan B. Chaires*

J. G. Brown Cancer Center, University of Louisville, 505 S. Hancock St., Louisville, KY 40202

Abstract

Circular dichroism and stopped-flow UV spectroscopies were used to investigate the thermodynamic stability and the folding pathway of d[TGAG₃TG₃TAG₃TG₃TA₂] at 25 °C in solutions containing 25 mM KCl. Under these conditions the oligonucleotide adopts a thermally stable, all-parallel G-quadruplex topography containing three stacked quartets. K⁺-induced folding shows three resolved relaxation times, each with distinctive spectral changes. Folding is complete within 200 s. These data indicate a folding pathway that involves at least two populated intermediates, one of which seems to be an antiparallel structure that rearranges to the final all-parallel conformation. Molecular dynamics reveals a stereochemically plausible folding pathway that does not involve complete unfolding of the intermediate. The rate of unfolding was determined using complementary DNA to trap transiently unfolded states to form a stable duplex. As assessed by 1D-¹H NMR and fluorescence spectroscopy, unfolding is extremely slow with only one observable rate-limiting relaxation time.

Keywords

c-Myc promoter; stopped-flow kinetics; circular dichroism; molecular dynamics simulation; thermodynamics

The folding mechanism of G-quadruplex (G4)-forming DNA has been the subject of numerous studies over the past decade. Most of the kinetic studies carried out either in solution^{1–5}, or by various single-molecule methods^{6–17}, have focused on oligonucleotide models of the human telomere G4 consisting of variations of the sequence d[(GGGTTA)_n] where n = 4^{18–19}. Depending on the exact sequence and identity of added monovalent cation, such telomeric oligonucleotides generally fold into either antiparallel “basket” or “chair” forms or into “hybrid” 3+1 antiparallel G4 topologies in multi-step reactions with spectroscopically distinct intermediates. For example, K⁺-induced folding of the human telomere sequence, d[A(G₃TTA)₃G₃] initially forms an ensemble of hairpins which collapse to an antiparallel structure within ~2 s at room temperature. Subsequently, the antiparallel structure slowly rearranges to a mixture of hybrid-1 and hybrid-2 topologies over a period of

*Corresponding Author: Phone: 502-852-1172. j.chaires@louisville.edu.

The authors declare no competing financial interests.

ASSOCIATED CONTENT

Supporting Information. The following files are available free of charge. A PDF file containing Experimental Methods, two Tables, seven Figures and one Scheme. Two mp4 files showing movies of the transition from the chair to a parallel G-quadruplex conformation derived from nudged elastic band simulations.

hours.^{1, 4–5} Unfolding of telomeric oligonucleotides in the presence of excess complementary DNA is multiphasic and relatively slow (time constant ~2000 s) at room temperature.^{5, 20–21} However, to date there are limited solution studies on the mechanism of folding of G4 DNA folded in an all-parallel topology. The purpose of the current investigation was to apply spectroscopic methods to determine the folding and unfolding pathways for a parallel G4 structure.

To accomplish this objective, we chose the G4-forming oligonucleotide d[**TGAG₃TG₃TAG₃TG₃TA₂**] (that we call “1XAV” after the Protein Data Bank entry), a well-characterized truncated sequence variant of the *c-MYC* promoter DNA. In the presence of K⁺, NMR studies show that this and related sequences fold into a G4 structure of exceptional thermal stability in which the four G₃-tracts are in a parallel configuration.^{3, 22–23} Recently a high-resolution (2.35 Å) crystal structure of the same sequence folded into a parallel G4 structure was reported.²⁴ A single-molecule study using magnetic tweezers was used to estimate an unfolding rate constant of ~10^{–6} s^{–1} for a related *c-MYC* sequence folded into a parallel G4 structure, with the assumption that folding was a two-state mechanism with no intermediates.²⁵ Hysteretic forward and reverse thermal denaturation transitions were used to estimate the folding and unfolding rate constants as a function of temperature for several *c-MYC* sequence variants, again assuming a simple two-state mechanism.³ For some variants that showed hysteresis (but not 1XAV), unfolding rates at physiological temperature of ~200–400 s^{–1} were reported in 2 mM KCl. No kinetic intermediates were considered. We report here thermodynamic and kinetic studies that describe the folding of 1XAV in more detail, revealing populated intermediates within its folding landscape.

Figures 1, S1 and S2 show the results of equilibrium titration and thermal denaturation studies of 1XAV. Upon addition of KCl, the initially unfolded oligonucleotide adopts a structure with a CD spectrum characterized by a maximum at 265 nm, a change in sign at 253.5 nm, and a minimum at 243 nm, consistent with formation of an all-parallel quadruplex topology.^{26–28} The titration curve shown in Figure 1A, derived from more complete spectral data in Figure S1, shows that folding is not a simple two-state process. A phenomenological fit of the titration curve to a biphasic dose-response curve yielded estimates of 67 [61.5;72.8] and 73 [51.2;104.3] μM for the KCl concentrations at half-maximal saturation of each phase, where the brackets indicate the 95% confidence interval. The corresponding Hill coefficients for the two transitions are 2.1 [1.5;2.8] and 0.6 [0.3;0.8]. The very low amplitude of the second transition contributes to the uncertainty of its parameter estimated, although inclusion of the second transition is required for an optimal fit to the experimental data as judged by the randomness of residuals for the fit.

Singular value decomposition was used to analyze thermal denaturation experiments²⁹ as shown in Figure S2 and summarized in Figure 1B. The results show that thermal unfolding is not a simple two-state process, and that a populated intermediate must be included in the reaction mechanism. The two steps in the unfolding process in 25 mM KCl derived by fitting the temperature-dependent family of CD spectra as previously described^{29–30} are characterized by transition melting temperatures of 57.6 ± 6.9 and 78.3 ± 2.1 °C, with van't Hoff enthalpy changes of 19.6 ± 8.7 and 53.9 ± 3.4 kcal mol^{–1}, respectively. From these

thermodynamic values, the overall free energy of folding of 1XAV at 20 °C is estimated to be -11.2 ± 1.3 kcal mol⁻¹ (assuming no heat capacity change). The errors are standard deviations of the mean for six independent replicate melting experiments. The complete thermodynamic profile for 1XAV folding is shown in Figure S2. Folding is driven by a favorable enthalpy and is opposed by an unfavorable entropic contribution.

The results of single-wavelength stopped-flow CD experiments for data collected at 295 nm are summarized in Figure 2. To capture the complete kinetic profile, kinetic traces using the same solutions were collected in separate experiments for 2 s (Figure 2A) and 60 s (Figure 2B). The 2-s data sets could be accurately fit to a single exponential from which a relaxation time of approximately 0.3 s could be determined. The amplitude of this step depended strongly on the wavelength monitored (Figure S3). As shown in Figure 2B, the 60 s data were clearly biphasic, showing the rapid increase in ellipticity during the initial 1–2 s, followed by a slower decrease in ellipticity over the ensuing 60 s with a relaxation time of approximately 9 s. Figure 2C shows a kinetic difference spectrum for the 2-s relaxation that was generated by measuring the kinetics of CD changes at different wavelengths (see Figure S3 for individual experiments). This kinetic difference CD spectrum shows that the folding intermediate(s) initially formed exhibit a spectrum with a maximum near 295 nm that is reminiscent of an antiparallel G4 structure, perhaps mixed with some parallel or hybrid structure. These data show unequivocally that folding is not a simple two-state process, and instead proceeds through at least one intermediate form with distinctive spectral properties. (It is interesting to note that since the duration of a single eye blink is 0.1–0.4 s,^{31–32} the initial folding event is literally over in a blink of the eye.)

An additional slower spectral change, indicative of a second intermediate form, was observed by hand-mixing kinetic experiments with rapid-wavelength scanning. Spectral scans were initiated within 10 s after KCl addition, and accumulated at 10 s intervals over 300 s. The results are shown in Figure 3A. These primary data show an initial transient spectral change near 295 nm that confirms the behavior seen in Figure 2A. Transient changes near 265 nm over a longer time scale are then observed. Singular value decomposition and nonlinear least-squares fitting of the amplitude vectors were used to analyze the data in Figure 3A. The best fit was obtained using a three-step sequential kinetic mechanism $U \leftrightarrow I_1 \leftrightarrow I_2 \leftrightarrow F$, where U and F are the unfolded initial and folded final states, respectively, and I_i are populated intermediate states. In addition to the relaxation times for the two steps seen in Figure 2, an additional slow step with a relaxation time ~60 s was revealed by these experiments. The calculated spectra of the four species are shown in Figure 3B. In the first step, forms with a transient increase I_1 in CD at 295 nm, which then decreases as I_2 forms, which has a maximum CD at 265 nm. Smaller CD changes at 265 nm then accompany the transition of to the final folded I_2 form F . The time-resolved species plot is shown in Figure 3C. These kinetics are fully consistent with the single wavelength stopped-flow experiments shown in Figure 2. The overall folding reaction is complete by 200 s, with three resolved kinetic relaxation times of approximately 0.3, 9 and ~60 s.

Rapid-scanning stopped-flow absorbance studies (Figures S4–S5) and stopped-flow fluorescence studies using a 2-aminopurine substituted oligonucleotide (Figure S6) provided additional kinetic data that are fully consistent with the picture described above.

The rate of the first relaxation time was found to have a hyperbolic dependence on KCl concentration, while the second and third relaxation times were independent of KCl concentration up to 300 mM (data not shown). The temperature dependence of the first relaxation showed (anomalously) an apparent negative activation energy of approximately $-11 \text{ kcal mol}^{-1}$ (data not shown). Such behavior was previously observed for the folding of antiparallel G4 structures¹ and indicates the presence of faster pre-equilibrium steps than can be observed by our methods.

We also characterized the kinetics of 1XAV unfolding by using the well-established complement trap method.^{5, 20–21} In the presence of complementary DNA, transiently unfolded G4 structures can be trapped as the more stable duplex, driving complete unfolding of the quadruplex. Since duplex formation is fast,³³ the observed slow apparent kinetics of duplex formation and G4 disappearance are assumed to arise from the rate-limiting unfolding of the G4 structure. We used ¹H NMR to track unfolding kinetics using the procedure of Lane et al.²¹ Figure 4A shows 1D-¹H NMR spectra of 1XAV obtained periodically during the course of complement-induced unfolding. The zero-time spectrum of folded 1XAV was identical to the 1-D proton NMR spectrum previously published by Ambrus et al.²⁶ Comparison of our spectrum with that of Ambrus et al. allowed assignment of individual resonances to specific G residues. NMR spectra collected hourly over a period of 98 h after addition of 5-fold excess of 1XAV complementary DNA revealed slow disappearance of quadruplex-specific resonances with concomitant appearance of overlapping resonances in the duplex DNA region. The integrated intensities of quadruplex-specific resonances due to imino protons of G residues in the 10.5–12 ppm region are shown in Figure 4B. It is apparent that there was no preferential loss of signal due to specific G residues, implying that no specific region of the quadruplex unfolds separately on this time scale. The unfolding reaction required days to complete and followed a single exponential decay. An average time constant of $2.3 (\pm 0.3) \times 10^5 \text{ s}$ (i.e. $\approx 63 \text{ h}$) was obtained by fits to the time-dependent changes in resonances for both individual G4 guanine residues and base pair formation. An independent experiment (Figure S7) using changes in fluorescence resonance energy transfer (FRET) for 5'-Fam, 3'-Tamra-labeled 1XAV, confirmed the slow unfolding kinetics seen by NMR.

Collectively, all of these kinetic experiments indicate the 1XAV folding is a complex process with populated intermediate steps. At a minimum, folding can be explained by a sequential three-step mechanism

where the initial and final states are U and F and $\langle I_i \rangle$ indicates populated ensembles of intermediates. Relaxation times of approximately 0.3, 2–9 and 60 s characterize the kinetic steps, and folding is complete within 200 s. Note that obligatory coupled K^+ binding events are not specified in this simplified mechanism, but must be present. In addition, the KCl- and temperature-dependence of first relaxation time indicate that faster kinetic steps, beyond the resolution of our methods, surely exist. Only a single, very slow unfolding rate is

observed. Because of the principle of microscopic reversibility, multiple additional unfolding steps must exist but are hidden by the slow rate-limiting process we observe.

A hypothetical model for G4 folding within the c-MYC promoter that is at least superficially consistent with the kinetic described here is shown in Scheme 1. In the context of the duplex promoter, a DNA bubble must form (perhaps driven by supercoiling) to separate the strand containing the G4 forming sequence from its complementary strand. An antiparallel “chair” G4 structure might then form by way of a transient hairpin stabilized by G-G base pairing, a pathway similar to that proposed for the early steps of the formation of telomeric G4 structures.^{1–2, 5} The spectral data in Figures 1 and 2 support the formation of such an intermediate. The final phase requires the unimolecular rearrangement of the antiparallel form to the final parallel structure.

We performed molecular dynamics simulations using the nudged elastic band (NEB) method to determine if there are stereochemically feasible routes from the chair to the final parallel form, following the strategy and methods previously described.⁵ For this simulation we set the starting point as an antiparallel “chair” G4 conformation, as the first experimentally observed intermediate has an antiparallel nature, and the final endpoint as the known 1XAV parallel structure. Therefore, the chair form of the 1XAV sequence d[G₃TG₃TAG₃TG₃] without 5′ and 3′ flanking bases was constructed and used as the starting structure. The corresponding truncated form of the known 1XAV parallel structure was the endpoint. The NEB simulation then located a low energy pathway between the two topological states. This exploration was not intended to be an extensive survey of the energy surface and all low energy pathways. Since this method has not been previously applied to the chair to parallel topological transition, we wanted only to determine if the NEB method could provide a feasible, possible pathway. The results show that it can.

The transition from chair to parallel structure is not a transformation that is intuitively easy to visualize because two diagonally-opposed runs of guanines must change their orientation and reverse direction in going from three lateral loops in the chair form to three chain-reversal (parallel) side loops in the final parallel form (Scheme S1, Supporting Information). If these two transitions were to occur simultaneously, the G-quartets would break and their CD signature would disappear. This was not observed experimentally, as strong G-quartet CD and UV signals were seen throughout the folding reaction. Therefore, the transition from chair to parallel topology must go via two possible pathways involving either parallel-parallel-lateral or lateral-lateral-parallel loop transitions (Scheme S1, Supporting Information). Both of these forms would have antiparallel or hybrid-like CD signatures with characteristic spectroscopic signals in the 295 nm region. These two forms were therefore created and inserted into the pathway in the nudged elastic band simulated annealing to guide the simulations. In both cases the NEB simulations were able to successfully map a pathway from the chair to the parallel form of 1XAV, the results of which are visualized in movies provided in Supporting Information. The general features of the pathway shown in Scheme 1 (and in more detail in Scheme S1) are stereochemically plausible. A recent computational study³⁴ simulated the folding pathway of a parallel RNA G4 structure and concluded, without any experimental verification, that the essence of the process is formation of compact coil-like structures stabilized by extensive cross-like interactions

among the four G-strands. While such interactions may govern the fastest initial steps of the folding process, our experimental results show that folded antiparallel structures are key intermediates along the folding pathway and must be taken into account. Kinetic and thermodynamic studies (using mass spectrometry) of the folding of a similar sequence (d[$\text{TGAG}_3\text{TG}_4\text{AG}_3\text{TG}_4\text{A}_2\text{G}_2$]) derived from the c-myc promoter concluded that folding into final parallel forms proceeded through a branched pathway with several intermediate states with antiparallel characteristics.¹⁸

The foregoing experiments may be relevant to the folding properties of other promoter GQs. Bioinformatic analysis of the human genome suggests that transcriptional promoters are enriched in sequence motifs capable of folding into GQs.³⁵ Based on characteristic CD spectra and DMS footprinting, many promoter sequences of oncogenes,³⁶ including *c-MYC*, *VEGF*, *KRAS*, *c-kit*, as well as the promoter for human telomerase reverse transcriptase (*hTert*),³⁷ contain substantial amounts of parallel GQ structures.

The c-*MYC* GQ consists of a 27-bp sequence (Pu27) with five G-tracts that lie upstream of the so-called P1 promoter which controls synthesis of the Myc protein, a transcription factor that is mutated in a variety of cancers.^{3, 23, 38–40} These mutations lead to constitutive expression of Myc and consequent dysregulation of cellular proliferation. It has been shown that GQ formation in Pu27 prevents *c-MYC* expression; thus characterization of factors influencing the stability of Pu27 model sequences may be relevant to strategies for controlling aberrant Myc activity. The timescale of the folding of 1XAV we observe is relevant since the cell cycle of a mammalian cell is roughly 1 day, and the transcription of a typical gene is roughly 10 minutes.⁴¹ In contrast, the extremely slow unfolding rate that we observe seems to be too slow to be of any physiological relevance, suggesting that disruption of G4 structures in vivo may need catalytic assistance. Indeed, such enzymes exist. Numerous G4 unwinding helicases have been identified that target and regulate specific G-quadruplex structures^{42–43} that would serve to accelerate the slow unfolding reaction we observe.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENT

Supported by grants CA35635 and GM077422 from the National Institutes of Health. We thank Dr. Valerie Gabelica for helpful comments on our manuscript.

REFERENCES

1. Gray RD; Chaires JB, Kinetics and mechanism of K^+ - and Na^+ -induced folding of models of human telomeric DNA into G-quadruplex structures. *Nucleic Acids Res* 2008, 36 (12), 4191–203. [PubMed: 18567908]
2. Gray RD; Li J; Chaires JB, Energetics and kinetics of a conformational switch in G-quadruplex DNA. *J Phys Chem B* 2009, 113 (9), 2676–83. [PubMed: 19708205]
3. Hatzakis E; Okamoto K; Yang D, Thermodynamic stability and folding kinetics of the major G-quadruplex and its loop isomers formed in the nuclease hypersensitive element in the human c-Myc

- promoter: effect of loops and flanking segments on the stability of parallel-stranded intramolecular G-quadruplexes. *Biochemistry* 2010, 49 (43), 9152–60. [PubMed: 20849082]
4. Zhang AY; Balasubramanian S, The kinetics and folding pathways of intramolecular G-quadruplex nucleic acids. *J Am Chem Soc* 2012, 134 (46), 19297–308. [PubMed: 23113843]
 5. Gray RD; Trent JO; Chaires JB, Folding and unfolding pathways of the human telomeric G-quadruplex. *J Mol Biol* 2014, 426 (8), 1629–50. [PubMed: 24487181]
 6. Ying L; Green JJ; Li H; Klenerman D; Balasubramanian S, Studies on the structure and dynamics of the human telomeric G quadruplex by single-molecule fluorescence resonance energy transfer. *Proc Natl Acad Sci U S A* 2003, 100 (25), 14629–34. [PubMed: 14645716]
 7. Lee JY; Okumus B; Kim DS; Ha T, Extreme conformational diversity in human telomeric DNA. *Proc Natl Acad Sci U S A* 2005, 102 (52), 18938–43. [PubMed: 16365301]
 8. Shirude PS; Okumus B; Ying L; Ha T; Balasubramanian S, Single-molecule conformational analysis of G-quadruplex formation in the promoter DNA duplex of the proto-oncogene c-kit. *J Am Chem Soc* 2007, 129 (24), 7484–5. [PubMed: 17523641]
 9. Okamoto K; Sannohe Y; Mashimo T; Sugiyama H; Terazima M, G-quadruplex structures of human telomere DNA examined by single molecule FRET and BrG-substitution. *Bioorg Med Chem* 2008, 16 (14), 6873–9. [PubMed: 18555689]
 10. Shirude PS; Ying L; Balasubramanian S, Single molecule conformational analysis of the biologically relevant DNA G-quadruplex in the promoter of the proto-oncogene c-MYC. *Chem Commun (Camb)* 2008, (17), 2007–9. [PubMed: 18536803]
 11. Lynch S; Baker H; Byker SG; Zhou D; Sinniah K, Single molecule force spectroscopy on G-quadruplex DNA. *Chemistry* 2009, 15 (33), 8113–6. [PubMed: 19603437]
 12. Okumus B; Ha T, Real-time observation of G-quadruplex dynamics using single-molecule FRET microscopy. *Methods Mol Biol* 2010, 608, 81–96. [PubMed: 20012417]
 13. de Messieres M; Chang JC; Brawn-Cinani B; La Porta A, Single-molecule study of G-quadruplex disruption using dynamic force spectroscopy. *Phys Rev Lett* 2012, 109 (5), 058101. [PubMed: 23006209]
 14. Shim J; Gu LQ, Single-molecule investigation of G-quadruplex using a nanopore sensor. *Methods* 2012, 57 (1), 40–6. [PubMed: 22487183]
 15. An N; Fleming AM; Middleton EG; Burrows CJ, Single-molecule investigation of G-quadruplex folds of the human telomere sequence in a protein nanocavity. *Proc Natl Acad Sci U S A* 2014, 111 (40), 14325–31. [PubMed: 25225404]
 16. You H; Zeng X; Xu Y; Lim CJ; Efremov AK; Phan AT; Yan J, Dynamics and stability of polymorphic human telomeric G-quadruplex under tension. *Nucleic Acids Res* 2014, 42 (13), 8789–95. [PubMed: 25013179]
 17. Noer SL; Preus S; Gudnason D; Aznauryan M; Mergny JL; Birkedal V, Folding dynamics and conformational heterogeneity of human telomeric G-quadruplex structures in Na⁺ solutions by single molecule FRET microscopy. *Nucleic Acids Res* 2016, 44 (1), 464–71. [PubMed: 26615192]
 18. Marchand A; Gabelica V, Folding and misfolding pathways of G-quadruplex DNA. *Nucleic Acids Res* 2016, 44 (22), 10999–11012. [PubMed: 27924036]
 19. Šponer J; Bussi G; Stadlbauer P; Kührová P; Banáš P; Islam B; Haider S; Neidle S; Otyepka M, Folding of guanine quadruplex molecules–funnel-like mechanism or kinetic partitioning? An overview from MD simulation studies. *Biochim Biophys Acta* 2017, (5, Part B), 1246–1263.
 20. Green JJ; Ying L; Klenerman D; Balasubramanian S, Kinetics of unfolding the human telomeric DNA quadruplex using a PNA trap. *J Am Chem Soc* 2003, 125 (13), 3763–7. [PubMed: 12656607]
 21. Lane AN; Chaires JB; Gray RD; Trent JO, Stability and kinetics of G-quadruplex structures. *Nucleic Acids Res* 2008, 36 (17), 5482–515. [PubMed: 18718931]
 22. Phan AT; Modi YS; Patel DJ, Propeller-type parallel-stranded G-quadruplexes in the human c-myc promoter. *J Am Chem Soc* 2004, 126 (28), 8710–6. [PubMed: 15250723]
 23. Yang D; Hurley LH, Structure of the biologically relevant G-quadruplex in the c-MYC promoter. *Nucleosides Nucleotides Nucleic Acids* 2006, 25 (8), 951–68. [PubMed: 16901825]

24. Stump S; Mou TC; Sprang SR; Natale NR; Beall HD, Crystal structure of the major quadruplex formed in the promoter region of the human c-MYC oncogene. *PLoS One* 2018, 13 (10), e0205584. [PubMed: 30312328]
25. You H; Wu J; Shao F; Yan J, Stability and kinetics of c-MYC promoter G-quadruplexes studied by single-molecule manipulation. *J Am Chem Soc* 2015, 137 (7), 2424–7. [PubMed: 25654467]
26. Ambrus A; Chen D; Dai J; Jones RA; Yang D, Solution structure of the biologically relevant G-quadruplex element in the human c-MYC promoter. Implications for G-quadruplex stabilization. *Biochemistry* 2005, 44 (6), 2048–58. [PubMed: 15697230]
27. Kypr J; Kejnovská I; Reniuk D; Vorlíková M, Circular dichroism and conformational polymorphism of DNA. *Nucleic Acids Res* 2009, 37 (6), 1713–1725. [PubMed: 19190094]
28. Karsisiotis AI; Hessari NM; Novellino E; Spada GP; Randazzo A; Webba da Silva M, Topological characterization of nucleic acid G-quadruplexes by UV absorption and circular dichroism. *Angew Chem Int Ed* 2011, 50 (45), 10645–8.
29. Gray RD; Chaires JB, Analysis of multidimensional G-quadruplex melting curves. *Curr Protoc Nucleic Acid Chem* 2011,
30. Gray RD; Buscaglia R; Chaires JB, Populated intermediates in the thermal unfolding of the human telomeric quadruplex. *J Am Chem Soc* 2012, 134 (40), 16834–44. [PubMed: 22989179]
31. Milo R; Jorgensen P; Moran U; Weber G; Springer M, BioNumbers--the database of key numbers in molecular and cell biology. *Nucleic Acids Res* 2010, 38 (Database issue), D750–3. [PubMed: 19854939]
32. Schiffman HR, Sensation and Perception. An Integrated Approach. John Wiley and Sons, Inc: New York, 2001.
33. Rauzan B; McMichael E; Cave R; Sevcik LR; Ostrosky K; Whitman E; Stegemann R; Sinclair AL; Serra MJ; Deckert AA, Kinetics and thermodynamics of DNA, RNA, and hybrid duplex formation. *Biochemistry* 2013, 52 (5), 765–72. [PubMed: 23356429]
34. Havrila M; Stadlbauer P; Kuhrova P; Banas P; Mergny JL; Otyepka M; Sponer J, Structural dynamics of propeller loop: towards folding of RNA G-quadruplex. *Nucleic Acids Res* 2018, 46 (17), 8754–8771. [PubMed: 30165550]
35. Huppert JL; Balasubramanian S, G-quadruplexes in promoters throughout the human genome. *Nucleic Acids Res* 2007, 35 (2), 406–413. [PubMed: 17169996]
36. Qin Y; Hurley LH, Structures, folding patterns, and functions of intramolecular DNA G-quadruplexes found in eukaryotic promoter regions. *Biochimie* 2008, 90 (8), 1149–71. [PubMed: 18355457]
37. Chaires JB; Trent JO; Gray RD; Dean WL; Buscaglia R; Thomas SD; Miller DM, An improved model for the hTERT promoter quadruplex. *PLoS One*. 2014, 9 (12), e115580 DOI:10.1371/journal.pone.0115580. [PubMed: 25526084]
38. Siddiqui-Jain A; Grand CL; Bearss DJ; Hurley LH, Direct evidence for a G-quadruplex in a promoter region and its targeting with a small molecule to repress c-MYC transcription. *Proc Natl Acad Sci U S A* 2002, 99 (18), 11593–8. [PubMed: 12195017]
39. Grand CL; Powell TJ; Nagle RB; Bearss DJ; Tye D; Gleason-Guzman M; Hurley LH, Mutations in the G-quadruplex silencer element and their relationship to c-MYC overexpression, NM23 repression, and therapeutic rescue. *Proc Natl Acad Sci U S A* 2004, 101 (16), 6140–5. [PubMed: 15079086]
40. Islam MA; Thomas SD; Murty VV; Sedoris KJ; Miller DM, c-Myc quadruplex-forming sequence Pu-27 induces extensive damage in both telomeric and nontelomeric regions of DNA. *J Biol Chem* 2014, 289 (12), 8521–31. [PubMed: 24464582]
41. Shamir M; Bar-On Y; Phillips R; Milo R, SnapShot: Timescales in cell biology. *Cell* 2016, 164 (6), 1302–1302 e1. [PubMed: 26967295]
42. Mendoza O; Bourdoncle A; Boule JB; Brosh RM Jr.; Mergny JL, G-quadruplexes and helicases. *Nucleic Acids Res* 2016, 44 (5), 1989–2006. [PubMed: 26883636]
43. Sauer M; Paeschke K, G-quadruplex unwinding helicases and their function in vivo. *Biochem Soc Trans* 2017, 45 (5), 1173–1182. [PubMed: 28939694]

Figure 1.

Equilibrium studies of 1XAV folding. (A) KCl titration of 1XAV. The data were fit to a double Hill equation (solid line) to obtain transition midpoints and Hill coefficients. The dashed line shows the fitted first transition, while the dot-dash line shows the second resolved transition. (B) Species distribution plot derived from SVD analysis of thermal denaturation experiments. The dashed line shows the disappearance of the initial folded form, the dot-dash line shows the transient formation of an intermediate, and the solid line shows the formation of the unfolded strand.

Figure 2.

Single-wavelength stopped-flow studies of 1XAV folding. (A). CD change at 295 nm observed for 2 seconds after KCl mixing. (B) CD change at 295 nm observed for 60 seconds after KCl mixing. (C) Kinetic difference spectrum derived from stopped-flow measurements observed 2 seconds after KCl mixing.

Figure 3.

Kinetic studies of 1XAV folding using rapid scanning circular dichroism. (A) Primary data collected at 10 second intervals beginning 10 seconds after KCl mixing. The inset shows the time course of CD changes at 265 nm. (B) Computed CD spectra of kinetic species derived by SVD analysis. The black line is the initial unfolded strand. The red line is the first intermediate. The blue line is the second intermediate. The magenta line is the final folded 1XAV form. (C) Amplitudes of the changes in species spectra as a function of time as derived by SVD analysis. The inset shows an expanded scale for the initial portion of the time course. Colors refer to the spectral species shown in panel (B).

Figure 4.

1XAV unfolding monitored by NMR. (Top) Selected NMR spectra obtained at the indicated times after addition of the complimentary trap strand. Resonances over the 10–12 ppm range are for guanines with the folded 1XAV structure. Resonances over the 12.5 – 14 ppm range are from duplex base pairs formed in the trap reaction. (Bottom) The time course of changes of selected NMR resonances. The black symbols are for the integrated area of GC duplex base pair resonances (left scale). The blue symbols refer to the integrated areas of G-quartet guanines 13 and 22 (right scale).

Scheme 1.

Hypothetical folding pathway within the c-myc promoter, as described in the text.