Published in final edited form as:

Mol Cell Endocrinol. 2018 March 05; 463: 116–130. doi:10.1016/j.mce.2017.03.022.

Regulation of Reproduction via Tight Control of Gonadotropin Hormone Levels

Djurdjica Coss

Division of Biomedical Sciences; School of Medicine, University of California, Riverside;
Riverside, CA 92521

Abstract

Mammalian reproduction is controlled by the hypothalamic-pituitary-gonadal axis. GnRH from the hypothalamus regulates synthesis and secretion of gonadotropins, LH and FSH, which then control steroidogenesis and gametogenesis. In females, serum LH and FSH levels exhibit rhythmic changes throughout the menstrual or estrous cycle that are correlated with pulse frequency of GnRH. Lack of gonadotropins leads to infertility or amenorrhea. Dysfunctions in the tightly controlled ratio due to levels slightly outside the normal range occur in a larger number of women and are correlated with polycystic ovaries and premature ovarian failure. Since the etiology of these disorders is largely unknown, studies in cell and mouse models may provide novel candidates for investigations in human population. Hence, understanding the mechanisms whereby GnRH regulates gonadotropin hormone levels will provide insight into the physiology and pathophysiology of the reproductive system. This review discusses recent advances in our understanding of GnRH regulation of gonadotropin synthesis.

Keywords

Luteinizing hormone; Follicle-stimulating hormone; Differential regulation; GnRH pulse frequency; Reproduction

1. Physiology and pathophysiology of LH and FSH

Gonadotropin hormones, luteinizing hormone (LH) and follicle-stimulating hormone (FSH) are synthesized by the anterior pituitary gonadotropes and secreted into the circulation to regulate gonadal function. Every secretory pulse of LH from the pituitary corresponds to a pulse of GnRH from the hypothalamus (Levine et al., 1982). On the other hand, FSH secretion is not entirely regulated by GnRH and most of FSH is constitutively released (Culler and Negro-Vilar, 1987; Levine and Duffy, 1988). In females, LH and FSH

Corresponding author: Djurdjica Coss, Division of Biomedical Sciences, School of Medicine, 303 SOM Research Building, University of California, Riverside; Riverside, CA 92521, Tel: 951 827-7791, Fax: 951 827-2477, djurdjica.coss@ucr.edu.

Disclosure

The author has nothing to disclose

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

concentration fluctuates throughout the menstrual or estrous cycle and the levels are firmly controlled to assure reproductive fitness and new waves of steroidogenesis and gametogenesis (Pierce, 1988). Pulsatile LH secretion maintains basal level of this hormone in the circulation and steroidogenesis throughout the cycle. LH increases steroid hormone production via induction of steroidogenic enzymes in interstitial Leydig cells and theca cells in males and females, respectively. In females, LH exhibits a 5–10 fold mid-cycle increase that causes ovulation. FSH in males induces androgen binding protein and contributes to spermatogenesis through interaction with testosterone. In females, FSH stimulates aromatase expression in follicle granulosa cells, which increases estrogen production and follicular growth. FSH increases together with LH during the preovulatory surge and exhibits a separate, second increase that is necessary for folliculogenesis.

LH and FSH are critical for reproductive fitness and thus, survival of the species. LH is required for pubertal development and gonadal function, and a lack results in hypogonadism and infertility in both sexes in humans and mouse models (Huhtaniemi, 2006; Huhtaniemi et al., 2006; Ma et al., 2004). FSH in humans is also critical for reproductive function in both genders, since its deficiency results in absent or incomplete pubertal development in women, relatively normal pubertal development but azoospermia in men, and infertility in both women and men (Lamminen et al., 2005). Female mice deficient in FSH have a block in folliculogenesis prior to the antral stage resulting in infertility, while males have impaired reproductive function due to lower sperm count, but are not infertile (Kumar et al., 1997).

LH and FSH are glycoprotein heterodimers, each comprised of a common α subunit (*Cga*, or α GSU) and a unique β subunit (Pierce and Parsons, 1981). α GSU is minimally regulated by hormones, since it is expressed at the sufficiently high basal level (Chin and Boime, 1990). Unique β -subunits provide biological specificity and are limiting components in the mature hormone synthesis, whose amounts are regulated at the transcriptional level of the β -subunits (Kaiser et al., 1997b; Papavasiliou et al., 1986). Pulsatile GnRH secretion increases transcription of LH β subunit (Burger et al., 2009; Haisenleder et al., 2008; Haisenleder et al., 1988). Although FSH can be secreted without a corresponding GnRH pulse, FSH β transcription is dependent on GnRH. In mice lacking GnRH (Mason et al., 1986) or GnRH receptor (Wu et al., 2010), FSH levels, as well as LH levels, are 60%–90% lower in males and females. GnRH injection in rats with low endogenous GnRH increased FSH β transcription 4 fold (Dalkin et al., 2001). This level of increase is comparable to the changes in FSH concentration in the circulation throughout the cycle. Therefore, while concentration of LH in the circulation is regulated at both the level of β subunit transcription and GnRH-stimulated secretion, FSH concentration is regulated primarily at the level of β subunit transcription. Regulation of gonadotropin synthesis and differential expression is not elucidated at the mechanistic level. Herein, we discuss recent findings since our previous review (Thackray et al., 2010), concentrating on the regulation of β -subunit transcription by GnRH. Our review may shed light on differential regulation and point to remaining gaps in our knowledge, providing basis for future investigations that may lead to novel therapeutic treatments of reproductive disorders discussed below.

1.1. Differential regulation of LH and FSH synthesis

Both gonadotropins are synthesized in the same gonadotrope cells, but differences in the relative amounts occur at specific stages of the menstrual or estrous cycle. As illustrated above, LH exhibits a single increase during the preovulatory surge, while FSH increases two times during the menstrual or estrous cycle. The question of how are these two hormones differentially regulated in the same gonadotrope population, by the same hormonal milieu, has been an area of intense investigation. Differential levels of LH and FSH may stem from differences in secretion, since LH is secreted strictly through the GnRH-regulated secretory pathway while FSH is constitutively secreted (Levine and Duffy, 1988).

Alternatively, post-translational modifications of these hormones contribute to differences in the half-life that would ultimately affect concentration in the circulation. Namely, LH and FSH are N-linked glycosylated proteins, containing N-acetylglucosamine residue linked to an amide group of an asparagine (Asn) amino acid in the peptide chain (Green et al., 1986). N-linked glycosylation of LH and FSH is critical for their bioactivity and non-glycosylated forms serve as antagonists (Fares, 2006). This is likely due to necessity of glycosylation for dimerization of α and β subunits and subsequent glycohormone secretion (Wang et al., 2016). LH and FSH have many molecular variants depending on the differences in their oligosaccharide chains. Alkalinity/acidity of variants, and their half-life in the circulation, depends in part on sialylation and sulfation of terminal galactose. There is a shift to less basic variants of LH at the evening of proestrus and to more basic variants of LH after ovariectomy (De Biasi et al., 2002), while FSH is shifted to less acidic variant at midcycle (Wide and Bakos, 1993), implying endocrine regulation of glycosylation enzymes' activity (Damian-Matsumura et al., 1999; Ulloa-Aguirre et al., 2001). Furthermore, human FSH is primarily sialylated while LH is sulfated, a difference that contributes to their variable clearance rates (Green et al., 1986). Since the enzyme that adds the sulfate, sulfotransferase, does not recognize specific peptide sequence but requires only tri-saccharides (Ulloa-Aguirre et al., 1999), enzyme synthesis or activity represents another layer of potential differential regulation of gonadotropin hormones synthesized in the same cell.

Lastly, β -subunit transcription precedes changes in hormone concentration in the circulation, and fluctuations in mRNA levels in the gonadotrope correlate with concentration of the hormones. Therefore, transcriptional regulation of β -subunits may be a critical point that determines levels of gonadotropin hormones. A midcycle surge of GnRH elicits a surge in both LH and FSH, resulting in ovulation of the mature follicle in response to LH (Jones, 1997). LH β gene is induced five fold and FSH β transcription increases four fold in the afternoon of proestrus prior to the onset of the surge (Butcher et al., 1974; Ortolano et al., 1988; Zmeili et al., 1986). In rodents, during the morning of estrus, a secondary FSH raise occurs without a concomitant increase in LH (Besecke et al., 1997; Woodruff et al., 1996). FSH β mRNA is induced three fold during this time (Halvorson et al., 1994; Ortolano et al., 1988). This secondary increase in FSH is essential for follicular development for the subsequent estrous cycle (DePaolo et al., 1979; Hoak and Schwartz, 1980; Papavasiliou et al., 1986). In humans, the FSH levels in the serum increase in the late luteal phase through the mid follicular phase of the menstrual cycle. This FSH increase corresponds to a time when the next cohort of follicles is being recruited during folliculogenesis. Thus, although

responding to the same hormonal milieu, synthesis of LH and FSH is differentially regulated during the estrous or menstrual cycle and this regulation appears to occur primarily at the level of the β subunits transcription. This differential expression of two gonadotropin hormones from the same cell in response to the same endocrine stimuli is critical for reproductive fitness.

1.2. Polycystic ovary syndrome (PCOS)

Higher than normal levels of gonadotropins in females also lead to reproductive pathologies, albeit less severe than a lack of hormone. Most notably, polycystic ovary syndrome (PCOS) is characterized by an increase specifically in LH (Chang, 2007; Hall et al., 1998; Morales et al., 1996), and premature ovarian failure (POF) or primary ovarian insufficiency is associated with an increase in FSH (Chand et al., 2010; Goswami and Conway, 2007). The etiology of PCOS and POF is not clear; it may stem from dysregulation of GnRH pulsatility, pituitary responsiveness, or ovarian hormone production. Endocrine feedback loops cause difficulty in pin-pointing the primary site of dysregulation.

PCOS affects 10% of reproductive age women, yet much is unknown regarding the cause and pathophysiology (De Leo et al., 2016). This is complicated by the fact that the syndrome is categorized with heterogeneous characteristics and patients present with a combination of several symptoms such as chronic or frequent anovulation, polycystic ovaries, hyperandrogenism, hirsutism, acne, obesity, and insulin resistance (Goodarzi et al., 2011). Elevated LH levels are observed in 55%–75% of the women with PCOS, leading to elevated LH/FSH ratio (De Leo et al., 2016). Since obesity is often comorbid with PCOS, and obesity causes reduced LH levels (Mihalca and Fica, 2014; Pasquali et al., 2002; Rosenfield and Bordini, 2010), PCOS women with normal LH levels may have competing influences of obesity and PCOS per se, that keep the LH within the normal range. In a support of this supposition, lean PCOS patients have elevated LH levels (Pangaribuan et al., 2011; Sahin et al., 2014). Recent genomic studies correlated polymorphism in the LH β gene to the occurrence of PCOS (El-Shal et al., 2016), implicating dysregulation in LH β transcription, and possibly higher LH levels, among the causes of PCOS. Surprisingly, another genome wide association study correlated polymorphism in FSH β promoter with PCOS, implicating changes in FSH levels with this disorder, maybe through altered estrogen synthesis (Hayes et al., 2015).

Preponderance of animal studies point to prenatal or early hyperandrogenemia as a contributing cause of PCOS (Abbott et al., 2008; Abbott et al., 2013; Kauffman et al., 2015; Manikkam et al., 2008; Roland et al., 2010; Veiga-Lopez et al., 2008). High insulin levels, with or without obesity, in PCOS patients may be a cause of hyperandrogenemia since insulin in synergy with LH elicits ovarian androgen production (recently reviewed in (Rosenfield, 2013)). Subsequently, hyperandrogenemia may diminish progesterone inhibition of GnRH pulse frequency, causing rapid pulsatile secretion of LH and further increase in ovarian androgen production, correlating regulation of LH frequency and amplitude to PCOS (Moore et al., 2015; Pielecka et al., 2006).

1.3. Premature ovarian failure (POF)

Premature ovarian failure (POF) or primary ovarian insufficiency is a disorder affecting 1% of reproductive age women that lose ovarian function before the age of 40. Majority of POF cases are idiopathic, while 10% are acquired due to chemotherapy, autoimmunity, or toxicity, and 20% have known genetic causes (Rossetti et al., 2016). Fragile X mutation or pre-mutation carriers comprise a bulk of individuals with known genetic causes of POF (Macpherson and Murray, 2016). The rest have mutations in autosomal genes, such as FOXL2, FSH receptor, aromatase and others, which were associated with POF (Laven, 2016).

However, as mentioned, in 70% of patients POF is idiopathic. They are diagnosed usually after ovarian function has ceased, since they seek care due to infertility or amenorrhea. At that time it is not clear if increased FSH stems from a deficiency in ovarian inhibin and is the result of a lack of negative feedback, or if increased FSH was a contributing factor (Laven, 2016). Thus, there is disagreement whether the high FSH level in POF is preceded by diminished ovarian inhibin level; or alternatively, high FSH leads to recruitment of larger number of follicles into a growing pool in every cycle in younger women resulting in early depletion. In the Fragile X syndrome mouse model, FSH is elevated prior to the loss of ovarian function, strengthening the possibility that high FSH may be a cause and not a consequence (Lu et al., 2012). Recently, it was determined that increased levels of FSH in the circulation in other mouse models, due to overexpression or re-trafficking of FSH to the GnRH-regulated secretory pathway, resulted in hemorrhagic ovaries and 5-fold more corpora lutea, implicating higher FSH levels in greater follicle recruitment (Kumar et al., 1999; Wang et al., 2014). Mouse model of constitutively active FSH receptor also experiences accelerated loss of follicular reserves (Peltoketo et al., 2010). Furthermore, polymorphism in the human FSH β promoter, which regulates β subunit expression, increases FSH levels and has been associated with early menarche and premature menopause in women (He et al., 2010). Another genome-wide association study reported that polymorphism in the 5' region of the FSH β gene is also associated with higher FSH levels in the circulation, premature puberty, early natural menopause, and dizygotic twinning (Mbarek et al., 2016).

Therefore, although feedbacks complicate pin-pointing the primary site of dysregulation, genomic evidence ties polymorphism in the FSH β gene promoter to POF, indicating that regulation of FSH β subunit expression may contribute significantly to the etiology, at least in some affected individuals. Alternatively, mutations in transcription factors that influence FSH β transcription, such as FOXL2, or signaling pathways that lead to induction of FSH β mRNA may contribute to POF. Given that 70% of POF patients develop this disorder from unknown causes, studies using mouse models or model cell lines, may provide potential candidates to investigate etiology in human population.

2. GnRH receptor signaling pathways that regulate gonadotropin gene expression

The complexity of GnRH signaling provides a variety of regulatory avenues for gonadotropin gene regulation (Fig. 1). Molecular studies that identified signal transduction pathways activated by GnRH receptor (GnRHR) binding have been facilitated with the development of model cells lines. Whole animal studies, although critical for understanding of the HPG axis within the endogenous hormonal setting, can be inconclusive as illustrated above, due to hormonal feedback and the possible effects on different levels of the axis. Primary pituitary cells are frequently employed to study hormonal effects on gonadotropin synthesis. However, they are not conducive to analyses at the molecular levels due to scarcity of gonadotropes that comprise around 10% of the mixed cells population in the anterior pituitary. Two immortalized clonal cell lines were developed in the Mellon laboratory using targeted tumorigenesis with SV40 T-antigen, which contributed greatly to delineating GnRH signaling pathways and transcriptional regulation of β -subunits (Alarid et al., 1998; Alarid et al., 1996). The α T3-1 cell line created using the 5' regulatory regions of α -GSU approximates an immature gonadotrope cell. It expresses α -GSU and GnRH receptor, but not β -subunits. On the other hand, the L β T2 cell line created using LH β promoter, displays a more mature phenotype since it expresses LH β and FSH β and secretes LH and FSH in response to hormonal cues (Graham et al., 1999; Pernasetti et al., 2001).

Because it may underlie molecular causes of subfertility, pituitary responsiveness to GnRH leading to induction of gonadotropin expression is an important area of investigation. GnRH binding to its G protein-coupled receptor (GPCR), expressed on the gonadotropes of the anterior pituitary, activates the synthesis of the gonadotropin hormones, LH and FSH, and stimulates their secretion (Seeburg et al., 1987; Vale et al., 1977). GnRH is secreted in a pulsatile fashion into the hypophysial-portal system, while long-term tonic exposure to GnRH adversely affects gonadotrope function (Belchetz et al., 1978; Burrin and Jameson, 1989; Shupnik, 1990). Frequency of GnRH pulses changes throughout the menstrual and estrous cycle, modulated by the upstream regulatory neuronal network. Kisspeptin, secreted by several neuronal populations, is a crucial neuropeptide that synchronizes GnRH neuron secretion and stimulates GnRH release (Semaan and Kauffman, 2010). GnRH exhibits high pulse frequency in proestrus during the preovulatory surge (Harris and Levine, 2003; Levine et al., 1985; Levine and Ramirez, 1982). Several seminal studies analyzing β subunits transcription in the rat in vivo determined that fast pulse frequency of GnRH favors LH β synthesis, while FSH β expression is more highly induced by low pulse frequency (Burger et al., 2002; Dalkin et al., 1989; Haisenleder et al., 1991; Kaiser et al., 1997b; and recently reviewed in Thompson and Kaiser, 2014). However, the mechanisms of gonadotrope responsiveness to varied pulse frequencies are only beginning to emerge.

Although a member of the G protein-coupled seven transmembrane domain receptors, GnRHR has a unique structure (Neill, 2002; Tsutsumi et al., 1992). Unlike the other family members, it lacks the intracellular carboxyl-terminal tail and contains several sequence differences in regions otherwise highly conserved (Kaiser et al., 1997a; Sealfon et al., 1997; Stojilkovic et al., 1994; Tsutsumi et al., 1992). The absence of the C-terminal tail that

normally mediates desensitization through β -arrestin association and encapsulation into clathrin-coated vesicles renders the receptor resistant to this regulatory mechanism. Thus, the receptor does not internalize as rapidly as other GPCRs (Blomenrohr et al., 1999; Hislop et al., 2001; McArdle et al., 2002; Pawson et al., 2008; Willars et al., 1999), leading to longer signaling, which raises a question of intracellular negative feedback that stops the signal. Since gonadotropin gene expression is dependent upon GnRH pulses, the gonadotropes must maintain sensitivity to subsequent GnRH pulses by resolving the effect of a prior pulse. Several steps of the GnRH signaling pathways therefore contain negative regulators, in addition to molecules that propagate the signal, which will be discussed below. This rapid adaptation and maintenance of responsiveness underlies the ability of gonadotropes to properly express LH and FSH and regulate reproductive function.

2.1. G-proteins

Ligand binding to the GnRHR causes activation of trimeric G proteins. $G_{\alpha q/11}$ isoform is activated in a model of immature gonadotrope, $\alpha T3$ cells, while both $G_{\alpha q/11}$ and $G_{\alpha s}$ are involved in GnRHR signaling in a model of mature gonadotrope L β T2 cells and in primary rat cultures. Some involvement of $G_{\alpha i}$ and $G_{\alpha 12/13}$ was shown as well, but without a clear downstream effect (Liu et al., 2002; Liu et al., 2003; Naor et al., 2000; Sim et al., 1995; Stanislaus et al., 1998). GnRHR coupling to various G proteins has been reviewed of late (Naor and Huhtaniemi, 2013) and it won't be discussed in depth here. Although GnRH signaling has been reviewed recently as well (Perrett and McArdle, 2013), in this review we will focus on how these various signaling pathways and downstream effectors contribute to differential expression of gonadotropin β subunits.

Recently the functional significance of activation of $G_{\alpha q/11}$ and $G_{\alpha s}$ trimeric G proteins by GnRH was assessed in L β T2 using siRNA. The study demonstrated that endogenous FSH β mRNA induction requires $G_{\alpha q/11}$, while maximal LH β involves $G_{\alpha s}$ (Choi et al., 2012). Surprisingly, $G_{\alpha s}$ activity may diminish FSH β expression, via induction of inhibin α (Choi et al., 2012). Thus, engagement of either $G_{\alpha q/11}$ or $G_{\alpha s}$ may lead to differential expression of gonadotropin β subunits.

Activation of $G_{\alpha s}$ normally leads to cAMP-PKA signaling, but downstream targets of this pathway at the level of the LH β promoter are unknown. Elegant study by Webster lab demonstrated that GnRH signal has a unique kinetics of cAMP generation and PKA activation compared to other stimuli that activate the same pathway (i.e. Pituitary adenylate cyclase-activating polypeptide (PACAP, (Tsutsumi et al., 2010)). More importantly, the study demonstrated that $G_{\alpha s}$ -cAMP-PKA module is not susceptible to desensitization (Tsutsumi et al., 2010).

More commonly, GnRHR couples to $G_{\alpha q/11}$ and activates phospholipase C β (PLC), causing a creation of inositol 1,4,5-triphosphate (IP3) and diacylglycerol (DAG; Stojilkovic et al., 1994). IP3 formation and IP3 receptor activation leads to an increase in intracellular calcium concentration via calcium mobilization from intracellular stores. Increase in DAG and intracellular calcium concentration leads to activation of PKC and calcium calmodulin kinase II (CamKII), respectively. $G_{\alpha q/11}$ -PLC-DAG/Calcium signaling module desensitizes at several steps after GnRH stimulation: at DAG generation and at calcium increase

(Tsutsumi et al., 2010). The latter may be due to the loss of IP₃ receptors via downregulation that are necessary for calcium mobilization (McArdle et al., 1999; McArdle et al., 2002; McArdle et al., 1996). Thus, fast pulse frequency of GnRH from the hypothalamus may desensitize Gα_q/11-DAG and lead to lower FSHβ expression, while Gα_s does not desensitize and leads to increased LHβ. The question whether Gα_s signal is less sensitive or not sustained during low pulse frequency, which results in lower LHβ than FSHβ needs to be elucidated. It has been postulated that GnRH may switch to Gα_s-cAMP signal only during preovulatory surge to engage new set of signaling targets (Lariviere et al., 2007). Indeed, fast pulse frequency increases receptor concentration on the cell surface (Bedecarrats and Kaiser, 2003). This higher receptor number may allow coupling to and engaging Gα_s in addition to Gα_q/11, which induces higher levels of LHβ necessary for a rapid increase of LH in the circulation and ovulation, while low pulse frequency and Gα_q/11 activation may be sufficient for lower level of LHβ induction, as well as FSHβ, throughout the rest of the cycle to stimulate steroidogenesis and folliculogenesis (Fig. 2).

2.2. Calcium

GnRH receptor activation causes an increase in intracellular calcium concentration. The calcium increase stems either from mobilization from intracellular calcium stores or from influx through the calcium channels (Mulvaney et al., 1999; Van Goor et al., 1999); recently reviewed in (Stojilkovic, 2012; Stojilkovic et al., 2010). As discussed above, increase in calcium from intracellular stores occurs through IP₃ formation, and is sensitive to desensitization via downregulation of IP₃ receptors (McArdle et al., 1999; McArdle et al., 2002; McArdle et al., 1996). Consistent with this, using αT3 cells, it was demonstrated that extracellular calcium influx leads to more sustained increase in calcium than mobilization from intracellular stores (Mulvaney and Roberson, 2000). Calcium channels may be activated by GnRH through PKC (Stojilkovic et al., 1991). Pulsatile stimulation of the calcium channel by a pharmacological activator in the rat pituitary cells exhibits the same effect on the gonadotropin gene expression as GnRH pulses, with higher pulse frequencies favoring LHβ transcription, and lower pulse frequencies preferring FSHβ transcription (Haisenleder et al., 2001).

Increase in calcium concentration activates CamKII, and CamKII is activated by GnRH irrespective of pulse frequency (Burger et al., 2008) and plays a role in induction of both β-subunits (Ely et al., 2011; Haisenleder et al., 2003b). Consistent with a lack of frequency dependence, CamKII activation can be achieved with either source of calcium, extracellular or from internal stores in LβT2 cells (Haisenleder et al., 2003b). Nuclear factor of activated T-cells (NFAT) is another putative target of calcium increase via calcineurin that may be activated by GnRH to induce immediate early gene promoters (Binder et al., 2012). Calcium causes calmodulin/calcineurin-dependent nuclear NFAT translocation, but a lack of desensitization of this pathway implies that while NFAT may mediate GnRH action, it is not a decoder of GnRH pulse frequency (Armstrong et al., 2009). Additionally, the effect of NFAT on gonadotropin β-subunit expression has not been demonstrated. Thus, CamKII maybe a critical signaling molecule activated by calcium.

2.3. MAPK

GnRHR ligand binding also leads to activation of three branches of the mitogen-activated protein kinase (MAPK) pathway: ERK1/2, JNK, and p38 (Liu et al., 2002; Roberson et al., 1995; Sundaresan et al., 1996). Mechanisms of MAPK activation are not completely clear. GnRHRs are found in low-density membrane microdomains called lipid rafts, which are enriched in cholesterol and sphingolipids and may allow assembly of multiprotein signaling complexes (Navratil et al., 2003). In α T3 cells and mouse pituitary cells, GnRHRs co-localize with a proportion of total cellular ERK2 in lipid rafts (Bliss et al., 2007), which may provide a partial explanation of ERK2 activation by GnRH. Activation of ERK1/2 by GnRH also involves calcium. ERK1/2 responds to pulsatile GnRH, and lower pulse frequency leads to longer, sustained activation in L β T2 mature gonadotrope cell line, which may correlate ERK1/2 activation to FSH β expression (Kanasaki et al., 2005). Additionally, low pulse frequency of GnRH causes high amplitude of ERK1/2 activation that is necessary for FSH β transcription in vivo in rats (Burger et al., 2008; Haisenleder et al., 1998). Extracellular calcium influx through voltage gated calcium channel may be necessary for ERK1/2 activation in α T3 cells (Mulvaney et al., 1999). The study demonstrating that PKC stimulation leads to voltage gated calcium channel activation (Stojilkovic et al., 1991), may explain why ERK1/2 phosphorylation is PKC dependent (Ely et al., 2011). However, ERK1/2 solely may not be a true frequency decoder due to different kinetics of activation and deactivation (Armstrong et al., 2010). The mathematical model postulated by this group (Armstrong et al., 2011) implies cooperative action of two or more pathways leading to two or more transcription factors that coordinately induce the expression of their target gene.

The p38 branch of the MAPK pathway is also activated by GnRH via PKC and calcium (Mugami et al., 2017; Roberson et al., 1999). p38 can be activated by both extracellular calcium influx and from intracellular stores in α T3 cells, while in L β T2 p38 activation requires calcium from intracellular stores. Pulse dependency of p38 activation has not been analyzed. p38 activation is involved in FSH β induction by GnRH in L β T2 cells (Bonfil et al., 2004; Coss et al., 2007; Ely et al., 2011; Wang et al., 2008), but it is not critical in vivo in rats (Haisenleder et al., 2008). Calcium from intracellular stores leads to JNK activation in α T3 cells (Kraus et al., 2001; Mulvaney and Roberson, 2000). Whether these distinct calcium sources: extracellular, leading to ERK1/2 activation and intracellular, leading to JNK activation play a role in more mature L β T2 cells or primary gonadotrope as well is yet to be determined. GnRH activation of ERK1/2, p38 and JNK involves several PKC isoforms in both α T3 and L β T2 cells, primarily PKC α , PKC ϵ and PKC δ (Dobkin-Bekman et al., 2010; Kratzmeier et al., 1996; Liu et al., 2003; Mugami et al., 2017). However, a specific isoform which selectively activates ERK1/2 but not JNK has not been identified, to indicate the isoform necessary for voltage gated calcium channel activation and extracellular calcium influx. Furthermore, understanding of the significance of the studies delineating G α s versus G α q/11 signaling modules is complicated by findings that these same PKC ϵ and PKC δ that are normally activated by G α q/11 maybe be involved in cAMP accumulation, in addition to cAMP accumulation by G α s activation (Lariviere et al., 2007). More importantly, PKC isoform-specific roles in gonadotropin gene expression were not examined.

While ERK1/2 may be activated longer or more highly with low pulse frequency of GnRH, regulation of MAPK activity occurs at the level of dephosphorylation as well. MAPKs are deactivated by a family of MAPK phosphatases (MKPs), which is a subfamily of a large dual-specificity phosphatase (DUSP) family. Studies analyzing MKPs/DUSPs are complicated due to multiple members that are ubiquitously expressed and lack substrate specificity. MKP1/DUSP1 and MKP2/DUSP4 are induced by GnRH in α T3 cells (Roberson et al., 1995; Zhang and Roberson, 2006). MKP1/DUSP1 is also phosphorylated by GnRH and, since localized specifically in the nucleus, it may provide rapid cessation of nuclear ERK1/2 activity specifically and maintain sensitivity to GnRH stimulation. (Nguyen et al., 2010). MKP1/DUSP1 is induced with high pulse frequency of GnRH (Purwana et al., 2011), which may explain sustained ERK1/2 activation with low pulse frequency.

In summary, GnRH-stimulated calcium increase from two different sources, intracellular stores or extracellular influx, activates different downstream signaling pathways. Since one of them, intracellular via IP₃, can be desensitized, calcium sources may lead to differential regulation of downstream gene targets; however this remains to be examined. Receptor concentration may also play a role. Fast pulse frequency leads to higher receptor concentration, which is correlated to higher LH β transcription. Therefore, at low pulse frequency, GnRHR couples to G α_q /11 and activates CamKII that induces both LH β and FSH β transcription. Activation of PKC additionally causes influx of extracellular calcium, which leads to sustained activation of ERK1/2 and preferential induction of FSH β to a higher level than LH β . At high pulse frequency, GnRHR number is augmented, and this high receptor number allows for coupling to both G α_s and G α_q /11, and while G α_q /11 desensitizes, sustained signaling via G α_s module induces higher expression of LH β via unknown mechanism (Fig. 2).

3. GnRH regulation of LH β and FSH β Transcription

The specific gonadotropin β subunits are expressed at low basal levels and are differentially induced through pulses of GnRH (Bedecarrats and Kaiser, 2003; Haisenleder et al., 1991; Shupnik, 1996; Weiss et al., 1990). GnRH regulates expression of the specific β subunits through immediate-early genes. Early growth response 1 (EGR1) is an intermediary gene that regulates GnRH induction of LH β , while activating protein 1 (AP1) transcription factor, comprised of a heterodimer of FOS and JUN isoforms, mediates induction of FSH β by GnRH (Fig. 3).

3.1. GnRH regulation of LH β transcription

Immediate-early gene that regulates LH β induction by GnRH is EGR1 and is the only transcription factor binding the LH β promoter that is dramatically induced by GnRH (Tremblay and Drouin, 1999). EGR1 is the most highly induced gene in the gonadotrope by GnRH in L β T2 model of mature gonadotrope when analyzed by microarrays (Kakar et al., 2003; Lawson et al., 2007; Wurmbach et al., 2001). EGR1 is also induced in rat primary pituitary cells following GnRH stimulation (Kucka et al., 2013). Confirming a role of EGR1 and lack of compensation by other family members (EGR2, EGR3, and EGR4), the EGR1 null mice are infertile and lack expression of the LH β gene (Lee et al., 1996). EGR1 also

Author Manuscript

Author Manuscript

Author Manuscript

organizes a complex on the LH β promoter to bring forth coordinated expression of this gene. In the proximal LH β promoter region, two EGR1 sites are located adjacent to sites for steroidogenic factor 1 (SF1; or NR5A1) on either side of a homeodomain element (HD) bound by Ptx1 (Halvorson et al., 1996; Quirk et al., 2001). This location allows for a complex formation that integrates pituitary specific (SF1), basal (Ptx1) and GnRH-induced (EGR1) expression. Synergistic interaction between SF1, EGR1 and Ptx1 is essential for maximal LH β gene expression (Dorn et al., 1999; Tremblay and Drouin, 1999). EGR1-organized complex interacts with the Sp1 sites in the distal region (Kaiser et al., 1998), in a spatially-dependent manner (Kaiser et al., 2000; Weck et al., 2000). Interaction is likely facilitated by the co-activators, SNURF (Curtin et al., 2004) and p300 (Mouillet et al., 2004). β -catenin may play a role in GnRH induction of LH β via interaction with this complex through SF-1 (Gardner et al., 2007; Salisbury et al., 2007). However, recent studies addressing a role of β -catenin in the gonadotrope *in vivo* demonstrated a lack of any role for β -catenin in LH β expression and LH levels in the circulation, in either gonadotrope specific deletion of β -catenin, or gonadotrope specific stabilization of β -catenin, which increases its levels in the cell (Boerboom et al., 2015).

Mechanisms involved in the responsiveness to pulsatile GnRH treatment and frequency-dependent expression of the LH β gene have also been analyzed. Namely, maximal LH β expression is achieved with high pulse frequency characteristic of pre-ovulatory period. This allows for a high level of LH β gene expression, which then dimerizes with non-limiting α -subunit and during preovulatory surge of GnRH increases rapidly LH levels in the circulation that are necessary for ovulation. EGR1 intermediate gene which induces LH β in response to GnRH, similarly to other immediate early genes, is not only rapidly activated, but is rapidly degraded as well. The short half-life allows for strong temporal regulation of gene targets. To address the importance of pulsatile GnRH treatment, it was postulated that proteasome function is critical for LH β responsiveness to GnRH treatment, since GnRH leads to ubiquitination of EGR1 and cycling of this factor on and off of the LH β promoter (Walsh and Shupnik, 2009). Cycling of EGR1 may explain the importance of pulsatile GnRH compared to tonic treatment that inhibits gonadotropin expression.

Explanation of higher induction with high pulse frequency of GnRH characteristic of the preovulatory proestrus period, may lie in the differential expression of intermediates. High pulse frequency upregulates SF1 mRNA in rat pituitary cells (Burger et al., 2011), which, although SF1 induction is much lower than of EGR1, combined with already high basal level, is sufficient to form a complex with highly induced EGR1 with any frequency. Another mechanisms is differential regulation of positive and negative regulators of transcription. Sustained induction of EGR1 specific repressor proteins from the Nab family at low pulse frequency may explain the necessity for high pulse frequency of GnRH for LH β responsiveness (Lawson et al., 2007). High pulse frequency of GnRH may be necessary to allow for accumulation of EGR1 protein in order to overcome repression by Nabs.

As mentioned previously, β -subunits are expressed at a very low basal level which may indicate active repression until a signal is applied that would de-repress the promoter at the appropriate time. Repressors that bind inactive promoters normally function via recruitment of histone deacetylase (HDAC) enzymes that remove active histone acetylation marks and

prevent transcription. GnRH stimulus leads to changes in histone acetylation in the α T3 model of immature gonadotrope that normally do not express β -subunits (Lim et al., 2007). This may designate developmental activation of β -subunit transcription via chromatin remodeling, which occurs after gonadotropes acquire responsiveness to GnRH. Developmental and prepubertal regulation of gonadotropin subunits by GnRH is supported by a recent thorough study that demonstrated strong correlation and synchronization of GnRHR expression with gonadotropin β -subunit expression in female and in male rats (Bjelobaba et al., 2015). We recently demonstrated that GnRH stimulus elicits an increase in histone acetylation in mature L β T2 cells (Jonak et al., 2017), which may suggest that chromatin remodeling occurs with each pulse. Remodeling of chromatin that modulates gene expression occurs upon recruitment of the coactivator proteins to the DNA. EGR1 interacts with SF1 and Ptx1 and forms the active transcriptional complex that recruits coactivator CBP. Coactivator CBP has histone acetyltransferase activity (HAT), which acetylates histones and leads to activation of transcription (Korzus et al., 1998). CBP is indeed necessary for LH β expression and normal fertility in vivo (Miller et al., 2012). This scenario points to rapid changes in repression and activation of the promoter via transcriptional repressors, chromatin remodeling and changes in histone acetylation (Fig. 3).

Signaling pathways triggered by GnRH receptor binding, discussed above, that activate gonadotropin genes transcription also induce transcription of immediate early genes. As demonstrated previously, GnRH induction of both gonadotropin β subunits is dependent on CamKII activation (Burger et al., 2008; Haisenleder et al., 2003a). We determined that GnRH induction of both EGR1 and cFOS immediate early genes occurs through an increase in intracellular calcium concentration and activation of CamKII. CamKII activity leads to phosphorylation of the SRF transcription factor (Ely et al., 2011). The EGR1 promoter contains four SRF binding sites (Witham et al., 2013), compared to the single site in the cFOS promoter that is responsive to GnRH (Ely et al., 2011). Higher induction of EGR1 than cFOS in response to GnRH may stem from four response elements in the promoter. Thus, this mechanism may be responsible for higher responsiveness of LH β gene, a target of EGR1, to GnRH during the high pulse frequency preovulatory period, since at that time interpulse interval is not lengthy enough to allow immediate early gene turnover and protein degradation (Fig. 4).

3.2. GnRH regulation of FSH β transcription

GnRH induces FSH β by upregulating immediate early genes that belong to AP1 family of transcription factors, which subsequently bind the only site in the FSH β promoter that conveys GnRH responsiveness (Coss et al., 2004). AP1 is a heterodimer of FOS and JUN transcription factors, which are activated rapidly and transiently by various stimuli and growth factors in numerous cell types (Karin et al., 1997). GnRH induces cFOS, FosB, cJUN and JunB but not JunD, and the combination of these factors bind the AP1 site in FSH β promoter (Coss et al., 2004). Transcriptome analysis revealed that AP1 members are highly induced by GnRH in L β T2 cells (Wurmbach et al., 2001) and primary rat gonadotrope cells (Yuen et al., 2012). Similarly to EGR1, AP1 organizes a complex to integrate basal expression through interaction with NFY, and activin induction through interaction with activin-regulated SMAD3 and FOXL2 (Coss et al., 2007; Roybal et al.,

2014; Wang et al., 2008). cFOS deficient mice survive until early adulthood indicating possible compensation by FosB for survival, but are infertile, hypogonadotropic, and have similar ovarian phenotype as FSH β knockouts, indicating necessity of cFOS for reproductive function (Xie et al., 2015). β -catenin may play a role in GnRH-induction of FSH β expression, via its cofactor, breast cancer metastasis-suppressor 1-like (Brms1L, (Wang et al., 2013). In vivo studies using mice with gonadotrope specific knockdown of β -catenin, mice failed to show the role of β -catenin in FSH β induction. On the other hand, male mice with β -catenin mutant that prevents its degradation and allows for accumulation in the gonadotrope, exhibit reduced FSH β mRNA expression and diminished FSH levels due to an increase in pituitary follistatin expression (Boerboom et al., 2015).

Since AP1 family members are induced by a variety of stimuli in other cell types, but are sufficient for activation of FSH β transcription, it is critical to understand specificity of GnRH induction of immediate early genes. Analysis of pathways and transcription factors activated solely by GnRH to induce cJUN in gonadotrope cells, provide critical insight into GnRH regulation of its target gene, the gonadotropin subunit FSH β . We determined that GnRH is the sole regulator of cJUN in the mature L β T2 cells, which occurs through GnRH phosphorylation of ATF2 transcription factor by p38, the same pathway responsible for GnRH induction of cJUN and FSH β in L β T2 cells (Bonfil et al., 2004; Lindaman et al., 2013). Of four p38 isoforms, p38 β and p38 δ are sufficient for cJUN induction, demonstrating additional specificity of signaling via this branch (Lindaman et al., 2013). On the contrary, FSH β induction by GnRH in primary rat pituitary cells does not involve p38 (Haisenleder et al., 2008), likely because cJUN component of the AP1 complex is present at some basal level in the gonadotrope cells, which is sufficient for heterodimerization with tightly regulated and highly induced cFOS. To induce cFOS, GnRH uses neurotransmitter-like signaling and the CamKII pathway, rather than a growth factor signaling via MAPK. This provides the specificity of GnRH signaling to regulate FSH β gene expression in the gonadotrope cells (Ely et al., 2011; Stojilkovic et al., 2010). The same signaling pathways are necessary for induction of both cFOS and FSH β , and cFOS induction precedes that of FSH β . As with EGR1 described above, GnRH induces cFOS primarily through an increase in intracellular calcium concentration and activation of CamKII, which then leads to phosphorylation of the SRF transcription factor. The necessity for CamKII in rat pituitary cultures for both LH β and FSH β (Haisenleder et al., 2003a), may stem from CamKII involvement in both EGR1 and cFOS induction (Ely et al., 2011; Witham et al., 2013). Although receptors for several growth factors that induce cFOS in other cell lines are expressed in the gonadotrope, these factors are not sufficient to induce cFOS, and therefore are not involved in the regulation of FSH β , likely because they do not cause an increase in intracellular calcium concentration that is necessary to activate CamKII and though it SRF. Presence of the single SRF site in the cFOS promoter, compared to four in the EGR1 promoter, may explain lower responsiveness of the cFOS gene than the EGR1 gene to GnRH, either when induction was analyzed using a reporter (Ely et al., 2011) or endogenous mRNA level (Wurmbach et al., 2001). Downstream, since both EGR1 and cFOS are induced rapidly and transiently, it may lead to lower FSH β , a target of cFOS, than LH β , a target of EGR1, at high pulse frequency. However, induction of immediate early genes does not explain higher induction of FSH β at low pulse frequency.

To answer dependence of FSH β expression on pulsatile GnRH, several studies analyzed repressors that may explain frequency responsiveness of the FSH β promoter. Skil and Tgif1 repressors that downregulate FSH β induction are induced at the higher level with high pulse frequency, thus antagonizing cFOS (Mistry et al., 2011). Another repressor, ICER is also induced by GnRH at the high pulse frequency and binds to the AP1 site in the FSH β promoter, thus repressing FSH β expression at high pulse frequency (Ciccone et al., 2010). Importance of either Skil, Tgif1 or ICER has not been examined in vivo. We recently identified Jun-dimerizing protein 2 (JDP2) as another repressor that binds the same AP1 site and interacts with cJUN thereby displacing cFOS (Jonak et al., 2017). JDP2 belongs to the AP1 superfamily since it contains homologous DNA binding domain and the leucine zipper through which it heterodimerizes with cJUN, but lacks a transactivation domain to form a transcriptionally active complex (Aronheim et al., 1997). JDP2 is expressed at the high basal level and is not highly induced transcriptionally, but its repressive function may be regulated by phosphorylation (Katz et al., 2001). However, frequency dependence of JDP2 phosphorylation status remains to be explored. Consistent with a JDP2 role in FSH β repression in the cells, female mice deficient in JDP2 exhibit higher expression of FSH β , augmented FSH concentration and consequently higher estrogen in the circulation, and increased recruitment of ovarian follicles to the growing pool. JDP2 nulls also exhibit early puberty, increased litter size and cessation of reproductive function at younger age, implicating higher FSH β expression to premature reproductive senescence (Jonak et al., 2017). Cessations of reproduction 70 days earlier in JDP2 null mouse is proportional to 8-year earlier menopause in women over respective reproductive lifespans. A complete lack of gonadotropins results in infertility in affected individuals. However, larger number of individuals have pathologies such as POF, with hormone levels slightly outside the physiological range. Since 70% of POF causes are idiopathic, repressors may provide candidates for investigation in affected individuals.

Repressors may underlie the lower levels of FSH β at high pulse frequency, but these findings still do not explain higher levels of FSH β than LH β at low pulse frequency. We uncovered a mechanism that may function during a period of low GnRH pulse frequency to extend the half-life of the cFOS protein specifically, allowing sufficient accumulation of this factor to increase transcription of its target gene FSH β (Reddy et al., 2013). This leads to selective induction of FSH and its secondary rise, which is necessary for follicular growth. GnRH induces rapidly both EGR1 and cFOS, but specifically decreases the rate of cFOS degradation, by triggering cFOS protein phosphorylation through the ERK1/2 pathway (Fig. 4). Role of ERK1/2 in GnRH induction of FSH β that has been shown in rat primary pituitary cells (Haisenleder et al., 1998), may stem not from direct role of ERK1/2 on FSH β transcription, but from increased half-life of the cFOS protein than induces FSH β (Reddy et al., 2013). Preferential activation of ERK1/2 in vivo in GnRH treated rats with low pulse frequency, also confirms its role in FSH β specifically (Burger et al., 2008). Phosphorylation extends the half-life of cFOS protein, which is normally rapidly degraded. Confirming the role of phosphorylation in promoting increased protein activity, cFOS mutant which cannot be phosphorylated by GnRH has a shorter half-life and induces lower expression of the FSH β promoter than wild-type, phosphorylatable cFOS (Reddy et al., 2013). Our study stressed the role of post-translational modification in the regulation of gonadotropin gene

expression by GnRH. This modification may cause higher levels of FSH during the time of low GnRH pulse frequency to stimulate folliculogenesis.

4. Crosstalk of GnRH with other hormones that regulate gonadotropins

Another possible mechanism of differential regulation is interaction of GnRH signaling pathways and other hormones that regulate gonadotropin levels. Gonadal steroid hormones regulate expression of gonadotropin β subunits and pituitary responsiveness to GnRH, by modulating the levels of induction by GnRH-induced signaling. Pituitary responsiveness to GnRH is regulated by paracrine factors expressed in the pituitary itself.

4.1. Gonadal hormones

Inhibin and activin are members of TGF β superfamily. Regulation of FSH secretion is their initially identified and thus most studied role (Makanji et al., 2014). Inhibin decreases FSH secretion, while activin stimulates it. Gonads are the primary source of circulating inhibin, since after gonadectomy inhibin levels are undetectable. Activin, although originally purified from follicular fluid, has low circulating level and primarily functions in the paracrine or autocrine manner. Thus, activin role in gonadotropin synthesis will be discussed below. Inhibin is a heterodimer of α and β subunits, whereas activin is $\beta\beta$ homodimers. Thus, they share β subunits and availability of α subunit and the preferential assembly of $\alpha\beta$ heterodimers over $\beta\beta$ dimers may be one mechanism of inhibin to antagonize activin. More likely, functional antagonism of activin signaling by inhibin is achieved through the competitive binding to the common receptor on the cell surface via the shared β subunit. Subsequently, inhibin binding prevents recruitment of activin signaling receptor, but instead recruits betaglycan, which prevents signaling (Chapman and Woodruff, 2003). Therefore, primary role of inhibin in FSH synthesis is to antagonize autocrine activin function.

Although gonadal steroid hormone feedback primarily targets the hypothalamus, modulation of pituitary responsiveness by steroid hormones has been reported. Estrogen likely does not directly regulate FSH β gene expression, but induction of LH β gene expression by estrogen in gonadotropes has been reported (Shupnik et al., 1989a; Shupnik et al., 1989b). The mechanism may involve either ER binding to an imperfect estrogen response element in the distal rat LH β promoter (Shupnik and Rosenzweig, 1991; Shupnik et al., 1989b); or ER recruitment to the LH β promoter through interactions with SF1 and Ptx1 (Luo et al., 2005). Estrogen has been shown to potentiate GnRH induction of LH β indirectly by enhancing the expression of EGR1 and suppressing the expression of ZEB, a transcriptional repressor that binds the rat LH β promoter (Kowase et al., 2007).

Progesterone and testosterone decrease LH β expression and increase FSH β induction (Thackray et al., 2009; Thackray et al., 2006; Thackray and Mellon, 2008). Testosterone and other androgens have been reported to repress LH β gene expression, in both castrated, GnRH antagonist-treated rats (Burger et al., 2004; Wierman and Wang, 1990) and primary pituitary cell culture (Winters et al., 1992). Androgens also reduced GnRH-stimulated LH secretion via alterations of calcium signaling (Ortmann et al., 1998). Androgen treatment suppressed GnRH-induced transcription of the rat and ovine LH β promoters, through a

direct protein-protein interaction between the androgen receptor (AR) and Sp1 (Curtin et al., 2001), or between AR and SF1 (Jorgensen and Nilson, 2001), respectively.

Both basal and GnRH-induced FSH β mRNA levels increased in rats in vivo and in primary cells treated with progesterone (Attardi and Fitzgerald, 1990; Kerrigan et al., 1993). Progesterone antagonists blocked FSH secretion and FSH β expression during the preovulatory FSH surge (Ringstrom et al., 1997) and the secondary FSH rise (Knox and Schwartz, 1992; Szabo et al., 1998). Like progesterone, androgens upregulate FSH β mRNA levels. Castrated GnRH antagonist-treated rats exhibited a selective increase in FSH β mRNA upon treatment with testosterone (Burger et al., 2004; Dalkin et al., 1992; Paul et al., 1990; Wierman and Wang, 1990). Suppression of LH β and induction of FSH β by progesterone and androgens may contribute to differential regulation of β -subunits.

4.2. Autocrine and paracrine regulation by pituitary hormones

Gonadotropes comprise 5–15% of the cells in the anterior pituitary (Ooi et al., 2004). Anterior pituitary contains four other types of secretory cells including thyrotropes, somatotropes, lactotropes and corticotropes, as well as folliculostellate cells. Many hormones secreted from these cells have paracrine effects on gonadotrope cells (reviewed in (Denef, 2008). Folliculostellate cells have been reported to rapidly overgrow other pituitary cells in culture (Kawakami et al., 2002) and can impact the experimental outcome since they produce paracrine factors such as follistatin and PACAP (reviewed in (Winters and Moore, 2007, 2011).

PACAP was initially identified as hypothalamic factor that increases pituitary adenylate cyclase signaling as its name suggests. PACAP is also secreted in the pituitary by gonadotropes and folliculostellate cells and works in the autocrine or paracrine manner to modulate response to GnRH (Counis et al., 2007). GnRH increases the expression of PACAP receptor in the gonadotrope and in turn, PACAP potentiates the effect of GnRH on β -subunit expression likely via cAMP pathway crosstalk (Purwana et al., 2010). Some inconsistencies that exist in the literature regarding a role of PACAP in the gonadotrope, maybe due to the level or PACAP receptor expression in different models (Fowkes et al., 2003b). Alternatively, they can possibly be explained by crosstalk of pathways, since with cotreatment the pathway may be augmented, while pretreatment may lead to downregulation and inability to respond to a subsequent signal (Fowkes et al., 2003a; Larivière et al., 2008). The fact that pretreatment with either PACAP or GnRH diminishes the effect of the other, can likely be explained by downregulation. This hypothesis is strengthened by a study in which transgenic male mice with permanently elevated PACAP levels have delayed puberty and decreased gonadotropin levels (Moore et al., 2012).

Prolactin from the lactotropes diminishes pituitary responsiveness to GnRH by decreasing LH secretion (De Paul et al., 2000; Tortonese et al., 1998). Lactotropes, which comprise 20% of pituitary cells, form a honey-comb pattern in the anterior pituitary surrounding the gonadotropes, allowing for a close contact and paracrine function by prolactin (Noda et al., 2001). Gonadotropes express prolactin receptor (Tortonese et al., 1998) and prolactin has been shown to diminish the LH secretion and expression in response to GnRH, with no effect on basal levels (Garcia et al., 1985; Hodson et al., 2012; Smith, 1982).

Follistatin is a functional antagonist of TGF β superfamily that binds activin member of the family with the highest activity (Makanji et al., 2014). It binds activin at the several residues that are necessary for activin binding to its receptor, thus inhibiting its function. Follistatin, synthesized by folliculostellate cells and gonadotropes themselves, represses FSH synthesis, likely by antagonizing endogenous, autocrine activin (Bilezikjian et al., 2004). Follistatin also diminishes GnRH induction of the FSH β gene (Coss et al., 2007), again likely due to synergistic effect of autocrine activin with GnRH as discussed below.

Activin stimulates secretion of FSH and activates FSH β transcription. Activin is synthesized by gonadotropes themselves and functions in the autocrine manner, while inhibin and follistatin, as discussed above, regulate FSH primarily by antagonizing activin. The acute GnRH-independent increase in FSH β transcription after ovariectomy is due to a lack of inhibin to antagonize autocrine activin (Burger et al., 2001). This is true as well in younger women with decreased ovarian function or menopausal women that experience an increase primarily in FSH, as discussed as the beginning of this manuscript. Since FSH is mostly constitutively secreted, transcriptional regulation may be a critical regulatory step of FSH levels by activin. Activin functions via SMAD transcription factors and induces FSH β transcription via SMAD3 phosphorylation. Activin-regulated SMAD3 and 4 are necessary for FSH β expression in vivo (Fortin et al., 2014a; Fortin et al., 2014b). Bernard lab (Lamba et al., 2009) and our lab (Corpuz et al., 2010) identified FOXL2 as a critical component in activin induction of the FSH β gene. FOXL2 overexpression in gonadotropes can potentiate activin induction and its expression can confer activin responsiveness to FSH β in heterologous cells where this promoter is normally refractory to activin induction. Mice deficient in FOXL2, both whole body knockout or gonadotrope-specific knockdown, have lower FSH β level (Justice et al., 2011; Roybal et al., 2014; Tran et al., 2013)). Various FOXL2 mutations have been identified in patients with an autosomal dominant disorder, BPES, which is characterized by distinctive eyelid abnormalities and POF (Crisponi et al., 2001; De Baere et al., 2001; De Baere et al., 2002). Considering that POF is caused by mutations of the FOXL2 gene, which induces FSH β , and that this reproductive defect is associated with higher FSH levels in the circulation, it is intriguing to postulate again a role of FSH in the development of POF.

Activin also increases pituitary responsiveness to GnRH (Coss et al., 2007; Coss et al., 2010; Fortin et al., 2015; Gregory and Kaiser, 2004). Cotreatment of activin and GnRH results in the synergistic induction of FSH β . p38 appears to be a point of crosstalk between different signaling pathways since it is more highly activated after GnRH and activin co-treatment of L β T2 cells than GnRH treatment alone (Coss et al., 2007; Zhang et al., 2006). Synergy is accomplished through a complex formation of GnRH-induced AP1 and activin-regulated SMAD3. We showed that FOXL2, with activin-phosphorylated SMAD3, also plays a role in synergistic induction of FSH β by GnRH and activin through interactions with the cJUN component of the AP1 complex that is necessary for GnRH responsiveness (Roybal et al., 2014). Another member of TGF β family, GDF9, previously considered specifically expressed in oocytes is expressed in primary pituitary cultures and L β T2 cells. GDF9 is implicated in induction of FSH β and synergism with GnRH, which may also contribute to differential expression (Choi et al., 2014). GDF9 also activates SMAD transcription factors, but whether the same complex plays a role in synergism with GnRH remains to be

determined. The synergistic induction is specific for FSH β gene and may contribute to the differential expression of the gonadotropin subunits. It may occur when ovarian inhibin levels decrease, allowing activin, or GDF9, to synergize with GnRH, at a time coinciding with the secondary peak of FSH, which is necessary for follicular development (Woodruff et al., 1996).

5. Conclusion

GnRH differentially regulates gonadotropin β -subunits expression, the limiting components of the mature hormone. Each pulse of GnRH induces a new wave of transcription of immediate early genes, EGR1 that leads to induction of LH β , and cFOS (and more stable, but less regulated, cJUN) that leads to induction of FSH β (Fig. 3 and Fig. 4). Both EGR1 and cFOS have labile mRNAs and equally unstable proteins, exhibiting rapid turnover and degradation to allow for a tight temporal regulation of gene targets. With increasing pulse frequencies of GnRH in the preovulatory period, EGR1 and cFOS accumulate to overcome repressors (Nabs and JDP2 for example, respectively) and allow expression of their respective targets. Enabled by EGR1 and cFOS complex organizing facilities, expression of β -subunits is augmented. Due to four response elements in its promoter, EGR1 is induced at the higher level than cFOS leading to higher induction of LH β than FSH β with fast pulse frequency. Responsiveness to GnRH fast pulse frequency is likely maintained by increased receptors concentration that allows for an engagement of G α s that is less susceptible to desensitization, in addition to G α q/11. Both LH β and FSH β heterodimerize with non-limiting α GSU and are secreted under pulsatile GnRH, increasing their concentration in the circulation. During the low pulse frequency, specific phosphorylation of the cFOS protein via ERK1/2 extends its half-life, while EGR1 is degraded during each interpulse interval preventing accumulation. Under the endogenous activin tone, activin-regulated factors, such as SMAD3 and FOXL2, interact with cFOS/AP1 complex to synergistically induce FSH β . FSH β dimerizes with abundant α GSU to form a mature hormone that is subsequently constitutively secreted to stimulate folliculogenesis. This mechanism may explain preferential induction of FSH β during the low pulse frequency of GnRH.

The conundrum of how a single hormone, GnRH, interacting with a single receptor can differentially activate gonadotropin β -subunit genes in the same cell has not been answered completely, but significant progress has been made in our understanding of molecular mechanisms that regulate differential expression. Given that pathologies such as PCOS or POF may arise due to aberrations in the narrowly controlled levels, further insight is crucial to our understanding of regulation of reproduction at the level of the gonadotrope. Future studies may lead to the development of potential therapeutics for pathologies caused by dysregulation of the tight differential regulation.

Acknowledgments

This work was supported by National Institute of Health Grant R01 HD057549 to DC. The author thanks Nancy M. Lainez and Carrie R. Jonak for editorial comments.

Abbreviations

AP1	activator protein 1
AR	androgen receptor
CamKII	calcium calmodulin kinase II
ChIP	chromatin immunoprecipitation
DAG	diacyl glycerol
EGR1	early growth response 1
ER	estrogen receptor
ERK	extracellular signal regulated kinase
FSH	follicle-stimulating hormone
GWAS	genome-wide association study
GSU	glycoprotein subunit
GnRH	gonadotropin-releasing hormone
HD	homeodomain element
IP3	inositol triphosphate
JNK	Jun N-terminal kinase
LH	luteinizing hormone
HPG	hypothalamic-pituitary-gonadal
MAPK	mitogen-activated protein kinase
MKP	MAP kinase phosphatase
PACAP	pituitary adenylate cyclase-activating polypeptide
PCOS	polycystic ovary syndrome
POF	premature ovarian failure
POI	primary ovarian insufficiency
PKC	protein kinase C
SBE	smad-binding element
SRF	serum response factor
SF1	steroidogenic factor 1

Literature

- Abbott DH, Barnett DK, Levine JE, Padmanabhan V, Dumesic DA, Jacoris S, Tarantal AF. 2008; Endocrine antecedents of Polycystic Ovary Syndrome in fetal and infant prenatally androgenized female rhesus monkeys. *Biol Reprod.* 79:154–163. [PubMed: 18385445]
- Abbott DH, Nicol LE, Levine JE, Xu N, Goodarzi MO, Dumesic DA. 2013; Nonhuman primate models of polycystic ovary syndrome. *Mol Cell Endocrinol.* 373:21–28. [PubMed: 23370180]
- Alarid ET, Holley S, Hayakawa M, Mellon PL. 1998; Discrete stages of anterior pituitary differentiation recapitulated in immortalized cell lines. *Mol Cell Endocrinol.* 140:25–30. [PubMed: 9722164]
- Alarid ET, Windle JJ, Whyte DB, Mellon PL. 1996; Immortalization of pituitary cells at discrete stages of development by directed oncogenesis in transgenic mice. *Development.* 122:3319–3329. [PubMed: 8898243]
- Armstrong SP, Caunt CJ, Finch AR, McArdle CA. 2011; Using automated imaging to interrogate gonadotrophin-releasing hormone receptor trafficking and function. *Mol Cell Endocrinol.* 331:194–204. [PubMed: 20688134]
- Armstrong SP, Caunt CJ, Fowkes RC, Tsaneva-Atanasova K, McArdle CA. 2009; Pulsatile and sustained gonadotropin-releasing hormone (GnRH) receptor signaling: does the Ca²⁺/NFAT signaling pathway decode GnRH pulse frequency? *The Journal of biological chemistry.* 284:35746–35757. [PubMed: 19858197]
- Armstrong SP, Caunt CJ, Fowkes RC, Tsaneva-Atanasova K, McArdle CA. 2010; Pulsatile and sustained gonadotropin-releasing hormone (GnRH) receptor signaling: does the ERK signaling pathway decode GnRH pulse frequency? *The Journal of biological chemistry.* 285:24360–24371. [PubMed: 20507982]
- Aronheim A, Zandi E, Hennemann H, Elledge SJ, Karin M. 1997; Isolation of an AP-1 repressor by a novel method for detecting protein-protein interactions. *Molecular and cellular biology.* 17:3094–3102. [PubMed: 9154808]
- Attardi B, Fitzgerald T. 1990; Effects of progesterone on the estradiol-induced follicle-stimulating hormone (FSH) surge and FSH beta messenger ribonucleic acid in the rat. *Endocrinology.* 126:2281–2287. [PubMed: 2109682]
- Bedecarrats GY, Kaiser UB. 2003; Differential regulation of gonadotropin subunit gene promoter activity by pulsatile gonadotropin-releasing hormone (GnRH) in perfused L beta T2 cells: role of GnRH receptor concentration. *Endocrinology.* 144:1802–1811. [PubMed: 12697686]
- Belchetz PE, Plant TM, Nakai Y, Keogh EJ, Knobil E. 1978; Hypophysial responses to continuous and intermittent delivery of hypophyseal gonadotropin-releasing hormone. *Science.* 202:631–633. [PubMed: 100883]
- Besecke LM, Guendner MJ, Sluss PA, Polak AG, Woodruff TK, Jameson JL, Bauer-Dantoin AC, Weiss J. 1997; Pituitary follistatin regulates activin-mediated production of follicle-stimulating hormone during the rat estrous cycle. *Endocrinology.* 138:2841–2848. [PubMed: 9202226]
- Bilezikjian LM, Blount AL, Leal AM, Donaldson CJ, Fischer WH, Vale WW. 2004; Autocrine/paracrine regulation of pituitary function by activin, inhibin and follistatin. *Mol Cell Endocrinol.* 225:29–36. [PubMed: 15451565]
- Binder AK, Grammer JC, Herndon MK, Stanton JD, Nilson JH. 2012; GnRH regulation of Jun and Atf3 requires calcium, calcineurin, and NFAT. *Mol Endocrinol.* 26:873–886. [PubMed: 22446101]
- Bjelobaba I, Janjic MM, Kucka M, Stojilkovic SS. 2015; Cell Type-Specific Sexual Dimorphism in Rat Pituitary Gene Expression During Maturation. *Biol Reprod.* 93:21. [PubMed: 26063874]
- Bliss SP, Navratil AM, Breed M, Skinner DC, Clay CM, Roberson MS. 2007; Signaling complexes associated with the type I gonadotropin-releasing hormone (GnRH) receptor: colocalization of extracellularly regulated kinase 2 and GnRH receptor within membrane rafts. *Mol Endocrinol.* 21:538–549. [PubMed: 17068198]
- Blomenrohr M, Heding A, Sellar R, Leurs R, Bogerd J, Eidne KA, Willars GB. 1999; Pivotal role for the cytoplasmic carboxyl-terminal tail of a nonmammalian gonadotropin-releasing hormone receptor in cell surface expression, ligand binding, and receptor phosphorylation and internalization. *Mol Pharmacol.* 56:1229–1237. [PubMed: 10570050]

- Boerboom D, Kumar V, Boyer A, Wang Y, Lambrot R, Zhou X, Rico C, Boehm U, Paquet M, Céleste C, et al. 2015; β -Catenin Stabilization in Gonadotropes Impairs FSH Synthesis in Male Mice In Vivo. *Endocrinology*. 156:323–333. [PubMed: 25343272]
- Bonfil D, Chuderland D, Kraus S, Shahbazian D, Friedberg I, Seger R, Naor Z. 2004; Extracellular signal-regulated kinase, Jun N-terminal kinase, p38, and c-Src are involved in gonadotropin-releasing hormone-stimulated activity of the glycoprotein hormone follicle-stimulating hormone beta-subunit promoter. *Endocrinology*. 145:2228–2244. [PubMed: 14736735]
- Burger LL, Dalkin AC, Aylor KW, Haisenleder DJ, Marshall JC. 2002; GnRH pulse frequency modulation of gonadotropin subunit gene transcription in normal gonadotropes-assessment by primary transcript assay provides evidence for roles of GnRH and follistatin. *Endocrinology*. 143:3243–3249. [PubMed: 12193535]
- Burger LL, Dalkin AC, Aylor KW, Workman LJ, Haisenleder DJ, Marshall JC. 2001; Regulation of gonadotropin subunit transcription after ovariectomy in the rat: measurement of subunit primary transcripts reveals differential roles of GnRH and inhibin. *Endocrinology*. 142:3435–3442. [PubMed: 11459788]
- Burger LL, Haisenleder DJ, Aylor KW, Dalkin AC, Prendergast KA, Marshall JC. 2004; Regulation of luteinizing hormone-beta and follicle-stimulating hormone (FSH)-beta gene transcription by androgens: testosterone directly stimulates FSH-beta transcription independent from its role on follistatin gene expression. *Endocrinology*. 145:71–78. [PubMed: 14512439]
- Burger LL, Haisenleder DJ, Aylor KW, Marshall JC. 2008; Regulation of intracellular signaling cascades by GNRH pulse frequency in the rat pituitary: roles for CaMK II, ERK, and JNK activation. *Biol Reprod*. 79:947–953. [PubMed: 18716286]
- Burger LL, Haisenleder DJ, Aylor KW, Marshall JC. 2009 Regulation of Lhb and Egr1 Gene Expression by GNRH Pulses in Rat Pituitaries Is Both c-Jun N-Terminal Kinase (JNK)- and Extracellular Signal-Regulated Kinase (ERK)-Dependent. *Biol Reprod*.
- Burger LL, Haisenleder DJ, Marshall JC. 2011; GnRH pulse frequency differentially regulates steroidogenic factor 1 (SF1), dosage-sensitive sex reversal-AHC critical region on the X chromosome gene 1 (DAX1), and serum response factor (SRF): potential mechanism for GnRH pulse frequency regulation of LH beta transcription in the rat. *Endocrine*. 39:212–219. [PubMed: 21409515]
- Burrin JM, Jameson JL. 1989; Regulation of transfected glycoprotein hormone α -gene expression in primary pituitary cell cultures. *Mol Endocrinol*. 3:1643–1651. [PubMed: 2558299]
- Butcher RL, Collins WE, Fugo NW. 1974; Plasma concentration of LH, FSH, prolactin, progesterone and estradiol-17beta throughout the 4-day estrous cycle of the rat. *Endocrinology*. 94:1704–1708. [PubMed: 4857496]
- Chand AL, Harrison CA, Shelling AN. 2010; Inhibin and premature ovarian failure. *Hum Reprod Update*. 16:39–50. [PubMed: 19752047]
- Chang RJ. 2007; The reproductive phenotype in polycystic ovary syndrome. *Nat Clin Pract Endocrinol Metab*. 3:688–695. [PubMed: 17893687]
- Chapman SC, Woodruff TK. 2003; Betaglycan localization in the female rat pituitary: implications for the regulation of follicle-stimulating hormone by inhibin. *Endocrinology*. 144:5640–5649. [PubMed: 14500575]
- Chin, W, Boime, I. Glycoprotein Hormones. Norwell, MA: Serono Symposia USA; 1990.
- Choi SG, Jia J, Pfeffer RL, Sealfon SC. 2012; G proteins and autocrine signaling differentially regulate gonadotropin subunit expression in pituitary gonadotrope. *The Journal of biological chemistry*. 287:21550–21560. [PubMed: 22549790]
- Choi SG, Wang Q, Jia J, Pincas H, Turgeon JL, Sealfon SC. 2014; Growth differentiation factor 9 (GDF9) forms an incoherent feed-forward loop modulating follicle-stimulating hormone beta-subunit (FSHbeta) gene expression. *The Journal of biological chemistry*. 289:16164–16175. [PubMed: 24778184]
- Ciccone NA, Xu S, Lacza CT, Carroll RS, Kaiser UB. 2010; Frequency-dependent regulation of follicle-stimulating hormone beta by pulsatile gonadotropin-releasing hormone is mediated by functional antagonism of bZIP transcription factors. *Molecular and cellular biology*. 30:1028–1040. [PubMed: 20008557]

- Corpuz PS, Lindaman LL, Mellon PL, Coss D. 2010; FoxL2 is required for activin induction of the mouse and human follicle-stimulating hormone β -subunit genes. *Mol Endocrinol.* 24:1037–1051. [PubMed: 20233786]
- Coss D, Hand CM, Yaphockun KK, Ely HA, Mellon PL. 2007; p38 mitogen-activated kinase is critical for synergistic induction of the FSH beta gene by gonadotropin-releasing hormone and activin through augmentation of c-Fos induction and Smad phosphorylation. *Mol Endocrinol.* 21:3071–3086. [PubMed: 17823303]
- Coss D, Jacobs SB, Bender CE, Mellon PL. 2004; A novel AP-1 site is critical for maximal induction of the follicle-stimulating hormone beta gene by gonadotropin-releasing hormone. *J Biol Chem.* 279:152–162. [PubMed: 14570911]
- Coss D, Mellon PL, Thackray VG. 2010; A FoxL in the Smad house: activin regulation of FSH. *Trends Endocrinol Metab.* 21:562–568. [PubMed: 20598900]
- Counis R, Laverriere JN, Garrel-Lazayres G, Cohen-Tannoudji J, Lariviere S, Bleux C, Magre S. 2007; What is the role of PACAP in gonadotrope function? *Peptides.* 28:1797–1804. [PubMed: 17601637]
- Crisponi L, Deiana M, Loi A, Chiappe F, Uda M, Amati P, Bisceglia L, Zelante L, Nagaraja R, Porcu S, et al. 2001; The putative forkhead transcription factor FOXL2 is mutated in blepharophimosis/ptosis/epicanthus inversus syndrome. *Nat Genet.* 27:159–166. [PubMed: 11175783]
- Culler MD, Negro-Vilar A. 1987; Pulsatile follicle-stimulating hormone secretion is independent of luteinizing hormone-releasing hormone (LHRH): pulsatile replacement of LHRH bioactivity in LHRH-immunoneutralized rats. *Endocrinology.* 120:2011–2021. [PubMed: 2436893]
- Curtin D, Ferris HA, Hakli M, Gibson M, Janne OA, Palvimo JJ, Shupnik MA. 2004; Small nuclear RING finger protein stimulates the rat luteinizing hormone-beta promoter by interacting with Sp1 and steroidogenic factor-1 and protects from androgen suppression. *Mol Endocrinol.* 18:1263–1276. [PubMed: 14988433]
- Curtin D, Jenkins S, Farmer N, Anderson AC, Haisenleder DJ, Rissman E, Wilson EM, Shupnik MA. 2001; Androgen suppression of GnRH-stimulated rat LHbeta gene transcription occurs through Sp1 sites in the distal GnRH-responsive promoter region. *Mol Endocrinol.* 15:1906–1917. [PubMed: 11682622]
- Dalkin AC, Burger LL, Aylor KW, Haisenleder DJ, Workman LJ, Cho S, Marshall JC. 2001; Regulation of gonadotropin subunit gene transcription by gonadotropin-releasing hormone: measurement of primary transcript ribonucleic acids by quantitative reverse transcription-polymerase chain reaction assays. *Endocrinology.* 142:139–146. [PubMed: 11145576]
- Dalkin AC, Haisenleder DJ, Ortolano GA, Ellis TR, Marshall JC. 1989; The frequency of gonadotropin-releasing-hormone stimulation differentially regulates gonadotropin subunit messenger ribonucleic acid expression. *Endocrinology.* 125:917–924. [PubMed: 2502379]
- Dalkin AC, Paul SJ, Haisenleder DJ, Ortolano GA, Yasin M, Marshall JC. 1992; Gonadal steroids effect similar regulation of gonadotrophin subunit mRNA expression in both male and female rats. *J Endocrinol.* 132:39–45. [PubMed: 1737957]
- Damian-Matsumura P, Zaga V, Maldonado A, Sanchez-Hernandez C, Timossi C, Ulloa-Aguirre A. 1999; Oestrogens regulate pituitary alpha2,3-sialyltransferase messenger ribonucleic acid levels in the female rat. *J Mol Endocrinol.* 23:153–165. [PubMed: 10514553]
- De Baere E, Dixon MJ, Small KW, Jabs EW, Leroy BP, Devriendt K, Gillerot Y, Mortier G, Meire F, Van Maldergem L, et al. 2001; Spectrum of FOXL2 gene mutations in blepharophimosis-ptosis-epicanthus inversus (BPES) families demonstrates a genotype-phenotype correlation. *Hum Mol Genet.* 10:1591–1600. [PubMed: 11468277]
- De Baere E, Lemerrier B, Christin-Maitre S, Durval D, Messiaen L, Fellous M, Veitia R. 2002; FOXL2 mutation screening in a large panel of POF patients and XX males. *J Med Genet.* 39:e43. [PubMed: 12161610]
- De Biasi SN, Perez GT, Apfelbaum LI, Apfelbaum ME. 2002; Pituitary luteinizing hormone microheterogeneity in the afternoon of proestrus in rats: some new insights. *Horm Res.* 58:8–15. [PubMed: 12169775]

- De Leo V, Musacchio MC, Cappelli V, Massaro MG, Morgante G, Petraglia F. 2016; Genetic, hormonal and metabolic aspects of PCOS: an update. *Reproductive biology and endocrinology: RB&E*. 14:38. [PubMed: 27423183]
- De Paul LA, Bonaterra M, Aoki A, Torres IA. 2000; Cellular and functional interactions between gonadotrophs and lactotrophs in pituitary cell cultures. *Medical Electron Microscopy*. 33:231–240. [PubMed: 11810480]
- Denef C. 2008; Paracrinicity: the story of 30 years of cellular pituitary crosstalk. *J Neuroendocrinol*. 20:1–70.
- DePaolo LV, Hirshfield AN, Anderson LD, Barraclough CA, Channing CP. 1979; Suppression of pituitary secretion of follicle-stimulating hormone by porcine follicular fluid during pro-oestrus and oestrus in the rat: effects on gonadotrophin and steroid secretion, follicular development and ovulation during the following cycle. *J Endocrinol*. 83:355–368. [PubMed: 575386]
- Dobkin-Bekman M, Rahamin-Ben Navi L, Shterental B, Sviridonov L, Przedeki F, Naidich-Exler M, Brodie C, Seger R, Naor Z. 2010; Differential role of PKC isoforms in GnRH and phorbol 12-myristate 13-acetate activation of extracellular signal-regulated kinase and Jun N-terminal kinase. *Endocrinology*. 151:4894–4907. [PubMed: 20810567]
- Dorn C, Ou Q, Svaren J, Crawford PA, Sadovsky Y. 1999; Activation of luteinizing hormone beta gene by gonadotropin-releasing hormone requires the synergy of early growth response-1 and steroidogenic factor-1. *J Biol Chem*. 274:13870–13876. [PubMed: 10318795]
- El-Shal AS, Zidan HE, Rashad NM, Abdelaziz AM, Harira MM. 2016; Association between genes encoding components of the Leutinizing hormone/Luteinizing hormone–choriogonadotrophin receptor pathway and polycystic ovary syndrome in Egyptian women. *IUBMB Life*. 68:23–36. [PubMed: 26662070]
- Ely HA, Mellon PL, Coss D. 2011; GnRH Induces the c-Fos gene via phosphorylation of SRF by the calcium/calmodulin kinase II pathway. *Mol Endocrinol*. 25:669–680. [PubMed: 21292826]
- Fares F. 2006; The role of O-linked and N-linked oligosaccharides on the structure-function of glycoprotein hormones: development of agonists and antagonists. *Biochimica et biophysica acta*. 1760:560–567. [PubMed: 16527410]
- Fortin J, Boehm U, Deng CX, Treier M, Bernard DJ. 2014a; Follicle-stimulating hormone synthesis and fertility depend on SMAD4 and FOXL2. *The FASEB Journal*. 28:3396–3410. [PubMed: 24739304]
- Fortin J, Boehm U, Weinstein MB, Graff JM, Bernard DJ. 2014b; Follicle-stimulating hormone synthesis and fertility are intact in mice lacking SMAD3 DNA binding activity and SMAD2 in gonadotrope cells. *FASEB journal: official publication of the Federation of American Societies for Experimental Biology*. 28:1474–1485. [PubMed: 24308975]
- Fortin J, Ongaro L, Li Y, Tran S, Lamba P, Wang Y, Zhou X, Bernard DJ. 2015; Minireview: Activin Signaling in Gonadotropes: What Does the FOX say. to the SMAD? *Mol Endocrinol*. 29:963–977. [PubMed: 25942106]
- Fowkes RC, Desclozeaux M, Patel MV, Aylwin SJ, King P, Ingraham HA, Burrin JM. 2003a; Steroidogenic factor-1 and the gonadotrope-specific element enhance basal and pituitary adenylate cyclase-activating polypeptide-stimulated transcription of the human glycoprotein hormone alpha-subunit gene in gonadotropes. *Mol Endocrinol*. 17:2177–2188. [PubMed: 12920232]
- Fowkes RC, Sidhu KK, Sosabowski JK, King P, Burrin JM. 2003b; Absence of pituitary adenylate cyclase-activating polypeptide-stimulated transcription of the human glycoprotein alpha-subunit gene in LbetaT2 gonadotrophs reveals disrupted cAMP-mediated gene transcription. *J Mol Endocrinol*. 31:263–278. [PubMed: 14519095]
- Garcia A, Herbon L, Barkan A, Papavasiliou S, Marshall JC. 1985; Hyperprolactinemia inhibits gonadotropin-releasing hormone (GnRH) stimulation of the number of pituitary GnRH receptors. *Endocrinology*. 117:954–959. [PubMed: 2990879]
- Gardner S, Maudsley S, Millar RP, Pawson AJ. 2007; Nuclear stabilization of beta-catenin and inactivation of glycogen synthase kinase-3beta by gonadotropin-releasing hormone: targeting Wnt signaling in the pituitary gonadotrope. *Mol Endocrinol*. 21:3028–3038. [PubMed: 17717075]
- Goodarzi MO, Dumesic DA, Chazenbalk G, Azziz R. 2011; Polycystic ovary syndrome: etiology, pathogenesis and diagnosis. *Nat Rev Endocrinol*. 7:219–231. [PubMed: 21263450]

- Goswami D, Conway GS. 2007; Premature Ovarian Failure. *Hormone Research*. 68:196–202. [PubMed: 17495481]
- Graham KE, Nusser KD, Low MJ. 1999; L β T2 gonadotroph cells secrete follicle stimulating hormone (FSH) in response to activin A. *J Endocrinol*. 162:R1–R5. [PubMed: 10467239]
- Green ED, Boime I, Baenziger JU. 1986; Differential processing of Asn-linked oligosaccharides on pituitary glycoprotein hormones: implications for biologic function. *Mol Cell Biochem*. 72:81–100. [PubMed: 3102943]
- Gregory SJ, Kaiser UB. 2004; Regulation of gonadotropins by inhibin and activin. *Semin Reprod Med*. 22:253–267. [PubMed: 15319828]
- Haisenleder D, Dalkin A, Ortolano G, Marshall J, Shupnik M. 1991; A pulsatile gonadotropin-releasing hormone stimulus is required to increase transcription of the gonadotropin subunit genes: Evidence for differential regulation of transcription by pulse frequency *in vivo*. *Endocrinology*. 128:509–517. [PubMed: 1702704]
- Haisenleder DJ, Burger LL, Aylor KW, Dalkin AC, Marshall JC. 2003a; Gonadotropin-releasing hormone stimulation of gonadotropin subunit transcription: evidence for the involvement of calcium/calmodulin-dependent kinase II (Ca/CAMK II) activation in rat pituitaries. *Endocrinology*. 144:2768–2774. [PubMed: 12810529]
- Haisenleder DJ, Burger LL, Walsh HE, Stevens J, Aylor KW, Shupnik MA, Marshall JC. 2008; Pulsatile gonadotropin-releasing hormone stimulation of gonadotropin subunit transcription in rat pituitaries: evidence for the involvement of Jun N-terminal kinase but not p38. *Endocrinology*. 149:139–145. [PubMed: 17932215]
- Haisenleder DJ, Cox ME, Parsons SJ, Marshall JC. 1998; Gonadotropin-releasing hormone pulses are required to maintain activation of mitogen-activated protein kinase: role in stimulation of gonadotrope gene expression. *Endocrinology*. 139:3104–3111. [PubMed: 9645682]
- Haisenleder DJ, Ferris HA, Shupnik MA. 2003b; The calcium component of gonadotropin-releasing hormone-stimulated luteinizing hormone subunit gene transcription is mediated by calcium/calmodulin-dependent protein kinase type II. *Endocrinology*. 144:2409–2416. [PubMed: 12746302]
- Haisenleder DJ, Katt JA, Ortolano GA, el-Gewely MR, Duncan JA, Dee C, Marshall JC. 1988; Influence of gonadotropin-releasing hormone pulse amplitude, frequency, and treatment duration on the regulation of luteinizing hormone (LH) subunit messenger ribonucleic acids and LH secretion. *Mol Endocrinol*. 2:338–343. [PubMed: 2454398]
- Haisenleder DJ, Workman LJ, Burger LL, Aylor KW, Dalkin AC, Marshall JC. 2001; Gonadotropin subunit transcriptional responses to calcium signals in the rat: evidence for regulation by pulse frequency. *Biol Reprod*. 65:1789–1793. [PubMed: 11717142]
- Hall JE, Taylor AE, Hayes FJ, Crowley WF Jr. 1998; Insights into hypothalamic-pituitary dysfunction in polycystic ovary syndrome. *J Endocrinol Invest*. 21:602–6011. [PubMed: 9856414]
- Halvorson LM, Kaiser UB, Chin WW. 1996; Stimulation of luteinizing hormone beta gene promoter activity by the orphan nuclear receptor, steroidogenic factor-1. *J Biol Chem*. 271:6645–6650. [PubMed: 8636081]
- Halvorson LM, Weiss J, Bauer-Dantoin AC, Jameson JL. 1994; Dynamic regulation of pituitary follistatin messenger ribonucleic acids during the rat estrous cycle. *Endocrinology*. 134:1247–1253. [PubMed: 8119165]
- Harris GC, Levine JE. 2003; Pubertal acceleration of pulsatile gonadotropin-releasing hormone release in male rats as revealed by microdialysis. *Endocrinology*. 144:163–171. [PubMed: 12488342]
- Hayes MG, Urbanek M, Ehrmann DA, Armstrong LL, Lee JY, Sisk R, Karaderi T, Barber TM, McCarthy MI, Franks S, et al. 2015; Genome-wide association of polycystic ovary syndrome implicates alterations in gonadotropin secretion in European ancestry populations. *Nat Commun*. 6:7502. [PubMed: 26284813]
- He C, Kraft P, Chasman D, Buring J, Chen C, Hankinson S, Paré G, Chanock S, Ridker P, Hunter D. 2010; A large-scale candidate gene association study of age at menarche and age at natural menopause. *Hum Genet*. 128:515–527. [PubMed: 20734064]
- Hislop JN, Everest HM, Flynn A, Harding T, Uney JB, Troskie BE, Millar RP, McArdle CA. 2001; Differential internalization of mammalian and non-mammalian gonadotropin-releasing hormone

- receptors. Uncoupling of dynamin-dependent internalization from mitogen-activated protein kinase signaling. *J Biol Chem.* 276:39685–39694. [PubMed: 11495905]
- Hoak DC, Schwartz NB. 1980; Blockade of recruitment of ovarian follicles by suppression of the secondary surge of follicle-stimulating hormone with porcine follicular field. *Proc Natl Acad Sci USA.* 77:4953–4956. [PubMed: 6776533]
- Hodson DJ, Henderson HL, Townsend J, Tortonesi DJ. 2012; Photoperiodic modulation of the suppressive actions of prolactin and dopamine on the pituitary gonadotropin responses to gonadotropin-releasing hormone in sheep. *Biol Reprod.* 86:122. [PubMed: 22302689]
- Huhtaniemi I. 2006; Mutations along the pituitary-gonadal axis affecting sexual maturation: novel information from transgenic and knockout mice. *Mol Cell Endocrinol.* 254–255:84–90.
- Huhtaniemi I, Ahtiainen P, Pakarainen T, Rulli SB, Zhang FP, Poutanen M. 2006; Genetically modified mouse models in studies of luteinising hormone action. *Mol Cell Endocrinol.* 252:126–135. [PubMed: 16675102]
- Jonak CR, Lainez NM, Roybal LL, Williamson AD, Coss D. 2017; c-JUN Dimerization Protein 2 (JDP2) Is a Transcriptional Repressor of Follicle-stimulating Hormone β (FSH β) and Is Required for Preventing Premature Reproductive Senescence in Female Mice. *Journal of Biological Chemistry.* 292:2646–2659. [PubMed: 28007961]
- Jones, RE. *Human Reproductive Biology.* 2. San Diego, CA: Academic Press; 1997.
- Jorgensen JS, Nilson JH. 2001; AR suppresses transcription of the LH β subunit by interacting with steroidogenic factor-1. *Mol Endocrinol.* 15:1505–1516. [PubMed: 11518799]
- Justice NJ, Blount AL, Pelosi E, Schlessinger D, Vale W, Bilezikjian LM. 2011; Impaired FSH β Expression in the Pituitaries of Foxl2 Mutant Animals. *Mol Endocrinol.* 25:1404–1415. [PubMed: 21700720]
- Kaiser UB, Conn PM, Chin WW. 1997a; Studies of gonadotropin-releasing hormone (GnRH) action using GnRH receptor-expressing pituitary cell lines. *Endocr Rev.* 18:46–70. [PubMed: 9034786]
- Kaiser UB, Halvorson LM, Chen MT. 2000; Sp1, steroidogenic factor 1 (SF-1), and early growth response protein 1 (egr-1) binding sites form a tripartite gonadotropin-releasing hormone response element in the rat luteinizing hormone-beta gene promoter: an integral role for SF-1. *Mol Endocrinol.* 14:1235–1245. [PubMed: 10935547]
- Kaiser UB, Jakubowiak A, Steinberger A, Chin WW. 1997b; Differential effects of gonadotropin-releasing hormone (GnRH) pulse frequency on gonadotropin subunit and GnRH receptor messenger ribonucleic acid levels in vitro. *Endocrinology.* 138:1224–1231. [PubMed: 9048630]
- Kaiser UB, Sabbagh E, Chen MT, Chin WW, Saunders BD. 1998; Sp1 binds to the rat luteinizing hormone beta (LH β) gene promoter and mediates gonadotropin-releasing hormone-stimulated expression of the LH β subunit gene. *J Biol Chem.* 273:12943–12951. [PubMed: 9582327]
- Kakar SS, Winters SJ, Zacharias W, Miller DM, Flynn S. 2003; Identification of distinct gene expression profiles associated with treatment of L β T2 cells with gonadotropin-releasing hormone agonist using microarray analysis. *Gene.* 308:67–77. [PubMed: 12711391]
- Kanasaki H, Bedecarrats GY, Kam KY, Xu S, Kaiser UB. 2005; Gonadotropin-releasing hormone pulse frequency-dependent activation of extracellular signal-regulated kinase pathways in perfused L β T2 cells. *Endocrinology.* 146:5503–5513. [PubMed: 16141398]
- Karin M, Liu Z, Zandi E. 1997; AP-1 function and regulation. *Curr Opin Cell Biol.* 9:240–246. [PubMed: 9069263]
- Katz S, Heinrich R, Aronheim A. 2001; The AP-1 repressor, JDP2, is a bona fide substrate for the c-Jun N-terminal kinase. *FEBS Lett.* 506:196–200. [PubMed: 11602244]
- Kauffman AS, Thackray VG, Ryan GE, Tolson KP, Glidewell-Kenney CA, Semaan SJ, Poling MC, Iwata N, Breen KM, Duleba AJ, et al. 2015; A Novel Letrozole Model Recapitulates Both the Reproductive and Metabolic Phenotypes of Polycystic Ovary Syndrome in Female Mice. *Biology of Reproduction.* 93:69–12. 61–12. [PubMed: 26203175]
- Kawakami S, Fujii Y, Okada Y, Winters SJ. 2002; Paracrine regulation of FSH by follistatin in folliculostellate cell- enriched primate pituitary cell cultures. *Endocrinology.* 143:2250–2258. [PubMed: 12021189]

- Kerrigan JR, Dalkin AC, Haisenleder DJ, Yasin M, Marshall JC. 1993; Failure of gonadotropin-releasing hormone (GnRH) pulses to increase luteinizing hormone beta messenger ribonucleic acid in GnRH-deficient female rats. *Endocrinology*. 133:2071–2079. [PubMed: 8404655]
- Knox KL, Schwartz NB. 1992; RU486 blocks the secondary surge of follicle-stimulating hormone in the rat without blocking the drop in serum inhibin. *Biol Reprod*. 46:220–225. [PubMed: 1536897]
- Korzus E, Torchia J, Rose DW, Xu L, Kurokawa R, McInerney EM, Mullen TM, Glass CK, Rosenfeld MG. 1998; Transcription factor-specific requirements for coactivators and their acetyltransferase functions. *Science*. 279:703–707. [PubMed: 9445475]
- Kowase T, Walsh HE, Darling DS, Shupnik MA. 2007; Estrogen enhances gonadotropin-releasing hormone-stimulated transcription of the luteinizing hormone subunit promoters via altered expression of stimulatory and suppressive transcription factors. *Endocrinology*. 148:6083–6091. [PubMed: 17823254]
- Kratzmeier M, Poch A, Mukhopadhyay AK, McArdle CA. 1996; Selective translocation of non-conventional protein kinase C isoenzymes by gonadotropin-releasing hormone (GnRH) in the gonadotrope-derived alpha T3-1 cell line. *Mol Cell Endocrinol*. 118:103–111. [PubMed: 8735596]
- Kraus S, Naor Z, Seger R. 2001; Intracellular signaling pathways mediated by the gonadotropin-releasing hormone (GnRH) receptor. *Arch Med Res*. 32:499–509. [PubMed: 11750725]
- Kucka M, Bjelobaba I, Clokie SJ, Klein DC, Stojilkovic SS. 2013; Female-specific induction of rat pituitary dentin matrix protein-1 by GnRH. *Mol Endocrinol*. 27:1840–1855. [PubMed: 24085820]
- Kumar TR, Palapattu G, Wang P, Woodruff TK, Boime I, Byrne MC, Matzuk MM. 1999; Transgenic Models to Study Gonadotropin Function: The Role of Follicle-Stimulating Hormone in Gonadal Growth and Tumorigenesis. *Molecular Endocrinology*. 13:851–865. [PubMed: 10379885]
- Kumar TR, Wang Y, Lu N, Matzuk MM. 1997; Follicle stimulating hormone is required for ovarian follicle maturation but not male fertility. *Nat Genet*. 15:201–204. [PubMed: 9020850]
- Lamba P, Fortin J, Tran S, Wang Y, Bernard DJ. 2009; A novel role for the forkhead transcription factor FOXL2 in activin A-regulated follicle-stimulating hormone beta subunit transcription. *Mol Endocrinol*. 23:1001–1013. [PubMed: 19324968]
- Lamminen T, Jokinen P, Jiang M, Pakarinen P, Simonsen H, Huhtaniemi I. 2005; Human FSH beta subunit gene is highly conserved. *Mol Hum Reprod*. 11:601–605. [PubMed: 16100240]
- Larivière S, Garrel-Lazayres G, Simon V, Shintani N, Baba A, Counis R, Cohen-Tannoudji J. 2008; Gonadotropin-Releasing Hormone Inhibits Pituitary Adenylyl Cyclase-Activating Polypeptide Coupling to 3',5'-Cyclic Adenosine-5'-Monophosphate Pathway in LβT2 Gonadotrope Cells through Novel Protein Kinase C Isoforms and Phosphorylation of Pituitary Adenylyl Cyclase-Activating Polypeptide Type I Receptor. *Endocrinology*. 149:6389–6398. [PubMed: 18755795]
- Lariviere S, Garrel G, Simon V, Soh JW, Laverriere JN, Counis R, Cohen-Tannoudji J. 2007; Gonadotropin-releasing hormone couples to 3',5'-cyclic adenosine-5'-monophosphate pathway through novel protein kinase Cdelta and -epsilon in LbetaT2 gonadotrope cells. *Endocrinology*. 148:1099–1107. [PubMed: 17185372]
- Laven JS. 2016; Primary Ovarian Insufficiency. *Seminars in reproductive medicine*. 34:230–234. [PubMed: 27513024]
- Lawson MA, Tsutsumi R, Zhang H, Talukdar I, Butler BK, Santos SJ, Mellon PL, Webster NJ. 2007; Pulse sensitivity of the luteinizing hormone beta promoter is determined by a negative feedback loop Involving early growth response-1 and Ngfi-A binding protein 1 and 2. *Mol Endocrinol*. 21:1175–1191. [PubMed: 17299135]
- Lee SL, Sadovsky Y, Swirnoff AH, Polish JA, Goda P, Gavriliina G, Milbrandt J. 1996; Luteinizing hormone deficiency and female infertility in mice lacking the transcription factor NGFI-A (Egr-1). *Science*. 273:1219–1221. [PubMed: 8703054]
- Levine JE, Duffy MT. 1988; Simultaneous measurement of luteinizing hormone (LH)-releasing hormone, LH, and follicle-stimulating hormone release in intact and short-term castrate rats. *Endocrinology*. 122:2211–2221. [PubMed: 3129282]

- Levine JE, Norman RL, Gliessman PM, Oyama TT, Bangsberg DR, Spies HG. 1985; In vivo gonadotropin-releasing hormone release and serum luteinizing hormone measurements in ovariectomized, estrogen-treated rhesus macaques. *Endocrinology*. 117:711–721. [PubMed: 3893989]
- Levine JE, Pau KY, Ramirez VD, Jackson GL. 1982; Simultaneous measurement of luteinizing hormone-releasing hormone and luteinizing hormone release in unanesthetized, ovariectomized sheep. *Endocrinology*. 111:1449–1455. [PubMed: 6751794]
- Levine JE, Ramirez VD. 1982; Luteinizing hormone-releasing hormone release during the rat estrous cycle and after ovariectomy, as estimated with push-pull cannulae. *Endocrinology*. 111:1439–1448. [PubMed: 6751793]
- Lim S, Luo M, Koh M, Yang M, bin Abdul Kadir MN, Tan JH, Ye Z, Wang W, Melamed P. 2007; Distinct mechanisms involving diverse histone deacetylases repress expression of the two gonadotropin beta-subunit genes in immature gonadotropes, and their actions are overcome by gonadotropin-releasing hormone. *Mol Cell Biol*. 27:4105–4120. [PubMed: 17371839]
- Lindaman LL, Yeh DM, Xie C, Breen KM, Coss D. 2013; Phosphorylation of ATF2 and interaction with NFY induces c-Jun in the gonadotrope. *Mol Cell Endocrinol*. 365:316–326. [PubMed: 23178797]
- Liu F, Austin DA, Mellon PL, Olefsky JM, Webster NJ. 2002; GnRH activates ERK1/2 leading to the induction of c-fos and LHbeta protein expression in LbetaT2 cells. *Mol Endocrinol*. 16:419–434. [PubMed: 11875099]
- Liu F, Austin DA, Webster NJ. 2003; Gonadotropin-releasing hormone-desensitized LbetaT2 gonadotrope cells are refractory to acute protein kinase C, cyclic AMP, and calcium-dependent signaling. *Endocrinology*. 144:4354–4365. [PubMed: 12960037]
- Lu C, Lin L, Tan H, Wu H, Sherman SL, Gao F, Jin P, Chen D. 2012; Fragile X premutation RNA is sufficient to cause primary ovarian insufficiency in mice. *Human Molecular Genetics*. 21:5039–5047. [PubMed: 22914733]
- Luo M, Koh M, Feng J, Wu Q, Melamed P. 2005; Cross talk in hormonally regulated gene transcription through induction of estrogen receptor ubiquitylation. *Mol Cell Biol*. 25:7386–7398. [PubMed: 16055746]
- Ma X, Dong Y, Matzuk MM, Kumar TR. 2004; Targeted disruption of luteinizing hormone beta-subunit leads to hypogonadism, defects in gonadal steroidogenesis, and infertility. *Proc Natl Acad Sci USA*. 101:17294–17299. [PubMed: 15569941]
- Macpherson JN, Murray A. 2016 Development of Genetic Testing for Fragile X Syndrome and Associated Disorders, and Estimates of the Prevalence of FMR1 Expansion Mutations. *Genes*. :7.
- Makanji Y, Zhu J, Mishra R, Holmquist C, Wong WPS, Schwartz NB, Mayo KE, Woodruff TK. 2014; Inhibin at 90: From Discovery to Clinical Application, a Historical Review. *Endocrine reviews*. 35:747–794. [PubMed: 25051334]
- Manikkam M, Thompson RC, Herkimer C, Welch KB, Flak J, Karsch FJ, Padmanabhan V. 2008; Developmental programming: impact of prenatal testosterone excess on pre- and postnatal gonadotropin regulation in sheep. *Biol Reprod*. 78:648–660. [PubMed: 18094361]
- Mason AJ, Hayflick JS, Zoeller RT, Young WS, Phillips HS, Nikolics K, Seeburg PH. 1986; A deletion truncating the gonadotropin-releasing hormone gene is responsible for hypogonadism in the hpg mouse. *Science*. 234:1366–1371. [PubMed: 3024317]
- Mbarek H, Steinberg S, Nyholt Dale R, Gordon Scott D, Miller Michael B, McRae Allan F, Hottenga Jouke J, Day Felix R, Willemsen G, de Geus Eco J, et al. 2016; Identification of Common Genetic Variants Influencing Spontaneous Dizygotic Twinning and Female Fertility. *The American Journal of Human Genetics*. 98:1–11.
- McArdle CA, Davidson JS, Willars GB. 1999; The tail of the gonadotrophin-releasing hormone receptor: desensitization at, and distal to, G protein-coupled receptors. *Mol Cell Endocrinol*. 151:129–136. [PubMed: 10411327]
- McArdle CA, Franklin J, Green L, Hislop JN. 2002; Signalling, cycling and desensitisation of gonadotrophin-releasing hormone receptors. *J Endocrinol*. 173:1–11. [PubMed: 11927379]
- McArdle CA, Willars GB, Fowkes RC, Nahorski SR, Davidson JS, Forrest-Owen W. 1996; Desensitization of gonadotropin-releasing hormone action in alphaT3-1 cells due to uncoupling

of inositol 1,4,5-trisphosphate generation and Ca^{2+} mobilization. *J Biol Chem.* 271:23711–23717. [PubMed: 8798594]

Mihalca R, Fica S. 2014; The impact of obesity on the male reproductive axis. *Journal of medicine and life.* 7:296–300. [PubMed: 25408743]

Miller RS, Wolfe A, He L, Radovick S, Wondisford FE. 2012; CREB Binding Protein (CBP) Activation Is Required for Luteinizing Hormone Beta Expression and Normal Fertility in Mice. *Molecular and cellular biology.* 32:2349–2358. [PubMed: 22508984]

Mistry DS, Tsutsumi R, Fernandez M, Sharma S, Cardenas SA, Lawson MA, Webster NJ. 2011; Gonadotropin-releasing hormone pulse sensitivity of follicle-stimulating hormone-beta gene is mediated by differential expression of positive regulatory activator protein 1 factors and corepressors SKIL and TGIF1. *Mol Endocrinol.* 25:1387–1403. [PubMed: 21659477]

Moore AM, Prescott M, Marshall CJ, Yip SH, Campbell RE. 2015; Enhancement of a robust arcuate GABAergic input to gonadotropin-releasing hormone neurons in a model of polycystic ovarian syndrome. *Proceedings of the National Academy of Sciences of the United States of America.* 112:596–601. [PubMed: 25550522]

Moore JP Jr, Yang RQ, Winters SJ. 2012; Targeted pituitary overexpression of pituitary adenylate-cyclase activating polypeptide alters postnatal sexual maturation in male mice. *Endocrinology.* 153:1421–1434. [PubMed: 22315445]

Morales AJ, Laughlin GA, Butzow T, Maheshwari H, Baumann G, Yen SS. 1996; Insulin, somatotrophic, and luteinizing hormone axes in lean and obese women with polycystic ovary syndrome: common and distinct features. *J Clin Endocrinol Metab.* 81:2854–2864. [PubMed: 8768842]

Mouillet JF, Sonnenberg-Hirche C, Yan X, Sadovsky Y. 2004; p300 regulates the synergy of steroidogenic factor-1 and early growth response-1 in activating luteinizing hormone-beta subunit gene. *J Biol Chem.* 279:7832–7839. [PubMed: 14681221]

Mugami S, Kravchook S, Rahamim-Ben Navi L, Seger R, Naor Z. 2017; Differential roles of PKC isoforms (PKCs) and Ca^{2+} in GnRH and phorbol 12-myristate 13-acetate (PMA) stimulation of p38MAPK phosphorylation in immortalized gonadotrope cells. *Mol Cell Endocrinol.* 439:141–154. [PubMed: 27810601]

Mulvaney JM, Roberson MS. 2000; Divergent signaling pathways requiring discrete calcium signals mediate concurrent activation of two mitogen-activated protein kinases by gonadotropin-releasing hormone. *J Biol Chem.* 275:14182–14189. [PubMed: 10799494]

Mulvaney JM, Zhang T, Fewtrell C, Roberson MS. 1999; Calcium influx through L-type channels is required for selective activation of extracellular signal-regulated kinase by gonadotropin-releasing hormone. *J Biol Chem.* 274:29796–29804. [PubMed: 10514457]

Naor Z, Benard O, Seger R. 2000; Activation of MAPK cascades by G-protein-coupled receptors: the case of gonadotropin-releasing hormone receptor. *Trends Endocrinol Metab.* 11:91–99. [PubMed: 10707049]

Naor Z, Huhtaniemi I. 2013; Interactions of the GnRH receptor with heterotrimeric G proteins. *Front Neuroendocrinol.* 34:88–94. [PubMed: 23200814]

Navratil AM, Bliss SP, Berghorn KA, Haughian JM, Farmerie TA, Graham JK, Clay CM, Roberson MS. 2003; Constitutive localization of the gonadotropin-releasing hormone (GnRH) receptor to low density membrane microdomains is necessary for GnRH signaling to ERK. *The Journal of biological chemistry.* 278:31593–31602. [PubMed: 12791688]

Neill JD. 2002; GnRH and GnRH receptor genes in the human genome. *Endocrinology.* 143:737–743. [PubMed: 11861490]

Nguyen KA, Intrigo RE, Upadhyay HC, Santos SJ, Webster NJ, Lawson MA. 2010; Modulation of gonadotropin-releasing hormone-induced extracellular signal-regulated kinase activation by dual-specificity protein phosphatase 1 in LbetaT2 gonadotropes. *Endocrinology.* 151:4882–4893. [PubMed: 20685880]

Noda T, Kaidzu S, Kikuchi M, Yashiro T. 2001; Topographic Affinities of Hormone-Producing Cells in the Rat Anterior Pituitary Gland. *ACTA HISTOCHEMICA ET CYTOCHEMICA.* 34:313–319.

- Ooi GT, Tawadros N, Escalona RM. 2004; Pituitary cell lines and their endocrine applications. *Mol Cell Endocrinol.* 228:1–21. [PubMed: 15541569]
- Ortmann O, Tomic M, Weiss JM, Diedrich K, Stojilkovic SS. 1998; Dual action of androgen on calcium signaling and luteinizing hormone secretion in pituitary gonadotrophs. *Cell Calcium.* 24:223–231. [PubMed: 9883276]
- Ortolano GA, Haisenleder DJ, Dalkin AC, Iliff-Sizemore SA, Landefeld TD, Maurer RA, Marshall JC. 1988; Follicle-stimulating hormone beta subunit messenger ribonucleic acid concentrations during the rat estrous cycle. *Endocrinology.* 123:2946–2948. [PubMed: 3143544]
- Pangaribuan B, Yusuf I, Mansyur M, Wijaya A. 2011; Serum adiponectin and resistin in relation to insulin resistance and markers of hyperandrogenism in lean and obese women with polycystic ovary syndrome. *Therapeutic advances in endocrinology and metabolism.* 2:235–245. [PubMed: 23148188]
- Papavasiliou SS, Zmeili S, Khoury S, Landefeld TD, Chin WW, Marshall JC. 1986; Gonadotropin-releasing hormone differentially regulates expression of the genes for luteinizing hormone alpha and beta subunits in male rats. *Proc Natl Acad Sci USA.* 83:4026–4029. [PubMed: 3012543]
- Pasquali R, Vicennati V, Gambineri A. 2002; Adrenal and gonadal function in obesity. *Journal of Endocrinological Investigation.* 25:893–898. [PubMed: 12508952]
- Paul SJ, Ortolano GA, Haisenleder DJ, Stewart JM, Shupnik MA, Marshall JC. 1990; Gonadotropin subunit messenger RNA concentrations after blockade of gonadotropin-releasing hormone action: testosterone selectively increases follicle-stimulating hormone beta-subunit messenger RNA by posttranscriptional mechanisms. *Mol Endocrinol.* 4:1943–1955. [PubMed: 1707128]
- Pawson AJ, Faccenda E, Maudsley S, Lu ZL, Naor Z, Millar RP. 2008; Mammalian type I gonadotropin-releasing hormone receptors undergo slow, constitutive, agonist-independent internalization. *Endocrinology.* 149:1415–1422. [PubMed: 18039780]
- Peltoketo H, Strauss L, Karjalainen R, Zhang M, Stamp GW, Segaloff DL, Poutanen M, Huhtaniemi IT. 2010; Female mice expressing constitutively active mutants of FSH receptor present with a phenotype of premature follicle depletion and estrogen excess. *Endocrinology.* 151:1872–1883. [PubMed: 20172968]
- Pernasetti F, Vasilyev VV, Rosenberg SB, Bailey JS, Huang HJ, Miller WL, Mellon PL. 2001; Cell-specific transcriptional regulation of FSH β by activin and GnRH in the L β T2 pituitary gonadotrope cell model. *Endocrinology.* 142:2284–2295. [PubMed: 11356674]
- Perrett RM, McArdle CA. 2013; Molecular mechanisms of gonadotropin-releasing hormone signaling: integrating cyclic nucleotides into the network. *Frontiers in endocrinology.* 4:180. [PubMed: 24312080]
- Pielecka J, Quaynor SD, Moenter SM. 2006; Androgens increase gonadotropin-releasing hormone neuron firing activity in females and interfere with progesterone negative feedback. *Endocrinology.* 147:1474–1479. [PubMed: 16339200]
- Pierce, JG. Gonadotropin chemistry and biosynthesis. In: Knobil, E, Neill, JD, editors. *In The Physiology of Reproduction.* New York: Raven Press; 1988. 1335–1348.
- Pierce JG, Parsons TF. 1981; Glycoprotein hormones: structure and function. *Ann Rev Biochem.* 50:465–495. [PubMed: 6267989]
- Purwana IN, Kanasaki H, Mijiddorj T, Oride A, Miyazaki K. 2011; Induction of dual-specificity phosphatase 1 (DUSP1) by pulsatile gonadotropin-releasing hormone stimulation: role for gonadotropin subunit expression in mouse pituitary LbetaT2 cells. *Biol Reprod.* 84:996–1004. [PubMed: 21228211]
- Purwana IN, Kanasaki H, Oride A, Mijiddorj T, Shintani N, Hashimoto H, Baba A, Miyazaki K. 2010; GnRH-induced PACAP and PAC1 receptor expression in pituitary gonadotrophs: a possible role in the regulation of gonadotropin subunit gene expression. *Peptides.* 31:1748–1755. [PubMed: 20553777]
- Quirk CC, Lozada KL, Keri RA, Nilson JH. 2001; A single Pitx1 binding site is essential for activity of the LHBeta promoter in transgenic mice. *Mol Endocrinol.* 15:734–746. [PubMed: 11328855]
- Reddy GR, Xie C, Lindaman LL, Coss D. 2013; GnRH increases c-Fos half-life contributing to higher FSHbeta induction. *Mol Endocrinol.* 27:253–265. [PubMed: 23275456]

- Ringstrom SJ, Szabo M, Kilen SM, Saberi S, Knox KL, Schwartz NB. 1997; The antiprogestins RU486 and ZK98299 affect follicle-stimulating hormone secretion differentially on estrus, but not on proestrus. *Endocrinology*. 138:2286–2290. [PubMed: 9165013]
- Roberson MS, Misra-Press A, Laurance ME, Stork PJ, Maurer RA. 1995; A role for mitogen-activated protein kinase in mediating activation of the glycoprotein hormone alpha-subunit promoter by gonadotropin-releasing hormone. *Mol Cell Biol*. 15:3531–3519. [PubMed: 7791760]
- Roberson MS, Zhang T, Li HL, Mulvaney JM. 1999; Activation of the p38 mitogen-activated protein kinase pathway by gonadotropin-releasing hormone. *Endocrinology*. 140:1310–1318. [PubMed: 10067858]
- Roland AV, Nunemaker CS, Keller SR, Moenter SM. 2010; Prenatal androgen exposure programs metabolic dysfunction in female mice. *J Endocrinol*. 207:213–223. [PubMed: 20713501]
- Rosenfield RL. 2013; Clinical review: Adolescent anovulation: maturational mechanisms and implications. *J Clin Endocrinol Metab*. 98:3572–3583. [PubMed: 23913942]
- Rosenfield RL, Bordini B. 2010; Evidence that obesity and androgens have independent and opposing effects on gonadotropin production from puberty to maturity. *Brain research*. 1364:186–197. [PubMed: 20816944]
- Rossetti R, Ferrari I, Bonomi M, Persani L. 2016 Genetics of primary ovarian insufficiency. *Clinical genetics*.
- Roybal LL, Hambarchyan A, Meadows JD, Barakat NH, Pepa PA, Breen KM, Mellon PL, Coss D. 2014; Roles of Binding Elements, FOXL2 Domains, and Interactions With cJUN and SMADs in Regulation of FSH β . *Molecular Endocrinology*. 28:1640–1655. [PubMed: 25105693]
- Sahin S, Eroglu M, Selcuk S, Turkgeldi L, Kozali S, Davutoglu S, Muhcu M. 2014; Intrinsic factors rather than vitamin D deficiency are related to insulin resistance in lean women with polycystic ovary syndrome. *European review for medical and pharmacological sciences*. 18:2851–2856. [PubMed: 25339479]
- Salisbury TB, Binder AK, Grammer JC, Nilson JH. 2007; Maximal activity of the luteinizing hormone beta-subunit gene requires beta-catenin. *Mol Endocrinol*. 21:963–971. [PubMed: 17244763]
- Sealfon SC, Weinstein H, Millar RP. 1997; Molecular mechanisms of ligand interaction with the gonadotropin-releasing hormone receptor. *Endocrine Rev*. 18:180–205. [PubMed: 9101136]
- Seeburg PH, Mason AJ, Stewart TA, Nikolics K. 1987; The mammalian GnRH gene and its pivotal role in reproduction. *Recent Prog Horm Res*. 43:69–98. [PubMed: 3306841]
- Semaan SJ, Kauffman AS. 2010; Sexual differentiation and development of forebrain reproductive circuits. *Current opinion in neurobiology*. 20:424–431. [PubMed: 20471241]
- Shupnik MA. 1990; Effects of gonadotropin-releasing hormone on rat gonadotropin gene transcription in vitro: Requirement for pulsatile administration for luteinizing hormone-beta gene stimulation. *Mol Endocrinol*. 4:1444–1450. [PubMed: 2126596]
- Shupnik MA. 1996; Gonadotropin gene modulation by steroids and gonadotropin-releasing hormone. *Biol Reprod*. 54:279–286. [PubMed: 8788177]
- Shupnik MA, Gharib SD, Chin WW. 1989a; Divergent effects of estradiol on gonadotropin gene transcription in pituitary fragments. *Mol Endocrinol*. 3:474–480. [PubMed: 2501667]
- Shupnik MA, Rosenzweig BA. 1991; Identification of an estrogen-responsive element in the rat LH beta gene. DNA-estrogen receptor interactions and functional analysis. *J Biol Chem*. 266:17084–17091. [PubMed: 1894604]
- Shupnik MA, Weinmann CM, Notides AC, Chin WW. 1989b; An upstream region of the rat luteinizing hormone beta gene binds estrogen receptor and confers estrogen responsiveness. *J Biol Chem*. 264:80–86. [PubMed: 2909546]
- Sim PJ, Wolbers WB, Mitchell R. 1995; Activation of MAP kinase by the LHRH receptor through a dual mechanism involving protein kinase C and a pertussis toxin-sensitive G protein. *Mol Cell Endocrinol*. 112:257–263. [PubMed: 7489830]
- Smith MS. 1982; Effect of Pulsatile Gonadotropin-Releasing Hormone on the Release of Luteinizing Hormone and Follicle-Stimulating Hormone in Vitro by Anterior Pituitaries from Lactating and Cycling Rats. *Endocrinology*. 110:882–891. [PubMed: 6799281]

- Stanislaus D, Ponder S, Ji TH, Conn PM. 1998; Gonadotropin-releasing hormone receptor couples to multiple G proteins in rat gonadotrophs and in GGH3 cells: evidence from palmitoylation and overexpression of G proteins. *Biol Reprod.* 59:579–586. [PubMed: 9716556]
- Stojilkovic SS. 2012; Molecular mechanisms of pituitary endocrine cell calcium handling. *Cell Calcium.* 51:212–221. [PubMed: 22138111]
- Stojilkovic SS, Iida T, Merelli F, Torsello A, Krsmanovic LZ, Catt KJ. 1991; Interactions between calcium and protein kinase C in the control of signaling and secretion in pituitary gonadotrophs. *The Journal of biological chemistry.* 266:10377–10384. [PubMed: 1645348]
- Stojilkovic SS, Reinhart J, Catt KJ. 1994; Gonadotropin-releasing hormone receptors: Structure and signal transduction pathways. *Endocrine Rev.* 15:462–499. [PubMed: 7988482]
- Stojilkovic SS, Tabak J, Bertram R. 2010; Ion channels and signaling in the pituitary gland. *Endocrine reviews.* 31:845–915. [PubMed: 20650859]
- Sundaresan S, Colin IM, Pestell RG, Jameson JL. 1996; Stimulation of mitogen-activated protein kinase by gonadotropin-releasing hormone: evidence for the involvement of protein kinase C. *Endocrinology.* 137:304–311. [PubMed: 8536629]
- Szabo M, Kilen SM, Saberi S, Ringstrom SJ, Schwartz NB. 1998; Antiprogestins suppress basal and activin-stimulated follicle-stimulating hormone secretion in an estrogen-dependent manner. *Endocrinology.* 139:2223–2228. [PubMed: 9564826]
- Thackray VG, Hunnicutt JL, Memon AK, Ghochani Y, Mellon PL. 2009; Progesterone inhibits basal and GnRH induction of luteinizing hormone β -subunit gene expression. *Endocrinology.* 150:2395–2403. [PubMed: 19106225]
- Thackray VG, McGillivray SM, Mellon PL. 2006; Androgens, progestins and glucocorticoids induce follicle-stimulating hormone β -subunit gene expression at the level of the gonadotrope. *Mol Endocrinol.* 20:2062–2079. [PubMed: 16675544]
- Thackray VG, Mellon PL. 2008; Synergistic induction of follicle-stimulating hormone beta-subunit gene expression by gonadal steroid hormone receptors and smad proteins. *Endocrinology.* 149:1091–1102. [PubMed: 18079204]
- Thackray VG, Mellon PL, Coss D. 2010; Hormones in synergy: Regulation of the pituitary gonadotropin genes. *Mol Cell Endocrinol.* 314:192–203. [PubMed: 19747958]
- Thompson IR, Kaiser UB. 2014; GnRH pulse frequency-dependent differential regulation of LH and FSH gene expression. *Mol Cell Endocrinol.* 385:28–35. [PubMed: 24056171]
- Tortorese DJ, Brooks J, Ingleton PM, McNeilly AS. 1998; Detection of Prolactin Receptor Gene Expression in the Sheep Pituitary Gland and Visualization of the Specific Translation of the Signal in Gonadotrophs. *Endocrinology.* 139:5215–5223. [PubMed: 9832462]
- Tran S, Zhou X, Lafleur C, Calderon MJ, Ellsworth BS, Kimmins S, Boehm U, Treier M, Boerboom D, Bernard DJ. 2013; Impaired fertility and FSH synthesis in gonadotrope-specific Foxl2 knockout mice. *Mol Endocrinol.* 27:407–421. [PubMed: 23340250]
- Tremblay JJ, Drouin J. 1999; Egr-1 is a downstream effector of GnRH and synergizes by direct interaction with Ptx1 and SF-1 to enhance luteinizing hormone β gene transcription. *Mol Cell Biol.* 19:2567–2576. [PubMed: 10082522]
- Tsutsumi M, Zhou W, Millar RP, Mellon PL, Roberts JL, Flanagan CA, Dong K, Gillo B, Sealfon SC. 1992; Cloning and functional expression of a mouse gonadotropin-releasing hormone receptor. *Mol Endocrinol.* 6:1163–1169. [PubMed: 1324422]
- Tsutsumi R, Mistry D, Webster NJ. 2010; Signaling responses to pulsatile gonadotropin-releasing hormone in LbetaT2 gonadotrope cells. *J Biol Chem.* 285:20262–20272. [PubMed: 20406815]
- Ulloa-Aguirre A, Maldonado A, Damian-Matsumura P, Timossi C. 2001; Endocrine regulation of gonadotropin glycosylation. *Arch Med Res.* 32:520–532. [PubMed: 11750727]
- Ulloa-Aguirre A, Timossi C, Damian-Matsumura P, Dias JA. 1999; Role of glycosylation in function of follicle-stimulating hormone. *Endocrine.* 11:205–215. [PubMed: 10786817]
- Vale W, Rivier C, Brown M. 1977; Regulatory peptides of the hypothalamus. *Ann Rev Physiol.* 39:473–527. [PubMed: 15499]
- Van Goor F, Krsmanovic LZ, Catt KJ, Stojilkovic SS. 1999; Coordinate regulation of gonadotropin-releasing hormone neuronal firing patterns by cytosolic calcium and store depletion. *Proc Natl Acad Sci USA.* 96:4101–4106. [PubMed: 10097170]

- Veiga-Lopez A, Ye W, Phillips DJ, Herkimer C, Knight PG, Padmanabhan V. 2008; Developmental programming: deficits in reproductive hormone dynamics and ovulatory outcomes in prenatal, testosterone-treated sheep. *Biol Reprod.* 78:636–647. [PubMed: 18094354]
- Walsh HE, Shupnik MA. 2009; Proteasome regulation of dynamic transcription factor occupancy on the GnRH-stimulated luteinizing hormone beta-subunit promoter. *Mol Endocrinol.* 23:237–250. [PubMed: 19095772]
- Wang H, Butnev V, Bousfield GR, Kumar TR. 2016; A human FSHB transgene encoding the double N-glycosylation mutant (Asn(7Delta) Asn(24Delta)) FSHbeta subunit fails to rescue Fshb null mice. *Mol Cell Endocrinol.* 426:113–124. [PubMed: 26911932]
- Wang H, Larson M, Jablonka-Shariff A, Pearl CA, Miller WL, Conn PM, Boime I, Kumar TR. 2014; Redirecting intracellular trafficking and the secretion pattern of FSH dramatically enhances ovarian function in mice. *Proceedings of the National Academy of Sciences.* 111:5735–5740.
- Wang Q, Chikina M, Zaslavsky E, Pincas H, Sealfon SC. 2013; β -Catenin Regulates GnRH-Induced FSH β Gene Expression. *Molecular Endocrinology.* 27:224–237. [PubMed: 23211523]
- Wang Y, Fortin J, Lamba P, Bonomi M, Persani L, Roberson MS, Bernard DJ. 2008; Activator protein-1 and smad proteins synergistically regulate human follicle-stimulating hormone beta-promoter activity. *Endocrinology.* 149:5577–5591. [PubMed: 18653705]
- Weck J, Anderson AC, Jenkins S, Fallest PC, Shupnik MA. 2000; Divergent and composite gonadotropin-releasing hormone-responsive elements in the rat luteinizing hormone subunit genes. *Mol Endocrinol.* 14:472–485. [PubMed: 10770486]
- Weiss J, Jameson JL, Burrin JM, Crowley WFJ. 1990; Divergent responses of gonadotropin subunit messenger RNAs to continuous *versus* pulsatile gonadotropin-releasing hormone *in vitro*. *Mol Endocrinol.* 4:557–564. [PubMed: 2126344]
- Wide L, Bakos O. 1993; More basic forms of both human follicle-stimulating hormone and luteinizing hormone in serum at midcycle compared with the follicular or luteal phase. *J Clin Endocrinol Metab.* 76:885–889. [PubMed: 8473400]
- Wierman ME, Wang C. 1990; Androgen selectively stimulates follicle-stimulating hormone-beta mRNA levels after gonadotropin-releasing hormone antagonist administration. *Biol Reprod.* 42:563–571. [PubMed: 2111188]
- Willars GB, Heding A, Vrecl M, Sellar R, Blumenröhr M, Nahorski SR, Eidne KA. 1999; Lack of a C-terminal tail in the mammalian gonadotropin-releasing hormone receptor confers resistance to agonist-dependent phosphorylation and rapid desensitization. *J Biol Chem.* 274:30146–30153. [PubMed: 10514504]
- Winters SJ, Ishizaka K, Kitahara S, Troen P, Attardi B. 1992; Effects of testosterone on gonadotropin subunit messenger ribonucleic acids in the presence or absence of gonadotropin-releasing hormone. *Endocrinology.* 130:726–734. [PubMed: 1370794]
- Winters SJ, Moore JP. 2007; Paracrine control of gonadotrophs. *Semin Reprod Med.* 25:379–387. [PubMed: 17710734]
- Winters SJ, Moore JP Jr. 2011; PACAP, an autocrine/paracrine regulator of gonadotrophs. *Biol Reprod.* 84:844–850. [PubMed: 21191111]
- Witham EA, Meadows JD, Hoffmann HM, Shojaei S, Coss D, Kauffman AS, Mellon PL. 2013; Kisspeptin Regulates Gonadotropin Genes via Immediate Early Gene Induction in Pituitary Gonadotrophs. *Molecular Endocrinology.* 27:1283–1294. [PubMed: 23770611]
- Woodruff TK, Besecke LM, Groome N, Draper LB, Schwartz NB, Weiss J. 1996; Inhibin A and inhibin B are inversely correlated to follicle-stimulating hormone, yet are discordant during the follicular phase of the rat estrous cycle, and inhibin A is expressed in a sexually dimorphic manner. *Endocrinology.* 137:5463–5467. [PubMed: 8940372]
- Wu S, Wilson MD, Busby ER, Isaac ER, Sherwood NM. 2010; Disruption of the single copy gonadotropin-releasing hormone receptor in mice by gene trap: severe reduction of reproductive organs and functions in developing and adult mice. *Endocrinology.* 151:1142–1152. [PubMed: 20068010]
- Wurnbach E, Yuen T, Ebersole BJ, Sealfon SC. 2001; Gonadotropin-releasing hormone receptor-coupled gene network organization. *The Journal of biological chemistry.* 276:47195–47201. [PubMed: 11581274]

- Xie C, Jonak CR, Kauffman AS, Coss D. 2015; Gonadotropin and kisspeptin gene expression, but not GnRH, are impaired in cFOS deficient mice. *Molecular and Cellular Endocrinology*. 411:223–231. [PubMed: 25958044]
- Yuen T, Choi SG, Pincas H, Waring DW, Sealfon SC, Turgeon JL. 2012; Optimized amplification and single-cell analysis identify GnRH-mediated activation of Rap1b in primary rat gonadotropes. *Molecular and Cellular Endocrinology*. 350:10–19. [PubMed: 22127306]
- Zhang H, Bailey JS, Coss D, Lin B, Tsutsumi R, Lawson MA, Mellon PL, Webster NJ. 2006; Activin modulates the transcriptional response of LbetaT2 cells to gonadotropin-releasing hormone and alters cellular proliferation. *Mol Endocrinol*. 20:2909–2930. [PubMed: 16772531]
- Zhang T, Roberson MS. 2006; Role of MAP kinase phosphatases in GnRH-dependent activation of MAP kinases. *J Mol Endocrinol*. 36:41–50. [PubMed: 16461925]
- Zmeili SM, Papavasiliou SS, Thorner MO, Evans WS, Marshall JC, Landefeld TD. 1986; Alpha and luteinizing hormone beta subunit messenger ribonucleic acids during the rat estrous cycle. *Endocrinology*. 119:1867–1869. [PubMed: 2428604]

Highlights

1. Gonadotropin hormone levels are regulated by variations in GnRH pulse frequency
2. GnRH-induced transcription of their unique β -subunits controls gonadotropin levels
3. GnRH signaling cross-talks with gonadal and pituitary factors to regulate β -subunits
4. GnRH induces LH β via induction of EGR1 factor and FSH β via induction of cFOS and cJUN
5. Signaling pathways and factors respond differently to varied GnRH pulse frequencies

Figure 1. GnRH signaling pathways

Following binding of GnRH, GnRH receptor (GnRHR) primarily couples to Gαq/11 and activates PLC signaling that leads to formation of inositol trisphosphate (IP₃) and diacylglycerol (DAG). IP₃ causes an increase in the intracellular calcium (i-Ca) by mobilization from intracellular stores, while DAG activates PKC, which in turn leads to ERK1/2 MAPK activation. Calcium influx (e-Ca) occurs through voltage-gated calcium channels (VGCC), whose activation may involve PKC. Increase in intracellular calcium from either source activates CamKII, which is necessary for induction of both LHβ and FSHβ. Sustained increase of extracellular calcium causes higher activation of ERK1/2 and preferentially increase of FSHβ.

Figure 2. GnRH signaling pathways that differentially regulate LH β and FSH β

A - Left, During low pulse frequency of GnRH, classical GnRH signaling, as described in Figure 1, activates both LH β and FSH β transcription. Higher activity of ERK1/2, likely caused by sustained influx of extracellular calcium, activates specifically FSH β and contributes to higher levels of FSH β during low pulse frequency. **B - Right,** During high pulse frequency of GnRH in proestrus, GnRHR engages Gas in addition to Gαq/11. This may occur due to increased receptor concentration that is correlated with higher LH β induction. Gas is necessary for maximal induction of LH β via unknown mechanisms, indicated with a question mark. Gas signaling module is resistant to desensitization, while both IP3 receptors and DAG formation in the Gαq/11 signaling module desensitize with high pulse frequency. This leads to higher LH β levels than FSH β with high pulse frequency of GnRH.

Figure 3. Transcriptional regulation of β -subunits by GnRH

GnRH activates transcription of gonadotropin β -subunits via intermediates, immediate early genes; EGR1 that induces LH β (top) and cFOS that induces FSH β following heterodimerization with cJUN that forms the AP1 transcription factor (bottom). These GnRH-induced genes organize complexes on their respective target gene promoters to integrate basal (Ptx1 for LH β and NFY for FSH β) and gonadotrope-specific factors (SF1). Under basal conditions (left panel), β subunit expression remains low due to repressors, Nab for LH β , and JDP2 or ICER for FSH β that recruit histone deacetylases (HDAC). Following EGR1 and cFOS induction by GnRH (middle panel), they replace Nab and JDP2 respectively, to induce β -subunit expression. Formation of the active transcriptional complex recruits histone acetyltransferase (HAT) to remodel histones (H). EGR1 and cFOS have very short half-lives and are rapidly degraded. This allows repressor recruitment and removal of active histone marks in the promoter until the following pulse of GnRH (right panel).

Figure 4. Differential regulation of β -subunits expression by GnRH

A – Left, During fast pulse frequency of GnRH in proestrus, EGR1 and cFOS accumulate to overcome repressors. Both EGR1 and cFOS induction by GnRH is dependent on CamKII activation of SRF transcription factor. Since the EGR1 promoter has four SRF response elements, compared to one in the cFOS promoter, EGR1 is more highly induced than cFOS, which leads to higher induction of LH β than FSH β during high pulse frequency of GnRH. **B - Right,** During low pulse frequency of GnRH, EGR1 is degraded throughout the interpulse interval due to its short half-life. GnRH signaling via ERK1/2 specifically phosphorylates cFOS protein (indicated with “p”) and increases its half-life, which allows accumulation of cFOS. GnRH and activin (Act) synergistically induce FSH β , indicated with “x”. GnRH-induced FOS/AP1 complex interacts with activin-regulated SMAD3/SMAD4/FOXL2 complex, which leads to synergistic induction of FSH β and increase of FSH in the circulation during low pulse frequency of GnRH.