Cochrane
Library

Cochrane Database of Systematic Reviews

Chinese herbal medicine for subfertile women with polycystic ovarian syndrome (Review)

Zhou K, Zhang J, Xu L, Wu T, Lim CED

Zhou K, Zhang J, Xu L, Wu T, Lim CED.
Chinese herbal medicine for subfertile women with polycystic ovarian syndrome.
Cochrane Database of Systematic Reviews 2016, Issue 10. Art. No.: CD007535.
DOI: [10.1002/14651858.CD007535.pub3](https://doi.org/10.1002/14651858.CD007535.pub3).

www.cochranelibrary.com

TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
SUMMARY OF FINDINGS	4
BACKGROUND	8
OBJECTIVES	9
METHODS	9
Figure 1.	11
RESULTS	12
Figure 2.	14
Figure 3.	15
Figure 4.	17
DISCUSSION	17
AUTHORS' CONCLUSIONS	18
ACKNOWLEDGEMENTS	18
REFERENCES	19
CHARACTERISTICS OF STUDIES	42
DATA AND ANALYSES	64
Analysis 1.1. Comparison 1 CHM versus clomiphene, Outcome 1 Pregnancy rate (per woman).	64
Analysis 1.2. Comparison 1 CHM versus clomiphene, Outcome 2 Ovulation rate (per woman).	65
Analysis 2.1. Comparison 2 CHM + clomiphene versus clomiphene, Outcome 1 Pregnancy rate (per woman).	65
Analysis 3.1. Comparison 3 CHM + follicle aspiration + ovulation induction versus follicle aspiration + ovulation induction, Outcome 1 Pregnancy rate (per woman).	66
Analysis 3.2. Comparison 3 CHM + follicle aspiration + ovulation induction versus follicle aspiration + ovulation induction, Outcome 2 LUFs (adverse events).	67
Analysis 3.3. Comparison 3 CHM + follicle aspiration + ovulation induction versus follicle aspiration + ovulation induction, Outcome 3 OHSS (adverse events).	67
Analysis 3.4. Comparison 3 CHM + follicle aspiration + ovulation induction versus follicle aspiration + ovulation induction, Outcome 4 Multiple pregnancy (adverse events).	67
Analysis 4.1. Comparison 4 CHM + LOD versus LOD, Outcome 1 Pregnancy rate (per woman).	68
Analysis 4.2. Comparison 4 CHM + LOD versus LOD, Outcome 2 Ovulation rate (per woman).	68
ADDITIONAL TABLES	68
APPENDICES	71
WHAT'S NEW	77
HISTORY	77
CONTRIBUTIONS OF AUTHORS	78
DECLARATIONS OF INTEREST	78
SOURCES OF SUPPORT	78
DIFFERENCES BETWEEN PROTOCOL AND REVIEW	78
NOTES	78
INDEX TERMS	79

[Intervention Review]

Chinese herbal medicine for subfertile women with polycystic ovarian syndrome

Kunyan Zhou^{1,2}, Jing Zhang^{1,2}, Liangzhi Xu^{1,2}, Taixiang Wu³, Chi Eung Danforn Lim⁴

¹Department of Obstetrics and Gynecology, West China Second University Hospital, Sichuan University, Chengdu, China. ²Key Laboratory of Birth Defects and Related Diseases of Women and Children (Sichuan University), Ministry of Education, Chengdu, China.

³Chinese Clinical Trial Registry, Chinese Ethics Committee of Registering Clinical Trials, West China Hospital, Sichuan University, Chengdu, China. ⁴Faculty of Science, University of Technology Sydney, Earlwood, Australia

Contact address: Liangzhi Xu, Department of Obstetrics and Gynecology, West China Second University Hospital, Sichuan University, Chengdu, Sichuan, China. liangzxi@126.com.

Editorial group: Cochrane Gynaecology and Fertility Group.

Publication status and date: New search for studies and content updated (no change to conclusions), published in Issue 10, 2016.

Citation: Zhou K, Zhang J, Xu L, Wu T, Lim CED. Chinese herbal medicine for subfertile women with polycystic ovarian syndrome. *Cochrane Database of Systematic Reviews* 2016, Issue 10. Art. No.: CD007535. DOI: [10.1002/14651858.CD007535.pub3](https://doi.org/10.1002/14651858.CD007535.pub3).

Copyright © 2016 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

Polycystic ovarian syndrome (PCOS) is one of the most common reproductive endocrinology abnormalities, and affects 5% to 10% of women of reproductive age. Western medicines, such as oral contraceptives, insulin sensitizers and laparoscopic ovarian drilling (LOD), have been used to treat PCOS. Recently, many studies have been published that consider Chinese herbal medicine (CHM) as an alternative treatment for women with PCOS.

Objectives

To assess the efficacy and safety of CHM for subfertile women with PCOS.

Search methods

We searched sources, including the following databases, from inception to 9 June 2016: the Cochrane Gynaecology and Fertility Group Specialized Register, the Cochrane Central Register of Controlled Trials (CENTRAL), MEDLINE, EMBASE, Allied and Complementary Medicine (AMED), PsycINFO, Chinese National Knowledge Infrastructure (CNKI), VIP, Wanfang and trial registries. In addition, we searched the reference lists of included trials and contacted experts in the field to locate trials.

Selection criteria

Randomized controlled trials (RCTs) that considered the use of CHM for the treatment of subfertile women with PCOS.

Data collection and analysis

Two review authors independently screened appropriate trials for inclusion, assessed the risk of bias in included studies and extracted data. We contacted primary study authors for additional information. We conducted meta-analyses. We used the odds ratios (ORs) to report dichotomous data, with 95% confidence intervals (CI). We assessed the quality of the evidence using the Grading of Recommendations Assessment, Development and Evaluation (GRADE) methods.

Main results

We included five RCTs with 414 participants. The comparisons in the included trials were as follows: CHM versus clomiphene, CHM plus clomiphene versus clomiphene (with or without ethinyloestradiol cyproterone acetate (CEA)), CHM plus follicle aspiration plus ovulation

induction versus follicle aspiration plus ovulation induction alone, and CHM plus laparoscopic ovarian drilling (LOD) versus LOD alone. The overall quality of the evidence for most comparisons was very low.

None of the included studies reported live birth rate, and only one study reported data on adverse events.

When CHM was compared with clomiphene (with or without LOD in both arms), there was no evidence of a difference between the groups in pregnancy rates (odds ratio (OR) 1.98, 95% confidence interval (CI) 0.78 to 5.06; two studies, 90 participants, I^2 statistic = 0%, *very low quality evidence*). No study reported data on adverse events. When CHM plus clomiphene was compared with clomiphene (with or without CEA), there was low quality evidence of a higher pregnancy rate in the CHM plus clomiphene group (OR 2.62, 95% CI 1.65 to 4.14; three RCTs, 300 women, I^2 statistic = 0%, *low quality evidence*). No data were reported on adverse events.

When CHM with follicle aspiration and ovulation induction was compared with follicle aspiration and ovulation induction alone, there was no evidence of a difference between the groups in pregnancy rates (OR 1.60, 95% CI 0.46 to 5.52; one study, 44 women, *very low quality evidence*), severe luteinized unruptured follicle syndrome (LUFS) (OR 0.60, 95% CI 0.06 to 6.14; one study, 44 women, *very low quality evidence*), ovarian hyperstimulation syndrome (OHSS) (OR 0.16, 95% CI 0.00 to 8.19; one study, 44 women, *very low quality evidence*) or multiple pregnancy (OR 0.60, 95% CI 0.06 to 6.14; one study, 44 women, *very low quality evidence*).

When CHM with LOD was compared with LOD alone, there was no evidence of a difference between the groups in rates of pregnancy (OR 3.50, 95% CI 0.72 to 17.09; one study, 30 women, *very low quality evidence*). No data were reported on adverse events.

There was no evidence of a difference between any of the comparison groups for any other outcomes. The quality of the evidence for all other comparisons and outcomes was very low. The main limitations in the evidence were failure to report live birth or adverse events, failure to describe study methods in adequate detail and imprecision due to very low event rates and wide CIs.

Authors' conclusions

There is insufficient evidence to support the use of CHM for women with PCOS and subfertility. No data are available on live birth, and there is no consistent evidence to indicate that CHM influences fertility outcomes. However there is very limited low quality evidence to suggest that the addition of CHM to clomiphene may improve pregnancy rates. There is insufficient evidence on adverse effects to indicate whether CHM is safe.

PLAIN LANGUAGE SUMMARY

Chinese herbal medicines for subfertile women with polycystic ovarian syndrome

Review question

We reviewed the evidence about the effect of Chinese herbal medicines (CHM) on rates of live birth pregnancy and adverse events in subfertile women with polycystic ovarian syndrome (PCOS).

Background

PCOS is a common and complex reproductive endocrine disorder, affecting 5% to 10% of women of reproductive age. Women with PCOS may present with irregular menstrual cycles, subfertility (failure to conceive), hirsutism (excessive hair growth), acne and obesity. Many western medical therapies have been used to manage PCOS, such as oral contraceptives, insulin sensitizers and laparoscopic ovarian drilling (LOD). CHM have been suggested as an alternative approach for subfertile women with PCOS. We wanted to investigate the effectiveness and safety of CHM compared to other therapies for subfertile women with PCOS.

Study characteristics

We searched evidence from commonly used databases and it is current to 9 June 2016. We included five RCTs with 414 participants. These included studies comparing CHM to western medicine, CHM plus western medicine versus western medicine, and CHM plus surgery versus surgery. All the included studies were in Chinese. All studies had fewer than six menstrual cycles treatment duration and less than one year follow-up duration. None of the included studies reported live birth, all reported pregnancy, two reported ovulation and only one reported adverse events.

Key results

There was insufficient evidence to support the use of CHM for women with PCOS and subfertility. No data were available on live birth, and there was no consistent evidence to indicate that CHM improves fertility outcomes. When CHM was compared with clomiphene (with or without laparoscopic ovarian drilling (LOD) in both arms), the pregnancy rates were no different between the groups. When CHM with follicle aspiration and ovulation induction was compared with follicle aspiration and ovulation induction alone, pregnancy rates were no different between the groups. When CHM with LOD was compared with LOD alone, the pregnancy rates were no different between the groups. However there was limited low quality evidence to suggest that the addition of CHM to clomiphene may improve pregnancy rates.

There was no evidence of a difference between any of the comparison groups for any other outcomes. There was insufficient evidence on adverse effects to indicate whether CHM is safe.

Quality of the evidence

The quality of the evidence was low or very low. The main limitations in the evidence were failure to report live birth or adverse events, failure to describe study methods in adequate detail, and imprecision, with very low event rates and wide confidence intervals.

SUMMARY OF FINDINGS

Summary of findings for the main comparison. Chinese herbal medicine (CHM) versus clomiphene for subfertile women with polycystic ovarian syndrome (PCOS)

CHM versus clomiphene for subfertile women with PCOS

Population: subfertile women with PCOS

Setting: fertility clinics

Intervention: Chinese herbal medicine (CHM)

Comparison: clomiphene

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	Number of participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Clomiphene	CHM				
Live birth rate	Not reported					
Pregnancy rate	250 per 1000	398 per 1000 (206 to 628)	OR 1.98 (0.78 to 5.06)	90 (2 RCTs)	⊕⊕⊕⊕ Very low ^{1,2}	
Adverse effects	Not reported					

*The basis for the assumed risk is the median control group risk across studies. The corresponding risk (and its 95% CI) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: CI: confidence interval; OR: odds ratio

GRADE Working Group grades of evidence

High quality: further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: we are very uncertain about the estimate.

¹Downgraded one level for serious risk of bias: study methods not described in sufficient detail.

²Downgraded two levels for very serious imprecision: small sample size, only 32 events altogether, CIs compatible with no effect or with substantial benefit from the intervention.

Summary of findings 2. Chinese herbal medicine (CHM) + clomiphene versus clomiphene for subfertile women with polycystic ovarian syndrome (PCOS)

CHM + clomiphene compared to clomiphene for subfertile women with PCOS

Population: subfertile women with PCOS

Setting: fertility clinics

Intervention: Chinese herbal medicine (CHM) + clomiphene

Comparison: clomiphene

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	Number of participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Clomiphene	CHM + clomiphene				
Live birth	Not reported					
Pregnancy rate (per woman)	387 per 1000	623 per 1000 (510 to 723)	OR 2.62 (1.65 to 4.14)	300 (3 RCTs)	⊕⊕⊕⊕ low ^{1,2}	
Adverse events	Not reported					

*The basis for the assumed risk is the median control group risk across studies. The corresponding risk (and its 95% CI) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: CI: confidence interval; OR: odds ratio.

GRADE Working Group grades of evidence

High quality: further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: we are very uncertain about the estimate.

¹Downgraded one level for serious risk of bias: study methods not described in sufficient detail.

²Downgraded one level for serious imprecision: small studies, low overall event rate.

Summary of findings 3. Chinese herbal medicine (CHM) + follicle aspiration + ovulation induction versus follicle aspiration + ovulation induction for subfertile women with polycystic ovarian syndrome (PCOS)

CHM + follicle aspiration + ovulation induction compared to follicle aspiration + ovulation induction for subfertile women with PCOS

Population: subfertile women with PCOS

Setting: fertility clinics

Intervention: Chinese herbal medicine + follicle aspiration + ovulation induction

Comparison: follicle aspiration + ovulation induction

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	Number of participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Follicle aspiration + ovulation induction	CHM + follicle aspiration + ovulation induction				
Live birth	Not reported					
Pregnancy rate	292 per 1000	397 per 1000 (159 to 694)	OR 1.60 (0.46 to 5.52)	44 (1 RCT)	⊕○○○ Very low ^{1,2}	
Luteinized unruptured follicle syndrome (adverse events)	83 per 1000	52 per 1000 (5 to 358)	OR 0.60 (0.06 to 6.14)	44 (1 RCT)	⊕○○○ Very low ^{1,2}	
Ovarian hyperstimulation syndrome (adverse events)	42 per 1000	7 per 1000 (0 to 263)	OR 0.16 (0.00 to 8.19)	44 (1 RCT)	⊕○○○ Very low ^{1,2}	
Multiple pregnancy (adverse events)	83 per 1000	52 per 1000 (5 to 358)	OR 0.60 (0.06 to 6.14)	44 (1 RCT)	⊕○○○ Very low ^{1,2}	

*The basis for the assumed risk is the median control group risk across studies. The corresponding risk (and its 95% CI) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: CI: confidence interval; OR: odds ratio

GRADE Working Group grades of evidence

High quality: further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: we are very uncertain about the estimate.

¹Downgraded one level for serious risk of bias: the study authors did not report the study methods in sufficient detail, and the study authors did not describe the allocation concealment method.

²Downgraded two levels for very serious imprecision: small study, few events, CIs compatible with no effect or with substantial harm or benefit in either arm.

Summary of findings 4. Chinese herbal medicine (CHM) + laparoscopic ovarian drilling (LOD) versus LOD for subfertile women with polycystic ovarian syndrome (PCOS)

CHM + LOD compared to LOD for subfertile women with PCOS

Population: subfertile women with PCOS

Setting: fertility clinics

Intervention: CHM + LOD

Comparison: LOD

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	Number of participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	LOD	CHM + LOD				
Live birth	Not reported					
Pregnancy rate (per woman)	400 per 1000	700 per 1000 (324 to 919)	OR 3.50 (0.72 to 17.09)	30 (1 RCT)	⊕○○○ Very low ^{1,2}	
Adverse events	Not reported					

*The basis for the assumed risk is the median control group risk across studies. The corresponding risk (and its 95% CI) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: CI: confidence interval; OR: odds ratio

GRADE Working Group grades of evidence

High quality: further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: we are very uncertain about the estimate.

¹Downgraded one level for serious risk of bias: the study authors did not report the study methods in sufficient detail.

²Downgraded two levels for very serious imprecision: small study, few events, and CIs were compatible with no effect or with substantial harm in the CHM group.

BACKGROUND

Description of the condition

Polycystic ovarian syndrome (PCOS) is a complex condition that affects 5% to 10% of women of reproductive age (Carmina 1999). PCOS is characterized by chronic anovulation (ongoing failure or absence of ovulation), hyperandrogenism (excessive production of androgen in women), dyslipidaemia (lipid metabolism disorder), and insulin resistance (a reduced glucose response to a given amount of insulin) leading to hyperinsulinaemia (compensatory serum insulin increase). Women with PCOS may present with irregular menstrual cycles, subfertility (failure to conceive), hirsutism (excessive hair growth), acne and obesity. The cause of PCOS remains unclear. It is proposed that high levels of androgen in serum may be the primary cause (Escobar-Morreale 2005). However, insulin resistance and obesity may also trigger the development of this hormonal defect (Dunaif 1997; Alvarez-Blasco 2006; Gambineri 2006). Other conditions associated with PCOS include type 2 diabetes mellitus (Ehrmann 1999; Legro 1999), gestational diabetes (Boomsma 2006; Lo 2006a), decreased high density lipoprotein cholesterol (HDL-C) (Rajkhowa 1997; Berneis 2007), increased triglycerides (TG) and low density lipoprotein cholesterol (LDL-C) (Talbot 1998; Legro 2001), increased risk of hypertension (high blood pressure) (Lo 2006b), and increased prevalence of metabolic syndrome.

Traditional Chinese Medicine (TCM) follows an independent theoretical and methodological pathway to assess the cause of the disease for making the diagnosis and treatment plan. Even though there is no classification for PCOS within TCM, the symptoms and signs of women with PCOS can be grouped as two disease classes within TCM: amenorrhoea (failure to menstruate) and infertility.

Studies of PCOS and TCM have been conducted since the 1980s (Sun 1981; Yv 1981; Wang 1982). The aetiology and clinical characteristics of PCOS still remain controversial but are believed to be related to the disorders of kidneys, liver and spleen, and from the TCM perspective reproductive function is regarded as being governed by kidneys. It is believed that kidney deficiency may be the main problem in PCOS (Ni 2007; Wang 2008). In addition, in TCM there is an association between the liver and the regulation of blood and the menstrual cycle, and the spleen is associated with body type, obesity and hirsutism (Liu 2009; Hou 2012).

Description of the intervention

Many Western medicine therapies have been used for PCOS, including oral contraceptives, insulin sensitizers, exercise, diet and laparoscopic ovarian drilling (LOD). Several Cochrane reviews have addressed different approaches to PCOS using Western medical treatments (Weiss 2015; Tang 2012; Costello 2007; Farquhar 2012; Sinawat 2012). The oral contraceptive pill (OCP) is believed to be more effective than insulin-sensitizing drugs in improving menstrual patterns and reducing serum androgen levels (Costello 2007). On the other hand, metformin, an insulin-sensitizing drug (ISD), has been found to be more effective than the OCP in reducing fasting insulin levels and not increasing triglyceride levels (Costello 2007). Metformin, either alone or in combination with clomiphene, increases ovulation in women with PCOS and may reduce health risks from insulin resistance and the effect of abnormal levels of androgen. However, the possible adverse effects from using metformin could include nausea and vomiting

(Tang 2012). The optimal duration for metformin pretreatment before initiation of clomiphene citrate is unknown (Sinawat 2012). Gonadotrophin is used for ovulation induction but it may also cause overstimulation of the ovaries. A reduced incidence of overstimulation was found with the use of more expensive urinary follicle stimulating hormone (uFSH) compared to human menopausal gonadotrophin (HMG). A higher overstimulation rate with the addition of gonadotrophin releasing hormone analogues (GnRH-a) to gonadotrophins is suggested (Weiss 2015). LOD followed by clomiphene or gonadotrophins, if necessary, are suggested to be as effective as gonadotrophin therapy alone in inducing ovulation. However, LOD is associated with a lower risk of multiple pregnancy (Farquhar 2012).

Chinese herbal medicines (CHM) are used broadly in various endocrinologic disorders. CHM is used with the aim of improving menstrual patterns, hirsutism, acne and pregnancy rate in women with PCOS (Cong 2006; Yang 2006; Ma 2010). In TCM, there are three different therapeutic strategies to treat PCOS by CHM. Firstly, only one special formula comprising of sovereign medicinal (the ingredient that provides the principal curative action on the main pattern/syndrome or primary symptom) is prescribed to women for the whole menstrual cycle. This formula is occasionally combined with some minister medicinal (the ingredient that helps strengthen the principal curative action) and assistant medicinal (the ingredient that treats the combined pattern/syndrome, relieves secondary symptoms or tempers the action of the sovereign ingredient when the latter is too potent) accordingly to one's individual symptoms and signs (Cui 2004; Ning 2004; Xia 2004; Zhang 2004; Liu 2005; Wang 2005; Cong 2006; Yang 2006). Secondly, different formulae are periodically prescribed to women with PCOS according to each individual's menstrual cycle. This strategy is aimed to restore normal reproductive endocrinological function (Yuan 2003; Xue 2004). Thirdly, Chinese herbal medicines are used in combination with Western medicines for treating PCOS (Li 2000; Li 2002; Ye 2004; Lin 2005; Li 2006).

How the intervention might work

Holistic therapy and multisystem regulation are the therapeutic characteristics of TCM. Many CHM used in treating PCOS are aimed to tone the kidneys to induce ovulation. The components of different formulae act synergistically in various ways. For example, it is proposed that Baishao (*Radix paeoniae alba*), Danggui (*Radix angelicae sinensis*) and Zaojiao (*Fructus gleditsiae sinensis*) reduce release of insulin and androgen (Li 2005), and that Luole (*Basil*) has an oestrogenic effect which prompts follicles to develop and mature (Jin 1986). In addition, it is reported that Dilong (*Lumbricus*), Sanqi (*Radix notoginseng*), Zelan (*Herba lycopi*) and Zexie (*Rhizoma alismatis*) can induce ovulation (Shao 2006), and that Gancao (*Radix glycythizae*) which possesses glucocorticoid effects, can improve ovulatory abnormality. It is reported that Zishiin (*Fluoritum*) can improve endometrial receptivity for embryo implantation and can regulate cervical mucus for sperm passing through the uterus (Wang 2008).

Why it is important to do this review

Various Western medicine therapies have been used for PCOS in recent decades. Their effectiveness varies and some are associated with adverse events. CHM has been used for thousands of years to treat PCOS, which has a different name in TCM. In both developed and developing countries, there is increasing public interest in, and

use of, a wide range of therapies which lie outside the 'mainstream' of traditional Western medical practice. Although CHM is generally considered safe when used properly by qualified practitioners, many herbs and formulae have contraindications, and some can be toxic. There are concerns about adverse events, including allergic reactions and Chinese herbal nephropathy (CHN) (Nortier 2000; Lord 2001; Lampert 2002).

As there is currently insufficient evidence about the safety and efficacy of CHM for the management of PCOS, a systematic review in this area was warranted. No systematic review on this topic has been done before. This is an update of a review first published in 2010 (Zhang 2010)

OBJECTIVES

To assess the efficacy and safety of Chinese herbal medicine (CHM) for subfertile women with polycystic ovarian syndrome (PCOS).

METHODS

Criteria for considering studies for this review

Types of studies

Randomized controlled trials (RCTs) studying the efficacy of Chinese herbal medicine (CHM) for subfertile women with polycystic ovarian syndrome (PCOS).

We excluded quasi-RCTs and non-RCTs.

Types of participants

Women with PCOS and subfertility wishing to conceive (18 to 44 years). We excluded trials that included both fertile and infertile women with PCOS unless there was a stratified analysis based on fertility.

We defined PCOS using the diagnostic criteria of the European Society of Human Reproduction and Embryology (ESHRE) and the American Society of Reproductive Medicine (ASRM) consensus in Rotterdam 2003 (ESHRE/ASRM 2004). PCOS can be diagnosed if a woman has two out of three criteria: oligo or anovulation, clinical or biochemical signs of hyperandrogenism and polycystic ovaries by ultrasonography. These diagnostic criteria exclude individuals who have other aetiologies of hyperandrogenism (such as androgen secreting tumour, hyperprolactinaemia, dysthyroid disease, Cushing syndrome and congenital adrenal cortical hyperplasia).

Ideally, the trials that we considered for inclusion in this review stated and described the diagnostic criteria of PCOS. If the primary study did not employ the Rotterdam criteria, we evaluated the stated diagnostic criteria in each individual study to confirm whether they met the Rotterdam criteria.

We excluded trials whose diagnostic criteria were inconsistent with Rotterdam criteria. If the trial did not clearly state the diagnostic criteria, we contacted the primary study authors for clarification. If clarification was unavailable, we also excluded these trials. Changes in diagnostic criteria might produce variability in the clinical characteristics of the women included and the results obtained. We considered and documented these changes. We plan to perform sensitivity analyses based on these changes when we

locate more RCTs that meet the inclusion criteria of this review in the future.

Types of interventions

1. CHM versus: placebo, no treatment, Western medicine, exercise plus diet control, laparoscopic surgery, another type of CHM, with or without co-medications.
2. CHM combined with another treatment versus another treatment, such as Western medicine, exercise plus diet control or laparoscopic surgery.
3. CHM alone or combined with another treatment versus CHM combined with another treatment.

We excluded trials that included ovarian wedge resection as the control intervention because physicians have not used this method since the application of laparoscopic ovarian drilling (LOD).

We excluded trials without CHM application.

Types of outcome measures

Primary outcomes

1. Live birth rate (per woman).

Secondary outcomes

1. Pregnancy rate per woman.
2. Ovulation rate (confirmed by ultrasound or increased progesterone) per woman.
3. Adverse events (severe or minor).

We defined serious adverse events according to the International Conference on Harmonisation of Technical Requirements for Registration of Pharmaceuticals for Human Use (ICH) Guidelines (ICHEWG 1997) and they included any event that led to death, was life-threatening, required inpatient hospitalization or prolongation of existing hospitalization, or resulted in persistent or significant disability, and any important medical event that might have jeopardized the patient or required intervention to prevent it. For example, we considered severe ovarian hyperstimulation syndrome (OHSS) and severe luteinized unruptured follicle syndrome (LUFS) that required in-patient hospitalization as serious adverse events in this review. We considered all other adverse events as non-serious.

We excluded trials that did not measure any of above outcomes.

Search methods for identification of studies

In consultation with the Cochrane Gynaecology and Fertility Group (CGF) Information Specialist, we formulated a comprehensive search strategy in order to identify all RCTs regardless of language or publication status (published, unpublished, in press or in progress).

Electronic searches

In this review update, we searched the following databases up to 9 June 2016.

1. The CGF Specialized Register.
2. The Cochrane Central Register of Controlled Trials (CENTRAL).
3. Ovid MEDLINE.

4. EMBASE.
5. Allied and Complementary Medicine (AMED).
6. PsycINFO.
7. Chinese National Knowledge Infrastructure (CNKI, including Chinese journal full-text database (CJFD), Chinese selected doctoral dissertations and Master's theses full-text databases (CDMD)).
8. Chinese important conference dissertations full-text database (from 2000 to June 2016), VIP (from 1989 to June 2016).
9. Wanfang database (from 1998 to June 2016).

We constructed search strategies using a combination of subject headings and text words relating to the use of traditional Chinese herbal medicines for the management of PCOS (Appendix 1; Appendix 2; Appendix 3; Appendix 4; Appendix 5; Appendix 6; Appendix 7; Appendix 8; Appendix 9; Appendix 10). We translated all of the search terms into Chinese terms when we conducted the searches in Chinese databases.

We combined the MEDLINE search with the Cochrane highly sensitive search strategy for identifying randomized trials which appears in the *Cochrane Handbook of Systematic Reviews of Interventions* (Version 5.0.2 chapter 6, 6.4.11) (Higgins 2011).

We combined the EMBASE search with trial filters developed by the Scottish Intercollegiate Guidelines Network (SIGN) (<http://sign.ac.uk/methodology/filters.html#random>).

We searched the following for ongoing trials (9 June 2016).

1. The ISRCTN Register (international); Action Medical Research (UK); NIHR Health Technology Assessment Programme (HTA)

(UK); The Wellcome Trust (UK); Medical Research Council (UK); UK trials (UK); NIH Clinical Trials.gov Register (International) (<http://www.controlled-trials.com/mrct/>).

2. The World Health Organization International Trials Registry Platform search portal (<http://apps.who.int/trialsearch/Default.aspx>).

3. The Chinese Clinical Trial Registry (<http://www.chictr.org/Site/Search.aspx?lang=CN>).

Searching other resources

We checked the reference lists of relevant trials, reviews and textbooks. We contacted experts in the field and pharmaceutical companies for relevant trials.

Data collection and analysis

Selection of studies

Two review authors independently performed the searches and retrieved articles (ZK, ZJ). We retrieved the searched trials that claimed to be randomized; two review authors (ZK, ZJ) then confirmed that these trials were correctly randomized by telephoning the original trial authors to evaluate the methodological quality. We judged trials to be adequately randomized if they met the set criteria (Schulz 1995; Jadad 1996; Moher 1998; Jüni 2001; Kjaergard 2001). Two review authors (ZK, ZJ) selected the trials to be included in the review and resolved any disagreements by discussion with a third review author (WT). Also, we listed the excluded studies and the reasons for exclusion in the 'Characteristics of excluded studies' table. See Figure 1 for details of the screening and selection process.

Figure 1. Study flow diagram.

Data extraction and management

Two review authors (ZK, XL) independently extracted data using a piloted data extraction form. We extracted data on study characteristics including methods, participants, interventions and outcomes (see the '[Characteristics of included studies](#)' table). We resolved any disagreements by discussion. We have listed the formulation contents of the included studies and herb names in three languages in [Table 1](#) and [Table 2](#).

Assessment of risk of bias in included studies

Two review authors (ZJ, XL) independently performed the 'Risk of bias' assessments using the Cochrane 'Risk of bias' tool, [Higgins 2011](#), to assess the following domains.

1. Sequence generation: randomized (for example, by computer, random number tables or drawing lots) or method of randomization not described (we excluded quasi-RCTs).
2. Allocation concealment: low risk of bias (for example, by third party, sealed opaque envelopes); high risk of bias (for example, open list of allocation codes); unclear risk of bias (for example, not stated, or 'envelopes' stated without further description).
3. Blinding of participants, personnel and outcome assessors.
4. Completeness of outcome data.
5. Selective outcome reporting.
6. Other sources of bias.

Measures of treatment effect

We only measured dichotomous data in this review. We compared the outcome measures for binary data by calculating Peto-odds ratios (Peto OR) with 95% confidence intervals (CI). To measure the treatment effect we conducted intention to treat (ITT) analyses.

Unit of analysis issues

We planned that we would assess any studies with non-standard designs, such as cluster-RCTs, to avoid unit-of-analysis errors including: recruitment bias, baseline imbalance, loss of clusters, incorrect analysis and comparability with individually RCTs.

Dealing with missing data

We attempted to contact the trial authors to ask for missing data, but were unsuccessful. We imputed outcomes where data were missing in the present review. We assumed failure for drop-outs in the treatment group and success for drop-outs in the control group.

Assessment of heterogeneity

We considered whether the clinical and methodological characteristics of the included studies were sufficiently similar for meta-analysis to provide a clinically meaningful summary. We assessed statistical heterogeneity by using the χ^2 test with a 10% level of statistical significance and by the I^2 statistic to estimate the total variation across studies that is due to heterogeneity rather than chance. We considered less than 25% to indicate low level heterogeneity; 25% to 50% as a moderate level; and greater than 50% to indicate substantial heterogeneity ([Higgins 2002](#); [Higgins 2011](#)).

Assessment of reporting biases

In view of the difficulty of detection of and correction for publication bias and other reporting biases, we planned to

minimize their potential impact by ensuring a comprehensive search for eligible studies and by being alert for duplication of data. If there were 10 or more studies in an analysis, we planned to use a funnel plot to explore the possibility of small study effects (a tendency for estimates of the intervention effect to be more beneficial in smaller studies).

Data synthesis

We planned that two review authors (ZK, ZJ) would pool data if studies were sufficiently similar, using Review Manager (RevMan) ([RevMan 2014](#)). If pooling was inappropriate, we planned to perform only descriptive analysis. We planned to use a fixed-effect model unless there was substantial heterogeneity, in which case we would use a random-effects model.

We planned to combine the data from primary studies in the following comparisons:

1. CHM versus clomiphene.
2. CHM + clomiphene versus clomiphene.
3. CHM + follicle aspiration + ovulation induction versus follicle aspiration + ovulation induction.
4. CHM + LOD versus LOD.

Subgroup analysis and investigation of heterogeneity

Where data were available, we planned to conduct subgroup analyses to determine the separate evidence within the following subgroups.

1. Different co-interventions.
2. Different treatment strategies.
3. The duration of intervention or follow-up.
4. Women with or without insulin resistance.
5. Women with or without obesity.
6. Ethnicity.

If we detected substantial heterogeneity, we planned to explore possible explanations in sensitivity analyses and to take any statistical heterogeneity into account when we interpreted the results, especially if there was any variation in the direction of effect.

Sensitivity analysis

We planned to conduct sensitivity analyses for the primary outcomes to determine whether the conclusions were robust to arbitrary decisions made regarding eligibility and analysis. These analyses would include consideration of whether the review conclusions would have differed under the following circumstances.

1. We restricted eligibility to studies without high risk of bias.
2. We adopted a random-effects model
3. We restricted eligibility to studies without commercial funding.

RESULTS

Description of studies

We have reported the characteristics of the included and excluded studies in the '[Characteristics of included studies](#)' table and the '[Characteristics of excluded studies](#)' table.

Results of the search

The search from inception to June 2016 retrieved 899 articles (excluding duplications): 163 in English and 736 in Chinese. Of these 899 articles, 146 were animal or experimental studies, 55 were non-polycystic ovarian syndrome (PCOS) studies, 69 were non-Chinese herbal medicine (CHM) studies, 92 included participants who were adolescent or had PCOS without infertility or had no wish to conceive, 19 were before-and-after studies, 18 were reviews, nine were case-control studies, nine were case reports, two were cross-sectional studies, three were parallel non-randomized controlled studies, 27 were systematic reviews, 17 were unrelated studies and 3 were duplications. Finally, 430 articles were potentially eligible and we retrieved the full-text of these articles. Seven studies met our inclusion criteria (Li 2007; Ye 2007; Liang 2008; Ma 2009a; Li 2012a; NCT01116167; ChiCTR-IOR-16008557); one was a new study to this review (Li 2012a), and two were ongoing studies (NCT01116167; ChiCTR-IOR-16008557). Finally, we included five studies in this review (Li 2007; Ye 2007; Liang 2008; Ma 2009a; Li 2012a), and excluded 426 articles. See the 'Characteristics of included studies' table and the 'Characteristics of excluded studies' table for further details.

We prepared a PRISMA flow diagram to describe the articles found from our searches (Figure 1).

Included studies

Study design

All five included studies were conducted and published in China. One was a double clinical centre design (Liang 2008), and the other four were single-centre studies. Three studies used two-arm parallel groups (Liang 2008; Ma 2009a; Li 2012a), and the other two studies used three-arm parallel groups (Li 2007; Ye 2007). The range of study duration was from one year to four years. Each included study reported the inclusion and exclusion criteria. Drop-outs and withdrawals occurred in three studies for different reasons (Li 2007; Liang 2008; Ma 2009a).

Participants

In this review update, the five studies included a total of 414 participants. Sample size ranged from 40 to 170. All participants were women of reproductive age, with PCOS (according to Rotterdam criteria) and subfertility. Furthermore, two included studies also had the inclusion criterion that participants were resistant to Western medicines for ovulation induction (Ye 2007; Liang 2008). The baseline characteristics among groups were comparable for each study.

Interventions

Two studies used Chinese patent drugs (Li 2007; Li 2012a), and the other three included studies used Chinese herbal formulas. We

have listed the contents of each CHM preparation in Table 1, and the names of each herbal medicinal in three languages in Table 2. The treatment duration was less than six menstrual cycles for all included studies. However, the duration of follow-up was three months (Liang 2008), one year (Ye 2007), and unclear (Li 2007; Ma 2009a; Li 2012a), respectively.

1. CHM versus clomiphene:
 - a. one study compared CHM versus clomiphene (Li 2007);
 - b. one study compared CHM plus laparoscopic ovarian drilling (LOD) and clomiphene plus LOD (Ye 2007).
2. CHM + clomiphene versus clomiphene:
 - a. two studies compared CHM + clomiphene versus clomiphene (Li 2007; Li 2012a);
 - b. one study compared CHM + ethinylloestradiol cyproterone acetate (CEA) + clomiphene versus CEA + clomiphene (Ma 2009a).
3. CHM + follicle aspiration + ovulation induction versus follicle aspiration + ovulation induction:
 - a. one study compared CHM+follicle aspiration+ovulation induction versus follicle aspiration+ovulation induction (Liang 2008).
4. CHM + LOD versus LOD:
 - a. one study compared CHM + LOD versus LOD (Ye 2007).

Outcomes

1. No study reported live birth rate.
2. All five included studies reported clinical pregnancy rate (Li 2007; Ye 2007; Liang 2008; Ma 2009a; Li 2012a).
3. One study reported ovulation rate (Ye 2007).
4. One study reported adverse events (luteinized unruptured follicle syndrome (LUFS), ovarian hyperstimulation syndrome (OHSS) and multiple pregnancy) (Liang 2008).

Excluded studies

We excluded 424 studies from the review for the following reasons (see the 'Characteristics of excluded studies' table for further details).

1. 122/424 were not RCTs.
2. 178/424 had participants that were not of interest to this review.
3. 50/424 reported interventions that were not of interest to this review.
4. 71/424 reported outcomes that were not of interest to this review.
5. 3/424 were duplicates of already excluded studies.

Risk of bias in included studies

We have summarized the risks of bias of the included studies in Figure 2 and Figure 3.

Figure 2. 'Risk of bias' graph: review authors' judgments about each 'Risk of bias' item presented as percentages across all included studies.

Figure 3. 'Risk of bias' summary: review authors' judgments about each 'Risk of bias' item for each included study.

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding (performance bias and detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias
Li 2007	?	?	+	+	?	+
Li 2012a	+	?	?	?	?	+
Liang 2008	+	-	-	?	?	+
Ma 2009a	+	?	?	+	?	+
Ye 2007	+	?	?	?	?	+

Allocation

Four studies were at low risk of selection bias related to sequence generation as they used random numbers tables (Ye 2007; Liang 2008; Ma 2009a; Li 2012a). One study did not describe the method used and was at unclear risk of this bias (Li 2007).

One study was at high risk of selection bias related to allocation concealment as the random number table was open (Liang 2008). Four studies were at unclear risk of selection bias as they did not report adequate details to establish whether an appropriate method of allocation concealment had been used (Li 2007; Ye 2007; Ma 2009a; Li 2012a).

Blinding

1. One study used placebo drugs and described blinding of participants and outcome assessors. We judged it to be at low risk of detection bias (Li 2007).
2. One study used no blinding, which the study authors confirmed. We judged it to be at high risk of bias (Liang 2008).
3. Three studies did not mention blinding. We judged them to be at unclear risk of bias (Ye 2007; Ma 2009a; Li 2012a).

Incomplete outcome data

Two studies analysed all or most (over 95%) women randomized and we judged them to be at low risk of bias (Li 2007; Ma 2009a). One study analysed only 91% of women randomized and we judged it to be at unclear risk of bias (Liang 2008). Two studies did not mention

drop-outs or withdrawals, and we judged them as at unclear risk of attrition bias (Ye 2007; Li 2012a). The reasons for attrition included moving to another place, pelvic inflammation and conversion to in vitro fertilization-embryo transfer (IVF-ET).

Selective reporting

The risk of selective reporting was unclear in each of the included studies, as the protocols of the included studies were unavailable. The five studies did not assess live birth rate. Only one study reported adverse events (Liang 2008). We were unable to obtain detailed information from the primary study authors. The outcomes of these five included studies might be influenced by the bias of selective reporting or publication bias, and we rated all as at unclear risk of selective reporting bias.

Other potential sources of bias

We did not identify any other potential sources of bias in the included studies, and judged each of the included studies at low risk of other potential sources of bias.

Publication bias

As there were fewer than 10 included studies, we did not assess potential publication bias using a funnel plot or other corrective analytical methods (Egger 1997).

Effects of interventions

See: **Summary of findings for the main comparison** Chinese herbal medicine (CHM) versus clomiphene for subfertile women with polycystic ovarian syndrome (PCOS); **Summary of findings 2** Chinese herbal medicine (CHM) + clomiphene versus clomiphene for subfertile women with polycystic ovarian syndrome (PCOS); **Summary of findings 3** Chinese herbal medicine (CHM) + follicle aspiration + ovulation induction versus follicle aspiration + ovulation induction for subfertile women with polycystic ovarian syndrome (PCOS); **Summary of findings 4** Chinese herbal medicine (CHM) + laparoscopic ovarian drilling (LOD) versus LOD for subfertile women with polycystic ovarian syndrome (PCOS)

We extracted summary data from the five included studies (Li 2007; Ye 2007; Liang 2008; Ma 2009a; Li 2012a). The clinical heterogeneity, which we have documented in the 'Characteristics of included studies' table, was high among these studies, especially regarding the interventions used. We therefore subgrouped the analyses by co-intervention (see Analysis 1.1 and Analysis 2.1).

1. CHM versus clomiphene

Two studies made this comparison (Li 2007; Ye 2007); one of these studies administered LOD in both arms (Ye 2007).

Primary outcome

1.1 Live birth rate

None of the included studies reported this outcome.

Secondary outcomes

1.2 Pregnancy rate

When we combined the studies, there was no evidence of a difference between CHM versus clomiphene (with or without LOD in both arms) (odds ratio (OR) 1.98, 95% confidence interval (CI) 0.78 to 5.06; two studies, 90 participants, I^2 statistic = 0%, *very low quality evidence*). See Analysis 1.1.

1.3 Ovulation rate

There was no evidence of a difference between CHM versus clomiphene (OR 1.42, 95% CI 0.19 to 10.49; one study, 30 participants). See Analysis 1.2.

1.4 Adverse events

None of the included studies reported this outcome.

2. CHM + clomiphene versus clomiphene

Three studies made this comparison (Li 2007; Ma 2009a; Li 2012a); one of the studies administered ethinyloestradiol cyproterone acetate in both study arms (Ma 2009a).

Primary outcome

2.1 Live birth rate

None of the included studies reported this outcome.

Secondary outcomes

2.2 Pregnancy rate

There was a higher rate of pregnancy in the CHM plus clomiphene group (OR 2.62, 95% CI 1.65 to 4.14; three studies, 300 participants, I^2 statistic = 0%, *low quality evidence*). See Analysis 2.1 and Figure 4.

Figure 4. Forest plot of comparison: 3 CHM + clomiphene versus clomiphene, outcome: 3.1 Pregnancy rate (per woman).

2.3 Ovulation rate

None of the included studies reported this outcome.

2.4 Adverse events

None of the included studies reported this outcome.

3. CHM + follicle aspiration + ovulation induction versus follicle aspiration + ovulation induction

One study made this comparison (Liang 2008).

Primary outcome

3.1 Live birth rate

Liang 2008 did not report this outcome.

Secondary outcomes

3.2 Pregnancy rate

There was no evidence of a difference between the two groups (OR 1.60, 95% CI 0.46 to 5.52; one study, 44 women, *very low quality evidence*). See Analysis 3.1.

3.3 Ovulation rate

Liang 2008 did not report this outcome.

3.4 Adverse events

Only one study reported adverse events (Liang 2008). There was no evidence of a difference between CHM plus follicle aspiration, ovulation induction and follicle aspiration plus ovulation induction for LUFs (OR 0.60, 95% CI 0.06 to 6.14; one study, 44 women, *very low quality evidence*), OHSS (OR 0.16, 95% CI 0.00 to 8.19; one study, 44 women, *very low quality evidence*) or multiple pregnancy (OR 0.60, 95% CI 0.06 to 6.14; one study, 44 women, *very low quality evidence*). See Analysis 3.2, Analysis 3.3 and Analysis 3.4. The severity of adverse events was not reported and no other data on adverse events were available.

4. CHM + LOD versus LOD

One study made this comparison (Ye 2007).

Primary outcome

4.1 Live birth rate

Ye 2007 did not report this outcome.

Secondary outcomes

4.2 Pregnancy rate

There was no evidence of a difference between the groups (OR 3.5, 95% CI 0.72 to 17.09; one study, 30 women, *very low quality evidence*). See Analysis 4.1.

4.3 Ovulation rate

There was no evidence of a difference between the groups (OR 2.43, 95% CI 0.39 to 15.08; one study, 30 women). See Analysis 4.2.

4.4 Adverse events

Ye 2007 did not report this outcome.

DISCUSSION

Summary of main results

There is insufficient evidence to support the use of CHM in treating women with polycystic ovarian syndrome (PCOS) and subfertility. None of the included studies reported live birth rate, and only very limited data were available for the other review outcomes.

There was very limited evidence to suggest the following.

1. For women with PCOS and infertility, the efficacy of clomiphene in improving pregnancy rate (per woman), with or without pretreatment of ethinyloestradiol cyproterone acetate, may be enhanced by Chinese herbal medicine (CHM). Inversely, that of CHM may not be enhanced by clomiphene.
2. For women with PCOS, infertility and resistance to Western drugs of ovulation induction, the efficacy of follicle aspiration or laparoscopic ovarian drilling (LOD) in improving pregnancy rate may not be strengthened by CHM. This may indicate that women resistant to Western drugs of ovulation induction may also be resistant to CHM of ovulation induction.

3. For women with PCOS, infertility and resistance to ovulation induction, there is not enough evidence to support the use of CHM in improving ovulation rate.
4. Only one included study reported adverse events, including luteinized unruptured follicle syndrome (LUFS), ovarian hyperstimulation syndrome (OHSS) and multiple pregnancy, but did not indicate the severity of the adverse events. None of the included studies reported some of the adverse events thought to be associated with CHM (e.g. impairment of liver and kidney, allergies). Therefore, the safety of CHM for women with PCOS and subfertility remains unclear.

There was very limited evidence that the addition of CHM to clomiphene was associated with improved clinical pregnancy outcomes but no other evidence of any other effect. This finding requires extremely cautious interpretation.

Overall completeness and applicability of evidence

The included studies only partially addressed the objectives of this review. We were unable to reach definite conclusions due to the lack of data for each comparison group. The high heterogeneity of CHM preparations in the included studies may limit the generalizability of the results regarding the effectiveness of CHM for subfertile women with PCOS in general. The included studies failed to report the most important outcome, which is live birth rate. Future studies should report major clinical outcomes such as live birth, clinical pregnancy and adverse events.

The included studies were clinically heterogeneous and differed in (or failed to report) factors such as the duration of treatment, CHM formula, dosage and length of follow-up. Moreover no studies compared CHM with the first-line interventions for PCOS, such as diet control and exercise. These interventions should be compared with CHM in future studies.

Quality of the evidence

The quality of the evidence for most comparisons was very low. The main limitations in the evidence were failure to report live birth or adverse events, failure to describe study methods in adequate detail and imprecision, with very low event rates and wide confidence intervals (CIs).

None of the included studies clearly reported blinding or described their method of allocation concealment, and some did not clearly report drop-out rates. Only one included study used placebo drugs (Li 2007), so the study may have used blinding. However, we were unable to obtain detailed information from the study authors.

Protocols were not registered in clinical trial registers, so we could not evaluate the risk of selective reporting bias.

Potential biases in the review process

In order to limit bias in the review process, the Cochrane Gynaecology and Fertility Group guided and developed the

literature search. We did not apply any restrictions on language to the searches. Two review authors (ZK, ZJ) independently performed study selection, 'Risk of bias' assessments and data collection but without blinding. We resolved any disagreements by discussion with a third review author (WT). We attempted to obtain missing information and data by contacting the primary study authors but were not always successful. Thus, we excluded those studies that we could not classify as randomized controlled trials (RCTs) due to lack of information.

In our review, we performed intention-to-treat (ITT) analyses by assuming failure for drop-outs in treatment group and success for drop-outs in the control group.

The review authors had no conflicts of interest.

Agreements and disagreements with other studies or reviews

In the future, well-designed RCTs with large sample size are warranted to confirm or refute the current evidence. There are no other systematic reviews on CHM for subfertile women with PCOS.

AUTHORS' CONCLUSIONS

Implications for practice

There is insufficient evidence to support the use of CHM for women with PCOS and subfertility. No data are available on live birth, and there is no consistent evidence to indicate that CHM improves fertility outcomes. However there is very limited low quality evidence to suggest that the addition of CHM to clomiphene may improve pregnancy rates. There is insufficient evidence on adverse effects to indicate whether CHM is safe.

Implications for research

Study authors should report methodology in detail, such as randomization and allocation concealment methods. Well-designed and well-conducted RCTs with double blinding should be conducted. The duration of follow-up for assessing outcomes should also be reported. The CHM formula and dosage should be reported. Future research should expand sample size, evaluate live birth rate and other safety indexes. Interventions for PCOS, such as diet control and exercise, should be compared with CHM in future studies.

ACKNOWLEDGEMENTS

We thank Jane Clarke (Managing Editor of the original review), Helen Nagels (Managing Editor of this updated review), Marian Showell (Information Specialist) and the editorial board of the Cochrane Gynaecology and Fertility Group (CGF) for their invaluable assistance in developing this review.

REFERENCES

References to studies included in this review

Li 2007 {published data only}

Li XB, Li LY, Fu F, Hu XD. Clinical research on Lingzhu infusion and Shenqi capsule sequential therapy for polycystic ovarian syndrome. *Journal of Traditional Chinese Medicine* 2007;**48**(12):1079-81.

Li 2012a {published data only}

Li Y. [Clinical observation of compound Xuanju capsule combined with clomiphene for polycystic ovary syndrome]. *Shanxi Journal of Traditional Chinese Medicine* 2012;**28**(9):21-2.

Liang 2008 {published data only}

Liang RN, Liu J, Lu J. Treatment of refractory polycystic ovary syndrome by Bushen Huoxue method combined with ultrasound guided follicle aspiration. *Zhongguo Zhong Xi Yi Jie He Za Zhi [Chinese Journal of Integrated Traditional and Western Medicine]* 2008;**28**(4):314-7.

Ma 2009a {published data only}

Ma HX, Lai MH, Liu H, Song CX, Pan SY. Clinical observation of integrated therapy of Chinese and western medicine on 85 patients with infertility caused by PCOS. *Journal of TCM University of Hunan* 2009;**29**(3):47-8, 78.

Ye 2007 {published data only}

Ye DM, Xu LM, Lu RL. Clinical observation on laparoscopic ovarian electrocauterization combined with cycle treatment of traditional Chinese medicine for type phlegm-damp refractory polycystic ovary syndrome. *Journal of Guangzhou University of TCM* 2007;**24**(6):445-8.

References to studies excluded from this review

An 2009 {published data only}

An XQ, Liu GJ, Zhang JY, Cai FJ, An XT, Jiao SF. Curative efficacy of letrozole combined with traditional chinese medicine for ovulation dysfunction caused by polycystic ovarian syndrome. *China Pharmacy* 2009;**20**(29):2289-91.

An 2012 {published data only}

An XQ. Clinical observation of compound XuanJu capsule combine with letrozole tablets treat 36 cases ovulation disorder in polycystic ovary syndrome. *Hebei Journal of Traditional Chinese Medicine* 2012;**34**(7):1053-4.

Arentz 2014 {published data only}

Arentz S, Abbott JA, Smith CA, Bensoussan A. Herbal medicine for the management of polycystic ovary syndrome (PCOS) and associated oligo/amenorrhoea and hyperandrogenism; a review of the laboratory evidence for effects with corroborative clinical findings. *BMC Complementary and Alternative Medicine* 2014;**14**:511.

Bablis 2006 {published data only}

Bablis P, Pollard H, Monti DA. Resolution of anovulation infertility using neuro emotional technique: a report of 3 cases. *Journal Of Chiropractic Medicine* 2006;**5**(1):13-21.

Bai 2011 {published data only}

Bai HY. Combine traditional Chinese and western medicine treatment of 94 cases of polycystic ovary syndrome. *Seek Medical And Ask The Medicine* 2011;**9**(6):169.

Bao 2009 {published data only}

Bao WY. Treatment of polycystic ovary syndrome with nourishing kidney and resolving phlegm combined with blood activating. *Tianjin Journal of Traditional Chinese Medicine* 2009;**26**(5):375-6.

Bao 2014 {published data only}

Bao Y, Bao Y, Piao CL, He Z. [The clinical efficacy of polycystic ovarian syndrome insulin resistance and insulin resistance in non-treated control analysis]. *Maternal & Child Health Care of China* 2014;**29**(30):4913-4.

Bei 2010 {published data only}

Bei GZ. Clinical observation of jianpiqushi therapy joint healthy life style treating obesity polycystic ovarian syndrome. Guang Zhou University of Chinese Medicine 2010.

Cai 2006 {published data only}

Cai LH. [Clinical observation on shugan qingjie decoction in treating adolescent PCOS]. *Beijing Journal of TCM* 2006;**25**(6):323-6.

Cai 2011 {published data only}

Cai XH. The kidney and liver soup treatment kidney empty stomach tone of childbearing age-to clinical observation of polycystic ovarian syndrome. Shandong University of Traditional Chinese Medicine 2011.

Cai 2012 {published data only}

Cai XQ, Chen HJ, Liu Y. Combine traditional Chinese and western medicine treatment of infertility in 35 cases of clinical observation. *Zhejiang Journal of Traditional Chinese Medicine* 2012;**47**(6):411.

Cai 2014 {published data only}

Cai W. [Integrative Medicine 74 cases of polycystic ovary syndrome]. *World Health Digest* 2014, (15):115-6.

Cao 2010 {published data only}

Cao WP. The Kidney and Liver Soup Treatment Kidney Empty Stomach Tone of Childbearing Age-to Clinical Observation of Polycystic Ovarian Syndrome. Shandong Traditional Chinese Medicine University 2010.

Cao 2012 {published data only}

Cao ZY, Zhang XF. The kidney and treatment of kidney Yang deficiency and phlegm wet expectorant method the clinical research of infertility caused by polycystic ovary syndrome. *Health Must-Read Magazine* 2012;**11**(7):1.

Chan 2006a {published data only}

Chan CC, Koo MW, Ng EH, Tang OS, Yeung WS, Ho PC. Effects of Chinese green tea on weight, and hormonal and biochemical profiles in obese patients with polycystic ovary syndrome-a

randomized placebo controlled trial. *Journal of the Society for Gynecologic Investigation* 2006;**13**(1):63-8.

Chan 2006b {published data only}

Chan LY, Lau TK, Fung TM, Chow KM. Re: Effects of rosiglitazone on hormonal profile and ovulatory function in Chinese women with polycystic ovary syndrome. *The Australian & New Zealand Journal Of Obstetrics & Gynaecology* 2006;**46**(2):170-1.

Chen 2005 {published data only}

Chen LS, Zhou JT. [Clinical research on combined therapy of Diane-35 with modified yougui pill on polycystic ovarian syndrome]. *Zhongguo Zhong Xi Yi Jie He Za Zhi [Chinese Journal of Integrated Traditional and Western Medicine]* 2005;**25**(9):794-6.

Chen 2006a {published data only}

Chen L. [Comparison of efficacy of two ovulation inducing methods in treating PCOS with infertility]. *Modern Journal of Integrated Traditional Chinese and Western Medicine* 2006;**15**(8):1035-6.

Chen 2006b {published data only}

Chen JL, Yang ZQ, Liu L. Clinical observation on 30 cases of hyperpimelic menostasia treated by Huatan tiaochong decoction. *Hunan Journal of TCM* 2006;**22**(6):10-1.

Chen 2007 {published data only}

Chen D, Chen SR, Shi XL, Guo FL, Zhu YK, Li S, et al. Clinical study on needle pricking therapy for treatment of polycystic ovarian syndrome. *Zhongguo Zhen Jiu [Chinese Acupuncture & Moxibustion]* 2007;**27**(2):99-102.

Chen 2008 {published data only}

Chen XP, Xie B, Zheng JP, Chen LF. Effect of nourishing kidney and dispersing stagnated hepatoqi decoction on symptoms and sex hormone of polycystic ovary syndrome patients with renal deficiency and hepatic stagnation. *Chinese Journal of Information on TCM* 2008;**15**(5):17-9.

Chen 2009 {published data only}

Chen J, Cui W, Li J, Sun W. Investigation on effect of electroacupuncture intervention on in vitro fertilization and embryo transfer of patients with polycystic ovarian syndrome. *Maternal and Child Health Care of China* 2009;**24**(30):4262-4.

Chen 2010a {published data only}

Chen XH, Huang BC, Kui Y. Abdominal needle combined with Chinese medicine treatment to insulin in patients with polycystic ovary syndrome. Guangdong institute of acupuncture and moxibustion academic conference. 2010.

Chen 2010b {published data only}

Chen H. Clinical observation of Yang Yin Qing Re Method for insulin resistance in polycystic ovary syndrome. Shanghai University of Chinese Medicine 2010.

Chen 2011a {published data only}

Chen PL. Clinical Research on Treatment of Polycystic Ovary Syndrome (Type of Phlegm Stagnation) with Modified Erchen Decoction. Guangzhou University of Chinese Medicine 2011.

Chen 2011b {published data only}

Chen RR. TKRPM and resolving phlegm, dispelling stasis, combined with LE on treating the PCOS patients with ovulatory dysfunction.. Nanjing University of Traditional Chinese Medicine 2011.

Chen 2011c {published data only}

Chen XF. [Clinical research on laparoscopic surgery supplemented by Luo's ovulation soup for the treatment of kidney Yang deficiency type polycystic ovary syndrome]. Fujian University of Traditional Chinese Medicine 2011.

Chen 2012a {published data only}

Chen N. Huo Xue Li Shi decoction in the treatment of phlegm and blood stasis type of PCOS clinical observation. Shandong University of Traditional Chinese Medicine 2012.

Chen 2012b {published data only}

Chen RJ. Clinical research about Xiao Nang Tiao Jing1Hao Fang on treatment of PCOS of stagnation of liver-qi type with the disease and to probe into Its mechanism of action. Hunan University of Chinese Medicine 2012.

Chen 2012c {published data only}

Chen WY. Clinical observation on treating obese women with PCOS infertility in the integrative medicine. *Clinical Journal of Chinese Medicine* 2012;**4**(16):96-7.

Chen 2012d {published data only}

Chen LS, Chen J, Li SM. [Combine traditional Chinese and western medicine treatment of refractory polycystic ovary syndrome infertility 31 cases of clinical analysis]. *Journal of Fujian University of Traditional Chinese Medicine* 2012;**22**(6):10-3.

Chen 2013 {published data only}

Chen FL. [41 cases of anovulatory infertility clinical observation of combined Chinese and Western medicine to treat]. *Chinese and Foreign Medical Research* 2013, (6):38-9.

Chen 2014a {published data only}

Chen QZ. [Due to polycystic ovary syndrome diagnosis and treatment of infertility]. *China Health Care & Nutrition* 2014, (6):3863.

Chen 2014b {published data only}

Chen Y. [Clinical observation of Chinese medicine treatment of polycystic ovary syndrome]. *Medical Information* 2014;**6**(20):477-8.

Chen 2014c {published data only}

Chen R, Wang C, Yan QY. Observation on therapeutic effect of polycystic ovary syndrome of damp-phlegm constitution treated with embedding therapy on back-shu points and front-mu points combined with needle-pricking therapy on Sifeng (EX-UE 10). *Zhongguo Zhen Jiu [Chinese Acupuncture & Moxibustion]* 2014;**34**(4):355-8.

Chen 2015 {published data only}

Chen XY. [The efficacy Huoxue Bushen decoction combined with western medicine treatment of polycystic ovary syndrome]. *Journal of Clinical Psychosomatic Diseases* 2015;**X21**(1):24-5.

Chen 2016 {published data only}

Chen J, Feng S, Zeng J, Wu X, Yang M, Tang H, et al. Effectiveness of electroacupuncture for polycystic ovary syndrome: study protocol for a randomized controlled trial. *Trials* 2016;**17**:256.

Cheng 2009 {published data only}

Cheng MX, Li XL, Tong Q, Xu C. Therapeutic effect of traditional Chinese medicine on polycystic ovary syndrome. *Journal of Difficult and Complicated Cases* 2009;**8**(1):26-7.

Cheng 2014 {published data only}

Cheng X, Guo J, Xie J. Association between levels of serum leptin and insulin resistance in patients with polycystic ovary syndrome. *Zhonghua Liu Xing Bing Xue Za Zhi* 2014;**35**(12):1389-91.

Cheng 2015 {published data only}

Cheng F, Zhang HL, Cheng L. Clinical research of Xiaonang Tiaoqing decoction combined with chloramphenicol in the treatment of polycystic ovarian syndrome. *China Journal of Chinese Medicine* 2015, (7):1027-9.

Chu 2013 {published data only}

Chu ES, Sze SC, Cheung HP, Liu Q, Ng TB, Tong Y. An in vitro and in vivo investigation of the antimetastatic effects of a Chinese medicinal decoction, Erxian Decoction, on human ovarian cancer models. *Integrative Cancer Therapies* 2013;**12**(4):336-46.

Craig 2015 {published data only}

Craig LB, Peck JD, Zhao D, Hansen KR. Clomid stair-step protocol may shorten the time to ovulation but not to pregnancy: A randomized clinical trial. *Fertility and Sterility Conference: 71st Annual Meeting of the American Society for Reproductive Medicine, ASRM 2015 Baltimore, MD United States* 2015;**104**(3):e97.

Cui 2012 {published data only}

Cui XF, Mo XY. [Kidney and eliminating phlegm method to treat high androgen levels, 50 cases of polycystic ovary syndrome]. *Jilin Journal of Traditional Chinese Medicine* 2012;**32**(4):378-9.

Dang 2012 {published data only}

Dang HM, Liu YQ, Liu RX, Chen W. ["Tiao jing zhu yun fang" combined with letrozole in women with polycystic ovary syndrome follicular development in patients with infertility and ovarian, uterine hemodynamic effects]. Eighth National Integrative Medicine Obstetrics and Gynecology Academic Conference. 2012.

Deng 2008 {published data only}

Deng AL, Zhou ZM, Jiang HZ. [Clinical research of integration of traditional Chinese medicine and western medicine in treating infertility caused by PCOS]. *Hubei Journal of Traditional Chinese Medicine* 2008;**30**(12):18-9.

Deveci 2015 {published data only}

Deveci C, Demir B, Sengul O, Dilbaz B, Goktolga U. Clomiphene citrate 'stair-step' protocol vs. traditional protocol in patients with polycystic ovary syndrome: a randomized controlled trial. *Archives of Gynecology & Obstetrics* 2015;**291**(1):179-84.

Ding 2015 {published data only}

Ding C-F, Chen W-Q, Zhu Y-T, Bo Y-L, Hu H-M, Zheng R-H. Circulating microRNAs in patients with polycystic ovary syndrome. *Human Fertility (Cambridge, England)* 2015;**18**(1):22-9.

Dong 2009 {published data only}

Dong CY, Tian YM. [Clinical observation on TCM combined with Western medicine in treating PCOS]. *Inner Mongol Journal of Traditional Chinese Medicine* 2009;**24**(7):84-5.

Dong 2010 {published data only}

Dong CY, Tian YM. [Clinical observation of "sheng ti gan qi fa" treat 30 cases of polycystic ovary syndrome]. *Hebei Journal of Traditional Chinese Medicine* 2010;**32**(1):38-9.

Du 2012 {published data only}

Du GH. Clinical study on the effect of kidney-nourishing and phlegm-resolving method combined with lifestyle adjustment treating obese women with polycystic ovary syndrome. Nanjing University of Chinese Medicine 2012.

Du 2013 {published data only}

Du AZ. Integrative Medicine 28 cases of polycystic ovary syndrome. *Chinese Medicine Morden Distance Education of China* 2013, (16):63-4.

Fang 2004 {published data only}

Fang RD, Li XR. [Clinical observation on Chinese herbal medicine combined with western medicine in treating 50 cases of PCOS]. *Chinese Journal of Traditional Medical Science and Technology* 2004;**11**(4):233-4.

Feng 2009a {published data only}

Feng CF. Treating 50 cases of phlegm-damp type polycystic ovarian syndrome with combination of Chinese medicine and western medicine. *Acta Chinese Medicine* 2009;**24**(6):60-1.

Feng 2009b {published data only}

Feng CF, Qin XJ, Xu JY. [Clinical research on Chinese herbal medicine in treating 20 cases with PCOS]. *Practical Clinical Journal of Integrated Traditional Chinese and Western Medicine* 2009;**9**(2):42-4.

Feng 2011a {published data only}

Feng X. The clinical observation of Chinese and Western integrative medicine treating infertility caused by polycystic ovarian syndrome (PCOS). Jilin University 2011.

Feng 2011b {published data only}

Feng J, Zhang XF. [Clinical observation of combine traditional Chinese and Western medicine treat polycystic ovary syndrome with infertility]. *Jilin Journal of Traditional Chinese Medicine* 2011;**31**(9):858-60.

Fu 2012 {published data only}

Fu SW. [Comprehensive treatment of polycystic ovary syndrome in TCM]. Xinli Yisheng 2012, issue 7:306-7.

Gao 2009 {published data only}

Gao TY, Wang J, Wang J, Wang WH, Wang WJ, Lun ZJ. Effect of acupuncture assist IUI treatment on the outcome of infertility patients with polycystic ovarian syndrome. *Reproduction and Contraception* 2009;**129**(10):680-2.

Gao 2011a {published data only}

Gao XL. [Integrative medicine clinical observation of 90 cases of infertility polycystic ovary syndrome]. *Harebin Medical Journal* 2011;**31**(2):123-4.

Gao 2011b {published data only}

Gao YH. Clinical study on treatment of polycystic ovarian syndrome in type of splenic deficiency and phlegm-damp with Wu Ji powder. Nanjing University of Chinese Medicine 2011.

Ghavi 2015 {published data only}

Ghavi F, Shakeri F. Effects of fennel on serum hormone levels in students with polycystic ovary syndrome. *Avicenna Journal of Phytomedicine* 2015;**5**:42-3.

Gong 2012 {published data only}

Gong XH. Observation on effect of therapy of catgut implantation at acupoint on PCOS. *Chinese Archives of Traditional Chinese Medicine* 2012;**30**(9):2001-2.

Grant 2010 {published data only}

Grant P. Spearmint herbal tea has significant anti-androgen effects in polycystic ovarian syndrome. A randomized controlled trial. *Phytotherapy Research* 2010;**24**(2):186-8.

Gu 2015 {published data only}

Gu LY, Wang Q, Du L, Zha QS, Gao H. [Metformin self side polycystic ovary syndrome phlegm clinical observation of 72 cases of amenorrhea]. *Chinese Journal for Clinicians* 2015, (3):72-4.

Guo 2008 {published data only}

Guo LC, Hu B. The study on traditional Chinese medicine in the treatment of 2cases of polycystic ovary syndrome complicated with 2-type diabetes mellitus. *Hebei Journal of TCM* 2008;**30**(2):130-1.

Guo 2009 {published data only}

Guo XY, Zhong HM, He HL, Liang HF. Clinical research of effect of wenshenhuatan capsule combined Diane-35 on polycystic ovarian syndrome. *Modern Medicine & Health* 2009;**25**(11):1606-7.

Guo 2011a {published data only}

Guo AP. [Integrative clinical observation on treatment of polycystic ovary syndrome]. *Journal of China Traditional Chinese Medicine Information* 2011;**3**(17):395, 410.

Guo 2011b {published data only}

Guo SX, Wang WJ, Gui SQ. Efficacy of Chinese patent medicine Tian Gui Capsule in patients with polycystic ovary syndrome: a randomized controlled trial. *Journal of Chinese Integrative Medicine* 2011;**9**(9):965-72.

Haj-Husein 2016 {published data only}

Haj-Husein I, Tukan S, Alkazaleh F. The effect of marjoram (*Origanum majorana*) tea on the hormonal profile of women with polycystic ovary syndrome: a randomised controlled pilot study. *Journal Of Human Nutrition And Dietetics: The Official Journal Of The British Dietetic Association* 2016;**29**(1):105-11.

Han 2008 {published data only}

Han YQ. [Effective comparison of TCM combined with western medicine ovulation inducing methods in treating PCOS with infertility]. *China Practical Medicine* 2008;**3**(9):49-50.

Han 2011 {published data only}

Han L, Gong W. [Clinical analysis of Guishen pill treat polycystic ovary syndrome]. *Chinese Journal of Experimental Traditional Medical Formulae* 2011;**17**(24):216-7.

Han 2013a {published data only}

Han M. Clinical observation of integrative treatment of polycystic ovary syndrome. *World Latest Medicine Information* 2013, (31):205-6.

Han 2013b {published data only}

Han SX. Chinese medicine for reinforcing of kidney combined with diane- 35 for polycystic ovary syndrome. *China Medicine and Pharmacy* 2013, (6):84-5.

Han 2015 {published data only}

Han ZY. TCM therapy clinical observation period infertility polycystic ovary syndrome. *Yiyao Qianyan* 2015, (34):317-8.

Hao 2012 {published data only}

Hao JJ. Clinical research on treating kidney deficiency and blood stasis type polycystic ovarian syndrome with Chinese medicine installments therapy and clomiphene citrate. Shandong Traditional Chinese Medicine University 2012.

Harman 2001 {published data only}

Harman J, Ward M. The role of nutritional therapy in the treatment of equine Cushing's syndrome and laminitis. *Alternative Medicine Review: A Journal Of Clinical Therapeutic* 2001;**6** Suppl:S4-16.

Hassanzadeh Bashtian 2013 {published data only}

Hassanzadeh Bashtian M, Emami SA, Mousavifar N, Esmaily HA, Mahmoudi M, Mohammad Poor AH. Evaluation of fenugreek (*Trigonella foenum-graceum* L.), effects seeds extract on insulin resistance in women with polycystic ovarian syndrome. *Iranian Journal of Pharmaceutical Research* 2013;**12**(2):475-81.

He 2009 {published data only}

He HL, Xiao XF, Guo XY. Study of Wenshenhuatan capsule combined with Diane-35 on kidney asthenia an phlegmatic hygrois type polycystic ovary syndrome. *Modern Journal of Integrated Traditional Chinese and Western Medicine* 2009;**18**(23):2749-50, 2753.

He 2010 {published data only}

He P, Ding L. Clinical analysis of the treatment of anovulation patients with polycystic ovary syndrome by letrozole combined

- with traditional Chinese medicine. *Yunnan Journal of Traditional Chinese Medicine and Materia Medica* 2010;**31**(1):8-9.
- He 2014** {published data only}
He YY. [Infertility clinical efficacy in combination therapy with polycystic ovary syndrome]. *Chinese Journal of Trauma and Disability Medicine* 2014;**66**(17):102-3.
- Hou 2000** {published data only}
Hou JW, Yu J, Wei MJ. [Study on treatment of hyperandrogenism and hyperinsulinism in polycystic ovary syndrome with Chinese herbal formula "tiangui fang"]. *Zhongguo Zhong Xi Yi Jie He Za Zhi [Chinese Journal of Integrated Traditional and Western Medicine]* 2000;**20**(8):589-92.
- Hu 2009a** {published data only}
Hu ZH, Wang Y, Shen HB. Study on impact of inserting Qihai and Sanyinjiao point on the endocrine changes in insulin resisting PCOS. *Journal of Clinical Acupuncture & Moxibustion* 2009;**25**(4):1-2.
- Hu 2009b** {published data only}
Hu ZH, Wang Y. Study of the mechanism of acupuncture action on polycystic ovary syndrome of non-insulin resistance type. *Shanghai Journal of Acupuncture and Moxibustion* 2009;**28**(4):197-8.
- Hu 2014** {published data only}
Hu YY. [Clinical observation of 35 cases of polycystic ovary syndrome TCM Tiaozhou therapy combined therapy of metformin hydrochloride tablets]. *World Latest Medicine Information* 2014, (16):132-8.
- Hua 2003** {published data only}
Hua L, Wu YN, Zhang JM. [Clinical study of Yishen jianpi yangxue tongli therapy in treating polycystic ovary syndrome]. *Zhongguo Zhong Xi Yi Jie He Za Zhi [Chinese Journal of Integrated Traditional and Western Medicine]* 2003;**23**(11):819-22.
- Huang 2004** {published data only}
Huang YB, Gu ZT, Pan JY, Ruan CL, Lin QE. Clinical study on the treatment of clomiphene-resistant polycystic ovarian syndrome with laparoscopic ovarian drilling and drugs. *Chinese Journal of Practical Gynecology and Obstetrics* 2004;**20**(8):480-1.
- Huang 2006a** {published data only}
Huang LY, Xu CH. [Yishen huoxue method in treating 50 cases of PCOS with infertility]. *Jilin Journal of Traditional Chinese Medicine* 2006;**26**(12):19-20.
- Huang 2006b** {published data only}
Huang YY, Yang XW, Deng MD. [Clinical research of integration of traditional chinese medicine and western medicine in treating unmarried PCOS]. *New Journal of Traditional Chinese Medicine* 2006;**38**(4):58-9.
- Huang 2007** {published data only}
Huang M, Lai H, Lv FB, Pan BQ. [Massage, moxibustion combined with western medicine in treating PCOS]. *Chinese Manipulation and Qi Gong Therapy* 2007;**23**(1):5-6.
- Huang 2008** {published data only}
Huang S, Chen AP. Traditional Chinese medicine and infertility. *Journal of Community Medicine* 2008;**20**(3):211-5.
- Huang 2010** {published data only}
Huang XO. Clinical effect of treatment of Chinese medicine for sterility caused by polycystic ovarian syndrome. *Clinical Medical Engineering* 2010;**17**(4):32-3.
- Huang 2011a** {published data only}
Huang DL, Xu F, Wen YL, Xie NX. [Acupuncture and Chinese medicine treatment of polycystic ovary syndrome-induced infertility in 30 cases]. *Journal of New Chinese Medicine* 2011;**43**(8):86-6.
- Huang 2011b** {published data only}
Huang LH. Kidney and phlegm and dispelling stasis party clinical observation for the treatment of polycystic ovary syndrome with insulin resistance. Hubei University of Chinese Medicine 2011.
- Huang 2012a** {published data only}
Huang L. Traditional Chinese medicine of invigorating the kidney and resolving phlegm on the inhibin B level in patients with polycystic ovary syndrome. Guangzhou University of Chinese Medicine 2012.
- Huang 2012b** {published data only}
Huang XT, Luo GL. [Clinical observation on combine traditional Chinese and western medicine treat polycystic ovary syndrome infertility]. *China Medical Engineering* 2012;**20**(1):65-6.
- Hung 2016** {published data only}
Hung YC, Kao CW, Lin CC, Liao YN, Wu BY, Hung IL, et al. Chinese herbal products for female infertility in Taiwan: a population-based cohort study. *Medicine* 2016;**95**(11):e3075.
- Huo 2008** {published data only}
Huo J, Yang SP, Xie BJ, Liao SG, Lin LP, Ding J, et al. Cytotoxic sesquiterpenoids from *Vernonia bockiana*. *Journal of Asian Natural Products Research* 2008;**10**(5-6):571-5.
- Jalilian 2013** {published data only}
Jalilian N, Modarresi M, Rezaie M, Ghaderi L, Bozorgmanesh M. Phytotherapeutic management of polycystic ovary syndrome: role of aerial parts of wood betony (*Stachys lavandulifolia*). *Phytotherapy Research: PTR* 2013;**27**(11):1708-13.
- Jia 2004** {published data only}
Jia YJ, Qv AP. Observation on the ultrasonogram of polycystic ovarian syndrome treated with traditional Chinese medicine, a report of 58 cases. *Shanxi Journal of TCM* 2004;**20**(6):42-3.
- Jia 2008** {published data only}
Jia Y, Lu XH. [Clinical research of integration of traditional Chinese medicine and western medicine in treating infertility caused by PCOS]. *Journal of Gansu College of Traditional Chinese Medicine* 2008;**25**(3):37-8.

Jia 2010 {published data only}

Jia Y, Jiang CL. ["bushen huoxue qinggan fa" clinical studies with clomiphene citrate treatment of polycystic ovary syndrome]. *Journal of Hubei University of Chinese Medicine* 2010;**12**(1):50-2.

Jia 2012a {published data only}

Jia CM. Clinical observation on treating 60 cases of spleen deficiency type PCOS infertility in TCM and moxibustion. *Clinical Journal of Chinese Medicine* 2012;**4**(16):47-8.

Jia 2012b {published data only}

Jia WH. [Chinese medicine sub-cycle therapy-based treatment of 42 cases of polycystic ovary syndrome]. *China Practical Medical* 2012;**7**(7):167-8.

Jian 2011 {published data only}

Jian QH. Clinical research on PCOS phlegmatic hygrois syndrome treatment with kidney-invigoration and sputum-elimination method and liver-clearing and sputum-elimination method. Guangzhou University of Chinese Medicine 2011.

Jiang 2007 {published data only}

Jiang WH, Xu XQ. Analysis of Yulin decoction combined with laparoscope on sterility caused by PCOS. *World Journal of Integrated Traditional and Western Medicine* 2007;**2**(7):408-9.

Jiang 2011a {published data only}

Jiang JH. Clinical study of nourishing kidney--yin method on polycystic ovarian syndrome of deficiency of kidney-yin. Nanjing University of Chinese Medicine 2011.

Jiang 2011b {published data only}

Jiang MF. Clinical research on polycystic ovarian syndrome's treatment with modified Cang Fu Dao Tan (CFDT) decoction. Guangzhou University of Chinese Medicine 2011.

Jiang 2014 {published data only}

Jiang DS, Wu XQ, Zhang YC. [Effects of warm needling combined with zhangmo decoction on endometrial receptivity in patient with ovulation induction]. *Zhongguo Zhen Jiu [Chinese Acupuncture & Moxibustion]* 2014;**34**(2):130-4.

Jiang 2015 {published data only}

Jiang D, Zhang Y, Wu X, Wu S. Infertility in polycystic ovary syndrome treated with acupuncture and clomiphene: a randomized controlled trial. *Zhongguo Zhen Jiu [Chinese Acupuncture & Moxibustion]* 2015;**35**(2):114-8.

Jin 2014a {published data only}

Jin XT, Ma K, Shan J. [Clinical efficacy observation on therapy ovulation failure infertility caused by PCOS with reinforcing kidney, activating blood circulation and ovarian stimulation compound recipe]. *Zhongguo Zhong Yao Za Zhi [China Journal of Chinese Materia Medica]* 2014;**39**(1):140-3.

Jin 2014b {published data only}

Jin CL, Wei LX, Zhao JP, Wu ZC. [Efficacy comparison between electroacupuncture and dyne-35 in treatment of polycystic ovary syndrome]. *Zhongguo Zhen Jiu [Chinese Acupuncture & Moxibustion]* 2014;**34**(12):1174-8.

Jin 2016 {published data only}

Jin JH. [Efficacy of integrative medicine obese women with polycystic ovary syndrome]. *Maternal & Child Health Care of China* 2016;**31**(1):187-8.

Johnson 2015 {published data only}

Johnson LK, Holven KB, Nordstrand N, Mellembakken JR, Tanbo T, Hjeltnes J. Fructose content of low calorie diets: effect on cardiometabolic risk factors in obese women with polycystic ovarian syndrome: a randomized controlled trial. *Endocrine Connections* 2015;**4**(3):144-54.

Kang 2012 {published data only}

Kang F, Kang Q, Zhang Q. [Combine traditional Chinese and western medicine in the treatment of polycystic ovary syndrome research]. *Chinese Community Doctors* 2012;**14**(32):169.

Kawakami 2011 {published data only}

Kawakami Z, Ikarashi Y, Kase Y. Isoliquiritigenin is a novel NMDA receptor antagonist in kampo medicine yokukansan. *Cellular and Molecular Neurobiology* 2011;**31**(8):1203-12.

Kitagawa 2015 {published data only}

Kitagawa H, Munekage M, Ichikawa K, Fukudome I, Munekage E, Takezaki Y, et al. Pharmacokinetics of active components of Yokukansan, a traditional Japanese herbal medicine after a single oral administration to healthy Japanese volunteers: a cross-over, randomized study. *PLoS One* 2015;**10**(7):e0131165.

Kort 2014 {published data only}

Kort DH, Lobo RA. Preliminary evidence that cinnamon improves menstrual cyclicity in women with polycystic ovary syndrome: a randomized controlled trial. *American Journal of Obstetrics and Gynecology* 2014;**211**(5):487.e1-6.

Kuang 2012 {published data only}

Kuang LJ, He B. Efficacy analysis of treating PCOS by clomiphene plus TCM artificial cycle. *Clinical Journal of Chinese Medicine* 2011;**4**(20):15-6.

Kuang 2013 {published data only}

Kuang H, Li Y, Wu X, Hou L, Wu T, Liu J, et al. Acupuncture and clomiphene citrate for live birth in polycystic ovary syndrome: study design of a randomized controlled trial. *Evidence-based Complementary and Alternative Medicine (eCAM)* 2013;**2013**:527303.

Kuang 2015 {published data only}

Kuang H, Jin S, Hansen KR, Diamond MP, Coutifaris C, Casson P, et al. Identification and replication of prediction models for ovulation, pregnancy and live birth in infertile women with polycystic ovary syndrome. *Human Reproduction* 2015;**30**(9):2222-33.

Kuek 2011 {published data only}

Kuek S, Wang WJ, Gui SQ. Efficacy of Chinese patent medicine Tian Gui Capsule in patients with polycystic ovary syndrome: a randomized controlled trial. *Zhong Xi Yi Jie He Xue Bao [Journal of Chinese Integrative Medicine]* 2011;**9**(9):965-72.

Lai 2006 {published data only}

Lai YQ, Chen H, Duan LJ. [Integration therapy of Chinese and western medicine in treating PCOS]. *Journal of Sichuan of Traditional Chinese Medicine* 2006;**24**(3):74-5.

Lai 2011 {published data only}

Lai MH, Ma HX, Liu H, Chen YL, Song XH, Ding T, et al. [Chinese medicine fumigation for ovulation induction treatment of polycystic ovary syndrome patients with endometrial development]. *Chinese Journal of Traditional Medical Science and Technology* 2011;**8**(4):335-6.

Lai 2014a {published data only}

Lai L, Flower A, Moore M, Lewith G. Chinese herbal medicine and polycystic ovary syndrome: a randomized feasibility and pilot study in the United Kingdom. *The Journal of Alternative and Complementary Medicine* 2014;**20**(5):A61-2.

Lai 2014b {published data only}

Lai L, Flower A, Moore M, Prescott P, Lewith G. Polycystic Ovary syndrome: a Randomised feasibility and pilot study using Chinese Herbal medicine to explore Impact on Dysfunction (ORCHID)—study protocol. *European Journal of Integrative Medicine* 2014;**6**(3):392-9.

Lai 2014c {published data only}

Lai L, Li X, Flower A, Moore M, Liu J. Chinese herbal medicine for oligomenorrhoea and amenorrhoea in polycystic ovary syndrome: a systematic review and meta-analysis. *Journal of Alternative & Complementary Medicine* 2014;**20**(5):A129.

Lai 2015a {published data only}

Lai L, Flower A, Moore M, Lewith G. Developing clinical practice guidelines for Chinese herbal treatment of polycystic ovary syndrome: A mixed-methods modified Delphi study. *Complementary Therapies in Medicine* 2015;**23**(3):430-8.

Lai 2015b {published data only}

Lai L, Flower A, Prescott P, Moore M, Lewith G. Chinese herbal medicine for oligomenorrhoea and amenorrhoea in polycystic ovary syndrome: A randomised feasibility study in the United Kingdom. *European Journal of Integrative Medicine* 2015;**7**(6):681-2.

Lai 2015c {published data only}

Lai L, Flower A, Prescott P, Moore M, Lewith G. Evaluating practitioner-blinding in Chinese herbal medicine research: Findings from a randomised feasibility study in the United Kingdom. *Integrative Medicine Research* 2015;**4**(1, Supplement):10.

Lai 2015d {published data only}

Lai L, Flower A, Prescott P, Moore M, Lewith G. Treatment adherence in Chinese herbal medicine: Findings from a randomised feasibility study in the United Kingdom. *Integrative Medicine Research* 2015;**4**(1, Supplement):126.

León-Gonzalez 2014 {published data only}

León-Gonzalez AJ, Acero N, Muñoz-Mingarro D, López-Lázaro M, Martín-Cordero C. Cytotoxic activity of hirsutanone, a diarylheptanoid isolated from *Alnus glutinosa* leaves.

Phytomedicine: International Journal Of Phytotherapy And Phytopharmacology 2014;**21**(6):866-70.

Li 2000 {published data only}

Li XB, Li LY. [Daotan Zhongzi formula combined with clomiphene in treating PCOS]. *Practical Medicine Journal* 2000;**16**(4):330-1.

Li 2002 {published data only}

Li XB, Li LY, Huang JL, Liang XF. Effect of operation under celioscopy combined with kidney tonifying and phlegm removing herbal medicine for polycystic ovarian disease syndrome. *Traditional Chinese Drug Research & Clinical Pharmacology* 2002;**13**(2):75-6, 131.

Li 2005 {published data only}

Li XP. Chinese drugs of reinforcing kidney, clearing liver, and activating blood for treatment of infertility caused by polycystic ovary syndrome. *Journal of Fujian College of TCM* 2005;**15**(4):9-11.

Li 2009a {published data only}

Li XW, Geng JF. [Clinical observation on Chinese herbal medicine combined with Western medicine in treating 95 cases of PCOS]. *Chinese Journal of Modern Drug Application* 2009;**3**(7):122-3.

Li 2009b {published data only}

Li L, Sun WF. [Clinical research on TCM Bushen huoxue method in treating endocrinological metabolism of adolescent PCOS]. *Practical Medicine Journal* 2009;**25**(13):2177-9.

Li 2009c {published data only}

Li L, Sun WF. Clinical research on kidney-tonifying and blood-activating therapy for the treatment of puberty polycystic ovarian syndrome. *Journal of Guangzhou University of TCM* 2009;**26**(3):221-4.

Li 2009d {published data only}

Li XL, Wu XQ. [Clinical research on Chinese herbal medicine combined with western medicine in treating 40 cases of PCOS]. *Practical Clinical Journal of Integrated Traditional Chinese and Western Medicine* 2009;**9**(2):40-2.

Li 2010a {published data only}

Li B, Han JR. [Clinical observation of combination of acupuncture with medicine treatment of polycystic ovary syndrome]. *Journal of Clinical Acupuncture and Moxibustion* 2010;**26**(1):20-1.

Li 2010b {published data only}

Li FY, Wang Q. [A warm nest soup treatment of 46 cases of kidney Yang deficiency type polycystic ovary syndrome curative effect]. *Journal of Shanxi College of Traditional Chinese Medicine* 2010;**11**(4):10-1.

Li 2010c {published data only}

Li SZ. ["bu shen tiao jing yu zi tang" combine with metformin treat 36 cases of infertility caused by polycystic ovary syndrome]. *Shanxi Journal of Traditional Chinese Medicine* 2010;**26**(6):26-7.

Li 2010d {published data only}

Li ZZ. The clinical study of using Cang Fu Dao Tan Wan in the treatment of PCOS of the phlegm stagnation syndrome. Guangzhou University of Chinese Medicine 2010.

Li 2011a {published data only}

Li HC, Ma YJ, Wang XQ. Combined treatment of traditional Chinese medicine and Western medicine for insulin resistant polycystic ovarian syndrome associated infertility. *Medical Journal of Qil* 2011;**26**(1):31-2.

Li 2011b {published data only}

Li XP, Lin S, Ye S. Therapeutic efficacy of modified zigui decoction in treatment of polycystic ovary syndrome of gan-shen yin deficiency syndrome. *Zhongguo Zhong Xi Yi Jie He Za Zhi [Chinese Journal of Integrated Traditional and Western Medicine]* 2011;**31**(8):1070.

Li 2011c {published data only}

Li XP, Ye S, Lin S, Zheng CS. Incites the last of the ten Heavenly Stems soup to add and subtract to the hepatorenal Yin deficient multi-pouch ovary syndrome high testosterone hormone blood's sickness influence research. *Guangming Journal of Chinese Medicine* 2011;**26**(2):242-4.

Li 2011d {published data only}

Li XB, Zhai JL, Li LY, HU XD. Influence of Lingzhu granules on endocrine and lipid metabolism of polycystic ovary syndrome patients. *Journal of Traditional Chinese Medicine* 2011;**52**(1):31-4.

Li 2011e {published data only}

Li C, Zhang SB. [Acupuncture treatment clinical observation of 30 cases of polycystic ovary syndrome]. *Beijing Journal of Traditional Chinese Medicine* 2011;**30**(2):128-9.

Li 2011f {published data only}

Li FY. [Combine traditional Chinese and Western medicine treatment of kidney Yang deficiency type PCOS clinical studies]. Guiyang College of Traditional Chinese Medicine 2011.

Li 2011g {published data only}

Li HX, Ma WM, Gao XA, Liu Y. [Chinese medicine PCOS patients in vitro fertilization - embryo transfer in assisted reproductive]. *Acta Chinese Medicine and Pharmacology* 2011;**39**(6):77-8.

Li 2011h {published data only}

Li SP. Clinical and empirical study of reproductive hormones and glucose and lipid metabolism with WuJiSan in the treatment of polycystic ovary syndrome with type of stagnation of phlegm-damp. Nanjing University of Chinese Medicine 2011.

Li 2011i {published data only}

Li XP, Lin S, Ye S, Cai YS, Chen CM, Zheng CS. Therapeutic efficacy of modified zigui decoction in treatment of polycystic ovary syndrome of gan-shen yin deficiency syndrome. *Zhongguo Zhong Xi Yi Jie He Za Zhi [Chinese Journal of Integrated Traditional and Western Medicine]* 2011;**31**(8):1070-3.

Li 2011j {published data only}

Li XH. [Clinical observation of Integrative Medicine treat 30 cases of polycystic ovary syndrome]. *Yunnan Journal of Traditional Chinese medicine and Material Medical* 2011;**32**(1):12-4.

Li 2011k {published data only}

Li YL, Lian F, Wang RX, Zhang N. Observation of Clomiphene citrate resistant polycystic ovary syndrome treated by combination of acupuncture, Chinese and Western medicine. *Hebei Journal of Traditional Chinese Medicine* 2011;**32**(7):244-6.

Li 2012b {published data only}

Li J. Combine traditional Chinese and Western medicine treatment of 62 cases of polycystic ovary syndrome combined insulin resistance. *Guiding Journal of Traditional Chinese Medicine and Pharmacy* 2012;**18**(7):60-1.

Li 2012c {published data only}

Li HX, Ma WM, Gao XA, Liu Y. Influence of kidney-nourishing and phlegm-resolving drugs on endocrine of polycystic ovarian syndrome. *Chinese Archives of Traditional Chinese Medicine* 2012;**30**(3):554-5.

Li 2013a {published data only}

Li N. [Efficacy and safety evaluation of acupuncture combined with auricular point sticking therapy in the treatment of polycystic ovary syndrome]. *Zhongguo Zhen Jiu [Chinese Acupuncture & Moxibustion]* 2013;**33**(11):961-4.

Li 2013b {published data only}

Li Y, Ma H, Zhang Y, Kuang H, Ng EH, Hou L, et al. Effect of berberine on insulin resistance in women with polycystic ovary syndrome: study protocol for a randomized multicenter controlled trial. *Trials* 2013;**14**:226.

Li 2015 {published data only}

Li QG. Clinical study on compound norethindrone tablet combined with metformin in the treatment of polycystic ovary syndrome. *Medical Recapitulate* 2015;**21**(2):322-4.

Li 2016 {published data only}

Li Q, Wang CY, Xu CH, Niu YH, Wang ZZ. [Clinical study of self yuan warm tone Yanxuesan treatment of polycystic ovary syndrome]. *World Latest Medical Information Digest (continuous electronic journals)* 2016;**16**(17):156-7.

Lian 2008 {published data only}

Lian F, Zhang YP, Liu YH, Ma FM, Sun ZG. [Influence of Bushen Tiaochong on PCOS serum, TNF-a and IL-6 in follicle fluid, quality of oocyte]. Colloquium of 8th National TCM Gynecological Conference. 2008.

Lian 2012 {published data only}

Lian F, Zhao S. [Effects of Shen invigorating and Chong-channel regulating method on anti-Müllerian hormone and oocyte quality in polycystic ovarian syndrome patients]. *Zhongguo Zhong Xi Yi Jie He Za Zhi [Chinese Journal Of Integrated Traditional And Western Medicine]* 2012;**32**(1):9-12.

Liang 2011 {published data only}

Liang RW, Wang XY. Combine traditional Chinese and Western medicine treat infertility with Insulin resistance in follicular developmental delays. *Medicinal Innovation of China* 2011;**8**(18):65-6.

Liao 2014 {published data only}

Liao Q. Clinical study of combination therapy with polycystic ovary syndrome. *Journal of Hei Longjiang Medicine* 2014, (12):1360-2.

Lim 2011 {published data only}

Lim CED, L JP. Traditional Chinese medicine for gynaecological diseases. *Journal of the Australian Traditional-Medicine Society* 2011;**17**(1):17-9.

Lin 2005 {published data only}

Lin Y. Combinative treatment of Chinese traditional and Western medicine in 48 patients with sterility due to polycystic ovarian syndrome. *Maternal and Child Health Care of China* 2005;**20**(13):1642-3.

Lin 2009a {published data only}

Lin HM, Wu T. [Clinical observation on TCM combined with Western medicine in treating PCOS with infertility]. *Journal of TCM University of Hunan* 2009;**29**(7):55-7.

Lin 2009b {published data only}

Lin Y, Zhen SQ, Liang JL, Kong S. Clinical study on treatment of polycystic ovary syndrome with nuangongzhongzi recipe. *Guangdong Medical Journal* 2009;**30**(4):635-7.

Lin 2011 {published data only}

Lin L, Chen J. Application of therapy of nourishing kidney essence in polycystic ovary syndrome patients with kidney deficiency after laparoscopic surgery. *Journal of New Chinese Medicine* 2011;**43**(9):63-5.

Lin 2013a {published data only}

Lin HM, He HZ, Ma PL. [Effect of insulin resistance in patients with polycystic ovary syndrome kidney phlegm on]. *Heilongjiang Journal of Traditional Chinese Medicine* 2013;**42**(3):50-2.

Lin 2013b {published data only}

Lin H, Wang P, Huang Q, Zhong MY, Yang Y. [Effects of kidney-tonifying and collateral-activating prescription on sex hormone and ovary in patients with polycystic ovary syndrome]. *Hunan Journal of Traditional Chinese Medicine* 2013;**29**(10):11-2, 18.

Lin 2013c {published data only}

Lin BQ, Zhang ZM, Lin ZX, Liu Y, Zhong HZ, Yang Cx, et al. Transvaginal color Doppler monitoring the ovulation value of warm acupuncture combined with Erchen soup treatment in patients with polycystic ovary syndrome. *International Medicine and Health Guidance News* 2013;**19**(4):452-4.

Liu 2007 {published data only}

Liu DM, Li SX. [Observation on efficacy of Wuji baifeng pill combined with Diane-35 in treating 40 cases of PCOS with

clomiphene resistance]. *Journal of New Chinese Medicine* 2007;**39**(6):34-5.

Liu 2008 {published data only}

Liu QC, Wu WH, Wu DY, Feng XW, Ma YH, Li JY, et al. Clinical observation on the treatment of childhood refractory idiopathic thrombocytopenic purpura with Dihuang Zhixue Capsule. *Chinese Journal Of Integrative Medicine* 2008;**14**(2):132-6.

Liu 2009 {published data only}

Liu X. [Observation on efficacy of herbs combined with clomiphene in treating PCOS]. *China Pharmacy Medicine* 2009;**4**(8):157-8.

Liu 2010a {published data only}

Liu YH, Li Y, Niu YH, Niu YL, Zhang WY. Study of kidney-reinforcing drugs combined with gonadotropin on infertile induced by polycystic ovary syndrome. *Modern Journal of Integrated Traditional Chinese and Western Medicine* 2010;**19**(1):13-4, 17.

Liu 2010b {published data only}

Liu XX, Li XY, Liu YM, Meng ZM. [Clinical observation of combine traditional Chinese and western medicine treat polycystic ovary syndrome with obesity]. *China Practical Medical* 2010;**5**(34):129-30.

Liu 2010c {published data only}

Liu RX, Shang HY, Liu YQ. ["tiao jing zhu yun fang" combine with metformin for polycystic ovary syndrome]. The Tenth National Gynecological Branch of China Association of Chinese Medicine Gynecology of TCM Academic Conference. 2010:192-4, 202.

Liu 2010d {published data only}

Liu GY, Tao LL, Xin J, Xie PP. The Impact of thread-embedding and Chinese medicine on hyperandrogenism of obese patients with polycystic ovary syndrome. *Liaoning Journal of Traditional Chinese Medicine* 2010;**37**(10):2027-8.

Liu 2011a {published data only}

Liu DP, Yan XJ. Integration of traditional Chinese and Western medicine in the treatment of 30 cases of infertility due to polycystic ovary syndrome. *Modern Traditional Chinese Medicine* 2011;**31**(4):14-6.

Liu 2011b {published data only}

Liu HL. [Integrative traditional and western medicine for 32 cases of polycystic ovary syndrome]. *Henan Traditional Chinese Medicine* 2011;**31**(12):1420-1.

Liu 2012a {published data only}

Liu XX. The clinical research on the treatment of invigorating the kidney and activating blood and resolving phlegm decoction to polycystic ovary syndrome and insulin resistance patients. Nanjing University Of Chinese Medicine 2012.

Liu 2012b {published data only}

Liu YQ. [LiuRunXia treatment of polycystic ovary syndrome, a professor at the academic thoughts and clinical experience in the research of infertility]. China Academy of Chinese Medical Sciences 2012.

Liu 2012c {published data only}

Liu YP, Wang QH, Huang SH. [TCM and Western medicine treatment of the phlegm polycystic ovary syndrome Infertility 38 cases of clinical observation]. *Guide of China Medicine* 2012;**10**(22):283-4.

Liu 2013 {published data only}

Liu Y, Mao LH. Effect of danzhi xiaoyao pill on ovulation induction of polycystic ovarian syndrome patients of pathogenic fire derived from stagnation of gan-qi. *Zhongguo Zhong Xi Yi Jie He Za Zhi [Chinese Journal of Integrated Traditional and Western Medicine]* 2013;**33**(9):1191-5.

Liu 2014a {published data only}

Liu Q. [Influence the phlegm scattered plot PCOS patients with lipid metabolism and reproductive hormones]. *Journal of Chinese Medicinal Materials* 2014;**37**(8):1502-4.

Liu 2014b {published data only}

Liu XL. [Clinical patients with polycystic ovary syndrome Integrative Medicine]. *Medical Information* 2014, (18):454.

Lu 2010 {published data only}

Lu J, Chen YX, Li FY, Chen Y. [Clomiphene joint mountain continued Fu Rong Dao Tan Tang treatment of polycystic ovary syndrome analysis of 48 cases]. *Maternal and Child Health Care of China* 2010;**25**(1):138-9.

Lu 2012 {published data only}

Lu LF. [Chinese cycle therapy combined with clomiphene citrate treatment of polycystic ovary syndrome infertility efficacy analysis]. *Health Must Read Magazine* 2012;**11**(9):91.

Luo 2010 {published data only}

Luo J, Guo R. Clinical observation on invigorating kidney to remove phlegm and stasis in treating PCOS-IR. *Hubei Journal of Traditional Chinese Medicine* 2010;**32**(6):24-5.

Luo 2014 {published data only}

Luo L, Chen J. [Polycystic ovary syndrome clinical observation type of treatment]. *Chinese and Foreign Medical Research* 2014, (29):5-7.

Lv 2007 {published data only}

Lv XP, Cong HF, Wang XB. [Observation on efficacy of acupuncture combined with medication in treating ovulation failure induced by PCOS]. *Journal of Clinical Acupuncture and Moxibustion* 2007;**23**(8):35-6.

Lv 2009 {published data only}

Lv LP, Chen GS. [Jiawei buzhong yiqi decoction combined with Diane-35 in treating 30 cases of PCOS]. *Jiangxi Journal of TCM* 2009;**40**(5):42-3.

Lv 2010 {published data only}

Lv JW. Clinical study on polycystic ovary syndrome patient's hyperplasia of mammary gland treated by method of tonifying-kidney and regulating-menstruation. Nanjing University of Chinese Medicine 2010.

Ma 2009b {published data only}

Ma HX, You ZL, Lai MH. [Clinical observation of integration of traditional Chinese medicine and Western medicine in treating infertility caused by obese PCOS]. *Guiding Journal of Traditional Chinese Medicine and Pharmacy* 2009;**15**(7):19-20.

Ma 2010 {published data only}

Ma MH, Wang XD. The application of method regulating kidney & clearing lung in the treatment of 26 cases with hyperandrogenism caused by PCOS. *Journal of Nanjing University of Traditional Chinese Medicine* 2010;**26**(4):311-2.

Madder 2013 {published data only}

Madder LS. Treating PCOS naturally. *HerbalGram (HERBALGRAM)* 2013;**Jun-May**(98):58-65.

Mao 2003 {published data only}

Mao YX. Clinical observation of 3 kinds different ovulation stimulating methods on polycystic ovary syndrome. *Heilongjiang Medical Journal* 2003;**27**(7):489-90.

Mao 2011a {published data only}

Mao XG, Fang SS, Mao AH. Observation research of 80 cases of polycystic ovary syndrome using combination of Chinese traditional and Western medicine. *Chinese Journal of Ethnomedicine and Ethnopharmacy* 2011;**20**(24):36-7.

Mao 2011b {published data only}

Mao XH. [Clinical observation of traditional Chinese combine with western medicine treatment of polycystic ovarian syndrome]. *Shanxi Journal of Traditional Chinese Medicine* 2011;**27**(7):27-8.

Mei 2010 {published data only}

Mei JL. Clinical study on the effect of kidney-nourishing and phlegm-resolving method combined with Diane-35 treating kidney-deficiency and phlegm-dampness type of polycystic ovary syndrome accompanied hyperandrogenism. Nanjing University of Traditional Chinese Medicine 2010.

Meng 2011 {published data only}

Meng J. Clinical and empirical study on therapeutic effects of Ling-Zhu granule in polycystic ovary syndrome and insulin resistance. Guangzhou University of Chinese Medicine 2011.

Miao 2012 {published data only}

Miao Y. [Clinical analysis of combine traditional Chinese and western medicine treatment of polycystic ovarian syndrome]. *Guangming Journal of Chinese Medicine* 2012;**27**(8):1619-20.

Ming-Wei 2011 {published data only}

Ming-wei X, Xin-yun L, Jun-qin H. Regulation and control of wenshen yangxue granule combined with clomifene citrate on INH-ACT-FS system in patients with follicular maldevelopment infertility. *Zhongguo Zhong Xi Yi Jie He Za Zhi [Chinese Journal of Integrated Traditional and Western Medicine]* 2011;**31**(12):1596-600.

Mohammad Hosseinzadeh 2016 {published data only}

Mohammad Hosseinzadeh F, Hosseinzadeh-Attar MJ, Yekaninejad MS, Rashidi B. Effects of selenium supplementation

- on glucose homeostasis and free androgen index in women with polycystic ovary syndrome: A randomized, double blinded, placebo controlled clinical trial. *Journal of Trace Elements in Medicine and Biology: organ of the Society for Minerals and Trace Elements (GMS)* 2016;**34**:56-61.
- Moradan 2012** {published data only}
Moradan S. Is there any role for acupuncture in infertility treatment?. *International Journal of Fertility & Sterility* 2012;**6**:114.
- Mosalanejad 2015** {published data only}
Mosalanejad L, Shakeri F, Ghavi F. The effect of asafoetida on hormone levels of students with polycystic ovary syndrome. *Avicenna Journal of Phytomedicine* 2015;**5**:106-7.
- Motoo 2014** {published data only}
Motoo Y, Arai I, Tsutani K. Use of Kampo diagnosis in randomized controlled trials of Kampo products in Japan: a systematic review. *PLoS One* 2014;**9**(8):e104422.
- Musumeci 2006** {published data only}
Musumeci ML, Schlecht K, West DP, West LE, Innocenzi D, Micali G. The spectrum of systemic therapy and procedural approaches in the management of moderate to severe acne vulgaris: A review. *Giornale Italiano di Dermatologia e Venereologia* 2006;**141**(4):379-91.
- O'Brien 2010** {published data only}
O'Brien PE. Bariatric surgery: mechanisms, indications and outcomes. *Journal of Gastroenterology and Hepatology* 2010;**25**(8):1358-65.
- Pan 2010** {published data only}
Pan H. The investigation simplify the assayed of the insulin resistance of polycystic ovary syndrome and interfering in clinical treatment of Chinese medicinal materials. Nanjing University Of Chinese Medicine 2010.
- Pan 2012** {published data only}
Pan LZ, Wang Y, He S. Clinical efficacy on 101 cases of intractable infertility of polycystic ovary syndrome treated with integrative Chinese and Western medicine. *World Journal of Integrated Traditional and Western Medicine* 2012;**7**(1):40-2, 65.
- Pastore 2011** {published data only}
Pastore LM, Williams CD, Jenkins J, Patrie JT. True and sham acupuncture produced similar frequency of ovulation and improved LH to FSH ratios in women with polycystic ovary syndrome. *The Journal Of Clinical Endocrinology And Metabolism* 2011;**96**(10):3143-50.
- Pazyar 2012** {published data only}
Pazyar N, Yaghoobi R. A rationale of ginseng as a novel addition to the antihirsutism armamentarium. *Journal Of Alternative And Complementary Medicine (New York, NY)* 2012;**18**(3):210-1.
- Pei 2012** {published data only}
Pei YX, Li JC, Ma YB, Hu LH. [Clinical observation of traditional Chinese and western medicine for polycystic ovary syndrome]. *Hebei Journal of Traditional Chinese Medicine* 2012;**34**(5):703-4.
- Peng 2012** {published data only}
Peng YT. Clinical observation on Gui-zhi-fu-ling-wan with catgut embedding therapy to treat blood stagnation type of PCOS. Guangzhou University of Chinese Medicine 2012.
- Qiao 2012** {published data only}
Qiao SX. The clinical research on Chinese medicine sequential with the acupuncture treat anovulation infertility caused by Pcos. Nanjing University of Chinese Medicine 2012.
- Qiu 2006** {published data only}
Qiu HN, Xu J. [Clinical observation on acupuncture in treating 46 cases of obese PCOS patients]. *Chinese Community Doctors* 2006;**8**(14):86-7.
- Qu 2015** {published data only}
Qu J, Che Y, Xu P, Xia Y, Wu X, Wang Y. The higher response of vascular endothelial growth factor and angiotensin-II to human chorionic gonadotropin in women with polycystic ovary syndrome. *International Journal of Fertility & Sterility* 2015;**8**(4):373-8.
- Qv 2011** {published data only}
Qv HY, Li MZ, Chen HS. [Clinical observation of combining traditional Chinese and Western medicine treatment of polycystic ovarian syndrome]. *Chinese Journal of Aesthetic Medicine* 2011;**20**(6):390.
- Ran 2007a** {published data only}
Ran MX, Zhang H, Qi XJ, Nie XX, Chen Q. An experimental research on endometrial receptivity is affected by bushenhuoxue Chinese herbs. *Chinese Journal of Birth Health and Heredity* 2007;**15**(4):56-7, 78.
- Ran 2007b** {published data only}
Ran XM, Zhang H, Tai GX, Song ZH, Chen Q. [The influence of Yulin formula on CTmRNA expression of endometrium after ovulation inducing]. *Traditional Chinese Medicinal Research* 2007;**20**(2):29-31.
- Ran 2008** {published data only}
Ran MX, Song ZH, Zhang H, Tai GX, Chen Q. An clinical observation and experimental research on the ovulation induction therapy of western medicines combined with Chinese herbs for the anovulatory patients with PCOS. *Guangming Journal of Chinese Medicine* 2008;**23**(2):175-6.
- Rao 2012** {published data only}
Rao XH, Zhang Y. A clinical analysis of treating the anovulation type of infertility in the integrative medicine. *Clinical Journal of Chinese Medicine* 2012, (22):71-2.
- Rashidi 2013** {published data only}
Rashidi BH, Tehrani ES, Hamedani NA, Pirzadeh L. Effects of acupuncture on the outcome of in vitro fertilisation and intracytoplasmic sperm injection in women with polycystic ovarian syndrome. *Acupuncture in Medicine: Journal of the British Medical Acupuncture Society* 2013;**31**(2):151-6.

Ren 2002a {published data only}

Ren XQ. [Comparison of effect of two ovulation inducing methods in treating refractory PCOS with infertility]. *Guangxi Medical Journal* 2002;**24**(8):1165-7.

Ren 2002b {published data only}

Ren XQ. [Comparison of effect of two ovulation inducing methods in treating refractory PCOS with infertility]. *Journal of Practical Obstetrics and Gynecology* 2002;**18**(6):351-2.

Ren 2006 {published data only}

Ren QL, Tan Y. [Influence of Ziyin buyang method on serum IGF-1 and sexual hormone in PCOS patients]. *Jiangsu Journal of Traditional Chinese* 2006;**27**(5):28-9.

Ren 2008 {published data only}

Ren QL, Tan Y, Sun LZ. [Clinical research of Chinese Ziyin buyang formula sequentially treating 31 cases with PCOS]. *Jiangsu Journal of Traditional Chinese* 2008;**40**(3):40-2.

Ren 2011 {published data only}

Ren WK. Exploration of follicle stimulating prescription and luteotrophic prescription in the co-treatment of polycystic ovarian syndrome with kidney deficiency and blood stasis. Shandong University of Traditional Chinese Medicine 2011.

Ren 2013 {published data only}

Ren MY, Zhu MC. Integrative medicine 150 cases of infertility salpingitic. *Guangming Journal of Chinese Medicine* 2013;**28**(11):2266-7.

Ren 2014 {published data only}

Ren LN, Guo LH, Ma WZ, Zhang R. [A meta-analysis on acupuncture treatment of polycystic ovary syndrome]. *Zhen Ci Yan Jiu [Acupuncture Research]* 2014;**39**(3):238-46.

Ried 2015 {published data only}

Ried K. Chinese herbal medicine for female infertility: an updated meta-analysis. *Complementary Therapies in Medicine* 2015;**23**(1):116-28.

Sadrefozalayi 2014 {published data only}

Sadrefozalayi S, Farokhi F. Effect of the aqueous extract of *Foeniculum vulgare* (fennel) on the kidney in experimental PCOS female rats. *Avicenna Journal of Phytomedicine* 2014;**4**(2):110-7.

Salah 2013 {published data only}

Salah IM. Office microlaparoscopic ovarian drilling (OMLOD) versus conventional laparoscopic ovarian drilling (LOD) for women with polycystic ovary syndrome. *Archives Of Gynecology and Obstetrics* 2013;**287**(2):361-7.

See 2011 {published data only}

See CJ, McCulloch M, Smikle C, Gao J. Chinese herbal medicine and clomiphene citrate for anovulation: a meta-analysis of randomized controlled trials. *The Journal of Alternative and Complementary Medicine* 2011;**17**(5):397-405.

Shah 2016 {published data only}

Shah KN, Patel SS. Phosphatidylinositol 3-kinase inhibition: A new potential target for the treatment of polycystic ovarian syndrome. *Pharmaceutical Biology* 2016;**54**(6):975-83.

Shao 2004 {published data only}

Shao RY, Lang FJ, Cai JF. Clinical observation on treatment of Stein-Leventhal syndrome caused sterility by combined use of clomiphene and Chinese nourishing shen and activating blood circulation drugs. *Zhongguo Zhong Xi Yi Jie He Za Zhi [Chinese Journal of Integrated Traditional and Western Medicine]* 2004;**24**(1):41-3.

Shao 2006 {published data only}

Shao MX, Zhang XL, Sun JJ, Yan YF. [Clinical observation of integration of traditional Chinese medicine and Western medicine in treating infertility caused by PCOS]. *Journal of Sichuan of Traditional Chinese Medicine* 2006;**24**(1):90-1.

Shen 2008 {published data only}

Shen GH. [Quyue tiaojing decoction in treating 150 cases of PCOS]. *Journal of New Chinese Medicine* 2008;**40**(2):82-3.

Shen 2013 {published data only}

Shen W, Zhang Y, Li W, Cong J, Zhou Y, Ng EHY, et al. Effects of tanshinone on hyperandrogenism and the quality of life in women with polycystic ovary syndrome: protocol of a double-blind, placebo-controlled, randomised trial. *BMJ Open* 2013;**3**(10):e003646.

Sheng 2010 {published data only}

Sheng WZ. Clinical study on cycle treatment of polycystic ovary syndrome with Bu Shen Huo Xue method. Shandong University of Traditional Chinese Medicine 2010.

Shi 2009a {published data only}

Shi Y, Xu WF, Ying XJ, Zhu D. Clinical study on acupuncture combined with kidney yin tonifying herbs for polycystic ovary syndrome of kidney in deficiency. *Shanghai Journal of TCM* 2009;**43**(10):33-5.

Shi 2009b {published data only}

Shi Y, Feng HJ, Liu HR, Zhu D. Observation on therapeutic effect of acupuncture combined with Chinese herbs on polycystic ovary syndrome of kidney deficiency and phlegm stasis type. *Zhongguo Zhen Jiu [Chinese Acupuncture & Moxibustion]* 2009;**29**(2):99-102.

Shi 2010a {published data only}

Shi LJ. Observation of treatment of kidney weakness and phlegm blocking polycystic ovary syndrome based on method of tonify the kidney and regulating the period of menses with moxibustion. Nanjing University of Chinese Medicine 2010.

Shi 2010b {published data only}

Shi F, Chen YF. [Clinical observation on the combine traditional Chinese and western medicine treatment of 60 patients with polycystic ovarian syndrome]. *Journal of Changchun University of Traditional Chinese Medicine* 2010;**26**(6):922-3.

Shi 2011 {published data only}

Shi Y. [Xiao Chai Hu Jia JianTang Treating polycystic ovary syndrome in 40 patients]. *Zhejiang Journal of Traditional Chinese Medicine* 2011;**46**(2):120.

Shu 2012 {published data only}

Shu RM. Clinical study on treating polycystic ovary syndrome of liver depression with adjustment of Jia Jian Xiao Yao San and chloramphenicol. Guangzhou University of Chinese Medicine 2012.

Si 2016 {published data only}

Si Q. [Integrative therapy of menstrual disorders caused by polycystic ovary syndrome]. *Practical Electronics Magazine Gynecological Endocrinology* 2016;**3**(2):113-5.

Song 2010 {published data only}

Song ZL, Xu J, Wang Y. Clinical observation on "Bushen Ruanjian Formula" in treating polycystic ovary syndrome with anovulation. *Shanghai Journal of Traditional Chinese Medicine* 2010;**44**(8):47-50.

Song 2011 {published data only}

Song ZH, Zhang HM, Zhang XS. [Clinical observation of combining traditional Chinese and Western medicine treatment of polycystic ovary syndrome]. *Chinese Journal of Rehabilitation* 2011;**26**(4):286-7.

Stone 2009 {published data only}

Stone JAM, Yoder KK, Case EA. Delivery of a full-term pregnancy after TCM treatment in a previously infertile patient diagnosed with polycystic ovary syndrome. *Alternative Therapies In Health And Medicine* 2009;**15**(1):50-2.

Su 2012 {published data only}

Su SP, Liu YW, Peng M, Gao BZ. [Clinical observation on metformin combine with Bu Shen Huo Xue Hua Tan Methods for polycystic ovary syndrome]. *China Practical Medicine* 2012;**17**(17):159-60.

Sui 2011 {published data only}

Sui MF. Treatment of 96 patients with polycystic ovary syndrome of two methods' comparison. *Value Engineering* 2011;**30**(7):192.

Sun 2009a {published data only}

Sun J. [Integrated therapy of Chinese and western medicine on 56 patients with infertility caused by PCOS]. *Chinese Journal of Practical Medicine* 2009;**36**(3):72.

Sun 2009b {published data only}

Sun Y. Combination of TCM and WM (Western medicine) treat sterility of polycystic ovary syndrome. *Journal of Zhejiang College of Traditional Chinese Medicine* 2009;**33**(3):344-5.

Sun 2010a {published data only}

Sun C. Clinical study on the effect of kidney-nourishing and phlegm-resolving method combined with metformin treating polycystic ovary syndrome with insulin resistance. Nanjing University Of Chinese Medicine 2010.

Sun 2010b {published data only}

Sun W, Cui W, Li J. Study of the mechanism of electro-acupuncture treatment on PCOS and asthenia of kidney. *Chinese Journal of Birth Health and Heredity* 2010;**18**(1):105-7.

Sun 2011 {published data only}

Sun WF, Liang J, Zhang XX, Feng SM. Clinical research on kidney-tonifying blood-activating and sputum-eliminating therapy for the treatment of polycystic ovary syndrome (PCOS) with insulin resistance (IR). *Chinese Archives of Traditional Chinese Medicine* 2011;**29**(9):2018-20.

Sun 2012 {published data only}

Sun H, Peng WJ. [Clinical observation on combine traditional Chinese and western medicine treat 51 cases of polycystic ovary syndrome with obesity]. *Xinjiang Journal of Traditional Chinese Medicine* 2012;**30**(3):47-9.

Sun 2014 {published data only}

Sun HL. Clinical study of acupuncture odd ovulation disorders through four veins treatment of polycystic ovary syndrome. Yunnan Chinese Medicine College 2014.

Tan 2005 {published data only}

Tan MH. Resistant clomiphene sterility with polycystic ovarian syndrome treated by prescription of Chinese traditional medicine benefiting to gravidity and clomiphene. *Maternal and Child Health Care of China* 2005;**20**(7):824-5.

Tan 2012 {published data only}

Tan L, Tong Y, Sze SCW, Xu M, Shi Y, Song XY, et al. Chinese herbal medicine for infertility with anovulation: a systematic review. *Journal of Alternative & Complementary Medicine* 2012;**18**(12):1087-100.

Tang 2012 {published data only}

Tang CY, Wu YH. Tonifying kidney and removing blood stasis combined with Western medicine treatment for ovulation induction in patients with polycystic ovary syndrome. *Journal of Traditional Chinese Medicine University of Hunan* 2012;**32**(6):46,73.

Tao 2003 {published data only}

Tao LL, Chen XP, Gu ZT. Study on treatment of polycystic ovarian syndrome with infertility by combined therapy of Chinese herbal medicine and compound cyproterone acetate. *Chinese Journal of Integrative Medicine* 2003;**9**(2):98-103.

Tao 2006 {published data only}

Tao LL, Zhang YZ, Sang X, Zeng L, Yu AQ, Chen LF. Effects of modified longdan xiegan decoction on hyperandrogenism in patients with polycystic ovary syndrome of stagnant fire in Gan channel type. *Zhongguo Zhong Xi Yi Jie He Za Zhi [Chinese Journal of Integrated Traditional and Western Medicine]* 2006;**26**(9):838-41.

Tao 2008 {published data only}

Tao LL, Long YL, Sang X, Zhang YZ, Chen XP, Niu XY, et al. Effects of point catgut-embedding therapy combined Chinese medicinal herbs of invigorating spleen to remove phlegm on insulin resistance and serum adiponectin level in obese patients

with polycystic ovary syndrome. *China Journal of Traditional Chinese Medicine and Pharmacy* 2008;**23**(5):434-7.

Tao 2009 {published data only}

Tao LL, Fu YH, Xie PP, Wu T, Sang X, Long YL, et al. Effect of acupoint catgut-embedding therapy combined with spleen strengthening and phlegm removing herbs on insulin resistance and adipocyte factors in patients with obese-type polycystic ovary syndrome. *Journal of Guangzhou University of TCM* 2009;**26**(2):134-7.

Tao 2010 {published data only}

Tao LL, Wang HY, Chen XP, Sang X, Zeng C, Xin J, et al. Effect of point thread-embedding plus herbal decoction for strengthening the spleen to dissolve phlegm on the glucolipide metabolism of the obese patients with polycystic ovary syndrome. *Journal of Traditional Chinese Medicine* 2010;**51**(3):239-42.

Tao 2011 {published data only}

Tao JF, Lu XH. [Chinese medicine cycle therapy for female college students of 30 cases of polycystic ovary syndrome clinical observation]. *Journal of New Chinese Medicine* 2011;**43**(7):75-7.

Ulbricht 2016 {published data only}

Ulbricht CE, Natural Standard Research Collaboration. An evidence-based systematic review of Yin Yang Huo (*Epimedium* spp.) by the Natural Standard Research Collaboration. *Journal Of Dietary Supplements* 2016;**13**(2):136-64.

Ushiroyama 2001 {published data only}

Ushiroyama T, Ikeda A, Sakai M, Hosotani T, Suxuki Y, Tsubokura S, et al. Effects of unkei-to, an herbal medicine, on endocrine function and ovulation in women with high basal levels of luteinizing hormone secretion. *The Journal of Reproductive Medicine* 2001;**46**(5):451-6.

Ushiroyama 2006 {published data only}

Ushiroyama T, Hosotani T, Mori K, Yamashita Y, Ikeda A, Ueki M. Effect of switching to Wen-jing-tang (Unkei-To) from preceding herbal preparations selected by eight-principle pattern identification on endocrinological status and ovulatory induction in women with polycystic ovary syndrome. *The American Journal of Chinese Medicine* 2006;**34**(2):177-87.

Vajda 2013 {published data only}

Vajda FJE, Dodd S, Horgan D. Lamotrigine in epilepsy, pregnancy and psychiatry--a drug for all seasons?. *Journal of Clinical Neuroscience: Official Journal of the Neurosurgical Society of Australasia* 2013;**20**(1):13-6.

van Oppen 2015 {published data only}

van Oppen JD, Daniel PS, Sovani MP. What is the potential role of transcutaneous carbon dioxide in guiding acute noninvasive ventilation?. *Respiratory Care* 2015;**60**(4):484-91.

Wan 2012 {published data only}

Wan PF. The clinical study Intherapies of obesity-PCOS by Chinese herbs combined with different frequency of

electroacupuncture. Shandong Traditional Chinese Medicine University 2012.

Wang 2005a {published data only}

Wang YL. [Clinical observation on Bushen Huoxue Huatan formula in treating PCOS with infertility]. *Chinese Archives of Traditional Chinese Medicine* 2005;**23**(8):1519-20.

Wang 2005b {published data only}

Wang YH, Yang YS, Zhang YL. Clinical study of Ganshao capsule in treating clomiphene resistant polycystic ovarian syndrome. *Zhongguo Zhong Xi Yi Jie He Za Zhi [Chinese Journal of Integrated Traditional and Western Medicine]* 2005;**25**(8):704-6.

Wang 2006a {published data only}

Wang J, Zhang DX, Cao JM. [Clinical observation on Yishen jianpi method combined with metformin in treating PCOS with insulin resistance]. *Journal of Sichuan of Traditional Chinese Medicine* 2006;**24**(9):67-8.

Wang 2006b {published data only}

Wang J, Chen YF, Xu SQ. [Observation of integration of traditional Chinese medicine and western medicine in treating infertility caused by PCOS]. *Public Medical Forum Magazine* 2006;**10**(8B):717-8.

Wang 2009 {published data only}

Wang JL, Zhang ZC. Observation of acupuncture and acupoint catgut-embedding therapy for PCOS patients of obesity. *Liaoning Journal of Traditional Chinese Medicine* 2009;**36**(9):1574-5.

Wang 2010a {published data only}

Wang YH. [Combine traditional Chinese and western medicine treatment of infertility caused by polycystic ovary syndrome]. *China Modern Medicine* 2010;**17**(20):93-4.

Wang 2010b {published data only}

Wang Q, Li FY. Clinical research of integrated Chinese and western medicine in treating polycystic ovarian syndrome of kidney yang deficiency. *Shanghai Journal of Traditional Chinese Medicine* 2010;**44**(12):66-8.

Wang 2011a {published data only}

Wang NS. GFW modified act on polycystic ovary syndrome in experimental and clinical efficacy of phlegm and blood stasis type. Guangzhou University of Chinese Medicine 2011.

Wang 2011b {published data only}

Wang Q. [Chinese and Western medicine treatment of 60 cases of polycystic ovary infertility]. *Beifang Yaoxue* 2011;**8**(9):51.

Wang 2011c {published data only}

Wang Y. Clinical observation on 45 cases of polycystic ovarian syndrome accompanied with emotional disturbance treated by Shugan Tiaojing formula. *Journal of Traditional Chinese Medicine* 2012;**52**(17):1479-82.

Wang 2011d {published data only}

Wang Y, Chen Y, Wang X. The effect of Cang-Fu-Dao-Tan decoction on insulin resistance, adiponectin and leptin levels in

patients with obese-type polycystic ovarian syndrome. *Chinese Archives of Traditional Chinese Medicine* 2011;**29**(11):2256-7.

Wang 2012a {published data only}

Wang Q, Li FY. [Nuan Gong Zhu Yun Tang combine with clomiphene citrate for polycystic ovary syndrome (the kidney yang deficiency type) follicular development]. *Liaoning Journal of Traditional Chinese Medicine* 2012;**39**(9):1752-4.

Wang 2012b {published data only}

Wang QH, Liu YP, Xie PP, Yang XP, Huang SH, Chen SW, et al. Clinical research of polycystic ovary syndrome treated with non-surgical integrated Chinese and Western medicine. *World Journal of Integrated Traditional and Western Medicine* 2012;**7**(11):968-71.

Wang 2013 {published data only}

Wang YR, Li CR. [Clinical observation on acupuncture treatment of polycystic ovary syndrome]. *China Healthcare & Nutrition* 2013;**23**(4):2112.

Wang 2016 {published data only}

Wang LL, Yang HW, Wu XK. [Acupuncture treatment of polycystic ovary syndrome RCTs syndrome and myofascial trigger point acupoints]. *Journal of Chinese Medicine* 2016;**66**(8):673-6.

Wei 2008 {published data only}

Wei X. Clinical observation of integration of traditional Chinese medicine and western medicine in treating infertility caused by PCOS. *Hebei Journal of TCM* 2008;**30**(7):732-4.

Wei 2011a {published data only}

Wei CL. [Chinese herbal compound joint Diane-35 on sex hormone levels in patients with polycystic ovary syndrome]. *Chinese Journal of Experimental Traditional Medical Formula* 2011;**17**(5):239.

Wei 2011b {published data only}

Wei XX. [Combine traditional Chinese and western medicine in the treatment of polycystic ovary syndrome curative effect observation]. *Guide of China Medicine* 2011;**9**(29):152-3.

Wei 2011c {published data only}

Wei YQ. [Clinical observation on combine traditional Chinese and Western medicine for polycystic ovarian syndrome]. *Seek Medical and Ask the Medicine* 2011;**9**(11):589.

Wu 2008 {published data only}

Wu N. Clinical study on Bushen Huayu Qutan treatment for PCOS. *The Chinese Journal of Modern Applied Pharmacy* 2008;**25**(2):167-70.

Wu 2010a {published data only}

Wu MY. [Clinical study of combining traditional Chinese and western medicine treatment of polycystic ovary syndrome]. *Medical Innovation of China* 2010;**7**(34):47-8.

Wu 2010b {published data only}

Wu XY, Liang GY, Yao L. Clinical observation on polycystic ovarian syndrome (PCOS) infertility treatment with Chinese

herbs combined with clomiphene. *Journal of Hubei University of Chinese Medicine* 2010;**12**(2):15-7.

Wu 2011 {published data only}

Wu KE. Random controlled prospective study of the treatment for PCOS by acupuncture and Chinese medicine combined. Guangzhou University of Chinese Medicine 2011.

Wu 2012a {published data only}

Wu D. Kidney and spleen frange with the clinical research of acupuncture on patients with polycystic ovary syndrome kaohsiung signs intervention. Heilongjiang University of Chinese Medicine 2012.

Wu 2012b {published data only}

Wu CC, Lei P, Ruan YM, Lin XM, Xiong YL, Yang GY. [Effects of oral contraceptive pretreatment on controlled ovarian hyperstimulation and outcomes of IVF-ET]. *Zhonghua Nan Ke Xue [National Journal Of Andrology]* 2012;**18**(7):623-6.

Wu 2013a {published data only}

Wu YY. Efficacy catgut embedding treatment of polycystic ovary syndrome. *Medical Information* 2013;**66**(20):422.

Wu 2013b {published data only}

Wu Y, Leonard C, Denz E, Haeberle M. The treatment of luteinised unruptured follicle syndrome with Chinese medicine. *Journal of Chinese Medicine* 2013;**Jun**(102):5-19.

Wuttke 2015 {published data only}

Wuttke W, Seidlova-Wuttke D. Herbal medicines for menopausal symptoms. *Maturitas Conference: 10th European Congress on Menopause and Andropause; Madrid, Spain* 2015;**81**(1):120.

Xia 2004 {published data only}

Xia XJ, Guo CY. [Clinical observation of integrated therapy of Chinese and western medicine on 86 patients with infertility caused by PCOS]. *The Chinese Journal of Human Sexuality* 2004;**13**(7):16-7, 19.

Xia 2007 {published data only}

Xia XJ. Treatment on anovulatory infertility induced by polycystic ovary syndrome through supplementing kidney and promoting blood circulation. *The Chinese Journal of Human Sexuality* 2007;**16**(10):23-6, 30.

Xia 2011 {published data only}

Xia FL, Si YR, Wen YN. [Clinical research on Wen Shen Jian Pi Xiao Tan Zhu Yun particles combined clomiphene treatment of polycystic ovary infertility]. *Shanxi Journal of Traditional Chinese Medicine* 2011;**32**(3):265-7.

Xiao 2014 {published data only}

Xiao SF, Xu YY, Zhu LJ, Hu JL. [Catgut embedding joint Diane and metformin in obese women with polycystic ovary syndrome in 36 cases]. *Jiangxi Journal of Traditional Chinese Medicine* 2014, (4):53-5.

Xie 2005 {published data only}

Xie GZ, Zhou ZX, Sun QH, Yang MC, Chen WY, Yang J, et al. Tiao jing gu chong tang and clomiphene treats the PCOS with the

result that clinical observation of the sterility. *Chinese Journal of the Practical Chinese with Modern Medicine* 2005;**18**(10):1538-9.

Xie 2010 {published data only}

Xie XY. [Clinical observation of combining traditional Chinese and western medicine treatment of polycystic ovary syndrome]. *Guide of China Medicine* 2010;**8**(34):103-4.

Xie 2012 {published data only}

Xie JH, Gao YM, Han R, Zhang Y. Yin-nourishing and heat-clearing therapy for 30 cases of polycystic ovary syndrome. *World Chinese Medicine* 2012;**7**(5):410-2.

Xiong 2012 {published data only}

Xiong XL. Treatment effect observation 115 cases patients with polycystic ovary syndrome. *Chinese Journal of Clinical Rational Drug Use* 2012;**5**(13):59-60.

Xu 2008a {published data only}

Xu JH. 30 cases of polycystic ovary syndrome treated in Chinese medicine combined with Diane-35. *Journal of Henan University of Chinese Medicine* 2008;**23**(2):46-7.

Xu 2008b {published data only}

Xu HO, Luo YL. [The application of Zishen yutai pill in ovulation inducing of PCOS]. *Chinese Journal of Information on TCM* 2008;**15**(5):68.

Xu 2009 {published data only}

Xu F. [The efficacy of Guishen pill in treating 40 cases of PCOS]. *Jiangxi Journal of TCM* 2009;**40**(7):36-7.

Xu 2010a {published data only}

Xu DW. [Observation on clinical efficacy of Chinese and Western medicine combined with acupuncture in treating PCOS with infertility]. *Journal of Sichuan of Traditional Chinese Medicine* 2010;**28**(1):93-4.

Xu 2010b {published data only}

Xu SQ. Effects of Bushenshugan method on hyperandrogenism and hyperinsulinemia in polycystic ovary syndrome. Hubei University of Chinese Medicine 2010.

Xu 2012 {published data only}

Xu LF. The clinical research of two Chentang traditional Chinese medicine combined cycle therapy treatment for obese type polycystic ovarian syndrome. Guangzhou University of Chinese Medicine 2012.

Xu 2016 {published data only}

Xu BH, Li MQ, Chen CL, Luo YJ. [Effects of TCM periodic therapy on sex hormone, glucose and lipid metabolism of patients with polycystic ovary syndrome]. *Chinese Journal of Information on Traditional Chinese Medicine* 2016, (1):35-8.

Xue 2004 {published data only}

Xue XW, Wang N. [Observation on efficacy of TCM combined with ultrasound in treating 56 cases of PCOS]. *Chinese General Practice* 2004;**7**(11):828.

Yan 2003 {published data only}

Yan YF, Feng XS, Jiang MF. [Clinical observation on Bushen Huoxue herbal medicine in treating 32 cases of PCOS with infertility]. *Journal of Traditional Chinese Medicine* 2003;**44**(1):39-40.

Yan 2005 {published data only}

Yan YF, Yan L. Clinical observation on 30 cases of polycystic ovarian syndrome treated by Bushen Huoxue oral liquid plus metformin. *Journal of Traditional Chinese Medicine* 2005;**46**(9):480-2.

Yan 2012 {published data only}

Yan XJ. [Clinical observation on combine traditional Chinese and western medicine treat 30 cases of polycystic ovary syndrome with infertility]. *Guiding Journal of Traditional Chinese Medicine and Pharmacology* 2012;**18**(5):53-4.

Yang 2005 {published data only}

Yang YH, Hong JY, Wei DY, Chen XM, Lin J. Acupuncture on polycystic ovarian syndrome-induced infertility. *Journal of Guangdong Medical College* 2005;**23**(4):377-8.

Yang 2008 {published data only}

Yang L, Qi B, Song JX, Hou LH, Wu XK. Detection and analysis of apoptosis related genes of insulin resistance granular cells induced by the traditional Chinese medicine by DNA microarray. *Science & Technology Review* 2008;**26**(16):42-6.

Yang 2010a {published data only}

Yang GM. [Clinical observation of combining traditional Chinese and Western medicine treatment of polycystic ovary syndrome]. *Journal of Clinical and Experimental Medicine* 2010;**9**(17):1331-2.

Yang 2010b {published data only}

Yang P. [Curative effect observation of combining traditional Chinese and Western medicine treatment of polycystic ovary syndrome]. *Journal of Guiyang College of Traditional Chinese Medicine* 2010;**32**(5):43-4.

Yang 2010c {published data only}

Yang JB, Xu PL, Yao F, Yang XY. [Combine traditional Chinese and western medicine clinical observation for the treatment of ovulation disorder polycystic ovary syndrome]. Association of Chinese Medicine Gynecology Branch of the Tenth National Gynecologic Academic Conference. 2010.

Yang 2011 {published data only}

Yang ZC. [Clinical observation on Zi Ni Wen Shen Tiao Jing Tang treatment of polycystic ovary syndrome]. *Chinese Journal of Misdiagnostics* 2011;**11**(34):8377.

Yang 2014a {published data only}

Yang D. [Effect of acupuncture and clomiphene citrate for kidney deficiency PCOS infertile women ovulation rate and pregnancy rate]. Hunan University of Traditional Chinese Medicine.

Yang 2014b {published data only}

Yang H. [Traditional Chinese medicine for the treatment of kidney and blood stasis 35 cases of polycystic ovary syndrome]. *Global Chinese Medicine* 2014, (5):382-4.

Yang 2015 {published data only}

Yang B-Z, Cui W, Li J. Effects of electroacupuncture intervention on changes of quality of ovum and pregnancy outcome in patients with polycystic ovarian syndrome. *Zhen Ci Yan Jiu [Acupuncture Research]* 2015;**40**(2):151-6.

Yao 2011 {published data only}

Yao FW. Clinical Research on the method of kidney-nourishing and phlegm-resolving method combined with Diane-35 on the effects of polycystic ovary syndrome with the high testosterone. Heilongjiang University of Chinese Medicine 2011.

Yao 2012a {published data only}

Yao Y. Clinical study on polycystic ovary syndrome accompanied anovulation by invigorating kidney and activating blood circulation and regulating chong drink. Nanjing University of Chinese Medicine 2012.

Yao 2012b {published data only}

Yao XY, Feng YL, Tang S. [Combine traditional Chinese and Western medicine treatment the clinical effect of the treatment of infertility caused by polycystic ovary syndrome]. *Health Must Read Magazine* 2012;**11**(10):31-2.

Ye 2004 {published data only}

Ye LQ. [Chinese periodical therapy combined with metformin in treating 62 cases with PCOS]. *Jiangxi Journal of TCM* 2004;**35**(5):22-3.

Ye 2010 {published data only}

Ye TZ, Cao LQ, Ye CC, Zhang QR, Shen XL, Guo M, et al. [Combine traditional Chinese and Western medicine treatment of insulin resistance of 40 cases of polycystic ovary syndrome were observed]. *Zhejiang Journal of Traditional Chinese Medicine* 2010;**45**(4):287.

Ye 2012a {published data only}

Ye YY. [Combine traditional Chinese and western medicine treatment of polycystic ovary syndrome curative effect analysis]. *The Journal of Medical Theory and Practice* 2012;**25**(5):560-2.

Ye 2012b {published data only}

Ye HJ, Jiang TJ, Li AP, Yv Y. Clinical observation on treating polycystic ovary syndrome with sterility with Guizhifulin capsule and Diane-35 and clomiphene citrate. *Chinese Journal of Clinical Pharmacology and Therapeutics* 2012;**17**(6):691-5.

Ye 2015 {published data only}

Ye LQ. [TCM periodic therapy with metformin efficacy of 120 cases of pubertal polycystic ovary syndrome]. *Journal of Heilongjiang Traditional Chinese Medicine* 2015;**44**(6):17-8.

Yi 2012 {published data only}

Yi LL. Clinical research of gubenquyu decoction in treating PCOS of Qi deficiency and phlegm and blood stasis type. Hubei University of Chinese Medicine 2012.

Yin 2007 {published data only}

Yin XP. [Laparoscopy combined with TCM in treating PCOS]. *Reproduction and Contraception* 2007;**27**(7):484-5.

Yu 2013 {published data only}

Yu L, Liao Y, Wu H, Zhao J, Wu L, Shi Y, et al. Effects of electroacupuncture and Chinese kidney-nourishing medicine on polycystic ovary syndrome in obese patients. *Chung i Tsah Chih Ying Wen Pan [Journal of Traditional Chinese Medicine]* 2013;**33**(3):287-93.

Yu 2015 {published data only}

Yu LJ. Traditional Chinese Medicine (yi xu huo wang) versus western medicine to treat women with polycystic ovary syndrome. *Modern Journal of Integrated Traditional Chinese and Western Medicine* 2015, (11):1206-8.

Yuan 2011 {published data only}

Yuan Q. Clinical research of AMH during the process of stimulation period of PCOS as well as the kidney sand sequential Chinese medicine. Nanjing University of Chinese Medicine 2011.

Yv 2011 {published data only}

Yv W. Clinical research on combination Chinese medicine and acupuncture in treating polycystic ovarian syndrome. Heilongjiang University of Chinese Medicine 2011.

Zeng 2007 {published data only}

Zeng BF, Xu AF. [Clinical observation on pregnancy prompting decoction combined with clomiphene in treating infertility induced by PCOS]. *China Medical Herald* 2007;**4**(12):68-9.

Zeng 2012 {published data only}

Zeng L, Zeng C, Tao LL. Comparative study on Chinese medical syndrome typing and treatment combined different surgical methods for treating clomiphene-resistant polycystic ovary syndrome. *Zhongguo Zhong Xi Yi Jie He Za Zhi [Chinese Journal of Integrated Traditional and Western Medicine]* 2012;**32**(11):1492-5.

Zhang 2003a {published data only}

Zhang LM, Hou LH. [Clinical observation on Bushen decoction combined with acupuncture in treating PCOS]. *Journal of Clinical Acupuncture and Moxibustion* 2003;**19**(4):23-4.

Zhang 2003b {published data only}

Zhang MM, Huang GY, Lu FE, Paulus WE, Sterzik K. Treatment of polycystic ovary syndrome with clomiphene resistance patients by herbs combined with clomiphene. *Chinese Journal of the Practical Chinese with Modern Medicine* 2003;**3**(16):157-9.

Zhang 2007a {published data only}

Zhang FC. Effect of qingfei jianpi xiaocuo decoction for the treatment of acne in polycystic ovary syndrome: an observation of 46 cases. *Journal of New Chinese Medicine* 2007;**39**(9):34-5.

Zhang 2007b {published data only}

Zhang H. [Research on Chinese herbal medicine combined with Western medicine in treating 40 cases of PCOS]. *Journal of Changchun University of TCM* 2007;**23**(5):68-9.

Zhang 2007c {published data only}

Zhang Y, Tu RQ, Tu DJ. The clinical effects of Diane-35 with Chinese traditional medicine in polycystic ovary syndrome therapy. *Clinical Medical Journal of China* 2007;**14**(5):711-2.

Zhang 2007d {published data only}

Zhang H, Ran MX, Chen Q, Tai GX, Song ZH. An experimental research on the expression of endometrial calcitonin affected by Bushen huoxue Chinese herbs. *Chinese Journal of Family Planning* 2007;**15**(3):171-3.

Zhang 2009 {published data only}

Zhang QM, Sun ZY, Chen X, Yuan JL. Clinical effect observation on treatment to 52 polycystic ovarian syndromes by YUAN-clan zishen huoxue decoction. *Journal of Liaoning University of TCM* 2009;**11**(6):151-2.

Zhang 2010a {published data only}

Zhang H. [Clinical observation on traditional Chinese and western medicine for polycystic ovary combine]. *Asia-Pacific Traditional Medicine* 2010;**6**(9):126-7.

Zhang 2010b {published data only}

Zhang M. Clinical study on the "JiaJian Wu Ji San" effect of phlegm-dampness type of polycystic ovary syndrome. Nanjing University of Chinese Medicine 2010.

Zhang 2010c {published data only}

Zhang L, Liu ZD, Ren YM. [From the traditional Chinese medicine ovulation decoction in treatment of polycystic ovary syndrome, 96 cases of clinical observation]. *Inner Mongol Journal of Traditional Chinese medicine* 2010;**29**(18):19-20.

Zhang 2010d {published data only}

Zhang Q. A clinic study of health-related quality-of-life in polycystic ovary syndrome and the impact of TCM method with invigorating kidney, dissipating phlegm and promoting blood flow. Guangzhou University of Chinese Medicine 2010.

Zhang 2010e {published data only}

Zhang T. Polycystic ovary syndrome treated with Chinese medicine "Culuan Pill". *Journal of Zhejiang University of Traditional Chinese Medicine* 2010;**34**(4):515-6.

Zhang 2011a {published data only}

Zhang N. Effect of invigorating kidney and removing phlegm herbs on women with polycystic ovary syndrome undergoing in vitro fertilization and embryo transplantation treatment. *Journal of Liaoning University of TCM* 2011;**13**(7):56-8.

Zhang 2011b {published data only}

Zhang N. Clinical observation of infertile patients with polycystic ovary syndrome treating with Qigongwan and immature follicle puncture technique through ultrasound. *Chinese Imaging Journal of Integrated Traditional and Western Medicine* 2011;**9**(3):202-3.

Zhang 2011c {published data only}

Zhang TH. [Clinical analysis of diagnosis and treatment of 118 cases of patients with polycystic ovary syndrome]. *Journal of China Traditional Chinese Medicine Information* 2011;**3**(20):230.

Zhang 2011d {published data only}

Zhang HH, Cui XP. [Combine traditional Chinese and western medicine treatment of 20 patients with polycystic ovary syndrome]. *Modern Traditional Chinese Medicine* 2011;**31**(4):12-3.

Zhang 2011e {published data only}

Zhang J, Qin JQ. [Combine traditional Chinese and Western medicine in the treatment of infertility caused by polycystic ovary syndrome]. *Journal of Qiqihar University of Medicine* 2011;**32**(5):320.

Zhang 2011f {published data only}

Zhang HM, Tan Y. Clinical study on therapy of nourishing yin and supplementing yang in decreasing androgen level of polycystic ovary syndrome patients with kidney asthenia. *Journal of New Chinese Medicine* 2011;**43**(1):50-2.

Zhang 2012a {published data only}

Zhang JJ. [Clinical observation of traditional Chinese medicine combined with dyne - 35 for polycystic ovary syndrome]. *Journal of Guangxi Traditional Chinese Medical University* 2012;**15**(2):16-7.

Zhang 2012b {published data only}

Zhang LX. The influence of Cang Fu Dao Tan Tang Jia Jian decoction on endocrine and glucose metabolism of polycystic ovary syndrome patients with insulin resistance. Zhejiang Chinese Medical University 2012.

Zhang 2012c {published data only}

Zhang YH, Peng X, Liu XF, Zhang YL, Zhang JS, Xu DJ, et al. [Chinese medicine internal and external washing treatment late onset acne with high androgen hematic disease related analysis]. *Practical Clinical Journal of Integrated Traditional Chinese and Western Medicine* 2012;**12**(6):1-2.

Zhang 2012d {published data only}

Zhang TY, Wang Q. [Phlegm, blood stasis by method of clinical trials for the treatment of ovulation disorder polycystic ovary syndrome]. *Modern Traditional Chinese Medicine* 2012;**32**(2):28-9.

Zhang 2014a {published data only}

Zhang XY. [TCM and Western medicine treatment of insulin resistance in polycystic ovary syndrome patients with clinical observation]. *Health Digest* 2014;**66**(8):189-90.

Zhang 2014b {published data only}

Zhang H, Ma Y, Wang ZL, Gao QQ. Treating 35 cases of polycystic ovary syndrome infertility in the integrative medicine. *Clinical Journal of Chinese Medicine* 2014, (7):33-4, 6.

Zhang 2015a {published data only}

Zhang JX. Efficacy of integrative medicine polycystic ovarian syndrome. *Modern Journal of Integrated Traditional Chinese and Western Medicine* 2015;**24**(14):1562-3.

Zhang 2015b {published data only}

Zhang T. [Effect of Qingre Yangyin recipe on endocrine and metabolism of polycystic ovary syndrome patients]. *Zhongguo Zhong Xi Yi Jie He Za Zhi [Chinese Journal Of Integrated Traditional And Western Medicine]* 2015;**35**(10):1175-80.

Zhang 2015c {published data only}

Zhang T. Affect the patient's endocrine and metabolic heat and nourishing on polycystic ovary syndrome. *Journal of Chinese Medicine* 2015, (10):1175-80.

Zhang 2015d {published data only}

Zhang T, Guo JY. [Effects of Lycii Cortex on express of PI3K/PKB in PCOS rats]. *Zhongguo Zhong Yao Za Zhi [China Journal Of Chinese Materia Medica]* 2015;**40**(10):2004-8.

Zhao 2006a {published data only}

Zhao CP. [Chinese herbal medicine combined with Western medicine in treating PCOS]. *Journal of Shanxi College of Traditional Chinese Medicine* 2006;**7**(6):44-5.

Zhao 2006b {published data only}

Zhao Y. [Clinical observation on acupoint injection in treating PCOS with infertility]. *Liaoning Journal of Traditional Chinese Medicine* 2006;**33**(1):101-2.

Zhao 2006c {published data only}

Zhao Y, Wang YP, Ding QL, Guan HY, Chen M. [Clinical research of acupoint injection of HMG in treating PCOS with infertility]. *Clinical Medicine of China* 2006;**22**(8):748-9.

Zhao 2007 {published data only}

Zhao YL, Li XH, Wen LJ. Observation the effect of polycystic ovarian syndrome by combination treatment of Chinese traditional and western medicine by ultrasonogram. *China Practical Medicine* 2007;**2**(17):22-3.

Zhao 2008a {published data only}

Zhao HB, He YR. Clinical observation of effect of clomiphene citrate combined with cassia tuckahoe capsule on infertile polycystic ovary syndrome patient. *China Tropical Medicine* 2008;**8**(11):1942-3.

Zhao 2008b {published data only}

Zhao H, Wang XE, Bao WF. The effect of metabolic status of blood sugar and lipid and reproductive endocrine of Chinese traditional medicine. *Journal of Zhejiang University of Traditional Chinese Medicine* 2008;**32**(4):458-60.

Zhao 2008c {published data only}

Zhao Y, Huang CL, Ding QL, Wang YP, Chen M. [Influence and efficacy of acupoint injection on endometrium and sexual hormone in PCOS patients]. *Lishizhen Medicine and Materia Medica Research* 2008;**19**(10):2400-1.

Zhao 2009 {published data only}

Zhao H, Bao WF, Zhang T. Advantages of Chinese medicine for treatment of blood sugar and lipid metabolic disorders in patients with polycystic ovarian syndrome. *Zhongguo Zhong Xi Yi Jie He Za Zhi [Chinese Journal of Integrated Traditional and Western Medicine]* 2009;**29**(7):595-8.

Zhao 2010a {published data only}

Zhao J. The clinical observation of the effects of ovulation induction in stimulation cycles of PCOS by sequential therapy of nourishing-yin and supplementing-yang. Nanjing University of Chinese Medicine 2010.

Zhao 2010b {published data only}

Zhao XL, Wang X. [Combine traditional Chinese and Western medicine treatment of polycystic ovary syndrome clinical observation to infertility]. *Journal of Changzhi Medical College* 2010;**24**(2):136-8.

Zhao 2014 {published data only}

Zhao Y, Du B, Jiang X, Ma M, Shi L, Zhang Q, et al. Effects of combining lowdose aspirin with a Chinese patent medicine on follicular blood flow and pregnancy outcome. *Molecular Medicine Reports* 2014;**10**(5):2372-6.

Zhao 2016 {published data only}

Zhao CJ, Jiang H, Zhuang CX. Efficacy of polycystic ovary syndrome new BREE Dodder pills. *Journal of New Chinese Medicine* 2016;**66**(02):148-51.

Zheng 2011 {published data only}

Zheng CC. Phlegm act joint metformin treatment of kidney phlegm type PCOS with IR clinical observation. Hubei University of Chinese Medicine 2011.

Zheng 2011a {published data only}

Zheng YX. Therapeutic effects on renal deficiency type of polycystic ovary syndrome with the periodic therapy of traditional Chinese medicine. Guangzhou University of Chinese Medicine 2011.

Zheng 2011b {published data only}

Zheng YX. The observation of Clinical Efficacy in treating polycystic ovary syndrome by traditional Chinese medicine and Western medicine. *Journal of Traditional Chinese Medicine University of Hunan* 2011;**31**(12):23-4.

Zheng 2011c {published data only}

Zheng GJ, Pan P. [Curative effect observation of combining traditional Chinese and Western medicine treatment of polycystic ovary syndrome]. *Guangming Journal of Chinese Medicine* 2011;**26**(12):2518-9.

Zheng 2014a {published data only}

Zheng YX. Clinical observation of Longdan Xiegan decoction and Bushen Huatan decoction in the treatment of polycystic ovary syndrome. *Chinese Journal Of Family Planning & Gynecotokology* 2014, (6):40-3.

Zheng 2014b {published data only}

Zheng YX. [Integrative medicine clinical observation ovulatory disorder infertility]. *Journal of Inner Mongolia Traditional Chinese Medicine* 2014;**66**(22):67-8.

Zheng 2015a {published data only}

Zheng SJ, Ding JZ, Li Y. [Integrative treatment of polycystic ovary syndrome efficacy]. *Practical Electronics Journal Gynecological Endocrinology* 2015, (9):86, 91.

Zheng 2015b {published data only}

Zheng XH, Zhao XD. Observation of infertility with polycystic ovarian syndrome treated by combination of Pailuan decoction and Clomifene citrate. *Hebei Journal of Traditional Chinese Medicine* 2015, (1):58-60.

Zhi 2012 {published data only}

Zhi J. Clinical observation of Diane-35, metformin combined with Chinese "Bushen Huoxue Tiaozhou therapy" in treatment of polycystic ovary syndrome. Hubei University of Chinese Medicine 2012.

Zhong 2006 {published data only}

Zhong XL, Zhuo XY, Zhang FC, Zhang Z. [Effective observation on nourishing kidney herbal medicine in treating 72 PCOS]. *China and Foreign Medical Journal* 2006;**4**(7):55-7.

Zhong 2008 {published data only}

Zhong XL, Cao DN, Lin WX, Zhang Z, Zheng QY. [Clinical research of Chinese herbal medicine in treating 82 cases with PCOS]. *Journal of New Chinese Medicine* 2008;**40**(11):58-9.

Zhong 2009a {published data only}

Zhong XL, Cao DN, Lin WX, Zheng QY, Zhang Z. [Clinical research of Yugui pill in treating PCOS with insulin resistance]. *Traditional Chinese Medicine Journal* 2009;**8**(3):50-2.

Zhong 2009b {published data only}

Zhong XC, Chen QX, Zhang J, Li XB. Clinical observation of shenqi capsule in treating polycystic ovary syndrome. *Journal of Liaoning University of TCM* 2009;**11**(3):118-20.

Zhong 2012 {published data only}

Zhong GT. The curative effect of Bushen Huoxue decoction to stimulate ovulation in treating polycystic ovary syndrome. *Medical Recapitulate* 2012;**18**(2):306-7.

Zhou 1996 {published data only}

Zhou P, Zhou XC, Liu ML. [Analysis of result of western medicine combined with or without TCM in treating PCOS]. *Practical Woman and Child Health* 1996;**7**(4):30-1.

Zhou 2010a {published data only}

Zhou YH, Huang YJ, Liu W, Yv H. Clinical research on combined therapy of traditional Chinese and Western medicine on polycystic ovarian syndrome. *Liaoning Journal of Traditional Chinese Medicine* 2010;**37**(7):1328-30.

Zhou 2010b {published data only}

Zhou YH, Tan Y. Clinical study on treatment of 21 cases of Stein-Leventhal syndrome with laparoscopic ovarian drilling

combined with Chinese medications. *Jiangsu Journal of Traditional Chinese Medicine* 2010;**42**(9):27-8.

Zhou 2012a {published data only}

Zhou LL. The spleen for eliminating dampness treatments for wet filled the spleen deficiency polycystic ovarian syndrome clinical observation. Heilongjiang University of Chinese Medicine 2012.

Zhou 2012b {published data only}

Zhou XL. [The sterility of polycystic ovary syndrome treatment analysis]. *Chinese Manipulation & Rehabilitation Medicine* 2012;**3**(12):177-8.

Zhou 2012c {published data only}

Zhou JH, Wang ZJ, Wang DJ, Gao WL. Clinical observation of combined acupuncture and herbs in treating polycystic ovary syndrome. *Shanghai Journal of Traditional Chinese Medicine* 2012;**46**(5):72-4.

Zhou 2012d {published data only}

Zhou XY. Clinical study on treatment of polycystic ovary syndrome with Yi-Jing-Bu-Shen decoction. Shandong University of Traditional Chinese Medicine 2012.

Zhou 2014a {published data only}

Zhou YX. Clinical research on combined traditional Chinese and Western medicine treatment of infertility with polycystic ovary syndrome. *China Journal of Chinese Medicine* 2014, (7):1032-3.

Zhou 2014b {published data only}

Zhou Z, Dang YM. [Wenjing decoction observation deficiency polycystic ovary syndrome patients]. *Traditional Chinese Medicine Journal* 2014, (2):56-8.

Zhou 2014c {published data only}

Zhou FB, Xu YL, Du CC. Self prescription treatment of infertility polycystic ovary syndrome phlegm observation of 40 cases. *Chinese Journal of Convalescent Medicine* 2014, (6):519-20.

Zhou 2015a {published data only}

Zhou F. 40 patients in the combination therapy of polycystic ovary syndrome. *Journal Of Medicine Information* 2015, (13):219-20.

Zhou 2015b {published data only}

Zhou M, Zheng L. [Hyperandrogenism clinical observation on 54 cases of integrative medicine polycystic ovary syndrome]. *Journal of Traditional Chinese Medicine* 2015;**66**(02):75-6.

Zhu 2009 {published data only}

Zhu Y, Wang XG. Clinical observation on polycystic ovary syndrome treated with integrated medicine. *Zhejiang Journal of Integrated Traditional Chinese and Western Medicine* 2009;**19**(11):665-7.

Zhu 2012a {published data only}

Zhu JQ. Clinical study on the effect of polycystic ovary syndrome with insulin resistance by the method of invigorating the kidney and regulating menstruation cycles. Nanjing University of Traditional Chinese Medicine 2012.

Zhu 2012b {published data only}

Zhu M. Clinical research on combination Ling-zhu granule and acupoint catgut embedding in treating polycystic ovary syndrome with insulin resistance. Guangzhou University of Chinese Medicine 2012.

Zhu 2013a {published data only}

Zhu TC. Integrative medicine 18 cases of polycystic ovary syndrome. *Guangming Journal of Chinese Medicine* 2013;**28**(7):1427-8.

Zhu 2013b {published data only}

Zhu Y, Wang XG. Infertility treatment in polycystic ovary syndrome with traditional Chinese medicine combined with Western medicine. *Zhejiang Journal of Integrated Traditional Chinese and Western Medicine* 2013, (11):872-5, 85.

Zhu 2014 {published data only}

Zhu XN, Zong LL, Hao S, Guan YC. [Integrative medicine 32 cases of infertility polycystic ovary syndrome]. *Henan Traditional Chinese Medicine* 2014, (7):1355-6.

Zhuang 2008 {published data only}

Zhuang XY, Chen MF. [Clinical research on Chinese herbal medicine combined with clomiphene in treating PCOS]. *Journal of New Chinese Medicine* 2008;**40**(9):61-2.

Zou 2012 {published data only}

Zou XQ. [The curative effect of nursing intervention combined with traditional Chinese and Western medicine treatment for polycystic ovary syndrome]. *Health Must Read Magazine* 2012, issue 4:54.

Zou 2014a {published data only}

Zou Y. [Integrative medicine 33 cases of polycystic ovary syndrome]. *Journal of Community Medicine* 2014, (3):34-5.

Zou 2014b {published data only}

Zou L, Tan AH. Polycystic ovary syndrome in combination therapy effectiveness analysis. *Journal of Medical Information* 2014, (21):611-2.

Zuo 2011 {published data only}

Zuo J. The clinical study on the "jiajian fang fen tong shen san" effect of overweight type of polycystic ovarian syndrome. Nanjing University of Chinese Medicine 2011.

References to ongoing studies
ChiCTR-IOR-16008557 {published data only}

ChiCTR-IOR-16008557. The Qilingwenshenxiaonang recipe in the treatment of patients with Polycystic ovary syndrome (PCOS): a multicenter, randomised, double-blind, placebo controlled trial. <http://www.chictr.org.cn/showproj.aspx?proj=14360> (accessed 9 June 2016). [ChiCTR-IOR-16008557]

NCT01116167 {published data only}

Li Y, Kuang H, Shen W, Ma H, Zhang Y, Stener-Victorin E, et al. Letrozole, berberine, or their combination for anovulatory infertility in women with polycystic ovary syndrome: study

design of a double-blind randomised controlled trial. *BMJ Open* 2013;**3**(11):e003934.

Additional references
Alvarez-Blasco 2006

Alvarez-Blasco F, Botella-Carretero JI, San Millán JL, Escobar-Morreale HF. Prevalence and characteristics of the polycystic ovary syndrome in overweight and obese women. *Archives of Internal Medicine* 2006;**166**(19):2081-6.

Berneis 2007

Berneis K, Rizzo M, Lazzaroni V, Fruzzetti F, Carmina E. Atherogenic lipoprotein phenotype and low-density lipoproteins size and subclasses in women with polycystic ovary syndrome. *Journal of Clinical Endocrinology and Metabolism* 2007;**92**(1):186-9.

Boomsma 2006

Boomsma CM, Eijkemans MJ, Hughes EG, Visser GH, Fauser BC, Macklon NS. A meta-analysis of pregnancy outcomes in women with polycystic ovary syndrome. *Human Reproduction Update* 2006;**12**(6):673-83.

Carmina 1999

Carmina E, Lobo RA. Polycystic ovary syndrome (PCOS): arguably the most common endocrinopathies associated with significant morbidity in women. *Journal of Clinical Endocrinology and Metabolism* 1999;**84**(6):1897-9.

Cong 2006

Cong LX. [Observation of Tiaojin Zhuyun Pellet combined with clomiphene in treating infertility caused by PCOS]. *Journal of Practical Traditional Chinese Medicine* 2006;**22**(5):290-1.

Costello 2007

Costello M, Shrestha B, Eden J, Sjoblom P, Johnson N. Insulin-sensitising drugs versus the combined oral contraceptive pill for hirsutism, acne and risk of diabetes, cardiovascular disease, and endometrial cancer in polycystic ovary syndrome. *Cochrane Database of Systematic Reviews* 2007, Issue 1. [DOI: [10.1002/14651858.CD005552.pub2](https://doi.org/10.1002/14651858.CD005552.pub2)]

Cui 2004

Cui FY, Liu XM. [Gaoshao soup in treating 60 cases of PCOS]. *Journal of Practical Traditional Chinese Medicine* 2004;**20**(12):686-7.

Dunaif 1997

Dunaif A. Insulin resistance and the polycystic ovary syndrome mechanism and implications for pathogenesis. *Endocrine Reviews* 1997;**18**(6):774-800.

Egger 1997

Egger M, Smith GD, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997;**315**(7109):629-34.

Ehrmann 1999

Ehrmann DA, Barnes RB, Rosenfield RL, Cavaghan MK, Imperial J. Prevalence of impaired glucose tolerance and diabetes in women with polycystic ovary syndrome. *Diabetes Care* 1999;**22**(1):141-6.

Escobar-Morreale 2005

Escobar-Morreale HF, Luque-Ramírez M, San Millán JL. The molecular genetic basis of functional hyperandrogenism and polycystic ovary syndrome. *Endocrine Reviews* 2005;**26**(2):251-82.

ESHRE/ASRM 2004

The Rotterdam ESHRE/ASRM Sponsored PCOS consensus workshop group. Revised 2003 consensus on diagnostic criteria and long-term health risks related to polycystic ovary syndrome (PCOS). *Human Reproduction* 2004;**19**(1):41-7.

Farquhar 2012

Farquhar C, Brown J, Marjoribanks J. Laparoscopic drilling by diathermy or laser for ovulation induction in anovulatory polycystic ovary syndrome. *Cochrane Database of Systematic Reviews* 2012, Issue 6. [DOI: [10.1002/14651858.CD001122.pub4](https://doi.org/10.1002/14651858.CD001122.pub4)]

Gambineri 2006

Gambineri A, Pasquali R. Insulin resistance, obesity and metabolic syndrome in polycystic ovary syndrome. *Endocrinología y Nutrición* 2006;**53**(S1):49-55.

Higgins 2002

Higgins JPT, Thompson SG. Quantifying heterogeneity in a metaanalysis. *Statistics in Medicine* 2002;**21**(11):1539-58.

Higgins 2011

Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions*. Version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011. Available from www.cochrane-handbook.org.

Hou 2012

Hou F, Shi YQ. [The progress on etiology of polycystic ovary syndrome in traditional Chinese medicine]. *Journal of Changchun University of Traditional Chinese Medicine* 2012;**28**(5):932-34.

ICHEWG 1997

International Conference on Harmonisation Expert Working Group. International conference on harmonisation of technical requirements for registration of pharmaceuticals for human use. ICH harmonised tripartite guideline. Guideline for good clinical practice. International conference on harmonisation of technical requirements for registration of pharmaceuticals for human use. ICH harmonised tripartite guideline. Guideline for good clinical practice. CFR & ICH Guidelines. 1st Edition. Needham, Massachusetts, USA: Barnett International/PAREXEL, 1997:19063-2043.

Jadad 1996

Jadad AR, Moore RA, Carroll D, Jenkinson C, Reynolds DJ, Gavaghan DJ, et al. Assessing the quality of reports of

randomized clinical trials: is blinding necessary?. *Controlled Clinical Trials* 1996;**17**(1):1-12.

Jin 1986

Jin WX, Sun SX, Wang CY. [Research on Luole in treating female ovulatory disorder]. *Journal of Fujian College of Traditional Chinese Medicine* 1986;**10**(2):11-3.

Jüni 2001

Jüni P, Altman DG, Egger M. Systematic reviews in health care: Assessing the quality of controlled clinical trials. *BMJ* 2001;**323**(7303):42-6.

Kjaergard 2001

Kjaergard LL, Villumsen J, Gluud C. Reported methodological quality and discrepancies between large and small randomized trials in meta-analyses. *Annals of Internal Medicine* 2001;**135**(11):982-9.

Lampert 2002

Lampert N, Xu Y. Chinese herbal nephropathy. *The Lancet* 2002;**359**(9308):796-7.

Legro 1999

Legro RS, Kunselman AR, Dodson WC, Dunaif A. Prevalence and predictors of risk for type 2 diabetes mellitus and impaired glucose tolerance in polycystic ovary syndrome a prospective, controlled study in 254 affected women. *Journal of Clinical Endocrinology and Metabolism* 1999;**84**(1):165-9.

Legro 2001

Legro RS, Kunselman AR, Dunaif A. Prevalence and predictors of dyslipidemia in women with polycystic ovary syndrome. *American Journal of Medicine* 2001;**111**(8):607-13.

Li 2006

Li CP. [Effective observation on Bushen Tiaozhou Method in treating 30 infertility patients with PCOS]. *New Journal of Traditional Chinese Medicine* 2006;**38**(1):50-1.

Liu 2005

Liu Y, Lu XY. [Traditional Chinese medicine in treating 12 PCOS]. *New Journal of Traditional Chinese Medicine* 2005;**37**(8):74-5.

Lo 2006a

Lo JC, Feigenbaum SL, Escobar GJ, Yang J, Crites YM, Ferrara A. Increased prevalence of gestational diabetes mellitus among women with diagnosed polycystic ovary syndrome a population based study. *Diabetes Care* 2006;**29**(8):1915-7.

Lo 2006b

Lo JC, Feigenbaum SL, Yang J, Pressman AR, Selby JV, Go AS. Epidemiology and adverse cardiovascular risk profile of diagnosed polycystic ovary syndrome. *Journal of Clinical Endocrinology and Metabolism* 2006;**91**(4):1357-63.

Lord 2001

Lord GM, Cook T, Arlt VM, Schmeiser HH, Williams G, Pusey CD. Urothelial malignant disease and Chinese herbal nephropathy. *The Lancet* 2001;**358**(9292):1515-6.

Moher 1998

Moher D, Pham B, Jones A, Cook DJ, Jadad A, Moher M, et al. Does quality of reports of randomised trials affect estimates of intervention efficacy reported in meta-analysis. *The Lancet* 1998;**352**(9128):609-13.

Ni 2007

Ni HY, Gong J. [Research progress on Chinese herbal medicine in treating PCOS]. *Liaoning Journal of Traditional Chinese Medicine* 2007;**34**(1):123-4.

Ning 2004

Ning MH, Liu YJ, Ning XG. Clinical observation on Yishenxiaozheng decoction for the treatment of 85 cases of polycystic ovarian disease. *Hunan Guiding Journal of TCM* 2004;**10**(3):27-8.

Nortier 2000

Nortier JL, Martinez MC, Schmeiser HH, Arlt VM, Bieler CA, Petein M, et al. Urothelial carcinoma associated with the use of a Chinese herb (*Aristolochia fangchi*). *The New England Journal of Medicine* 2000;**342**(23):1686-92.

Rajkhowa 1997

Rajkhowa M, Neary RH, Kumpatla P, Garne FL, Jones PW, Obhrai MS, et al. Altered composition of high density lipoproteins in women with the polycystic ovary syndrome. *Journal of Clinical Endocrinology and Metabolism* 1997;**82**(10):3389-94.

RevMan 2014 [Computer program]

The Nordic Cochrane Centre, The Cochrane Collaboration. Review Manager (RevMan). Version 5.3. Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration, 2014.

Schulz 1995

Schulz KF, Chalmers I, Hayes RJ, Altman DG. Empirical evidence of bias. Dimensions of methodological quality associated with estimates of treatment effects in controlled trials. *JAMA* 1995;**273**(5):408-12.

Sinawat 2012

Sinawat S, Buppasiri P, Lumbiganon P, Pattanittum P. Long versus short course treatment with metformin and clomiphene citrate for ovulation induction in women with PCOS. *Cochrane Database of Systematic Reviews* 2012, Issue 10. [DOI: [10.1002/14651858.CD006226.pub3](https://doi.org/10.1002/14651858.CD006226.pub3)]

Sun 1981

Sun YL, Guo HR, Yu J, Li CJ. [133 cases of polycystic ovarian syndrome treated by the method 'nourishing kidney and sputum-reducing']. *Shanghai Journal of TCM* 1981;**15**(06):14.

Talbott 1998

Talbott I, Clerici A, Berga SL, Kuller L, Guzick D, Detre K, et al. Adverse lipid and coronary heart disease risk profiles in young women with polycystic ovary syndrome results of a case control study. *Journal of Clinical Epidemiology* 1998;**51**(5):415-22.

Tang 2012

Tang T, Lord JM, Norman RJ, Yasmin E, Balen AH. Insulin-sensitising drugs (metformin, rosiglitazone, pioglitazone, D-chiro-inositol) for women with polycystic ovary syndrome, oligo amenorrhoea and subfertility. *Cochrane Database of Systematic Reviews* 2012, Issue 5. [DOI: [10.1002/14651858.CD003053.pub5](https://doi.org/10.1002/14651858.CD003053.pub5)]

Wang 1982

Wang ZQ, Shi LY, Wang DZ. [The effect of "Long Dan Xie Gan Tang" treat polycystic ovary syndrome]. *Shanghai Journal of Traditional Chinese Medicine* 1982, (12):16-7.

Wang 2005

Wang ZH, Yang YS, Zhang YL. Clinical study of Ganshao capsule in treating clomiphene-resistant polycystic ovarian syndrome. *Zhongguo Zhong Xi Yi Jie He Za Zhi [Chinese Journal of Integrated Traditional and Western Medicine]* 2005;**25**(8):704-6.

Wang 2008

Wang BQ, Ling M. [Research development of Chinese herbal medicine for PCOS]. *Shandong Journal of Traditional Chinese Medicine* 2008;**27**(2):138-40.

Weiss 2015

Weiss NS, Nahuis M, Bayram N, Mol BWJ, Van der Veen F, van Wely M. Gonadotrophins for ovulation induction in women with polycystic ovarian syndrome. *Cochrane Database of Systematic Reviews* 2015, Issue 9. [DOI: [10.1002/14651858.CD010290.pub2](https://doi.org/10.1002/14651858.CD010290.pub2)]

Yang 2006

Yang ZW, You ZL, Zhang XH, Wang Y, Zeng M. [Research on influence of Bushen Huoxue Method on menstrual cyclicity and reproductive hormone in PCOS]. *Chinese Journal of Traditional Medical Science and Technology* 2006;**13**(1):5-6.

Yuan 2003

Yuan XF. [TCM periodical treatment on 38 PCOS]. *Fujian Journal of Traditional Chinese Medicine* 2003;**34**(2):22.

Yv 1981

Yv XL, Li YA. [99 cases of Chinese and Western treatment of polycystic ovary syndrome]. *Shanghai Medical Journal* 1981;**4**(1):50.

Zhang 2004

Zhang QP. [Bushen Huoxue Method in treating PCOS]. *Chinese Journal of Information on TCM* 2004;**11**(11):1014-5.

References to other published versions of this review

Zhang 2009

Zhang J, Zhou L, Tang L, Wu T, Lim DCE. Chinese herbal medicine for subfertile women with polycystic ovarian syndrome. *Cochrane Database of Systematic Reviews* 2009, Issue 1. [DOI: [10.1002/14651858.CD007535](https://doi.org/10.1002/14651858.CD007535)]

Zhang 2010

Zhang J, Li T, Zhou L, Tang L, Xu L, Wu T, et al. Chinese herbal medicine for subfertile women with polycystic ovarian

syndrome. *Cochrane Database of Systematic Reviews* 2010, Issue 9. [DOI: [10.1002/14651858.CD007535.pub2](https://doi.org/10.1002/14651858.CD007535.pub2)]

CHARACTERISTICS OF STUDIES

Characteristics of included studies [ordered by study ID]

Li 2007

Methods	RCT, single centre, 90 participants, 3 years' duration
Participants	<p>90 enrolled: CHM 1 = 30, CHM 2 = 30, control = 30, 21 to 38 years, baseline were comparable</p> <p>87 analysed/evaluated: CHM1 = 29 (1 converted to IVF-ET after CHM treating for 1 month), CHM2 = 30, control = 28 (1 moved to other place, 1 discontinued therapy because of pelvic inflammation)</p> <p>Obesity: CHM1 = 7, CHM2 = 6, control = 6</p> <p>Hirsutism: CHM1 = 19, CHM2 = 18, control = 21</p> <p>LH/FSH > 2.5: CHM1 = 20, CHM2 = 19, control = 21</p> <p>High testosterone: CHM1 = 16, CHM2 = 17, control = 15</p> <p>Follicle number > 10: CHM1 = 25, CHM2 = 24, control = 22</p> <p>Enlarged ovary: CHM1 = 5, CHM2 = 6, control = 7</p> <p>PCOS DC: consistent with Rotterdam criteria (evaluated by review authors)</p> <p>In: PCOS and infertility</p> <p>Ex: using other drugs of ovulation induction, participants unable to follow-up, tumour patients, adrenal diseases, other hyper androgenic diseases</p>
Interventions	<p>CHM1: clomiphene simulacrum (5 to 9 day of menstrual cycle, 1 pill, once a day, 5 days), Lingzhu infusion (5 to 14 day of menstrual cycle, 1 bag, tid, 10 days), Shenqi capsule (from 14th day of menstrual cycle or after ovulation, 4 grains, tid, until menstrual onset or pregnancy or the 45th day of menstrual cycle), if amenorrhoea for 45 days then MPA would be prescribed (10 mg, once a day, 5 days)</p> <p>CHM2: clomiphene (5 to 9 day of menstrual cycle, 50 mg, once a day, 5 days), Lingzhu infusion, Shenqi capsule, and MPA</p> <p>Control: clomiphene, Lingzhu simulacrum, Shenqi simulacrum, and MPA</p> <p>Duration: treated no more than 6 menstrual cycles, follow-up time was unclear.</p>
Outcomes	<p>LH, testosterone, LH/FSH, estradiol, insulin, BMI, cervical mucus</p> <p>Pregnancy rate (per woman)</p> <p>Ovulation rate (per cycle)</p>
Notes	<p>Clomiphene: Codal Synto Ltd. batch number: H20020325</p> <p>Lingzhu infusion: hospital preparation, batch number Z03020211, 6 g/bag</p> <p>Shenqi capsule: hospital preparation, batch number Z03020212, 0.5 g/pill</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Li 2007 (Continued)

Random sequence generation (selection bias)	Unclear risk	We were unable to contact the study authors for more information
Allocation concealment (selection bias)	Unclear risk	We were unable to contact the study authors for more information
Blinding (performance bias and detection bias) All outcomes	Low risk	The study used mimic drugs. Participants and the outcome assessor were blinded
Incomplete outcome data (attrition bias) All outcomes	Low risk	No intention-to-treat (ITT) analysis. The analysis rate was 96.7% (87/90)
Selective reporting (reporting bias)	Unclear risk	The study protocol was unavailable
Other bias	Low risk	No other potential risk of bias identified

Li 2012a

Methods	RCT, single centre, 70 participants, 1 year duration
Participants	<p>70 enrolled: CHM = 35, control = 35, 22 to 39 years, baseline were comparable</p> <p>70 analysed/evaluated: CHM = 35, control = 35</p> <p>Age (years): CHM 28.5 ± 3.8, control 26.2 ± 3.6</p> <p>Subfertility time (years): CHM 5 ± 2.7, control 4.6 ± 2.4</p> <p>PCOS DC: consistent with Rotterdam criteria (evaluated by review authors)</p> <p>In: PCOS, infertility, 20 to 40 years</p> <p>Ex: using hormone or drugs of ovulation induction in the last 3 months, tubal infertility, uterine infertility, male sterility</p>
Interventions	<p>CHM: Xuanju capsule (day 3 of menstrual cycle, 3 pills, tid, 4 weeks), clomiphene (day 3 of menstrual cycle, 50 mg, once a day, 5 days), HCG was injected when dominant follicle was present, if dominant follicle was absent until the 20 day of menstrual cycle, progesterone was injected 20 mg, once a day, 5 days</p> <p>Control: clomiphene (day 3 of menstrual cycle, 50 mg, once a day, 5 days), HCG was injected when dominant follicle was present, if dominant follicle was absent until the 20 day of menstrual cycle, progesterone was injected 20 mg, once a day, 5 days</p> <p>Duration: treatment until pregnancy but no more than 3 cycles; follow-up duration was unclear</p>
Outcomes	<p>Pregnancy rate (per woman)</p> <p>Ovulation rate (per cycle)</p>
Notes	Xuanju capsule: Zhejiang Shiqiang Pharmaceutical Company, batch number: Z20060462

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Li 2012a (Continued)

Random sequence generation (selection bias)	Low risk	Random number table
Allocation concealment (selection bias)	Unclear risk	We were unable to contact the study authors for more information
Blinding (performance bias and detection bias) All outcomes	Unclear risk	We were unable to contact the study authors for more information
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	No withdrawal or drop-out was reported.
Selective reporting (reporting bias)	Unclear risk	The study protocol was unavailable.
Other bias	Low risk	No other potential risk of bias identified

Liang 2008

Methods	RCT, 2 clinical centres, 44 participants, 1 year duration
Participants	<p>44 enrolled: CHM = 20, control = 24, baseline was comparable</p> <p>44 analysed/evaluated: CHM = 20, control = 24</p> <p>40 ovulation induction: CHM = 18, control = 22 (follicle aspiration was ineffective for 4)</p> <p>Age (years): CHM 27.4 ± 2.7, control 27.1 ± 3.2</p> <p>Subfertility time (years): CHM 2.10 ± 0.97, control 2.0 ± 0.84</p> <p>BMI (kg/m^2): CHM 24.2 ± 2.9, control 25.2 ± 3.1</p> <p>PCOS DC: 2003 Rotterdam criteria</p> <p>In: PCOS patients with infertility and clomiphene resistance (clomiphene 150 mg/d, 5 d/month, 3 months, but without follicle growth)</p> <p>Ex: other endocrinology diseases, tubal infertility, male sterility</p>
Interventions	<p>CHM interventions: Bushen Huoxue formula combined with ultrasound guided follicle aspiration and ovulation induction</p> <p>Control interventions: ultrasound guided follicle aspiration and ovulation induction</p> <p>Ultrasound guided follicle aspiration: on 10th to 12th day of menstrual cycle, 36 hours after HCG (10000 IU) injection, bilateral ovaries, 2 to 4 times of inserting per ovary, once a month until presence of efficacy but no more than 3 months (efficacy was defined as testosterone $< 1.6 \text{ nmol/L}$, LH/FSH < 2, number of antral follicle in each ovary were less than 10 at early follicle phase of the following menstrual cycle)</p> <p>Bushen Huoxue formula: from 5th day of menstrual cycle, 1 dose/day, 14 days</p> <p>Ovulation induction: after effective follicle aspiration, no more than 3 cycles, human menopausal gonadotrophin (HMG) (from 5th day of menstrual cycle, 15 to 150IU/d, until presence of dominant follicle), then HCG (5000 to 10000 IU)</p> <p>Duration: treatment: no more than 6 menstrual cycles, follow-up: 3 months after ovulation induction</p>

Liang 2008 (Continued)

Outcomes	FSH, LH, testosterone Number of antral follicle Pregnancy rate (per woman) Dosage of HMG Side effects: LUFS, OHSS, multiple pregnancy Number of mature follicles
Notes	Blood hormone level and ultrasound were usually measured at 3rd day of menstrual cycle.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Random number table
Allocation concealment (selection bias)	High risk	Random number was open
Blinding (performance bias and detection bias) All outcomes	High risk	No blinding. We contacted the study author for this information
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	No ITT analysis. The analysis rate was 90.9% (40/44)
Selective reporting (reporting bias)	Unclear risk	The study protocol was unavailable
Other bias	Low risk	No other potential risk of bias identified

Ma 2009a

Methods	RCT, single centre, 170 participants, 4 years' duration
Participants	170 enrolled: CHM = 85, control = 85, baseline was comparable 165 analysed/evaluated: CHM = 85, control = 80 (5 withdrawals for personal reasons) Age (years): CHM: 28.4 ± 5.3, control: 27.9 ± 4.9 Infertility time (years): CHM: 3.8 ± 2.1, control: 3.6 ± 1.9 PCOS DC: 2003 Rotterdam criteria In: PCOS and infertility Ex: other endocrinology diseases, hormone user in the last 3 months, male infertility, tubal infertility
Interventions	CHM: CHM combined with ethinyloestradiol cyproterone acetate and ovulation induction Control: CEA following with ovulation induction

Ma 2009a (Continued)

CHM: basic formula in CEA therapy duration, CHM periodic therapy in ovulation induction phase (gui shao di huang soup in 5th to 14th day of menstrual cycle, tao hong si wu soup in 12th to 16th day of menstrual cycle, shou tai pellet after ovulation)

CEA: from 5th day of menstrual cycle, 1 pill, once a day, 21 days/m, treated for 3 cycles and then ovulation induction

ovulation induction: clomiphene (from 5th day of menstrual cycle, 50 mg, once a day, 5 days/m), 5000 to 10000 IU HCG was injected when dominant follicle was present, ovulation induction until pregnancy but no more than 3 cycles

Duration: treatment: no more than 6 menstrual cycles, follow-up time was unclear.

Outcomes	Ovulation rate (per cycle) Pregnancy rate (per woman) Miscarriage rate
Notes	Ethinylestradiol cyproterone acetate: Germany Schering company, batch number: G20040104

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Random number table
Allocation concealment (selection bias)	Unclear risk	We contacted the study author who declined to provide related information
Blinding (performance bias and detection bias) All outcomes	Unclear risk	We contacted the study author who declined to provide related information
Incomplete outcome data (attrition bias) All outcomes	Low risk	No ITT analysis. The analysis rate was 97.1% (165/170)
Selective reporting (reporting bias)	Unclear risk	The study protocol was unavailable
Other bias	Low risk	No other potential risk of bias identified

Ye 2007

Methods	RCT, single centre, 40 participants, 20 months' duration
Participants	40 enrolled: CHM = 20, control 1 = 10, control 2 = 10, baseline were comparable, 27.4 ± 2.7 years 40 analysed/evaluated: CHM = 20, control 1 = 10, control 2 = 10 PCOS DC: 2003 Rotterdam criteria In: PCOS and infertility and resistance to ovulation induction drugs Ex: tubal infertility, male infertility, malformation of genital organ, immunological infertility
Interventions	CHM: CHM periodic therapy combined with laparoscopic ovary drilling

Chinese herbal medicine for subfertile women with polycystic ovarian syndrome (Review)

Ye 2007 (Continued)

Control 1: clomiphene combined with laparoscopic ovary drilling

Control 2: laparoscopic ovary drilling

Duration: treatment: 6 months, follow-up: 1 year

Clomiphene: 50 mg, once a day, 5d/m (if without efficacy, add 50 mg, maximum 150 mg/d)

Outcomes	LH, FSH, testosterone Ovulation rate (per woman) Pregnancy rate (per woman)
Notes	Laparoscopic ovary drilling: 8 to 10 holes per ovary, injected 300 mL of low molecular dextran or 4 mL of sodium hyaluronate in abdomen after surgery.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Random number table
Allocation concealment (selection bias)	Unclear risk	We contacted the study author who refused to provide related information
Blinding (performance bias and detection bias) All outcomes	Unclear risk	We contacted the study author who refused to provide related information
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	No withdrawal or drop-out was reported.
Selective reporting (reporting bias)	Unclear risk	The study protocol was unavailable.
Other bias	Low risk	No other potential risk of bias identified

Abbreviations: CHM: Chinese herbal medicine, BMI: body mass index, PCOS: polycystic ovarian syndrome, DC: diagnosis criteria, In: inclusion criteria, Ex: exclusion criteria, HCG: human chorionic gonadotrophin, HMG: human menopausal gonadotropin, LH: luteinizing hormone, FSH: follicle stimulating hormone, LUFS: luteinized unruptured follicle syndrome, OHSS: ovarian hyperstimulation syndrome, ITT: intention to treat, MPA: medroxyprogesterone acetate, CEA: ethinyloestradiol cyproterone acetate.

Characteristics of excluded studies [ordered by study ID]

Study	Reason for exclusion
An 2009	No outcomes of interest
An 2012	Diagnosis is inconsistent with Rotterdam criteria
Arentz 2014	Review
Bablis 2006	Case report

Study	Reason for exclusion
Bai 2011	Non-randomized controlled trials (RCTs)
Bao 2009	No outcomes of interest; polycystic ovarian syndrome (PCOS) with or without infertility in this study
Bao 2014	No Chinese herbal medicine (CHM) intervention
Bei 2010	Non-RCT, which the primary study authors confirmed
Cai 2006	Adolescent PCOS without infertility; no outcomes of interest
Cai 2011	Quasi-RCT, which the primary study authors confirmed
Cai 2012	PCOS diagnosis is inconsistent with Rotterdam criteria
Cai 2014	Not a RCT
Cao 2010	Non-RCT, which the primary study authors confirmed
Cao 2012	Non-RCT, which the primary study authors confirmed
Chan 2006a	No outcomes of interest; PCOS with or without infertility
Chan 2006b	Non-CHM intervention
Chen 2005	PCOS with or without infertility in this study
Chen 2006a	PCOS diagnosis is inconsistent with Rotterdam criteria; quasi-RCT
Chen 2006b	PCOS with or without infertility in this study
Chen 2007	Intervention with acupuncture but without herbal medicine
Chen 2008	No outcomes of interest; PCOS with or without infertility in this study
Chen 2009	Intervention with acupuncture but without herbal medicine
Chen 2010a	No outcomes of interest
Chen 2010b	No outcomes of interest
Chen 2011a	No outcomes of interest; PCOS with or without infertility
Chen 2011b	No outcomes of interest; PCOS with or without infertility
Chen 2011c	Quasi-RCT
Chen 2012a	No outcomes of interest; PCOS with or without infertility
Chen 2012b	No outcomes of interest; PCOS with or without infertility
Chen 2012c	Quasi-RCT
Chen 2012d	PCOS diagnosis is inconsistent with Rotterdam criteria; quasi-RCT

Study	Reason for exclusion
Chen 2013	Non-PCOS participants
Chen 2014a	Quasi-RCT
Chen 2014b	Not a RCT
Chen 2014c	Non-CHM intervention
Chen 2015	Participants had no wish to conceive
Chen 2016	Non-CHM intervention
Cheng 2009	No outcomes of interest; PCOS with or without infertility in this study
Cheng 2014	Non-CHM intervention
Cheng 2015	Participants had no wish to conceive
Chu 2013	Non-PCOS participants
Craig 2015	Non-CHM intervention
Cui 2012	PCOS with or without infertility; no outcomes of interest
Dang 2012	Non-RCT, which the primary authors confirmed
Deng 2008	PCOS diagnosis is inconsistent with Rotterdam criteria; quasi-RCT
Deveci 2015	Non-CHM intervention
Ding 2015	Unrelated study
Dong 2009	No outcomes of interest
Dong 2010	No outcomes of interest
Du 2012	PCOS with or without infertility; no outcomes of interest
Du 2013	Participants had no wish to conceive
Fang 2004	No outcomes of interest
Feng 2009a	No outcomes of interest; PCOS with or without infertility in this study
Feng 2009b	No outcomes of interest; PCOS with or without infertility
Feng 2011a	Non-RCT confirmed by primary authors
Feng 2011b	Quasi-RCT
Fu 2012	PCOS diagnosis is inconsistent with Rotterdam criteria; PCOS with or without infertility in this study; no outcomes of interest
Gao 2009	Intervention with acupuncture but without herbal medicine

Study	Reason for exclusion
Gao 2011a	PCOS diagnosis is inconsistent with Rotterdam criteria; quasi-RCT
Gao 2011b	PCOS with or without infertility in this study
Ghavi 2015	With or without subfertility
Gong 2012	PCOS with or without infertility in this study
Grant 2010	No outcomes of interest; PCOS with or without infertility
Gu 2015	Participants had no wish to conceive
Guo 2008	No outcomes of interest; PCOS with or without infertility
Guo 2009	PCOS with or without infertility in this study
Guo 2011a	PCOS diagnosis is inconsistent with Rotterdam criteria; quasi-RCT
Guo 2011b	No outcomes of interest; PCOS with or without infertility in this study
Haj-Husein 2016	With or without subfertility
Han 2008	PCOS diagnosis is inconsistent with Rotterdam criteria; quasi-RCT
Han 2011	PCOS diagnosis is inconsistent with Rotterdam criteria; quasi-RCT
Han 2013a	Participants had no wish to conceive
Han 2013b	Not a RCT
Han 2015	Not a RCT
Hao 2012	Quasi-RCT, which the primary authors confirmed
Harman 2001	Non-PCOS
Hassanzadeh Bashtian 2013	Participants had no wish to conceive
He 2009	PCOS with or without infertility in this study
He 2010	Quasi-RCT
He 2014	Quasi-RCT
Hou 2000	No outcomes of interest
Hu 2009a	Intervention with acupuncture but without herbal medicine; PCOS with and without infertility
Hu 2009b	Intervention with acupuncture but without herbal medicine
Hu 2014	Participants had no wish to conceive
Hua 2003	Case control study
Huang 2004	Quasi-RCT confirmed by primary authors

Study	Reason for exclusion
Huang 2006a	Non-RCT confirmed by primary authors
Huang 2006b	PCOS without infertility; quasi-RCT; No outcomes of interest
Huang 2007	PCOS with or without infertility in this study
Huang 2008	Review
Huang 2010	Quasi-RCT
Huang 2011a	No outcomes of interest
Huang 2011b	No outcomes of interest
Huang 2012a	No outcomes of interest; PCOS with or without infertility
Huang 2012b	PCOS diagnosis is inconsistent with Rotterdam criteria
Hung 2016	Cohort study
Huo 2008	Unrelated
Jalilian 2013	Non-CHM intervention
Jia 2004	Diagnosis inconsistent with Rotterdam; quasi-RCT
Jia 2008	Concurrent control study
Jia 2010	PCOS with or without infertility; PCOS diagnosis is inconsistent with Rotterdam criteria
Jia 2012a	No outcomes of interest
Jia 2012b	PCOS diagnosis is inconsistent with Rotterdam criteria; quasi-RCT; no outcomes of interest; PCOS with or without infertility
Jian 2011	No outcomes of interest
Jiang 2007	PCOS diagnosis is inconsistent with Rotterdam criteria; quasi-RCT
Jiang 2011a	No outcomes of interest; PCOS with or without infertility
Jiang 2011b	No outcomes of interest; PCOS with or without infertility
Jiang 2014	Non-PCOS participants
Jiang 2015	Non-CHM intervention
Jin 2014a	Quasi-RCT
Jin 2014b	Non-CHM intervention
Jin 2016	Participants had no wish to conceive
Johnson 2015	Non-CHM intervention

Study	Reason for exclusion
Kang 2012	No outcomes of interest; PCOS with or without infertility
Kawakami 2011	Unrelated
Kitagawa 2015	Non-PCOS participants
Kort 2014	With or without subfertility
Kuang 2012	Quasi-RCT, which the primary authors confirmed
Kuang 2013	Non-CHM intervention
Kuang 2015	Non-CHM intervention
Kuek 2011	No outcomes of interest; PCOS with or without infertility in this study
Lai 2006	No outcomes of interest; PCOS with or without infertility; PCOS diagnosis is inconsistent with Rotterdam criteria
Lai 2011	Quasi-RCT
Lai 2014a	Participants had no wish to conceive
Lai 2014b	Participants had no wish to conceive
Lai 2014c	Participants had no wish to conceive
Lai 2015a	Review
Lai 2015b	Participants had no wish to conceive
Lai 2015c	Participants had no wish to conceive
Lai 2015d	Participants had no wish to conceive
León-Gonzalez 2014	Unrelated
Li 2000	PCOS with or without infertility in this study
Li 2002	PCOS diagnosis is inconsistent with Rotterdam criteria
Li 2005	PCOS diagnosis is inconsistent with Rotterdam criteria; quasi-RCT
Li 2009a	No outcomes of interest
Li 2009b	PCOS without infertility
Li 2009c	Duplication
Li 2009d	No outcomes of interest; PCOS with or without infertility
Li 2010a	No outcomes of interest; PCOS with or without infertility; PCOS diagnosis is inconsistent with Rotterdam criteria
Li 2010b	No outcomes of interest; PCOS with or without infertility

Study	Reason for exclusion
Li 2010c	PCOS diagnosis is inconsistent with Rotterdam criteria
Li 2010d	No outcomes of interest; PCOS with or without infertility
Li 2011a	No outcomes of interest
Li 2011b	Participants had no wish to conceive
Li 2011c	PCOS with or without infertility; no outcomes of interest
Li 2011d	No outcomes of interest; PCOS with or without infertility
Li 2011e	No outcomes of interest; PCOS with or without infertility; PCOS diagnosis is inconsistent with Rotterdam criteria
Li 2011f	PCOS with or without infertility in this study
Li 2011g	No outcomes of interest
Li 2011h	No outcomes of interest; PCOS with or without infertility
Li 2011i	No outcomes of interest; PCOS with or without infertility
Li 2011j	PCOS diagnosis is inconsistent with Rotterdam criteria
Li 2011k	No outcomes of interest
Li 2012b	No outcomes of interest
Li 2012c	No outcomes of interest; PCOS with or without infertility
Li 2013a	Non-CHM intervention
Li 2013b	Participants had no wish to conceive; protocol
Li 2015	Non-CHM intervention
Li 2016	Participants had no wish to conceive
Lian 2008	Quasi-RCT
Lian 2012	Quasi-RCT
Liang 2011	Quasi-RCT
Liao 2014	Non-CHM intervention
Lim 2011	Review
Lin 2005	Non-RCT, which the primary authors confirmed
Lin 2009a	Non-RCT, which was confirmed by the author
Lin 2009b	Quasi-RCT

Study	Reason for exclusion
Lin 2011	We were unable to contact the study authors for the detailed information about the laparoscopic surgery method
Lin 2013a	Participants had no wish to conceive
Lin 2013b	Participants had no wish to conceive
Lin 2013c	Participants had no wish to conceive
Liu 2007	PCOS diagnosis is inconsistent with Rotterdam criteria; quasi-RCT
Liu 2008	Unrelated
Liu 2009	Diagnosis inconsistent with Rotterdam; quasi-RCT
Liu 2010a	Quasi-RCT
Liu 2010b	Non-RCT, which the primary study authors confirmed
Liu 2010c	No outcomes of interest; PCOS with or without infertility
Liu 2010d	No outcomes of interest; PCOS with or without infertility
Liu 2011a	Quasi-RCT
Liu 2011b	Quasi-RCT; no outcomes of interest; PCOS with or without infertility
Liu 2012a	No outcomes of interest
Liu 2012b	No outcomes of interest
Liu 2012c	No outcomes of interest
Liu 2013	Quasi-RCT
Liu 2014a	Participants had no wish to conceive
Liu 2014b	Participants had no wish to conceive
Lu 2010	PCOS diagnosis is inconsistent with Rotterdam criteria
Lu 2012	Diagnosis inconsistent with Rotterdam; quasi-RCT
Luo 2010	No outcomes of interest
Luo 2014	Not a RCT
Lv 2007	Intervention with acupuncture but without herbal medicine
Lv 2009	No outcomes of interest; PCOS with or without infertility in this study
Lv 2010	No outcomes of interest; PCOS with or without infertility in this study
Ma 2009b	Quasi-RCT

Study	Reason for exclusion
Ma 2010	No outcomes of interest
Madder 2013	Review
Mao 2003	PCOS diagnosis is inconsistent with Rotterdam criteria; quasi-RCT
Mao 2011a	No outcomes of interest
Mao 2011b	Non-RCT, which the primary authors confirmed
Mei 2010	PCOS with or without infertility
Meng 2011	No outcomes of interest; part of the study was about animals
Miao 2012	PCOS diagnosis is inconsistent with Rotterdam criteria; quasi-RCT; PCOS diagnosis is inconsistent with Rotterdam criteria
Ming-Wei 2011	Non-PCOS participants
Mohammad Hosseinzadeh 2016	Non-CHM intervention
Moradan 2012	Review
Mosalanejad 2015	Not a RCT
Motoo 2014	SR
Musumeci 2006	Review
O'Brien 2010	Unrelated
Pan 2010	No outcomes of interest
Pan 2012	No outcomes of interest
Pastore 2011	Non-CHM intervention
Pazyar 2012	Unrelated study
Pei 2012	PCOS with or without infertility in this study
Peng 2012	No outcomes of interest
Qiao 2012	Quasi-RCT
Qiu 2006	No outcomes of interest; PCOS with or without infertility in this study
Qu 2015	Unrelated
Qv 2011	PCOS with or without infertility
Ran 2007a	No outcomes of interest
Ran 2007b	Case control study

Study	Reason for exclusion
Ran 2008	Case control study
Rao 2012	No PCOS
Rashidi 2013	Non CHM intervention
Ren 2002a	PCOS diagnosis is inconsistent with Rotterdam criteria; quasi-RCT
Ren 2002b	Duplication
Ren 2006	No outcomes of interest; PCOS with or without infertility
Ren 2008	No outcomes of interest; PCOS with or without infertility
Ren 2011	PCOS with or without infertility
Ren 2013	No PCOS
Ren 2014	SR
Ried 2015	SR
Sadrefozalayi 2014	Animals
Salah 2013	Non-CHM intervention
See 2011	SR
Shah 2016	Non-CHM intervention
Shao 2004	PCOS diagnosis is inconsistent with Rotterdam criteria
Shao 2006	PCOS diagnosis is inconsistent with Rotterdam criteria
Shen 2008	PCOS diagnosis is inconsistent with Rotterdam criteria
Shen 2013	Participants were not subfertile
Sheng 2010	No outcomes of interest; PCOS with or without infertility
Shi 2009a	No outcomes of interest; PCOS with or without infertility in this study
Shi 2009b	No outcomes of interest; PCOS with or without infertility in this study
Shi 2010a	PCOS with or without infertility
Shi 2010b	No outcomes of interest
Shi 2011	PCOS with or without infertility; PCOS diagnosis is inconsistent with Rotterdam criteria
Shu 2012	No outcomes of interest
Si 2016	Participants had no wish to conceive
Song 2010	No outcomes of interest; PCOS with or without infertility

Study	Reason for exclusion
Song 2011	PCOS diagnosis is inconsistent with Rotterdam criteria; no outcomes of interest
Stone 2009	Case report
Su 2012	No outcomes of interest; PCOS with or without infertility
Sui 2011	PCOS diagnosis is inconsistent with Rotterdam criteria; no outcomes of interest
Sun 2009a	PCOS diagnosis is inconsistent with Rotterdam criteria; quasi-RCT
Sun 2009b	No outcomes of interest
Sun 2010a	PCOS with or without infertility
Sun 2010b	Intervention with acupuncture but without herbal medicine
Sun 2011	PCOS with or without infertility
Sun 2012	PCOS with or without infertility; quasi-RCT
Sun 2014	No CHM intervention
Tan 2005	Diagnosis inconsistent with Rotterdam; quasi-RCT
Tan 2012	SR
Tang 2012	No outcomes of interest; PCOS with or without infertility
Tao 2003	Diagnosis inconsistent with Rotterdam; PCOS with or without infertility
Tao 2006	Concurrent control study
Tao 2008	No outcomes of interest; PCOS with or without infertility in this study
Tao 2009	No outcomes of interest; PCOS with or without infertility in this study
Tao 2010	No outcomes of interest; PCOS with or without infertility
Tao 2011	No outcomes of interest; PCOS with or without infertility
Ulbricht 2016	SR
Ushiroyama 2001	Diagnosis inconsistent with Rotterdam, participants including PCOS and non-PCOS
Ushiroyama 2006	Diagnosis inconsistent with Rotterdam
Vajda 2013	Not a RCT
van Oppen 2015	Non-PCOS
Wan 2012	No outcomes of interest; PCOS with or without infertility
Wang 2005a	PCOS diagnosis is inconsistent with Rotterdam criteria; quasi-RCT
Wang 2005b	Before-and-after study

Study	Reason for exclusion
Wang 2006a	PCOS with or without infertility in this study
Wang 2006b	Quasi-RCT
Wang 2009	Intervention with acupuncture but without herbal medicine
Wang 2010a	PCOS with or without infertility; quasi-RCT
Wang 2010b	No outcomes of interest; PCOS with or without infertility
Wang 2011a	PCOS with or without infertility
Wang 2011b	Quasi-RCT
Wang 2011c	No outcomes of interest; PCOS with or without infertility
Wang 2011d	No outcomes of interest
Wang 2012a	PCOS with or without infertility
Wang 2012b	PCOS with or without infertility; quasi-RCT
Wang 2013	No CHM intervention
Wang 2016	No CHM intervention
Wei 2008	PCOS diagnosis is inconsistent with Rotterdam criteria
Wei 2011a	No outcomes of interest; PCOS with or without infertility
Wei 2011b	PCOS diagnosis is inconsistent with Rotterdam criteria; PCOS with or without infertility in this study
Wei 2011c	No outcomes of interest; PCOS with or without infertility; diagnosis inconsistent with Rotterdam criteria
Wu 2008	PCOS with or without infertility in this study
Wu 2010a	No outcomes of interest; PCOS with or without infertility in this study
Wu 2010b	PCOS diagnosis is inconsistent with Rotterdam criteria
Wu 2011	PCOS with or without infertility in this study
Wu 2012a	PCOS with or without infertility; no outcomes of interest
Wu 2012b	Non-CHM intervention
Wu 2013a	No CHM intervention
Wu 2013b	Non-PCOS participants
Wuttke 2015	Non-PCOS participants
Xia 2004	No outcomes of interest

Study	Reason for exclusion
Xia 2007	Quasi-RCT
Xia 2011	PCOS diagnosis is inconsistent with Rotterdam criteria
Xiao 2014	No CHM intervention
Xie 2005	Diagnosis inconsistent with Rotterdam criteria
Xie 2010	PCOS with or without infertility
Xie 2012	PCOS with or without infertility
Xiong 2012	No outcomes of interest; PCOS with or without infertility
Xu 2008a	PCOS with or without infertility
Xu 2008b	Diagnosis inconsistent with Rotterdam criteria; quasi-RCT
Xu 2009	PCOS with or without infertility in this study
Xu 2010a	No outcomes of interest
Xu 2010b	PCOS with or without infertility
Xu 2012	No outcomes of interest; PCOS with or without infertility
Xu 2016	Participants had no wish to conceive
Xue 2004	Diagnosis inconsistent with Rotterdam criteria; quasi-RCT
Yan 2003	Duplication
Yan 2005	PCOS diagnosis is inconsistent with Rotterdam criteria
Yan 2012	PCOS diagnosis is inconsistent with Rotterdam criteria; quasi-RCT
Yang 2005	Intervention with acupuncture but without herbal medicine
Yang 2008	Animal study
Yang 2010a	No outcomes of interest; PCOS with or without infertility in this study
Yang 2010b	PCOS diagnosis is inconsistent with Rotterdam criteria; no outcomes of interest; PCOS with or without infertility
Yang 2010c	No outcomes of interest; quasi-RCT
Yang 2011	PCOS diagnosis is inconsistent with Rotterdam criteria; PCOS with or without infertility in this study
Yang 2014a	No CHM intervention
Yang 2014b	Participants had no wish to conceive
Yang 2015	No CHM intervention

Study	Reason for exclusion
Yao 2011	No outcomes of interest; PCOS with or without infertility
Yao 2012a	PCOS with or without infertility
Yao 2012b	Quasi-RCT; no outcomes of interest
Ye 2004	PCOS with or without infertility; diagnosis inconsistent with Rotterdam criteria
Ye 2010	PCOS with or without infertility; no outcomes of interest
Ye 2012a	PCOS diagnosis is inconsistent with Rotterdam criteria; no outcomes of interest; quasi-RCT
Ye 2012b	No outcomes of interest
Ye 2015	Participants had no wish to conceive
Yi 2012	PCOS with or without infertility
Yin 2007	Laparoscopic ovary wedgedshaped resection was used in this study
Yu 2013	With or without subfertility
Yu 2015	Participants had no wish to conceive
Yuan 2011	No outcomes of interest
Yv 2011	No outcomes of interest; PCOS with or without infertility
Zeng 2007	PCOS diagnosis is inconsistent with Rotterdam criteria; quasi-RCT
Zeng 2012	No interventions of interest
Zhang 2003a	No outcomes of interest; PCOS with or without infertility;diagnosis inconsistent with Rotterdam;quasi-RCT
Zhang 2003b	Concurrent control study
Zhang 2007a	No outcomes of interest; PCOS with or without infertility in this study
Zhang 2007b	No outcomes of interest; PCOS with or without infertility in this study
Zhang 2007c	PCOS with or without infertility in this study
Zhang 2007d	Duplication
Zhang 2009	No outcomes of interest; PCOS with or without infertility in this study
Zhang 2010a	PCOS diagnosis is inconsistent with Rotterdam criteria; no outcomes of interest
Zhang 2010b	No outcomes of interest; PCOS with or without infertility
Zhang 2010c	No outcomes of interest
Zhang 2010d	No outcomes of interest; PCOS with or without infertility

Study	Reason for exclusion
Zhang 2010e	PCOS diagnosis is inconsistent with Rotterdam criteria; no outcomes of interest
Zhang 2011a	Not a RCT
Zhang 2011b	No outcomes of interest
Zhang 2011c	No outcomes of interest; PCOS with or without infertility
Zhang 2011d	No outcomes of interest; PCOS with or without infertility
Zhang 2011e	Non-RCT, which the primary authors confirmed
Zhang 2011f	No outcomes of interest; PCOS with or without infertility
Zhang 2012a	No outcomes of interest; PCOS with or without infertility
Zhang 2012b	PCOS with or without infertility; no outcomes of interest
Zhang 2012c	PCOS with or without infertility; no outcomes of interest
Zhang 2012d	No outcomes of interest; quasi-RCT
Zhang 2014a	Participants had no wish to conceive
Zhang 2014b	Participants had no wish to conceive
Zhang 2015a	Participants had no wish to conceive
Zhang 2015b	With or without subfertility
Zhang 2015c	Participants had no wish to conceive
Zhang 2015d	Animals
Zhao 2006a	No outcomes of interest; PCOS with or without infertility in this study
Zhao 2006b	Intervention without herbal medicine
Zhao 2006c	Intervention without herbal medicine
Zhao 2007	PCOS diagnosis is inconsistent with Rotterdam criteria
Zhao 2008a	Quasi-RCT
Zhao 2008b	Duplication
Zhao 2008c	Intervention without herbal medicine
Zhao 2009	Concurrent control study; PCOS with or without infertility
Zhao 2010a	No outcomes of interest
Zhao 2010b	Non-RCT confirmed by primary authors
Zhao 2014	No interventions of interest

Study	Reason for exclusion
Zhao 2016	Participants had no wish to conceive
Zheng 2011	No outcomes of interest; PCOS with or without infertility
Zheng 2011a	PCOS with or without infertility
Zheng 2011b	No outcomes of interest; PCOS with or without infertility
Zheng 2011c	PCOS with or without infertility
Zheng 2014a	Quasi-RCT
Zheng 2014b	Participants had no wish to conceive
Zheng 2015a	Quasi-RCT
Zheng 2015b	Participants had no wish to conceive
Zhi 2012	No outcomes of interest; PCOS with or without infertility
Zhong 2006	PCOS with or without infertility in this study
Zhong 2008	PCOS with or without infertility
Zhong 2009a	No outcomes of interest; PCOS with or without infertility in this study
Zhong 2009b	PCOS with or without infertility in this study
Zhong 2012	Non-RCT, which the primary study authors confirmed
Zhou 1996	PCOS diagnosis is inconsistent with Rotterdam criteria
Zhou 2010a	PCOS with or without infertility
Zhou 2010b	Non-RCT, which the primary authors confirmed
Zhou 2012a	No outcomes of interest
Zhou 2012b	No outcomes of interest
Zhou 2012c	No outcomes of interest; PCOS with or without infertility
Zhou 2012d	PCOS with or without infertility
Zhou 2014a	Quasi-RCT
Zhou 2014b	Quasi-RCT
Zhou 2014c	Quasi-RCT
Zhou 2015a	Quasi-RCT
Zhou 2015b	Quasi-RCT
Zhu 2009	Concurrent control study

Study	Reason for exclusion
Zhu 2012a	PCOS with or without infertility; no outcomes of interest
Zhu 2012b	No outcomes of interest
Zhu 2013a	Participants had no wish to conceive
Zhu 2013b	Participants had no wish to conceive
Zhu 2014	Quasi-RCT
Zhuang 2008	PCOS diagnosis is inconsistent with Rotterdam criteria
Zou 2012	No outcomes of interest; PCOS with or without infertility; PCOS diagnosis is inconsistent with Rotterdam criteria
Zou 2014a	Quasi-RCT
Zou 2014b	Participants had no wish to conceive
Zuo 2011	No outcomes of interest; PCOS with or without infertility

Abbreviations: RCT: randomized controlled trial, CHM: Chinese herbal medicine, PCOS: polycystic ovarian syndrome, SR: systematic review.

Characteristics of ongoing studies *[ordered by study ID]*

[ChiCTR-IOR-16008557](#)

Trial name or title	The Qilingwenshenxiaoanang recipe in the treatment of patients with Polycystic ovary syndrome (PCOS): a multicenter, randomized, double-blind, placebo controlled trial
Methods	Randomized parallel controlled trial
Participants	Women with polycystic ovarian syndrome (PCOS)
Interventions	Qilingwenshenxiaoanang recipe
Outcomes	Ovulation rate
Starting date	Unknown
Contact information	ChiCTR-IOR-16008557
Notes	

[NCT01116167](#)

Trial name or title	Letrozole, berberine, or their combination for anovulatory infertility in women with polycystic ovary syndrome: study design of a double-blind randomized controlled trial
Methods	A multicentre randomized, double-blind trial
Participants	Women with PCOS who desire pregnancy

NCT01116167 (Continued)

Interventions	1. Letrozole and berberine. 2. Letrozole and berberine placebo. 3. Letrozole placebo and berberine.
Outcomes	Live birth rate
Starting date	October 2009
Contact information	ClinicalTrials.gov identifier: NCT01116167
Notes	

Abbreviations:PCOS: polycystic ovarian syndrome.

DATA AND ANALYSES

Comparison 1. CHM versus clomiphene

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Pregnancy rate (per woman)	2	90	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.98 [0.78, 5.06]
1.1 CHM versus clomiphene	1	60	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.44 [0.44, 4.73]
1.2 CHM + LOD versus clomiphene + LOD	1	30	Peto Odds Ratio (Peto, Fixed, 95% CI)	3.35 [0.73, 15.36]
2 Ovulation rate (per woman)	1	30	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.42 [0.19, 10.49]

Analysis 1.1. Comparison 1 CHM versus clomiphene, Outcome 1 Pregnancy rate (per woman).

Study or subgroup	CHM n/N	Clomiphene n/N	Peto Odds Ratio Peto, Fixed, 95% CI	Weight	Peto Odds Ratio Peto, Fixed, 95% CI
1.1.1 CHM versus clomiphene					
Li 2007	8/30	6/30		62.25%	1.44[0.44,4.73]
Subtotal (95% CI)	30	30		62.25%	1.44[0.44,4.73]
Total events: 8 (CHM), 6 (Clomiphene)					
Heterogeneity: Not applicable					
Test for overall effect: Z=0.61(P=0.54)					
1.1.2 CHM + LOD versus clomiphene + LOD					
Ye 2007	14/20	4/10		37.75%	3.35[0.73,15.36]
Subtotal (95% CI)	20	10		37.75%	3.35[0.73,15.36]
Total events: 14 (CHM), 4 (Clomiphene)					
<div> <div>Favours clomiphene</div> <div>0.01 0.1 1 10 100 Favours CHM</div> </div>

Analysis 1.2. Comparison 1 CHM versus clomiphene, Outcome 2 Ovulation rate (per woman).

Comparison 2. CHM + clomiphene versus clomiphene

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Pregnancy rate (per woman)	3	300	Peto Odds Ratio (Peto, Fixed, 95% CI)	2.62 [1.65, 4.14]
1.1 CHM + clomiphene versus clomiphene	2	130	Peto Odds Ratio (Peto, Fixed, 95% CI)	2.45 [1.22, 4.95]
1.2 CHM + CEA + clomiphene versus CEA + clomiphene	1	170	Peto Odds Ratio (Peto, Fixed, 95% CI)	2.75 [1.49, 5.04]

Analysis 2.1. Comparison 2 CHM + clomiphene versus clomiphene, Outcome 1 Pregnancy rate (per woman).

Comparison 3. CHM + follicle aspiration + ovulation induction versus follicle aspiration + ovulation induction

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Pregnancy rate (per woman)	1	44	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.60 [0.46, 5.52]
2 LUFS (adverse events)	1	44	Peto Odds Ratio (Peto, Fixed, 95% CI)	0.60 [0.06, 6.14]
3 OHSS (adverse events)	1	44	Peto Odds Ratio (Peto, Fixed, 95% CI)	0.16 [0.00, 8.19]
4 Multiple pregnancy (adverse events)	1	44	Peto Odds Ratio (Peto, Fixed, 95% CI)	0.60 [0.06, 6.14]

Analysis 3.1. Comparison 3 CHM + follicle aspiration + ovulation induction versus follicle aspiration + ovulation induction, Outcome 1 Pregnancy rate (per woman).

Analysis 3.2. Comparison 3 CHM + follicle aspiration + ovulation induction versus follicle aspiration + ovulation induction, Outcome 2 LUFS (adverse events).

Analysis 3.3. Comparison 3 CHM + follicle aspiration + ovulation induction versus follicle aspiration + ovulation induction, Outcome 3 OHSS (adverse events).

Analysis 3.4. Comparison 3 CHM + follicle aspiration + ovulation induction versus follicle aspiration + ovulation induction, Outcome 4 Multiple pregnancy (adverse events).

Comparison 4. CHM + LOD versus LOD

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Pregnancy rate (per woman)	1	30	Odds Ratio (Peto, Fixed, 95% CI)	3.5 [0.72, 17.09]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
2 Ovulation rate (per woman)	1	30	Odds Ratio (Peto, Fixed, 95% CI)	2.43 [0.39, 15.08]

Analysis 4.1. Comparison 4 CHM + LOD versus LOD, Outcome 1 Pregnancy rate (per woman).

Analysis 4.2. Comparison 4 CHM + LOD versus LOD, Outcome 2 Ovulation rate (per woman).

ADDITIONAL TABLES

Table 1. Contents of the formulations used in included studies

Study	Type of intervention	Formula
Liang 2008	Bushen Huoxue formula	Basic formula: tu si zi 20 g, shu di 10 g, sang ji sheng 20 g, xian ling pi 15 g, bu gu zhi 10 g, huang jing 10 g, zao jiao ci 15 g, tao ren 10 g, shan ci gu 10 g, dan shen 10 g, gan cao 6 g plus huang qi 20 g, shan zha 10 g, fa ban xia 10 g in obese patients plus zhi mu 10 g, huang qin 10 g in hirsutism or acne patients
Li 2007	CHM preparations	Shenqi capsule: tu si zi 15 g, dang shen 20 g, ji xue teng 20 g, fu ling 15 g, dang gui 9 g, dan shen 15 g Ling zhu infusion: yin yang huo 9 g, xian mao 9 g, dan nan xing 9 g, bai zhu 15 g, dang gui 9 g, fa ban xia 9 g, fu ling 15 g

Table 1. Contents of the formulations used in included studies (Continued)

Ye 2007	CHM periodic therapy	<p>Basic formula: kang zhu 10 g, bai zhu 10 g, zhe bei mu 15 g, shi chang pu 15 g, dan shen 10 g, xiang fu 10 g.</p> <ol style="list-style-type: none"> 1. Menstrual phase: basic formula plus tao ren 10 g, san qi 10 g, yi mu cao 15 g, for 3 to 5 days. 2. Late follicular phase: basic formula plus tu si zi 15 g, dang gui 9 g, shi di 10 g, shan yu rou 10 g, fu ling 15 g, for 7 to 10 days. 3. Ovulation phase: basic formula plus lu lu tong 20 g, e zhu 10 g, bei qi 20 g, gui zhi 9 g, for 3 days. 4. Luteinizing phase: basic formula plus tu si zi 15 g, dang gui 10 g, yin yang huo 10 g, rou gui 6 g, for 7 to 10 days.
Ma 2009a	CHM formula	<p>Basic formula in ethinylloestradiol cyproterone acetate therapy phase.</p> <ol style="list-style-type: none"> 1. Yin deficiency of liver and kidney: shu di 30 g, dang gui 15 g, bai shao 15 g, shan yu rou 15 g. 2. Deficiency of spleen and kidney: shu di 30 g, ba ji 30 g, fried bai zhu 30 g, ren shen 15 g, raw huang qi 15 g, shan yu rou 9 g, gou qi zi 6 g, chai hu 1.5 g. <p>Periodic formula</p> <ol style="list-style-type: none"> 1. Gui shao di huang soup at day 5 to 14 of menstrual cycle: dang gui 10 g, bai shao 15 g, shu di 15 g, shan yu rou 10 g, shan yao 10 g, fu ling 15 g, dan pi 10 g, ze xie 15 g. 2. Tao hong si wu soup at day 12 to 16 of menstrual cycle: shu di 10 g, dang gui 15 g, chi shao 15 g, chuan xiong 10 g, tao ren 10 g, hong hua 10 g. 3. Shou tao pellet after ovulation: tu si zi 20 g, sang ji sheng 15 g, e jiao 10 g, xu duan.
Li 2012a	Compound Xuanju capsule	<p>Ingredients: hei ma yi , yin yang huo, gou qi zi, she chuang zi (patent medicine, detailed prescription is not open)</p>

Table 2. The name of CHM in different languages

Pingying name	Latin name	English name
Tu Si Zi	<i>Semen cuscutae dodder</i>	The seed of Chinese dodder
Shu Di	<i>Radix rehmanniae preparata</i>	Prepared rehmannia root
Sang Ji Sheng	<i>Parasitic loranthus</i>	Chinese taxillus twig
Xian Ling Pi	<i>Herba epimedii</i>	Herba epimedii
Bu Gu Zhi	<i>Fructus psoraleae</i>	Malaytea scurfpea fruit
Huang Jing	<i>Rhizoma polygonati</i>	Solomon's seal
Zao Jiao Ci	<i>Spina gleditsiae</i>	Chinese honeylocust spine
Tao Ren	<i>Semen persicae</i>	Peach seed
Shan Ci Gu	<i>Pseudobulbus cremastrae seu pleiones</i>	Pseudobulb of appendiculate cremastra
Dan Shen	<i>Salvia miltiorrhiza</i>	The root of red-rooted salvia
Gan Cao	<i>Radix glycyrrhizae</i>	Licorice roots northwest origin

Table 2. The name of CHM in different languages (Continued)

Huang Qi	<i>Stragalus membranaceus</i>	Membranous milk vetch root
Shan Zha	<i>Crataegus pinnatifida bge</i>	The fruit of a hawthorn
Fa Ban Xia	<i>Rhizoma pinellinae praeparata</i>	Pinellia tuber
Zhi Mu	<i>Rhizoma anemarrhenae</i>	Common anemarrhena rhizome
Huang Qin	<i>Scutellaria baicalensis</i>	Baical skullcap root
Dang Shen	<i>Codonopsis pilosula</i>	Root of hairy asiabell
Ji Xue Teng	<i>Lignum millettiae</i>	Caulis spatholobi
Fu Ling	<i>Tuckahoe</i>	Poria cocos
Dang Gui	<i>Radix angelica</i>	Chinese angelica
Yin Yang Huo	<i>Herba epimedii</i>	Epimedium herb
Xian Mao	<i>Rhizoma curculiginis</i>	Common curculigo rhizome
Dan Nan Xing	<i>Bile arisaema</i>	Arisaema cum bile
Bai Zhu	<i>Atractylodes macrocephala</i>	The rhizome of large headed atractylodes
Cang Zhu	<i>Rhizoma atractylodis</i>	The rhizome of Chinese atractylode
Zhe Bei Mu	<i>Fritillaria thunbergii</i>	Thunberg fritillary bulb
Shi Chang Pu	<i>Rhizoma acori graminei</i>	Grassleaf sweetflag rhizome
Xiang Fu	<i>Rhizoma cyperi</i>	Nutgrass galingale rhizome
San Qi	<i>Pseudo-ginseng</i>	Panax notoginseng
Yi Mu Cao	<i>Leenurus heterophyllus</i>	Motherwort
Shan Yu Rou	<i>Fructus corni</i>	Common macrocarpium fruit
Lu Lu Tong	<i>Liquidambar formosana hance</i>	Beautiful sweetgum fruit
E Zhu	<i>Curcuma zedoary</i>	Zedoray rhizome
Bei Qi	<i>Radix astragali</i>	Northeast milkvetch root
Gui Zhi	<i>Ramulus cinnamomi</i>	Cassia twig
Rou Gui	<i>Cinnamomum cassia</i>	Chinese cinnamon
Bai Shao	<i>Radices paeoniae alba</i>	Root of herbaceous peony
Ba Ji	<i>Radix morindae officinalis</i>	Medicinal Indianmulberry root
Ren Shen	<i>Panax, gen-seng</i>	Ginseng

Table 2. The name of CHM in different languages (Continued)

Shan Yao	<i>Rhizoma dioscoreae</i>	Common yam rhizome
Dan Pi	<i>Paeoniasuffruticosa</i>	The root bark of the peony tree
Ze Xie	<i>Rhizoma alismatic</i>	Oriental waterplantain rhizome
Chi Shao	<i>Radix paeoniae rubrathe</i>	Root of common peony
Chuan Xiong	<i>Ligusticum wallichii</i>	The rhizome of chuanxiong
Hong Hua	<i>Carthamus tinctorious</i>	Red flower
E Jiao	<i>Colla dorii asini</i>	Donkey hide gelatin
Xu Duan	<i>Radix dipsaci</i>	Teasel root
Hei Ma Yi	<i>Formicae populus infirmus quae nigra</i>	Black Ants
Yin Yang Huo	<i>Herba epimedii</i>	Epimedium Herb
Gou Qi Zi	<i>Fructus lycii</i>	Barbary Wolfberry Fruit
She Chuang Zi	<i>Fructus cnidii</i>	Common Cnidium Fruit

APPENDICES

Appendix 1. CGF search string

PROCITE platform

From inception until 9 June 2016

Keywords CONTAINS "Polycystic Ovary Syndrome" or "polycystic ovary morphology" or "PCOS" or "anovulation" or Title CONTAINS "PCOS" or "Polycystic Ovary Syndrome" or "polycystic ovary morphology" or "anovulation"

AND

Keywords CONTAINS "Chinese" or "Chinese herbal medicine" or "Chinese drugs" or "Chinese herbal remedy" or "Chinese traditional medicine" or "traditional Chinese medicine" or "traditional medicine" or "herbal preparations" or "herbal remedy" or "herbal supplement" or "herbal supplements" or Title CONTAINS "Chinese" or "Chinese herbal medicine" or "Chinese drugs" or "Chinese herbal remedy" or "Chinese traditional medicine" or "traditional Chinese medicine" or "traditional medicine" or "herbal preparations" or "herbal remedy" or "herbal supplement" or "herbal supplements" (16 hits)

Appendix 2. CENTRAL CRSO search strategy

Web platform

From inception until 9 June 2016

#1 MESH DESCRIPTOR Polycystic Ovary Syndrome EXPLODE ALL TREES (857)
 #2 (Polycystic Ovar*):TI,AB,KY (1599)
 #3 (stein leventhal):TI,AB,KY (3)
 #4 PCOS:TI,AB,KY (1212)
 #5 hirsut*:TI,AB,KY (451)
 #6 anovulat*:TI,AB,KY (435)
 #7 #1 OR #2 OR #3 OR #4 OR #5 OR #6 (2265)
 #8 MESH DESCRIPTOR Drugs, Chinese Herbal EXPLODE ALL TREES (2809)

#9 MESH DESCRIPTOR Medicine, Chinese Traditional EXPLODE ALL TREES (872)
 #10 MESH DESCRIPTOR Medicine, East Asian Traditional EXPLODE ALL TREES (952)
 #11 (Chinese adj5 Tradition*):TI,AB,KY (1004)
 #12 (Oriental medicine*):TI,AB,KY (26)
 #13 herb*:TI,AB,KY (5850)
 #14 TCM:TI,AB,KY (1515)
 #15 (traditional adj5 medicine*):TI,AB,KY (2850)
 #16 (Chinese adj5 medicine*):TI,AB,KY (4043)
 #17 #8 OR #9 OR #10 OR #11 OR #12 OR #13 OR #14 OR #15 OR #16 (9987)
 #18 #7 AND #17 (57)

Appendix 3. MEDLINE search strategy

MEDLINE® Epub Ahead of Print, In Process & Other Non-Indexed Citations, Ovid MEDLINE® Daily, and Ovid MEDLINE®

OVID platform

from inception until 9 June 2016

1 exp Polycystic Ovary Syndrome/ (11497)

2 Polycystic Ovar\$.tw. (12565)

3 stein leventhal.tw. (607)

4 PCOS.tw. (7822)

5 hirsut\$.tw. (7717)

6 anovulat\$.tw. (4937)

7 or/1-6 (24480)

8 exp Drugs, Chinese Herbal/ or exp Medicine, Chinese Traditional/ or exp Medicine, Oriental Traditional/ (46673)

9 (Chinese adj5 Traditional).tw. (18109)

10 Oriental medicine\$.tw. (830)

11 herb\$.tw. (74874)

12 TCM.tw. (6589)

13 (traditional adj5 medicine\$).tw. (25786)

14 (Chinese adj5 medicine\$).tw. (20908)

15 or/8-14 (128194)

16 randomized controlled trial.pt. (420793)

17 controlled clinical trial.pt. (91006)

18 randomized.ab. (358742)

19 placebo.tw. (178518)

20 clinical trials as topic.sh. (177498)

21 randomly.ab. (256600)

22 trial.ti. (156177)

23 (crossover or cross-over or cross over).tw. (69506)

24 or/16-23 (1065847)

25 (animals not (humans and animals)).sh. (4229204)

26 24 not 25 (981472)

27 7 and 15 and 26 (53)

Appendix 4. EMBASE search strategy

Web platform

From inception until 9 June 2016

1 exp ovary polycystic disease/ or exp stein leventhal syndrome/ (19944)

2 (polycystic adj5 ovar\$).tw. (16641)

3 stein leventhal.tw. (547)

4 PCOS.tw. (11206)

5 hirsut\$.tw. (8657)

6 anovulat\$.tw. (5539)

7 or/1-6 (32581)

8 exp Chinese Drug/ or exp Herbal Medicine/ or exp Chinese Medicine/ or exp Chinese Herb/ (52208)

9 (Chinese Drug\$ or Herbal Medicine\$ or Chinese Medicine or Chinese Herb\$).tw. (38642)

10 Oriental medicine\$.tw. (1196)

11 traditional.tw. (269441)

12 TCM.tw. (9137)

13 or/8-12 (314777)

14 7 and 13 (497)

15 Clinical Trial/ (858094)

16 Randomized Controlled Trial/ (404847)

17 exp randomization/ (70570)

18 Single Blind Procedure/ (22197)

19 Double Blind Procedure/ (128704)

20 Crossover Procedure/ (47202)

21 Placebo/ (275451)

22 Randomi?ed controlled trial\$.tw. (136119)

23 Rct.tw. (20323)

24 random allocation.tw. (1528)

Chinese herbal medicine for subfertile women with polycystic ovarian syndrome (Review)

Copyright © 2016 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

25 randomly allocated.tw. (24807)

26 allocated randomly.tw. (2113)

27 (allocated adj2 random).tw. (757)

28 Single blind\$.tw. (17385)

29 Double blind\$.tw. (162039)

30 ((treble or triple) adj blind\$.tw. (555)

31 placebo\$.tw. (232908)

32 prospective study/ (334857)

33 or/15-32 (1574988)

34 case study/ (37932)

35 case report.tw. (305862)

36 abstract report/ or letter/ (961247)

37 or/34-36 (1298029)

38 33 not 37 (1533942)

39 14 and 38 (121)

Appendix 5. PsycINFO search strategy

Web platform

From inception until 9 June 2016

1 Polycystic Ovar\$.tw. (225)

2 stein leventhal.tw. (2)

3 PCOS.tw. (132)

4 hirsut\$.tw. (107)

5 or/1-4 (320)

6 exp "medicinal herbs and plants"/ or exp "plants (botanical)"/ (1996)

7 Chinese Herb\$.tw. (136)

8 (Chinese adj5 Traditional).tw. (940)

9 Oriental medicine\$.tw. (47)

10 (herbal adj5 medicine\$.tw. (389)

11 herb\$.tw. (4889)

12 or/6-11 (7150)

13 5 and 12 (2)

Appendix 6. CINAHL search strategy

EBSCO platform

from inception until 9 June 2016

#	Query	Results
S28	S15 AND S27	46

(Continued)

S27	S16 OR S17 OR S18 OR S19 OR S20 OR S21 OR S22 OR S23 OR S24 OR S25 OR S26	1,053,818
S26	TX allocat* random*	5,059
S25	(MH "Quantitative Studies")	14,585
S24	(MH "Placebos")	9,730
S23	TX placebo*	38,559
S22	TX random* allocat*	5,059
S21	(MH "Random Assignment")	41,063
S20	TX randomi* control* trial*	105,940
S19	TX ((singl* n1 blind*) or (singl* n1 mask*)) or TX ((doubl* n1 blind*) or (doubl* n1 mask*)) or TX ((tripl* n1 blind*) or (tripl* n1 mask*)) or TX ((trebl* n1 blind*) or (trebl* n1 mask*))	835,812
S18	TX clinic* n1 trial*	186,430
S17	PT Clinical trial	79,707
S16	(MH "Clinical Trials+")	198,930
S15	S7 AND S14	126
S14	S8 OR S9 OR S10 OR S11 OR S12 OR S13	66,100
S13	TX traditional N2 medicine*	18,439
S12	TX CHM	274
S11	TX TCM	2,035
S10	TX oriental	4,154
S9	TX chinese	51,134
S8	(MM "Medicine, Chinese Traditional") OR (MM "Drugs, Chinese Herbal") OR (MM "Medicine, Herbal") OR (MM "Medicine, Oriental Traditional")	12,296
S7	S1 OR S2 OR S3 OR S4 OR S5 OR S6	3,528
S6	TX anovulat*	396
S5	TX hirsut*	513
S4	TX PCOS	1,591
S3	TX stein leventhal	10
S2	TX Polycystic Ovar*	2,226
S1	(MM "Polycystic Ovary Syndrome")	1,381

(Continued)

Appendix 7. AMED search strategy

OVID platform

From inception until 9 June 2016

1 exp Ovarian disease/ (194)

2 (polycystic adj5 ovar\$).tw. (60)

3 stein leventhal.tw. (1)

4 PCOS.tw. (28)

5 anovulat\$.tw. (22)

6 hirsut\$.tw. (66)

7 or/1-5 (225)

8 exp Traditional medicine chinese/ or exp Drugs chinese herbal/ (7255)

9 (Chinese Drug\$ or Herbal Medicine\$ or Chinese Medicine\$ or Chinese Herb\$).tw. (4909)

10 traditional.tw. (12454)

11 or/8-10 (14994)

12 7 and 11 (28)

Appendix 8. CNKI search strategy

From inception until 9 June 2016

1.polycystic ovary syndrome (12274)

2.polycystic ovary (11405)

3.1~2/or (12980)

4.Chinese herbal medicine (56602)

5.herbal medicine (6671)

6.traditional medicine (13798)

7.traditional Chinese medicine (420522)

8.traditional Chinese medicine combined with western medicine (149311)

9.4~8/or (594003)

10.random* (1492056)

11.3 and 9 and 10 (573)

Appendix 9. Wanfang search strategy

From inception until 9 June 2016

1.polycystic ovary syndrome (11011)

Chinese herbal medicine for subfertile women with polycystic ovarian syndrome (Review)

Copyright © 2016 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

2.polycystic ovary (11862)
3.1~2/or (11862)
4.Chinese herbal medicine (26898)
5.herbal medicine (32132)
6.traditional medicine (8347)
7.traditional Chinese medicine (408902)
8.traditional Chinese medicine combined with western medicine (126458)
9.4~8/or (755982)
10.random* (1410377)
11.3 and 9 and 10 (665)

Appendix 10. VIP search strategy

From inception until 9 June 2016

1.polycystic ovary syndrome (8587)
2.polycystic ovary (9001)
3.1~2/or (9001)
4.Chinese herbal medicine (10559)
5.herbal medicine (12013)
6.traditional medicine (2874)
7.traditional Chinese medicine (167976)
8.traditional Chinese medicine combined with western medicine (118451)
9.4~8/or (289985)
10.random* (23253)
11.3 and 9 and 10 (5)

WHAT'S NEW

Date	Event	Description
10 July 2016	New search has been performed	The updated search found two ongoing studies (ChiCTR-IOR-16008557 ; NCT01116167), and one new study (Li 2012a).
10 July 2016	New citation required but conclusions have not changed	There is insufficient evidence for the conclusions of this review to be changed.

HISTORY

Protocol first published: Issue 1, 2009
Review first published: Issue 9, 2010

Date	Event	Description
20 September 2010	Amended	Contact details updated.
25 September 2008	Amended	Title changed from 'Chinese herbal medicine for polycystic ovarian syndrome' to 'Chinese herbal medicine for subfertile women with polycystic ovary syndrome'; objectives were also changed.
22 September 2008	Amended	Title changed from 'Chinese herbal medicine for the management of polycystic ovarian syndrome' to 'Chinese herbal medicine for polycystic ovarian syndrome'.

CONTRIBUTIONS OF AUTHORS

Kunyan Zhou updated the review, searched for trials, entered data entry into RevMan ([RevMan 2014](#)), extracted data, selected trials for inclusion or exclusion, and contacted the primary study authors.

Jing Zhang drafted the protocol and original review; entered data into RevMan ([RevMan 2014](#)), and screened trials for inclusion or exclusion.

Liangzhi Xu extracted data.

Taixiang Wu screened trials for inclusion or exclusion.

Danfor Lim revised and corrected the text.

DECLARATIONS OF INTEREST

Kuanyan Zhou has no known conflicts of interest.

Jing Zhan has no known conflicts of interest.

Liangzhi Xu has no known conflicts of interest.

Taixiang Wu has no known conflicts of interest.

Chi Eung Danfor Lim has no known conflicts of interest.

SOURCES OF SUPPORT

Internal sources

- West China Second University Hospital, Sichuan Universtiy, China.
- Key Laboratory of Birth Defects and Related Diseases of Women and Children (Sichuan University), Ministry of Education, China.
- Chinese Cochrane Center, West China Hospital, Sichuan University, China.
- National Natural Science Foundation of China (81270665), China.
- National Natural Science Foundation of China (41473097), China.

External sources

- None, Other.

DIFFERENCES BETWEEN PROTOCOL AND REVIEW

We searched more electronic databases in this review update than we listed in the original protocol ([Zhang 2009](#)).

We added CHM+clomiphene versus clomiphene as a comparison in last review ([Zhang 2010](#)) that was not listed in the protocol ([Zhang 2009](#)). In this 2016 updated review we deleted this comparison as it was a duplicate.

NOTES

None.

INDEX TERMS

Medical Subject Headings (MeSH)

Clomiphene [therapeutic use]; Cyproterone Acetate [therapeutic use]; Drugs, Chinese Herbal [*therapeutic use]; Ethinyl Estradiol [therapeutic use]; Fertility Agents, Female [therapeutic use]; Infertility [*drug therapy] [etiology]; Laparoscopy; Ovulation Induction [methods]; Polycystic Ovary Syndrome [complications] [*therapy]; Pregnancy Rate; Randomized Controlled Trials as Topic; Suction

MeSH check words

Adult; Female; Humans; Pregnancy