Published in final edited form as:

Virus Res. 2019 May ; 265: 68–73. doi:10.1016/j.virusres.2019.03.002.

To assemble or not to assemble: The changing rules of pneumovirus transmission

Nicolás Cifuentes-Muñoz^{1,2} and Rebecca Ellis Dutch^{1,#}

¹Department of Molecular and Cellular Biochemistry, University of Kentucky College of Medicine, Lexington, KY 40536, USA.

²Instituto de Ciencias Biomédicas, Facultad de Ciencias de la Salud, Universidad Autónoma de Chile, El Llano Subercaseaux 2801, San Miguel, Santiago 8910060, Chile.

Abstract

Pneumoviruses represent a major public health burden across the world. Respiratory syncytial virus (RSV) and human metapneumovirus (HMPV), two of the most recognizable pediatric infectious agents, belong to this family. These viruses are enveloped with a non-segmented negative-sense RNA genome, and their replication occurs in specialized cytosolic organelles named inclusion bodies (IB). The critical role of IBs in replication of pneumoviruses has begun to be elucidated, and our current understanding suggests they are highly dynamic structures. From IBs, newly synthesized nucleocapsids are transported to assembly sites, potentially via the actin cytoskeleton, to be incorporated into nascent virions. Released virions, which generally contain one genome, can then diffuse in the extracellular environment to target new cells and reinitiate the process of infection. This is a challenging business for virions, which must face several risks including the extracellular immune responses. In addition, several recent studies suggest that successful infection may be achieved more rapidly by multiple, rather than single, genomic copies being deposited into a target cell. Interestingly, recent data indicate that pneumoviruses have several mechanisms that permit their transmission en bloc, i.e. transmission of multiple genomes at the same time. These mechanisms include the well-studied syncytia formation as well as the newly described formation of long actin-based intercellular extensions. These not only permit en bloc viral transmission, but also bypass assembly of complete virions. In this review we describe several aspects of en bloc viral transmission and how these mechanisms are reshaping our understanding of pneumovirus replication, assembly and spread.

Keywords

Pneumoviruses; spread; en bloc; inclusion bodies; nucleocapsids

[#]Author to whom correspondence should be addressed; rdut2@uky.edu; Tel.: +1-859-323-1795; Fax: +1-859-257-2283.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1. Introduction

Pneumoviruses are a family of enveloped, non-segmented negative-sense RNA viruses (nsNSVs) (1). Two of the most significant pediatric respiratory viruses, human metapneumovirus (HMPV) and respiratory syncytial virus (RSV) belong to this family. Both viruses can cause upper and lower respiratory tract infections, which can lead to bronchiolitis and pneumonia. According to data surveillance obtained for children under the age of five, RSV infections account for >57,000 hospitalizations and over 2 million outpatient visits per year in the US (2). For the same pediatric population, HMPV associated infections are responsible for an estimated 1 in 1,000 hospitalizations, similar to those associated with influenza (3). However, no FDA approved vaccines are available against RSV or HMPV. A humanized monoclonal antibody that targets the RSV fusion protein (F), palivizumab, is available for use in premature or immunocompromised high-risk infant populations. No effective antivirals have been reported against HMPV, and the use *in vitro* of ribavirin, heparin, and the sulfated sialyl lipid NMSO3 have had only limited success (4, 5).

As with other enveloped viruses, the process of budding and release of pneumovirus virions involves the coalescence of viral components at specific locations on cellular membranes, followed by membrane deformation and pinching-off of a viral particle (6). Critical roles for the viral matrix protein (M) and the glycoproteins have been described in these processes, which implicate a high energetic cost. Though the release of particles into the extracellular environment is required for host-to-host viral spread, as a mechanism of spread within the host it does present several disadvantages. Examples of these disadvantages are the low stability of the particles in the extracellular environment and the susceptibility of virions to be targeted by immune cells. While the model of target cell infection resulting from primarily single particle entry is prevalent, this concept has been challenged by novel reports suggesting that infection of target cells may often be the result of infection by multiple particles. In this context, pneumoviruses have developed alternative mechanisms of spread that potentially allow the virus to directly deposit multiple copies of the viral genome into target cells simultaneously. This review discusses some of the mechanisms for transport of multiple genome copies and their implications for virus spread.

2. Pneumovirus genome organization and replication

2.1 Pneumovirus nucleocapsids

Pneumoviruses possess a non-segmented, negative-sense single-stranded RNA genome that encodes between 9-11 proteins. Respiratory syncytial virus genomic RNA (vRNA) is about 15.2 Kb in size, while for HMPV a slightly smaller RNA molecule of about 13.2 Kb serves as the genome (Figure 1a). Pneumovirus vRNA is entirely coated by the nucleoprotein (N), giving rise to long, flexible, left-handed helical nucleocapsids (7, 8). Each nucleoprotein subunit binds to seven bases of the RNA (9) (10), in contrast to the paramyxovirus nucleoprotein subunits which bind to six RNA bases (8). Pneumoviral nucleocapsids are additionally decorated by copies of the viral phosphoprotein (P), which interacts with N and serves as a polymerase cofactor (11). The viral phosphoprotein also recruits the large RNA-dependent RNA-polymerase (L) and the M2-1 protein onto the nucleocapsids. It is widely accepted that viral nucleocapsids not only protect the vRNA from degradation, but also serve

as the template for viral replication and transcription (12). Sequences located at the 3' (leader, le) and 5' (trailer, tr) ends of the vRNA contain promoter signals that are important for viral genome replication and transcription (13). Replication of the vRNA starts after the viral polymerase is recruited to the +1 site at the 3' le region (14) and occurs through synthesis of a full-length positive-sense copy of the genome, also known as antigenome (Figure 1a). The antigenome and genome are simultaneously coated by the nucleoprotein (12). Transcription of the genome starts at the +3 site in the promoter of the 3' le region (14) and proceeds through a sequential termination-reinitiation mechanism (13). This mechanism is controlled by the presence of gene start (GS) and gene end (GE) signals for each individual gene, which ultimately results in the synthesis of sub-genomic, capped and poly-adenylated viral mRNAs (Figure 1a)(12). Unlike the vRNA or the antigenome, viral mRNAs are not coated by N, but a recent study suggested they might contain copies of the M2-1 protein (15).

2.2 Inclusion bodies: The viral genome factories

Reference to inclusion body (IB) formation upon infection of Vero cells by RSV was reported early in 1970 by using electron microscopy (16). Inclusion bodies induced by RSV are described as cytosolic, compact, membraneless organelles that incorporate replicative viral proteins such as N, P, and L, while excluding others such as M or the glycoproteins G and F (16-18). The minimal components necessary for formation of RSV IBs were defined to be the RSV N and P proteins (17). More recently, the HMPV N and P proteins were also shown to be the minimal components required for assembly of inclusion bodies (19). Though initially IBs were believed to represent a station for accumulation of misfolded proteins, several recent reports have highlighted their role as the sites of active viral genome replication (Figure 1b). Combining metabolic labelling with fluorescence *in situ* hybridization (FISH) and high resolution microscopy, viral genome replication was shown to occur within RSV IBs (15). Different compartments within IBs, denominated IB-associated granules (IBAGs) were shown to contain newly synthesized RNA, most likely viral mRNAs (15). HMPV IBs were also shown to contain vRNA as well as viral mRNAs, plus the N and P replicative proteins, while excluding assembly proteins such as M or F (20). Interestingly, HMPV IBs were shown to result from the coalescence of multiple smaller replicative spots during the early steps of HMPV infection, revealing an intrinsic plasticity for pneumovirus IBs (Figure 1b)(20). Remarkably, formation of IBs represents a replication strategy used beyond the *Pneumoviridae*. Although quite different in their host range and symptoms, the role of IBs in replication of the viral genome has been recently reported for rhabdoviruses (21, 22), filoviruses (23-25) and for some paramyxoviruses (26, 27).

2.3 Transport of viral nucleocapsids inside the cells

Inclusion bodies are a station from where newly synthesized nucleocapsids must be transported to viral assembly sites at the plasma membrane. Though the exact mechanism of how this occurs has remained elusive for pneumoviruses, several reports point out an important role of the actin cytoskeleton in this process (Figure 1b). In early times post-infection, when large inclusion bodies have not formed, incoming -as well as newly synthesized- HMPV nucleocapsids coalesce in a process that is dependent on actin polymerization (20). Coalescence of nucleocapsids into larger IBs was shown to be

important for efficient replication and transcription of the HMPV genome, likely due to the concentration of replicative components within these organelles. At later times post infection, coalescence of large IBs still occurs (Figure 1b)(20). Coalescence of IBs has been additionally observed for other nsNSVs including rabies virus (28), Zaire Ebola virus (ZEBOV)(25), and the paramyxovirus parainfluenza virus type 3 (PIV3)(27). Interestingly, coalescence of PIV3 IBs was shown to be dependent on acetylated α -tubulin, through a direct interaction with the N-P complex (27). Interaction of the RSV N-P complex was not found with acetylated α -tubulin (27), most likely because the RSV N-P complex interacts mainly with components of the actin cytoskeleton (29). Purified RSV particles contain actin, but exclude other cytoskeletal components such as vimentin, cytokeratin and tubulin (30, 31). Human metapneumovirus purified particles also contain actin (32), and a major role of the actin cytoskeleton, but not microtubules, in the replication and assembly of both HMPV and RSV has been well documented (20, 29-34). In this context, pneumovirus nucleocapsid transport may operate in a similar way to filovirus nucleocapsid transport. Marburg virus (MARV) and EBOV nucleocapsid cytoplasmic transport was shown to be dependent on actin polymerization and the actin nucleating complex Arp2/3, but independent of microtubules (35, 36). Transport of nucleocapsids over long distances from IBs to the tip of filopodia, the budding sites for filoviruses, was detected (35, 36). Interestingly, HMPV IBs have been observed far from their usual perinuclear localization, which suggests their movement as an independent organelle over long distances (32). The mechanisms and cellular components involved in their movement remain to be addressed.

3. En bloc transmission of pneumoviruses

Recent evidence suggests that novel mechanisms of direct cell-to-cell spread might be an important avenue for pneumovirus transmission (20, 37). These mechanisms, and others described previously, would allow pneumovirus spread to operate at least partially en bloc, defined here as spread of multiple genomes occurring at the same time. This notion challenges the more accepted concept of one infectious particle infecting one cell at the time. Based on evidence primarily obtained for pneumoviruses but also for other related viruses, we discuss several mechanisms for how en bloc viral transmission could occur.

3.1 Cell-to-cell fusion

Cell to cell fusion results in the appearance of large multinucleated cells, named syncytia. Syncytia formation has been reported for HMPV and RSV *in vitro* (38), but their presence *in vivo* is less clear (39). However, lung autopsy specimens have revealed the presence of large syncytia from both HMPV and RSV patients, which strongly argues for their formation *in vivo* (40, 41). Mechanistically, syncytia formation is facilitated by the accumulation of the viral fusion protein (F) in the plasma membrane of infected cells. F binds to the membranes of adjacent cells, and through the aperture of a fusion pore that enlarges it completely fuses both membranes. As a result, cytosolic content mixing occurs, and consequently infection is spread directly from cell-to-cell (42). Once an infected cell has completely merged with an uninfected cell, a massive transmission of nucleocapsids and IBs would have occurred, therefore rapidly spreading infection (Figure 2a). Interestingly, an RSV F with a hyperfusogenic phenotype has been associated with increased pathogenesis in mice (43).

Future studies will elucidate other essential aspects of this mechanism of en bloc viral transmission for pneumoviruses and how they impact viral pathogenesis.

3.2 Intercellular extensions

Recent reports have described a novel route of direct cell-to-cell spread by pneumoviruses that involves the formation of filamentous structures that extend towards distant cells. Respiratory syncytial virus induces formation of filopodia in lung epithelial A549 cells, which facilitate viral spread (37). Induction of filopodia was shown to be dependent on the RSV F protein and the cellular actin-related protein 2 (Arp2), an important actin-nucleation factor. At the surface of filopodia, clusters of filamentous structures most likely corresponding to RSV virions were shown to contact neighboring uninfected cells. This phenotype suggests a mechanism of spread through which multiple virions can be shuttled from one cell to another without the need for release. Additionally, formation of filopodia conferred increased motility to RSV infected A549 cells, which was shown to also facilitate cell-to-cell spread (37). Formation of filopodia was not as prominent in HMPV-infected A549 cells, but human bronchial epithelial BEAS2B cells infected with HMPV have been shown to result in conspicuous formation of intercellular extensions (32). Cellular factors including F-actin and the Rho GTPases Cdc42 and Rac1 were shown to be important for the induction of HMPV intercellular extensions. Similar to RSV-induced filopodia, the surface of HMPV-induced extensions is covered by clusters of branching filaments that contain viral proteins and RNA, which we hypothesize correspond to filamentous virions. Hence, the potential massive shuttle of HMPV virions directly from cell-to-cell arises as an efficient mechanism of spread (32). Interestingly, HMPV cell-to-cell spread was shown to be at least partially independent of the appropriate attachment factors and of the presence of neutralizing antibodies. This suggests that cell-to-cell spread might occur through a direct connection between the cytoplasm of an HMPV infected cell and a target cell. In support of this, vRNA as well as IB-like structures were shown to be present in the HMPV-induced intercellular extensions, potentially being transported towards neighboring cells (Figure 2b) (32). Transmission of nucleocapsids through long intercellular structures has been reported for influenza virus in lung epithelial cells (44, 45). Though the open-ended nature of pneumovirus intercellular extensions remains to be addressed, this mechanism of en bloc viral transmission represents an advantageous way to rapidly spread infection.

3.3 Polyploid virions

Pneumovirus purified particles display a wide size variability, ranging from 100 nm up to 1,000 nm, and from 150 nm up to 600 nm, for RSV and HMPV, respectively. Not only is the pneumovirus particle size quite heterogeneous, but also their shape, and usually two major forms, pleomorphic and filamentous, can be observed (46-49). Some of the filamentous RSV particles have been observed to grow up to 10 μ m in length (48). In the case of HMPV, which grows predominantly cell-associated *in vitro*, filaments budding from infected cells, potentially corresponding to viral particles, are usually above 2 μ m in length (32, 50). Though the number of pneumovirus nucleocapsids packaged within filamentous virions could not be determined from the above mentioned studies, it is possible, based on the size of the particles as well as their heterogeneity, that polyploid virions are present. Polyploid virions have been described for paramyxoviruses including Newcastle disease virus (NDV),

measles virus and Sendai virus (SeV) (51-53). More important, the presence of multiple genomes -up to 6 nucleocapsids in the case of SeV- within one viral particle has been shown not to be detrimental for viral infection (51). In the context of mass transport of genomes, the presence of multiple nucleocapsids within particles fits with the proposal that infection is more successfully initiated by multiple particles rather than by single particles (Figure 2c).

3.4 En bloc transmission of pneumoviruses: A double edged sword

An important feature during replication of different RNA viruses *in vitro* is the synthesis of truncated forms of the vRNA denominated defective viral genomes (DVGs), initially named defective viral particles, as they can interfere with viral replication (54). Defective viral genomes exist in different conformations and, as they are truncated forms of the genome, in order to propagate they need a helper virus (reviewed in (55)). Early studies with measles virus DVGs demonstrated that their presence correlated with the establishment of persistent infections in cell lines (56). Moreover, other nsNSV DVGs have been described as strong inducers of the type-I IFN response (57). When passaged at high titers in cell lines, RSV as well as HMPV infections generate defective interfering genomes (58, 59). Similar to measles, the presence of RSV DVGs has been associated with the establishment of viral persistence in cell lines (60). More recently, it was shown that RSV DVGs can stimulate the expression of antiviral genes in mice, leading to decreased viral titers and symptoms (61). A direct correlation was found in human lung tissues between the amount of DVGs and the quality of the antiviral response mounted, which suggests DVGs have a role *in vivo* (61). Considering the previous mechanisms described for en bloc pneumoviral transmission, and the large presence of DVGs upon infection, a valid question that arises is whether these mechanisms exist as a viral strategy for rapid spread or as a cellular antiviral strategy to counteract the infection. Future work will elucidate how both possibilities might operate simultaneously.

4. En bloc transmission of Pneumoviruses: Implications for viral dynamics

The mechanisms of en bloc viral transmission described here permit the potential transfer of many viral genomes into a target cell at the same time. In the context of pneumoviruses, this might represent a crucial mechanism for successful infection. Pneumoviruses are within the group of nsRNA viruses, and estimations indicate that mutation frequencies for these viruses can range from 10^{-5} to 10^{-6} per nucleotide per infectious cycle (62). In the absence of proofreading mechanisms, the high mutation frequency for these viruses results in multiple variations of vRNA populations, which are usually referred to as quasispecies (63). The infection of a cell with viral quasispecies could result in a complementation of a defect in one genome by the presence of additional genomes. This cooperativity would be less likely in the context of an individual viral particle carrying an attenuated genome. Additionally, the massive transfer of genomes into a target cell ensures a higher initial concentration of viral proteins and RNA, which we have previously shown to rapidly coalesce to form large IBs (20). A higher concentration of these replicative components would facilitate infection and would allow the virus to bypass the critical first hours of infection, potentially allowing high levels of viral replication before antiviral responses are in place. Though some of these

hypotheses remain merely speculative for pneumoviruses, a deeper understanding of the virus spread mechanisms will help us improve the way we approach the design of antiviral strategies.

Acknowledgements

This work was supported in part by NIH grants AI051517 and AI140758 to R.E.D. as well as Fondecyt Inicio grant 11180269 to N.C.M.

References

1. Afonso CL, Amarasinghe GK, Banyai K, Bao Y, Basler CF, Bavari S, Bejerman N, Blasdel KR, Briand FX, Briese T, Bukreyev A, Calisher CH, Chandran K, Cheng J, Clawson AN, Collins PL, Dietzgen RG, Dolnik O, Domier LL, Durrwald R, Dye JM, Easton AJ, Ebihara H, Farkas SL, Freitas-Astua J, Formenty P, Fouchier RA, Fu Y, Ghedin E, Goodin MM, Hewson R, Horie M, Hyndman TH, Jiang D, Kitajima EW, Kobinger GP, Kondo H, Kurath G, Lamb RA, Lenardon S, Leroy EM, Li CX, Lin XD, Liu L, Longdon B, Marton S, Maisner A, Muhlberger E, Netesov SV, Nowotny N, Patterson JL, Payne SL, Paweska JT, Randall RE, Rima BK, Rota P, Rubbenstroth D, Schwemmle M, Shi M, Smither SJ, Stenglein MD, Stone DM, Takada A, Terregino C, Tesh RB, Tian JH, Tomonaga K, Tordo N, Towner JS, Vasilakis N, Verbeek M, Volchkov VE, Wahl-Jensen V, Walsh JA, Walker PJ, Wang D, Wang LF, Wetzel T, Whitfield AE, Xie JT, Yuen KY, Zhang YZ, Kuhn JH. 2016 Taxonomy of the order Mononegavirales: update 2016. *Archives of virology* 161:2351–2360. [PubMed: 27216929]
2. Hall CB, Weinberg GA, Iwane MK, Blumkin AK, Edwards KM, Staat MA, Auinger P, Griffin MR, Poehling KA, Erdman D, Grijalva CG, Zhu Y, Szilagyi P. 2009 The burden of respiratory syncytial virus infection in young children. *The New England journal of medicine* 360:588–598. [PubMed: 19196675]
3. Edwards KM, Zhu Y, Griffin MR, Weinberg GA, Hall CB, Szilagyi PG, Staat MA, Iwane M, Prill MM, Williams JV, New Vaccine Surveillance N. 2013 Burden of human metapneumovirus infection in young children. *The New England journal of medicine* 368:633–643. [PubMed: 23406028]
4. Wyde PR, Chetty SN, Jewell AM, Boivin G, Piedra PA. 2003 Comparison of the inhibition of human metapneumovirus and respiratory syncytial virus by ribavirin and immune serum globulin in vitro. *Antiviral research* 60:51–59. [PubMed: 14516921]
5. Wyde PR, Moylett EH, Chetty SN, Jewell A, Bowlin TL, Piedra PA. 2004 Comparison of the inhibition of human metapneumovirus and respiratory syncytial virus by NMSO3 in tissue culture assays. *Antiviral research* 63:51–59. [PubMed: 15196820]
6. El Najjar F, Schmitt AP, Dutch RE. 2014 Paramyxovirus glycoprotein incorporation, assembly and budding: a three way dance for infectious particle production. *Viruses* 6:3019–3054. [PubMed: 25105277]
7. Bakker SE, Duquerroy S, Galloux M, Loney C, Conner E, Eleouet JF, Rey FA, Bhella D. 2013 The respiratory syncytial virus nucleoprotein-RNA complex forms a left-handed helical nucleocapsid. *The Journal of general virology* 94:1734–1738. [PubMed: 23677789]
8. Ruigrok RW, Crepin T, Kolakofsky D. 2011 Nucleoproteins and nucleocapsids of negative-strand RNA viruses. *Current opinion in microbiology* 14:504–510. [PubMed: 21824806]
9. Renner M, Bertinelli M, Leyrat C, Paesen GC, Saraiva de Oliveira LF, Huiskonen JT, Grimes JM. 2016 Nucleocapsid assembly in pneumoviruses is regulated by conformational switching of the N protein. *eLife* 5:e12627. [PubMed: 26880565]
10. Tawar RG, Duquerroy S, Vonnrhein C, Varela PF, Damier-Piolle L, Castagne N, MacLellan K, Bedouelle H, Bricogne G, Bhella D, Eleouet JF, Rey FA. 2009 Crystal structure of a nucleocapsid-like nucleoprotein-RNA complex of respiratory syncytial virus. *Science* 326:1279–1283. [PubMed: 19965480]
11. Garcia-Barreno B, Delgado T, Melero JA. 1996 Identification of protein regions involved in the interaction of human respiratory syncytial virus phosphoprotein and nucleoprotein: significance

- for nucleocapsid assembly and formation of cytoplasmic inclusions. *Journal of virology* 70:801–808. [PubMed: 8551618]
12. Fields BN, Knipe DMDM-, Howley PM. 2013 Paramyxoviridae in *Fields virology* / editors-in-chief, Knipe David M., Howley Peter M. ; associate editors, Cohen Jeffrey I. ... [et al.]. *Fields virology*, 6th ed. ed. Philadelphia : Wolters Kluwer Health/Lippincott Williams & Wilkins, c2013., United States.
 13. Noton SL, Fearn R. 2015 Initiation and regulation of paramyxovirus transcription and replication. *Virology* 479-480:545–554. [PubMed: 25683441]
 14. Cressey TN, Noton SL, Nagendra K, Braun MR, Fearn R. 2018 Mechanism for de novo initiation at two sites in the respiratory syncytial virus promoter. *Nucleic acids research* 46:6785–6796. [PubMed: 29873775]
 15. Rincheval V, Lelek M, Gault E, Bouillier C, Sitterlin D, Blouquit-Laye S, Galloux M, Zimmer C, Eleouet JF, Rameix-Welti MA. 2017 Functional organization of cytoplasmic inclusion bodies in cells infected by respiratory syncytial virus. *Nature communications* 8:563.
 16. Norrby E, Marusyk H, Orvell C. 1970 Morphogenesis of respiratory syncytial virus in a green monkey kidney cell line (Vero). *Journal of virology* 6:237–242. [PubMed: 4100527]
 17. Garcia J, Garcia-Barreno B, Vivo A, Melero JA. 1993 Cytoplasmic inclusions of respiratory syncytial virus-infected cells: formation of inclusion bodies in transfected cells that coexpress the nucleoprotein, the phosphoprotein, and the 22K protein. *Virology* 195:243–247. [PubMed: 8317099]
 18. Carroumeu C, Simabuco FM, Tamura RE, Farinha Arcieri LE, Ventura AM. 2007 Intracellular localization of human respiratory syncytial virus L protein. *Archives of virology* 152:2259–2263. [PubMed: 17703289]
 19. Derdowski A, Peters TR, Glover N, Qian R, Utley TJ, Burnett A, Williams JV, Spearman P, Crowe JE Jr. 2008 Human metapneumovirus nucleoprotein and phosphoprotein interact and provide the minimal requirements for inclusion body formation. *The Journal of general virology* 89:2698–2708. [PubMed: 18931065]
 20. Cifuentes-Munoz N, Brantje J, Slaughter KB, Dutch RE. 2017 Human Metapneumovirus Induces Formation of Inclusion Bodies for Efficient Genome Replication and Transcription. *Journal of virology* 91.
 21. Heinrich BS, Cureton DK, Rahmeh AA, Whelan SP. 2010 Protein expression redirects vesicular stomatitis virus RNA synthesis to cytoplasmic inclusions. *PLoS pathogens* 6:e1000958. [PubMed: 20585632]
 22. Lahaye X, Vidy A, Pomier C, Obiang L, Harper F, Gaudin Y, Blondel D. 2009 Functional characterization of Negri bodies (NBs) in rabies virus-infected cells: Evidence that NBs are sites of viral transcription and replication. *Journal of virology* 83:7948–7958. [PubMed: 19494013]
 23. Becker S, Rinne C, Hofsass U, Klenk HD, Muhlberger E. 1998 Interactions of Marburg virus nucleocapsid proteins. *Virology* 249:406–417. [PubMed: 9791031]
 24. Kolesnikova L, Muhlberger E, Ryabchikova E, Becker S. 2000 Ultrastructural organization of recombinant Marburg virus nucleoprotein: comparison with Marburg virus inclusions. *Journal of virology* 74:3899–3904. [PubMed: 10729166]
 25. Hoenen T, Shabman RS, Groseth A, Herwig A, Weber M, Schudt G, Dolnik O, Basler CF, Becker S, Feldmann H. 2012 Inclusion bodies are a site of ebolavirus replication. *Journal of virology* 86:11779–11788. [PubMed: 22915810]
 26. Carlos TS, Young DF, Schneider M, Simas JP, Randall RE. 2009 Parainfluenza virus 5 genomes are located in viral cytoplasmic bodies whilst the virus dismantles the interferon-induced antiviral state of cells. *The Journal of general virology* 90:2147–2156. [PubMed: 19458173]
 27. Zhang S, Jiang Y, Cheng Q, Zhong Y, Qin Y, Chen M. 2017 Inclusion Body Fusion of Human Parainfluenza Virus Type 3 Regulated by Acetylated alpha-Tubulin Enhances Viral Replication. *Journal of virology* 91.
 28. Nikolic J, Le Bars R, Lama Z, Scrima N, Lagaudriere-Gesbert C, Gaudin Y, Blondel D. 2017 Negri bodies are viral factories with properties of liquid organelles. *Nature communications* 8:58.
 29. Santangelo PJ, Bao G. 2007 Dynamics of filamentous viral RNPs prior to egress. *Nucleic acids research* 35:3602–3611. [PubMed: 17485480]

30. Burke E, Dupuy L, Wall C, Barik S. 1998 Role of cellular actin in the gene expression and morphogenesis of human respiratory syncytial virus. *Virology* 252:137–148. [PubMed: 9875324]
31. Ulloa L, Serra R, Asenjo A, Villanueva N. 1998 Interactions between cellular actin and human respiratory syncytial virus (HRSV). *Virus research* 53:13–25. [PubMed: 9617766]
32. El Najjar F, Cifuentes-Munoz N, Chen J, Zhu H, Buchholz UJ, Moncman CL, Dutch RE. 2016 Human metapneumovirus Induces Reorganization of the Actin Cytoskeleton for Direct Cell-to-Cell Spread. *PLoS pathogens* 12:e1005922. [PubMed: 27683250]
33. Barik S 1992 Transcription of human respiratory syncytial virus genome RNA in vitro: requirement of cellular factor(s). *Journal of virology* 66:6813–6818. [PubMed: 1404620]
34. Huang YT, Romito RR, De BP, Banerjee AK. 1993 Characterization of the in vitro system for the synthesis of mRNA from human respiratory syncytial virus. *Virology* 193:862–867. [PubMed: 8460489]
35. Schudt G, Dolnik O, Kolesnikova L, Biedenkopf N, Herwig A, Becker S. 2015 Transport of Ebolavirus Nucleocapsids Is Dependent on Actin Polymerization: Live-Cell Imaging Analysis of Ebolavirus-Infected Cells. *The Journal of infectious diseases* 212 Suppl 2:S160–166. [PubMed: 26038396]
36. Schudt G, Kolesnikova L, Dolnik O, Sodeik B, Becker S. 2013 Live-cell imaging of Marburg virus-infected cells uncovers actin-dependent transport of nucleocapsids over long distances. *Proceedings of the National Academy of Sciences of the United States of America* 110:14402–14407. [PubMed: 23940347]
37. Mehedi M, McCarty T, Martin SE, Le Nouen C, Buehler E, Chen YC, Smelkinson M, Ganesan S, Fischer ER, Brock LG, Liang B, Munir S, Collins PL, Buchholz UJ. 2016 Actin-Related Protein 2 (ARP2) and Virus-Induced Filopodia Facilitate Human Respiratory Syncytial Virus Spread. *PLoS pathogens* 12:e1006062. [PubMed: 27926942]
38. Hamelin ME, Abed Y, Boivin G. 2004 Human metapneumovirus: a new player among respiratory viruses. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America* 38:983–990. [PubMed: 15034830]
39. Zhang L, Peeples ME, Boucher RC, Collins PL, Pickles RJ. 2002 Respiratory syncytial virus infection of human airway epithelial cells is polarized, specific to ciliated cells, and without obvious cytopathology. *Journal of virology* 76:5654–5666. [PubMed: 11991994]
40. Neilson KA, Yunis EJ. 1990 Demonstration of respiratory syncytial virus in an autopsy series. *Pediatric pathology* 10:491–502. [PubMed: 1695371]
41. Vargas SO, Kozakewich HP, Perez-Atayde AR, McAdam AJ. 2004 Pathology of human metapneumovirus infection: insights into the pathogenesis of a newly identified respiratory virus. *Pediatric and developmental pathology : the official journal of the Society for Pediatric Pathology and the Paediatric Pathology Society* 7:478–486; discussion 421.
42. Sattentau Q. 2008 Avoiding the void: cell-to-cell spread of human viruses. *Nature reviews. Microbiology* 6:815–826. [PubMed: 18923409]
43. Hotard AL, Lee S, Currier MG, Crowe JE Jr., Sakamoto K, Newcomb DC, Peebles RS Jr., Plemper RK, Moore ML. 2015 Identification of residues in the human respiratory syncytial virus fusion protein that modulate fusion activity and pathogenesis. *Journal of virology* 89:512–522. [PubMed: 25339762]
44. Roberts KL, Manicassamy B, Lamb RA. 2015 Influenza A virus uses intercellular connections to spread to neighboring cells. *Journal of virology* 89:1537–1549. [PubMed: 25428869]
45. Kumar A, Kim JH, Ranjan P, Metcalfe MG, Cao W, Mishina M, Gangappa S, Guo Z, Boyden ES, Zaki S, York I, Garcia-Sastre A, Shaw M, Sambhara S. 2017 Influenza virus exploits tunneling nanotubes for cell-to-cell spread. *Scientific reports* 7:40360. [PubMed: 28059146]
46. Liljeroos L, Krzyzaniak MA, Helenius A, Butcher SJ. 2013 Architecture of respiratory syncytial virus revealed by electron cryotomography. *Proceedings of the National Academy of Sciences of the United States of America* 110:11133–11138. [PubMed: 23776214]
47. van den Hoogen BG, de Jong JC, Groen J, Kuiken T, de Groot R, Fouchier RA, Osterhaus AD. 2001 A newly discovered human pneumovirus isolated from young children with respiratory tract disease. *Nature medicine* 7:719–724.

48. Bachi T, Howe C. 1973 Morphogenesis and ultrastructure of respiratory syncytial virus. *Journal of virology* 12:1173–1180. [PubMed: 4128827]
49. Peret TC, Boivin G, Li Y, Couillard M, Humphrey C, Osterhaus AD, Erdman DD, Anderson LJ. 2002 Characterization of human metapneumoviruses isolated from patients in North America. *The Journal of infectious diseases* 185:1660–1663. [PubMed: 12023774]
50. Loo LH, Jumaat MR, Fu Y, Ayi TC, Wong PS, Tee NW, Tan BH, Sugrue RJ. 2013 Evidence for the interaction of the human metapneumovirus G and F proteins during virus-like particle formation. *Virology journal* 10:294. [PubMed: 24067107]
51. Rager M, Vongpunsawad S, Duprex WP, Cattaneo R. 2002 Polyploid measles virus with hexameric genome length. *The EMBO journal* 21:2364–2372. [PubMed: 12006489]
52. Loney C, Mottet-Osman G, Roux L, Bhella D. 2009 Paramyxovirus ultrastructure and genome packaging: cryo-electron tomography of sendai virus. *Journal of virology* 83:8191–8197. [PubMed: 19493999]
53. Dahlberg JE, Simon EH. 1969 Physical and genetic studies of Newcastle disease virus: evidence for multiploid particles. *Virology* 38:666–678. [PubMed: 5816954]
54. Huang AS, Baltimore D. 1970 Defective viral particles and viral disease processes. *Nature* 226:325–327. [PubMed: 5439728]
55. Lazzarini RA, Keene JD, Schubert M. 1981 The origins of defective interfering particles of the negative-strand RNA viruses. *Cell* 26:145–154. [PubMed: 7037195]
56. Rima BK, Davidson WB, Martin SJ. 1977 The role of defective interfering particles in persistent infection of Vero cells by measles virus. *The Journal of general virology* 35:89–97. [PubMed: 404394]
57. Lopez CB. 2014 Defective viral genomes: critical danger signals of viral infections. *Journal of virology* 88:8720–8723. [PubMed: 24872580]
58. Treuhaft MW, Beem MO. 1982 Defective interfering particles of respiratory syncytial virus. *Infection and immunity* 37:439–444. [PubMed: 6288562]
59. van den Hoogen BG, van Boheemen S, de Rijck J, van Nieuwkoop S, Smith DJ, Laksono B, Gultyaev A, Osterhaus AD, Fouchier RA. 2014 Excessive production and extreme editing of human metapneumovirus defective interfering RNA is associated with type I IFN induction. *The Journal of general virology* 95:1625–1633. [PubMed: 24760760]
60. Valdovinos MR, Gomez B. 2003 Establishment of respiratory syncytial virus persistence in cell lines: association with defective interfering particles. *Intervirology* 46:190–198. [PubMed: 12867758]
61. Sun Y, Jain D, Koziol-White CJ, Genoyer E, Gilbert M, Tapia K, Panettieri RA Jr., Hodinka RL, Lopez CB. 2015 Immunostimulatory Defective Viral Genomes from Respiratory Syncytial Virus Promote a Strong Innate Antiviral Response during Infection in Mice and Humans. *PLoS pathogens* 11:e1005122. [PubMed: 26336095]
62. Sanjuan R, Nebot MR, Chirico N, Mansky LM, Belshaw R. 2010 Viral mutation rates. *Journal of virology* 84:9733–9748. [PubMed: 20660197]
63. Borderia AV, Stapleford KA, Vignuzzi M. 2011 RNA virus population diversity: implications for inter-species transmission. *Current opinion in virology* 1:643–648. [PubMed: 22440922]

Highlights

- Pneumoviruses are a major public health burden
- Recent evidence confirms pneumovirus replication occurs in inclusion bodies
- Pneumoviruses can spread directly from cell-to-cell
- Pneumoviruses potentially spread en bloc

Figure 1. Schematic representation of the pneumovirus genome and its replication.

A) The HMPV genome (red) contains different elements including the 3' leader (le) and 5' trailer (tr) regions, and gene start (dark green) and gene end (maroon) signals within each of the individual genes. RSV possess a larger genome that includes two additional proteins, NS1 and NS2 (not depicted). B) Inclusion bodies (IB, light blue) are the central replication station for pneumoviruses, and they result from the coalescence of nucleocapsids and smaller IBs, a process dependent on actin polymerization. From IBs, nucleocapsids are transported to assembly sites where they will be incorporated into nascent virions. Smaller IBs can also be transported long distances from the cell body.

Figure 2. En bloc mechanisms of spread proposed for pneumoviruses.

A) Cell-to-cell fusion is mediated by the viral fusion protein and results in formation of syncytia. The cytosolic contents of the infected cell, including IBs and viral mRNAs, are then transferred to the newly formed giant cell. B) Direct cell-to-cell spread mediated by actin-based intercellular extensions results in mass transport of genomes between cells. We propose this could happen either by sequential transport of individual nucleocapsids, or by transport of IBs, which would result in a fast avenue to establish infection in the target cell. C) Newly assembled virions can contain one or multiple copies of the viral genome. The advantage of carrying multiple genomes is that after entry, these will act in a complementary way to establish rapidly infection by formation of IBs. Red arrowheads represent efficiency for establishment of large IBs.