Published in final edited form as:

Mol Cancer Ther. 2018 September ; 17(9): 2034–2048. doi:10.1158/1535-7163.MCT-17-0421.

Targeting Notch1 and IKK α enhanced NF- κ B activation in CD133⁺ Skin Cancer Stem Cells

Xin Xin Quan¹, Nga Voong Hawk², Weiping Chen³, Jamie Coupar¹, Steven Lee¹, David W Petersen⁴, Paul S Meltzer⁴, Andrew Montemarano⁵, Martin Braun⁶, Zhong Chen^{1,7}, and Carter Van Waes^{1,7}

¹Tumor Biology Section, Head and Neck Surgery Branch, National Institute on Deafness and Other Communication Disorders, NIH;

²Experimental Transplantation & Immunology Branch, National Cancer Institute, NIH;

³Microarray Core Facility, National Institute of Diabetes & Digestive & Kidney Diseases, NIH;

⁴Cancer Genetics Branch, National Cancer Institute, NIH;

⁵Skin Cancer Surgery Center, Bethesda, MD, 20817;

⁶Braun Dermatology Associate, Washington D.C., 20037.

⁷Contributed equally as senior authors.

Abstract

Cancer stem-like cells are hypothesized to be the major tumor initiating cell population of human cutaneous squamous cell carcinoma (cSCC), but the landscape of molecular alterations underpinning their signaling and cellular phenotypes as drug targets remain undefined. In this study, we developed an experimental pipeline to isolate a highly enriched CD133⁺CD31⁻CD45⁻CD61⁻CD24⁻ (CD133⁺) cell population from primary cSCC specimens by flow cytometry. The CD133⁺ cells show enhanced stem-like phenotypes, which were verified by spheroid and colony formation *in vitro* and tumor generation *in vivo*. Gene expression profiling of CD133⁺ cells was compared and validated, and differentially expressed gene signatures and top pathways were identified. CD133⁺ cells expressed a repertoire of stemness and cancer related genes, including NOTCH and NOTCH1-mediated NF- κ B pathway signaling. Other cancer-related genes from WNT, growth factor receptors, PI3K/mTOR, STAT pathways, and chromatin modifiers were also identified. Pharmacologic and genetic targeting Notch1, IKK α , RELA and RELB modulated NF- κ B transactivation, the CD133⁺ population, and cellular and stemness phenotypes. Immunofluorescent staining confirmed co-localization of CD133⁺ and IKK α expression in SCC tumor specimens. Our functional, genetic and pharmacologic studies uncovered a novel linkage between Notch1, IKK α and NF- κ B pathway activation in maintaining the CD133⁺ stem SCC phenotypes. Studies investigating markers of activation and modulators of NOTCH, IKK/NF- κ B

Correspondence: Zhong Chen, MD, PhD; Phone: 301-435-2073; Fax: 301-402-1140; chen@nidcd.nih.gov; Head and Neck Surgery Branch, NIDCD/NIH, 10/7N240, Bethesda, MD 20892. Or Carter Van Waes, MD, PhD; Phone: 301-402-4216; Fax: 301-402-1140; vanwaesc@nidcd.nih.gov; Head and Neck Surgery Branch, NIDCD/NIH, 10/7N240, Bethesda, MD 20892.

Conflict of Interest Statement: The authors have declared that no conflict of interest exists.

and other pathways regulating these cancer stem gene signatures could further accelerate the development of effective therapeutic strategies to treat cSCC recurrence and metastasis.

Keywords

CD133⁺; IKK; NF- κ B; Notch; drug targets; skin cancer; cancer stem cells

Introduction

Human cutaneous squamous cell carcinoma (cSCC) has been increasing over the past several decades, with more than 700,000 cases in the United States annually (1,2). Despite surgery, radiation, and chemotherapy, SCC cells can escape treatment and reform tumors and metastasis, increasing morbidity and mortality. This gives prominence to the existence of a subpopulation of SCC cells capable of tumor initiation and therapeutic resistance, and the importance of characterizing and targeting the molecular alterations mediating their maintenance for cancer prevention and treatment.

The cancer stem cell (CSC) hypothesis holds that tumors are a hierarchical organ derived from cell sub-population(s) capable of self-renewal, called cancer stem-like cells or tumor initiating cells (TICs)(3). TICs exhibit stem-like and tumor initiating properties, including increased self-renewal/colony formation, tumor forming capacity, as well as altered migration, differentiation and therapeutic sensitivity. Evidence for the existence of TICs has been obtained in different types of human solid tumors through identification of sub-populations enriched for surface determinants or enzymatic markers, such as CD133, CD44, CXCR4, and ALDH1(3). We previously demonstrated that cell membrane protein CD133 (also called prominin-1) is specifically expressed by cSCC cells enriched for a TIC phenotype, and not by CD45 and other non-epithelial subpopulations overlapping CD44 (4). CD133⁺ cSCC cells exhibited long-term proliferative ability, self-renewal, sphere formation, regeneration of differentiated SCC tumor cells, and enriched tumor initiating capacity for xenografts in immunodeficient mice. CD133⁺ has also been identified as a biologically and clinically relevant marker for sub-populations of lung and head and neck SCC (HNSCC), where it is associated with tumor initiating capacity, aggressive clinical features, and resistance to cytotoxic therapies (5,6).

Recent studies of cSCC, HNSCC and lung SCC tumors have identified significant genetic alterations involving components of several common and distinct pathways important in cell growth, death or survival, migration, and epithelial/mesenchymal differentiation (7–10). These alterations are implicated in inactivation or activation of several canonical pathways including Notch, Wnt, Hedgehog, NF- κ B, growth factor receptors, RAS-mitogen activated protein kinase, PI3K-Akt-mTOR, and TP53, but their expression and role in TIC versus other populations in established cSCC has not been dissected. Among these are numbers of genes linked to stem cell maintenance or differentiation (11–14). Notch signaling is important in epithelial differentiation and as a suppressor of tumor development in a subset of HNSCC, but is activated in others (8, 15). IKK and NF- κ B signaling has been implicated in promoting tumor cell survival, inflammatory and angiogenesis responses (9). However,

how these signaling pathways and function of corresponding molecular components contribute to the regulation of phenotype of the CSC/TIC sub-population in cSCC tumors are not well understood. Identification of deregulated components of pathways critical to maintenance of CD133⁺ CSC phenotype could potentially help identify targets for precision medicine for prevention and therapy.

In this study, we integrated molecular profiling, signaling pathway, mechanistic, and pharmacological studies of the CD133⁺ sub-population in primary human cSCC tumors and cell line models. Through cell sorting with multiple positive and negative selection markers by flow cytometry, we successfully isolated live CD133⁺ cells that form spheroid colonies *in vitro* and tumors *in vivo*. This small distinct CD133⁺ population differentially expresses stem-like and cancer gene signatures linked to Notch1-mediated NF- κ B modulation, NF- κ B, and WNT pathways. Characterization of the landscape of gene signatures in these CD133⁺ stem cells revealed activation of multiple pathways, which were linked to NOTCH and NF- κ B signaling networks and showed sensitivity to genetic and pharmacologic inhibitors of NOTCH and NF- κ B. Our functional, genetic and pharmacologic studies uncovered a linkage between Notch1, IKK α and NF- κ B pathway activation in maintaining the CD133⁺ population and its self-renewal ability in established primary cSCC and cell lines.

Materials and Methods

Human tumor tissue samples.

De-identified primary human skin cSCC tissue samples were obtained under an exemption from IRB approval by the Office of Human Subjects Research, National Institutes of Health, from Potomac Ambulatory Surgery Center, Bethesda, MD, and Braun Dermatology Associates, Washington D.C.

Preparation of single cell suspension from cSCC specimens

Tumor tissues were gently chopped into small pieces and incubated with collagenase III and dispase, then further dissociated using GentleMACS Dissociator. The dissociated tissues were spun down, incubated with trypsin and EDTA and filtered, spun down, and the cSCC single-cell suspension was then ready for staining and FACS sorting for sphere, tumorigenicity assays, and expression profiling.

Flow cytometry analysis and FACS sorting.

cSCC single cell suspensions were labeled with multiple antibodies and sorted by BD FACSARIA-II following standard protocol as detailed in Supplemental Methods. Analysis was performed with BD LSR II and FlowJo7.6.5.

Sphere formation assay.

Fifteen cSCC samples were collected and used for sphere experiments after preparation of single cell suspension. The cSCC single cell suspension of unsorted or sorted CD133⁺ or – cells was seeded on NIH 3T3 fibroblast cells feeder layer, and cultured for 14–21 days. Spheres $\geq 50\mu\text{m}$ were counted under the microscope.

***In vivo* mouse model.**

Animal studies were performed under a protocol approved by the Institutional Animal Care and Use Committee (IACUC) of the National Cancer Institute. 6–8wk old nude mice were obtained from NCI Frederick. To establish a niche, 10^6 human endothelium and fibroblast cells with 120 μ l Matrigel were injected into 0.5×0.5 cm GelFoam sponges implanted subcutaneously 2 weeks prior to tumor cell transplantation. Freshly obtained, dissociated and sorted CD133⁺, CD133⁻ and unsorted cells at doses of 10^6 , 10^5 , 10^4 , 10^3 , or 10^2 cells were mixed with 10^6 human endothelium and fibroblast cells in Matrigel, for injection into the established GelFoam niche. Each dose level included 15 mice with 5 mice each that received freshly isolated primary sorted CD133⁺, CD133⁻, or unsorted SCC tumor cells. To obtain 10^6 rare CD133⁺ and matched CD133⁻ cells, we sorted dissociated suspensions from pools of 5, 5, 6, 6, 6 (28 total) fresh tumors for inoculation of 5 pairs of mice. CD133⁺ and – cells sorted from 5 tumors were sufficient to inoculate 5 paired recipients at 10^2 , 10^3 , 10^4 and 3 pairs at 10^5 , and cells pooled from 3 and 4 additional tumors each (12 total) were required to sort 10^5 CD133⁺ and – cells for 2 additional pairs. Five tumors were freshly dissociated and 10^6 , 10^5 , 10^4 , 10^3 , and 10^2 cells from each were inoculated into paired recipients as unsorted controls for each dose level.

Affymetrix GeneChip Human Gene 1.0 ST Array.

Total RNA was isolated after FACS sorting and fifteen samples were used with RNA integrity numbers (RINs) equal or bigger than 7.0, as the control for RNA quality. Single stranded cDNA was generated, fragmented, labeled and hybridized with standard Affymetrix protocol. Data were scanned with an Affymetrix GeneChip Scanner 3000, analyzed with Partek and GeneGo. The microarray data has been submitted to NCBI with GEO Submission number (GSE84588).

Nanostring nCounter gene expression assay.

100 ng total RNAs from sorted CD133⁺ and CD133⁻ cells from seven cSCC samples were hybridized, purified and immobilized in the Prep Station. The images were measured and further analyzed with nSolver.

NF- κ B family transcription factor DNA-binding assays.

Nuclear fraction of SCC13 cells was placed on a plate coated with immobilized NF- κ B consensus oligonucleotides. Activated NF- κ B is recognized by antibodies, following with the HRP-conjugated second antibody and quantified by spectrophotometry.

Drugs and treatment.

The following compounds were used for inhibition of gamma-secretase-NOTCH and Inhibitor-kappaB Kinase in experiments: DAPT (<https://pubchem.ncbi.nlm.nih.gov/compound/dapt>), catalog D5942, Sigma Aldrich, St Louis, MO); RO4929097 (<https://pubchem.ncbi.nlm.nih.gov/compound/49867930#section=Top>), catalog No. ADV465749148, Sigma Aldrich, St Louis, MO); Wedelolactone (<https://pubchem.ncbi.nlm.nih.gov/compound/Wedelolactone#section=Top>), catalog #56639, Sigma Aldrich, St Louis, MO).

SCC13 cells were treated with two different Notch inhibitors, DAPT or R04929097 in different doses and measured by FACS analysis. 10 μ M wedelolactone was used to treat SCC13 cells for 24 hours. Cells then harvested and stained with CD133 antibody for FACS analysis. Spheres or colonies were counted after 5 days drug treatment. Medium with drug were changed every three days.

Luciferase reporter gene assays.

SCC cells were co-transfected with NF- κ B Luciferase and β -Gal and then treated with wedelolactone and TNF- α at different time points. Relative luciferase activity was normalized to the cells without treatment.

Western blotting.

15 μ g whole cell lysates were used for electrophoresis, then protein was transferred onto a Nitrocellulose membrane, and incubated with first and second antibodies following standard protocol.

Statistical analysis.

Mean and standard deviation (SD) were calculated for different experiments, and the statistical significance was calculated using Student's t-test. P value <0.05 was considered a statistically significant difference. To analyze microarray data, a two-way analysis of variance (ANOVA) for the paired samples was used.

Additional information of human SCC specimens, antibodies, and instrumentation settings are presented in the Supplemental Information.

Results

Clinical characteristics of human primary cutaneous squamous cell carcinomas (cSCC).

Eighty-two primary cSCC were collected fresh after Mohs micrographic surgery from patients. The diagnosis and the tumor's grade were confirmed by a dermatopathologist, and the clinical characteristics are summarized in Supplemental Table 1. The majority of subjects were men (78%), and tumors were predominantly located on sun-exposed skin from head (62.2%) and limbs (34.2%). Most tumors were well or moderately differentiated (85.4+12.2=97.6%) cSCC of less than 2cm² in size. The collected tumors were freshly delivered from the clinics and processed on the same day, to isolate stem cell subpopulations for *in vitro* and *in vivo* characterization of purity, stemness phenotype, molecular profiling, and functional validation studies (Supplemental Table 1, Supplemental Figure 1). The clinical characteristics of tumor samples used in different experimental studies below are summarized Supplemental Table 2.

CD133⁺CD31⁻CD45⁻CD61⁻CD24⁻ cells isolated from cSCC exhibit enriched CSC/TIC features by *in vitro* and *in vivo* assays.

New protocols to isolate, dissociate and separate cSCC cells from other populations in tumor specimens were optimized, as summarized in Supplemental Methods and Supplemental Figure 1. Each cSCC sample was microdissected to carefully remove excess surrounding

stroma, and regions with >90% malignant epithelia were obtained. We developed protocols for cell sorting and improved purification of CD133⁺ and CD133⁻ tumor cells after gating out differentiating keratinocytes (CD24), stromal endothelial cells (CD31), leukocytes (CD45), and fibroblasts (CD61). The percentage of cells expressing these CD markers in cSCC varied among individual tumors by FACS analysis (Figure 1A). CD133⁺ cells represent a rare population that resides within a relatively stable range from 0.14–1.7% (mean 0.62%)(Figure 1A, B), and displayed a slight increase with tumor size that was not significant ($R^2=0.037$, $p>0.05$). Sorted CD133⁺CD31⁻CD45⁻CD61⁻CD24⁻ (hereafter called CD133⁺) cells, have a purity as high as 95% as detected by FACS analysis (Figure 1C). More than 99% of FACS sorted CD133⁺ cells are Pan-Keratin positive (Figure 1D), indicating that the CD133⁺ subpopulation represents highly purified CD133⁺ keratinocytes from cSCC tissues. Thus, we isolated highly purified CD133⁺ CSC and compared with those relatively pure CD133⁻ keratinocytes without stemness characteristics using this improved sorting and selection protocol with the five different cellular markers.

To confirm whether the CD133⁺ population exhibits TIC features including self-renewal and tumorigenic ability established previously (3), we compared them with both CD133⁻ and unsorted cells for sphere formation efficiency *in vitro* and tumor formation *in vivo*. Unlike normal human keratinocytes growing as monolayer colonies, dissociated cSCC cells grew as tethered spheres on NIH 3T3 feeder layers (Figure 1E). Further, 1×10^3 sorted CD133⁺ cells formed 55 ± 8 spheres, while the same number of unsorted and CD133⁻ cSCC cells grew only 23 ± 3 and 23 ± 4 spheres, respectively (Figure 1F). The sphere formation in CD133⁺ was significantly increased ($p<0.0001$) compared with unsorted cSCC or CD133⁻ cells, whereas there is no difference between the unsorted and the CD133⁻ tumor cell groups. The sphere numbers increased corresponding to increases in the seeded cell numbers, indicating that cSCC cell aggregation alone is unlikely to account for sphere formation.

To confirm the enrichment of TICs within the purified CD133⁺ population, we used the cSCC tumor xenograft assay in a mouse model established previously (3), modified to replenish the normal human fibroblasts and endothelial cells depleted by sorting. Varying concentrations of sorted CD133⁺, CD133⁻, and unsorted cells were injected subcutaneously, to analyze tumor formation frequency by limiting dilution assay. A minimum of 10^5 unsorted cSCC cells was required for initiating tumor in this model (Figure 1G, blue bar). By contrast, as few as one hundred CD133⁺ cells generated tumors *in vivo* (Figure 1G, green bars). At the same time, the CD133⁻ subpopulation did not form any visible tumors with as high as 10^6 cells (Figure 1G, yellow bar). Thus, the estimated tumor initiating efficiency of the CD133⁺ subset is $1/45$, which is dramatically enriched ~340 fold compared to unsorted cSCC cells ($1/83,333$).

Gene expression profiling of CD133⁺ cells display gene enrichment for stem cell markers and NOTCH and NF- κ B signaling pathways.

To investigate genes differentially expressed in CD133⁺ cSCC population, we compared the whole genome expression profile between sorted CD133⁺ and CD133⁻ cells after gating for CD31⁻CD45⁻CD61⁻CD24⁻ population from 15 fresh cSCC tumor samples by using Affymetrix Whole Genome RNA microarray. The microarray data were normalized by the

robust multichip average (RMA) method and analyzed using the ANOVA paired test in Partek Genomics Suite, with p value 0.05 and expression level change > 1.5 folds. There were 2,089 genes significantly differentially expressed in CD133⁺ when compared to CD133⁻ tumor cells. We further compared this gene list with 247 stem-cell related genes from major publications of human stem cell microarray data (2,11–14). We identified 80 overlapped genes between 2,089 genes identified in this study versus 247 stem-cell related genes from major publications (Supplemental Figure 2A). After subtraction of the 80 genes, the Venn diagram of pink cycle shows total of 2009 genes with significantly altered expression in CD133⁺ cells compared with CD133⁻ cells. There are 167 stem genes from literatures after subtraction of 80 genes (green circle). Our data indicate that the sorted CD133⁺ cells are enriched for a stem cell gene expression signature. Moreover, a heat map of 80 signature genes shows that the genes are mainly distributed in two clusters (Figure 2A). Cluster I includes 32 up-regulated stem-related genes in CD133⁺ cells, including several Notch signaling pathway genes *NOTCH1*, *JAG2* and *MAML1*; Wnt pathway genes *WNT3*, *FZD1* and *CTNGB1*; and Sonic Hedgehog pathway genes *GLI2*, *SUFU*, *PTCH1*, and *GSCNF* (Germ Cell Nuclear Factor). Cluster II shows 48 down-regulated genes in CD133⁺ group, including Notch signaling related genes *JAG1*, *NOTCH4* and downstream components *RBPI*, and *NUMB*; Wnt signaling negative regulator gene *GSK3B* and hedgehog signaling regulator *HHIP*. Unexpectedly, we also detected increased expression of hematopoietic cell determinant *CD8A* in several CD133⁺ samples. CD8a has been associated with CD133⁺ lympho-myeloid stem cells and cytotoxic T lymphocytes, which could potentially infiltrate cSCC and be detected among a subset of samples (16–19).

GeneGo analysis of this 80-stem gene signature revealed the Notch pathway, Notch1-mediated NF- κ B pathway, Wnt pathway, Hedgehog, and Notch-mediated EMT signaling pathways are the top five most significantly ranked pathways in CD133⁺ cSCC cells (Supplemental Figure 2B, Supplemental Table 3). To further investigate what kinds of signaling pathways contributed in the CD133⁺ population, we analyzed pathways ranking for all 2089 significantly different expressed genes based on the ANOVA paired test results (Supplemental Figure 2C). The broader analysis showed that molecular mechanisms of cancer, regulation of EMT pathways, PI3K/AKT, mTOR, NF- κ B, ERK/MAPK signaling, and RAR activation are within the top twenty pathways in the CD133⁺ cell subset.

Since Notch, Notch1-mediated NF- κ B, and Wnt signaling appeared in top ranked pathways in both stem cell supervised and unsupervised analyses, we generated heat maps for genes included in the top ranked signaling/stem gene signatures from the 2089 differentially expressed genes in cSCC (Figure 2B, C, Supplemental Figure 2D). The heat map of Notch pathway showed two distinct clusters (Figure 2B): cluster I displayed relative increase in gene expression, of ligand *JAG2*, receptors *NOTCH1*, *NOTCH2*, *NOTCH3*, and downstream transcription factor *MAML1*; cluster II and III revealed down-regulated gene expression, including ligands *JAG1*, *DLL1*, receptors *NOTCH4*, glycosylation modification factor *POFUT2*, and downstream transcription factor component *RBPI*. CD133⁺ cSCC did not exhibit significant differential expression of canonical Notch target genes *HES* and *HEY1*, consistent with a recent study which reported defective transcription of *HES* and *HEY1* in subsets of human HNSCC with wild-type Notch genes (15).

We next examined the profile of NF- κ B related pathway genes in cSCC based on our hypothesis that Notch and other mediators could contribute to regulation of the NF- κ B signaling network (20–23). The clustering data revealed increased gene expression of several key signal and transcription factor subunits for the NF- κ B pathway in CD133⁺ cells (Figure 2C). These include *IL1R1*, *IRAK2/4*, *TRAF2/3*, *TRAF3IP2*, *CHUK (IKKalpha)*, *IKBKB*, *REL*, *NFKB2*, *RELB*, as well as *AKT*, implicated in NF- κ B activation by PI3K-Akt signaling. In addition, the heat map of Wnt pathway showed increase in *WNT3*, *DVL2* and *CTNNB1*, and decrease in expression of the repressors such as *AXIN2* and *GSK3B*, in CD133⁺ SCC cells (Supplemental Figure 2D).

We hypothesized Notch, NF- κ B, and Wnt pathways could form interactive networks in cSCC CD133⁺ cancer stem cell populations, and used Ingenuity Pathway Analysis (IPA) to reveal the potential regulatory relationships annotated among these three pathways. The results showed experimental evidence for altered expression and annotated links between NF- κ B with the Notch (Figure 2D) or Wnt pathway (Figure 2E). The network map suggests the hypothesis that non-canonical NOTCH1 signaling could contribute to the activation of REL and RELB in cSCC (Figure 2D). The interaction of NF- κ B with Wnt- β -catenin pathway is more indirectly through other nodes and pathways (Figure 2E).

Independent validation of stem and pathway signature gene expression using Nanostring in a different set of cSCC tissue samples.

To independently validate the microarray results, we verified gene expression levels with sorted CD133⁺ and CD133⁻ cells from seven additional SCC tumors using the Nanostring nCounter Gene Expression assay, for 149 selected genes from the stem- and top-ranked cancer-related pathway gene signatures. Notch signaling pathway components *NOTCH1*, *NOTCH2* and *MAML1* genes were confirmed to have higher expression levels, while *JAG1* and *EP300* showed decreased expression (Figure 3A). The genes involved in NF- κ B pathway also showed up-regulated expression, including *IL1R1*, *IRAK2/4*, *CHUK*, and *RELB*, as well as *FAS*, *TLR6*, *PRKACA*, *MMP8* and *ICAM4*. In the Wnt pathway, increased *WNT3* and *DVL2* expression and decreased *GSK3B* and *DVL3* expression in CD133⁺ cells were independently verified using Nanostring.

Furthermore, genes in several other signaling pathways displaying genomic or expression alterations and implicated in pathogenesis of SCC (8–10) were confirmed by Nanostring validation (Figure 3B, 3C). In CD133⁺ cells, increased gene expression was observed in EGFR/ERBB, *IGF1R*, *AKT* and *mTOR*, and STAT family genes, as well as those implicated in FGF, MAPK, TGF- β signaling pathways (Figure 3B). Among other validated genes, cell cycle related genes *FBXW11*, *PPP2R5C* are overexpressed, while *YAP1* and *CTNNA1* related to cell growth and differentiation, are decreased (Figure 3C). Additionally, chromatin regulatory factors including *DNMT1* and *EZH2* genes were decreased while *HDAC6* is overexpressed. The differential expression of mRNA for several of these genes in CD133⁺ vs CD133⁻ cells is not always congruent with the predicted effects of gene mutations (*NOTCH1*), copy number alterations (*YAP1*), or overall expression (*EZH2*) observed in SCC subsets vs non-malignant epithelia (7–9, 19, 20), underscoring the importance of functional validation.

Inhibiting Notch reduces the CD133⁺ cell population.

As Notch signaling has been implicated in suppressing differentiation or promoting growth in different SCC (7, 15, 21), we first examined the functional effects of two different γ -secretase inhibitors (DAPT and RO4929097) that inhibit Notch, upon the CD133⁺ subset in the human cSCC cell line SCC13 (23). We confirmed that SCC13 exhibits a similar small percentage of CD133⁺ cells varying between 0.2–6%, depending on culture conditions, and displays clonogenic and sphere forming potential, as observed by primary cSCC tumors. Three concentrations of DAPT (1, 5, 10 μ M) and RO4929097 (1, 3, 5 μ M) were selected based on ranges found to include IC₅₀ for other tumor cell lines (24). FACS analysis shows that the CD133⁺ cell population is significantly inhibited by 5–10 μ M DAPT and 1 μ M RO4929097, supporting a functional role of Notch1 in maintenance of the CD133⁺ cells (Figure 4A). We next tested if these two Notch inhibitors could affect CD133⁺ cell colony formation of SCC13, as well as UM-SCC-46, an HNSCC line verified to express wtNotch1 and pathway genes by whole exome and RNA sequencing (Hui Cheng, unpublished observations). Colony formation of both SCC13 and UM-SCC46 cells was significantly decreased by DAPT or RO4929097 (Figure 4B; blue, red bars).

To independently examine the specificity of these findings for Notch1 or 2 differentially expressed in tumor CD133⁺ cells in Fig. 3A, we examined the expression of Notch1 and 2 protein, and tested the effects of small interfering (si)RNAs silencing Notch1 and Notch2 on the CD133⁺ cell population in two cell lines. We confirmed that Notch1 protein of expected size is expressed and knocked down efficiently by siRNA in both SCC13 and UM-SCC46 cells, while weak expression of Notch2 protein is only detected with prolonged exposure, and weakly inhibited (Figure 4C, upper panel). Consistent with the differential expression and knockdown of Notch proteins, the relative CD133⁺ cell number decreased ~40% of control with Notch1 knockdown alone, while no effect was attributable to the limited expression and knockdown of Notch 2 alone or when combined with Notch1 siRNA (Figure 4C, lower panel). Together the functional effects observed with pharmacologic and genetic approaches suggest Notch1 signaling contributes to CD133⁺ population and colony forming capability in a cSCC as well as a HNSCC cell line expressing wild-type Notch1.

Inhibiting Notch reduces NF- κ B transcriptional and nuclear DNA binding activity in cSCC cells.

Since Notch1-mediated NF- κ B activation is the second top-ranked GeneGo analysis signature for the CD133⁺ cell population, and aberrant activation of NF- κ B has been observed in SCC (8, 25), we explored the possibility that Notch signaling enhances NF- κ B pathway signaling, by assay for NF- κ B reporter gene activity after Notch inhibition. As shown in Figure 4D, inhibitors of γ -secretase and Notch activation, DAPT and RO4929097, decrease TNF- α induced NF- κ B reporter activity in human SCC13 cells in a dose-dependent manner. To further confirm the above results, small interfering RNA (siRNA) was used to specifically knock down Notch1 and/or Notch2, and nuclear DNA binding of five NF- κ B nuclear transcription factor subunits was assayed and quantified. The results showed that after transfection with Notch1 siRNA, the DNA-binding activity of canonical NF- κ B pathway subunit p65 (RelA), and non-canonical pathway subunits p52, and RelB, were significantly decreased to 64%, 80% and 77% respectively ($p < 0.001$) compared to the

control siRNA group (Figure 4E left panel). Notch2 siRNA showed similar but weaker inhibitory effects on nuclear DNA binding activity of these three transcription factor components, and combination of Notch2 with Notch1 siRNA did not further inhibit NF- κ B nuclear binding activity (Figure 4E middle and right panels). These results provide evidence for Notch modulated NF- κ B pathway activity, which could potentially promote the CD133⁺ population in cSCC cells.

IKK-NF- κ B pathway signaling promotes the CD133⁺ cell population in cSCC.

As IKK α (CHUK) and other NF- κ B pathway components were differentially overexpressed in CD133⁺ SCC tumor cells, we screened SCC13 cells for the effects of wedelolactone, an inhibitor shown to inhibit IKK α and β kinases, and IKK α dependent NF- κ B activation in SCC cells (25, 26). Wedelolactone inhibits TNF- α inducible NF- κ B reporter activity assayed 24h and 48 after treatment in SCC13 cells (Figure 5A). To determine the effects of NF- κ B inhibition on SCC CD133⁺ cells, we analyzed the CD133⁺ cell population with or without wedelolactone treatment. The FACS results showed lower concentrations of 1–5 μ M wedelolactone reduce CD133⁺ cells, while 10 μ M wedelolactone significantly decreases the CD133⁺ population (Figure 5B, $p < 0.001$). The effect of wedelolactone on CD133⁺ population is not due to general cytotoxicity. The above results suggest the NF- κ B pathway contributes to maintaining the cSCC CD133⁺ cancer stem cell population *in vitro*.

Expression and activation of IKK α promotes the CD133⁺ cell population in cSCC.

As both our microarray and Nanostring verification data showed there is an increased *CHUK* (*IKK α*) gene expression in CD133⁺ cells, and IKK inhibitor wedelolactone could decrease CD133⁺ population in human SCC13 cells, we independently investigated the specific role and function of IKK α for maintenance of CD133⁺ CSC in cSCC. To achieve this, we first tested the effect of siRNA knockdown of endogenous IKK α on the CD133⁺ cells. Knockdown by siRNA efficiently inhibited cytoplasmic and nuclear of IKK α (Figure 5C), and reduced the CD133⁺ population by ~50% compared to the control group in SCC13 cells (Figure 5D). Similar effects were also observed in the HNSCC cell line UMSSC46. Next we examined the effects of introducing IKK α wild type (WT) or activated or inactivated genetic IKK α mutants upon the CD133⁺ population. These include IKK α with serine phospho-acceptor sites substituted with constitutively activating glutamate residues (EE); inactivating alanines (AA); or Lysine44 substituted with Alanine that produces a catalytically inactive form (KA) (25). Cells expressing IKK α WT or EE protein significantly increased the CD133⁺ population 2.1 ($p < 0.01$) and 3.5 folds ($p < 0.001$) in SCC13 cells, whereas the inactive IKK α forms AA or KA transfection exhibited no increase of CD133⁺ subset (Figure 5E). Furthermore, we confirmed that the increase in the CD133⁺ population by overexpression wild type IKK α (WT) or its active form (EE), can be significantly inhibited by IKK α siRNA knockdown (Figure 5F).

To further examine the potential relationship between IKK α and CD133⁺ population in cSCC *in vivo*, we performed immunofluorescent (IF) staining of cSCC specimens. Figure 5G shows colocalization of IKK α and CD133⁺ immunostaining in cell sub-populations in two different primary cSCC tissues. The tumor specimens from two patients in Figure 5G and Supplemental Figure 3C/D show merged staining of CD133 and IKK α , as well as with

DAPI, at the basal layer and leading edge of epithelial tumor cell nests shown in adjacent H&E sections. The co-stained population of cells is seen along the interface of malignant epithelia with intervening inflammatory (Supplementary Figure 3C & 3D) or fibrovascular stroma (Figure 5G). In addition, Supplemental Figure 3A and 3B show merged staining of CD133 and IKK α for cells in two additional SCC tumors, over a wider region at 100X magnification. These data are consistent with the differential co-expression of these detected in CD133⁺ cells from multiple tumors, and functional effects of IKK α genomic mutants on CD133⁺ cells in cell lines. Together, these observations support a role for increased IKK α expression in the CD133⁺ population in cSCC cells and tumors.

Knockdown of RELA and/or RELB inhibits cSCC CD133⁺ population and its colony/spheroid formation phenotype.

IKK α can mediate downstream canonical and alternative NF- κ B pathway activation through RELA and RELB, that are found overexpressed in CD133⁺ cells. Next, we examined the knockdown effects of RELA and/or RELB on the CD133⁺ population in cSCC SCC13 and HNSCC UMSCC46 cells (Figure 6A). The CD133⁺ cell populations were clearly reduced to 54% with RELA siRNA, 43% with RELB siRNA, and 43% with combined knockdown in SCC13 cells (Figure 6B, blue bars). Similar results were observed in UMSCC46 cells (Figure 6B, green bars). Next, we quantified the colony/spheroid formation ability in SCC13 and UMSCC46 cells after knockdown of RELA and/or RELB. Clonogenic colonies from CD133⁺ cells were reduced to 78%, 63% and 55% by RELA or RELB siRNA knockdown alone, or in combination in SCC13 cells (Figure 6C, pink bars). Spheroid formation in UMSCC46 cells also decreased to 76%, 69% and 68% respectively (Figure 6C, purple bars, 6D). Together, the above data indicate both IKK α and NF- κ B components RELA and RELB play an important role in maintaining the CD133⁺ population and its cancer stem-like features in human SCC.

Discussion

To our knowledge, this is the first study to demonstrate the CD133⁺ gene expression signature and its top ranked signaling pathways in a purified cancer stem cell-like subset from primary cSCC tumors. There are four important findings from our research. First, our optimized isolation and separation further enriched for high purity CD133⁺ cytokeratin⁺ cells, overcoming a significant technical obstacle and enabling whole gene expression profiling detection and validation of stem and cancer genes in the small CD133⁺ population from clinical tumor samples. Second, our study provides a global overview of the gene signatures of CD133⁺ CSC in the human cSCC. Using microarray, nanostring validation and bioinformatics analyses, these data provide a valuable resource for hypothesis generation and testing for functional and clinical role of different signaling components and pathways in cSCC CD133⁺ CSC. Our stem gene expression profiling data identified ~2000 candidate genes differentially expressed in CD133⁺ cSCC cells, including an 80-gene set of known stem cell signatures. Third, we demonstrated upregulated expression of genes of key signaling pathways and networks from the cancer stem cell population by analyzing the top ranked pathways involved with GeneGO and IPA platform for the CD133⁺ transcript signature. The CD133⁺ transcript signatures represent pathways implicated in self-renewal

and epithelial differentiation, such as Notch, Wnt, Shh, and NF- κ B, and cancer-related pathways PTEN, PI3K/AKT/mTOR, NGF, MAPK, EGFR/ERBB, FGF, JAK/STAT, TP53, and TGF- β . Genomics and experimental studies provide evidence supporting the biologic and clinical significance of several of these pathways to SCC biology (7,8). Complementing our observations, recent whole exome sequencing studies of cSCC and HNSCC provide evidence for gene copy alterations or mutations in components or co-regulators of the Notch (*Notch 1,2*), Wnt (*AJUBA*), cancer-related MAPK and PI3K growth (*HRAS, RASA1, BRAF, PIK3CA*), NF- κ B pro-survival and death pathways (*FADD, BIRC2/3, CASP8, RIPK4*). Lastly, our findings revealed a link between the Notch and NF- κ B signal pathways and expression of several of their key components, Notch1, IKK α , RELA and RELB, contributing to the maintenance of the CD133⁺ cancer stem cell population and its stem-like phenotypic features (Figure 6E).

We uncovered a NOTCH cSCC cancer stem cell gene signature and showed that NOTCH plays a role in maintaining the CD133⁺ cell population. Blocking NOTCH by either of two different γ -secretase inhibitors, or NOTCH1 by siRNA, reduced the percentage of CD133⁺ SCC cells, and their capacity for clonogenic sphere formation, which are established surface and phenotypic markers of CSC/TIC (3). The NOTCH pathway plays a key role in cell-cell communication via interaction of membrane bound JAG ligands and Notch receptors (27). The NOTCH receptor family is comprised of 4 family members (27). In HNSCC and cSCC, inactivating and missense mutations in NOTCH1 and 2 have been observed in a subset of tumors, suggesting that the NOTCH may function as a tumor suppressor in those contexts (8, 9, 28, 29). Complexes of NICD with canonical Notch transcriptional co-factors can activate HES1/HEY1 repressors that trigger epithelial differentiation. Intriguingly, differential expression of HES1/HEY1 was not detected among CD133⁺ cSCC overexpressing NOTCH1 and other signal components, and there was decreased expression of key canonical transcription factor RBPJ. These findings are consistent with recent studies in human HNSCC, where tumors expressing wild-type NOTCH1 included subsets without evidence of HES1/HEY1 activation, and where NOTCH inhibitors reduced proliferation, through an unknown mechanism (15).

In this regard, we provide evidence that NOTCH can promote non-canonical cross-activation of Nuclear-Factor- κ B (NF- κ B/REL), a family of signal activated transcription factors which are often aberrantly activated in cSCC, HNSCC, and cervical SCC (25,30–33). The activation of canonical NF- κ B1/RELA and alternate NF- κ B2/RELB transcription factors is mediated by Inhibitor-kappaB Kinase (IKK) $\alpha/\beta/\gamma$ or IKK α complexes, respectively (25). Our profiling of CD133⁺ cSCC cells unveiled increased expression of multiple components of the NOTCH and NF- κ B signaling pathways, including key components NOTCH1, IKK α (CHUK), RELA, and RELB (Figure 6E). We showed that inhibition of Notch by γ -secretase or Notch1 by siRNA functionally partially inhibits NF- κ B reporter and nuclear DNA binding of NF- κ B p52, RelA and RelB along with the CD133⁺ phenotype, and that knocking down overexpressed IKK α , RelA and RelB similarly inhibited the CD133⁺ phenotype. The partial reduction of NF- κ B activity by inhibition of Notch signaling is expected, because Notch are not the only molecules that directly modulate NF- κ B activity in SCC (8, 25).

Interestingly, our study also establishes that IKK α is an important regulator for maintaining the CD133⁺ CSC population in cSCC cells. Notably, Notch1 and IKK α have previously been shown to physically and functionally interact in normal keratinocytes and cervical SCC cells (33,34). Notch1 knockdown modulated TNF α -induced NF- κ B activity and promoter binding in cervical SCC cells (33) similar to our findings for Notch γ -secretase pharmacologic inhibition of TNF-induced NF- κ B activity in cSCC cells. In cervical SCC, Notch1 and IKK α also promoted chemotherapeutic resistance, a feature often attributed to CSC. Our data indicate that wild type or activating phosphor-acceptor site mutants for TNF-mediated IKK α activation increase CD133⁺ cells, whereas inactivated IKK α mutants did not, implicating the role of IKK α in the expansion of CD133⁺ CSC. Further supporting this, IKK α colocalized in CD133⁺ cells in malignant epithelia of cSCC tumors. Besides its role in NF- κ B activation, IKK α may also function as a co-factor to regulate other key transcription factor genes that control cell proliferation and differentiation, such as E2F1/Bmi1 (35). Intriguingly, IKK α has also been reported to bind the promoter and reciprocally repress Notch canonical target gene HES1 (36), potentially helping to explain the lack of HES1 induction among Notch related genes we detected, and the paradoxical role of Notch1 and IKK α in promoting CSC expansion without triggering Notch-mediated terminal differentiation.

Although the involvement of the IKK-NF- κ B signaling pathway in promoting SCC has been demonstrated (25,30–33), its role in the maintenance of a CD133⁺ CSCs population in cSCC has not previously been reported. Our gene expression profiling and functional analysis reveals the important role of increased NF- κ B signaling pathway and key transcription factors RELA and RELB in promoting CD133⁺ CSCs in cSCC. Our finding that elevated NF- κ B signaling is important to maintain CSCs is supported by microarray expression profiling in human embryonic stem cell (ESC), which demonstrates that RelA is essential to maintain ESC pluripotency (37).

The evidence for altered expression of components of the PI3K/AKT/mTOR, MAPK, FGF, ERBB and STAT signaling pathways is consistent with evidence for frequent genomic alterations activating these pathways in HNSCC and cSCC (38). These pathways are also important for controlling the self-renewal and growth of cancer stem cells (39–43). PI3K and MAPK signaling is implicated in SCC growth and CSC (44–48). The regulatory role of STAT pathway on CSCs self-renewal was reported in HNSCC (49). Hence, our profiling data suggest that multiple signaling pathways may contribute to the stemness signature in cSCC. Consistent with this, we observed only partial inhibition of CSC phenotypes when blocking NOTCH and IKK-NF- κ B signal. Thus, combination therapy inhibiting CSCs may be more effective, since pharmacologic inhibition of one pathway may only partially reduce CD133⁺ cell population.

Several stemness makers have been studied in SCC and other cancers, including CD44, ALDH1, and CD133 (3, 4, 5, 50). Among these, CD44 and ALDH1 are expressed in relatively broader populations in tumor and cell lines, and additional markers or criteria are often used to enrich and characterize the CSC sub-populations. In our prior (3) and current study, the CD133⁺ population showed relatively lower expression of CD44, and <1.5 fold difference in ALDH1 compared with the CD133⁻ population. As ALDH1, CD44 and

CD133 populations do not always overlap, they may contain distinct stem cell sub-populations. Our prior and current study of skin SCC tumors established that the relatively rare CD133⁺ population is a biologically and therapeutically relevant subset enriched for TIC and sphere forming stem phenotypes. While our studies did not preclude that there are CSC bearing overlapping or distinct markers, this study explores and experimentally validates a role for NOTCH and IKK-NF- κ B signaling in the CD133⁺ population CSC phenotype. This study provides information about gene expression patterns and signaling pathways for this CSC population in human primary cSCC, which could help identify potential drug targets for CSC to complement current therapies for cSCC.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This study has been supported by the Intramural Research Program of the National Institute of Health, National Cancer Institute, The National Institute on Deafness and Other Communication Disorders (X. Quan, J. Coupar, S. Lee, Z. Chen, and C. Van Waes were supported by intramural projects ZIA-DC-000016, ZIA-DC-000073, ZIA-DC-000074, which were awarded to C. Van Waes). We specially thank Dr. Jonathan C. Vogel, who is deceased, for his guidance in initiating this project and helpful discussions and comments on the purification of CD133⁺ cell subpopulation and the design and setup of microarray experiment. We thank Dr. Mark Udey in the Dermatology Branch, National Cancer Institute, NIH, Bethesda, Maryland for many comments and strong support for the project. We thank Dr. Thomas Hornyak, in Dermatology Department, University of Maryland, School of Medicine, Baltimore, Maryland for his comments and support. We thank Dr. William G. Telford, in Experimental Transplantation and Immunology Branch, National Cancer Institute, NIH, Bethesda for his advice on cell flow cytometry experiments. This study utilized the high-performance computational capabilities of the Biowulf Linux cluster at the National Institutes of Health, Bethesda, MD. (<http://biowulf.nih.gov>).

References

1. Kauvar AN, Arpey CJ, Hruza G, Olbricht SM, Bennett R. Consensus for Nonmelanoma Skin Cancer Treatment, Part II: Squamous Cell Carcinoma, Including a Cost Analysis of Treatment Methods. *Dermatol Surg* 2015;41(11):1214–40 doi 10.1097/DSS.0000000000000478. [PubMed: 26445288]
2. Phillips RL, Ernst RE, Brunk B, Ivanova N, Mahan MA, Deanehan JK, et al. The genetic program of hematopoietic stem cells. *Science* 2000;288(5471):1635–40. [PubMed: 10834841]
3. D'Angelo RC, Wicha MS. Stem cells in normal development and cancer. *Prog Mol Biol Transl Sci* 2010;95:113–58 doi 10.1016/B978-0-12-385071-3.00006-X. [PubMed: 21075331]
4. Patel GK, Yee CL, Terunuma A, Telford WG, Voong N, Yuspa SH, et al. Identification and characterization of tumor-initiating cells in human primary cutaneous squamous cell carcinoma. *J Invest Dermatol* 2012;132(2):401–9 doi 10.1038/jid.2011.317. [PubMed: 22011906]
5. Zhang Q, Shi S, Yen Y, Brown J, Ta JQ, Le AD. A subpopulation of CD133(+) cancer stem-like cells characterized in human oral squamous cell carcinoma confer resistance to chemotherapy. *Cancer Lett* 2010;289(2):151–60 doi 10.1016/j.canlet.2009.08.010. [PubMed: 19748175]
6. Eramo A, Lotti F, Sette G, Pilozi E, Biffoni M, Di Virgilio A, et al. Identification and expansion of the tumorigenic lung cancer stem cell population. *Cell Death Differ* 2008;15(3):504–14 doi 10.1038/sj.cdd.4402283. [PubMed: 18049477]
7. Durinck S, Ho C, Wang NJ, Liao W, Jakkula LR, Collisson EA, et al. Temporal dissection of tumorigenesis in primary cancers. *Cancer Discov* 2011;1(2):137–43 doi 10.1158/2159-8290.CD-11-0028. [PubMed: 21984974]
8. Pickering CR, Zhou JH, Lee JJ, Drummond JA, Peng SA, Saade RE, et al. Mutational landscape of aggressive cutaneous squamous cell carcinoma. *Clin Cancer Res* 2014;20(24):6582–92 doi 10.1158/1078-0432.CCR-14-1768. [PubMed: 25303977]

9. Cancer Genome Atlas N. Comprehensive genomic characterization of head and neck squamous cell carcinomas. *Nature* 2015;517(7536):576–82 doi 10.1038/nature14129. [PubMed: 25631445]
10. Hoadley KA, Yau C, Wolf DM, Cherniack AD, Tamborero D, Ng S, et al. Multiplatform analysis of 12 cancer types reveals molecular classification within and across tissues of origin. *Cell* 2014;158(4):929–44 doi 10.1016/j.cell.2014.06.049. [PubMed: 25109877]
11. Ivanova NB, Dimos JT, Schaniel C, Hackney JA, Moore KA, Lemischka IR. A stem cell molecular signature. *Science* 2002;298(5593):601–4 doi 10.1126/science.1073823. [PubMed: 12228721]
12. Hofmann WK, de Vos S, Komor M, Hoelzer D, Wachsman W, Koeffler HP. Characterization of gene expression of CD34+ cells from normal and myelodysplastic bone marrow. *Blood* 2002;100(10):3553–60 doi 10.1182/blood.V100.10.3553. [PubMed: 12411319]
13. Sperger JM, Chen X, Draper JS, Antosiewicz JE, Chon CH, Jones SB, et al. Gene expression patterns in human embryonic stem cells and human pluripotent germ cell tumors. *Proc Natl Acad Sci U S A* 2003;100(23):13350–5 doi 10.1073/pnas.2235735100. [PubMed: 14595015]
14. Ramalho-Santos M, Yoon S, Matsuzaki Y, Mulligan RC, Melton DA. “Stemness”: transcriptional profiling of embryonic and adult stem cells. *Science* 2002;298(5593):597–600 doi 10.1126/science.1072530. [PubMed: 12228720]
15. Sun W, Gaykalova DA, Ochs MF, Mambo E, Arnaoutakis D, Liu Y, et al. Activation of the NOTCH pathway in head and neck cancer. *Cancer Res* 2014;74(4):1091–104 doi 10.1158/0008-5472.CAN-13-1259. [PubMed: 24351288]
16. Radtke S, Gorgens A, Kordelas L, Schmidt M, Kimmig KR, Koninger A, et al. CD133 allows elaborated discrimination and quantification of haematopoietic progenitor subsets in human haematopoietic stem cell transplants. *British journal of haematology* 2015;169(6):868–78 doi 10.1111/bjh.13362. [PubMed: 25819405]
17. Forget MA, Huon Y, Reuben A, Grange C, Liberman M, Martin J, et al. Stimulation of Wnt/ss-catenin pathway in human CD8+ T lymphocytes from blood and lung tumors leads to a shared young/memory phenotype. *PLoS One* 2012;7(7):e41074 doi 10.1371/journal.pone.0041074. [PubMed: 22859966]
18. Lai C, August S, Albibas A, Behar R, Cho SY, Polak ME, et al. OX40+ Regulatory T Cells in Cutaneous Squamous Cell Carcinoma Suppress Effector T-Cell Responses and Associate with Metastatic Potential. *Clin Cancer Res* 2016;22(16):4236–48 doi 10.1158/1078-0432.CCR-15-2614. [PubMed: 27034329]
19. Freeman A, Bridge JA, Maruthayanar P, Overgaard NH, Jung JW, Simpson F, et al. Comparative immune phenotypic analysis of cutaneous Squamous Cell Carcinoma and Intraepidermal Carcinoma in immune-competent individuals: proportional representation of CD8+ T-cells but not FoxP3+ Regulatory T-cells is associated with disease stage. *PLoS One* 2014;9(10):e110928 doi 10.1371/journal.pone.0110928. [PubMed: 25340823]
20. Xie Q, Wang H, Heilman ER, Walsh MG, Haseeb MA, Gupta R. Increased expression of enhancer of Zeste Homolog 2 (EZH2) differentiates squamous cell carcinoma from normal skin and actinic keratosis. *Eur J Dermatol* 2014;24(1):41–5. [PubMed: 24413452]
21. Adhikary G, Grun D, Balasubramanian S, Kerr C, Huang JM, Eckert RL. Survival of skin cancer stem cells requires the Ezh2 polycomb group protein. *Carcinogenesis* 2015;36(7):800–10. [PubMed: 25969142]
22. Egloff AM, and Grandis JR. Molecular pathways: context-dependent approaches to Notch targeting as cancer therapy. *Clin Cancer Res* 2012;18(19):5188–95. [PubMed: 22773520]
23. Rheinwald JG and Beckett MA. Tumorigenic keratinocyte lines requiring anchorage and fibroblast support cultured from human squamous cell carcinomas. *Cancer Res.* 1981;41(5):1657–63. [PubMed: 7214336]
24. Saito N, Fu J, Zheng S, Yao J, Wang S, Liu DD, et al. A high Notch pathway activation predicts response to gamma secretase inhibitors in proneural subtype of glioma tumor-initiating cells. *Stem cells* 2014;32(1):301–12 doi 10.1002/stem.1528. [PubMed: 24038660]
25. Nottingham LK, Yan CH, Yang X, Si H, Coupar J, Bian Y, et al. Aberrant IKKalpha and IKKbeta cooperatively activate NF-kappaB and induce EGFR/AP1 signaling to promote survival and migration of head and neck cancer. *Oncogene* 2014;33(9):1135–47 doi 10.1038/ncr.2013.49. [PubMed: 23455325]

26. Kobori M, Yang Z, Gong D, Heissmeyer V, Zhu H, Jung YK, et al. Wedelolactone suppresses LPS-induced caspase-11 expression by directly inhibiting the IKK complex. *Cell Death Differ* 2004;11(1):123–30 doi 10.1038/sj.cdd.4401325. [PubMed: 14526390]
27. Katoh M, Katoh M. Notch signaling in gastrointestinal tract (review). *Int J Oncol* 2007;30(1):247–51. [PubMed: 17143535]
28. Wang WM, Zhao ZL, Ma SR, Yu GT, Liu B, Zhang L, et al. Epidermal growth factor receptor inhibition reduces angiogenesis via hypoxia-inducible factor-1alpha and Notch1 in head neck squamous cell carcinoma. *PLoS One* 2015;10(2):e0119723 doi 10.1371/journal.pone.0119723. [PubMed: 25723392]
29. Martin A, Salvador F, Moreno-Bueno G, Floristan A, Ruiz-Herguido C, Cuevas EP, et al. Lysyl oxidase-like 2 represses Notch1 expression in the skin to promote squamous cell carcinoma progression. *EMBO J* 2015;34(8):1090–109 doi 10.15252/embj.201489975. [PubMed: 25759215]
30. Duffey DC, Chen Z, Dong G, Ondrey FG, Wolf JS, Brown K, et al. Expression of a dominant-negative mutant inhibitor-kappaBalpha of nuclear factor-kappaB in human head and neck squamous cell carcinoma inhibits survival, proinflammatory cytokine expression, and tumor growth in vivo. *Cancer Res* 1999;59(14):3468–74. [PubMed: 10416612]
31. Kobiela A, Fuchs E. Links between alpha-catenin, NF-kappaB, and squamous cell carcinoma in skin. *Proc Natl Acad Sci U S A* 2006;103(7):2322–7 doi 10.1073/pnas.0510422103. [PubMed: 16452166]
32. Lun M, Zhang PL, Pellitteri PK, Law A, Kennedy TL, Brown RE. Nuclear factor-kappaB pathway as a therapeutic target in head and neck squamous cell carcinoma: pharmaceutical and molecular validation in human cell lines using Velcade and siRNA/NF-kappaB. *Ann Clin Lab Sci* 2005;35(3):251–8. [PubMed: 16081580]
33. Song LL, Peng Y, Yun J, Rizzo P, Chaturvedi V, Weijzen S, et al. Notch-1 associates with IKKalpha and regulates IKK activity in cervical cancer cells. *Oncogene* 2008;27(44):5833–44 doi 10.1038/onc.2008.190. [PubMed: 18560356]
34. Yugawa T, Narisawa-Saito M, Yoshimatsu Y, Haga K, Ohno S, Egawa N, et al. DeltaNp63alpha repression of the Notch1 gene supports the proliferative capacity of normal human keratinocytes and cervical cancer cells. *Cancer Res* 2010;70(10):4034–44 doi 10.1158/0008-5472.CAN-09-4063. [PubMed: 20442293]
35. Ammirante M, Kuraishy AI, Shalapour S, Strasner A, Ramirez-Sanchez C, Zhang W, et al. An IKKalpha-E2F1-BMI1 cascade activated by infiltrating B cells controls prostate regeneration and tumor recurrence. *Genes Dev* 2013;27(13):1435–40 doi 10.1101/gad.220202.113. [PubMed: 23796898]
36. Aguilera C, Hoya-Arias R, Haegeman G, Espinosa L, Bigas A. Recruitment of IkappaBalpha to the hes1 promoter is associated with transcriptional repression. *Proc Natl Acad Sci U S A* 2004;101(47):16537–42 doi 10.1073/pnas.0404429101. [PubMed: 15536134]
37. Armstrong L, Hughes O, Yung S, Hyslop L, Stewart R, Wappler I, et al. The role of PI3K/AKT, MAPK/ERK and NFkappabeta signalling in the maintenance of human embryonic stem cell pluripotency and viability highlighted by transcriptional profiling and functional analysis. *Hum Mol Genet* 2006;15(11):1894–913 doi 10.1093/hmg/ddl112. [PubMed: 16644866]
38. Li YY, Hanna GJ, Laga AC, Haddad RI, Lorch JH, Hammerman PS. Genomic analysis of metastatic cutaneous squamous cell carcinoma. *Clin Cancer Res* 2015;21(6):1447–56 doi 10.1158/1078-0432.CCR-14-1773. [PubMed: 25589618]
39. Pattabiraman DR, Weinberg RA. Tackling the cancer stem cells - what challenges do they pose? *Nat Rev Drug Discov* 2014;13(7):497–512 doi 10.1038/nrd4253. [PubMed: 24981363]
40. Vermeulen L, Snippert HJ. Stem cell dynamics in homeostasis and cancer of the intestine. *Nat Rev Cancer* 2014;14(7):468–80 doi 10.1038/nrc3744. [PubMed: 24920463]
41. Schneider MR, Yarden Y. The EGFR-HER2 module: a stem cell approach to understanding a prime target and driver of solid tumors. *Oncogene* 2015 doi 10.1038/onc.2015.372.
42. Zhou BB, Zhang H, Damelin M, Geles KG, Grindley JC, Dirks PB. Tumour-initiating cells: challenges and opportunities for anticancer drug discovery. *Nat Rev Drug Discov* 2009;8(10):806–23 doi 10.1038/nrd2137. [PubMed: 19794444]

43. Pardal R, Clarke MF, Morrison SJ. Applying the principles of stem-cell biology to cancer. *Nat Rev Cancer* 2003;3(12):895–902 doi 10.1038/nrc1232. [PubMed: 14737120]
44. Li Q, Rycak K, Chen X, Tang DG. Cancer stem cells and cell size: A causal link? *Semin Cancer Biol* 2015;35:191–9 doi 10.1016/j.semcancer.2015.07.002. [PubMed: 26241348]
45. Xia P, Xu XY. PI3K/Akt/mTOR signaling pathway in cancer stem cells: from basic research to clinical application. *Am J Cancer Res* 2015;5(5):1602–9. [PubMed: 26175931]
46. Mohan S, Vander Broek R, Shah S, Eytan DF, Pierce ML, Carlson SG, et al. MEK Inhibitor PD-0325901 Overcomes Resistance to PI3K/mTOR Inhibitor PF-5212384 and Potentiates Antitumor Effects in Human Head and Neck Squamous Cell Carcinoma. *Clin Cancer Res* 2015;21(17):3946–56 doi 10.1158/1078-0432.CCR-14-3377. [PubMed: 25977343]
47. Li H, Gao Q, Guo L, Lu SH. The PTEN/PI3K/Akt pathway regulates stem-like cells in primary esophageal carcinoma cells. *Cancer Biol Ther* 2011;11(11):950–8. [PubMed: 21467840]
48. Sun S, Liu S, Duan SZ, Zhang L, Zhou H, Hu Y, et al. Targeting the c-Met/FZD8 signaling axis eliminates patient-derived cancer stem-like cells in head and neck squamous carcinomas. *Cancer Res* 2014;74(24):7546–59 doi 10.1158/0008-5472.CAN-14-0826. [PubMed: 25320014]
49. Bourguignon LY, Earle C, Wong G, Spevak CC, Krueger K. Stem cell marker (Nanog) and Stat-3 signaling promote MicroRNA-21 expression and chemoresistance in hyaluronan/CD44-activated head and neck squamous cell carcinoma cells. *Oncogene* 2012;31(2):149–60 doi 10.1038/onc.2011.222. [PubMed: 21685938]
50. Pan Q, Li Q, Liu S, Ning N, Zhang X, Xu Y, et al. Concise Review: Targeting Cancer Stem Cells Using Immunologic Approaches. *Stem cells* 2015;33(7):2085–92 doi 10.1002/stem.2039. [PubMed: 25873269]

Figure 1. Isolated CD133⁺CD61⁻CD31⁻CD24⁻CD45⁻ cell subpopulation exhibits cancer stem cell (CSCs) features in human cutaneous SCC (hsSCC).

(A) Distribution of cell surface markers in human SCC cells isolated from tumor specimens. (B) FACS detection of CD133⁺ CSC population in hsSCC. The red square gate was used to quantify the CD133⁺ population after gating out the other four CD marker positive cells. (C) Purity detection after FACS sorting for CD133⁺ cells after sorting from CD133⁻ cells, and gated CD61⁻CD31⁻CD24⁻CD45⁻ cell subpopulation from hsSCC. (D) Cytokeratin staining for CD133⁺ cell subpopulation. 99% of CD133⁺ cells were pan-cytokeratin positive as shown in the right square gate in the right panel. (E) Human SCC CD133⁺ cells are enriched

for spheroid colony formation *in vitro*. Normal human keratinocytes grew as adherent monolayer on the culture dishes with NIH 3T3 feeder layer (left upper panel, 40X), whereas the dissociated cSCC cells grew as tethered spheres on the NIH 3T3 feeder layer (right upper panel, 40X). The lower two panels show enlarged spheroid images under microscope (100X). (F) CD133⁺ cells exhibit statistically significantly greater spheroid formation numbers than CD133⁻ ($p<0.05$) or unsorted SCC cells ($p<0.05$) when seeded the same number cells (1×10^3). (G) In vivo tumor formation assay in nude mice for comparison of transplant of different numbers of primary cSCC cells. Each group includes 15 mice with 5 mice each that received the indicated number of primary SCC tumor cells from unsorted (green bars), sorted CD133⁻ (yellow bars) and CD133⁺ (blue bars) cells, as described in Methods.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 2. Characteristics of transcriptional profile of CD133⁺ CSC subpopulation and its stem gene expression signature.

(A) Hierarchical clustering of altered expression of 80 up- or down-regulated stem gene signature in sorted CD133⁺ cells compared with CD133⁻ cells after gating for CD61⁻CD31⁻CD24⁻CD45⁻ population (from 15 tumor specimens using Affymetrics microarray (fold change >1.5, ANOVA analysis, $p < 0.05$). Section I: up-regulated gene clustering and section II: down-regulated gene clustering. Hierarchical cluster analysis of Notch (B), and NF-κB (C) pathways from 2089 significantly different expressed genes in CD133⁺ CSCs compared with CD133⁻ tumor cells. IPA analysis of the network of NF-κB and Notch pathway interaction (D), and NF-κB and Wnt pathway interaction (E). The solid line indicates direct interaction and dotted line indicates indirect interaction. The pink color

indicates up-regulated genes and the blue color indicates down-regulated genes. The different shapes indicate different function category of the molecules as indicated.

Figure 3. Validation of differentially expressed genes in CD133⁺ CSC population using Nanostring expression assay.

Totally significantly altered 149 genes involved in stem gene signature or signaling pathways were selected and validated using Nanostring nCounter gene expression assay. The consistently altered genes detected by both microarray and Nanostring are shown in NF- κ B, Notch, Wnt, and Shh pathways (A), EGFR/ERBB, FGF, MAPK, PI3K/AKT/MTOR, JAK/STAT, TP53/P73, TGF β pathways (B), and Cell cycle, Chromatin Regulation and other function categories (C). Blue bar: Microarray; pink bar: Nanostring nCounter assay.

Figure 4. Blocking Notch inhibits CD133⁺ cell population and reduces NF-κB reporter activity and p65, p52 and RelB nuclear activation.

(A) Relative CD133⁺ cell population in human skin SCC13 cells were treated with two different Notch inhibitors, DAPT or R04929097 in different doses and measured by FACS analysis. (B) The relative colony formation ratio in both SCC13 and UMSSC46 cells were examined after DAPT or R04929097 treatment. (C) The specific Notch1 siRNA were transfected either alone or in combination into SCC13 or UMSSC 46 cells for 24 hours, and proteins after knockdown were shown by Western blot (upper panel). The CD133⁺ cell population of SCC13 was examined by FACS. Non-targeting siRNA control was used as the

control and its CD133⁺ population was set as 100% (lower panel). (D) NF- κ B reporter activity of SCC13 cells was measured with Notch inhibitors DAPT or R04929097 treatment for 24, 48 and 72 hours. NF- κ B activity was stimulated with TNF- α (1 ng/ml) and normalized with β -gal. (E) Silencing Notch 1, Notch2, or combined Notch 1 and 2 siRNA partially decreases p65, p52 and RelB nuclear activation by binding assay. All results are shown as mean $\pm$ SD; * p <0.05, ** p <0.01, *** p <0.001. The experiments were done in triplicate.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 5. Modulation of CD133⁺ CSCs by IKK inhibitor and IKKα expression, which co-localized with CD133 expression in cSCC tissues.

(A) IKK inhibitor Wedelolactone inhibits NF-κB activity in SCC13 cells. Cells were transfected with NF-κB luciferase reporter and β-Gal plasmids and treated with Wedelolactone and TNF-α alone, or in combination for 24 and 48 hours. Red, no TNF-α; blue, TNF-α treatment. * p<0.05, compared with TNF-α treated, non-wedelolactone group; #p<0.05, compared with non-TNF-α treated group; (B) Wedelolactone reduces CD133⁺ cancer stem cell population in SCC13 cells. Cells were treated with wedelolactone in different doses for 24 hours, stained with CD133-PE antibody, and CD133⁺ population was analyzed with FACS. (C) IKKα (CHUK) siRNA exhibited high knockdown efficiency in both cytoplasmic and nuclear compartment of SCC13 cells. (D) IKKα (CHUK) knockdown

significantly decreased CD133⁺ cancer stem cell population both in SCC13 and UMSCC46 cells. (E) Overexpression IKK α (CHUK) active forms dramatically increases CD133⁺ cancer stem cell population in SCC13 cells. (F) Decreased CD133⁺ population caused by IKK α (CHUK) siRNA knockdown could be reversed by introducing wild type IKK α or IKK α active forms (EE). Cells were co-transfected with different active forms of IKK α (WT, EE, AA and KA) plus IKK α siRNA with control siRNA. CD133⁺ population were analyzed after 48 hours transfection. * $p < 0.05$, compared with introduced WT or EE with control siRNA groups; # $p < 0.05$, compared with vector with control siRNA group. (G) The frozen section of cSCC tissues were co-immunostained with anti-IKK α and CD133 primary antibodies followed by Alexa-Fluor- 488 (green) and Alexa-Fluor-555 (red) secondary antibodies. DAPI (blue) was used for visualizing nuclei, and HE staining is showing morphology of tumor specimens. Magnification, 200X. All Data represent mean and standard deviation (s.d.) of one representative of three independent experiments, and each conducted in triplicate.

Figure 6. Knockdown NF- κ B RelA and/or RelB inhibit CD133⁺ population and its colony/spheroid formation abilities, and the summary of Notch and NF- κ B pathway activation. (A) RelA and RelB siRNA exhibited high efficiency of siRNA knockdown. (B) Knockdown of RelA and/or RelB inhibited CD133⁺ population. After 48 hours transfection, CD133⁺ population was analyzed by FACS. (C) Knockdown of RelA or RelB inhibited colony and spheroid formation abilities. After transfection for 24 hours, cells were seeded for colony formation of SCC13 cells (left panel) and spheroid formation of UMSCC46 cells (right panel). Results were shown as mean $\pm$ s.d. and data represent three independent experiments. * $p < 0.05$; ** $p < 0.01$ and *** $p < 0.001$. (D) The images of UMSCC46 spheroids

with different siRNA knockdown after 14 days of culture (100X). (E) Schematic illustration of activated Notch enhances NF- κ B pathway activation in cSCC cancer stem cell population. The CD133⁺ cSCC cancer stem cell gene signature is demonstrated with activated Notch signaling which includes NOTCH1, NOTCH2 and JAG2 increased expressing. The activated Notching pathway then enhances CHUK(IKK α), one of the key upstream components of NF- κ B signaling pathway. The active CHUK(IKK α) forms complexes with other IKKs and translocate to the nucleus, where further increased the NF- κ B2 p52/RelB complexes activation and induces target gene expression. The increased NOTCH1 and NOTCH2 might transfer signal to the nucleus and regulate transcription factor MAML1 and RBPJ to control the downstream target gene expression.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript