Published in final edited form as:

J Immunol Methods. 2018 September ; 460: 93–100. doi:10.1016/j.jim.2018.06.014.

***In vitro* analysis of antigen induced T cell-monocyte conjugates by imaging flow cytometry**

Meseret Habtamu^{a,b}, Markos Abebe^b, Abraham Aseffa^b, Anne Margarita Dyrhol-Riise^{c,d,e}, Anne Spurkland^a, and Greger Abrahamsen^{a,*}

^aDepartment of Molecular Medicine, Institute of Basic Medical Sciences, University of Oslo, Norway

^bArmauer Hansen Research Institute, Addis Ababa, Ethiopia

^cDepartment of Infectious Disease, Oslo University Hospital, N-0424 Oslo, Norway

^dInstitute of Clinical Medicine, Faculty of Medicine, University of Oslo, N-0424 Oslo, Norway

^eDepartment of Clinical Science, Faculty of Medicine, University of Bergen, N-5020 Bergen, Norway

Abstract

There is a lack of suitable correlates of immune protection against *Mycobacterium tuberculosis* (*Mtb*) infection. T cells and monocytes play key roles in host immunity against *Mtb*. Thus, a method that allows assessing their interaction would contribute to the understanding of immune regulation in tuberculosis (TB). We have established imaging flow cytometer (IFC) based *in vitro* assay for the analysis of early events in T cell-monocyte interaction, upstream of cytokine production and T cell proliferation. This was achieved through short term stimulation of peripheral blood mononuclear cells (PBMC) from healthy Norwegian blood donors with *Mycobacterium bovis* Bacille Calmette-Guérin (BCG). In our assay, we examined the kinetics of BCG uptake by monocytes using fluorescently labeled BCG and T cell-monocyte interaction based on synapse formation (CD3/ TCR polarization). Our results showed that BCG stimulation induced a gradual increase in the proportion of conjugated T cells displaying NF- κ B translocation to the nucleus in a time dependent manner, with the highest frequency observed at 6 h. We subsequently tested PBMC from a small cohort of active TB patients ($n = 7$) and observed a similar BCG induced NF- κ B translocation in T cells conjugated with monocytes. The method allowed for simultaneous evaluation of T cell-monocyte conjugates and T cell activation as measured by NF- κ B translocation, following short-term challenge of human PBMC with BCG. Whether this novel approach could serve as a diagnostic or prognostic marker needs to be investigated using a wide array of *Mtb* specific antigens in a larger cohort of patients with different TB infection status.

*Corresponding author at: Institute of Basic Medical Sciences, University of Oslo, Pb 1105 Blindern, 0317 Oslo, Norway., greger.abrahamsen@medisin.uio.no (G. Abrahamsen).

Conflict of interest

The authors declare that there is no conflict of interest.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jim.2018.06.014>.

Keywords

Imaging flow cytometry; BCG; Monocyte; Conjugate; Nuclear factor kappa-B

---

1. Introduction

Prominent progress has been made in the fight against tuberculosis (TB) in the past decades. Yet, it is still among the leading infectious causes of morbidity and mortality worldwide with an estimated 10.4 million new cases and 1.8 million associated deaths in 2015 (WHO, 2016). The development of novel approaches for the control and elimination of TB is challenged by the lack of tools to better understand the mechanisms and outcome of the host-pathogen interaction.

T cell mediated immune response of the Th-1 type is widely regarded as protective against TB (Kaufmann, 2006). The Th-1 cytokine, interferon gamma (IFN- γ) is considered a key protective cytokine based on earlier studies, in both mice and humans, which showed that defects in the IFN- γ signaling pathways provoked higher susceptibility to *Mycobacterium tuberculosis* (*Mtb*) infection and severe disease (Cooper et al., 1993; Newport et al., 1996; Flynn et al., 2003). Thus, many *in vitro* studies have focused on determining the frequency of *Mtb* specific T cells secreting IFN- γ alone, in combination with other Th-1 cytokines; IL-2 and TNF α measuring their level of secretion, and extent of T cell proliferation as markers of protection (McShane et al., 2004; Beveridge et al., 2008; Hoft et al., 2008; Smith et al., 2016). On the other hand, studies have shown that cytokine levels do not necessarily correlate with protective immunity nor predict risk of progression to active TB disease (Pai et al., 2004; Elias et al., 2005; Majlessi et al., 2006; Mittrucker et al., 2007; Panteleev et al., 2017). Developing suitable assays for the evaluation of protective immunity, risk of disease progression or immunogenicity of a vaccine is a priority in TB research (Centis et al., 2017).

It has long been established that T cell activation requires direct interaction between T cell surface receptors (TCR) and cognate peptide-major histocompatibility complex (pMHC) on the surface of antigen presenting cells (APCs) such as monocytes, macrophages, dendritic cells and B cells (Rossjohn et al., 2015; Jakubzick et al., 2017). Efficient TCR and pMHC interaction triggers signaling from the TCR-CD3 complex. Together with signals from costimulatory molecules, this results in the reorganization of T cell cytoskeletal elements with polarization of TCR as well as other signaling molecules towards the contact site resulting in the formation of an immunological synapse (IS) (Grakoui et al., 1999). The IS formation ensured sustained intracellular signaling in T cells, which involves translocation of transcription factors from the cytoplasm to the nucleus (Kane et al., 2000). Nuclear factor kappa-B (NF- κ B) is a transcription factor that regulates various genes involved in the host immune response; including cell proliferation and cytokine production (review in (Lucas et al., 2004; Tripathi and Aggarwal, 2006)). In the resting state, NF- κ B is localized in the cytoplasm and translocates to the nucleus upon activation, where it regulates gene transcription. It has been shown that these early events are among the factors that influence the quality of host immune response (Grakoui et al., 1999; Kane et al., 2002). Thus, assessing T cell-APC interactions and NF- κ B nuclear translocation could be a better

estimate in monitoring efficiency of immune response against TB, than measuring individual effector molecules.

Here we describe the use of imaging flow cytometer (IFC) to simultaneously evaluate T cell-monocyte conjugates and T cell activation as measured by NF- κ B translocation, following short-term BCG challenge *in vitro*. We used PBMC from buffy coats of healthy Norwegian blood donors as well as from TB patients residing in Norway.

2. Materials and methods

2.1.1. Antibodies, reagents and antigens

Antibodies used include anti-CD3 (clone OKT3), anti-CD28 (clone CD28.2, Cat# 555725, BD Pharmingen), anti-CD3-FITC (clone UCHT1, Cat# 21620033, Immunotools), anti-CD14-APC (clone MEM-15, Cat# 21279146, Immunotools), anti-IFN- γ PerCP Cy-5.5 (Clone B27, Cat # 560704, BD biosciences), rabbit polyclonal IgG (C-20) anti-human NF- κ B primary antibody (SC-372R, Santa Cruz Biotechnology), Alexa Fluor 594, goat anti-rabbit IgG (H + L) secondary antibody (Cat# R37117, Molecular Probes, Invitrogen). Fluorescent reagents include LIVE/ DEAD Fixable Near-IR (Cat# L34976, Molecular Probes, Invitrogen), and DAPI nuclear stain (Cat# 00-4959-52, Thermo Fisher) and CellTrace Violet (CTV) (Cat# C34557, Molecular probes, Invitrogen). RPMI 1640 (Gibco, Life Technologies) complete medium supplemented with 10% fetal calf serum (FCS), 100 units/ml penicillin, 100 μ g/ml streptomycin (all from GIBCOBRL®, Life Technologies™), 1 M HEPES buffer solution (Gibco, Life Technologies), 100 mM Sodium-pyruvate (Gibco, Life Technologies), MEM NEAA (100x) (Gibco, Life Technologies), 2-Mercaptoethanol (Sigma), Mycobacterial antigens used include: PPD for *in vitro* use (Statens Serum Institute, Copenhagen, Denmark) and ds-Red fluorescent protein expressing BCG (strain Myc3305, provided by Dr. Brigitte Gicquel) (Abadie et al., 2005).

2.1.2. Cell isolation and culture

Peripheral blood mononuclear cells (PBMC) from buffy coats, Oslo University Hospital, Norway, were isolated in high numbers by density gradient using Lymph prep (Axis-shield PoC AS, Cat # 1114545, Oslo, Norway) according to the manufacturer's protocol, and cryopreserved at a final concentration of 10% DMSO and 20% FCS until analysis. Prior to experiments, PBMC were thawed, washed and re-suspended in RPMI 1640 complete medium supplemented with 10% FCS, and allowed to rest overnight at 37 °C and 5% CO₂. PBMC (1×10^6 cells/mL) were then incubated at 37 °C and 5% CO₂ with ds-Red expressing BCG, at a final concentration of 3×10^6 CFU/mL for 6 h. PBMC incubated in culture medium alone under the same conditions were used as unstimulated control. Cells were harvested after careful scraping and resuspension, and then transferred to microcentrifuge tubes for fluorochrome staining and acquired by IFC.

For the pilot experiment PBMC from treatment-naïve clinically and/ or bacteriologically confirmed patients with active TB (n = 7) were obtained from the Norwegian biobank for infectious diseases, Oslo University Hospital, Ulleval. The research biobank is approved by the Regional Ethics Committee (REK Helse-sør-øst 2008.173) and written informed consent

was obtained from all participants prior to sample collection. The culture condition for cells from patients was similar to the buffy coats mentioned above.

2.1.3. In vitro T cell proliferation and IFN- γ releasing T cells

PBMCs were thawed and rested as described above and fluorescently labeled using CTV, according to the manufacturer's instructions. CTV labeled cells were re-suspended in complete media to a concentration of 1×10^6 cells/mL, in the presence of IL-2 and cultured at 37° C in a 5% CO₂ atmosphere with the following antibodies or antigens added: plate bound anti-CD3 (5 μ g/ml) and anti-CD28 (1 μ g/ml), PPD (10 μ g/ml) or ds-Red BCG (3×10^6 CFU/ml).

Unstimulated PBMC were used to assess background proliferation. Cells were then harvested on days 0, 4, 5, 6 and fluorochrome stained. In the preparation of IFN- γ production assay, brefeldin A (Sigma-Aldrich) was added to the five day culture prior to harvest and incubated for 4 h. Following anti-CD3 surface staining, cells were fixed and IFN- γ intracellular staining was performed. Stained cells were acquired with a FACSCantoII flow cytometer using FACS Diva software (BD Biosciences) to determine CTV dye dilution on proliferating T cells and IFN- γ releasing T cells. Data analysis was performed using FlowJo version 10.0 (Tree Star Inc.).

2.1.4. Imaging flow cytometry

Throughout the protocol, wide bore pipette tips (VWR) were used. Harvested cells were washed once with cold FACS buffer (2% FCS, 0.1% sodium azide in PBS). To block nonspecific binding, cells were incubated with 10% heat inactivated human serum in FACS buffer for 10 min on ice and surface stained with anti-CD3-FITC and anti-CD14-APC antibodies by incubating for 20 min on ice and in the dark.

For detection of intracellular NF- κ B, cells were fixed in 2% paraformaldehyde for 10 min at room temperature and incubated with 0.1% Triton-X permeabilization buffer (Sigma, USA) for 5 min. Cells were then incubated with rabbit antihuman NF- κ B primary antibody for 30 min followed by AF594 labeled goat anti-rabbit secondary antibody together with DAPI nuclear stain for 30 min on ice and in the dark. Single colour controls were prepared from PBMC and used for compensation.

An Amnis Image stream MkII 12 channel system fitted with 4 lasers was used for acquisition (Amnis Corporation, Seattle, WA). All acquisitions were performed at 40 $\times$ magnification on a low stream flow rate setting using INSPIRE software (Amnis Corporation). Up to 100,000 events were acquired from each sample. Images acquired include bright field (Channels 01 and 09), CD3-FITC (Ch-02), ds-Red BCG (Ch-03), NF- κ B AF594 (Ch-04), DAPI (Ch-07), CD14-APC (Ch-11) and live-dead fixative (Ch-12). Cellular events were acquired by setting a gate using a scatter plot that allowed discriminating cells from speed beads, which were used as an internal calibrator for the machine. The gated cells were plotted as a histogram to exclude non-cellular images that are DAPI negative. Single colour controls of 10,000 events were acquired for each colour in every experiment with all relevant lasers "on" and with the bright-field illumination and scatter laser "off", to acquire spectral compensation samples.

2.1.5. Imaging flow cytometry data analysis

The imaging flow cytometry data was analyzed using the IDEAS v6.1 software (Amnis Corporation, Seattle, WA). A compensation matrix was created using 'raw image files' of single colour controls and the IDEAS compensation wizard. Events *in camera* focus were selected on the basis of 'Gradient RMS' feature. Following this, an 'Area' *versus* 'Aspect ratio' plot was used to identify cellular aggregates and single cells where single cells have higher aspect ratio while aggregates have lower aspect ratio (Ahmed et al., 2009). T cells and monocytes were defined within each image using individual "object masks" based on the respective fluorescence intensity of CD3 and CD14. Two cell conjugates were identified from the aggregates on the basis of low 'nuclear Aspect ratio' based on DAPI signal and two cell nuclear content. From the two cell nuclear content, those adjacent with one T cell were gated on the basis of high 'CD3 Aspect ratio intensity'. This subpopulation was further gated to identify subpopulation of one T cell conjugated with one monocyte based on high 'CD14 Aspect ratio intensity'. We used TCR/CD3 polarization at the interface between a T cell and conjugating monocyte as marker for IS formation (Ahmed et al., 2009). To define the synapse within a T cell-monocyte conjugate, 'interface' mask was generated within the conjugating T cell. The T cell-monocyte conjugate subpopulation was further separated by plotting ds-Red BCG fluorescent signal in CD14+ cells gated as BCG positive (BCG+) or BCG negative (BCG-). In order to determine NF- κ B nuclear translocation in T cells, a software generated 'morphology' mask based on the nuclear image was applied to the 'Similarity' feature comparing the intensities of NF- κ B and nuclear (DAPI) image, as previously described (George et al., 2006) within the T cells.

2.1.6. Statistical analysis

The GraphPad Prism 6 (GraphPad Software, La Jolla, CA, USA, www.graphpad.com) data analysis software was used to do statistical comparisons, summarizations and data presentation. Non-parametric, two-tailed tests were used for all comparisons. Two-way analysis of variance was applied to comparisons with multiple time points. The Wilcoxon matched-pairs test was used for matched data sets while Mann-Whitney *U* test to determine mean differences between TB cases and the healthy control group. P-values ≤ 0.05 were considered statistically significant.

3. Results

3.1.1. Fluorescent labeled BCG induced T cell proliferation in vitro

In order to assess the ability of ds-Red expressing BCG to induce *in vitro* lymphocyte proliferation, PBMC from Norwegian healthy blood donors ($n = 6$) were analyzed in a fluorescence dilution assay using CTV at different time points (days 0,4,5 and 6) as a benchmark for cellular immune response against tuberculosis (Magnani et al., 2000).

Overlays of representative histograms showing CD4+ T cells with CTV dye dilution (CTV^{Low} events) in response to BCG, PPD, anti-CD3/ CD28 or unstimulated cells (UNS), at various time points are shown in Fig. 1A. Proportion of proliferating CD4+ T cells in response to various stimulants at the indicated times are summarized in Fig. 1B as line graphs. Anti-CD3/CD28 stimulation, used as positive control, induced massive proliferation

of CD4⁺ T cells after 4 days of incubation, and reached close to 100% by day 6. PPD and BCG stimulation resulted in on average 47% and 36% CTV^{Low} CD4⁺ T cells at day 6, respectively (Fig. 1B). In parallel, we also analyzed the Th-1 cytokine IFN- γ secretion, measured at day 5 (Fig. 1C). We observed that PPD (positive control) and BCG stimulation induced IFN- γ production in 14% and 11% of CD4⁺ T cells, respectively (Fig. 1D). Taken together, our data indicated that presence of ds-Red expressing BCG in the cell culture induced a T cell response. We thus proceeded using the ds-Red BCG for further investigation.

3.1.2. IFC allowed monitoring the kinetics of BCG uptake and T cell-monocyte conjugate formation

We used TCR/CD3 polarization at the interface between T cells and monocytes as marker for immune synapse formation and NF- κ B nuclear translocation as downstream effect after treating PBMC from healthy blood donors with ds-Red BCG and analyzed by IFC (Fig. 2A). In order to define the optimal culture condition allowing simultaneous assessment of T cell-monocyte conjugates and NF- κ B translocation, BCG titration with concentration up to 4×10^6 CFU/ml (Supplementary Fig. 1) and kinetics experiment up to 24 h (Supplementary Fig. 2) were performed. The results showed a decline in the frequency of T cell-monocyte conjugates as the concentration increases, which was considerable at the highest concentration (Supplementary Fig. 1A). A similar effect was observed in the proportions of T cell-monocyte conjugates and CD14⁺ cells as incubation time extended (Supplementary Fig. 2A and B). Conversely, more NF- κ B translocation was induced in T cells with increased BCG concentration and incubation time. We thus limited further experiments to BCG concentration of 3×10^6 CFU/ml, which we found optimal.

T cell-monocyte conjugates were identified as outlined in the gating strategy (Fig. 2B) and described in detail in the methods section. Briefly, two cell conjugates were selected by their low aspect ratio of the nuclear images based on DAPI signal. Of the two cell conjugates subpopulation, events of one T cell engaged with one monocyte were identified by their high CD3 and CD14 aspect ratio plotted consecutively (Fig. 2B vi, vii and viii). An image gallery is presented showing individual images and T cell-monocyte conjugates (Fig. 3A). Immunological synapse (IS) formation was determined by applying 'Interface mask' with TCR/CD3 polarization at the contact site between the two cells as shown in the image gallery (Fig. 3A, last column) and gated as in Fig. 3B. Although there was a trend towards higher frequency of T cell-monocyte conjugates with polarized TCR/CD3 in BCG stimulated compared to unstimulated cells during the early hours of stimulation, it was terminated after 3 h incubation (Fig. 3C). We also examined conjugates formed between T cell and BCG infected monocyte based on ds-Red fluorescent detection (Fig. 3A, upper panel) and gated in Fig. 3D and found that the frequency of T cells conjugated with BCG positive monocytes increase over time, from 11% after 1 h reaching to 25% after 6 h (Fig. 3E).

3.1.3. NF- κ B translocation in T cells increased with time following in vitro BCG stimulation

Having found that monocytes readily take up BCG (Fig. 3), we moved on to analyze its impact on T cell activation. Taking advantage of the capacity of IFC to monitor intracellular localization of fluorescently labeled molecules, we examined NF- κ B nuclear translocation in T cells in response to BCG stimulation and the optimal time point for this was determined. Cells were probed with fluorescently labeled antibody to visualize NF- κ B and stained with DAPI to visualize the nucleus. The NF- κ B signal, the nuclear mask within the T cell along with their composite are presented in Fig. 4A. We applied 'Similarity feature' to quantify the level of NF- κ B nuclear localization based on correlation between the two images.

Representative similarity plot histogram overlay of cells treated with BCG, PMA/ionomycin (positive control) and unstimulated cells (negative control) is presented to show how subpopulation of T cells with NF- κ B translocation were identified (Fig. 4B). The proportion of T cells display NF- κ B translocation increased over time in response to BCG stimulation. The frequency of T cell conjugates displaying NF- κ B nuclear translocation was an average of 4% at 1 h and increased to 14% at 6 h of incubation (Fig. 4C). Thus, for subsequent experiments, we considered the 6 h incubation to be optimal for measuring T cell-monocyte conjugate formation and induced NF- κ B nuclear translocation.

3.1.4. T cell-monocyte conjugate formation in active TB cases following in vitro BCG stimulation

The majority of Norwegian healthy blood donors have been BCG vaccinated as teenagers. However, the BCG vaccination status of the blood donors tested here was unknown. In order to assess whether *Mtb*-infected individuals respond differently to BCG in the IFC conjugate assay, we challenged PBMC from confirmed active treatment naïve TB patients (n = 7) with BCG for 6 h and compared this to the results obtained using PBMCs from healthy blood donors (n = 6). Our data showed that there was no significant difference in the frequency of T cell-monocyte conjugates in response to BCG stimulation, neither in TB cases nor in healthy controls, compared to unstimulated control cells (Fig. 5A). In addition, no significant difference was observed among TB cases and healthy controls in the frequency of total BCG induced T cell-monocyte conjugates as well as the frequency of conjugates of T cell-BCG infected monocytes (Fig. 5B and C, respectively).

3.1.5. NF- κ B nuclear translocation in T cells from active TB cases

Once we identified the T cell-monocyte conjugates in IFC analysis, we proceeded to consider T cell conjugates for NF- κ B translocation analysis. Our data showed that the proportion of conjugates with NF- κ B translocation was significantly higher upon BCG stimulation both in TB cases and healthy controls than in the spontaneously formed conjugates of unstimulated cells (UNS) (Fig. 6A and B). A potential advantage of IFC evaluation of nuclear translocation of a cytosolic protein is the ability to detect and quantify this event in a heterogeneous population of cells at a single cell level or in the cell of interest interacting with another cell. When we compared NF- κ B nuclear translocation in single T cells to T cells which were engaged in conjugate with monocytes, we observed that the proportion of T cells displaying NF- κ B translocation was significantly higher in conjugated

T cells than in single T cells both in active TB patients and in controls (Fig. 6B). Although we observed a trend towards a higher proportion of T cells conjugated with BCG infected monocytes and displaying NF- κ B translocation in TB cases than in healthy controls, this difference did not reach statistical significance in this limited number of tested individuals (Fig. 6C).

4. Discussion

The immune response against *Mtb* is complex and involves innate and adaptive immune cells (Zuniga et al., 2012; Liu et al., 2017). Although studies have reported controversial findings, monitoring host immunity to TB mainly relies on assessing antigen induced T cell proliferation and cytokine secretion (Mittrucker et al., 2007; Fletcher et al., 2016; Sakai et al., 2016; Smith et al., 2016). Here we established an imaging flow cytometer (IFC) based *in vitro* assay to evaluate multiple parameters of the early events in T cell activation, upstream of IFN- γ production and T cell proliferation in primary human cells.

The IFC platform, used here, is being well recognized in its application to assess large numbers of rare events and for analyzing co-localization of intracellular molecules (McFarlin and Gary, 2017) and thus providing statistically robust results. In recent years, various applications of IFC have been reported in TB research; including assessment of phagocytosis, phagosome maturation and *Mtb* replication inside macrophage differentiated THP-1 cells (Johansson et al., 2015; Ranjbar et al., 2015).

Being interested in biological markers for susceptibility to TB, we set out to establish an IFC-method for analyzing the early steps of T cell response to TB antigens as a correlate of host immunity. If feasible, such a method could have the potential in providing information on T cell response not given by the currently available assays, such as lymphocytes proliferation, IFN- γ release assay or enumeration of antigen specific T cells using tetramer staining.

In our assay, we monitored T cell-monocyte interaction using TCR/ CD3 polarization as a marker for synapse formation. Using the IFC based assay, we also monitored antigen induced NF- κ B translocation in interacting T cells with an IS. We observed a clear effect of BCG stimulation in terms of higher frequency of NF- κ B translocation in T cell conjugates compared to spontaneously formed conjugates in un-stimulated cells, especially from three to six hour incubation. Although the effect we observed was significantly lower compared to that observed in T cells conjugated with monocytes, NF- κ B translocation was also induced in single T cells that were exposed to BCG, though no difference was measured between patient cohorts. This finding could be explained by Toll-like receptor (TLR) mediated response, which has been reported be expressed in T cells and directly influence effector function upon exposure to bacterial products (Reynolds and Dong, 2013).

The main aim of this work was to establish the method itself using PBMC from healthy Norwegian blood donors, many of whom had previously been immunized with BCG, and thus were expected to harbor T cells responsive to BCG. After the method was established, we also tested a small number of active TB cases. Although we noticed a trend towards an

increase in stimulation induced response in TB cases compared to Norwegian healthy controls, the number of individuals included here was too small for this difference to reach statistical significance.

IFC has previously been used to assess conjugates formed between T cells with anti-CD3 coated macrophages (Ahmed et al., 2009) or anti-CD3 coated beads (Burbach et al., 2008). Another group has established a method in defining immune synapse formed between purified CD4⁺ T cells and dendritic cells (Markey et al., 2015). Here, we established an IFC method to identify and analyze signaling events in antigen induced T cell-monocyte conjugates in PBMC based on TCR/CD3 polarization at the synapse. Overall, no significant difference was found in the proportions of T cell-monocyte conjugates between BCG stimulated and unstimulated cells. Although we observed an increase in conjugate formation at the early time points, we also experienced a gradual decline in the number of CD14⁺ cells over time upon BCG challenge (Supplementary Fig. 3), which potentially could affect the number of conjugates identified. A similar observation was reported by Shey and colleagues that *in vitro* BCG challenge lowers the frequency of CD14⁺ cell (monocytes) (Shey et al., 2012) where phagocytosis induced shedding from the cell surface, micropinocytosis and monocyte death has been reported as possible causes (Mollen et al., 2008; Webster et al., 2010).

To our knowledge, this is the first report on the use of PBMC from humans in an IFC based assay to monitor the interaction between antigen presenting cell and T cell in response to live pathogen challenge (ds-Red BCG). The use of primary cells from healthy donors and active TB patients without sorting particular cell population, or using any additional inducement in the culture system, has the added value of measuring antigen-induced effect with minimal external influence. Monocytes are relatively rare in the circulation, and manipulations to enrich for the number of monocytes including purification by sorting has been reported to cause phenotypic and functional alterations (Delirezh et al., 2013), which could possibly influence on the results.

In conclusion, we have established an IFC based method using heterogeneous cell population to monitor early signaling events in T cells following antigen stimulation leading to various effector functions. The method allowed for simultaneous evaluation of T cell-monocyte conjugates and T cell activation as measured by NF- κ B translocation, following short-term BCG challenge of PBMC samples. This high throughput setting could help to address issues related to tetramer staining in assessing antigen specific T cell sub-populations (Vollers and Stern, 2008; Rius et al., 2018). Characterizing various aspects of immune cell interaction, combined with the consequence on the responding effector cell, might be used to address the current challenge of identifying correlate immune bio-markers of protection in TB or evaluation of vaccine induced responses. To substantiate if this platform would be instrumental in the search for bio-markers of diagnostic and/or prognostic value; further investigation with a larger cohort including TB patients with active disease as well as latent infection will be required. Adding more parameters for identifying IS may help improve assay specificity and sensitivity, allowing for differentiation of TB immunity in clinical samples. Moreover, the issue of antigen specific response and loss of CD14⁺ signal

on monocytes that we observed with BCG stimulation could be addressed by using fluorescently labeled *Mtb* antigens or peptides.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This study was supported by grants from Johs Svanholms grant, Henrik Homans Mindes grant, University of Oslo, Quota scheme of the Norwegian State, and Armauer Hansen Research Institute (AHRI). AHRI receives core support from Norad, Sida and the government of Ethiopia. The ds-Red fluorescent BCG strain Myc3305 is the property of Institut Pasteur, 25 rue du Docteur Roux, France 75015.

References

- Abadie V, Badell E, Douillard P, Ensergueix D, Leenen PJ, Tanguy M, Fiette L, Saeland S, Gicquel B, Winter N, 2005 Neutrophils rapidly migrate via lymphatics after Mycobacterium bovis BCG intradermal vaccination and shuttle live bacilli to the draining lymph nodes. *Blood* 106, 1843–1850. [PubMed: 15886329]
- Ahmed F, Friend S, George TC, Barteneva N, Lieberman J, 2009 Numbers matter: quantitative and dynamic analysis of the formation of an immunological synapse using imaging flow cytometry. *J. Immunol. Methods* 347, 79–86. [PubMed: 19524586]
- Beveridge NE, Fletcher HA, Hughes J, Pathan AA, Scriba TJ, Minassian A, Sander CR, Whelan KT, Dockrell HM, Hill AV, Hanekom WA, McShane H, 2008 A comparison of IFN γ detection methods used in tuberculosis vaccine trials. *Tuberculosis* 88, 631–640 (Edinb). [PubMed: 18801705]
- Burbach BJ, Srivastava R, Medeiros RB, Gorman WE, Peterson EJ, Shimizu Y, 2008 Distinct regulation of integrin-dependent T cell conjugate formation and NF- κ B activation by the adapter protein ADAP. *J. Immunol* 181, 4840. [PubMed: 18802088]
- Centis R, D'Ambrosio L, Zumla A, Migliori GB, 2017 Shifting from tuberculosis control to elimination: where are we? What are the variables and limitations? Is it achievable? *Int. J. Infect. Dis* 56, 30–33. [PubMed: 27916675]
- Cooper AM, Dalton DK, Stewart TA, Griffin J, Russell D, Orme I, 1993 Disseminated tuberculosis in interferon gamma gene-disrupted mice. *J. Exp. Med* 178, 2243–2247. [PubMed: 8245795]
- Delirez N, Shojaeefar E, Parvin P, Asadi B, 2013 Comparison the effects of two monocyte isolation methods, plastic adherence and magnetic activated cell sorting methods, on phagocytic activity of generated dendritic cells. *Cell J* 15, 218–223 (Yakhteh). [PubMed: 24027662]
- Elias D, Akuffo H, Britton S, 2005 PPD induced in vitro interferon gamma production is not a reliable correlate of protection against Mycobacterium tuberculosis. *Trans. R. Soc. Trop. Med. Hyg* 99, 363–368. [PubMed: 15780343]
- Fletcher HA, Snowden MA, Landry B, Rida W, Satti I, Harris SA, Matsumiya M, Tanner R, O'Shea MK, Dheenadhayalan V, Bogardus L, Stockdale L, Marsay L, Chomka A, Harrington-Kandt R, Manjaly-Thomas ZR, Naranbhai V, Stylianou E, Darboe F, Penn-Nicholson A, Nemes E, Hatheril M, Hussey G, Mahomed H, Tameris M, McClain JB, Evans TG, Hanekom WA, Scriba TJ, McShane H, 2016 T-cell activation is an immune correlate of risk in BCG vaccinated infants. *Nat. Commun* 7, 11290. [PubMed: 27068708]
- Flynn J, Capuano S, Croix D, Pawar S, Myers A, Zinovik A, Klein E, 2003 Non-human primates: a model for tuberculosis research. *Tuberculosis* 83, 116–118. [PubMed: 12758200]
- George TC, Fanning SL, Fitzgerald-Bocarsly P, Medeiros RB, Highfill S, Shimizu Y, Hall BE, Frost K, Basiji D, Ortyen WE, Morrissey PJ, Lynch DH, 2006 Quantitative measurement of nuclear translocation events using similarity analysis of multispectral cellular images obtained in flow. *J. Immunol. Methods* 311, 117–129. [PubMed: 16563425]

- Grakoui A, Bromley SK, Sumen C, Davis MM, Shaw AS, Allen PM, Dustin ML, 1999 The immunological synapse: a molecular machine controlling T cell activation. *Science* 285, 221–227 (New York, N.Y.). [PubMed: 10398592]
- Hoft DF, Blazevic A, Abate G, Hanekom WA, Kaplan G, Soler JH, Weichold F, Geiter L, Sadoff JC, Horwitz MA, 2008 A new recombinant bacille Calmette-Guerin vaccine safely induces significantly enhanced tuberculosis-specific immunity in human volunteers. *J. Infect. Dis* 198, 1491–1501. [PubMed: 18808333]
- Jakubzick CV, Randolph GJ, Henson PM, 2017 Monocyte differentiation and antigen-presenting functions. *Nat. Rev. Immunol* 17, 349–362. [PubMed: 28436425]
- Johansson J, Karlsson A, Bylund J, Welin A, 2015 Phagocyte interactions with *Mycobacterium tuberculosis* — simultaneous analysis of phagocytosis, phagosome maturation and intracellular replication by imaging flow cytometry. *J. Immunol. Methods* 427, 73–84. [PubMed: 26476130]
- Kane LP, Lin J, Weiss A, 2000 Signal transduction by the TCR for antigen. *Curr. Opin. Immunol* 12, 242–249. [PubMed: 10781399]
- Kane LP, Lin J, Weiss A, 2002 It's all Rel-ative: NF-kappaB and CD28 costimulation of T-cell activation. *Trends Immunol* 23, 413–420. [PubMed: 12133805]
- Kaufmann SH, 2006 Envisioning future strategies for vaccination against tuberculosis. *Nat. Rev. Immunol* 6, 699–704. [PubMed: 16917508]
- Liu CH, Liu H, Ge B, 2017 Innate immunity in tuberculosis: host defense vs pathogen evasion. *Cell. Mol. Immunol* 14, 1–13. [PubMed: 27545072]
- Lucas PC, McAllister-Lucas LM, Nunez G, 2004 NF-kappaB signaling in lymphocytes: a new cast of characters. *J. Cell Sci* 117, 31–39. [PubMed: 14657271]
- Magnani ZI, Confetti C, Besozzi G, Codecasa LR, Panina-Bordignon P, Lang R, Rossi GA, Pardi R, Burastero SE, 2000 Circulating, *Mycobacterium tuberculosis*-specific lymphocytes from PPD skin test-negative patients with tuberculosis do not secrete interferon-gamma (IFN- γ) and lack the cutaneous lymphocyte antigen skin-selective homing receptor. *Clin. Exp. Immunol* 119, 99–106. [PubMed: 10606970]
- Majlessi L, Simsova M, Jarvis Z, Brodin P, Rojas M-J, Bauche C, Nouzé C, Ladant D, Cole ST, Sebo P, Leclerc C, 2006 An increase in antimycobacterial Th1-cell responses by prime-boost protocols of immunization does not enhance protection against tuberculosis. *Infect. Immun* 74, 2128–2137. [PubMed: 16552042]
- Markey KA, Gartlan KH, Kuns RD, MacDonald KPA, Hill GR, 2015 Imaging the immunological synapse between dendritic cells and T cells. *J. Immunol. Methods* 423, 40–44. [PubMed: 25967948]
- McFarlin BK, Gary MA, 2017 Flow cytometry what you see matters: enhanced clinical detection using image-based flow cytometry. *Methods* 112, 1–8. [PubMed: 27620330]
- McShane H, Pathan AA, Sander CR, Keating SM, Gilbert SC, Huygen K, Fletcher HA, Hill AV, 2004 Recombinant modified vaccinia virus Ankara expressing antigen 85A boosts BCG-primed and naturally acquired antimycobacterial immunity in humans. *Nat. Med* 10, 1240–1244. [PubMed: 15502839]
- Mittrucker HW, Steinhoff U, Kohler A, Krause M, Lazar D, Mex P, Miekley D, Kaufmann SH, 2007 Poor correlation between BCG vaccination-induced T cell responses and protection against tuberculosis. *Proc. Natl. Acad. Sci. U. S. A* 104, 12434–12439. [PubMed: 17640915]
- Mollen KP, Gribar SC, Anand RJ, Kaczorowski DJ, Kohler JW, Branca MF, Dubowski TD, Sodhi CP, Hackam DJ, 2008 Increased expression and internalization of the endotoxin co-receptor CD14 in enterocytes occurs as an early event in the development of experimental necrotizing enterocolitis. *J. Pediatr. Surg* 43, 1175–1181. [PubMed: 18558203]
- Newport MJ, Huxley CM, Huston S, Hawrylowicz CM, Oostra BA, Williamson R, Levin M, 1996 A mutation in the interferon- γ -receptor gene and susceptibility to mycobacterial infection. *N. Engl. J. Med* 335, 1941–1949. [PubMed: 8960473]
- Pai M, Riley LW, Colford JM, 2004 Interferon- γ assays in the immunodiagnosis of tuberculosis: a systematic review. *Lancet Infect. Dis* 4, 761–776. [PubMed: 15567126]
- Pantelev AV, Nikitina IY, Burmistrova IA, Kosmiadi GA, Radaeva TV, Amansahedov RB, Sadikov PV, Serdyuk YV, Larionova EE, Bagdasarian TR, Chernousova LN, Ganusov VV, Lyadova IV,

- 2017 Severe tuberculosis in humans correlates best with neutrophil abundance and lymphocyte deficiency and does not correlate with antigen-specific CD4 T-cell response. *Front. Immunol* 8.
- Ranjbar S, Haridas V, Jasenosky LD, Falvo JV, Goldfeld AE, 2015 A role for IFITM proteins in restriction of *Mycobacterium tuberculosis* infection. *Cell Rep* 13, 874–883. [PubMed: 26565900]
- Reynolds JM, Dong C, 2013 Toll-like receptor regulation of effector T lymphocyte function. *Trends Immunol* 34, 511–519. [PubMed: 23886621]
- Rius C, Attaf M, Tungatt K, Bianchi V, Legut M, Bovay A, Donia M, Thor Straten P., Peakman M, Svane IM, Ott S, Connor T, Szomolay B, Dolton G, Sewell AK, 2018 Peptide–MHC Class I tetramers can fail to detect relevant functional T cell clonotypes and underestimate antigen-reactive T cell populations. *J. Immunol* 200, 2263. [PubMed: 29483360]
- Rossjohn J, Gras S, Miles JJ, Turner SJ, Godfrey DI, McCluskey J, 2015 T cell antigen receptor recognition of antigen-presenting molecules. *Annu. Rev. Immunol* 33, 169–200. [PubMed: 25493333]
- Sakai S, Kauffman KD, Sallin MA, Sharpe AH, Young HA, Ganusov VV, Barber DL, 2016 CD4 T cell-derived IFN-gamma plays a minimal role in control of pulmonary *Mycobacterium tuberculosis* infection and must be actively repressed by PD-1 to prevent lethal disease. *PLoS Pathog* 12, e1005667. [PubMed: 27244558]
- Shey MS, Hughes EJ, de Kock M, Barnard C, Stone L, Kollmann TR, Hanekom WA, Scriba TJ, 2012 Optimization of a whole blood intracellular cytokine assay for measuring innate cell responses to mycobacteria. *J. Immunol. Methods* 376, 79–88. [PubMed: 22155193]
- Smith SG, Zelmer A, Blitz R, Fletcher HA, Dockrell HM, 2016 Polyfunctional CD4 T-cells correlate with in vitro mycobacterial growth inhibition following *Mycobacterium bovis* BCG-vaccination of infants. *Vaccine* 34, 5298–5305. [PubMed: 27622301]
- Tripathi P, Aggarwal A, 2006 NF- κ B transcription factor: a key player in the generation of immune response. *Curr. Sci* 90, 519 (BANGALORE).
- Vollers SS, Stern LJ, 2008 Class II major histocompatibility complex tetramer staining: progress, problems, and prospects. *Immunology* 123, 305–313. [PubMed: 18251991]
- Webster SJ, Daigneault M, Bewley MA, Preston JA, Marriott HM, Walmsley SR, Read RC, Whyte MK, Dockrell DH, 2010 Distinct cell death programs in monocytes regulate innate responses following challenge with common causes of invasive bacterial disease. *J. Immunol* 190 (185), 2968–2979 (Baltimore, MD).
- WHO, 2016 Global Tuberculosis Report World Health Organization.
- Zuniga J, Torres-Garcia D, Santos-Mendoza T, Rodriguez-Reyna TS, Granados J, Yunis EJ, 2012 Cellular and humoral mechanisms involved in the control of tuberculosis. *Clin. Dev. Immunol* 2012, 193923. [PubMed: 22666281]

Fig. 1.

Antigen induced lymphocyte proliferation and IFN- γ production *in vitro*: PBMC from healthy blood donors ($n = 6$) labeled with CellTrace Violet (CTV) were cultured for 6 days in the presence of PPD, ds-Red BCG, anti-CD3/CD28 or media alone (UNS). Cells were harvested at various time points, fluorochrome stained and analyzed by flow cytometer for CTV dilution. (A) Representative histogram overlays; 0, 4, 5 and 6 days *in vitro* are shown here with the gate indicating CTV_{Low} CD4⁺ events. (B) Frequency of CD4⁺ T cells with CTV dye dilution (CTV_{Low} events) in response to anti-CD3/CD28 (dotted line with triangles), PPD (dashed line with circles), ds-Red BCG (solid line with squares), or UNS (gray line with diamond) at days 0, 4, 5 & 6. Error bars show mean $\pm$ SD. (C) Representative dot plots of IFN- γ secreting CD4⁺ T cell. (D) Proportion of IFN- γ secreting CD4⁺ T cells in response to PPD (circles), ds-Red BCG (squares), anti-CD3/CD28 (inverted triangle) or UNS (triangles). Horizontal bars indicate median values.

Fig. 2.

Assay outline and gating strategy applied to identify single cells, conjugates and NF- κ B translocation in T cells: (A) Schematic diagram summarizing cell preparation and analysis. PBMC from healthy donors ($n = 6$) were incubated with ds-Red BCG (BCG) for 6 h, fluorochrome stained and analyzed by IFC. (B) Cells *in camera* focus were selected from all events on the basis of gradient RMS of the bright field image (i). Single cells and cellular aggregates were identified by plotting 'Area' versus 'Aspect ratio' in which events with higher 'Aspect ratio' are likely to be single cells while those with lower 'Aspect ratio' are aggregates (ii). Live cells were identified from the single cell sub-population based on live/dead stain (iii). Single T cells and monocytes were plotted from live cell sub-population based on CD3 and CD14 fluorescent intensities in Ch-2 and Ch-11, respectively (iv). As for single cells, live cells from cellular aggregates were identified on the basis of live/dead stain (v). Two-cell nuclei were defined by plotting 'Intensity of DAPI' against 'Aspect ratio intensity of DAPI' from the aggregates where those with low 'Aspect ratio' are gated (vi). From the subpopulation of cells with two nuclei, those with one T cell were identified by plotting 'Intensity of Ch-2 (CD3_FITC)' against 'Aspect ratio intensity of Ch-2' where those with high 'aspect ratio intensity' are gated (vii). Conjugates with one T cell and one monocyte were identified by plotting "intensity of Ch-11" (CD14_APC) against "Aspect ratio intensity of Ch-11" (viii), events with high 'Aspect ratio intensity' are gated as population of interest.

Fig. 3.

Kinetics of BCG uptake and T cell-monocyte conjugate formation. (A) Image gallery, from left to right; showing bright field (BF) images in the first column, followed by T cells and monocytes based on CD3_FITC (green), and CD14_APC (red) fluorescent signal intensities, respectively. Masked images are shown in the right hand side of the respective images. BCG infected monocytes were identified based on detection of ds-Red fluorescent signal within CD14+ cells. Merged imageries are shown as; T cell conjugated with ds-Red BCG positive monocyte (upper panel) or monocyte with no ds-Red BCG (lower panel). Last column presents DAPI images of T cell-monocyte conjugates with 'interface' mask. (B) Representative histogram overlay for TCR/CD3 polarization at 6 h incubation of cells with BCG (red) or left unstimulated (UNS) (black). (C) Frequency of T cell-monocyte conjugates following BCG challenge (solid line) or UNS (dashed line), monitored for 6 subsequent time points in an hour interval. (D) Representative histograms defining BCG positive and BCG

negative T cell-monocyte conjugates. (E), Proportion of ds-Red BCG positive conjugates relative to all T cell-monocyte conjugates in BCG stimulated cells, analyzed sequentially for 6 time points. Line graphs represent mean values from 6 healthy blood donors and error bars show mean $\pm$ SD. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Fig. 4.

Kinetics of BCG induced NF- κ B nuclear translocation in T cells: PBMC from healthy donors ($n = 6$) were co-cultured with ds-Red BCG and analyzed as outlined in Fig. 2. (A). A gallery showing bright field (BF) images of two cell conjugates in the first column followed by T cell-monocyte conjugates as CD3+ (green)-CD14+ (red), ds-Red BCG in the third column followed by NF- κ B (golden brown) followed by DAPI mask. The last two columns present nuclear mask within T cell and merged images determining NF- κ B nuclear localization: 'high similarity' between fluorescent intensities of NF- κ B and DAPI images represents translocation (top panel) and 'low similarity' is for no translocation (bottom panel). The first and third rows represent T cells conjugated with BCG infected monocytes while second and fourth rows show no BCG. (B) Representative histogram overlay for NF- κ B translocation in T cells in response to BCG (black), PMA/ionomycin (red) or UNS (gray) at 6 h incubation. (C) Proportion of NF- κ B translocation in T cell conjugates at the indicated time points. Solid lines with circles represent BCG stimulated cells while dashed lines with squares indicated UNS. Line graphs represent mean values from 6 healthy blood donors. Error bars show mean $\pm$ SD. Statistical significance was determined by two-way analysis of variance. * $P < 0.05$, *** $P < 0.001$, **** $P < 0.0001$. (For interpretation of the

references to colour in this figure legend, the reader is referred to the web version of this article.)

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Fig. 5.

T cell-monocyte conjugates in active TB cases and healthy controls: T cell-monocyte conjugates were analyzed in confirmed TB cases (n = 7) and healthy controls (HC) (n = 6) following PBMC challenged for 6 h with BCG or media (UNS). (A) Frequency of T cell-monocyte conjugates with synapse relative to all T cell-monocyte conjugates in BCG stimulated and UNS pairs of TB cases and healthy controls. (B) Frequency of T cell-monocyte conjugates with synapse relative to all T cell-monocyte conjugates in TB cases and healthy controls upon BCG stimulation. (C) Proportion of ds-Red BCG positive T cell-monocyte conjugates in TB cases and healthy controls. Median values are indicated by horizontal lines (B and C).

Fig. 6.

NF- κ B nuclear translocation in T cells of active TB cases and healthy controls: PBMC from confirmed TB cases ($n = 7$) and healthy controls (HC) ($n = 6$) were incubated for 6 h. (A) Frequency of T cell conjugates displaying NF- κ B translocation in response to BCG stimulation in active TB cases and HC relative to all T cell-monocyte conjugates. (B) Proportion of single T cells (s-CD3; squares) compared with T cell conjugates (c-CD3; circles) displaying NF- κ B translocation in PBMC from active TB cases and HC in response to BCG stimulation. (C) Frequency of BCG positive (BCG+) T cell conjugates displaying NF- κ B translocation in cells from active TB cases and HC. Horizontal lines (B and C) indicate median values. Wilcoxon matched-pairs signed rank test was used to assess changes after stimulation and Mann-Whitney test was used to compare median differences between TB cases and HC, * $P < 0.05$.