Cochrane
Library

Cochrane Database of Systematic Reviews

Exercise for haemophilia (Review)

Strike K, Mulder K, Michael R

Strike K, Mulder K, Michael R.

Exercise for haemophilia.

Cochrane Database of Systematic Reviews 2016, Issue 12. Art. No.: CD011180.

DOI: [10.1002/14651858.CD011180.pub2](https://doi.org/10.1002/14651858.CD011180.pub2).

www.cochranelibrary.com

Exercise for haemophilia (Review)

Copyright © 2016 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

WILEY

TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
SUMMARY OF FINDINGS	4
BACKGROUND	11
OBJECTIVES	12
METHODS	12
RESULTS	15
Figure 1.	16
Figure 2.	18
Figure 3.	19
DISCUSSION	23
AUTHORS' CONCLUSIONS	25
ACKNOWLEDGEMENTS	26
REFERENCES	27
CHARACTERISTICS OF STUDIES	29
DATA AND ANALYSES	38
Analysis 1.1. Comparison 1 Exercise versus no intervention, Outcome 1 Colorado score.	38
Analysis 2.1. Comparison 2 Exercise A versus Exercise B, Outcome 1 Balance (Biodex Stability System).	39
Analysis 2.2. Comparison 2 Exercise A versus Exercise B, Outcome 2 Pain.	40
Analysis 2.3. Comparison 2 Exercise A versus Exercise B, Outcome 3 6-min walk: distance.	40
Analysis 2.4. Comparison 2 Exercise A versus Exercise B, Outcome 4 Range of motion: flexion.	40
Analysis 2.5. Comparison 2 Exercise A versus Exercise B, Outcome 5 Muscle strength.	40
Analysis 2.6. Comparison 2 Exercise A versus Exercise B, Outcome 6 Knee circumference.	41
Analysis 3.1. Comparison 3 Exercise versus Exercise plus electrophysical modality, Outcome 1 Colorado score.	42
Analysis 4.1. Comparison 4 Hydrotherapy versus no intervention, Outcome 1 Pain.	42
Analysis 4.2. Comparison 4 Hydrotherapy versus no intervention, Outcome 2 Range of motion.	42
ADDITIONAL TABLES	43
APPENDICES	43
CONTRIBUTIONS OF AUTHORS	45
DECLARATIONS OF INTEREST	45
SOURCES OF SUPPORT	45
DIFFERENCES BETWEEN PROTOCOL AND REVIEW	45
INDEX TERMS	45

[Intervention Review]

Exercise for haemophilia

Karen Strike¹, Kathy Mulder², Rojer Michael³

¹Department of Physiotherapy, Hamilton Health Sciences, McMaster Children's Hospital, Hamilton, Canada. ²Physiotherapy-Child Health, Health Sciences Centre, Winnipeg, Canada. ³School of Allied Health Sciences, Manipal Hospital, Bangalore, India

Contact address: Karen Strike, Department of Physiotherapy, Hamilton Health Sciences, McMaster Children's Hospital, 1200 Main Street West, Hamilton, ON, L8N 3Z5, Canada. strike@HHSC.CA.

Editorial group: Cochrane Cystic Fibrosis and Genetic Disorders Group.

Publication status and date: New, published in Issue 12, 2016.

Citation: Strike K, Mulder K, Michael R. Exercise for haemophilia. *Cochrane Database of Systematic Reviews* 2016, Issue 12. Art. No.: CD011180. DOI: [10.1002/14651858.CD011180.pub2](https://doi.org/10.1002/14651858.CD011180.pub2).

Copyright © 2016 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

Haemophilia is a bleeding disorder associated with haemorrhaging into joints and muscles. Exercise is often used to aid recovery after bleeds, and to improve joint function in the presence of arthropathy.

Objectives

Our objective was to systematically review the available evidence on the safety and effectiveness of exercise for people with haemophilia.

Search methods

We searched the Cochrane Cystic Fibrosis and Genetic Disorders Group's Coagulopathies Trials Register and electronic databases PubMed, OVID-Embase, and CINAHL. We hand searched abstracts from congresses of the World Federation of Hemophilia and the European Hematology Association, trial registries and the reference lists of relevant articles.

Date of the last search of the Cochrane Cystic Fibrosis and Genetic Disorders Group's Coagulopathies Trials Register: 14 December 2016.

Selection criteria

Randomized or quasi-randomized controlled studies comparing any exercise intervention considered relevant in haemophilia management including supervised, unsupervised, aquatic, strengthening, aerobic or cardiovascular, stretching, proprioceptive and balance training exercise programs in males of any age with haemophilia A or B of any severity (those with co-morbidities were not excluded).

Data collection and analysis

Two authors reviewed the identified abstracts to determine their eligibility. For studies meeting the inclusion criteria, full articles were obtained. The two authors extracted data and assessed the risk of bias. Any disagreements were resolved by discussion. The authors contacted study investigators to obtain any missing data.

Main results

Eight studies were included, which represented 233 males with all severities of haemophilia A and B, ranging in age from eight years to 49 years. Study duration ranged from four to 12 weeks. Exercise interventions varied greatly and included resistance exercises, isometric exercises, bicycle ergometry, treadmill walking and hydrotherapy; therefore, comparison between studies was difficult.

None of the studies measured or reported adverse effects from the interventions. None of the studies reported outcomes regarding bleed frequency, quality of life or aerobic activity. Overall risk of bias across all studies was assessed as unclear.

Exercise for haemophilia (Review)

Copyright © 2016 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

Very few studies provided sufficient information for comparison. None of the studies reported data that favoured the control group. One study reported that six weeks of resistance training improved joint health status (Colorado score) compared to controls. The addition of pulsed electromagnetic fields also improved ankle scores compared to exercises alone, but this was not seen in the elbows or knees.

Two studies reported statistically significant improvements in pain intensity after exercise interventions compared to controls. Hydrotherapy exercises produced significant decreases in pain compared to controls and land-based exercise groups.

Two studies found improvement in joint motion in the exercise group compared to controls. One study compared land- and water-based exercises; there was no difference in the range of motion between the two groups; however, the water-based exercise group did show improvement over the control group.

One study, comparing joint traction and proprioceptive neuromuscular facilitation for the elbow to a control group, showed no differences in biceps girth or strength after 12 weeks of intervention.

Some studies reported comparisons between interventions. In one study, treadmill training significantly improved balance in children compared to bicycle ergometry. Another study added partial weight bearing exercises to quadriceps exercises and showed improved walking tolerance.

Four studies evaluated quadriceps or hamstring strength (or both). The addition of bicycle ergometry and exercises with weights was more effective than static exercises and treadmill walking for strengthening knee flexors and extensors. Partial weight-bearing exercises through range were more effective than static and short arc exercises for improving knee extensor strength. The addition of treadmill walking to ultrasound, stretching and strengthening exercises showed increased peak torque of knee flexors and extensors and decrease in knee effusion.

The results should be interpreted with caution due to the quality of evidence (GRADE) as outlined in the summary of findings tables, which demonstrates that all but one of the outcomes assessed were rated as low or very low due to the small sample sizes and potential bias.

Authors' conclusions

These results must be considered with caution. There is a lack of confidence in the results due to the small number of included studies and the inability to pool the results due to the heterogeneity of outcome measures. Most exercise interventions produced improvement in one or more of the measured outcomes including pain, range of motion, strength and walking tolerance. Hydrotherapy may be more effective than land exercises for pain relief in adults. Functional exercises such as treadmill walking and partial weight bearing exercises seem to be more effective than static or short arc exercises for improving muscle strength. These findings are consistent with the many non-controlled intervention reports in the haemophilia literature. No adverse effects were reported as a result of any of the interventions. However, some groups used prophylactic factor prior to exercise and other groups studied only subjects with moderate haemophilia. Therefore, the safety of these techniques for persons with severe haemophilia remains unclear.

PLAIN LANGUAGE SUMMARY

Exercise for people with haemophilia

Review question

We reviewed the evidence about the safety and usefulness of exercise for men with haemophilia.

Background

Haemophilia is a group of disorders in which one of the blood clotting proteins does not work properly. It mainly affects men, although women can also be affected. People with haemophilia bleed into their joints and muscles which can lead to painful chronic arthritis.

Exercise is often used during recovery from a joint or muscle bleed and to maintain or improve a person's ability to function and participate in daily activities despite joint damage.

We wanted to know if exercise was safe (that is, exercise does not cause additional bleeding) and whether exercise was able to improve muscle strength, joint mobility, pain, balance, gait, fitness, and overall functioning.

Search date

The evidence is current to: 14 December 2016.

Study characteristics

We included eight studies with 233 male participants with haemophilia A or B (of any severity), aged eight to 49 years. Length of study ranged from four to 12 weeks.

Key results

Several types of exercise programs were studied, including stretching, strengthening with weights, exercise in water, treadmill walking, and exercise bicycle. Some studies compared participants who did one type of exercise with those who did another type of exercise; other studies compared an exercise group with a control group that did no exercise.

There were no data relating to our primary outcomes which indicated whether bleed frequency changed after an exercise program. There were no adverse effects measured or reported. Quality of life was not measured.

Regarding our secondary outcomes, improvements were seen in balance, joint health, and pain. Walking distance was the only functional status measured.

In an unplanned additional analysis, improvements were seen in the range of motion, biceps perimeter; strength, and knee circumference.

These small studies showed more improvements in pain, muscle strength and joint range of motion in exercise groups than in control groups. Studies that included functional activity, such as walking on a treadmill, showed more improvement than exercise alone. Exercise in water seems to be more effective than land exercise in relieving joint pain in adults.

Quality of the evidence

Four studies included only males with moderate haemophilia. Three studies included all severities of haemophilia and in one, participants used clotting factor prior to participating. Two studies included males with both haemophilia A and B; three studies did not specify type. Only one study limited their participants to those with severe haemophilia, and these also had osteoporosis. It is not clear whether the same results would be achieved if only males with severe haemophilia A were studied.

The results should be interpreted with caution due to the quality of the evidence; we judged that all but one of the outcomes assessed were low or very low quality, due to small sample sizes and potential bias.

SUMMARY OF FINDINGS

Summary of findings for the main comparison. Summary of findings - Exercise compared with no intervention

Exercise compared with no intervention for haemophilia

Patient or population: participants with haemophilia

Settings: outpatients

Intervention: exercise

Comparison: no intervention

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	No intervention	Exercise				
Adverse events	Outcome not reported				NA	
Quality of life	Outcome not reported				NA	
Joint health: Modified Colorado Score Follow up: 6 weeks	See comment	See comment	NA	25 (1 study)	⊕⊕⊕⊕ low ^{1,2}	Statistically significant improvements were demonstrated in the exercise group compared to the no intervention group in the joint health status of the elbows, MD -1.21 points (95% CI -2.14 to -0.28 points), knees, MD -3.42 points (95% CI -4.82 to -2.02 points) and ankles, MD -1.65 points (95% CI -2.51 to -0.79).
Pain: reduction of pain intensity by visual analogue scale Follow up: 12 weeks	See comment	See comment	NA	18 (1 study)	⊕⊕⊕⊕ very low ^{1,2,3}	Data were presented as number of joints (n = 30) rather than number of participants so could not be entered into the analysis. There was "observed improvement (P < 0.05)" in pain perception of the elbow in exercise group compared to the no intervention group.
Functional Status	Outcome not reported				NA	

Range of Motion: joint flexion and joint extension Follow up: 12 weeks	See comment	See comment	NA	18 (1 study)	⊕⊕⊕⊕ very low ^{1,2,3}	Data were presented as number of joints (n = 30) rather than number of participants so could not be entered into the analysis. There was "observed improvement (P < 0.05)" in flexion of the elbow in exercise group compared to the no intervention group but no statistically significant difference between treatment groups for joint extension.
Strength: bicep strength Follow up: 12 weeks	See comment	See comment	NA	18 (1 study)	⊕⊕⊕⊕ very low ^{1,2,3}	Data were presented as number of joints (n = 30) rather than number of participants so could not be entered into the analysis. There was no statistically significant difference between treatment groups.

*The basis for the **assumed risk** is provided in the comments. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).
CI: confidence interval; **MD:** mean difference; **NA:** not applicable.

GRADE Working Group grades of evidence

High quality: further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: we are very uncertain about the estimate.

1. Downgraded once due to potential risk of bias: limited details of study design provided making assessment of study quality difficult; presence of bias cannot be ruled out.
2. Downgraded once due to imprecision: evidence available from only a single study recruiting a small sample of participants.
3. Downgraded once due to applicability: the single study contributing evidence recruits only participants with severe haemophilia; results are not applicable to participants with mild or moderate haemophilia

Summary of findings 2. Summary of findings - Exercise A compared with Exercise B

Exercise A compared with Exercise B for haemophilia

Patient or population: participants with haemophilia

Settings: outpatients

Intervention: Exercise A

Comparison: Exercise B

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Exercise B	Exercise A				
Adverse events	Outcome not reported					
Quality of life	Outcome not reported					
Balance: Biodex stability parameters Follow up: 12 weeks	See comment	See comment	NA	30 (1 study)	⊕⊕⊕⊕ very low^{1,2,3}	Exercise B group (treadmill) significantly improved compared to Exercise A group (bicycle) in terms of overall stability index, MD 0.25 points (95% CI 0.19 to 0.31 points), anterior-posterior stability index, MD 0.35 points (95% CI 0.27 to 0.43 points) and medio-lateral stability index, MD 0.24 points (CI 95% 0.17 to 0.31 points).
Pain: reduction of pain intensity by visual analogue scale Follow up: 4 weeks	The mean reduction in pain from baseline was 1.7 on the VAS scale in the Exercise B group.	The mean reduction in pain from baseline was 0.8 lower (0.41 to 1.19 lower) on the VAS scale in the Exercise A group.	NA	27 (1 study)	⊕⊕⊕⊕ low^{1,2}	Exercise A is a land-based exercise program and Exercise B is an aquatic-based exercise program.
Functional Status: 6MWT (metres) Follow up: 6 weeks	The mean 6MWT was 32 metres in Exercise group B.	The mean 6MWT was 2.6 metres greater (0.08 metres to 5.12 metres greater) in Exercise group A.	NA	30 (1 study)	⊕⊕⊕⊕ very low^{1,2,3}	Exercise A is a partial weight bearing exercise program plus a quadriceps exercise training program and Exercise B is a quadriceps exercise training program alone.
Range of Motion: flexion and extension Follow up: 4 weeks	See comment	See comment	NA	27 (1 study)	⊕⊕⊕⊕ low^{1,2}	Exercise A is a land-based exercise program and Exercise B is an aquatic-based exercise program. No significant difference was demonstrated between the treatment groups; flexion, MD 0.20 degrees (95% CI -5.61 to 6.01 degrees) and extension MD -0.10 (95% CI -1.59 to 1.39).
Strength: flexors and extensors	See comment	See comment	NA	90	⊕⊕⊕⊕ moderate¹	Strength outcomes demonstrated a significant improvement for the study groups over the control

Follow up: 12 weeks (3 studies) groups in all outcome measures of strength including knee extensor and flexor strength.⁴

*The basis for the **assumed risk** is the mean control group risk or provided in the comments. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

6MWT: six-minute walk test; **CI**: confidence interval; **MD**: mean difference; **NA**: not applicable.

GRADE Working Group grades of evidence

High quality: further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: we are very uncertain about the estimate.

1. Downgraded once due to potential risk of bias: limited details of study design provided making assessment of study quality difficult; presence of bias cannot be ruled out.
2. Downgraded once due to imprecision: evidence available from only a single study recruiting a small sample of participants.
3. Downgraded once due to applicability: the single studies contributing evidence recruit only individuals with mild or moderate haemophilia; results are not applicable to participants with severe haemophilia.
4. Comparisons were partial weight bearing exercise program plus a quadriceps exercise training program (study group) compared to quadriceps exercise training program alone (control group), physical therapy program of stretching, strengthening and aerobic activity (control group) compared to the same program with the addition of bicycle ergometry (study group) and weight resistance and exercise therapy program including ultrasound, stretching and strengthening exercises (control group) compared to the same program plus treadmill training (study group). Also see [Table 2](#) for further details of interventions.

Summary of findings 3. Summary of findings - Exercise alone compared with exercise plus electrophysical modality

Exercise alone compared with exercise plus electrophysical modality for haemophilia

Patient or population: participants with haemophilia

Settings: outpatients

Intervention: exercise alone

Comparison: exercise plus electrophysical modality

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Exercise plus electrophysical	Exercise alone				

	modality					
Adverse events	Outcome not reported		NA			
Quality of life	Outcome not reported		NA			
Pain	Outcome not reported		NA			
Functional Status	Outcome not reported		NA			
Joint health: Modified Col-orado Score Follow up: 12 weeks	See comment	See comment	NA	25 (1 study)	⊕⊕⊕⊕ very low ^{1,2,3}	A statistically significant improvement was noted in the ankle scores, MD 0.90 points (95% CI 0.07 to 1.73 points) favouring the exercise plus electrophysical modality group over exercise alone group. There was no statistically significant difference between groups in elbow scores, MD 0.35 points (95% CI -0.70 to 1.40 points) or knee scores, MD 0.75 points (95% CI -0.47 to 1.97 points).

*The basis for the **assumed risk** is provided in the comments. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: confidence interval; **MD:** mean difference; **NA:** not applicable.

GRADE Working Group grades of evidence

High quality: further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: we are very uncertain about the estimate.

1. Downgraded once due to potential risk of bias: limited details of study design provided making assessment of study quality difficult; presence of bias cannot be ruled out.
2. Downgraded once due to imprecision: evidence available from only a single study recruiting a small sample of participants.
3. Downgraded once due to applicability: the single study contributing evidence recruits only participants with severe haemophilia; results are not applicable to participants with mild or moderate haemophilia.

Summary of findings 4. Summary of findings - Hydrotherapy compared with no exercise

Hydrotherapy compared with no exercise for haemophilia

Patient or population: participants with haemophilia

Settings: outpatients

Intervention: hydrotherapy

Comparison: no exercise

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	No exercise	Hydrotherapy				
Adverse events	Outcome not reported				NA	
Quality of life	Outcome not reported				NA	
Pain	Outcome not reported				NA	
Functional status	Outcome not reported				NA	
Range of motion: flexion and extension Follow up: 8 weeks	See comment	See comment	NA	20 (1 study)	⊕⊕⊕⊕ very low ^{1,2,3}	Data were presented via an analysis of covariance (post intervention scores adjusted for differences in pre intervention scores), so data could not be entered into the analysis. There was a statistically significant improvement in range of motion of the hydrotherapy group over the no exercise group.
Strength: knee flexor and extensor strength Follow up: 8 weeks	See comment	See comment	NA	20 (1 study)	⊕⊕⊕⊕ very low ^{1,2,3}	Data were presented via an analysis of covariance (post intervention scores adjusted for differences in pre intervention scores), so data could not be entered into analysis. There was a statistically significant improvement in both knee flexor and extensor strength of the hydrotherapy group over the no exercise group.

*The basis for the **assumed risk** is provided in the comments. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).
CI: confidence interval; **NA:** not applicable.

GRADE Working Group grades of evidence

High quality: further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: we are very uncertain about the estimate.

1. Downgraded once due to potential risk of bias: limited details of study design provided making assessment of study quality difficult and suspected selective reporting bias and post-intervention scores are not presented.
2. Downgraded once due to imprecision: evidence available from only a single study recruiting a small sample of participants.
3. Downgraded once due to applicability: the single study contributing evidence recruits only participants with moderate haemophilia; results are not applicable to participants with mild or severe haemophilia.

BACKGROUND

Description of the condition

Haemophilia is a congenital condition in which blood cannot clot due to the absence or deficiency of a clotting factor protein. The two most common types are haemophilia A (factor VIII deficiency) and haemophilia B (factor IX deficiency or Christmas disease) (WFH 2012). Haemophilia has an estimated frequency of approximately one per 10,000 births and there are approximately 400,000 people with haemophilia worldwide. Haemophilia A is more common than haemophilia B, representing 80% to 85% of the total haemophilia population (WFH 2012).

The frequency and severity of bleeding in haemophilia generally correlates to the clotting factor level, as measured by lab assay. People with severe haemophilia have clotting factor levels of less than 1% of normal and experience bleeding into joints and muscles, often in the absence of an identifiable haemostatic challenge, whereas individuals with mild haemophilia (levels of 5% to 40%) have severe bleeding only after major trauma or surgery (WFH 2012).

Haemophilia is caused by an inherited X-linked recessive trait. Males with the defective X chromosome are affected; females are carriers and may also exhibit signs of coagulopathy. Each child of a carrier has a 50% chance of inheriting the defective gene. In approximately one third of new cases, there is no family history, and the condition arises due to a spontaneous genetic mutation in either the mother or the son (WFH 2012).

Internal and prolonged bleeding are the main symptoms of haemophilia, and haemorrhagic episodes can occur anywhere in the body. For both haemophilia A and B, haemarthrosis accounts for 70% to 80% of all bleeding episodes; muscle bleeds account for 10% to 20% of bleeding episodes. Although any joint may be affected, hinge joints, particularly the ankles, knees and elbows, are the most frequently affected (WFH 2012).

Bleeding occurs from the small vessels in the synovial membrane. If the bleeding continues, the joint capsule becomes distended. This is accompanied by pain, loss of range of motion, muscle spasm or muscle inhibition, and loss of joint proprioception. An acute inflammatory response follows, and the blood is removed by synoviocytes. Articular cartilage is damaged directly by the blood as well as by the inflammatory enzymes (Rodríguez-Merchan 2012; Roosendaal 2008).

Repeated haemarthroses are common and, in the absence of adequate clotting factor replacement, frequently occur before the previous haemarthrosis has resolved. This creates a vicious cycle of bleed-inflammation-re-bleed, known as a target joint. The synovium is in a chronic state of inflammation, and becomes hypervascularized, hypertrophied and villous (De La Corte-Rodríguez 2013).

Recurrent hemarthroses eventually lead to epiphyseal changes and cartilage destruction with joint space narrowing, subchondral cysts, osteophyte formation, and incongruence of joint surfaces (Pettersson 1980).

Joint disease affects 90% of people with severe haemophilia and contributes the greatest cost and morbidity in the haemophilia population (Manco-Johnson 2004).

Muscle bleeds can occur in any muscle of the body, and usually result from a direct blow or a sudden stretch. Muscle bleeds are associated with pain, swelling, and loss of function. Bleeds into deep muscular compartments may result in neurovascular compromise, a serious complication. The deep flexor muscle groups, such as iliopsoas, forearm flexors and toe flexors are particularly at risk of developing compartment syndromes (Beyer 2010; WFH 2012).

Recurrent haemarthrosis and muscular bleedings have serious consequences on the musculoskeletal system and the functional health status and quality of life of individuals (Rodríguez-Merchan 2010).

Description of the intervention

For the purpose of this review, exercise will refer to a diverse set of interventions prescribed or planned by a health professional that includes conducting specific activities, postures or movements. Examples of such interventions are aerobic exercise, strengthening of specific muscles or groups of muscles, joint mobility and stretching exercises, and exercises designed to improve balance and proprioception, including exercise in water (Blamey 2010; Gomis 2009).

How the intervention might work

Exercise programs for people with haemophilia are usually designed and implemented to help manage the recovery after a hemarthrosis or a muscle bleed, to maintain or improve function in the presence of chronic arthropathy, or as a tool to help prevent frequent bleeding episodes.

The aims of exercise that are considered to be important for people with haemophilia are to (Gomis 2009; Iorio 2010; Negrier 2013):

- promote normal neuromuscular development;
- preserve or restore joint range of motion (ROM) and muscular flexibility;
- increase muscular strength and endurance;
- maintain or improve co-ordination and balance;
- maintain a healthy body weight thus decreasing stress on joints;
- promote development and maintenance of good bone density;
- improve aerobic conditioning;
- improve function and participation, and thus improve quality of life;
- reduce health risks associated with a sedentary lifestyle.

Why it is important to do this review

Over the past 50 years, the medical treatment for people with haemophilia has evolved from transfusions of whole blood to the use of fractionated blood products and to the use of recombinant technology. The availability and safety of coagulation factor products has allowed a philosophical change in medical care that previously limited activities and exercise for people with haemophilia because of concerns about the induction of bleeding (Gomis 2009; Riske 2007).

Since therapeutic exercise is an important component of the management of other forms of arthropathy (e.g. osteoarthritis, rheumatoid arthritis), it would appear logical that exercise would

be effective for people with haemophilia (Hunter 2009; Stenstrom 2003).

Thus, this review will seek to evaluate the safety and effectiveness of exercise in preventing bleeding episodes, maintaining and restoring joint and muscle function, and in maintaining the ability of the individual with haemophilia to participate in society.

OBJECTIVES

To systematically assess the available evidence on the safety and effectiveness of exercise interventions for people with haemophilia.

METHODS

Criteria for considering studies for this review

Types of studies

Randomized or quasi-randomized controlled studies.

Types of participants

Males of any age diagnosed with haemophilia A or B of any severity (severe, moderate or mild). Males with inhibitors or other co-morbidities were not excluded.

Types of interventions

For the purpose of this review, exercise will encompass "a diverse set of interventions prescribed or planned by a health professional that included conducting specific activities, postures or movements" (Blamey 2010). Any exercise intervention considered relevant to the management of haemophilia was included, such as aerobic exercise, strengthening of specific muscles, joint mobility and muscle stretching exercises, exercises designed to improve balance and proprioception, weight-bearing exercises, and exercise in water (hydrotherapy).

Types of outcome measures

Primary outcomes

1. Bleed frequency (number of major bleeds reported per year, month, or week)
2. Adverse events (e.g. bleed following exercise, worsening of symptoms)
3. Quality of life (e.g. assessed through self-administered questionnaires such as the 'Hemo-QOL and SF-36')

Secondary outcomes

1. Balance or proprioception (or both)
2. Aerobic activity
3. Joint health status measured via a validated outcome measure
4. Pain
5. Functional status

A post hoc analysis resulted in the addition of the following clinically relevant outcomes to the analysis. These outcomes were deemed important to healthcare providers and consumers as primary impairments that are assessed by musculoskeletal specialists and that may be impacted by an exercise intervention.

1. Range of motion

2. Biceps perimeter
3. Strength
4. Knee circumference

Search methods for identification of studies

There were no language or publication restrictions. Abstracts and full articles were included.

Electronic searches

We searched for relevant studies from the Cystic Fibrosis and Genetic Disorders Group's Coagulopathies Trials Register using the terms: haemophilia* AND exercise.

The Coagulopathies Trials Register is compiled from electronic searches of the Cochrane Central Register of Controlled Trials (CENTRAL) (updated each new issue of *The Cochrane Library*) and quarterly searches of MEDLINE and the prospective handsearching of one journal - *Haemophilia*. Unpublished work is identified by searching the abstract books of major conferences: the European Haematology Association conference; the American Society of Hematology conference; the British Society for Haematology Annual Scientific Meeting; and the Congress of the World Federation of Hemophilia. For full details of all searching activities for the register, please see the relevant section of the [Cochrane Cystic Fibrosis and Genetic Disorders Group's website](#).

Date of the most recent search of the Cochrane Cystic Fibrosis and Genetic Disorders Group's Coagulopathies Trials Register: 14 December 2016.

We performed additional searches on:

- PubMed (www.ncbi.nlm.nih.gov/pubmed) (searched from conception to 25 November 2014);
- Embase (Ovid) (searched from conception to 25 November 2014);
- CINAHL (EBSCO) (searched from conception to 30 April 2014).

We searched ClinicalTrials.gov (www.clinicaltrials.gov/); the International Clinical Trials Registry Platform (<http://apps.who.int/trialsearch/>); and the EU Clinical Trials Register (www.clinicaltrialsregister.eu). Date of most recent search of trials registers: 01 December 2014.

For full search strategies please refer to the appendices (Appendix 1).

Unpublished work was identified by the authors by searching the (online) abstract books of the following major conferences.

- European Haematology Association conference (2002 to 2013)
- American Society of Hematology conference (2006 to 2013)
- British Society for Haematology Annual Scientific Meeting (2007 to 2014)
- Congress of the World Federation of Hemophilia (1996 to 2014)
- Musculoskeletal Congresses of World Federation of Hemophilia (2005 to 2013)

For further details please refer to the appendices (Appendix 2).

Searching other resources

In addition, we searched the reference lists of all publications found by the above methods for any other potentially relevant studies.

Data collection and analysis

Selection of studies

Two authors (KS and KM) independently checked the titles and abstracts of the studies identified from the searches. We obtained the full text of all studies of possible relevance for assessment. We decided which studies fulfilled the inclusion criteria and resolved any disagreement by discussion. We also contacted study authors for clarification where necessary. Each of these two authors assessed studies published in languages other than English and included them in the review when feasible using English language abstracts, translation tools and services, or review by co-authors and colleagues familiar with the language of publication.

Data extraction and management

Two authors (KS and KM) independently extracted data from the included studies using forms provided by the Cochrane Cystic Fibrosis and Genetic Disorders Group.

We extracted population characteristics (patient population, source and setting, study inclusion criteria, symptom characteristics, severity, mean age, prophylaxis), intervention characteristics (description and types of exercise, duration and number of treatment sessions, intervention delivery type and co-interventions), comparison characteristics and outcome data. We extracted results for primary and secondary outcomes as change scores or final value scores for inclusion in the meta-analyses. We assessed the clinical relevance of each study with four items: participants described in detail to assess clinical comparability; interventions and treatment settings adequately described to allow repetition; clinically relevant outcomes measured and reported; and whether the likely treatment benefits are worth the potential harms. We also extracted data on adverse events, if reported, from each paper.

When information regarding any of the above was unclear, we contacted the study authors for further details. We entered the data into the Review Manager software and pooled where appropriate ([RevMan 2014](#)).

Assessment of risk of bias in included studies

The two authors assessed the risk of bias of each included study using Cochrane's risk of bias assessment tool ([Higgins 2011a](#)).

We assessed the following domains as having either a low, unclear or high risk of bias.

- Randomization ('low risk': randomization list generated using a computer, random number table, or similar methods; 'unclear risk': described as randomized, but no details given; 'high risk': non-random approach, e.g. alternation, use of case record numbers, dates of birth or day of the week).
- Concealment of allocation ('low risk': list from a central independent unit, on-site locked computer, identically appearing treatment; 'unclear risk': not described; 'high risk': if allocation sequence was known to, or could be deciphered

by the investigators who assigned participants or if an open allocation schedule was used.

- Blinding ('low risk': if participants, investigators and outcome assessors were blinded, or if any of these were not blinded but outcome assessment was judged not to influence the outcome; 'unclear risk': if this issue was not discussed; 'high risk': if none of the parties involved in the study were blinded).
- Incomplete outcome data ('low risk': if any withdrawals were described in full and were equal across groups; 'unclear risk': if insufficient information was given; 'high risk': if the missing data were likely to be directly related to the outcome or if they were uneven across groups).
- Selective outcome reporting.
- Other potential sources of bias.

Measures of treatment effect

Two authors (KS and KM) analysed the continuous outcomes by calculating the mean difference (MD) and the corresponding 95% confidence intervals (CI). For future updates, we will calculate the standardised mean difference (SMD) and corresponding 95% CIs if different scales of measurement have been used. Variation on measurement was reported as positive or negative providing information about effect. For example, a reduction in pain as measured by the visual analog scale (VAS) was reported as a negative effect as lower numbers on the VAS indicate improvement. Similarly, an improvement in range of motion (flexion) was reported as positive effect as higher numbers indicate improvement. For future updates, we will calculate the risk ratio (RR) and 95% CIs for dichotomous outcomes and for categorical data the results will be presented as a RR with 95% CI.

Since none of the included studies reported measures of intra-subject variability (e.g. within-group correlation co-efficients for the post versus baseline measures) we were unable to account for it in assessing variability of the variation over time. We measured the effect across randomization by comparing within-group differences, for which we calculated variability assuming independence. This likely led to a conservative approach (missing an effect where the effect was there, but not claiming a spurious effect when this was not true), which we considered an acceptable limitation. We will try to gather data from the study authors in order to properly update the calculation and publish an update if successful.

Unit of analysis issues

Cluster-randomized studies

If, for future updates, we identify cluster-randomized studies, we will include these in the analyses along with individually randomized studies. In an attempt to account for any unit of analysis error, we will use the methods described in the *Cochrane Handbook for Systematic Reviews of Interventions* using an estimate of the intra-cluster correlation co-efficient (ICC) derived from the study (if possible), from a similar (in design) study or from a study of a similar population ([Higgins 2011b](#)). If we use ICCs from other sources, we will report them and conduct sensitivity analyses to investigate the effect of variation in the ICC. If we identify both cluster-randomized studies and individually-randomized studies, we plan to pool the relevant information. If there is little heterogeneity between the study designs and the interaction between the effect of intervention and the choice of

randomization unit is considered to be unlikely, we will consider it reasonable to combine the results from both. If appropriate, we will also acknowledge heterogeneity in the randomization unit and perform a sensitivity or subgroup analysis to investigate the effects of the randomization unit.

Cross-over studies

No cross-over studies were identified in the literature search and therefore were not included in the review. Given that an effective washout period is almost impossible to achieve with an exercise program, in future versions of the review, if we identify cross-over studies, we will only analyse first-arm data (collected prior to the cross-over of the intervention).

In the first instance, we extracted outcome data with the unit of analysis being the participant (i.e. number of participants). However, for one study, 18 participants (nine in the intervention and nine in the control group) the results were reported at the unit of analysis of joints only (16 elbows included in the intervention group and 14 elbows included in the control group) (Cuesta-Barriuso 2013). No data reported using different units of analysis were pooled and the difference in unit of analysis must be considered when interpreting the results.

Dealing with missing data

We reported the numbers and reasons for dropouts and withdrawals in all intervention groups, if described, and also reported if the papers specified that there were no dropouts or withdrawals. We contacted the original Investigators for clarification on any missing information.

In the event that we were unable to contact the original authors or they were unable to supply missing data, we imputed missing variance scores using the mean variance from studies with similar populations. If data were reported as a median and interquartile range (IQR), we used the median to estimate the mean for studies with moderate to large sample sizes ($n > 25$); for studies with small sample sizes, we used the formula proposed by Hozo (Hozo 2005). When appropriate we calculated the standard deviation (SD) with the width of the IQR equivalent to 1.35 times the SD (Higgins 2002). In studies where a range was presented along with the median instead of an IQR, we estimated the SD to be one quarter of the range (Higgins 2011c).

For included studies, we noted levels of attrition. For all outcomes, we carried out analyses, as far as possible, on an intention-to-treat basis, i.e. we attempted to include all participants randomized to each group in the analyses, and analysed all participants in the group to which they were allocated, and regardless of whether or not they received the allocated intervention. We analysed missing data using the Review Manager software (RevMan 2014).

Assessment of heterogeneity

We assessed heterogeneity between studies by inspecting the forest plots and using the χ^2 test and I^2 statistic for heterogeneity with a statistical significance level of $P < 0.10$ and an interpretation of I^2 as follows:

- 0% to 40%: might not be important;
- 30% to 60%: may represent moderate heterogeneity;

- 50% to 90%: may represent substantial heterogeneity;

- 75% to 100%: considerable heterogeneity.

Assessment of reporting biases

For study protocols that were available, we compared these to the published reports. For primary outcomes, we were planning on investigating any potential reporting biases using a funnel plot; however, a sufficient number of studies were not available (Begg 1994). We performed comprehensive searches to locate all randomized controlled studies and also sought out all relevant unpublished studies to minimize reporting bias.

Data synthesis

Two authors (KS and KM) performed statistical analysis in accordance with Cochrane guidelines (Deeks 2011) using the Review Manager software (RevMan 2014). If there was no substantial heterogeneity (less than 50%) we used the fixed-effect model. In the presence of at least moderate heterogeneity (over 50%) we used the random-effects model and sensitivity analyses as described below to investigate the source of heterogeneity. We planned to analyse the different interventions separately and planned to group outcome data as follows: four weeks to three months; three months to six months; and longer than six months; however, we did not obtain sufficient data from the included studies.

Subgroup analysis and investigation of heterogeneity

We intended to undertake the following subgroup analyses:

1. exercise versus other interventions (including no intervention);
2. comparison of various modes of exercise in the prophylaxis group;
3. comparison of various modes of exercise in the on-demand group;
4. long-term and short-term effects of exercise on patients with haemophilia.

We did not find sufficient data to perform the pre-planned subgroup analysis.

Sensitivity analysis

If there are sufficient comparable studies, i.e. 10 or more, in future updates we will perform sensitivity analyses excluding studies with clearly inadequate allocation of concealment, randomization, blinding or incomplete outcome data (high risk of bias). As reported above, in reference to cluster randomized studies, if we use ICCs from other sources we will conduct sensitivity analyses to investigate the effect of variation in the ICC.

Summary of findings and quality of the evidence (GRADE)

In a post hoc change from the protocol, we have presented a summary of findings table for each comparison in the review (Summary of findings for the main comparison; Summary of findings 2; Summary of findings 3; Summary of findings 4).

1. Exercise compared to no intervention
2. Exercise A compared to Exercise B
3. Exercise alone compared to exercise plus electrophysical modality

4. Hydrotherapy compared to no exercise

The following outcomes were reported in all tables (chosen based on relevance to clinicians and consumers): quality of life; adverse events; functional status; and the following outcomes were reported in at least one table (again chosen based on clinical relevance to the comparison): pain; joint health; range of motion; strength; and balance.

We determined the quality of the evidence using the GRADE approach; and downgraded evidence in the presence of a high risk of bias in at least one study, indirectness of the evidence, unexplained heterogeneity or inconsistency, imprecision of results, high probability of publication bias. We downgraded evidence by one level if we considered the limitation to be serious and by two levels if very serious.

For clarity in the tables, where outcomes were presented using different measures (e.g. balance) or different domains (e.g. joint health according to modified Colorado score), a general statement is made in the table regarding the summary of findings for these outcomes and the evidence is graded based on all of the measures or subdomains combined.

RESULTS

Description of studies

See: [Characteristics of included studies](#); [Characteristics of excluded studies](#).

Results of the search

Database searches (PubMed, CINAHL and Embase) and searches of the Group's Coagulopathies Trials Register yielded 231 references. A search of a clinical trials registries identified one study. Handsearches of conference proceedings from the World and Musculoskeletal Congresses of the World Federation of Hemophilia identified 77 references. Searches of European, British and American Hematology meetings did not yield any results. A total of 22 additional references were identified from reference lists, for a total of 328 references.

After duplicates were removed, 250 potential references remained. Of these, 129 described patient education strategies, gait analysis, population characteristics, measures of participation in activity, and laboratory training 'exercises'.

The remaining 121 described some type of exercise intervention, but most were single case studies or had no control group. A total of 75 of these references described clinical studies; however, after inspection, only 18 studies (22 references) were relevant. These were assessed by both authors for eligibility: after full text review, eight studies were deemed eligible for the review, seven studies were excluded and three are awaiting assessment.

See figure 1 'Study Flow Diagram' ([Figure 1](#)).

Figure 1. Study flow diagram.

Included studies

Eight studies met the inclusion criteria. The number of participants per study were ranged from 18 to 48, comprising a total of 233 males (Abd-Elmonem 2014; Cuesta-Barriuso 2013; Eid 2014; Kargarfard 2013; Mazloun 2014; Mohamed 2015; Parhampour 2014; Zaky 2013). Note: there were an additional 13 participants in the Mazloun study but these related to a third non-randomized arm and have not been included in this review.

Design

All studies were randomized or quasi-randomized with control groups.

Sample sizes

Samples sizes ranged from 18 to 48 participants (four studies each had 30 participants, with the remaining four having 18, 20, 40 and 48 participants).

Location

All studies were conducted in a physical therapy clinic at either a university or a large hospital. Four studies were conducted at the University of Cairo (Abd-Elmonem 2014; Eid 2014; Zaky 2013; Mohamed 2015), three in Iran (Kargarfard 2013; Mazloun 2014; Parhampour 2014), and one in Spain (Cuesta-Barriuso 2013).

Participants

Five studies selected participants with haemophilia A (Abd-Elmonem 2014; Cuesta-Barriuso 2013; Eid 2014; Mazloun 2014; Parhampour 2014) and two of these also included participants with haemophilia B (Abd-Elmonem 2014; Cuesta-Barriuso 2013). Three authors did not specify type of haemophilia (Kargarfard 2013; Mohamed 2015; Zaky 2013). The participants in four studies had moderate haemophilia (Abd-Elmonem 2014; Eid 2014; Kargarfard 2013; Zaky 2013). Two studies included participants with all severities (Mazloun 2014; Mohamed 2015) and one study combined participants with mild and severe haemophilia (Cuesta-Barriuso 2013). One study selected participants with severe haemophilia and osteoporosis (Parhampour 2014). Four studies included only children (aged eight years to 14 years) (Abd-Elmonem 2014; Eid 2014; Mohamed 2015; Zaky 2013) while the other four included adults aged 20 years to 49 years (Cuesta-Barriuso 2013; Kargarfard 2013; Mazloun 2014; Parhampour 2014). Participants in two studies infused factor prior to participation in the exercise programs (Mazloun 2014; Parhampour 2014).

Interventions

Study periods ranged from four to 12 weeks. Interventions included: exercises in water (Kargarfard 2013); land-based exercises including resisted exercises with weights (Eid 2014; Mazloun 2014; Parhampour 2014); static exercises (Abd-Elmonem 2014; Zaky 2013); stretching exercises (Abd-Elmonem 2014; Eid 2014); bicycle ergometry (Eid 2014; Mohamed 2015); treadmill walking (Abd-Elmonem 2014); partial weight-bearing (Zaky 2013);

and proprioceptive neuromuscular facilitation (PNF) techniques (Cuesta-Barriuso 2013). Other non-exercise interventions included: ultrasound (Abd-Elmonem 2014); pulsed electromagnetic fields (Parhampour 2014); and joint traction (Cuesta-Barriuso 2013).

Outcomes

Primary outcomes

1. Bleed frequency (number of major bleeds reported per year, month, or week) - not measured
2. Adverse events (e.g. bleed following exercise, worsening of symptoms) - not measured
3. Quality of life - not measured

Secondary outcomes

1. Balance or proprioception (or both): Biodex stability system (Mohamed 2015)
2. Aerobic activity - not measured
3. Joint health status measured via a validated outcome measure: Modified Colorado Score (Parhampour 2014)
4. Pain (visual analog scale (VAS)) (Cuesta-Barriuso 2013; Mazloun 2014)
5. Functional status: Six Minute Walk test (Eid 2014)
6. Range of motion (Kargarfard 2013; Mazloun 2014)
7. Biceps perimeter (Cuesta-Barriuso 2013)
8. Strength: peak torque using Biodex Isokinetic Dynamometer (Abd-Elmonem 2014; Eid 2014; Kargarfard 2013) and Lafayette Manual Muscle Test (Zaky 2013)
9. Knee circumference (Abd-Elmonem 2014)

See also: 'Characteristics of included studies' table

Excluded studies

See: 'Characteristics of Excluded Studies' table (Characteristics of excluded studies).

We excluded seven studies; four studies had no control group (Greene 1983; Khriesat 2000; Mulvany 2010; Vallejo 2010) and the remaining three had control groups but participants were not randomized (Czepa 2013; Hilberg 2003; Von Mackensen 2012).

Studies awaiting classification

Three additional studies are awaiting classification and will be assessed for possible inclusion in the next update (Cuesta-Barriuso 2014; Cuesta-Barriuso 2015; Firoozabadi 2012). Further details are provided in the tables (Studies awaiting classification).

Risk of bias in included studies

We assessed the included studies for general risk of bias and overall, we assessed these as having a low risk of bias for some aspects and unclear for others. The authors' judgements of risk of bias for each study are outlined in each of the studies risk of bias tables and are summarized in the risk of bias summary tables (Figure 2; Figure 3).

Figure 2. Risk of bias graph: review authors' judgements about each risk of bias item presented as percentages across all included studies.

Figure 3. Risk of bias summary: review authors' judgements about each risk of bias item for each included study.

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias
Abd-Elmonem 2014	?	?	?	?	?	?	+
Cuesta-Barriuso 2013	?	?	?	+	?	?	+
Eid 2014	?	?	?	?	?	?	+
Kargarfard 2013	?	?	?	?	?	-	+
Mazloun 2014	?	?	?	?	?	?	+
Mohamed 2015	+	+	?	?	?	?	+
Parhampour 2014	+	+	?	?	+	?	+
Zaky 2013	?	?	?	?	?	?	+

Allocation

Sequence generation

We assessed two studies as having a low risk of bias for random sequence generation as one used a random draw

of closed envelopes ([Mohamed 2015](#)) and the other used computerized blocks indicating group allocation ([Parhampour 2014](#)). We assessed six studies as having an unclear risk of bias as the method of sequence generation was not described ([Abd-Elmonem 2014](#); [Cuesta-Barriuso 2013](#); [Eid 2014](#); [Kargarfard 2013](#);

Mazloun 2014; Zaky 2013). One study did report that participants were selected using "accessible sampling method" (Kargarfard 2013).

Allocation concealment

Two studies received a rating of low risk of selection bias (Mohamed 2015; Parhampour 2014). Allocation was concealed in one study by using "closed envelopes" (Mohamed 2015) and in another by computer block randomization carried out by an independent person not involved in the research team or project (Parhampour 2014). For six studies the method of concealment was not stated and these were judged to have an unclear risk of bias (Abd-Elmonem 2014; Cuesta-Barriuso 2013; Eid 2014; Kargarfard 2013; Mazloun 2014; Zaky 2013).

Blinding

In all eight included studies, blinding of the participants was not possible. Blinding of study personnel was demonstrated in only one study (Cuesta-Barriuso 2013). Specifically "blinded evaluators" assessed outcomes at all time points and this study was rated to have a low risk of detection bias (Cuesta-Barriuso 2013). Overall, the risk of blinding was unclear in seven studies as insufficient information was given as to the extent to which the unblinded assessment may have affected outcomes (Abd-Elmonem 2014; Eid 2014; Kargarfard 2013; Mazloun 2014; Mohamed 2015; Parhampour 2014; Zaky 2013).

Incomplete outcome data

One study received a rating of low risk of attrition bias as the study reported the number, reason for the dropouts and from which treatment group the dropouts were from (Parhampour 2014). Overall, the proportion of dropouts in the study population was 10% (n = 5) for the Parhampour study (three from the resistance training group and two from the resistance training with pulsed electromagnetic fields group) (Parhampour 2014). The Mazloun study did indicate the number of dropouts and the reason for withdrawing from the study, but did not report from which intervention group the dropouts occurred, therefore, this was classified as having an unclear risk of bias for this domain (Mazloun 2014). The remaining six studies received an unclear rating as none indicated any dropouts nor did the authors directly state that all participants completed the study (Abd-Elmonem 2014; Cuesta-Barriuso 2013; Eid 2014; Kargarfard 2013; Mohamed 2015; Zaky 2013).

Selective reporting

Seven studies were determined to have an unclear risk of selective reporting bias. One study was deemed to have a high risk of selective reporting bias as reference is made to administration of a questionnaire post-treatment but results were not reported (Kargarfard 2013).

Other potential sources of bias

We rated all eight included studies as having a low risk of other potential sources of bias.

Effects of interventions

See: [Summary of findings for the main comparison](#) Summary of findings - Exercise compared with no intervention; [Summary of findings 2](#) Summary of findings - Exercise A compared with Exercise

[B](#); [Summary of findings 3](#) Summary of findings - Exercise alone compared with exercise plus electrophysical modality; [Summary of findings 4](#) Summary of findings - Hydrotherapy compared with no exercise

A total of eight studies have been included in the analysis.

Exercise versus no intervention

Two studies were included in this comparison (Cuesta-Barriuso 2013; Parhampour 2014). Cuesta-Barriuso included 18 participants with 16 elbows as the unit of analysis and Parhampour included 25 participants (13 in the exercise group and 12 in the control group) (Cuesta-Barriuso 2013; Parhampour 2014).

Primary outcomes

1. Bleed frequency

Neither included study reported bleed frequency before or after the interventions.

2. Adverse events

Neither included study reported adverse events related to the study interventions.

3. Quality of life

Neither included study reported quality of life.

Secondary outcomes

1. Balance or proprioception (or both)

Neither included study reported on balance or proprioception.

2. Aerobic activity

Neither included study reported aerobic activity as a result of the interventions.

3. Joint health status

One study used the 'Modified Colorado Score' to assess status of elbows, knees and ankles in 25 participants (Parhampour 2014).

Interventions consisted of a six-week study of resistance training (n = 13) compared to the control group (no intervention) (n = 12).

A positive effect using the 'Modified Colorado Questionnaire' is reported as a reduction in score. Statistically significant improvements were demonstrated in the resistance training group as compared to the control in measures of joint health status of the elbows, MD -1.21 points (95% CI -2.14 to -0.28 points), knees, MD -3.42 points (95% CI -4.82 to -2.02 points) and ankles, MD -1.65 points (95% CI -2.51 to -0.79) ([Analysis 1.1](#)) (Parhampour 2014).

4. Pain intensity

Pain intensity was measured using a visual analog scale with an improvement in pain reported as a decreased in VAS score.

The Cuesta-Barriuso study (n = 18 - but data relate to 30 joints (16 in exercise group and 14 in the control group)), reported the effects of exercise as compared to no intervention on pain intensity after 12 weeks of treatment (Cuesta-Barriuso 2013). The study compared an exercise intervention which consisted of proprioceptive neuromuscular facilitation of the elbow joint (combined with joint traction) versus no exercise. We are unable

to enter the data into the 'Data and analyses' section, given the unit of analysis issue (data on number of elbows available, not the number of participants) but the abstract described an "observed improvement ($P < 0.05$) on the pain perception of elbow" favouring the exercise group (Cuesta-Barriuso 2013). The data reported in the abstract for the perception of elbow pain (VAS) are presented in an additional table (Table 1) (Cuesta-Barriuso 2013).

5. Functional status

Neither included study reported on functional status as a result of the interventions.

6. Range of motion

Joint flexion

The Cuesta-Barriuso study ($n = 18$ - but data relate to 30 joints (16 in exercise group and 14 in the control group)) reported on this outcome. The exercise intervention consisted of joint traction and proprioceptive neuromuscular facilitation of the elbow joint and reported the effects of exercise ($n = 9$) as compared to no intervention (control, $n = 9$) on range of motion or joint flexion. We are unable to enter the data into the 'Data and analyses' section, given the unit of analysis issue (data on number of elbows available, not the number of participants) but it was reported in the abstract that there was an "observed improvement ($P < 0.05$) on the flexion of elbow". The data reported in the abstract are presented in an additional table (Table 1) (Cuesta-Barriuso 2013).

Joint extension

The Cuesta-Barriuso study also reported on the effects of exercise as compared to no intervention on range of motion or joint extension after 12 weeks of treatment (Cuesta-Barriuso 2013). We are unable to enter the data into RevMan given the unit of analysis issue (data on number of elbows available, not the number of participants). The data reported in the abstract are presented in an additional table (Table 1) (Cuesta-Barriuso 2013).

7. Biceps perimeter

The Cuesta-Barriuso study ($n = 18$ participants with 30 elbows as the unit of analysis) reported the effectiveness of joint traction and proprioceptive neuromuscular facilitation ($n = 16$ elbows) as compared to no intervention (control, $n = 14$ elbows) on the perimeter of the biceps muscle following a 12-week intervention period (Cuesta-Barriuso 2013). We are unable to enter the data into the 'Data and analyses' section, given the unit of analysis issue (data on number of elbows available, not the number of participants). The data reported in the abstract are presented in an additional table (Table 1) (Cuesta-Barriuso 2013).

8. Strength

One study ($n = 18$ participants with 30 elbows as the unit of analysis) compared joint traction and proprioceptive neuromuscular facilitation ($n = 16$ elbows) to no intervention (control, $n = 14$ elbows) to investigate the effects on biceps strength (Cuesta-Barriuso 2013). We are unable to enter the data into the 'Data and analyses' section, given the unit of analysis issue (data on number of elbows available, not the number of participants). The data reported in the abstract are presented in an additional table (Table 1) (Cuesta-Barriuso 2013).

9. Knee circumference

Neither included study reported on knee joint circumference as a result of the interventions.

Exercise A versus Exercise B

Five studies ($n = 147$) were included in this comparison (Abd-Elmonem 2014; Eid 2014; Mazloun 2014; Mohamed 2015; Zaky 2013).

Three studies used a standardized exercise program for both groups ("Exercise A") then added additional exercises for one of the groups ("Exercise B") (Abd-Elmonem 2014; Eid 2014; Zaky 2013).

Two studies used a different exercise program for each group (Mazloun 2014; Mohamed 2015).

The components of all of the exercise programs and the differences between the groups are summarized in an additional table (Table 2).

Primary outcomes

1. Bleed frequency

None of the included studies reported bleed frequency.

2. Adverse events

None of the included studies reported on adverse effects

3. Quality of life

None of the included studies reported on quality of life.

Secondary outcomes

1. Balance or proprioception (or both)

The Mohamed study ($n = 30$) reported the effects of bicycle ergometry and treadmill training on balance and gait parameters in children with haemophilia (Mohamed 2015). Both groups received the same physical activity program and were then randomized into either the bicycle ($n = 15$) or the treadmill group ($n = 15$) (Mohamed 2015). The group that performed aerobic exercise training on the treadmill combined with the physical activity program significantly improved balance after three months of successive treatment in terms of overall stability index, MD 0.25 points (95% CI 0.19 to 0.31 points), anterior-posterior stability index, MD 0.35 points (95% CI 0.27 to 0.43 points) and medio-lateral stability index, MD 0.24 points (CI 95% 0.17 to 0.31 points) (Analysis 2.1) (Mohamed 2015).

2. Aerobic activity

None of the included studies reported on aerobic activity.

3. Joint health status measured via a validated outcome measure

None of the included studies reported joint health status.

4. Pain

The Mazloun study investigated the effects of the land-based exercise program ($n = 13$) described above compared to an aquatic-based exercise program ($n = 14$) on pain intensity (Mazloun 2014). The aquatic-based exercise program consisted of five minutes of warm up involving rhythmic movement of the lower limb in water, followed by 30 to 45 minutes of hamstring stretching and quadriceps strengthening in water followed by a five-minute

cool down of stretching exercises in water. The aquatic-based exercise program group demonstrated a statistically significant improvement in pain intensity following the four-week study period, MD 0.80 VAS (95% CI 0.41 to 1.19 VAS) ([Analysis 2.2](#)) ([Mazloun 2014](#)).

5. Functional status

The Zaky study ($n = 30$) investigated the effects of a partial weight bearing exercise program plus a quadriceps exercise training program ($n = 15$) compared to the quadriceps exercise training program alone ($n = 15$) using the six-minute walk test as a measure of functional status ([Zaky 2013](#)). An increase in the distance walked in six minutes is indicative of improvement. The six-minute walk test was measured before and after a six-week intervention period. A statistically significant improvement favouring the partial weight bearing program plus the quadriceps training program was demonstrated, MD 2.60 metres (95% CI 0.08 to 5.12 metres) ([Analysis 2.3](#)) ([Zaky 2013](#)).

6. Range of motion

The Mazloun study also investigated the effects of the land-based exercise program ($n = 13$) described above compared to an aquatic-based exercise program ($n = 14$) on range of motion or joint flexion. The aquatic-based exercise program consisted of five minutes of warm up involving rhythmic movement of the lower limb in water, followed by 30 to 45 minutes of hamstring stretching and quadriceps strengthening in water followed by a five-minute cool down of stretching exercises in water ([Mazloun 2014](#)). No significant difference was demonstrated between the land-based exercise program and the aquatic program on range of motion or joint flexion and extension following the four-week study period with the following results: flexion, MD 0.20 degrees (95% CI -5.61 to 6.01 degrees); and extension MD -0.10 (95% CI -1.59 to 1.39) ([Analysis 2.4](#)) ([Mazloun 2014](#)).

7. Biceps perimeter

None of the included studies reported on biceps perimeter.

8. Strength

Muscle strength was evaluated in three studies ($n = 90$) in which an exercise intervention ($n = 45$) was compared to another exercise intervention (control $n = 45$) ([Abd-Elmonem 2014](#); [Eid 2014](#); [Zaky 2013](#)).

In the Zaky study, the effects of a partial weight bearing exercise program plus a quadriceps exercise training program ($n = 15$) compared to the quadriceps exercise training program alone ($n = 15$) on knee extensor muscle strength was investigated ([Zaky 2013](#)). The Eid study investigated the effects on knee extensor muscle strength of a physical therapy program including stretching, strengthening and aerobic activity ($n = 15$) versus the same program with the addition of bicycle ergometry and weight resistance ($n = 15$) ([Eid 2014](#)). The Abd-Elmonem study compared a control group ($n = 15$) who received an exercise therapy program including ultrasound, stretching and strengthening exercises with a study group ($n = 15$) who received the same program plus treadmill training.

Strength outcomes demonstrated a significant improvement for the study groups over the control groups in all outcome measures of strength including knee extensor and flexor strength ([Analysis](#)

[2.5](#)) ([Abd-Elmonem 2014](#); [Eid 2014](#); [Zaky 2013](#)). In the Zaky study, as measured in force in kg, MD 6.40 (95% CI 4.81 to 7.99); in the Abd-Elmonem study, peak torque of the knee extensors at 60 degrees per second, MD 4.31 (95% CI 2.5 to 5.87); in the Abd-Elmonem and Eid studies combined at 120 degrees per second, MD 12.75 (95% CI 1.46 to 24.04) ([Analysis 2.5](#)).

A statistically significant improvement of knee flexor strength was also demonstrated in favour of the exercise groups ([Analysis 2.5](#)) ([Abd-Elmonem 2014](#); [Eid 2014](#)). In the Abd-Elmonem study, peak torque of the knee flexors at 60 degrees per second, MD 4.31 (95% CI 3.20 to 5.42); in the Abd-Elmonem and Eid studies combined at 120 degrees per second, MD 9.12 (95% CI 6.74 to 11.51) ([Analysis 2.5](#)).

9. Knee circumference

In the Abd-Elmonem study ($n = 30$) a exercise program consisting of stretching, strengthening and therapeutic ultrasound ($n = 15$) was compared to the same exercise program plus the addition of treadmill walking ($n = 15$) on knee circumference which was used as a measure of knee effusion ([Abd-Elmonem 2014](#)). A significant improvement in knee circumference was demonstrated in the treadmill walking group as compared to the exercise group, MD 2.70 (95% CI 2.20 to 3.20) ([Analysis 2.6](#)) ([Abd-Elmonem 2014](#)). Units of measurement for knee circumference were not provided by the study authors.

Exercise alone versus exercise plus electrophysical modality

One study ($n = 25$) was included in this comparison ([Parhampour 2014](#))

Primary outcomes

1. Bleed frequency (number of major bleeds reported per year, month, or week)

The included study did not report on bleed frequency.

2. Adverse events (e.g. bleed following exercise, worsening of symptoms)

The included study did not report on adverse events.

3. Quality of life

The included study did not report on quality of life.

Secondary outcomes

1. Balance or proprioception (or both)

The included study did not report on balance or proprioception.

2. Aerobic activity

The included study did not report on aerobic activity.

3. Joint health status measured via a validated outcome measure

The included study used the Modified Colorado Score to assess joint health status ([Parhampour 2014](#)).

A statistically significant improvement was noted in the ankle scores favouring the resistance training plus pulsed electromagnetic fields group over the resistance training group, MD 0.90 points (95% CI 0.07 to 1.73 points) ([Analysis 3.1](#)) ([Parhampour 2014](#)).

No significant differences were found between the resistance training group and the resistance training plus pulsed electromagnetic fields group for the elbows, MD 0.35 points (95% CI -0.70 to 1.40 points) or knees, MD 0.75 points (95% CI -0.47 to 1.97 points) ([Analysis 3.1](#)) ([Parhampour 2014](#)).

4. Pain

The included study did not report on pain.

5. Functional status

The included study did not report functional status.

6. Range of motion

The included study did not report on range of motion.

7. Biceps perimeter

The included study did not report on biceps perimeter.

8. Strength

The included study did not report on strength.

9. Knee circumference

The included study did not report on knee circumference.

Hydrotherapy versus no exercise

One study ($n = 20$) was included in this comparison ([Kargarfard 2013](#)). The study authors completed an analysis of covariance (ANCOVA) to make comparisons, between experimental and control groups on post-test scores adjusted for differences in pre-test scores ([Kargarfard 2013](#)). Unfortunately, we were unable to calculate differences as the baseline data for the control group were not available in the publication. We have contacted the study authors in an attempt to obtain the baseline data and if successful we will include these in the next update of this review.

Primary outcomes

1. Bleed frequency

The included study did not report on bleed frequency.

2. Adverse events

The included study did not report on adverse events.

3. Quality of life

The included study did not report on quality of life.

Secondary outcomes

1. Balance or proprioception (or both)

The included study did not report on balance or proprioception.

2. Aerobic activity

The included study did not report on aerobic activity.

3. Joint health status measured via a validated outcome measure

The included study did not report on joint health status.

4. Pain

The included study did not report on pain.

5. Functional status

The included study did not report functional status.

6. Range of motion

The study presented data for range of motion (flexion and extension) comparing an aquatic exercise therapy group ($n = 10$) (40 to 60 minutes per week for eight weeks) to a control group ($n = 10$) who did not participate in any physical activity for a period of eight weeks. The covariance analysis test in this study demonstrated a statistically significant improvement in range of motion in the experimental compared to the control group; however, as mentioned above we were unable to calculate the MD, SD and 95% CI since data were missing ([Kargarfard 2013](#)).

7. Biceps perimeter

The included study did not report on biceps perimeter.

8. Strength

The Kargarfard study presented data for knee flexor and extensor strength comparing an aquatic exercise therapy group ($n = 10$) (40 to 60 minutes per week for eight weeks) to a control group ($n = 10$) who did not participate in any physical activity for a period of eight weeks. The covariance analysis test in this study demonstrated a statistically significant improvement in both knee flexor and extensor strength in the experimental compared to the control group; however, as mentioned above we were unable to calculate MD, SD and 95% CI since data were missing ([Kargarfard 2013](#)).

9. Knee circumference

The included study did not report on knee circumference.

DISCUSSION

Summary of main results

Eight studies were included, which represented 233 males with all severities of haemophilia A and B, ranging in age from eight to 49 years. Study durations were from four to 12 weeks. Exercise interventions varied greatly and included resisted exercises, isometric exercises, bicycle ergometry, treadmill walking and hydrotherapy, making comparisons between studies difficult.

The general risk of bias was unclear. None of these studies measured or reported outcomes regarding bleed frequency or quality of life. No adverse events related to the interventions were measured or reported. None of the studies reported outcomes regarding aerobic activity.

All studies favoured exercise to no intervention for most of the outcome measures that were assessed. One study reported that six weeks of resistance training improved joint health status (Colorado score) compared to controls. The addition of pulsed electromagnetic fields also improved ankle scores compared to exercises alone, but this was not seen in the elbows or knees.

One study reported that hydrotherapy exercises produced statistically significant improvements in joint range of motion and knee flexor and extensor strength in the exercise group compared to controls.

One study, comparing joint traction and the proprioceptive neuromuscular facilitation (PNF) technique for the elbow to a control group, showed no differences in biceps girth or biceps strength after 12 weeks of intervention.

Some studies reported comparisons between interventions. In one study, treadmill training significantly improved balance in children compared to bicycle ergometry. Another study added a partial weight bearing exercise program to a quadriceps exercise program and showed improved walking tolerance.

Four studies evaluated quadriceps or hamstring strength (or both). Addition of bicycle ergometry and exercises with weights was more effective than static exercises and treadmill walking for strengthening knee flexors and extensors. Partial weight bearing exercises through range (leg presses in supine against a wall or therapy ball, bridging exercises in supine) were more effective than static and short arc exercises for improving knee extensor strength. The addition of treadmill walking to ultrasound, stretching and strengthening exercises showed increased peak torque of knee flexors and extensors and decrease in knee effusion (circumference).

Most exercise interventions produced improvement in one or more measured outcomes including pain, range of motion, strength and walking tolerance. Hydrotherapy may be more effective than land exercises for pain relief in adults. Functional exercises such as treadmill walking and partial weight bearing exercises seem to be more effective than static or short arc exercises for improving muscle strength. These findings would be consistent with the many non-controlled intervention reports in the haemophilia literature.

No adverse effects, new bleeds or dropouts due to injury were reported due to any of the interventions. However, it should be noted that some groups used prophylactic factor prior to exercise and other groups studied only subjects with moderate haemophilia. Therefore, application of these techniques to persons with severe haemophilia, especially to those without access to prophylactic clotting factor, should still be approached with caution.

Overall completeness and applicability of evidence

Overall, the results demonstrate a large degree of heterogeneity in interventions and outcomes and limit the ability to demonstrate a clear preference of one type of exercise program over another.

Outcome measures that were reported upon measured impairment or function of the musculoskeletal system and none of the included studies provided reports of quality of life, effect on bleeding frequency or participation in age-appropriate societal activities. Further, there was a lack of information regarding prophylaxis or on-demand treatment regimens and the impact that access to factor replacement treatment has on the ability or willingness to exercise. The general risk of bias was assessed as unclear. No adverse events related to the interventions were reported. None of the studies reported outcomes regarding aerobic activity.

Exercise interventions varied greatly and included resisted exercises, isometric exercises, partial weight-bearing exercises, bicycle ergometry, treadmill walking and hydrotherapy, making comparisons between studies difficult. Also, variability in the outcome measures used to demonstrate treatment effect negatively affects the confidence of our findings due to the inability

to pool amongst studies. Lastly, due to the lack of reported data on intra-subject variability we were unable to account for this when assessing the variability of the variation over time. This limitation results in the over estimation of the variability of participant change scores and wider confidence intervals of the pooled results which may have impacted statistical significance. However, the authors believe that this method is more conservative and we may have missed an effect where one is there, but would not have resulted in finding a spurious effect, which we considered an acceptable limitation.

All studies in this review favoured exercise to no intervention or control for all secondary outcome measures with the exception of biceps perimeter and biceps strength. One study reported that six weeks of resistance training improved joint health status (Colorado score) compared to controls (Parhampour 2014). The addition of pulsed electromagnetic fields also improved ankle scores compared to exercises alone, but this was not seen in the elbows or knees (Parhampour 2014). One study reported statistically significant improvements in pain intensity, favouring the exercise group over controls after the exercise interventions (Cuesta-Barriuso 2013). This study also reported statistically significant improvements on range of motion or joint flexion, and range of motion or joint extension, favouring the exercise groups (Cuesta-Barriuso 2013). In this study the exercise intervention consisted of joint traction and PNF of the elbow joint (Cuesta-Barriuso 2013). These results indicate that exercise interventions have a positive impact on pain palliation, joint range of motion and joint health status over that of no intervention or control.

When comparing one type of exercise intervention to another type of exercise intervention, all secondary outcome measures with the exception of range of motion or joint flexion or joint extension demonstrated statistical significance of one exercise program over the other as described below.

When analyzing the results of the comparison of one exercise intervention to an exercise intervention plus an add-on modality this review demonstrated an improvement in joint health status of the ankles, but not knees or elbows, and also demonstrated an improvement in secondary outcome measures of strength and knee circumference.

Again, these results demonstrate a large degree of heterogeneity in interventions and outcomes and limit the interpretation of effects.

The fourth and final comparison of hydrotherapy versus no intervention or control reported a statistically significant improvement in range of motion and knee flexor and extensor strength via covariant analysis; however, as mentioned above, we were unable to calculate the mean difference, standard deviation and 95% confidence interval due to the missing data (Kargarfard 2013).

Quality of the evidence

There are few randomized controlled studies addressing exercise for people with haemophilia. The quality of the eight studies included in this review was graded according to the standardized risk of bias tool as described in the *Cochrane Handbook of Systematic Reviews of Interventions* (Higgins 2011a). As a result of the heterogeneity of study populations, interventions and outcomes it was difficult to accurately determine study quality. This

is further illustrated by the inability to pool data for quantitative analysis of the primary outcomes.

Three studies did not specify type or severity of haemophilia (Kargarfard 2013; Mohamed 2015; Zaky 2013). Four studies included only people with moderate haemophilia (Abd-Elmonem 2014; Eid 2014; Kargarfard 2013; Zaky 2013). Three studies included all (Mazloun 2014; Mohamed 2015) or mixed (Cuesta-Barriuso 2013) severities of haemophilia. Two studies included participants with both haemophilia A and B (Abd-Elmonem 2014; Cuesta-Barriuso 2013). In one study participants were asked "to take their clotting factor prior to participating" (Mazloun 2014). Only one study limited their participants to severe haemophilia, and these also had osteoporosis (Parhampour 2014). Therefore, it is not clear whether the same results would be achieved if, for example, only individuals with severe haemophilia A were studied.

Some of the outcome measures that were chosen were not used appropriately. For example, one study used the Modified Colorado Hemophilia PEDIATRIC Physical Examination Scale for an ADULT cohort (Parhampour 2014), another study evaluated an exercise intervention using biceps perimeter as a measure of muscle strength (Cuesta-Barriuso 2013). However, biceps perimeter does not correlate with muscle strength or function. Also units of measurement for the biceps perimeter (Cuesta-Barriuso 2013) and knee circumference (Abd-Elmonem 2014) were not provided by the study authors.

The results should be interpreted with caution due to the Quality of Evidence (GRADE) as outlined in the summary of finding tables which demonstrate that all but one of the outcomes assessed were rated as "low" or "very low" due to small sample sizes and potential bias (Summary of findings for the main comparison; Summary of findings 2; Summary of findings 3; Summary of findings 4).

Potential biases in the review process

Many different types of exercise interventions and outcomes were included in this review. This may produce potential biases in the review process as some methods of exercise were better described to allow for replication, and many outcome assessments were not completed using valid and reliable outcome measures. For example, in one of the studies a measure of joint health, the Modified Colorado Hemophilia Pediatric Physical examination Scale was used to evaluate an adult population; however, this outcome measure has only been validated in children.

Agreements and disagreements with other studies or reviews

The findings of this review, which was limited to full-text or abstracts of randomized, controlled studies only, are consistent with the findings of two recent comprehensive review articles which included a wider range of study designs (Gomis 2009; Negrier 2013). These authors reviewed the benefits and effectiveness of exercise, and also concluded that exercise can have a positive effect on physical well-being. The authors further commented on emotional and social well-being, but this current review was unable to draw any conclusions in this regard due to lack of data. The authors also commented on the lack of rigorously designed trials, methodological drawbacks, and low levels of evidence in the existing literature.

AUTHORS' CONCLUSIONS

Implications for practice

These results must be considered with caution. There is a lack of confidence in the results due to the small number of included studies and the inability to pool the results due to the heterogeneity of interventions and outcome measures used. Very few studies provided sufficient information for comparison.

All studies favoured some form of exercise to no intervention for most of the assessed outcome measures and none of the studies reported data favoured the control group.

Hydrotherapy may be more effective than land exercises for pain relief in adults. Functional exercises such as treadmill walking and partial weight bearing exercises seem to be more effective than static or short arc exercises for improving muscle strength.

No adverse effects, changes in bleeding frequency or bleeding due to the exercise interventions used, were reported in any of the studies. However, it is impossible to conclude whether these interventions are safe for all people with severe haemophilia. Only one study specifically selected participants with severe haemophilia (who also had osteoporosis). The other studies looked at children with moderate haemophilia, or mixed participants of all severities within the same groups. Therefore, it is our contention that caution should be used when prescribing exercise for individuals with severe haemophilia, especially where factor replacement is limited. Further, optimal timing, dosage and types of exercise are yet to be determined.

Implications for research

The challenges of conducting randomized controlled studies with this population are large; haemophilia is a rare condition, best managed in large comprehensive treatment centres. The ordinary challenges of daily living are compounded greatly when haemophilia is added to the picture: financial burden, time lost from work or school, the stress of pain and limited ability are all significant factors. Researchers must be sensitive to the increased burden associated with participation in an ongoing research project. The authors of the selected studies should be commended for their efforts but it is apparent that there is a need for more well-designed clinical studies with larger numbers of participants.

Future studies should aim to:

- continue to evaluate safety of different forms of exercise, especially for those with severe haemophilia and inhibitors and those with access only to factor on demand (versus access to prophylactic factor);
- homogenize participants by disease (A and B) and severity (mild, moderate, severe), as well as joint status and age;
- have clear intervention protocols, including frequency and intensity of exercise, and matched control groups;
- ensure that comparisons between groups are comparing equivalent items;
- use standardized outcome measures that have established psychometric properties and apply them correctly;
- develop a core set of internationally accepted outcome assessments that evaluate the musculoskeletal complications of the disease (e.g. achieve an international consensus on

which measure of muscular strength to apply and encourage all researchers to use this same measure to allow for future meta-analyses).

ACKNOWLEDGEMENTS

The authors would like to acknowledge the assistance of Tracey Remington.

REFERENCES

References to studies included in this review

Abd-Elmonem 2014 {published data only}

Abd-Elmonem AM, Abonour AA, Elnaggar RK. Effect of treadmill training on Quadriceps and hamstring muscles strength in Children with Knee Haemarthrosis. *International Journal of Physiotherapy and Research* 2014;**2**(4):591-8.

Cuesta-Barriuso 2013 {published data only}

Cuesta-Barriuso R, Moren-Moreno M, Garcia-Diego D. Effectiveness of joint traction and PNF in hemophilic arthropathy of the elbow [abstract]. *Haemophilia* 2013;**19**(4):e206.

Eid 2014 {published data only}

Eid M, Ibrahim M, Aly SM. Effect of resistance and aerobic exercises on bone mineral density, muscle strength and functional ability in children with hemophilia. *Egyptian Journal of Medical Human Genetics* 2014;**15**:139-47.

Kargarfard 2013 {published data only}

Kargarfard M, Dehghadani M, Ghias R. The effect of aquatic exercise therapy on muscle strength and joint's range of motion in hemophilia patients. *International Journal of Preventive Medicine* 2013;**4**(1):50-6. [CENTRAL: 908835; CRS: 5500050000000025; EMBASE: 2013152750]

Kargarfard M, Dehghani M, Heidari A. Effect of a period of aquatic exercise therapy on the quality of life, anxiety and depression in patients with hemophilia. *Koomesh* 2011;**12**(4):364-71. [CENTRAL: 1016052; CRS: 55000500000000511; EMBASE: 2011352770]

Soltani M, Kargarfard M, Nadi M, Hoseini M. The effect of 8-weeks exercise in water on factor VIII and partial thromboplastin time (PTT) of men with hemophilia. [Persian]. *Journal of Isfahan Medical School* 2016;**33**(357):1878-83. [CENTRAL: 1134151; CRS: 55000500000000419; EMBASE: 20160049552]

Mazloun 2014 {published data only}

Mazloun V, Khayambashi KH, Rahnama N. Comparison of the effect of aquatic exercise therapy and land-based therapeutic exercise on knee muscles' strength and quality of life in patients with knee joint arthropathy due to hemophilia. *Journal of Babol University of Medical Sciences* 2014;**16**(6):26-32.

* Mazloun V, Rahnama N, Khayambashi K. Effects of therapeutic exercise and hydrotherapy on pain severity and knee range of motion in patients with hemophilia: A randomized controlled trial. *International Journal of Preventive Medicine* 2014;**5**(1):83-8. [CENTRAL: 978360; CRS: 5500050000000044; EMBASE: 2014078882]

Mohamed 2015 {published data only}

Mohamed RA, Sherief AE-AA. Bicycle ergometer versus treadmill on balance and gait parameters in children with hemophilia. *Egypt Journal of Medical Human Genetics* 2015;**16**(2):181-7.

Parhampour 2014 {published data only}

Parhampour B, Torkaman G, Hoorfar H, Hedayati M, Ravanbod R. Effects of short-term resistance training and pulsed electromagnetic fields on bone metabolism and joint function in severe haemophilia A patients with osteoporosis: a randomized controlled trial. *Clinical Rehabilitation* 2014;**28**(5):440-50. [CENTRAL: 1035543; CRS: 5500131000000301; DOI: [10.1177/0269215513505299](https://doi.org/10.1177/0269215513505299); JID:: 8802181; PUBMED: 24249841]

Parhampour B, Torkaman G, Hoorfar H, Hedayati M, Ravanbod R. Effects of six weeks resistance training and pulsed electromagnetic fields (PEMFs) on static and dynamic balance and quality of life in hemophilia A patients with osteoporosis [abstract]. *Haemophilia* 2014;**20** Suppl 3:154. [CENTRAL: 1000901; CRS: 5500131000000112]

Zaky 2013 {published data only}

Zaky LA, Hassan WF. Effect of partial weight bearing program on functional ability and quadriceps muscle performance in hemophilic knee arthritis. *Egyptian Journal of Medical Human Genetics* 2013;**14**(4):413-8. [CENTRAL: 913753; CRS: 5500050000000042; EMBASE: 2013643065]

References to studies excluded from this review

Czepa 2013 {published data only}

Czepa D, Von Mackensen S, Hilberg T. Haemophilia & Exercise Project (HEP): The impact of 1-year sports therapy programme on physical performance in adult haemophilia patients. *Haemophilia* 2013;**19**:194-9. [DOI: [10.1111/hae.12031](https://doi.org/10.1111/hae.12031)]

Greene 1983 {published data only}

Greene W, Strickler E. A Modified Isokinetic Strengthening Program of Patients with Severe Hemophilia. *Developmental Medicine and Child Neurology* 1983;**25**:189-96.

Hilberg 2003 {published data only}

Hilberg T, Herbsleb M, Puta C, Gabriel HH, Schramm W. Physical training increases isometric muscular strength and proprioceptive performance in haemophilic subjects. *Haemophilia* 2003;**19**(1):86-93.

Khriesat 2000 {published data only}

Khriesat I, Thonaibat W, Hammaury M. Haemophilic Knee: role of Physiotherapy. *Bahrain Medical Bulletin* 2000;**22**(4):164-66.

Mulvany 2010 {published data only}

Mulvany R, Zucker-Levin A, Jeng M, Joyce C, Tuller J, Rose JM, et al. Effects of a 6-week, Individualized, Supervised Exercise Program for People with Bleeding Disorders and Hemophilic Arthritis. *Physical Therapy* 2010;**90**:509-26.

Vallejo 2010 {published data only}

Vallejo L, Pardo A, Gomis M, Gallach JE, Perez s, Querol F. Influence of aquatic training on the motor performance of patients with haemophilic arthropathy. *Haemophilia* 2010;**16**:155-61.

Von Mackensen 2012 {published data only}

* Von Mackensen S, Eifrig B, Zach D, Kalnins J, Wieloch A, Zeller W. The impact of a specific aqua-training for adult haemophilic patients – results of the WATERCISE study (WAT-QoL). *Haemophilia* 2012 Sep;**18**(5):714-21. [DOI: [10.1111/j.1365-2516.2012.02819.x](https://doi.org/10.1111/j.1365-2516.2012.02819.x)]

References to studies awaiting assessment

Cuesta-Barriuso 2014 {published data only}

Cuesta-Barriuso R, Gomez-Conesa A, Lopez-Pina J-A. Effectiveness of two modalities of physiotherapy in the treatment of haemophilic arthropathy of the ankle: A randomized pilot study. *Haemophilia* 2014;**20**(1):e71-8. [CENTRAL: 961298; CRS: 5500050000000043; EMBASE: 2013812438]

Cuesta-Barriuso 2015 {published data only}

Cuesta-Barriuso R, Torres-Ortuno A, Nieto-Munuera J. Effectiveness of physiotherapy on quality of life and illness behaviour in patients with haemophilic arthropathy. A randomized clinical trial [abstract]. *Haemophilia* 2015;**21**(Suppl):68, Abstract no: PP092. [CENTRAL: 1073620; CRS: 55000500000000241; EMBASE: 71814497]

Firoozabadi 2012 {published data only}

Firoozabadi MD, Mahdavejad R, Ghias M, Rouzbehani R, Dehghani M. The effects of an exercise therapy program on joint range of motion, aerobic fitness, and anxiety of hemophilia A patients. *Journal of Isfahan Medical School* 2012;**30**(177):1-9.

Additional references

Begg 1994

Begg CB, Mazumdar M. Operating characteristics of a rank correlation test for publication bias. *Biometrics* 1994;**50**(4):1088-101.

Beyer 2010

Beyer R, Ingerslev J, Sørensen B. Muscle bleeds in professional athletes--diagnosis, classification, treatment and potential impact in patients with haemophilia. *Haemophilia* 2010;**16**(6):858-65. [DOI: [10.1111/j.1365-2516.2010.02278.x](https://doi.org/10.1111/j.1365-2516.2010.02278.x); PUBMED: PMID: 20491962]

Blamey 2010

Blamey G, Forsyth A, Zourikian N, Short L, Jankovic N, De Kleijn P, et al. Comprehensive elements of a physiotherapy exercise programme in haemophilia – a global perspective. *Haemophilia* 2010;**16** Suppl 5:136–45.

De La Corte-Rodriguez 2013

De La Corte-Rodriguez H, Rodriguez-Merchan EC. The role of physical medicine and rehabilitation in haemophilia patients. *Blood Coagulation and Fibrinolysis* 2013;**24**(1):1-9.

Deeks 2011

Deeks J, Higgins J, Altman D. Chapter 9 Analysing data and undertaking meta-analysis. In: Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of*

Interventions Version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011. Available from www.cochrane-handbook.org.

Gomis 2009

Gomis M, Querol F, Gallach JE, González LM, Aznar JA. Exercise and sport in the treatment of haemophilic patients: a systematic review. *Haemophilia* 2009;**15**(1):43-54.

Higgins 2002

Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Statistics in Medicine* 2002;**21**(11):1539-58. [PUBMED: 12111919]

Higgins 2011a

Higgins JPT, Altman DG (editors). Chapter 8: Assessing risk of bias in included studies. In: Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011. Available from www.cochrane-handbook.org.

Higgins 2011b

Higgins JPT, Deeks JJ, Altman DG, on behalf of the CSMG. Chapter 16: Special topics in statistics. In: Higgins JPT, Green S (editors). *Cochrane Handbook of Systematic Reviews of Interventions*. Version 5.1 [updated March 2011]. The Cochrane Collaboration, 2011. Available from www.cochrane-handbook.org.

Higgins 2011c

Higgins JPT, Green S (Editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011. Available from www.cochrane-handbook.org.

Hozo 2005

Hozo SP, Djulbegovic B, Hozo I. Estimating the mean and variance from the median, range, and the size of a sample. *BMC Medical Research Methodology* 2005;**5**:13.

Hunter 2009

Hunter DJ, Eckstein F. Exercise and osteoarthritis. *Journal of Anatomy* 2009;**214**(2):197-207.

Iorio 2010

Iorio A, Fabbriani G, Marcucci M, Brozzetti M, Filipponi P. Bone mineral density in haemophilia patients. A meta-analysis. *Thrombosis and haemostasis* 2010;**103**(3):596-603.

Manco-Johnson 2004

Manco-Johnson MJ, Pettersson H, Petrini P, Babyn PS, Bergstrom BM, Bradley CS, et al. Physical Therapy and imaging outcome measures in a hemophilia population treated with factor prophylaxis: current status and future directions. *Haemophilia* 2004;**10** Suppl 4:88-93.

Negrier 2013

Negrier C, Seuser A, Forsyth A, Lobet S, Llinas A, Rosas M, Heijnen L. The benefits of exercise for patients with haemophilia and recommendations for safe and effective

physical activity. *Haemophilia* 2013;**19**(4):487-98. [DOI: [10.1111/hae.12118](https://doi.org/10.1111/hae.12118)]

Pettersson 1980

Pettersson H, Ahlberg A, Nilsson IM. A radiologic classification of hemophilic arthropathy. *Clinical Orthopedics and Related Research* 1980;**149**:153-9.

RevMan 2014 [Computer program]

The Nordic Cochrane Centre, The Cochrane Collaboration. Review Manager (RevMan). Version 5.3. Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration, 2014.

Riske 2007

Riske B. Sports and exercise in haemophilia: benefits and challenges. *Haemophilia* 2007;**13** Suppl 2:29-30.

Rodriguez-Merchan 2010

Rodriguez-Merchan EC. Musculoskeletal complications of hemophilia. *Hospital for Special Surgery Journal*. 2010;**6**(1):37-42. [doi: 10.1007/s11420-009-9140-9]

Rodriguez-Merchan 2012

Rodriguez-Merchan EC. Cartilage damage in the hemophilic joints: pathophysiology, diagnosis and management. *Blood Coagulation & Fibrinolysis* 2012;**23**(3):179-83.

Roosendaal 2008

Roosendaal G, Jansen NW, Schutgens R, Lafeber FP. Hemophilic arthropathy: the importance of the earliest hemarthrosis and consequences for treatment. *Haemophilia* 2008;**14** Suppl 6:4-10.

Stenstrom 2003

Stenstrom Christina H, Minor Marian A. Evidence for the benefit of aerobic and strengthening exercise in rheumatoid arthritis. *Arthritis and Rheumatism* 2003;**49**(3):428-34.

WFH 2012

Srivastava A, Brewer AK, Mahlangu JN, Mauser-Bunschoten EP, Mulder K, Key NS. Guidelines For The Management Of Hemophilia (2nd edition). <http://www1.wfh.org/publications/files/pdf-1472.pdf> (accessed 01 April 2013).

* Indicates the major publication for the study

CHARACTERISTICS OF STUDIES

Characteristics of included studies [ordered by study ID]

Abd-Elmonem 2014

Methods	Randomized controlled study.	
Participants	30 boys with moderate A and B, unilateral knee haemarthrosis, 8 - 12 years of age. Exclusion criteria: acute bleeds, advanced radiologic changes, congenital or acquired skeletal deformities.	
Interventions	<p>Both groups: exercise stretching and strengthening, ultrasound for 1 hour, 5 days per week.</p> <p>Exercise group: in addition to the above performed ambulation on treadmill: speed 1.5 km/hr no incline for 5 min as a warm up, 20 minutes of exercise 3 km/hr 10 degree incline, cool down 1.5 km/hr no incline for 30 min.</p> <p>Duration of study: 3 months.</p>	
Outcomes	Strength assessment with the Biodex Isokinetic Dynamometer: isokinetic pre- and post-test at 2 different velocities, peak torque of knee flexors and peak torque of knee extensors; knee circumference.	
Notes		
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	<p>Quote: "They were divided randomly into two groups of equal numbers."</p> <p>Comment: method of randomization was not reported.</p>
Allocation concealment (selection bias)	Unclear risk	Comment: no information was reported re allocation concealment.

Abd-Elmonem 2014 (Continued)

Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: not reported whether personnel or participants were blinded.
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: not reported whether assessors of outcome were blinded.
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Quote: "Isokinetic pre-test and post-test measurements....were performed on every child...". Comment: not reported whether joint circumference was measured on every child.
Selective reporting (reporting bias)	Unclear risk	Insufficient information provided to make judgement of low or high risk.
Other bias	Low risk	No other sources of bias were identified.

Cuesta-Barriuso 2013

Methods	Randomized controlled study.
Participants	18 participants, 14 haemophilia A, 4 haemophilia B; 11 with severe haemophilia, 7 with mild; 9 participants in intervention group, 9 participants in control resulting in 16 elbows in the intervention and 14 elbows in the control.
Interventions	Intervention group: traction and PNF, control group: no intervention. 12 week intervention, 2 times per week, 1 hour session , evaluation pre-, post- and 6-month follow up.
Outcomes	ROM, strength, pain, circumference.
Notes	Elbows only.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "randomized trial with two groups". Comment: not reported how participants were randomized.
Allocation concealment (selection bias)	Unclear risk	Comment: not reported if or how randomization was concealed.
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: not reported whether participants were blinded.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "Three blind evaluators assessed: ROM, biceps perimeter, biceps strength, pain."

Exercise for haemophilia (Review)

Cuesta-Barriuso 2013 (Continued)

Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Comment: not reported whether all participants completed all measurements.
Selective reporting (reporting bias)	Unclear risk	Insufficient information provided to make judgement of low or high risk.
Other bias	Low risk	No other sources of bias were identified.

Eid 2014

Methods	Randomized controlled study.
Participants	30 boys with moderate haemophilia A, 10 - 14 years. Participants equally divided between intervention and control.
Interventions	Control and experimental group: gentle stretching for tight muscle groups around elbow, knee, ankle for 15 min, isometric contractions knees, ankles, elbows 15 min, aerobic exercise on treadmill 30 min. 1 hr, 3 times per week. In addition to this, the experimental group also received bicycle ergometry 20 min, resistance training with sand bags 20 min. Total time for experimental group was 1 hr 40 min, control group 1 hr. Intervention period: 3 successive months.
Outcomes	BMD, muscle strength and function, Biodex Isokinetic Dynamometer (knee flexor/extensor peak torque) 6MWT, DEXA.
Notes	Lots of detail of intervention protocol.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Comment: participants were "assigned randomly" not reported how allocation sequence was generated.
Allocation concealment (selection bias)	Unclear risk	Comment: not reported how allocation was concealed.
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: not reported whether personnel or participants were blinded.
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: not reported whether assessors of outcome were blinded.
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Comment: not reported whether all participants completed each outcome assessment.
Selective reporting (reporting bias)	Unclear risk	Insufficient information provided to make judgement of low or high risk.

Exercise for haemophilia (Review)

Eid 2014 (Continued)

Other bias	Low risk	No other sources of bias were identified.
------------	----------	---

Kargarfard 2013

Methods	Semi-experimental, 8-week intervention.
Participants	20 men (ages not specified, although mean ages are given) with moderate haemophilia.
Interventions	<p>Experimental group: 24 sessions 3x per week, 40 - 60 min, increase time and quality of exercise at each session. Sessions consisted of 3 sections: warm up, main part and cool down. Warm up aerobic activity 50% - 74% of HR predicted. Main part: 10 simple movements (5UE, 5LE) in water: 1 min for each movement maximum power as pain allowed. Cool down: individualized stretching program.</p> <p>Control group: no intervention.</p> <p>Each session (3 times per week) lasted 45 - 60 min.</p> <p>Duration of study: 8 weeks.</p>
Outcomes	Muscle strength (Biodex Isokinetic Dynamometer, knees only), ROM (standard goniometer, inactive range: elbows, knees, ankles), heart rate before and during.
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Comment: not reported how allocation sequence was generated.
Allocation concealment (selection bias)	Unclear risk	Comment: not reported how allocation was concealed.
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: not reported whether personnel or participants were blinded.
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: not reported whether assessors of outcome were blinded.
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Comment: not reported whether all participants completed each outcome assessment.
Selective reporting (reporting bias)	High risk	Comment: reference is made to administration of a questionnaire post-treatment; results were not reported.
Other bias	Low risk	No other sources of bias were identified.

Mazloun 2014

Methods	Quasi-experimental, prospective study design, randomized into 2 groups, with the addition of a non-randomized control group (data from this arm (n = 13) not used in this review).
Participants	<p>40 participants, haemophilia A, all severities, < 50 years of age. 13 exercise, 14 hydro, 13 control (not randomized). 43 started in protocol, 3 dropouts.</p> <p>Inclusion criteria: haemophilia, ability to participate, impaired knee ROM.</p> <p>Exclusion criteria: surgery 6 weeks prior, participation in sports or other exercises, severe haemorrhage despite factors, open wounds, sensibility to aquatic therapy.</p>
Interventions	<p>4 week hydrotherapy program vs exercise vs control.</p> <p>Hydrotherapy: co-ordinated and rhythmic movement of lower limb in water (warm-up), hamstring stretches, quad strengthening.</p> <p>Exercise: stretches for muscles around the knee joint (warm up), isometrics progressing to isotonic, hamstring stretches, quad strengthening.</p> <p>Control: regular lifestyle. NOTE: this group was not randomized, therefore data from this arm not used in the review.</p> <p>** Only difference between intervention groups is the warm up: hydrotherapy and stretches for muscles around the knee joint.</p> <p>Duration of study: 4 weeks.</p>
Outcomes	Pain before and after interventions, ROM knee flexion and extension.

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Comment: not reported how participants were randomized.
Allocation concealment (selection bias)	Unclear risk	Comment: not reported how allocation was concealed.
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: not reported whether personnel or participants were blinded.
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: not reported whether assessors of outcome were blinded.
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Comment: data for 3 participants who did not complete the intervention were not included in the analysis and were reported by the authors but they did not report from which group the drop outs occurred, therefore characterized as unclear risk.
Selective reporting (reporting bias)	Unclear risk	Insufficient information provided to make judgement of low or high risk.
Other bias	Low risk	No other sources of bias were identified.

Exercise for haemophilia (Review)

Mohamed 2015

Methods	Prospective randomized controlled study, no blinding.
Participants	30 boys with mild or moderate haemophilia, ages ranged from 10 - 14 years. Participants were randomly assigned to group A (bicycle ergometer + exercise, n = 15) or group B (treadmill training + exercise, n = 15).
Interventions	Group A received exercise program and bicycle ergometry, Group B received exercise program and treadmill training. Both groups received treatment sessions 3 times per week for 3 successive months.
Outcomes	Stability or balance and kinematic gait parameters were evaluated before and after 3 successive months of treatment using the Biodex Stability System and the Biodex Gait trainer 2TM.
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Comment: randomization was done by having the participants draw an envelope which contained a card indicating either group A or group B
Allocation concealment (selection bias)	Low risk	Comment: each participant drew a sealed envelope.
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: not reported whether personnel or participants were blinded.
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: not reported whether assessors of outcome were blinded.
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Comment: not reported whether all participants completed each outcome assessment.
Selective reporting (reporting bias)	Unclear risk	Insufficient information provided to make judgement of low or high risk.
Other bias	Low risk	No other sources of bias were identified.

Parhampour 2014

Methods	Randomized controlled study.
Participants	<p>Inclusion criteria: severe haemophilia A, BMI 22 - 29, no inhibitor, t-score < 2.5, willingness to comply with the design, regular patient at clinic. Age 20 - 35 years.</p> <p>Exclusion criteria: target joint preventing resistance training, hepatitis B or C, thyroid disorder, steroids, regular physical training more than 2x per week in preceding 6 months, hormone therapy, drugs that influence bone metabolism, hypertension, other orthopedic that might limit ability to participate in resistance training.</p>

Parhampour 2014 (Continued)

70 enrolled in study; 22 excluded based on inclusion criteria; 48 remaining and were randomly assigned to 4 groups.

1. RT only: 13 participants.
2. RT with PEMF: 12 participants.
3. PEMF only: 11 participants.
4. Control no resistance and no PEMF: 12 participants.

Within random allocation participants with low t-scores were evenly distributed within the groups.

Duration of study: 6 weeks.

Interventions	<p>Resistance training involved 18 sessions over 6 weeks, 3 times a week, 30 - 40 min, resistance was set based on 1 repetition max (started at 50%, increased to 55%, finished at 60%), 10 reps during weeks 1, 3, 5. 15 reps during weeks 2, 4, 6, amount of rest time was specified.</p> <p>RT only: after exercises received 30 min of placebo PEMF, very good description of exercises provided.</p> <p>RT and PEMF group: 5 reps of each exercise in 1, 3, 5 and 10 reps in weeks 2, 4, 6. After the training program participants received 30 min of PEMF, parameters described.</p> <p>PEMF: no exercise training, 6 weeks of PEMF 1 hour per session over 6 weeks.</p> <p>Control group: no intervention.</p>
Outcomes	<p>Body fat percentage.</p> <p>Modified Colorado Scale, Pediatric (ROM, strength, pain, etc).</p> <p>Blood work for bone biomarkers.</p>
Notes	Conclude that pain is improved but unable to find raw data in results.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "randomization (using computerized blocks) was carried out....".
Allocation concealment (selection bias)	Low risk	Comment: randomization was carried out by an independent person, not involved in the research project.
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	<p>Quote: "all patients were blinded to the groupings but the investigator for the modified Colorado Questionnaire was not masked to the group assignment."</p> <p>Comment: extent to which this unblinded assessment may have influenced outcomes is unclear.</p>
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	<p>Quote: "the investigator for the modified Colorado Questionnaire was not masked to the group assignment."</p> <p>Comment: extent to which this unblinded assessment may have influenced outcomes is unclear.</p>
Incomplete outcome data (attrition bias) All outcomes	Low risk	<p>Quote: "Five patients did not complete the project."</p> <p>Comment: 10% overall dropout rate. Not reported how incomplete data were handled.</p>

Exercise for haemophilia (Review)

Parhampour 2014 (Continued)

Selective reporting (reporting bias)	Unclear risk	Insufficient information provided to make judgement of low or high risk.
Other bias	Low risk	No other sources of bias were identified.

Zaky 2013

Methods	Participants were randomly assigned into 2 equal groups.
Participants	30 boys with moderate haemophilia aged 8 - 12 years; randomly assigned into 2 equal groups.
Interventions	Control group: quadriceps training exercise program. Treatment group: as above plus partial weight bearing program. Study duration: 3 time a week for 6 weeks.
Outcomes	Manual muscle strength testing of quadriceps (Lafayette Manual Muscle Test system); 6MWT.
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Comment: not reported how allocation sequence was generated.
Allocation concealment (selection bias)	Unclear risk	Comment: not reported how allocation was concealed.
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: not reported whether personnel or participants were blinded.
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: not reported whether assessors of outcome were blinded.
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Comment: not reported whether all participants completed each outcome assessment.
Selective reporting (reporting bias)	Unclear risk	Insufficient information provided to make judgement of low or high risk.
Other bias	Low risk	No other sources of bias were identified.

PEMF: pulsed electromagnetic fields but no resistance training

PNF: proprioceptive neuromuscular facilitation

ROM: range of movement

RT: resistance training

6MWT: 6-minute walk test

Characteristics of excluded studies *[ordered by study ID]*

Study	Reason for exclusion
Czepa 2013	Not randomized.
Greene 1983	No control group.
Hilberg 2003	No control group.
Khriesat 2000	No control group.
Mulvany 2010	No control group.
Vallejo 2010	No control group.
Von Mackensen 2012	Not randomized.

Characteristics of studies awaiting assessment *[ordered by study ID]*

[Cuesta-Barriuso 2014](#)

Methods	Randomized study.
Participants	31 participants with haemophilia A or B and with haemophilic arthropathy in one or both ankles.
Interventions	Manual therapy group (articular traction, passive stretching of the gastrocnemius muscles, and exercises for muscle strength and proprioception), an educational group (educational sessions and home exercises) and a control group.
Outcomes	Range of motion, gastrocnemius muscle circumference, muscular strength of the gastrocnemius muscles, pain perception, radiological assessment of joint deterioration.
Notes	

[Cuesta-Barriuso 2015](#)

Methods	Randomized study.
Participants	People with haemophilia.
Interventions	Experimental group: received training sessions on the clinical features of haemophilia and the management of musculoskeletal limitations of haemophilic arthropathy, plus home exercises. Control group: no intervention.
Outcomes	Physical state (Gilbert score), A36 haemophilia-QoL and IBQ.
Notes	

Firoozabadi 2012

Methods	Quasi-experimental, cross-sectional study.
Participants	20 participants with haemophilia A from the Isfahan Hemophiliac Center.
Interventions	Exercise therapy program versus a "control group".
Outcomes	6MWT; Beck Anxiety Inventory; range of movement measurements were done on 10 joints.
Notes	

IBQ: illness behaviour questionnaire

QoL: quality of life

6MWT: 6-minute walk test

DATA AND ANALYSES

Comparison 1. Exercise versus no intervention

Outcome or sub-group title	No. of studies	No. of participants	Statistical method	Effect size
1 Colorado score	1		Mean Difference (IV, Fixed, 95% CI)	Totals not selected
1.1 elbow	1		Mean Difference (IV, Fixed, 95% CI)	0.0 [0.0, 0.0]
1.2 knee	1		Mean Difference (IV, Fixed, 95% CI)	0.0 [0.0, 0.0]
1.3 ankles	1		Mean Difference (IV, Fixed, 95% CI)	0.0 [0.0, 0.0]

Analysis 1.1. Comparison 1 Exercise versus no intervention, Outcome 1 Colorado score.

Study or subgroup	Exercise		No intervention		Mean Difference Fixed, 95% CI	Mean Difference Fixed, 95% CI
	N	Mean(SD)	N	Mean(SD)		
1.1.1 elbow						
Parhampour 2014	13	-1.2 (1.2)	12	-0 (1.2)		-1.21[-2.14,-0.28]
1.1.2 knee						
Parhampour 2014	13	-3.8 (1.6)	12	-0.4 (2)		-3.42[-4.82,-2.02]
1.1.3 ankles						
Parhampour 2014	13	-1.6 (1.1)	12	0 (1.1)		-1.65[-2.51,-0.79]
					-5 -2.5 0 2.5 5	
					Favours exercise	Favours no intervention

Comparison 2. Exercise A versus Exercise B

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Balance (Biodex Stability System)	1		Mean Difference (IV, Fixed, 95% CI)	Totals not selected
1.1 overall stability index	1		Mean Difference (IV, Fixed, 95% CI)	0.0 [0.0, 0.0]
1.2 anterior-posterior stability index	1		Mean Difference (IV, Fixed, 95% CI)	0.0 [0.0, 0.0]
1.3 medio-lateral stability index	1		Mean Difference (IV, Fixed, 95% CI)	0.0 [0.0, 0.0]
2 Pain	1		Mean Difference (IV, Fixed, 95% CI)	Totals not selected
3 6-min walk: distance	1		Mean Difference (IV, Fixed, 95% CI)	Totals not selected
4 Range of motion: flexion	1		Mean Difference (IV, Fixed, 95% CI)	Totals not selected
4.1 flexion	1		Mean Difference (IV, Fixed, 95% CI)	0.0 [0.0, 0.0]
4.2 extension	1		Mean Difference (IV, Fixed, 95% CI)	0.0 [0.0, 0.0]
5 Muscle strength	3		Mean Difference (IV, Random, 95% CI)	Subtotals only
5.1 extensors (force in kg)	1	30	Mean Difference (IV, Random, 95% CI)	6.40 [4.81, 7.99]
5.2 extensors 60 degrees per second	1	30	Mean Difference (IV, Random, 95% CI)	4.31 [2.75, 5.87]
5.3 extensors 120 degrees per second	2	60	Mean Difference (IV, Random, 95% CI)	12.75 [1.46, 24.04]
5.4 flexors speed 60 degrees per second	1	30	Mean Difference (IV, Random, 95% CI)	4.31 [3.20, 5.42]
5.5 flexors speed 120 degrees per second	2	60	Mean Difference (IV, Random, 95% CI)	9.12 [6.74, 11.51]
6 Knee circumference	1		Mean Difference (IV, Fixed, 95% CI)	Totals not selected

Analysis 2.1. Comparison 2 Exercise A versus Exercise B, Outcome 1 Balance (Biodex Stability System).

Study or subgroup	Exercise A (bicycle)		Exercise B (treadmill)		Mean Difference	Mean Difference
	N	Mean(SD)	N	Mean(SD)	Fixed, 95% CI	Fixed, 95% CI
2.1.1 overall stability index						
Mohamed 2015	15	-0.4 (0.1)	15	-0.6 (0.1)		0.25[0.19,0.31]
2.1.2 anterior-posterior stability index						
Mohamed 2015	15	-0.4 (0.1)	15	-0.8 (0.1)		0.35[0.27,0.43]
					Favours Exercise A -0.5 -0.25 0 0.25 0.5 Favours Exercise B	

Study or subgroup	Exercise A (bicycle)		Exercise B (treadmill)		Mean Difference	Mean Difference
	N	Mean(SD)	N	Mean(SD)	Fixed, 95% CI	Fixed, 95% CI
2.1.3 medio-lateral stability index						
Mohamed 2015	15	-0.4 (0.1)	15	-0.6 (0.1)		0.24[0.17,0.31]
					Favours Exercise A	Favours Exercise B

Analysis 2.2. Comparison 2 Exercise A versus Exercise B, Outcome 2 Pain.

Study or subgroup	Exercise A		Exercise B		Mean Difference	Mean Difference
	N	Mean(SD)	N	Mean(SD)	Fixed, 95% CI	Fixed, 95% CI
Mazloun 2014	13	-0.9 (0.5)	14	-1.7 (0.5)		0.8[0.41,1.19]
					Favours Exercise A	Favours Exercise B

Analysis 2.3. Comparison 2 Exercise A versus Exercise B, Outcome 3 6-min walk: distance.

Study or subgroup	Exercise A		Exercise B		Mean Difference	Mean Difference
	N	Mean(SD)	N	Mean(SD)	Fixed, 95% CI	Fixed, 95% CI
Zaky 2013	15	34.6 (3)	15	32 (4)		2.6[0.08,5.12]
					Favours Exercise B	Favours Exercise A

Analysis 2.4. Comparison 2 Exercise A versus Exercise B, Outcome 4 Range of motion: flexion.

Study or subgroup	Exercise A		Exercise B		Mean Difference	Mean Difference
	N	Mean(SD)	N	Mean(SD)	Fixed, 95% CI	Fixed, 95% CI
2.4.1 flexion						
Mazloun 2014	13	7.2 (9.4)	14	7 (5.3)		0.2[-5.61,6.01]
2.4.2 extension						
Mazloun 2014	13	1.9 (2)	14	2 (2)		-0.1[-1.59,1.39]
					Favours Exercise B	Favours Exercise A

Analysis 2.5. Comparison 2 Exercise A versus Exercise B, Outcome 5 Muscle strength.

Study or subgroup	Exercise A		Exercise B		Mean Difference	Weight	Mean Difference
	N	Mean(SD)	N	Mean(SD)	Random, 95% CI		Random, 95% CI
2.5.1 extensors (force in kg)							
Zaky 2013	15	17.6 (2.6)	15	11.2 (1.8)		100%	6.4[4.81,7.99]
Subtotal ***	15		15			100%	6.4[4.81,7.99]
Heterogeneity: Not applicable							
Test for overall effect: Z=7.87(P<0.0001)							
2.5.2 extensors 60 degrees per second							
Abd-Elmonem 2014	15	9.2 (2.7)	15	4.9 (1.5)		100%	4.31[2.75,5.87]
Subtotal ***	15		15			100%	4.31[2.75,5.87]
					Favours Exercise B		Favours Exercise A

Analysis 2.6. Comparison 2 Exercise A versus Exercise B, Outcome 6 Knee circumference.

Comparison 3. Exercise versus Exercise plus electrophysical modality

Outcome or sub-group title	No. of studies	No. of participants	Statistical method	Effect size
1 Colorado score	1		Mean Difference (IV, Fixed, 95% CI)	Totals not selected
1.1 ankle	1		Mean Difference (IV, Fixed, 95% CI)	0.0 [0.0, 0.0]
1.2 elbow	1		Mean Difference (IV, Fixed, 95% CI)	0.0 [0.0, 0.0]
1.3 knee	1		Mean Difference (IV, Fixed, 95% CI)	0.0 [0.0, 0.0]

Analysis 3.1. Comparison 3 Exercise versus Exercise plus electrophysical modality, Outcome 1 Colorado score.

Comparison 4. Hydrotherapy versus no intervention

Outcome or sub-group title	No. of studies	No. of participants	Statistical method	Effect size
1 Pain	1		Mean Difference (IV, Fixed, 95% CI)	Totals not selected
2 Range of motion	1		Mean Difference (IV, Fixed, 95% CI)	Totals not selected
2.1 flexion	1		Mean Difference (IV, Fixed, 95% CI)	0.0 [0.0, 0.0]
2.2 extension	1		Mean Difference (IV, Fixed, 95% CI)	0.0 [0.0, 0.0]

Analysis 4.1. Comparison 4 Hydrotherapy versus no intervention, Outcome 1 Pain.

Analysis 4.2. Comparison 4 Hydrotherapy versus no intervention, Outcome 2 Range of motion.

ADDITIONAL TABLES

Table 1. Exercise versus no intervention - Additional data (Cuesta-Barriuso 2013)

Outcome	Assessment	Exercise group: mean (SD) (n = 9 participants, 16 elbows)	Control group: mean (SD) (n = 9 participants, 14 elbows)
Flexion	Pre-treatment:	136.130 (14.818)	138.000 (19.958)
	Post-treatment (12 weeks):	140.310 (11.247)	138.430 (18.475)
Extension	Pre-treatment:	10.870 (14.207)	8.640 (15.310)
	Post-treatment (12 weeks):	11.880 (14.917)	8.290 (15.364)
Pain (VAS)	Pre-treatment:	0.719 (0.752)	0.143 (0.305)
	Post-treatment (12 weeks):	0.344 (0.436)	0.071 (0.267)
Biceps perimeter	Pre-treatment:	31.331 (3.474)	32.007 (3.837)
	Post-treatment (12 weeks):	31.725 (3.205)	31.857 (3.566)
Bicep strength	Pre-treatment:	0.094 (0.272)	0.143 (0.363)
	Post-treatment (12 weeks):	0.000 (0.000)	0.071 (0.181)

Number of participants in each group = 9. Data relate to 16 elbows in the exercise group and 14 in the control group.
VAS: visual analog scale

Table 2. Exercise A vs Exercise B: types of exercise programs used

Author	"Exercise A"	"Exercise B"
Abd-Elmonem 2014	Ultrasound, stretching, strengthening 5 days per week	Exercise A plus treadmill training
Eid 2014	Stretching, static exercises, treadmill	Exercise A plus bicycle ergometer and resisted isotonic exercises
Zaky 2013	Static and short-arc quadriceps, straight leg raises	Exercise A plus partial weight-bearing exercises in supine
Mazloun 2014	Stretching, isometric and isotonic strengthening	Rhythmic movement in water, isometric and isotonic strengthening
Mohamed 2015	Stretching, isometrics, balance or gait training; bicycle ergometry	Stretching, isometrics, balance or gait training; treadmill

APPENDICES

Appendix 1. Search Strategies

Database/ Register	Years of Coverage	Search terms
PubMed (www.ncbi.nlm.nih.gov/pubmed)	1966 - present	#1 hemophilia* AND exercise [Filters: Clinical Trial; Humans]
Embase (Ovid)	1976 - present	#1 exercise.mp. OR exercise/ #2 exp hemophilia A/ OR hemophilia.mp. OR exp hemophilia/ OR exp hemophilia B/ #3 1 AND 2
CINAHL (Ebsco) (www.ebscohost.com/)	1961 - present	#1Exercise AND Hemophilia
ClinicalTrials.gov (http://www.clinicaltrials.gov/)	Conception - present	hemophilia* AND exercise
International Clinical Trials Registry Platform (http://apps.who.int/trialsearch/)		
EU Clinical Trials Register (www.clinicaltrialsregister.eu)		

Appendix 2. Conference Abstract Book Searches

Hand searches were done by the authors in the Abstract Documents of the Congress of the World Federation of Hemophilia and the Musculoskeletal Congresses of WFH as described below.

WFH World Congresses:

1996 - Haemophilia Volume 2 supplement 1. Plenary abstracts published. No musculoskeletal (MSK) topics included.

1998 - Haemophilia Volume 4:3. Abstract headings listed in Table of Contents. No MSK topics included.

2000 - Haemophilia Volume 6:4 Medium: printed abstracts. Manually searched.

2002 - Medium: Abstracts on CD-ROM in PDF format. Search strategy: Find "exercise". Results: a few with exercise in the title. Several with 'exercise' in the body. Not all relevant to physical exercise.

2004 - Haemophilia 10 Supplement 3. Medium: printed abstracts. Hand Search of Abstracts in sections 16- Musculoskeletal, 21- Physiotherapy, and 25-Rehabilitation.

2006 - Haemophilia 12: Supplement 2. Medium: printed abstracts. Manually searched Sections 18- Musculoskeletal Issues, 22- Physiotherapy, and 27- Rehabilitation.

2008 - Haemophilia 14: Supplement 2 Medium: printed abstracts. Manually searched Musculoskeletal Section. (There were no PT or Rehab sections this year). 3 abstracts selected.

2010 - Medium: Flash Drive, Adobe PDF document. Keyword Search "exercise" found 58 instances in 23 abstracts. Only 3 articles described studies of exercise interventions.

2012 - Medium: Flash Drive, Adobe PDF document. Keyword Search "exercise" found 109 instances in 46 abstracts. Six described studies of exercise interventions.

2014 - Haemophilia 20 Supplement 3. Medium: PDF document, accessed on line at www.WFH.org. Search strategy: Find "exercise". Seven articles described exercise interventions.

WFH - Musculoskeletal Congresses:

2005 - Medium: final program abstract booklet. Strategy: Hand search. Results: no suitable abstracts.

2007 - Medium: Word Document, Abstracts. Search strategy: Find "exercise". Results: 20 matches in 7 articles. Only one article remotely relevant.

Exercise for haemophilia (Review)

Copyright © 2016 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

2009 - Medium: final program abstract booklet. Strategy: Hand search. Results: no studies. (Only reviews and case reports).

2011 - Medium: PDF of final program and abstract booklet. Strategy: hand search and Find "exercise". Results: 6 possible studies.

2013 - Medium: PDF of final program and abstract booklet. Strategy: hand search and Find "exercise". Results: 7 possible studies.

CONTRIBUTIONS OF AUTHORS

RM and KM developed the protocol. KS and KM conducted the search for studies and selected included trials; assessed study quality; extracted, entered and analyzed the data. KS interpreted the results. KS and KM wrote the review and RM reviewed the final manuscript.

DECLARATIONS OF INTEREST

Karen Strike: none known.

Kathy Mulder: none known.

Roger Michael: none known.

SOURCES OF SUPPORT

Internal sources

- No sources of support supplied

External sources

- National Institute for Health Research, UK.

This systematic review was supported by the National Institute for Health Research, via Cochrane Infrastructure funding to the Cochrane Cystic Fibrosis and Genetic Disorders Group.

DIFFERENCES BETWEEN PROTOCOL AND REVIEW

Post hoc analysis resulted in the addition of the following clinically relevant outcomes. These outcomes were deemed important to healthcare providers and consumers as primary impairments that are assessed by musculoskeletal specialists and may be impacted by an exercise intervention.

1. Range of motion ([Kargarfard 2013](#); [Mazloun 2014](#)).
2. Biceps perimeter ([Cuesta-Barriuso 2013](#)).
3. Strength: peak torque using the Biodex Isokinetic Dynamometer ([Abd-Elmonem 2014](#); [Eid 2014](#); [Kargarfard 2013](#)) and the Lafayette Manual Muscle Test ([Zaky 2013](#)).
4. Knee circumference ([Abd-Elmonem 2014](#)).

INDEX TERMS

Medical Subject Headings (MeSH)

Exercise Therapy [*methods]; Hemarthrosis [rehabilitation]; Hemophilia A [*rehabilitation]; Hemophilia B [*rehabilitation]; Hydrotherapy; Muscle Strength; Muscle Stretching Exercises; Pain [rehabilitation]; Pain Measurement; Randomized Controlled Trials as Topic; Range of Motion, Articular; Resistance Training

MeSH check words

Adolescent; Adult; Child; Humans; Male; Middle Aged