Cochrane
Library

Cochrane Database of Systematic Reviews

Oxygen for breathlessness in patients with chronic obstructive pulmonary disease who do not qualify for home oxygen therapy (Review)

Ekström M, Ahmadi Z, Bornefalk-Hermansson A, Abernethy A, Currow D

Ekström M, Ahmadi Z, Bornefalk-Hermansson A, Abernethy A, Currow D.
Oxygen for breathlessness in patients with chronic obstructive pulmonary disease who do not qualify for home oxygen therapy.
Cochrane Database of Systematic Reviews 2016, Issue 11. Art. No.: CD006429.
DOI: [10.1002/14651858.CD006429.pub3](https://doi.org/10.1002/14651858.CD006429.pub3).

www.cochranelibrary.com

Oxygen for breathlessness in patients with chronic obstructive pulmonary disease who do not qualify for home oxygen therapy (Review)

Copyright © 2017 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

WILEY

TABLE OF CONTENTS

ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
SUMMARY OF FINDINGS	3
BACKGROUND	5
OBJECTIVES	5
METHODS	5
RESULTS	7
Figure 1.	9
Figure 2.	11
Figure 3.	13
Figure 4.	15
Figure 5.	16
Figure 6.	17
DISCUSSION	19
AUTHORS' CONCLUSIONS	21
ACKNOWLEDGEMENTS	21
REFERENCES	22
CHARACTERISTICS OF STUDIES	28
DATA AND ANALYSES	85
Analysis 1.1. Comparison 1 Oxygen versus air, Outcome 1 Breathlessness - all trials.	86
Analysis 1.2. Comparison 1 Oxygen versus air, Outcome 2 Breathlessness - subgroup analysis - short-burst oxygen vs not.	87
Analysis 1.3. Comparison 1 Oxygen versus air, Outcome 3 Breathlessness - subgroup analysis - exertional desaturation vs not. .	88
Analysis 1.4. Comparison 1 Oxygen versus air, Outcome 4 Breathlessness - subgroup analysis - mean PaO ₂ < 9.3 kPa vs higher. .	89
Analysis 1.5. Comparison 1 Oxygen versus air, Outcome 5 Breathlessness - subgroup analysis - measured during exercise test vs not.	90
Analysis 1.6. Comparison 1 Oxygen versus air, Outcome 6 Breathlessness - subgroup analysis - laboratory vs other.	91
Analysis 1.7. Comparison 1 Oxygen versus air, Outcome 7 Breathlessness - subgroup analysis - short-term vs long-term (training) effect of oxygen.	92
Analysis 1.8. Comparison 1 Oxygen versus air, Outcome 8 Breathlessness - subgroup analysis - mean oxygen dose > 2 L/min vs lower.	93
Analysis 1.9. Comparison 1 Oxygen versus air, Outcome 9 Breathlessness - sensitivity analysis - excluding measurements at peak exertion.	94
Analysis 1.10. Comparison 1 Oxygen versus air, Outcome 10 Breathlessness - sensitivity analysis - excluding studies with high risk of bias.	95
Analysis 1.11. Comparison 1 Oxygen versus air, Outcome 11 Breathlessness - sensitivity analysis - excluding outliers.	96
Analysis 1.12. Comparison 1 Oxygen versus air, Outcome 12 Breathlessness - sensitivity analysis - post hoc - excluding short-burst and outliers.	97
Analysis 1.13. Comparison 1 Oxygen versus air, Outcome 13 Health-related quality of life - all trials.	97
ADDITIONAL TABLES	99
APPENDICES	104
WHAT'S NEW	108
HISTORY	108
CONTRIBUTIONS OF AUTHORS	109
DECLARATIONS OF INTEREST	109
SOURCES OF SUPPORT	109
DIFFERENCES BETWEEN PROTOCOL AND REVIEW	109
INDEX TERMS	109

[Intervention Review]

Oxygen for breathlessness in patients with chronic obstructive pulmonary disease who do not qualify for home oxygen therapy

Magnus Ekström¹, Zainab Ahmadi¹, Anna Bornefalk-Hermansson², Amy Abernethy³, David Currow⁴

¹Department of Clinical Sciences, Division of Respiratory Medicine & Allergology, Lund University, Lund, Sweden. ²Department of Statistics, Uppsala University, Uppsala, Sweden. ³CLHC, Duke University Medical Center, Durham, North Carolina, USA. ⁴Department of Palliative and Supportive Services, Flinders University, Daw Park, Australia

Contact: Magnus Ekström, Department of Clinical Sciences, Division of Respiratory Medicine & Allergology, Lund University, Lund, Sweden. pmekstrom@gmail.com.

Editorial group: Cochrane Airways Group.

Publication status and date: Edited (no change to conclusions), published in Issue 8, 2017.

Citation: Ekström M, Ahmadi Z, Bornefalk-Hermansson A, Abernethy A, Currow D. Oxygen for breathlessness in patients with chronic obstructive pulmonary disease who do not qualify for home oxygen therapy. *Cochrane Database of Systematic Reviews* 2016, Issue 11. Art. No.: CD006429. DOI: [10.1002/14651858.CD006429.pub3](https://doi.org/10.1002/14651858.CD006429.pub3).

Copyright © 2017 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

Breathlessness is a cardinal symptom of chronic obstructive pulmonary disease (COPD). Long-term oxygen therapy (LTOT) is given to improve survival time in people with COPD and severe chronic hypoxaemia at rest. The efficacy of oxygen therapy for breathlessness and health-related quality of life (HRQOL) in people with COPD and mild or no hypoxaemia who do not meet the criteria for LTOT has not been established.

Objectives

To determine the efficacy of oxygen versus air in mildly hypoxaemic or non-hypoxaemic patients with COPD in terms of (1) breathlessness; (2) HRQOL; (3) patient preference whether to continue therapy; and (4) oxygen-related adverse events.

Search methods

We searched the Cochrane Airways Group Register, the Cochrane Central Register of Controlled Trials (CENTRAL), MEDLINE and Embase, to 12 July 2016, for randomised controlled trials (RCTs). We handsearched the reference lists of included articles.

Selection criteria

We included RCTs of the effects of non-invasive oxygen versus air on breathlessness, HRQOL or patient preference to continue therapy among people with COPD and mild or no hypoxaemia (partial pressure of oxygen (PaO₂) > 7.3 kPa) who were not already receiving LTOT. Two review authors independently assessed articles for inclusion in the review.

Data collection and analysis

Two review authors independently collected and analysed data. We assessed risk of bias by using the Cochrane 'Risk of bias tool'. We pooled effects recorded on different scales as standardised mean differences (SMDs) with 95% confidence intervals (CIs) using random-effects models. Lower SMDs indicated decreased breathlessness and reduced HRQOL. We performed subanalyses and sensitivity analyses and assessed the quality of evidence according to the Grading of Recommendations, Assessment, Development and Evaluations (GRADE) approach.

Main results

Compared with the previous review, which was published in 2011, we included 14 additional studies (493 participants), excluded one study and included data for meta-analysis of HRQOL. In total, we included in this review 44 studies including 1195 participants, and we included 33 of these (901 participants) in the meta-analysis.

We found that breathlessness during exercise or daily activities was reduced by oxygen compared with air (32 studies; 865 participants; SMD -0.31, 95% CI -0.43 to -0.20; $I^2 = 29\%$; low-quality evidence). This translates to a decrease in breathlessness of about 0.7 points on a 0 to 10 numerical rating scale. In contrast, we found no effect of short-burst oxygen given before exercise (four studies; 90 participants; SMD -0.03, 95% CI -0.28 to 0.22; $I^2 = 0\%$; low-quality evidence). Oxygen reduced breathlessness measured during exercise tests (30 studies; 591 participants; SMD -0.34, 95% CI -0.46 to -0.22; $I^2 = 29\%$; moderate-quality evidence), whereas evidence of an effect on breathlessness measured in daily life was limited (two studies; 274 participants; SMD -0.13, 95% CI, -0.37 to 0.11; $I^2 = 0\%$; low-quality evidence).

Oxygen did not clearly affect HRQOL (five studies; 267 participants; SMD 0.12, 95% CI -0.04 to 0.28; $I^2 = 0\%$; low-quality evidence). Patient preference and adverse events could not be analysed owing to insufficient data.

Authors' conclusions

We are moderately confident that oxygen can relieve breathlessness when given during exercise to mildly hypoxaemic and non-hypoxaemic people with chronic obstructive pulmonary disease who would not otherwise qualify for home oxygen therapy. Most evidence pertains to acute effects during exercise tests, and no evidence indicates that oxygen decreases breathlessness in the daily life setting. Findings show that oxygen does not affect health-related quality of life.

PLAIN LANGUAGE SUMMARY

Oxygen therapy for breathless people with chronic obstructive pulmonary disease with only mildly or moderately decreased oxygen in the blood

Review question

We reviewed the evidence regarding effects of oxygen compared with air on breathlessness in people with chronic obstructive pulmonary disease (COPD) with only mildly or moderately decreased blood oxygen levels.

Background

People with COPD are sometimes prescribed oxygen therapy to reduce the severity of breathlessness. However, the use of oxygen in people who do not have severely reduced levels of oxygen in their bloodstream remains controversial, as little is known about its effectiveness. Additionally, oxygen is relatively costly and is not given without risk, particularly to smokers because of the risk of fire.

Study characteristics

We examined the research published to 12 July 2016. We included studies of oxygen therapy versus air delivered through nasal prongs or mask during exertion, continuously, 'as needed' over a defined period or as short-burst oxygen before exertion. Study participants were 18 years of age or older, had received a diagnosis of COPD, had low oxygen levels in the blood and did not receive long-term oxygen therapy. We included a total of 44 studies (1195 participants) in this review. Compared with the previous review, which was published in 2011, we have added 14 studies (493 participants) to this review.

Key results

We found that oxygen can modestly reduce breathlessness. To be effective, oxygen has to be given during exercise. Most studies evaluated oxygen given during exercise testing in the laboratory. Oxygen therapy during daily life had uncertain effects on breathlessness and did not clearly change patient quality of life.

Quality of the evidence

We rated the quality of evidence using one of the following grades: very low, low, moderate or high. For very low-quality evidence, we were uncertain about the results. With high-quality evidence, we were very certain about the results. We found that evidence for oxygen given for breathlessness was moderate to low. We are moderately confident that oxygen can relieve breathlessness when given during exercise to people with COPD with mildly or moderately decreased blood oxygen levels. However, most studies reported acute effects during an exercise test, and no evidence suggests that oxygen decreases breathlessness during daily life. Findings indicate that oxygen does not affect health-related quality of life.

SUMMARY OF FINDINGS

Summary of findings for the main comparison. Summary of findings table

Oxygen for breathlessness in patients with chronic obstructive pulmonary disease who do not qualify for home oxygen therapy

Patient or population: patients with chronic obstructive pulmonary disease who do not qualify for home oxygen therapy

Intervention: oxygen delivered through a non-invasive method

Comparison: air delivered through the same non-invasive method

Outcomes	Difference (95% CI)	Number of participants (studies)	Quality of the evidence (GRADE)	Comments
Breathlessness - all trials Lower score indicates improvement in breathlessness	SMD 0.31, SD lower (0.43 lower to 0.2 lower)	865 (32)	⊕⊕⊕⊕ Low ^a	This corresponds to 0.65 points lower (0.90 lower to 0.42 lower) on a 0-10 NRS.*
Breathlessness - subgroup analysis - studies using short-burst oxygen Lower score indicates improvement in breathlessness	SMD 0.03, SD lower (0.28 lower to 0.22 higher)	90 (4)	⊕⊕⊕⊕ Low ^b	This corresponds to 0.06 points lower (0.59 lower to 0.46 higher) on a 0-10 NRS.*
Breathlessness - subgroup analysis - studies not using short-burst oxygen Lower score indicates improvement in breathlessness	SMD 0.36, SD lower (0.48 lower to 0.24 lower)	775 (28)	⊕⊕⊕⊕ Low ^a	This corresponds to 0.76 points lower (1.01 lower to 0.50 lower) on a 0-10 NRS.*
Breathlessness - subgroup analysis - studies measuring during exercise test Lower score indicates improvement in breathlessness	SMD 0.34, SD lower (0.46 lower to 0.22 lower)	591 (30)	⊕⊕⊕⊕ Moderate ^c	This corresponds to 0.71 points lower (0.97 lower to 0.46 lower) on a 0-10 NRS.*
Breathlessness - subgroup analysis - studies not measuring during exercise test Lower score indicates improvement in breathlessness	SMD 0.13, SD lower (0.37 lower to 0.11 higher)	274 (2)	⊕⊕⊕⊕ Low ^b	This corresponds to 0.27 points lower (0.78 lower to 0.23 higher) on a 0-10 NRS.*
Health-related quality of life - all trials Higher scores indicate improved health-related quality of life	SMD 0.12, SD higher (0.04 lower to 0.28 higher)	267 (5)	⊕⊕⊕⊕ Low ^b	This corresponds to 0.25 points higher (0.09 lower to 0.59 higher) on a 0-10 NRS.*

*Difference on a 0-10 NRS calculated using the SD of 2.1 for the COPD group in [Abernethy 2010](#) for individual participant data.

CI: confidence interval; OR: odds ratio; RR: risk ratio.

GRADE Working Group grades of evidence

High quality: We are very confident that the true effect lies close to that of the estimate of effect.

Moderate quality: We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of effect but may be substantially different.

Low quality: Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of effect.

Very low quality: We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect.

^aAlthough the effect was consistent in exercise tests in the laboratory setting, evidence of an effect was limited for domiciliary oxygen in daily life.

^bFew studies.

^cGraded as moderate, as most evidence pertained to breathlessness during exercise tests.

BACKGROUND

Description of the condition

Breathlessness, a cardinal symptom of chronic obstructive pulmonary disease (COPD), is distressing to both patients and caregivers. Breathlessness is defined as "a subjective experience of breathing discomfort that consists of qualitatively distinct sensations that vary in intensity" (Parshall 2012). Breathlessness is multi-factorial (Laviolette 2014) and arises as a combination of underlying pathologies involving neural pathways and subjective central perception (Parshall 2012).

Description of the intervention

Treatment of patients with breathlessness includes management of the underlying cause(s). However, breathlessness often persists despite optimal management of underlying disease(s), that is, chronic breathlessness. Pharmacological treatment with the strongest evidence base for relieving breathlessness consists of low-dose opioids (Ekstrom 2015a). People with insufficiently relieved breathlessness are then left to try any of a number of interventions for which evidence is limited, such as supplemental oxygen therapy.

Supplemental oxygen therapy most often is administered through nasal prongs or a mask and may be prescribed continuously, during exercise or as a short burst before exercise (Hardinge 2015). Oxygen sources include oxygen concentrators (which concentrate oxygen from ambient air), cylinders of compressed oxygen and flasks of liquid oxygen. Equipment can be stationary in the home and/or portable (Hardinge 2015).

Long-term oxygen therapy (LTOT) refers to oxygen given for 15 or more hours per day to prolong survival time in people with COPD and chronic severe resting hypoxaemia (partial pressure of oxygen (PaO_2) < 7.4 kPa) or moderate hypoxaemia (PaO_2 7.4 to 7.8 kPa), together with signs of right-sided heart failure or secondary polycythemia (Cranston 2005; MRCWP 1981; NOTTG 1980). Studies in severe hypoxaemia have not evaluated whether LTOT relieves breathlessness or improves health-related quality of life (HRQOL) (Cranston 2008).

Palliative oxygen therapy refers to supplemental oxygen given to relieve symptoms in people with COPD who have no to moderate hypoxaemia.

How the intervention might work

Oxygen given before or during physical activity might increase oxygen content of and oxygen delivery to exercising muscles. This might prolong aerobic metabolism during exertion and may decrease muscle fatigue and formation of lactic acid (O'Donnell 2001), which could lead to a decreased level of ventilatory drive and increased ventilatory capacity at a given work rate, thereby decreasing the mismatch between ventilatory drive and work and perceived severity of breathlessness (O'Donnell 2001; Parshall 2012).

Why it is important to do this review

A Cochrane review of the efficacy of palliative oxygen for breathlessness was published in 2011 (Uronis 2011; Uronis 2015). A meta-analysis of 18 trials (431 participants) reported

that supplemental oxygen during exercise and activities reduced breathlessness to a greater extent than air. Short-burst oxygen before exercise did not decrease breathlessness. Review authors could not analyse HRQOL because data were insufficient. Most evidence pertained to exercise testing performed in the laboratory setting (Uronis 2011; Uronis 2015). A subsequent Cochrane review of four studies (331 participants) suggested that longer-term treatment with ambulatory oxygen relieves breathlessness more effectively than air after exercise in people with COPD (Ameer 2014). Current guidelines do not recommend palliative oxygen therapy in COPD, as evidence on treatment in the domiciliary and daily life setting remains insufficient (Ekstrom 2015b; Hardinge 2015). Despite this, palliative oxygen therapy might be commonly prescribed in clinical practice (Abernethy 2005; Stringer 2004).

The discrepancy between current clinical practice and available evidence has important implications. First, patients may be prescribed ineffective treatment. Second, oxygen therapy is not a benign intervention. Functional restriction caused by tubing, concentrators or cylinders and the "sick role" may limit quality of life. People also express concern about being reliant on a machine. Nasal cannulae can lead to nasal irritation and can increase the risk of nosebleeds. Oxygen therapy carries a fire risk, particularly for smokers (Robb 2003). Serious burn injuries seem infrequent in LTOT with strict adherence to eligibility criteria and contraindications, including smoking cessation (Tanash 2015). A small but increased risk of exacerbated hypercarbia is possible, and home oxygen therapy is relatively expensive. If patients do not meet the criteria for LTOT, they may have to pay for oxygen therapy themselves or may receive it on compassionate use grounds (Guyatt 2000). Improved evidence regarding optimal use of palliative oxygen therapy is needed.

Since the previous Cochrane reviews were published (Uronis 2011; Uronis 2015), several studies of palliative oxygen therapy have been reported, including some larger trials (Abernethy 2010 and Moore 2011).

This is an update of the Cochrane review published in 2011 (Uronis 2011; Uronis 2015).

OBJECTIVES

To determine the efficacy of oxygen versus air in mildly hypoxaemic or non-hypoxaemic patients with COPD in terms of (1) breathlessness; (2) HRQOL; (3) patient preference whether to continue therapy; and (4) oxygen-related adverse events.

METHODS

Criteria for considering studies for this review

Types of studies

We included randomised controlled trials (RCTs).

Types of participants

We included studies with participants 18 years of age or older who had COPD, had mild or no hypoxaemia (mean PaO_2 > 7.3 kPa) and did not receive LTOT. For studies that also included participants without COPD, we obtained from study authors individual participant data for those with COPD and included only these data in the analyses.

Types of interventions

We included studies of oxygen therapy versus air delivered by a non-invasive method at any inspired dose above that of ambient air (> 21%). Oxygen/air should have been delivered during exertion, continuously or 'as needed' over a defined period, or as short-burst oxygen before exertion. Short-burst oxygen was defined as therapy given during a short, defined period just before exertion. We did not include short-burst oxygen given only after exertion.

Types of outcome measures

Primary outcomes

1. Level of breathlessness measured on any validated scale

Secondary outcomes

1. HRQOL measured on any validated scale
2. Blinded participant preference to continue therapy
3. Adverse events

Search methods for identification of studies

Electronic searches

We searched the Cochrane Airways Group Specialised Register of trials, which is derived from systematic searches of multiple bibliographic databases and from handsearching of respiratory journals and meeting abstracts (see [Appendix 1](#) for details). We conducted additional searches of the Cochrane Central Register of Controlled Trials (CENTRAL; 2016, Issue 6), MEDLINE (to 12 July 2016) and Embase (to 12 July 2016). Search strategies are shown in [Appendix 2](#); [Appendix 3](#); [Appendix 4](#); and [Appendix 5](#). We also searched ClinicalTrials.gov (www.ClinicalTrials.gov) and the World Health Organization (WHO) trials portal (www.who.int/ictrp/en/) for ongoing trials by using appropriately adapted search terms. We searched all databases from their inception to 12 July 2016, with no restriction on language of publication.

Searching other resources

We handsearched relevant review articles and reference lists of identified articles.

Data collection and analysis

Selection of studies

Two review authors (ME and ZA) independently selected studies that fulfilled all inclusion criteria.

1. RCT.
2. Included participants 18 years of age or older.
3. At least one participant with COPD.
4. Mean PaO₂ > 7.3 kPa at baseline.
5. Only participants not already receiving LTOT.
6. Comparison of oxygen versus air at any dose delivered through a non-invasive method.
7. Endpoint of breathlessness or HRQOL.

Data extraction and management

Two review authors (ME and ZA) independently extracted data and resolved disagreements by consensus and discussion with a third

review author (DC), if needed. We contacted the authors of primary studies to obtain additional information when necessary.

Assessment of risk of bias in included studies

Two review authors (ME and ZA) independently assessed each study for risk of bias in terms of allocation sequence generation, allocation concealment, blinding of participants and outcome assessors, handling of missing data, selective outcome reporting and other threats to the validity of studies, in line with recommendations provided in the *Cochrane Handbook for Systematic Reviews of Interventions* ([Higgins 2011](#)). We conducted a retrospective risk of bias assessment using the above method applied to all studies from the review published in 2011 ([Uronis 2011](#)).

Measures of treatment effect

We performed meta-analyses as appropriate and possible, given available data. We analysed outcomes on different scales as standardised mean differences (SMDs) using random-effects models. We compared within-patient effects from both periods of cross-over trials. Meta-analyses included post scores only. For studies for which post scores were not available, we reported changes from baseline scores separately. For breathlessness during exercise, we used scores measured at a similar time point in both groups (iso-scores) when available. For studies evaluating different oxygen doses, we included only the lowest dose in the analysis. We reversed St George's Respiratory Quotient (SGRQ) scores (as increasing scores indicate worse quality of life), so that higher scores on all HRQOL measures indicated better quality of life in all analyses.

Unit of analysis issues

We estimated standard errors for paired outcome data from cross-over trials, according to [Follmann 1992](#). We estimated correlations between repeated outcomes from P values when available, and for studies for which little evidence was available to impute a correlation coefficient from another trial, we assumed a value of 0.5 ([Follmann 1992](#)). We performed sensitivity analyses using different imputed correlations.

Dealing with missing data

When statistics essential for analysis were missing (e.g. group means and standard deviations for both groups were not reported) and could not be calculated from other data, we attempted to contact study authors to obtain the missing data. We assumed that loss of participants that occurred before baseline measurements were taken had no effect on eventual outcome data provided by the study. We assessed any losses that occurred after baseline measurements had been taken and discussed them on an intention-to-treat basis.

Assessment of heterogeneity

We assessed statistical heterogeneity by using the I² statistic and by inspecting funnel plots.

Assessment of reporting biases

We assessed possible publication bias by using funnel plots.

Data synthesis

We performed meta-analyses by using Review Manager software version 5.3 (RevMan 2014).

Subgroup analysis and investigation of heterogeneity

We conducted the following a priori subgroup analyses on the presence/absence of:

1. short-burst oxygen therapy;
2. exertional desaturation (oxygen saturation (SaO₂) < 88% as entry criteria, or mean PaO₂ < 8 kPa on exertion);
3. baseline mean PaO₂ breathing air < 9.3 kPa;
4. measurement during exertion;
5. laboratory setting (compared with domiciliary setting);
6. measured acute oxygen effect (test on oxygen vs air) compared with long-term oxygen effect (test on the same inhaled gas after a period of oxygen or air); and
7. mean oxygen dose > 2 L/min. For studies that reported only administered fraction of inspired oxygen (FiO₂), an oxygen dose > 2 L/min was defined as FiO₂ > 0.27.

Sensitivity analysis

We conducted the following a priori sensitivity analyses by excluding studies with:

1. measurement at peak exertion (compared with iso-time);
2. high risk of bias for any bias category;
3. any participant without COPD; and
4. outlier findings (based on forest and funnel plots).

RESULTS

Description of studies

See [Characteristics of included studies](#) and [Characteristics of excluded studies](#) tables.

Participant characteristics

All 1195 participants included in the analysis had COPD (Table 1). Baseline PaO₂ was provided in 30 of 44 studies, ranging from 7.7 to 11.3 kPa. Twelve of the remaining studies provided baseline oxygen saturation ranging from 90% to 97% (Table 1).

Twenty studies reported mean baseline breathlessness at rest as follows: visual analogue score (VAS) score 5 to 40 mm (100 mm VAS); Borg score 0.1 to 1 point (Table 1).

Intervention characteristics

All studies compared oxygen versus air, delivered via the same non-invasive method in both groups. The most frequent mode of administration was the nasal cannula (22 studies) (Abernethy 2010; Davidson 1988; Dyer 2012; Eaton 2002; Eaton 2006; Haidl 2004; Jolly 2001a; Jolly 2001b; Knebel 2000; Kurihara 1989; Lewis 2003; McDonald 1995; McKeon 1988a; McKeon 1988b; Moore 2011; Nonoyama 2007; Ringbaek 2013; Rooyackers 1997a; Rooyackers 1997b; Spielmanns 2014; Wadell 2001; Woodcock 1981) followed by mouthpiece/valve (13 studies) (Bruni 2012a; Bruni 2012b; Dean 1992; Emtner 2003a; Emtner 2003b; Eves 2006; Laude 2006; Maltais 2001; Moore 2009; O'Donnell 1997; Scorsone 2010; Somfay 2001;

Swinburn 1984) and mask (eight studies) (Killen 2000; Leach 1992; Miki 2012; Nandi 2003; O'Driscoll 2011; Oliveira 2012a; Oliveira 2012b; Voduc 2010) or unknown (one study) (Ishimine 1995).

Doses of oxygen provided ranged from 2 to 6 L/min via nasal cannula, and FiO₂ ranged from 24% to 75% via mask/mouthpiece (Table 1).

A total of 32 studies (Bruni 2012a; Bruni 2012b; Davidson 1988; Dean 1992; Eaton 2006; Emtner 2003a; Emtner 2003b; Eves 2006; Haidl 2004; Ishimine 1995; Jolly 2001a; Jolly 2001b; Killen 2000; Knebel 2000; Kurihara 1989; Laude 2006; Leach 1992; Maltais 2001; McKeon 1988b; Miki 2012; O'Donnell 1997; Oliveira 2012a; Oliveira 2012b; Rooyackers 1997a; Rooyackers 1997b; Scorsone 2010; Somfay 2001; Spielmanns 2014; Swinburn 1984; Voduc 2010; Wadell 2001; Woodcock 1981) provided continuous oxygen during exercise testing - six-minute walk test (6MWT), endurance shuttle walk test, incremental shuttle walk test, step test or cycle exercise test. .

Six studies (Abernethy 2010; Eaton 2002; McDonald 1995; Moore 2011; Nonoyama 2007; Ringbaek 2013) provided domiciliary oxygen during daily life and activities.

Four studies (Killen 2000; Lewis 2003; McKeon 1988a; Nandi 2003) gave participants short-burst oxygen therapy before exertion.

Outcome characteristics

Breathlessness

Twenty-nine studies (Bruni 2012a; Bruni 2012b; Dean 1992; Eaton 2002; Emtner 2003a; Emtner 2003b; Eves 2006; Haidl 2004; Jolly 2001a; Jolly 2001b; Kurihara 1989; Laude 2006; Lewis 2003; Maltais 2001; McDonald 1995; Miki 2012; Moore 2009; Nonoyama 2007; O'Donnell 1997; O'Driscoll 2011; Oliveira 2012a; Oliveira 2012b; Ringbaek 2013; Rooyackers 1997a; Rooyackers 1997b; Scorsone 2010; Somfay 2001; Voduc 2010; Wadell 2001) measured breathlessness using a modified Borg scale, nine studies (Davidson 1988; Killen 2000; Knebel 2000; Leach 1992; McKeon 1988a; McKeon 1988b; Nandi 2003; Swinburn 1984; Woodcock 1981) used a VAS and five studies (Abernethy 2010; Dyer 2012; Eaton 2006; Ishimine 1995; Moore 2011) used other scales (Table 1).

HRQOL

Seven studies (Eaton 2002; Eaton 2006; Emtner 2003a; Emtner 2003b; Moore 2011; Rooyackers 1997a; Rooyackers 1997b) measured HRQOL as Chronic Respiratory Questionnaire (CRQ) total score, 10 studies (Dyer 2012; Eaton 2002; Eaton 2006; Emtner 2003a; Emtner 2003b; McDonald 1995; Moore 2011; Nonoyama 2007; Rooyackers 1997a; Rooyackers 1997b) as CRQ subdomains, five studies (Eaton 2002; Eaton 2006; Emtner 2003a; Emtner 2003b; Spielmanns 2014) as Short Form-36 (SF-36) total score and two studies (Nonoyama 2007; Ringbaek 2013) as SGRQ total score. One study (Abernethy 2010) measured perceived overall well-being on a 0 to 10 VAS and HRQOL over weeks or months (Table 1), with higher scores indicating better HRQOL.

Results of the search

Of an identified 866 unique records, we included 44 studies (1195 participants) in this update review (Figure 1). Compared with the previous review, which was published in 2011 (Uronis 2011), we included 14 additional studies (Abernethy 2010; Bruni 2012a;

Bruni 2012b; Dyer 2012; Miki 2012; Moore 2011; Nonoyama 2007; O'Driscoll 2011; Oliveira 2012a; Oliveira 2012b; Ringbaek 2013; Scorsone 2010; Spielmanns 2014; Voduc 2010), and we excluded one additional trial (Garrod 2000).

Figure 1. Study flow diagram.

Included studies

We have provided characteristics of the 44 included studies in the [Characteristics of included studies](#) table and in [Table 1](#). Of 44 included studies, 31 were cross-over trials ([Bruni 2012a](#); [Bruni 2012b](#); [Davidson 1988](#); [Dean 1992](#); [Eaton 2002](#); [Eves 2006](#); [Ishimine 1995](#); [Jolly 2001a](#); [Jolly 2001b](#); [Killen 2000](#); [Knebel 2000](#); [Kurihara 1989](#); [Laude 2006](#); [Leach 1992](#); [Lewis 2003](#); [Maltais 2001](#); [McDonald 1995](#); [McKeon 1988a](#); [McKeon 1988b](#); [Miki 2012](#); [Moore 2009](#); [Nandi 2003](#); [Nonoyama 2007](#); [O'Donnell 1997](#); [O'Driscoll 2011](#); [Oliveira 2012a](#); [Oliveira 2012b](#); [Somfay 2001](#); [Swinburn 1984](#); [Voduc 2010](#); [Woodcock 1981](#)) and 13 were parallel-group trials ([Abernethy 2010](#); [Dyer 2012](#); [Eaton 2006](#); [Emtner 2003a](#); [Emtner 2003b](#); [Haidl 2004](#); [Moore 2011](#); [Ringbaek 2013](#); [Rooyackers 1997a](#); [Rooyackers 1997b](#); [Scorsone 2010](#); [Spielmans 2014](#); [Wadell 2001](#)).

Ten studies ([Bruni 2012a](#); [Bruni 2012b](#); [Emtner 2003a](#); [Emtner 2003b](#); [Jolly 2001a](#); [Jolly 2001b](#); [Oliveira 2012a](#); [Oliveira 2012b](#); [Rooyackers 1997a](#); [Rooyackers 1997b](#)) included two different comparisons of independent groups, which we included in the analysis.

Excluded studies

We have presented excluded trials with reasons for exclusion in the [Characteristics of excluded studies](#) table. We identified two

ongoing or recently completed trials but did not include them in this review, as data were unavailable: the Long-Term Oxygen Therapy Trial (LOTT; registered at ClinicalTrials.gov: NCT00692198) of supplemental oxygen in 738 participants with COPD and moderate hypoxaemia at rest and/or exertion, which included measures of breathlessness and HRQOL; and another recently completed cross-over study comparing effects of supplemental oxygen versus air on breathlessness during exercise testing in 11 participants with COPD, which was presented as an abstract ([Vesteng 2015](#)).

Risk of bias in included studies

We have provided risk of bias judgements for each study at the end of each [Characteristics of included studies](#) table and have summarised each risk of bias category in [Figure 2](#). Methods were poorly reported in most of the included studies. We assessed risk of bias as mostly unclear regarding selection bias, but as low for more than half of studies regarding performance, detection, attrition and reporting bias, and as mostly unclear regarding other biases ([Figure 2](#)).

Figure 2. Methodological quality summary: review authors' judgements about each methodological quality item for each included study.

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias
Abernethy 2010	+	+	+	+	?	+	+
Bruni 2012a	?	?	+	+	+	+	?
Bruni 2012b	?	?	+	+	+	+	?
Davidson 1988	?	?	?	?	?	?	?
Dean 1992	?	?	+	+	+	+	?
Dyer 2012	+	-	?	-	?	?	-
Eaton 2002	?	?	+	+	+	+	?
Eaton 2006	+	+	+	+	?	+	+
Emtner 2003a	+	+	+	+	+	+	+
Emtner 2003b	+	+	+	+	+	+	+
Eves 2006	?	?	+	+	+	+	?
Haidl 2004	?	+	-	-	-	+	?
Ishimine 1995	?	?	?	-	?	?	?
Jolly 2001a	?	-	+	+	+	+	?
Jolly 2001b	?	-	+	+	+	+	?
Killen 2000	?	?	-	?	+	+	?
Knebel 2000	?	?	+	+	?	+	?
Kurihara 1989	?	?	-	?	?	?	?
Laude 2006	?	?	+	?	?	+	?
Leach 1992	?	?	-	-	+	+	?

Figure 2. (Continued)

Leach 1992	?	?	-	-	+	+	?
Lewis 2003	?	?	-	-	?	+	?
Maltais 2001	?	?	+	+	+	+	?
McDonald 1995	?	?	+	+	?	?	?
McKeon 1988a	?	?	+	+	+	+	?
McKeon 1988b	?	?	+	+	+	+	?
Miki 2012	?	?	?	?	?	+	?
Moore 2009	?	?	+	+	+	+	?
Moore 2011	+	+	+	+	+	+	?
Nandi 2003	?	?	+	+	+	+	?
Nonoyama 2007	+	+	+	+	?	+	?
O'Donnell 1997	?	?	+	+	+	+	?
O'Driscoll 2011	+	+	-	?	?	+	?
Oliveira 2012a	?	?	+	+	+	+	+
Oliveira 2012b	?	?	+	+	+	+	+
Ringbaek 2013	+	+	-	-	-	+	+
Rooyackers 1997	?	?	-	-	+	+	?
Rooyackers 1997a	?	?	-	-	+	+	?
Rooyackers 1997b	?	?	-	-	+	+	?
Scorsone 2010	?	?	+	+	+	+	?
Somfay 2001	?	?	-	-	+	+	?
Spielmanns 2014	+	+	?	?	-	+	+
Swinburn 1984	?	?	-	?	+	+	?
Voduc 2010	?	?	+	+	-	+	?
Wadell 2001	+	?	-	?	?	+	+
Woodcock 1981	?	?	+	+	+	+	+

Allocation

Although all studies were described as randomised, we could verify that sequence generation was adequate in only 11 studies (Abernethy 2010; Dyer 2012; Eaton 2006; Emtner 2003a; Emtner 2003b; Moore 2011; Nonoyama 2007; O'Driscoll 2011; Ringbaek 2013; Spielmanns 2014; Wadell 2001). Concealment of allocation was adequate in 10 studies and inadequate in three studies

(Figure 2). For remaining studies, we had insufficient information to determine risk of bias for allocation procedures.

Blinding

Masking of treatment was undertaken in several studies. For five studies, we were unable to determine how blinding of study participants or investigators had been achieved. For 12 studies (Haidl 2004; Killen 2000; Kurihara 1989; Leach 1992; Lewis 2003; O'Driscoll 2011; Ringbaek 2013; Rooyackers 1997a; Rooyackers

1997b; Somfay 2001; Swinburn 1984; Wadell 2001), we found that blinding was not undertaken, or investigators knew which containers contained oxygen. The remaining 27 studies attempted blinding of both study participants and study investigators (Figure 2).

Incomplete outcome data

Twenty-six studies (Bruni 2012a; Bruni 2012b; Dean 1992; Eaton 2002; Emtner 2003a; Emtner 2003b; Eves 2006; Jolly 2001a; Jolly 2001b; Killen 2000; Leach 1992; Maltais 2001; McKeon 1988a; McKeon 1988b; Moore 2009; Moore 2011; Nandi 2003; O'Donnell 1997; Oliveira 2012a; Oliveira 2012b; Rooyackers 1997a; Rooyackers 1997b; Scorsone 2010; Somfay 2001; Swinburn 1984;

Woodcock 1981) reported no withdrawals. Cross-over studies analyse within-participant differences; therefore, participants who did not complete both treatment periods were not included in the analyses. For the remaining studies, we could not reliably ascertain how missing data were handled (Figure 2).

Selective reporting

Possible publication bias on the effect of oxygen on breathlessness was indicated by the funnel plot shown in Figure 3. Studies showing a larger positive effect of oxygen on breathlessness were more likely to have lower precision. Asymmetry was apparent, with few studies with low precision reporting no or non-beneficial effects of oxygen on breathlessness (Figure 3).

Figure 3. Funnel plot of comparison: 1 Oxygen versus air, outcome: 1.1 Breathlessness - all trials.

Other potential sources of bias

We categorised the risk of other sources of bias as high for Dyer 2012 and as unclear for most studies.

Effects of interventions

See: [Summary of findings for the main comparison](#) Summary of findings table

All comparisons consisted of oxygen versus air delivered by the same mechanism among participants with COPD and no or mild hypoxaemia.

Primary outcome: breathlessness

The meta-analysis of breathlessness included 32 studies with 865 participants (Abernethy 2010; Bruni 2012a; Bruni 2012b; Davidson 1988; Dean 1992; Eaton 2002; Emtner 2003a; Emtner 2003b; Eves 2006; Jolly 2001a; Jolly 2001b; Killen 2000; Knebel 2000; Kurihara 1989; Laude 2006; Lewis 2003; McDonald 1995; McKeon 1988a; Miki 2012; Moore 2011; Nandi 2003; Nonoyama 2007; O'Donnell 1997; Oliveira 2012a; Oliveira 2012b; Ringbaek 2013; Rooyackers 1997a; Rooyackers 1997b; Scorsone 2010; Somfay 2001; Voduc 2010; Woodcock 1981).

In pooled analysis of all trials, we found that oxygen reduced breathlessness (SMD -0.31, 95% CI -0.43 to -0.20; $I^2 = 29\%$; Analysis 1.1; low-quality evidence).

A priori subgroup analyses of breathlessness

Short-burst oxygen therapy

Short-burst oxygen therapy before exertion did not reduce breathlessness (SMD -0.03, 95% CI -0.28 to 0.22; four studies; $I^2 = 0\%$), whereas oxygen given during exercise or daily activities did reduce breathlessness (SMD -0.36, 95% CI -0.48 to -0.24; 28 studies; $I^2 = 27\%$; [Analysis 1.2](#); low-quality evidence). Differences between groups were statistically significant ($P = 0.02$).

Exertional desaturation

In studies with desaturation during exercise, defined as $\text{SaO}_2 < 88\%$ among entry criteria or mean $\text{PaO}_2 < 8$ kPa on exertion,

oxygen reduced breathlessness (SMD -0.28, 95% CI -0.39 to -0.16; 16 studies), but the effect tended to be greater in studies without exertional desaturation (SMD -0.47, 95% CI -0.69 to -0.24; 15 studies; [Analysis 1.3](#)). $P = 0.14$ for differences between groups.

Baseline mean PaO_2 (air) < 9.3 kPa

The effect of oxygen was similar among studies with baseline mean PaO_2 (air) < 9.3 kPa (SMD -0.28, 95% CI -0.48 to -0.07; seven studies) and studies with PaO_2 (air) ≥ 9.3 kPa (SMD -0.33, 95% CI -0.47 to -0.20; 25 studies; [Analysis 1.4](#)), as shown in [Figure 4](#). $P = 0.65$ for differences between groups.

Figure 4. Forest plot of comparison: 1 Oxygen versus air, outcome: 1.2 Breathlessness - subgroup analysis - short-burst oxygen versus not.

Measurement during exercise test

Most studies measured breathlessness during exertion (SMD -0.34, 95% CI -0.46 to -0.22; 30 studies); only two studies measured

breathlessness in daily life and found a smaller and less precise effect of oxygen (SMD -0.13, 95% CI -0.37 to 0.11; two studies; Analysis 1.5; moderate-quality evidence), as shown in Figure 5. $P = 0.12$ for differences between groups.

Figure 5. Forest plot of comparison: 1 Oxygen versus air, outcome: 1.5 Breathlessness - subgroup analysis - measured during exercise test versus not.

Laboratory setting

Most studies measured breathlessness during exercise testing in a laboratory setting (SMD -0.37, 95% CI -0.52 to -0.22; 25 studies).

The effect of oxygen was less in non-laboratory domiciliary settings (SMD -0.23, 95% CI -0.36 to -0.09; seven studies; [Analysis 1.6](#)). $P = 0.16$ for differences between groups.

Short-term versus long-term (training) oxygen effects

Most studies evaluated acute effects of oxygen by comparing an exercise test on oxygen versus an exercise test on air (SMD -0.34, 95% CI -0.46 to -0.22; 29 studies). Investigators evaluated a potential longer-term (training) effect of oxygen by performing a test on the same gas after a (training) period with oxygen or air. Only three studies performed this evaluation and found no evidence of a long-term effect of oxygen (SMD -0.09, 95% CI -0.37 to 0.19; three studies; [Analysis 1.7](#)). $P = 0.11$ for differences between groups.

Mean oxygen dose > 2 L/min

Most studies evaluated oxygen at a dose > 2 L/min, which tended to reduce breathlessness more (SMD -0.35, 95% CI -0.49 to -0.21; 25 studies) than an oxygen dose ≤ 2 L/min (SMD -0.19, 95% CI -0.39 to 0.01; five studies; [Analysis 1.8](#)). $P = 0.19$ for differences between groups.

A priori sensitivity analyses of breathlessness

Analysis excluding trials measuring breathlessness at peak exertion

Oxygen reduced breathlessness measured at iso-time (SMD -0.37, 95% CI -0.50 to -0.24; 26 studies; $I^2 = 32\%$) but had no effect on breathlessness measured at peak exertion (SMD -0.14, 95% CI -0.33 to 0.05; six studies; $I^2 = 0\%$; [Analysis 1.9](#)).

Analysis excluding trials with high risk of bias

When we excluded trials with high risk of bias for any bias category (risk of bias shown in [Figure 2](#)), the effect of oxygen on breathlessness remained unchanged (SMD -0.30, 95% CI -0.41 to -0.20; 25 studies; $I^2 = 14\%$; [Analysis 1.10](#)).

Analysis excluding trials with any participant without COPD

All participants included in the meta-analysis had COPD. For studies that included non-COPD participants, we used individual participant data for the COPD subgroup ([Abernethy 2010](#)).

Analysis excluding outlier findings

Upon inspecting forest and funnel plots ([Figure 3](#); [Figure 4](#)), we identified five outlier findings ([Bruni 2012a](#); [Bruni 2012b](#); [Dean 1992](#); [Jolly 2001a](#); [Jolly 2001b](#); [Somfay 2001](#)). Exclusion of the outliers revealed that oxygen reduced breathlessness, but the effect became slightly smaller (SMD -0.27, 95% CI -0.35 to -0.18; 26 studies; $I^2 = 0\%$; [Analysis 1.11](#)). The effect of continuous oxygen during exertion or daily activities (no short-burst therapy) was similar after outliers were excluded (SMD -0.30, 95% CI -0.39 to -0.20; 22 studies; $I^2 = 0\%$; [Analysis 1.12](#)).

Secondary outcome: HRQOL

Meta-analysis

Twelve studies ([Abernethy 2010](#); [Dyer 2012](#); [Eaton 2002](#); [Eaton 2006](#); [Emtner 2003a](#); [Emtner 2003b](#); [McDonald 1995](#); [Moore 2011](#); [Nonoyama 2007](#); [Ringbaek 2013](#); [Rooyackers 1997](#); [Spielmanns 2014](#)) examined changes in HRQOL. Of these, we were able to include five studies (267 participants) ([Eaton 2002](#); [Moore 2011](#); [Nonoyama 2007](#); [Rooyackers 1997](#); [Spielmanns 2014](#)) in the meta-analysis. Oxygen had no clear effect on HRQOL (SMD 0.12, 95% CI -0.04 to 0.28; five studies; $I^2 = 0\%$; [Analysis 1.13](#); low-quality evidence), as shown in [Figure 6](#). In the analysis, higher scores indicated better HRQOL.

Figure 6. Forest plot of comparison: 1 Oxygen versus air, outcome: 1.13 Health-related quality of life - all trials.

Descriptive analysis of studies not included in meta-analysis

We did not include seven studies in the meta-analysis, as they did not use a validated scale of HRQOL ([Abernethy 2010](#)), reported only HRQOL subdomain scores ([Dyer 2012](#); [McDonald 1995](#)), reported change scores ([Emtner 2003a](#); [Emtner 2003b](#); [Ringbaek 2013](#)) or provided data that could not be combined in the analysis ([Eaton 2006](#)). For all scales used, higher scores indicated better quality of life.

[Abernethy 2010](#) measured overall perceived quality of life over the past two days on a 0 to 10 numerical rating scale (NRS) based on a daily diary kept over seven days. In the subgroup with COPD ($n = 158$), improvement from baseline at day 7 was 0.8 points (standard deviation (SD) 2.0) in the oxygen group and 0.3 points (SD 1.9) in the air group.

[Dyer 2012](#) measured change from baseline in CRQ subdomain scores after six to seven weeks of pulmonary rehabilitation using

oxygen or air. The only statistically and clinically significant change was found in the CRQ mastery score. For oxygen compared with air, the mean difference was -0.1 (95% CI -0.9 to 0.7; $P = 0.76$) for the domain CRQ breathlessness, 0.5 (95% CI -0.2 to 1.2; $P = 0.15$) for CRQ emotion, 0.7 (95% CI -0.1 to 1.5; $P = 0.10$) for CRQ fatigue and 0.7 (95% CI 0.0 to 1.4; $P = 0.006$) for CRQ mastery scores.

Eaton 2006 was a randomised, controlled, parallel-group trial that included three arms as follows: oxygen, air and usual care. The only domain of the CRQ to show statistical significance ($P = 0.045$) was emotional function; the greatest improvement was noted in the usual care group, which received neither oxygen nor air.

For Emtner 2003a and Emtner 2003b, CRQ total score and subscores increased significantly in both groups. Only in mastery was a statistically significantly greater improvement detected in the oxygen-trained group ($P < 0.05$). For the Short Form-36, a significant difference (improvement) in general health was seen in the oxygen compared with the air group ($P < 0.05$).

McDonald 1995 measured subdomains of CRQ and noted statistically significant improvement in all CRQ subdomains for the comparison of baseline scores versus scores after six weeks of oxygen therapy ($P < 0.02$ for all domains). However, when scores after oxygen therapy were compared with scores after air, they reported no statistically or clinically significant differences.

Ringbaek 2013 measured total SGRQ score at 7 weeks compared with baseline. Mean changes from baseline were -1.8 (SD 8) for the oxygen group and -3.2 (SD 7.2) for no oxygen group; the difference between groups was not statistically significant ($P = 0.80$).

Secondary outcome: blinded patient preference

Five studies (Abernethy 2010; Eaton 2002; Killen 2000; McDonald 1995; Moore 2011) assessed patient preference to continue therapy (oxygen vs air delivered by the same mechanism) at a time when participants were still blinded. Owing to heterogeneity of the reported data, we could not perform a meta-analysis.

Abernethy 2010 assessed patient preference after seven days of blinded oxygen/air by asking the question, "Reviewing the benefits and burdens of your experience with the treatment over the past week, would you want to continue this treatment?" Among participants with COPD who answered ($N = 139$), preferences to continue treatment were as follows in the oxygen group: yes 50%, maybe 27% and no 23%. In the air group, preferences were yes 41%, maybe 23% and no 36%. The difference in preference between groups was not statistically significant ($P = 0.25$).

Eaton 2002 asked participants ($N = 41$) if they were interested in clinical provision of oxygen at study completion. Among participants identified as having a short-term response to oxygen, 14 participants did not wish to be considered for continued therapy. Eleven of these 14 (76%) cited poor tolerability or acceptability as the reason.

Killen 2000 studied short-burst oxygen given immediately before and after walking up a flight of steps, asking participants which blinded gas they preferred. Of 18 participants, five preferred oxygen before ascending the stairs, three preferred air and three had no preference. The remaining seven participants preferred to receive oxygen at the top of the stairs. As a group, participants expressed no statistically significant preference for oxygen therapy ($P = 0.12$).

McDonald 1995 included both acute assessments and a domiciliary portion that lasted six weeks with each gas, asking participants ($N = 26$) at the end of the study which six-week period they preferred. Fifty percent preferred the period when they received oxygen; the remaining 50% preferred air or had no preference.

Moore 2011 asked participants whether they wanted to continue or discontinue the provision of domiciliary cylinders after the 12-week study period. Preferences were similar between groups; 35 of 65 (54%) wanted to continue in the oxygen group compared with 37 of 73 (51%) in the air group.

Secondary outcome: adverse events

Adverse events were insufficiently and inconsistently reported; therefore, meta-analysis was not possible.

Abernethy 2010 included a total of 65 participants in the oxygen group and 70 in the air group. Serious adverse events were rare, with no clinically meaningful differences between groups for moderate to extreme drowsiness (47% oxygen vs 51% air); moderate to extreme nasal irritation (29% oxygen vs 35% air); moderately to extremely troublesome nosebleeds (3% oxygen vs 3% air) and moderate to extreme anxiousness (26% oxygen vs 40% air).

Dyer 2012 included a total of 47 participants in the study. Investigators reported seven dropouts; three participants withdrew because of an exacerbation of their condition, one because of other medical problems and three for social reasons.

Eaton 2002 included a total of 41 participants in the study. Investigators reported nine dropouts; causes of dropout included morbidity for two participants and cancer for one participant in both study groups. Three in each group withdrew for personal reasons.

Eaton 2006 included a total of 25 participants in the oxygen group and 26 in the air group. Five participants in the oxygen group died, one was admitted to the hospital and three were prescribed LTOT. In the air group, two participants died, one was prescribed LTOT, one was admitted to a rest home and four were admitted to the hospital.

Emtner 2003a and Emtner 2003b included a total of 14 participants in the oxygen group and 15 in the air group. We excluded one participant from the oxygen-trained group because of illness post intervention.

Haidl 2004 included a total of 26 participants in the study. During three-year follow-up, four participants in the oxygen group died and one was given a diagnosis of cancer.

Knebel 2000 included a total of 31 participants in the study. Two participants were unable to complete all of the walks in the study because of unrelated problems (fever and migraine headache).

Laude 2006 included a total of 76 participants in the study. After randomisation, seven participants withdrew: five because of exacerbations and two for other reasons.

Lewis 2003 included a total of 18 participants in the study and reported four dropouts. Two participants failed to complete the first visit and attend the second visit (chest pain during 6MWT ($n = 1$), personal reasons ($n = 1$)). Furthermore, two participants failed to attend the second visit (clinically unstable ($n = 1$), other ($n = 1$)).

[McDonald 1995](#) included a total of 26 participants in the study and reported seven dropouts; three participants withdrew during the first six weeks of domiciliary therapy (acute gout, muscular pain related to pulling the gas cylinder, unwillingness to continue in the study, respectively), each of whom had received cylinder air in the first six weeks. A total of four participants were withdrawn during the second six weeks: One participant died of a cerebrovascular accident and another of overwhelming pneumonia, a third was hospitalised with an acute exacerbation of COPD and the last incurred an incidental injury. All four participants had received cylinder oxygen during this second six-week trial period.

[Moore 2011](#) included a total of 66 participants in the oxygen group and 73 in the air group. After randomisation, among those allocated to cylinder oxygen, one participant was deceased and one got unwell.

[Nonoyama 2007](#) included a total of 27 participants in the study and reported 11 dropouts. Five were reluctant to continue, three developed resting hypoxaemia with a PaO_2 less than 55 mmHg, two died, and one was non-compliant, utilising the test mixtures for less than one hour per day.

[Ringbaek 2013](#) included a total of 15 participants in the oxygen group and 22 in the air group. At study end (33 weeks), the mean number of adverse events did not differ significantly between treatment groups in terms of acute COPD exacerbations ($P = 0.30$) or number of participants with hospital admission or dropout ($P = 0.59$).

[Spielmans 2014](#) included a total of 19 participants in the oxygen group and 17 in the air group. After randomisation, in the oxygen group, five participants discontinued owing to comorbidities, and in the air group, seven discontinued because of comorbidities.

[Voduc 2010](#) included a total of 15 participants in the study. After randomisation, three participants developed COPD exacerbations and one developed worsening of arthritis that limited exercise and thus was excluded.

[Bruni 2012a](#); [Bruni 2012b](#); [Davidson 1988](#); [Dean 1992](#); [Eves 2006](#); [Ishimine 1995](#); [Jolly 2001a](#); [Jolly 2001b](#); [Killen 2000](#); [Kurihara 1989](#); [Leach 1992](#); [Maltais 2001](#); [McKeon 1988a](#); [McKeon 1988b](#); [Miki 2012](#); [Moore 2009](#); [Nandi 2003](#); [O'Donnell 1997](#); [O'Driscoll 2011](#); [Oliveira 2012a](#); [Oliveira 2012b](#); [Rooyackers 1997a](#); [Rooyackers 1997b](#); [Scorsone 2010](#); [Somfay 2001](#); [Swinburn 1984](#); [Wadell 2001](#); and [Woodcock 1981](#) did not report adverse events.

DISCUSSION

Summary of main results

Since the previous version of this review was published in 2011 ([Uronis 2011](#)), we have included an additional 14 studies of the effects of oxygen versus air on breathlessness, we have excluded one study, and data for meta-analysis of health-related quality of life (HRQOL) are now available.

When compared with air, continuous oxygen (not short-burst oxygen) reduced breathlessness more during exercise or activity (standard mean difference (SMD) -0.36, 95% confidence interval (CI) -0.48 to -0.24), and most data pertaining to breathlessness were obtained during exercise testing in the laboratory setting. This effect translated to a mean decrease in breathlessness of 0.7

points on a 0 to 10 numerical rating scale (NRS) based on data from a recent large study ([Abernethy 2010](#)). The minimal clinically important difference in chronic breathlessness has been reported to be 1 point, with small, moderate and large effects seen at about 0.6, 1.1 and 1.8 points, respectively ([Johnson 2013](#)). It is unclear whether the effect of oxygen on breathlessness could be clinically meaningful for many participants in this setting.

Evidence for breathlessness pertains mostly to acute effects of exercise testing on breathlessness in the laboratory setting. Effects on breathlessness during daily life (not measured during an exercise test) in the domiciliary setting were smaller and were statistically non-significant (SMD -0.13, 95% CI -0.37 to 0.11).

Short-burst oxygen therapy before exertion did not affect breathlessness.

Oxygen did not clearly affect HRQOL (SMD 0.10, 95% CI -0.06 to 0.26). Blinded patient preference to continue therapy did not differ between groups given oxygen and groups given air. Investigators insufficiently reported adverse events, and we could not meta-analyse adverse event data.

Overall completeness and applicability of evidence

After an extensive database and literature search, we included 44 studies (1195 participants) in this update review. Although two relatively large studies of ambulatory oxygen have been published in recent years ([Abernethy 2010](#); [Moore 2011](#)), additional high-quality data on effects of ambulatory oxygen in daily life, including effects on HRQOL, are needed. Good methodological rigour would include publication of a prespecified protocol outlining trial design, adequate sequence generation, randomisation, blinding (for participants and for outcome assessors) and transparent reporting of appropriate outcomes at baseline in a format suitable for meta-analysis (e.g. mean with standard deviation and sample size).

Mechanism of effect on breathlessness

The likely mechanism underpinning an effect on breathlessness is that supplemental oxygen prevents or decreases hypoxaemia during exercise, thereby reducing a hypoxaemia-related increase in ventilatory demand, dynamic hyperinflation and increased exertional breathlessness in some patients ([O'Donnell 2001](#)). This mechanism is supported by the present finding that a higher oxygen dose (> 2 L/min) was associated with a larger decrease in breathlessness. The finding that the oxygen effect was not stronger in studies of patients with exertional desaturation might reflect that the analysis was based on the study mean, and that heterogeneity in the level of exertional hypoxaemia among individual participants might have attenuated the association. Some people developed hypoxaemia during exercise despite supplemental oxygen therapy that could have increased their breathlessness scores. The issue is further complicated in that some patients might have adapted to exertional hypoxaemia, with a reduced ventilatory response to hypoxaemia and therefore less benefit of oxygen during exercise. Another hypothesis is that airflow to the face and upper airways could relieve breathlessness, possibly through increased afferent feedback mimicking increased ventilatory work ([Johnson 2016](#); [Schwartzstein 1987](#)). This could reduce the imbalance between ventilatory demand and perceived ventilatory work and thus the level of breathlessness, and may explain why participants did not prefer to go on, or why no significant change in HRQOL was evident

when oxygen was compared with air delivered through the same method.

Evidence for use in clinical care

Most evidence of the benefit of oxygen for breathlessness pertains to symptoms noted during standardised exercise testing in the laboratory setting. The oxygen effect during domiciliary treatment in daily life was smaller and more uncertain (SMD -0.12, 95% CI -0.39 to 0.15). More recent double-blinded randomised controlled trials (RCTs) of oxygen compared with medical air have found no statistically or clinically relevant effects of domiciliary/ambulatory oxygen on breathlessness in daily life ([Abernethy 2010](#); [Moore 2011](#)). Less efficacy of domiciliary oxygen might reflect insufficient adherence to therapy or low physical activity due to deconditioning, or the fact that breathlessness in daily life is not assessed at a comparable time point or after a standardised workload. A patient who benefits from oxygen (decreased breathlessness at a given workload and improved exercise capacity) might increase his/her activity up to maximal tolerable levels of breathlessness. We found that the level of breathlessness at peak exercise did not seem to be affected by oxygen. When breathlessness is recalled in daily life, reported symptom level might therefore be similar between people receiving oxygen and those given air, even if oxygen would in fact be beneficial in some patients. Whether palliative oxygen in the domiciliary setting is beneficial remains unclear. We found no evidence to show that oxygen improved HRQOL, but the data were limited.

Quality of the evidence

We rated the quality of evidence for the effect of oxygen compared with air on breathlessness as low overall upon completing a GRADE assessment ([Summary of findings for the main comparison](#)). Most evidence pertained to breathlessness measured during a standardised exercise test (moderate-quality evidence), whereas evidence was limited for breathlessness during daily life (low-quality evidence). We rated the quality of evidence showing an effect of oxygen on HRQOL as low ([Summary of findings for the main comparison](#)).

Potential biases in the review process

Limitations of this systematic review and meta-analysis mainly reflect the heterogeneity and methodological limitations of the currently available body of literature. Although we excluded studies of participants already qualifying for home oxygen therapy according to current guidelines, the review population still included a wide range of baseline oxygen saturation/partial pressure of oxygen (PaO₂). This variability could affect our results if a relationship is found between oxygen saturation and effects of oxygen on breathlessness. We addressed this issue by performing a subgroup analysis based on baseline PaO₂ (≥ 9.3 kPa) and obtained consistent findings. Additionally, baseline breathlessness and physical capacity were often insufficiently reported. The review population likely included participants with varied perceptions of breathlessness and physical capacity.

We explored potential bias due to methodological limitations by excluding studies with high risk for any bias category; this yielded consistent findings. A possible limitation of this review is that many studies did not report data on participants dropping out or

withdrawing from the study. Reporting of data only for participants who completed the trial might have introduced selection bias.

Palliative oxygen is prescribed most commonly for seriously ill patients nearing the end of life; however, these patients are not likely to participate in randomised trials, especially studies involving an exercise test. The applicability of review findings to all individuals with chronic obstructive pulmonary disease (COPD) who require palliative oxygen is unclear.

Effects of oxygen on breathlessness were driven in part by five outlier findings among studies with small sample sizes/low levels of precision. This asymmetry in the funnel plot might indicate the presence of publication bias. When we excluded the outliers, the oxygen effect became somewhat smaller (SMD -0.24, 95% CI -0.34 to -0.15).

Although the funnel plot indicated potential publication bias, we did not downgrade the quality of the evidence, as a sensitivity analysis excluding outlying studies yielded similar findings.

Agreements and disagreements with other studies or reviews

Compared with the 2011 version of this review ([Uronis 2011](#); [Uronis 2015](#)), we have included an additional 14 studies (493 participants) in this update. The finding of the main analysis of breathlessness is consistent with the previous estimate but is more precise because we included more studies. Subgroup analyses by short burst and by oxygen dose were also in agreement with those of the previous review. Novel subgroup analyses in the present review show that evidence of effects of oxygen pertains mostly to breathlessness during exercise tests performed in the laboratory standardised rehabilitation training setting. Evidence is less consistent and of low quality for effects of supplemental oxygen during daily life. The novel analysis of HRQOL showed no clear effect of supplemental oxygen compared with air on quality of life.

[Ameer 2014](#) performed a Cochrane meta-analysis (four studies; 331 participants) of the effects of ambulatory oxygen therapy given for two weeks or longer on breathlessness and HRQOL in the home setting. When compared with air or no therapy, ambulatory oxygen therapy was associated with a small reduction in breathlessness as measured on the Borg scale (mean difference 0.28, 95% CI, 0.10 to 0.45). This finding is consistent with our finding in the non-laboratory setting. The analysis of breathlessness performed by [Ameer 2014](#) included only three studies ([McDonald 1995](#); [Eaton 2002](#); [Nonoyama 2007](#)), which we also included in the present review. Effects on HRQOL as noted in [Ameer 2014](#) were inconsistent, with mean improvements in breathlessness and fatigue domains but were not related to emotional function or mastery. Small or inconsistent effects of supplemental oxygen on HRQOL and lack of clear effect on exercise capacity in [Ameer 2014](#) support the present conclusion that the evidence base for supplemental oxygen therapy during daily life is limited.

Findings of this review are consistent with recent British Thoracic Society (BTS) guidelines stating that although ambulatory oxygen therapy should be offered to patients for use during exercise in a pulmonary rehabilitation programme, it "should not be routinely offered to patients who are not eligible for LTOT" ([Harding 2015](#)).

AUTHORS' CONCLUSIONS

Implications for practice

Oxygen given continuously during exercise or activity can relieve breathlessness in patients with COPD who have no or mild hypoxaemia and would not qualify for long-term oxygen therapy. Evidence was of moderate quality for breathlessness during exercise testing in the laboratory setting and of low quality for effects on breathlessness during daily life. It is not clear whether the reduction in breathlessness shown in the laboratory setting translates into a clinically important benefit, and no evidence supports a clinically important benefit from oxygen for breathlessness in daily life. Short-burst oxygen therapy given before exercise had no effect and should not be used.

Clinical recommendations

Based on the present findings, supplemental oxygen therapy is not a first-line treatment for patients with breathlessness. Management of underlying disease(s) should be optimised. Evidence-based symptomatic treatment should include individualised rehabilitation training (McCarthy 2015), the fan and low-dose opioids (Ekstrom 2015a; Ekstrom 2015b; Hardinge 2015). In patients with intractable breathlessness despite these interventions, a trial of supplemental oxygen could be considered. Guidelines recommend that palliative oxygen should be evaluated only in patients with hypoxaemia (saturation < 92%) at rest (Hardinge 2015). We found that the oxygen effect was similar in studies with mean $\text{PaO}_2 \geq 9.3$ kPa compared with $\text{PaO}_2 < 9.3$ kPa, and that a trial of supplemental oxygen might be reasonable in patients without resting hypoxaemia, especially in the presence of hypoxaemia during limited exertion. Evaluation of the potential benefit of ambulatory oxygen should involve the patient's levels of physical capacity and activity, along with the benefits of oxygen for breathlessness and physical capacity in a standardised test such as the six-minute walking test (6MWT) when compared with air (Hardinge 2015). Patients who perceive decreased breathlessness when receiving oxygen most often experience this effect within two to three days of the start of treatment (Abernethy 2010). Therefore, palliative oxygen therapy should be evaluated after a few days and withdrawn if patients do not perceive benefit.

Patient quality of life (QOL) factors such as convenience and adverse consequences should be considered in the decision whether to prescribe oxygen as treatment for patients with breathlessness who may already be burdened by their illness and other life changes prominent in the advanced illness setting.

Finally, in considering these results, one must be sure to remember the downsides of administering oxygen. Oxygen is costly, and, with current stresses on healthcare systems in many countries, this needs to be taken into account. Evidence of the effect of supplemental oxygen on HRQOL is lacking.

In conclusion, we are moderately confident that oxygen can relieve breathlessness when given during exercise to mildly hypoxaemic

and non-hypoxaemic people with COPD who would not otherwise qualify for home oxygen therapy. Most evidence pertains to acute effects during exercise testing; evidence of long-term effects of oxygen during daily life is less consistent. Findings show that oxygen did not affect HRQOL.

Implications for research

Current findings on the effect of oxygen therapy versus air have several important implications for research. First, oxygen affected breathlessness measured only at iso-time, not at peak exercise. This confirms the importance of measuring breathlessness at iso-time with comparable workloads in mechanistic studies and interventional trials.

Although this review confirms that oxygen can decrease exertional breathlessness in the laboratory setting, studies should focus on effects of domiciliary and ambulatory oxygen in clinical and daily life settings. Studies on effects of oxygen on HRQOL and on different aspects (dimensions) of breathlessness (Laviolette 2014) are needed, and investigators should identify optimal doses and routes of oxygen administration, while determining which patients are most likely to receive beneficial symptomatic effects of oxygen. Studies of symptomatic benefit should also account for changes in physical capacity and activity to explore whether oxygen given during daily living could allow increased exercise activity, although it might not affect reported breathlessness scores. Tools for measuring physical activity in COPD were recently validated (Gimeno-Santos 2015).

We found that adverse events were insufficiently assessed and/or reported. The first population-based longitudinal study of the risk of burn injury during LTOT was published only recently (Tanash 2015). Studies quantifying adverse events, patient burden and potential risks associated with supplemental oxygen therapy are needed to determine the net clinical benefit of palliative oxygen.

ACKNOWLEDGEMENTS

We would like to thank Dr Hope Uronis, Duke Medical Center, Durham, North Carolina, USA, for her work with the review published in 2011, as well as Dr Emma J Dennett and Mrs Liz Stovold, Cochrane Airways Group, for their assistance with subsequent MEDLINE/Embase/CENTRAL searches and the review process.

Professor Phillippa Poole was the Editor for this review and commented critically on the review.

The Background and Methods sections of this review are based on a standard template used by Cochrane Airways.

This project was supported by the National Institute for Health Research (NIHR) via Cochrane Infrastructure funding to the Cochrane Airways Group. The views and opinions expressed therein are those of the review authors and do not necessarily reflect those of the Systematic Reviews Programme, NIHR, NHS or the Department of Health.

REFERENCES

References to studies included in this review

Abernethy 2010 {published and unpublished data}

Abernethy AP, McDonald CF, Frith PA, Clark K, Herndon JE 2nd, Marcello J, et al. Effect of palliative oxygen versus room air in relief of breathlessness in patients with refractory dyspnoea: a double-blind, randomised controlled trial. *Lancet* 2010;**376**(9743):784-93.

Bruni 2012a {published data only}

Bruni GI, Gigliotti F, Binazzi B, Romagnoli I, Duranti R, Scano G. Dyspnea, chest wall hyperinflation, and rib cage distortion in exercising patients with chronic obstructive pulmonary disease. *Medicine and Science in Sports and Exercise* 2012;**44**(6):1049-56.

Bruni 2012b {published data only}

Bruni GI, Gigliotti F, Binazzi B, Romagnoli I, Duranti R, Scano G. Dyspnea, chest wall hyperinflation, and rib cage distortion in exercising patients with chronic obstructive pulmonary disease. *Medicine and Science in Sports and Exercise* 2012;**44**(6):1049-56.

Davidson 1988 {published data only}

Davidson AC, Leach R, George RJD, Geddes DM. Supplemental oxygen and exercise ability in chronic obstructive pulmonary disease. *Thorax* 1988;**43**:965-71.

Dean 1992 {published data only}

Dean NC, Brown JK, Himelman RB, Doherty JJ, Gold WM, Stulbarg MS. Oxygen may improve dyspnea and endurance in patients with chronic obstructive pulmonary disease and only mild hypoxemia. *American Review of Respiratory Disease* 1992;**146**:941-5.

Dyer 2012 {published data only}

Dyer F, Bott J. Ambulatory oxygen improves the effectiveness of pulmonary rehabilitation (PR) in selected patients [Abstract]. American Thoracic Society International Conference; 2010 May 14-19; New Orleans. 2010:A3763.

* Dyer F, Callaghan J, Cheema K, Bott J. Ambulatory oxygen improves the effectiveness of pulmonary rehabilitation in selected patients with chronic obstructive pulmonary disease. *Chronic Respiratory Disease* 2012;**9**(2):83-91.

Dyer FJ. Ambulatory oxygen improves the effectiveness of pulmonary rehabilitation (PR) in selected patients [Abstract]. *Thorax* 2010;**65**:S71.

Eaton 2002 {published data only}

Eaton T, Garret JE, Young P, Fergusson W, Kolbe J, Rudkin S, et al. Ambulatory oxygen improves quality of life of COPD patients: a randomised controlled study. *European Respiratory Journal* 2002;**20**(2):306-12.

Eaton 2006 {published data only}

Eaton T, Fergusson W, Kolbe J, Lewis CA, West T. Short-burst oxygen therapy for COPD patients: a 6-month randomised, controlled trial. *European Respiratory Journal* 2006;**27**(4):697-704.

Emtner 2003a {published data only}

* Emtner M, Porszasz J, Burns M, Somfay A, Casaburi R. Benefits of supplemental oxygen in exercise training in non-hypoxemic chronic obstructive pulmonary disease patients. *American Journal of Respiratory and Critical Care Medicine* 2003;**168**(9):1034-42.

Emtner M, Porszasz J, Burns M, Somfay A, Casaburi R. Online supplement to 'Benefits of supplemental oxygen in exercise training in non-hypoxemic chronic obstructive pulmonary disease patients'. <http://ajrccm.atsjournals.org/cgi/data/168/9/1034/DC1/1> (accessed 21 August 2008).

Emtner 2003b {published data only}

* Emtner M, Porszasz J, Burns M, Somfay A, Casaburi R. Benefits of supplemental oxygen in exercise training in non-hypoxemic chronic obstructive pulmonary disease patients. *American Journal of Respiratory and Critical Care Medicine* 2003;**168**(9):1034-42.

Emtner M, Porszasz J, Burns M, Somfay A, Casaburi R. Online supplement to 'Benefits of supplemental oxygen in exercise training in non-hypoxemic chronic obstructive pulmonary disease patients'. <http://ajrccm.atsjournals.org/cgi/data/168/9/1034/DC1/1> (accessed 21 August 2008).

Eves 2006 {published data only}

Eves ND, Petersen SR, Haykowsky MJ, Wong EY, Jones RL. Helium-hyperoxia, exercise, and respiratory mechanics in chronic obstructive pulmonary disease. *American Journal of Respiratory and Critical Care Medicine* 2006;**174**(7):763-71.

Haidl 2004 {published data only}

Haidl P, Clement C, Wiese C, Dellweg D, Kohler D. Long-term oxygen therapy stops the natural decline of endurance in COPD patients with reversible hypercapnia. *Respiration* 2004;**71**:342-7.

Ishimine 1995 {published data only}

Ishimine A, Saito T, Nishimura S, Nakano G, Miyamoto K, Kawakami Y. The effect of oxygen supplementation during exercise in COPD patients with PaO₂ over 60 Torr. *Japanese Journal of Chest Disease* 1995;**33**:510-8.

Jolly 2001a {published data only}

Jolly EC, Di Boscio V, Aguirre L, Luna CM, Berensztejn S, Gene RJ. Effects of supplemental oxygen during activity in patients with advanced COPD without severe resting hypoxemia. *Chest* 2001;**120**:437-43.

Jolly 2001b {published data only}

Jolly EC, Di Boscio V, Aguirre L, Luna CM, Berensztejn S, Gene RJ. Effects of supplemental oxygen during activity in patients with advanced COPD without severe resting hypoxemia. *Chest* 2001;**120**:437-43.

Killen 2000 {published data only}

Killen JWW, Corris PA. A pragmatic assessment of the placement of oxygen when given for exercise induced dyspnea. *Thorax* 2000;**55**:544-6.

Knebel 2000 {published data only}

Knebel AR, Bentz E, Barnes P. Dyspnea management in alpha-1 antitrypsin deficiency: effect of oxygen administration. *Nursing Research* 2000;**49**(6):333-8.

Kurihara 1989 {published data only}

Kurihara N, Fujimoto S, Kuono M, Futoda K, Hirata K, Takeda C. Exercise induced hypoxemia and exercise tolerance in patients with COPD and the benefits of oxygen supplementation. *Japanese Journal of Chest Disease* 1989;**26**:155-62.

Laude 2006 {published data only}

Laude EA, Duffy NC, Baveystock C, Dougill B, Campbell MJ, Lawson R, et al. The effect of helium and oxygen on exercise performance in chronic obstructive pulmonary disease. *American Journal of Respiratory and Critical Care Medicine* 2006;**173**(8):865-70.

Leach 1992 {published data only}

Leach RM, Davidson AC, Chinn S, Twort CHC, Cameron IR, Bateman NT. Portable liquid oxygen and exercise ability in severe respiratory disability. *Thorax* 1992;**47**:781-9.

Lewis 2003 {published data only}

Lewis CA, Eaton TE, Young P, Kolbe J. Short-burst oxygen immediately before and after exercise is ineffective in non-hypoxic COPD patients. *European Respiratory Journal* 2003;**22**(4):584-8.

Maltais 2001 {published data only}

Maltais F, Simon M, Jobin J, Desmeules M, Sullivan M, Belanger M, et al. Effects of oxygen on lower limb blood flow and O₂ uptake during exercise in COPD. *Medicine and Science in Sports and Exercise* 2001;**33**(6):916-22.

McDonald 1995 {published data only}

McDonald CF, Blyth DM, Lazarus MD, Marschner I, Barter CE. Exertional oxygen of limited benefit in patients with chronic obstructive pulmonary disease and mild hypoxemia. *American Journal of Respiratory and Critical Care Medicine* 1995;**152**:1616-9.

McKeon 1988a {published data only}

McKeon JL, Murree-Allen K, Saunders NA. Effects of breathing supplemental oxygen before progressive exercise in patients with chronic obstructive lung disease. *Thorax* 1988;**43**(1):53-6.

McKeon 1988b {published data only}

McKeon JL, Tomlinson JC, Tarrant EP, Mitchell CA. Portable oxygen in patients with severe chronic obstructive pulmonary disease. *Australia and New Zealand Journal of Medicine* 1988;**18**(2):125-9.

Miki 2012 {published data only}

Miki K, Maekura R, Hiraga T, Kitada S, Miki M, Yoshimura K, et al. Effects of oxygen on exertional dyspnoea and exercise performance in patients with chronic obstructive pulmonary disease. *Respirology (Carlton, Vic)* 2012;**17**(1):149-54.

Moore 2009 {published data only}

Moore R, Berlowitz D, Pretto J, Brazzale D, Denehy L, Jackson B, et al. Acute effects of hyperoxia on resting pattern of ventilation and dyspnoea in COPD. *Respirology* 2009;**14**(4):545-50.

Moore 2011 {published data only}

* Moore RP, Berlowitz DJ, Denehy L, Pretto JJ, Brazzale DJ, Sharpe K, et al. A randomised trial of domiciliary, ambulatory oxygen in patients with COPD and dyspnoea but without resting hypoxaemia. *Thorax* 2011;**66**(1):32-7.

Moore RP, Berlowitz R, Denehy D, Pretto L, Brazzale J, Sharpe D, et al. Domiciliary ambulatory oxygen is not beneficial in patients with COPD and mild resting hypoxaemia [Abstract]. *American Journal of Respiratory and Critical Care Medicine* 2010;**181**:P4380.

Nandi 2003 {published data only}

* Nandi K, Smith AA, Crawford A, MacRae KD, Garrod R, Seed WA, et al. Oxygen supplementation before or after submaximal exercise in patients with chronic obstructive pulmonary disease. *Thorax* 2003;**58**:670-3.

Smith AA, Nandi K, Crawford A, Garrod R, MacRae KD, Roberts CM, et al. Exercise intolerance, hypoxaemia and breathlessness in COPD: does pre-dosing with oxygen actually help?. *American Journal of Respiratory and Critical Care Medicine* 2001;**163**(Suppl 5):A501.

Nonoyama 2007 {published data only}

Nonoyama ML. Effect of oxygen in COPD patients with transient exertional hypoxemia [Dissertation]. Dissertation Abstracts International 2010; Vol. 70, issue 10-B:6153.

Nonoyama ML, Brooks D, Guyatt GH, Goldstein RS. Ambulatory gas usage in patients with chronic obstructive pulmonary disease and exertional hypoxemia [Abstract]. *Journal of Cardiopulmonary Rehabilitation and Prevention*. 2008; Vol. 28, issue 5:323-9.

* Nonoyama ML, Brooks D, Guyatt GH, Goldstein RS. Effect of oxygen on health quality of life in patients with chronic obstructive pulmonary disease with transient exertional hypoxemia. *American Journal of Respiratory and Critical Care Medicine* 2007;**176**(4):343-9.

O'Donnell 1997 {published data only}

O'Donnell DE, Bain DJ, Webb KA. Factors contributing to relief of exertional breathlessness during hyperoxia in chronic airflow limitation. *American Journal of Respiratory and Critical Care Medicine* 1997;**155**(2):530-5.

O'Driscoll 2011 {published data only}

O'Driscoll BR, Neill J, Pulakal S, Turkington PM. A crossover study of short burst oxygen therapy (SBOT) for the relief of exercise-induced breathlessness in severe COPD. *BMC Pulmonary Medicine* 2011;**11**:23.

Oliveira 2012a {published and unpublished data}

Oliveira MF, Rodrigues MK, Treptow E, Cunha TM, Ferreira EM, Neder JA. Effects of oxygen supplementation on cerebral oxygenation during exercise in chronic obstructive pulmonary

disease patients not entitled to long-term oxygen therapy. *Clinical Physiology and Functional Imaging* 2012;**32**(1):52-8.

Oliveira 2012b {published and unpublished data}

Oliveira MF, Rodrigues MK, Treptow E, Cunha TM, Ferreira EM, Neder JA. Effects of oxygen supplementation on cerebral oxygenation during exercise in chronic obstructive pulmonary disease patients not entitled to long-term oxygen therapy. *Clinical Physiology and Functional Imaging* 2012;**32**(1):52-8.

Ringbaek 2013 {published and unpublished data}

Ringbaek T, Martinez G, Lange P. The long-term effect of ambulatory oxygen in normoxaemic COPD patients: a randomised study. *Chronic Respiratory Disease* 2013;**10**(2):77-84.

Rooyackers 1997 {published data only}

Rooyackers JM, Dekhuijzen PNR, Van Herwaarden CLA, Folgering HTM. Training with supplemental oxygen in patients with COPD and hypoxaemia at peak exercise. *European Respiratory Journal* 1997;**10**:1278-84.

Rooyackers 1997a {published data only}

Rooyackers JM, Dekhuijzen PNR, Van Herwaarden CLA, Folgering HTM. Training with supplemental oxygen in patients with COPD and hypoxaemia at peak exercise. *European Respiratory Journal* 1997;**10**(6):1278-84.

Rooyackers 1997b {published data only}

Rooyackers JM, Dekhuijzen PNR, Van Herwaarden CLA, Folgering HTM. Training with supplemental oxygen in patients with COPD and hypoxaemia at peak exercise. *European Respiratory Journal* 1997;**10**:1278-84.

Scorsone 2010 {published data only}

Scorsone D, Bartolini S, Saporiti R, Braido F, Baroffio M, Pellegrino R, et al. Does a low-density gas mixture or oxygen supplementation improve exercise training in COPD?. *Chest* 2010;**138**(5):1133-9.

Somfay 2001 {published data only}

Somfay A, Porszasz J, Lee SM, Casaburi R. Dose-response effect of oxygen on hyperinflation and exercise endurance in non-hypoxemic COPD patients. *European Respiratory Journal* 2001;**18**(1):77-84.

Spielmanns 2014 {published and unpublished data}

Spielmanns M, Fuchs-Bergsma C, Winkler A, Fox G, Krüger S, Baum K. Effects of oxygen supply during training on subjects with COPD who are normoxemic at rest and during exercise: a blinded randomized controlled trial. *Respiratory Care* 2015;**60**:540-8.

Swinburn 1984 {published data only}

Swinburn CR, Wakefield JM, Jones PW. Relationship between ventilation and breathlessness during exercise in chronic obstructive airways disease is not altered by prevention of hypoxaemia. *Clinical Sciences* 1984;**67**:515-9.

Voduc 2010 {published data only}

Voduc N, Tessier C, Sabri E, Fergusson D, Lavalée L, Aaron SD. Effects of oxygen on exercise duration in chronic obstructive pulmonary disease patients before and after pulmonary rehabilitation. *Canadian Respiratory Journal [Revue canadienne de pneumologie]* 2010;**17**(1):e14-e9.

Wadell 2001 {published data only}

Janaudis-Ferreira T, Henriksson-Larsen K, Lundgren R, Wadell K. Differences in training effects following training with and without supplemental oxygen in patients with COPD. *Advances in Physiotherapy* 2009;**11**(4):186-92.

* Wadell K, Henriksson-Larsen K, Lundgren R. Physical training with and without oxygen in patients with chronic obstructive pulmonary disease and exercise-induced hypoxemia. *Journal of Rehabilitation Medicine* 2001;**33**(5):200-5.

Woodcock 1981 {published data only}

Woodcock AA, Gross ER, Geddes DM. Oxygen relieves breathlessness in "pink puffers". *Lancet* 1981;**1**(8226):907-9.

References to studies excluded from this review

Alexandre 2010 {published data only}

Alexandre P, Jaume S, Angel R, Aina N, Amanda, I, Alvar A. Oxygen therapy decreases systemic inflammation in patients with chronic obstructive pulmonary disease and hypoxemia [Abstract]. *European Respiratory Journal*. 2010:[376].

Arnold 2008 {published data only}

Arnold B. Oxygen for COPD patients [Abstract]. *Journal of Pain & Palliative Care Pharmacotherapy*. 2008; Vol. 22, issue 3:228-9.

Balkissoon 2006 {published data only}

Balkissoon R. The effect of helium and oxygen on exercise performance in chronic obstructive pulmonary disease: a randomized crossover trial. *COPD: Journal of Chronic Obstructive Pulmonary Disease* 2006;**3**:173-4.

Bjørgen 2009 {published data only}

Bjørgen S, Helgerud J, Husby V, Steinshamn S, Richadson RR, Hoff J. Aerobic high intensity one-legged interval cycling improves peak oxygen uptake in chronic obstructive pulmonary disease patients; no additional effect from hyperoxia. *International Journal of Sports Medicine* 2009;**30**(12):872-8.

Bradley 1978 {published data only}

Bradley BL, Garner AE, Billiu D, Mestas JM, Forman J. Oxygen-assisted exercise in chronic obstructive lung disease. *American Review of Respiratory Disease* 1978;**118**:239-43.

Bye 1985 {published data only}

Bye PTP, Esau SA, Levy RD, Shiner RJ, Macklem PT, Martin JG, et al. Ventilatory muscle function during exercise in air and oxygen in patients with chronic air-flow limitation. *American Review of Respiratory Disease* 1985;**132**:236-40.

Campbell 2013 {published data only}

Campbell M, Yarandi H. The benefit of routine oxygen administration for terminally ill patients who are near death [Abstract]. *Journal of Pain and Symptom Management*. Conference: Annual Assembly of the American Academy of Hospice and Palliative Medicine and the Hospice and Palliative Nurses Association, 2012, Denver, CO, United States. 2012; Vol. 43, issue 2:336-7.

* Campbell ML, Yarandi H, Dove-Medows E. Oxygen is nonbeneficial for most patients who are near death. *Journal of Pain and Symptom Management* 2013;**45**(3):517-23.

Caspersen 2012 {published data only}

Caspersen NF, Karlsen CG, Edvardsen A, Skumlien S. Reduced exertional dyspnea with supplemental oxygen in patients with COPD - Characteristics of responders and non-responders [Abstract]. *European Respiratory Journal* 2012;**40**(Suppl 56):548s [3094].

Criner 1987 {published data only}

Criner GJ, Celli BR. Ventilatory muscle recruitment in exercise with O₂ in obstructed patients with mild hypoxemia. *Journal of Applied Physiology* 1987;**63**:195-200.

Cuvelier 2002 {published data only}

Cuvelier A, Nuir J-F, Chakroun N. Refillable oxygen cylinders may be an alternative for ambulatory oxygen therapy in COPD. *Chest* 2002;**122**:451-6.

Edvardsen 2007 {published data only}

Edvardsen A, Christensen CC, Skjongsberg OH, Ryg M. Effect of high dose oxygen on dyspnea and exercise tolerance in patients with COPD given LTOT. *European Respiratory Journal* 2007;**30**(51):513s [E3078].

Evans 1986 {published data only}

Evans TW, Waterhouse JC, Carter A, Nicholl JF, Howard P. Short burst oxygen treatment for breathlessness in chronic obstructive airways disease. *Thorax* 1986;**41**:611-5.

Fraser 2013 {published data only}

Fraser JF, Corley A, Spooner A, Dunster KR. Nasal high flow with supplemental oxygen modifies ventilation and reduces tissue CO₂ In COPD patients. *American Journal of Respiratory and Critical Care Medicine* 2013;**187**:A4381.

Fujimoto 2002 {published data only}

Fujimoto K, Matsuzawa Y, Yamaguchi S, Koizumi T, Kubo K. Benefits of oxygen on exercise performance and pulmonary hemodynamics in patients with COPD with mild hypoxemia. *Chest* 2002;**122**:457-63.

Garrod 2000 {published data only}

Garrod R, Paul EA, Wedzicha JA. Supplemental oxygen during pulmonary rehabilitation in patients with COPD with exercise hypoxaemia. *Thorax* 2000;**55**:539-43.

Gosselin 2004 {published data only}

Gosselin N, Durand F, Poulain M, Lambert K, Ceugniet F, Prefaut C, et al. Effect of acute hyperoxia during exercise on

quadriceps electrical activity in active COPD patients. *Acta Physiologica Scandinavica* 2004;**181**:333-43.

Heinzelmann 2011 {published data only}

Heinzelmann I, Kenn K. Characteristics of 'oxygen responder' before and after rehabilitation [Abstract]. *American Journal of Respiratory and Critical Care Medicine* 2010;**181**:A3580.

* Heinzelmann I, Kenn K. The influence of O₂ on exercise capacity (6-minute-walking-distance, 6MWD) in COPD [Abstract]. *European Respiratory Journal* 2011;**38**(Suppl 55).

Helgerud 2010 {published data only}

Helgerud J, Bjorgen S, Karlsen T, Husby VS, Steinshamn S, Richardson RS, et al. Hyperoxic interval training in chronic obstructive pulmonary disease patients with oxygen desaturation at peak exercise. *Scandinavian Journal of Medicine & Science in Sports* 2010;**20**(1):e170-6.

King 1973 {published data only}

King AJ, Cooke NJ, Leitch AG, Flenley DC. The effects of 30% oxygen on the respiratory response to treadmill exercise in chronic respiratory failure. *Clinical Science* 1973;**44**:151-62.

King 2012 {published data only}

King C, Young P, Fergusson W, Eaton T, Kolbe J. No benefit from short-burst oxygen therapy for dyspnoeic COPD patients after activities of daily living [Abstract]. 2012 Annual Scientific Meetings of the Thoracic Society of Australia and New Zealand and the Australian and New Zealand Society of Respiratory Science; 2012 Mar 30-Apr 4; Canberra. 2012, issue Suppl 1:59.

Lane 1987 {published data only}

Lane R, Cockcroft A, Adams L, Guz A. Arterial oxygen saturation and breathlessness in patients with chronic obstructive airways disease. *Clinical Science* 1987;**72**:693-8.

Leggett 1977 {published data only}

Leggett RJE, Flenley DC. Portable oxygen and exercise tolerance in patients with chronic hypoxic cor pulmonale. *British Medical Journal* 1977;**2**(6079):84-6.

Lellouche 2011 {published data only}

Lellouche F, Maltais F, Bouchard PA, Brouillard C, L'Her E. Free O₂: closed-loop automatic titration of oxygen flow based on SpO₂ evaluation in COPD patients during endurance shuttle walking [Abstract]. *Canadian Respiratory Journal [Revue canadienne de pneumologie]*. 2011; Vol. 18:22A.

Light 1989 {published data only}

Light RW, Mahutte CK, Stansbury DW, Fischer CE, Brown SE. Relationship between improvement in exercise performance with supplemental oxygen and hypoxic ventilatory drive in patients with chronic airflow obstruction. *Chest* 1989;**95**:751-6.

Liss 1988 {published data only}

Liss HP, Grant BJB. The effect of nasal flow on breathlessness in patients with chronic obstructive pulmonary disease. *American Review of Respiratory Disease* 1988;**137**:1285-8.

Lock 1992 {published data only}

Lock SH, Blower G, Prynn M, Wedzicha JA. Comparison of liquid and gaseous oxygen for domiciliary portable use. *Thorax* 1992;**47**:98-100.

Maldonado 2014 {published data only}

Maldonado D, Gonzalez-Garcia M, Barrero M, Jaramillo C, Casas A. Exercise endurance in chronic obstructive pulmonary disease patients at an altitude of 2640 meters breathing air and oxygen (FIO₂ 28% and 35%): a randomized crossover trial. *COPD* 2014;**11**(4):401-6.

Mannix 1992 {published data only}

Mannix ET, Manfredi F, Palange P, Dowdeswell IR, Farber M. Oxygen may lower the O₂ cost of ventilation in chronic obstructive lung disease. *Chest* 1992;**101**:910-5.

Marques-Magallanes 1998 {published data only}

Marques-Magallanes JA, Storer TW, Cooper CB. Treadmill exercise duration and dyspnea recovery time in chronic obstructive pulmonary disease: effects of oxygen breathing and repeated testing. *Respiratory Medicine* 1998;**92**:735-8.

Matsuzawa {published data only}

Matsuzawa Y, Kubo K, Fujimoto K, Eda S, Yamaguchi S, Koizumi T, et al. Acute effects of oxygen on dyspnea and exercise tolerance in patients with pulmonary emphysema with only mild exercise-induced oxyhemoglobin desaturation [Japanese]. *Nihon Kokyuki Gakkai Zasshi* 2000;**38**(11):831-5.

Nasilowski 2008 {published data only}

Nasilowski J, Przybylowski T, Zielinski J, Chazan R. Comparing supplementary oxygen benefits from a portable oxygen concentrator and a liquid oxygen portable device during a walk test in COPD patients on long-term oxygen therapy. *Respiratory Medicine* 2008;**102**(7):1021-5.

Nguyen 2008 {published data only}

Nguyen HQ, Donesky-Cuenca D, Wolpin S, Reinke LF, Benditt JO, Paul SM, et al. Randomized controlled trial of an internet-based versus face-to-face dyspnea self-management program for patients with chronic obstructive pulmonary disease: pilot study. *Journal of Medical Internet Research* 2008;**10**(2):e9.

Nosedá 1997 {published data only}

Nosedá A, Carpioux JP, Markstein C, Meyvaert A, de Maertelaer V. Disabling dyspnoea in patients with advanced disease: lack of effect of nebulized morphine. *European Respiratory Journal* 1997;**10**:1079-83.

O'Donnell 2001 {published data only}

O'Donnell DE, D'Arsigny C, Webb KA. Effects of hyperoxia on ventilatory limitation during exercise in advanced chronic obstructive pulmonary disease. *American Journal of Respiratory and Critical Care Medicine* 2001;**163**:892-8.

O'Driscoll 2003 {published data only}

O'Driscoll R, Wolstenholme RJ, Pilling A, Bassett C, Singer M, Bellingan G, et al. The use of oxygen in acute exacerbations of chronic obstructive pulmonary disease: a prospective audit

of pre-hospital and hospital emergency management. *Clinical Medicine* 2003;**3**:183-6.

O'Driscoll 2007 {published data only}

O'Driscoll BR, Siddiq Pulakal A, Neill J, Turkington P. Does short burst oxygen therapy (SBOT) relieve breathlessness after exercise in patients with severe COPD who are not hypoxemic at rest. *European Respiratory Journal* 2007;**30**(5):72s [P579].

O'Neill 2006 {published data only}

O'Neill B, Bradley JM, Mckeivitt AM, Bradbury I, Riley M, MacMahon J. Do patients prescribed short-burst oxygen therapy meet criteria for ambulatory oxygen?. *International Journal of Clinical Practice* 2006;**60**:146-9.

Ouyang 2006 {published data only}

Ouyang XH, Yin R, Zhang Y. Effects of shenmai injection on the clinical efficacy of noninvasive ventilation in patients with severe respiratory failure caused by chronic obstructive pulmonary disease. *Chinese Journal of Integrated Traditional and Western Medicine* 2006;**26**(7):608-11.

Palwai 2010 {published data only}

Palwai A, Skowronski M, Coreno A, Drummond C, McFadden ER Jr. Critical comparisons of the clinical performance of oxygen-conserving devices. *American Journal of Respiratory and Critical Care Medicine* 2010;**181**(10):1061-71.

Peters 2006 {published data only}

* Peters MM, Webb KA, O'Donnell DE. Combined physiological effects of bronchodilators and hyperoxia on exertional dyspnoea in normoxic COPD. *Thorax* 2006;**61**:559-67.

Peters MM, Webb KA, O'Donnell DE. The effects of supplemental oxygen and bronchodilators on exertional dyspnea and exercise performance in COPD. American Thoracic Society International Conference; 2005 May 20-25; San Diego. 2005:Poster A58.

Raimondi 1970 {published data only}

Raimondi AC, Edwards RHT, Denison DM, Leaver DG, Spencer RG, Siddorn JA. Exercise tolerance breathing a low density gas mixture, 35% oxygen and air in patients with chronic obstructive bronchitis. *Clinical Science* 1970;**39**:675-85.

Roberts 1996 {published data only}

Roberts CM, Bell J, Wedzicha JA. Comparison of the efficacy of a demand oxygen delivery system with continuous low flow oxygen in subjects with stable COPD and severe oxygen desaturation on walking. *Thorax* 1996;**51**:831-4.

Sandland 2008a {published data only}

Sandland CJ, Morgan MD, Singh SJ. Patterns of domestic activity and ambulatory oxygen usage in COPD. *Chest* 2008;**134**(4):753-60.

Sandland 2008b {published data only}

Sandland CJ, Morgan MD, Singh SJ. Detecting oxygen desaturation in patients with COPD: incremental versus endurance shuttle walking. *Respiratory Medicine* 2008;**102**(8):1148-52.

Sforza 2010 {published data only}

Riario Sforza GG, Incorvaia C, Carminati A, Paterniti F, Pessina L, Caligiuri R, et al. Effects of oxygen supplementation during pulmonary rehabilitation in COPD normoxemic patients [Abstract]. *European Respiratory Journal*. 2010:[P626].

Stein 1982 {published data only}

Stein DA, Bradley BL, Miller WC. Mechanisms of oxygen effects on exercise in patients with chronic obstructive pulmonary disease. *Chest* 1982;**81**:6-10.

Stevenson 2004 {published data only}

Stevenson NJ, Calverley PM. Effect of oxygen on recovery from maximal exercise in patients with chronic obstructive pulmonary disease. *Thorax* 2004;**59**:668-72.

Swinburn 1991 {published data only}

Swinburn CR, Mould H, Stone TN, Corris PA, Gibson GJ. Symptomatic benefit of supplemental oxygen in hypoxemic patients with chronic lung disease. *American Review of Respiratory Disease* 1991;**143**:913-5.

Vyas 1971 {published data only}

Vyas MN, Banister EW, Morton JW, Grzybowski S. Response to exercise in patients with chronic airway obstruction. *American Review of Respiratory Disease* 1971;**103**:401-12.

Waterhouse 1983 {published data only}

Waterhouse JC, Howard P. Breathlessness and portable oxygen in chronic obstructive airways disease. *Thorax* 1983;**38**:302-6.

Wedzicha 2006 {published data only}

Wedzicha JA. Heliox in chronic obstructive pulmonary disease: lightening the airflow. *American Journal of Respiratory and Critical Care Medicine* 2006;**173**:825-6.

Womble 2010 {published data only}

Womble HM, Johnston RP, Schwartzstein RM, Roberts DH. The effect of supplemental oxygen on maximal consumption of oxygen and ventilatory parameters in chronic obstructive pulmonary disease [Abstract]. *American Journal of Respiratory and Critical Care Medicine* 2010;**181**:A3572.

Womble 2012 {published data only}

Womble HM, Schwartzstein RM, Johnston RP, Roberts DH. Effect of therapeutic hyperoxia on maximal oxygen consumption and perioperative risk stratification in chronic obstructive pulmonary disease. *Lung* 2012;**190**:263-9.

Additional references
Abernethy 2005

Abernethy AP, Currow DC, Frith P, Fazekas B. Prescribing palliative oxygen: a clinician survey of expected benefit and patterns of use. *Palliative Medicine* 2005;**19**(2):168-70.

Ameer 2014

Ameer F, Carson KV, Usmani ZA, Smith BJ. Ambulatory oxygen for people with chronic obstructive pulmonary disease who

are not hypoxaemic at rest. *Cochrane Database of Systematic Reviews* 2014, Issue 6. [DOI: [10.1002/14651858.CD000238.pub2](https://doi.org/10.1002/14651858.CD000238.pub2)]

Cranston 2005

Cranston JM, Crockett AJ, Moss JR, Alpers JH. Domiciliary oxygen for chronic obstructive pulmonary disease. *Cochrane Database of Systematic Reviews* 2005, Issue 4. [DOI: [10.1002/14651858.CD001744.pub2](https://doi.org/10.1002/14651858.CD001744.pub2)]

Cranston 2008

Cranston JM, Crockett A, Currow D. Oxygen therapy for dyspnoea in adults. *Cochrane Database of Systematic Reviews* 2008;**3**:CD004769.

Ekstrom 2015a

Ekstrom M, Nilsson F, Abernethy AA, Currow DC. Effects of opioids on breathlessness and exercise capacity in chronic obstructive pulmonary disease: a systematic review. *Annals of the American Thoracic Society* 2015;**12**(7):1079-92.

Ekstrom 2015b

Ekstrom MP, Abernethy AP, Currow DC. The management of chronic breathlessness in patients with advanced and terminal illness. *BMJ* 2015;**349**:g7617.

Follmann 1992

Follmann D, Elliott P, Suh I, Cutler J. Variance imputation for overviews of clinical trials with continuous response. *Journal of Clinical Epidemiology* 1992;**45**(7):769-73.

Gimeno-Santos 2015

Gimeno-Santos E, Raste Y, Demeyer H, Louvaris Z, de Jong C, Rabinovich RA, et al. The PROactive instruments to measure physical activity in patients with chronic obstructive pulmonary disease. *The European Respiratory Journal* 2015;**46**(4):988-1000.

Guyatt 2000

Guyatt GH, McKim DA, Austin P, Bryan R, Norgren J, Weaver B, et al. Appropriateness of domiciliary oxygen delivery. *Chest* 2000;**118**(5):1303-8.

Hardinge 2015

Hardinge M, Annandale J, Bourne S, Cooper B, Evans A, Freeman D, et al. British Thoracic Society guidelines for home oxygen use in adults. *Thorax* 2015;**70** Suppl 1:i1-43.

Higgins 2011

Higgins JP, Altman DG, Gotzsche PC, Juni P, Moher D, Oxman AD, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ* 2011;**343**:d5928.

Johnson 2013

Johnson MJ, Bland JM, Oxberry SG, Abernethy AP, Currow DC. Clinically important differences in the intensity of chronic refractory breathlessness. *Journal of Pain and Symptom Management* 2013;**46**(6):957-63.

Johnson 2016

Johnson MJ, Booth S, Currow DC, Lam LT, Phillips JL. A mixed-methods, randomized, controlled feasibility trial to inform the design of a phase III trial to test the effect of the handheld fan

on physical activity and carer anxiety in patients with refractory breathlessness. *Journal of Pain and Symptom Management* 2016;**51**(5):807-15.

Laviolette 2014

Laviolette L, Laveneziana P, Faculty ERSRS. Dyspnoea: a multidimensional and multidisciplinary approach. *The European Respiratory Journal* 2014;**43**(6):1750-62.

McCarthy 2015

McCarthy B, Casey D, Devane D, Murphy K, Murphy E, Lacasse Y. Pulmonary rehabilitation for chronic obstructive pulmonary disease. *Cochrane Database of Systematic Reviews* 2015, Issue 2. [DOI: [10.1002/14651858.CD003793.pub3](https://doi.org/10.1002/14651858.CD003793.pub3)]

MRCWP 1981

Medical Research Council Working Party. Long term domiciliary oxygen therapy in chronic hypoxic cor pulmonale complicating chronic bronchitis and emphysema. Report of the Medical Research Council Working Party. *Lancet* 1981;**1**(8222):681-6.

NOTTG 1980

Nocturnal Oxygen Therapy Trial Group. Continuous or nocturnal oxygen therapy in hypoxemic chronic obstructive lung disease: a clinical trial. *Annals of Internal Medicine* 1980;**93**(3):391-8.

Parshall 2012

Parshall MB, Schwartzstein RM, Adams L, Banzett RB, Manning HL, Bourbeau J, et al. An official American Thoracic Society statement: update on the mechanisms, assessment, and management of dyspnea. *American Journal of Respiratory and Critical Care Medicine* 2012;**185**(4):435-52.

RevMan 2014 [Computer program]

The Nordic Cochrane Centre, The Cochrane Collaboration. Review Manager (RevMan). Version 5.3. Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration, 2014.

Robb 2003

Robb BW, Hungness ES, Hershko DD, Warden GD, Kagan RJ. Home oxygen therapy: adjunct or risk factor?. *Journal of Burn Care and Rehabilitation* 2003;**24**:403-6.

Schwartzstein 1987

Schwartzstein RM, Lahive K, Pope A, Weinberger SE, Weiss JW. Cold facial stimulation reduces breathlessness induced in normal subjects. *American Review of Respiratory Disease* 1987;**136**(1):58-61.

Stringer 2004

Stringer E, McParland C, Hernandez P. Physician practices for prescribing supplemental oxygen in the palliative care setting. *Journal of Palliative Care* 2004;**20**(4):303-7.

Tanash 2015

Tanash HA, Huss F, Ekstrom M. The risk of burn injury during long-term oxygen therapy: a 17-year longitudinal national study in Sweden. *International Journal of Chronic Obstructive Pulmonary Disease* 2015;**10**:2479-84.

Uronis 2015

Uronis HE, Ekstrom MP, Currow DC, McCrory DC, Samsa GP, Abernethy AP. Oxygen for relief of dyspnoea in people with chronic obstructive pulmonary disease who would not qualify for home oxygen: a systematic review and meta-analysis. *Thorax* 2015;**70**(5):492-4.

Vesteng 2015

Vesteng LK, Skjørten I, Melsom M, Edvardsen A. COPD and oxygen supplementation: improvement in inspiratory capacity and dyspnea during exercise. *European Respiratory Journal* 2015;**46**:PA4603.

References to other published versions of this review

Uronis 2011

Uronis H, McCrory DC, Samsa G, Currow D, Abernethy A. Symptomatic oxygen for non-hypoxaemic chronic obstructive pulmonary disease. *Cochrane Database of Systematic Reviews* 2011, Issue 6. [DOI: [10.1002/14651858.CD006429.pub2](https://doi.org/10.1002/14651858.CD006429.pub2)]

* Indicates the major publication for the study

CHARACTERISTICS OF STUDIES

Characteristics of included studies [ordered by study ID]

Abernethy 2010

Methods	Randomised, controlled, triple-blinded (participants, practitioners, investigators), parallel-group trial
Participants	<p>Inclusion criteria: patients with COPD older than 18 years; PaO₂ > 7.3 kPa; refractory breathlessness related to life-limiting illness (as determined by referring physicians); maximum treatment for underlying disease; breathlessness at rest or with negligible exertion ≥ 3 on the MRC categorical breathlessness scale; on stable medications during the previous week; expected survival ≥ 1 month (as judged by the responsible physician))</p> <p>Exclusion criteria: met criteria for long-term oxygen therapy; history of hypercarbic respiratory failure related to oxygen; anaemia (haemoglobin < 100 g/L); hypercapnia (PaCO₂ > 6.7 kPa); cognitive impairment (Folstein mini-mental status score < 24/30); smoking; respiratory or cardiac event in the previous week</p>

Abernethy 2010 (Continued)

Setting: outpatient pulmonary, palliative care, oncology and primary care clinics at 5 sites in Australia, 2 in the USA and 2 in the UK

Method of recruitment of participants: patients referred to the clinics

239 participants randomised (211 completed)

152 participants with COPD

Male 103 (68%)

Mean age: 73.2 (SD 9.9)

Lung function: no data available

Mean baseline SaO₂ (%): no data available

Mean baseline PaO₂ (kPa): 10.0 (SD 1.5)

Mean PaCO₂ (kPa): 5.2 (SD 0.6)

Mean breathlessness morning and evening NRS scores at baseline: 4.7 (SD 2.0) overall; 4.6 (SD 2.0) in the oxygen group; 4.8 (SD 2.0) in the air group

Interventions	Domiciliary oxygen (2 L/min) versus room air (2 L/min) delivered continuously from concentrator by nasal cannula; prescribed for at least 15 hours per day. Duration 7 days
Outcomes	<p>Breathlessness measured by NRS (0-10, 0 "none" to 10 "maximal"), defined as breathlessness "right now" during morning and evening. Mean of morning and evening scores used in the meta-analysis</p> <p>Overall well-being on NRS (0-10) scores in the evening day 7</p> <p>Treatment preference - categorical at day 7</p> <p>Adverse effects - day 3 telephone contact, reported drowsiness, nasal irritation, nasal bleeds, anxiousness</p>

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"Randomised to treatment by a central computer-generated system available (via web or telephone) through the pharmacy service at Repatriation General Hospital (Adelaide, Australia) with balanced blocks of four patients per stratum, on the basis of Fisher and Yates' statistical tables."
Allocation concealment (selection bias)	Low risk	"Patients, individuals delivering the interventions, investigators, and nurses were masked to treatment assignments. Oxygen and room air concentrators were identical in appearance. By use of a standard protocol, the medical gas company serving each site modified half the concentrators to dispense room air without setting off the internal alarm that sounds when oxygen concentrations are low."
Blinding of participants and personnel (performance bias) All outcomes	Low risk	As above
Blinding of outcome assessment (detection bias)	Low risk	As above

Abernethy 2010 (Continued)

All outcomes

Incomplete outcome data (attrition bias) All outcomes	Unclear risk	7% attrition for oxygen (overall) and 17% for air (overall)
Selective reporting (reporting bias)	Low risk	According to the aims, adequately reported variables
Other bias	Low risk	We judged that this trial appeared to be free of other sources of bias.

Bruni 2012a

Methods	Randomised, controlled, double-blinded (participants and investigators), cross-over trial
Participants	<p>Inclusion criteria: no formal criteria reported. Male patients with severe airway obstruction and hyperinflation participated in the study. COPD was diagnosed on the basis of history, physical examination, chest radiograph and results of pulmonary function studies. All patients had a long history of smoking and were clinically stable and taking appropriate medication</p> <p>Exclusion criteria: not stated</p> <p>Setting: laboratory (Italy)</p> <p>Method of recruitment of participants: not described</p> <p>10 participants with COPD classified as "hyperinflators" who exhibited a 0.47 (SD 0.35 L) increase in chest wall volume</p> <p>All male</p> <p>Mean age: 68.1 (SD 6.6)</p> <p>Mean FEV₁ (L): 0.91 (SD 0.3)</p> <p>Mean FEV₁ (% predicted): 32.7 (SD 8.7)</p> <p>FEV₁/FVC: 0.29 (SD 0.1)</p> <p>Mean baseline SaO₂ (%): no data available</p> <p>Mean baseline PaO₂ (kPa): 9.5 (SD 1.2)</p> <p>Mean PaCO₂ (kPa): 5.6 (SD 0.6)</p> <p>Mean breathlessness at baseline: no data available</p>
Interventions	<p>Oxygen (FiO₂ 0.5) vs room air during 1 symptom-limited cycle test at 75% of peak work rate</p> <p>Participants wore a nose-clip and breathed through a low-dead-space mouthpiece; Douglas bag</p>
Outcomes	<p>Breathlessness measured by modified Borg at iso-time (mBorg 0-10, 0 "none" to 10 "maximal")</p> <p>HRQOL – no data available</p> <p>Treatment preference - no data available</p> <p>Adverse effects - no data available</p>
Notes	

Bruni 2012a (Continued)

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	"In randomised order"; other information not available
Allocation concealment (selection bias)	Unclear risk	Information not available
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"Subjects were blinded to the oxygen concentration being breathed as was the investigator evaluating subjective responses and performing data analysis."
Blinding of outcome assessment (detection bias) All outcomes	Low risk	No data on whether saturation was measured; but self-rated
Incomplete outcome data (attrition bias) All outcomes	Low risk	All participants completed the study.
Selective reporting (reporting bias)	Low risk	Important outcomes were reported.
Other bias	Unclear risk	Unclear how participants were selected and whether participants withdrew; data reported only for participants completing the study

Bruni 2012b

Methods	Randomised, controlled, double-blinded (participants, investigators), cross-over trial
Participants	<p>Inclusion criteria: no formal criteria reported. Male patients with severe airway obstruction and hyperinflation participated in the study. COPD was diagnosed on the basis of history, physical examination, chest radiograph and results of pulmonary function studies. All patients had a long history of smoking and were clinically stable and taking appropriate medication.</p> <p>Exclusion criteria: not stated</p> <p>Setting: laboratory (Italy)</p> <p>Method of recruitment of participants: not described</p> <p>6 participants with COPD classified as "non-hyperinflators" because of constant chest wall volume</p> <p>Mean age: 68.1 (SD 6.6)</p> <p>Mean FEV₁ (L): 0.91 (SD 0.3)</p> <p>Mean FEV₁ (% predicted): 32.7 (SD 8.7)</p> <p>FEV₁/FVC: 0.29 (SD 0.1)</p> <p>Mean baseline SaO₂ (%): no data available</p> <p>Mean baseline PaO₂ (kPa): 10.3 (SD 0.9)</p>

Bruni 2012b (Continued)

Mean PaCO₂ (kPa): 5.2 (SD 0.1)

Mean breathlessness at baseline: no data available

Interventions	Oxygen (FiO ₂ 0.5) vs room air during 1 symptom-limited cycle test at 75% of peak work rate Participants wore a nose-clip and breathed through a low-dead-space mouthpiece; Douglas bag
Outcomes	Breathlessness measured by modified Borg at iso-time (mBorg 0-10, 0 "none" to 10 "maximal") HRQOL – no data available Treatment preference - no data available Adverse effects - no data available
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	"In randomised order"; other information not available
Allocation concealment (selection bias)	Unclear risk	Information not available
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"Subjects were blinded to the oxygen concentration being breathed as was the investigator evaluating subjective responses and performing data analysis."
Blinding of outcome assessment (detection bias) All outcomes	Low risk	No data on whether saturation was measured; but self-rated
Incomplete outcome data (attrition bias) All outcomes	Low risk	All participants completed the study.
Selective reporting (reporting bias)	Low risk	Important outcomes were reported.
Other bias	Unclear risk	It is unclear whether or not there was risk of other bias due to the lack of detail in the methods.

Davidson 1988

Methods	Randomised, double-blinded (participants and investigators), cross-over trial
Participants	Inclusion criteria: exercise tolerance limited by breathlessness secondary to severe chronic airflow obstruction Exclusion criteria: angina; impaired cardiac function or locomotor disability that might contribute to exercise limitation Setting: laboratory

Davidson 1988 (Continued)

17 participants (14 completed)
Gender not specified

Mean age: 64.4 (SEM 2.1)

Mean FEV₁ (L): 0.79 (SEM 0.0)
Mean FVC (L): 2.1 (SEM 0.11)

Mean baseline SaO₂ (%): no data available
Mean baseline PaO₂ (kPa): 8.6 (SEM 0.3)

Mean PaCO₂ (kPa): 6.0 (SEM 0.4)

Mean breathlessness at baseline: no data available

Interventions	Compressed air (4 L/min) vs oxygen (2, 4 or 6 L/min) during 6MWT, cycle ergometer test or endurance shuttle walk test
Outcomes	Breathlessness measured by VAS at the end of exercise (10 cm VAS, "not at all breathless" to "extremely breathless") HRQOL – no data available Treatment preference - no data available Adverse effects - no data available

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Described as randomised: "The order of testing in both the oxygen dosing study and the walking tests was allocated randomly"; other information not available
Allocation concealment (selection bias)	Unclear risk	Information not available
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Described as double-blind (walking test) and single-blind (cycle test); other information not available "For walking exercise the tests were performed double blind. For cycling endurance it was necessary for the investigator to determine the bias flow composition and this part of the study was therefore single (patient) blind. To ensure that this did not affect performance, care was taken to provide a similar degree of encouragement as detailed above."
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	As above
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Data reported only for participants completing the study. "Not all patients completed six minutes of exercise when breathing air or the low concentration of oxygen." Reported VAS scores only for 4 L/min for 14 participants after 6 minutes of exercise

Davidson 1988 (Continued)

Selective reporting (reporting bias)	Unclear risk	As above
Other bias	Unclear risk	It is unclear whether or not there was risk of other bias due to the lack of detail in the methods.

Dean 1992

Methods	Randomised, double-blinded (participants and investigators), cross-over trial
Participants	<p>Inclusion criteria: age > 50; DLCO < 80% predicted; extensive smoking history; resting PaO₂ > 55 mmHg</p> <p>Exclusion criteria: active coronary artery disease; congestive heart failure; vascular, orthopaedic or neurological problems that would interfere with cycling</p> <p>Setting: laboratory</p> <p>12 participants All male Age > 50, but specifics not reported Mean FEV₁ (L): 0.89 (SEM 0.1) Mean FVC (L): 2.4 (SEM 0.2) Mean DLCO (mL/min/mmHg): 9.8 (SEM 1.5)</p> <p>Mean baseline SaO₂ (%): no data available Mean baseline PaO₂ (kPa): 9.5 (SE 0.3)</p> <p>Mean baseline PaCO₂ (kPa): no data available</p> <p>Mean breathlessness at baseline: no data available</p>
Interventions	Compressed air vs 40% oxygen during incremental and endurance exercise shuttle walk tests
Outcomes	<p>Breathlessness measured by modified Borg at iso-time (mBorg)</p> <p>HRQOL – no data available</p> <p>Treatment preference - no data available Adverse effects - no data available</p>
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Described as randomised; other information not available
Allocation concealment (selection bias)	Unclear risk	Information not available
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Described as double-blind (participants, study personnel)

Dean 1992 (Continued)

Blinding of outcome assessment (detection bias) All outcomes	Low risk	"Oximetry values were not available to the patient or supervising physician during endurance testing to preserve blinding."
Incomplete outcome data (attrition bias) All outcomes	Low risk	All participants completed the study.
Selective reporting (reporting bias)	Low risk	Important outcomes were reported.
Other bias	Unclear risk	It is unclear whether or not there was risk of other bias due to the lack of detail in the methods.

Dyer 2012

Methods	Randomised, controlled, single-blinded (investigators), parallel-group trial
Participants	<p>Inclusion criteria: COPD; desaturated by > 4% to < 90% on exertion at baseline; improved $\geq 10\%$ in ESWT with ambulatory O₂ (mean 280 m)</p> <p>Exclusion criteria: any existing home oxygen</p> <p>Setting: laboratory (Surrey, UK)</p> <p>Method of recruitment of participants: consecutive patients with COPD referred to 3 PR services in Surrey, UK</p> <p>55 participants with COPD (47 completed)</p> <p>Male 30 (64% of 47 completing)</p> <p>Mean age: air 70 (SD 7); oxygen 68 (SD 8)</p> <p>Mean FEV₁ (L): 1.12 (SD 0.5); oxygen 0.96 (SD 0.4)</p> <p>Mean FEV₁ (% predicted): room air 44 (SD 11); oxygen 39 (SD 16)</p> <p>Mean baseline SaO₂ (%): 94 (SD 2)</p> <p>Mean baseline PaO₂ (kPa): no data available</p> <p>Mean baseline PaCO₂ (kPa): no data available</p> <p>Mean breathlessness at baseline: CRQ dyspnoea score 3 (SD 1); MRC 3 (SD 1)</p>
Interventions	Ambulatory oxygen (2-6 L/min) vs room air during PR and during exertion at home; delivered via nasal prongs; most participants had cylinders and a few were supplied with liquid oxygen. Duration of treatment: 6-7 weeks of training, as well as at home during all activities that induced breathlessness but not at rest
Outcomes	<p>Breathlessness measured by CRQ dyspnoea subdomain on a 7-point Likert scale (higher = good)</p> <p>HRQOL - CRQ subdomains</p> <p>Treatment preference - no data available</p> <p>Adverse effects - number of dropouts</p>

Dyer 2012 (Continued)

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	By random number tables and sealed envelopes
Allocation concealment (selection bias)	High risk	Measured SaO ₂ during PR; participants were unblinded; limited blinding for staff. "It was not possible to blind participants to group allocation as sham O ₂ was not available [...] due to limited study resources."
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Single-blinded (staff) "Exercise tolerance at the end of PR for all participants was reassessed by [author], with blinding maintained by all participants wearing nasal cannulae with an oxygen cylinder attached, carried in a backpack by an assistant, which was switched off for the RA group. At the end of PR, exercise tolerance was reassessed by [author] (blind to group allocation)."
Blinding of outcome assessment (detection bias) All outcomes	High risk	As above
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Data reported only for participants completing the study. The number of dropouts was equal in both groups.
Selective reporting (reporting bias)	Unclear risk	No protocol. Did not report breathlessness during exercise test after PR
Other bias	High risk	Per-protocol analysis only

Eaton 2002

Methods	Randomised, controlled, double-blind (participants, investigators), cross-over trial
Participants	<p>Inclusion criteria: COPD as defined by ATS criteria, exertional breathlessness impacting daily activities, not fulfilling criteria for LTOT, exertional desaturation (O₂ saturation ≤ 88%), ex-smoker, clinically stable for 2 months with standard optimal medical care, completion of a formal 6-week pulmonary rehabilitation programme</p> <p>Exclusion criteria: "important co-morbidities (e.g. limiting angina or significant musculoskeletal disability)"</p> <p>Setting: non-laboratory</p> <p>50 participants (41 completed) 70% male Mean age: 57.1 (SD 9.3) Mean FEV₁ (% predicted): 25.9 (SD 8) Mean baseline SaO₂ (%): 94 (SD 1.9) Mean baseline PaO₂ (kPa): 9.2 (SD 1.0) Mean baseline PaCO₂ (kPa): 5.8 (SD 0.7)</p>

Eaton 2002 (Continued)

Mean breathlessness at baseline: 0.7 (SD 1.0)

Interventions	Compressed air (4 L/min) vs oxygen (4 L/min) during both 6MWT and 6-week period at home during which participants were instructed "to use flow rate of 4 L/min intranasally for any activity during which they would normally experience breathlessness"
Outcomes	Breathlessness measured by modified Borg (mBorg 0-10, 0 "none" to 10 "maximal") HRQOL – CRQ total score, CRQ subdomains, SF-36 Treatment preference - no data available Adverse effects - number of dropouts
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Described as randomised; other information not available
Allocation concealment (selection bias)	Unclear risk	Information not available
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"Patients were randomly assigned in a double blinded manner to cylinder air or O ₂ . All cylinders were painted pink, prefilled with either air or O ₂ and identifiable only by a unique cylinder number, ensuring blinding of both participants and observers."
Blinding of outcome assessment (detection bias) All outcomes	Low risk	As above
Incomplete outcome data (attrition bias) All outcomes	Low risk	"A mixed model approach to crossover trials was employed, which used information from all patients, including those who did not complete both time periods."
Selective reporting (reporting bias)	Low risk	Important outcomes were reported
Other bias	Unclear risk	It is unclear whether or not there was risk of other bias due to the lack of detail in the methods.

Eaton 2006

Methods	Randomised, controlled, double-blind (participants, investigators), parallel-group trial
Participants	<p>Inclusion criteria: 78 hospital inpatients with an acute exacerbation of COPD; moderate or severe COPD as defined by British Thoracic Society criteria; exertional dyspnoea interfering with daily activity; resting PaO₂ > 60 mmHg at discharge; ability to complete HRQOL questionnaires</p> <p>Exclusion criteria: current smoker; severe comorbidity likely to cause death within the 6-month study period; resident of a long-term facility in which SBOT is available; hypercapnia (PaCO₂ > 45 mmHg)</p> <p>Setting: non-laboratory</p> <p>51 participants (34 completed); 25 in oxygen group; 26 in air group 56% male in oxygen group; 50% male in air group</p>

Eaton 2006 (Continued)

Mean age: oxygen 77.4 (SD 7.2); air 77.6 (SD 5.7)

Mean FEV₁ (L): oxygen 0.9 (SD 0.5); air 0.9 (SD 0.4)

Mean FEV₁ (% predicted): oxygen 44 (SD 22); air 39 (SD 17)

Mean baseline SaO₂ (%): oxygen 95 (SD 1.9); air 95 (SD 1.6)

Mean baseline PaO₂ (kPa): oxygen 9.6 (SD 1.3); air 10.1 (SD 1.7)

Mean baseline PaCO₂ (kPa): oxygen 5.3 (SD 0.6); air 5.4 (SD 0.4)

Mean breathlessness at baseline, CRD dyspnoea: oxygen 17.8 (SD 5.0); air 17.5 (SD 4.2)

Interventions	Cylinder oxygen (2 L/min) vs cylinder air (2 L/min) vs usual care during 6-month domiciliary period. Participants were given standardised instructions "to use they cylinder gas at 2 L/min via nasal prongs, as necessary for distressing or limiting breathlessness." No short-term assessments were performed.
Outcomes	Breathlessness measured by CRQ dyspnoea score HRQOL – CRQ total score, CRQ subdomains, SF-36 Treatment preference - no data available Adverse effects - number of dropouts

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"Patients were randomised using computer-generated randomisation numbers."
Allocation concealment (selection bias)	Low risk	"Allocation of cylinders was by a separate member of the research team not involved in patient assessment."
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"To ensure double-blinding, cylinders, prefilled with air or oxygen, were identifiable only by a unique cylinder number. Cylinders were painted pink to ensure they would not be used in routine clinical care."
Blinding of outcome assessment (detection bias) All outcomes	Low risk	As above
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Data reported only for participants completing the study. 17 dropouts in total; number of dropouts was equal in both groups
Selective reporting (reporting bias)	Low risk	Important outcomes were reported.
Other bias	Low risk	We judged that this trial appeared to be free of other sources of bias.

Emtner 2003a

Methods	Randomised, controlled, double-blind (participants, investigators), parallel-group trial
---------	--

Emtner 2003a (Continued)

Participants

Inclusion criteria: COPD that is clinically stable with no recent exacerbations; FEV₁ < 50% predicted; FEV₁/VC < 0.65; resting PaO₂ > 55 mmHg; SpO₂ > 88% during constant work rate test while breathing room air

Exclusion criteria: symptomatic cardiovascular comorbidity or other disease that might contribute to exercise limitation; regular participation in a formal exercise programme or participation in a formal rehabilitation programme within the past 2 years

Setting: laboratory

30 participants overall (29 completed)

15 participants categorised as "air trained group"

10 male and 5 female

Mean age: 67 (SD 10)

Mean FEV₁ (L): 1.01 (SD 0.27)

Mean FVC (L): 2.58 (SD 0.7)

Mean TLC (L): 7.3 (SD 1.2)

Mean RV (L): 4.6 (SD 1.3)

Mean DLCO (mL/min/mmHg): 11.4 (SD 4.4)

Mean baseline SaO₂ (%): no data available

Mean baseline PaO₂ (kPa): 9.8 (SD 1.0)

Mean baseline PaCO₂ (kPa): 5.5 (SD 0.5)

Mean breathlessness at baseline: no data available

Interventions

Supplemental oxygen (FiO₂ 0.30) vs compressed air during constant work rate exercise

During training sessions, gas was delivered at 3 L/min via nasal cannula, connected to the appropriate tank (compressed air or oxygen) by an unblinded investigator

Incremental and constant work rate tests were performed by inhaling compressed air or 30% oxygen (4 tests). Participants were blinded to the inhaled mixture and breathed through mouthpiece and low-resistance valve with a nose-clip in place.

Outcomes

Breathlessness measured by modified Borg (mBorg 0-10, 0 "none" to 10 "maximal")

HRQOL – CRQ total score, CRQ subdomains, SF-36

Treatment preference - no data available

Adverse effects - number of dropouts

Notes
Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"The randomization sequence was generated using numbered sealed envelopes."
Allocation concealment (selection bias)	Low risk	Sealed envelopes were used to conceal generated randomisation sequence from study investigators.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"The nasal cannula tubing was connected to the appropriate tank (compressed air or oxygen) by an unblinded investigator. Patient and staff did not know which gas mixture the patient received (...). Exercise intensity was subsequently adjusted, considering the subject's breathlessness and fatigue sensations, by blinded therapists."

Emtner 2003a (Continued)

Blinding of outcome assessment (detection bias) All outcomes	Low risk	"An unblinded investigator measured oxygen saturation periodically; the findings were never reported to the patients or therapists."
Incomplete outcome data (attrition bias) All outcomes	Low risk	No participants withdrew from this arm of treatment.
Selective reporting (reporting bias)	Low risk	According to the aims, adequately reported variables
Other bias	Low risk	We judged that this trial appeared to be free of other sources of bias.

Emtner 2003b

Methods	Randomised, controlled, double-blind (participants, investigators), parallel-group trial
Participants	<p>Inclusion criteria: COPD that is clinically stable with no recent exacerbations; FEV₁ < 50% predicted; FEV₁/VC < 0.65; resting PaO₂ > 55 mmHg; SpO₂ ≥ 88% during constant work rate test while breathing room air</p> <p>Exclusion criteria: symptomatic cardiovascular comorbidity or other disease that might contribute to exercise limitation; regular participation in a formal exercise programme or participation in a formal rehabilitation programme within the past 2 years</p> <p>Setting: laboratory</p> <p>30 participants overall (29 completed)</p> <p>14 participants categorised as "oxygen-trained group"</p> <p>8 male and 6 female</p> <p>Mean age: 66 (SD 7)</p> <p>Mean FEV₁ (L): 1.01 (SD 0.33)</p> <p>Mean FVC (L): 2.7 (SD 0.8)</p> <p>Mean TLC (L): 7.2 (SD 1.6)</p> <p>Mean RV (L): 4.1 (SD 0.9)</p> <p>Mean DLCO (mL/min/mmHg): 10.9 (SD 3.5)</p> <p>Mean baseline SaO₂ (%): no data available</p> <p>Mean baseline PaO₂ (kPa): 9.5 (SD 1.1)</p> <p>Mean baseline PaCO₂ (kPa): 5.4 (SD 0.7)</p> <p>Mean breathlessness at baseline: no data available</p>
Interventions	<p>Supplemental oxygen (FiO₂ 0.30) vs compressed air during constant work rate exercise</p> <p>During training sessions, gas was delivered at 3 L/min via nasal cannula, connected to the appropriate tank (compressed air or oxygen) by an unblinded investigator.</p> <p>Incremental and constant work rate tests were performed by inhaling compressed air or 30% oxygen (4 tests). Participants were blinded to the inhaled mixture and breathed through mouthpiece and low-resistance valve with a nose-clip in place.</p>
Outcomes	<p>Breathlessness measured by modified Borg (mBorg 0-10, 0 "none" to 10 "maximal")</p> <p>HRQOL – CRQ total score, CRQ subdomains, SF-36</p>

Emtner 2003b (Continued)

Treatment preference - no data available
Adverse effects - number of dropouts

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"The randomization sequence was generated using numbered sealed envelopes."
Allocation concealment (selection bias)	Low risk	Sealed envelopes were used to conceal generated randomisation sequence from study investigators.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"The nasal cannula tubing was connected to the appropriate tank (compressed air or oxygen) by an unblinded investigator. Patient and staff did not know which gas mixture the patient received (...). Exercise intensity was subsequently adjusted, considering the subject's breathlessness and fatigue sensations, by blinded therapists."
Blinding of outcome assessment (detection bias) All outcomes	Low risk	"An unblinded investigator measured oxygen saturation periodically; the findings were never reported to the patients or therapists."
Incomplete outcome data (attrition bias) All outcomes	Low risk	"One patient from the oxygen-trained group was excluded because of illness during the postintervention testing period."
Selective reporting (reporting bias)	Low risk	According to the aims, adequately reported variables
Other bias	Low risk	We judged that this trial appeared to be free of other sources of bias.

Eves 2006

Methods	Randomised, controlled, double-blinded (participants, investigators), cross-over trial
Participants	<p>Inclusion criteria: clinically stable moderate to severe COPD</p> <p>Exclusion criteria: dependence on supplemental oxygen, cardiovascular disease and/or musculoskeletal abnormality</p> <p>Setting: laboratory</p> <p>10 participants All men Mean age: 65 (SD 11) Mean FEV₁ (L): 1.66 (0.59) Mean FVC (L): 3.81 (0.99) Mean baseline SaO₂ (%): no data available Mean baseline PaO₂ (kPa): 9.1 (SD 0.9)</p> <p>Mean baseline PaCO₂ (kPa): 4.9 (SD 0.4)</p> <p>Mean breathlessness at baseline: no data available</p>

Eves 2006 (Continued)

Interventions	Medical air vs 40% oxygen vs heliox vs heliox/oxygen during constant-load cycling. Throughout exercise, humidified gases were passed into a reservoir bag and supplied through a low-resistance 2-way breathing valve.	
Outcomes	Breathlessness measured by modified Borg (mBorg 0-10, 0 "none" to 10 "maximal") HRQOL – no data available Treatment preference - no data available Adverse effects - no data available	
Notes		
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	"During the other two visits, four constant-load symptom-limited exercise trials were performed in a random order"; other information not available
Allocation concealment (selection bias)	Unclear risk	Information on concealment of allocation not available
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"Throughout exercise, humidified gases were passed into a reservoir bag and supplied through a low-resistance two-way breathing valve. The patients were blinded to the gas mixture used and were asked not to talk during, or for a short period after, exercise due to the change in vocal tone with helium."
Blinding of outcome assessment (detection bias) All outcomes	Low risk	"Two research assistants, blinded to the gas mixture used, consistently encouraged patients to exercise for as long as possible."
Incomplete outcome data (attrition bias) All outcomes	Low risk	All participants completed the study.
Selective reporting (reporting bias)	Low risk	According to the aims, adequately reported variables
Other bias	Unclear risk	It is unclear whether or not there was risk of other bias due to the lack of detail in the methods.

Haidl 2004

Methods	Randomised, controlled, unblinded, parallel-group trial
Participants	<p>Inclusion criteria: COPD diagnosis (per "current clinical guidelines"); FEV₁/FVC < 0.70; PCO₂ > 45 mmHg at rest on 2 different days or increase in PCO₂ after cycle testing > 45 mmHg; PO₂ at rest > 55 mmHg; mean nocturnal oxygen saturation > 90%</p> <p>Exclusion criteria: malignant disease; left heart failure or other significant comorbidities (e.g. severe renal failure, severe diabetes)</p> <p>Setting: non-laboratory</p> <p>28 participants (15 dropouts during 3-year follow-up) 26 (93%) male Mean age: oxygen group 65.7 (SD 6.7); controls 64.5 (SD 6.4)</p>

Haidl 2004 (Continued)

Mean FEV₁ (% predicted): oxygen group 38.8 (8.4); controls 42.7 (11.8)

Mean baseline SaO₂ (%): no data

Mean baseline PaO₂ (kPa): oxygen group 9.0 (SD 0.9); controls 8.7 (SD 0.8)

Mean baseline PaCO₂ (kPa): oxygen group 5.3 (SD 0.4); controls 5.6 (SD 0.5)

Mean breathlessness at baseline: oxygen group 5.0 (SD 2.1); controls 5.0 (SD 1.5)

Interventions	LTOT (2 L/min ≥ 15 hours/d) vs control (no treatment) delivered through nasal cannula connected to gas concentrators
Outcomes	Breathlessness measured by modified Borg (mBorg 0-10, 0 "none" to 10 "maximal") HRQOL - no data available Treatment preference - no data available Adverse effects - number of dropouts
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Information not available
Allocation concealment (selection bias)	Low risk	Participants' randomisation status was unknown to staff performing their tests.
Blinding of participants and personnel (performance bias) All outcomes	High risk	Unblinded. Control group received usual care (no sham oxygen).
Blinding of outcome assessment (detection bias) All outcomes	High risk	See above.
Incomplete outcome data (attrition bias) All outcomes	High risk	Data reported only for participants completing the study. Both treatment groups were complete at 1 year, when assessment was undertaken. Study planned to measure differences at 3 years, but attrition rates prevented this (15 dropouts during 3-year follow-up).
Selective reporting (reporting bias)	Low risk	Important outcomes were reported.
Other bias	Unclear risk	It is unclear whether or not there was risk of other bias due to the lack of detail in the methods.

Ishimine 1995

Methods	Randomised, controlled, double-blinded (participants, investigators), cross-over study
Participants	Inclusion criteria: male with "stable" COPD or chronic bronchitis; moderate to severe obstruction; PaO ₂ > 60 torr at rest Exclusion criteria: not available (not specified in Cochrane translation)

Ishimine 1995 (Continued)

Setting: laboratory

22 participants

All men

Mean age: 69 (SD 7)

Mean FEV₁ (L): 1.02 (0.51)

Mean FVC (L): 2.26 (0.57)

Mean baseline SaO₂ (%): no data

Mean baseline PaO₂ (kPa): 10.1 (SD 1.1)

Mean baseline PaCO₂ (kPa): no data

Mean breathlessness at baseline: no data

Interventions	Room air vs compressed air (3 L/min) vs oxygen (3 L/min) during 6MWT
Outcomes	Breathlessness measured by a questionnaire involving 8 questions; each question was answered on a 100-mm horizontal line with anchors from the modified Borg HRQOL – no data available Treatment preference - no data available Adverse effects - no data available
Notes	Translated from Japanese and assessed with the translation sheet

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Described as randomised; other information not available
Allocation concealment (selection bias)	Unclear risk	Information not available
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Placebo controlled; blinding of assessors could not be ascertained
Blinding of outcome assessment (detection bias) All outcomes	High risk	Blinding of assessors could not be ascertained.
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Data reported only for participants completing the study
Selective reporting (reporting bias)	Unclear risk	Information not available
Other bias	Unclear risk	Unclear how participants were selected and whether they withdrew; blinding unclear

Jolly 2001a

Methods	Randomised, double-blinded (participants, investigators), placebo-controlled, cross-over trial
---------	--

Jolly 2001a (Continued)

Participants	<p>Inclusion criteria: participants with COPD (ATS grade II/III) with ≥ 30 days of clinical stability; participating in respiratory rehab programme; FEV₁ < 55% and/or FEV₁ ratio < 50%; resting PaO₂ > 60 mmHg</p> <p>Exclusion criteria: peripheral vascular disease; cardiac failure; active CAD</p> <p>Setting: laboratory</p> <p>9 participants, classified as "non-de-saturators"</p> <p>All male</p> <p>Mean age: 70 (SEM 3)</p> <p>Mean FEV₁ (L): 0.9 (SEM 0.8)</p> <p>Mean FVC (% predicted): 63 (SEM 6)</p> <p>Mean TLC (L): 7.4 (SEM 0.4)</p> <p>Mean RV (L): 4.4 (SEM 0.39)</p> <p>Mean baseline SaO₂ (%): 96 (SEM 0.5))</p> <p>Mean baseline PaO₂ (kPa): 10.5 (SE 0.4)</p> <p>Mean baseline PaCO₂ (kPa): 5.3 (SE 0.2)</p> <p>Mean breathlessness at baseline: 0.56 (SE 0.3)</p>
Interventions	Room air vs compressed air (3, 6, 9, 12 L/min) vs oxygen (3, 6, 9, 12 L/min) during 6MWT - amount of oxygen increased on the basis of desaturation during exercise. Two indistinguishable cylinders located at the middle of the corridor - 1 with compressed air (CA) and 1 with oxygen - were connected by a Y-piece to a 15-m tube, ending in a nasal cannula.
Outcomes	Breathlessness measured by modified Borg (mBorg 0-10, 0 "none" to 10 "maximal") HRQOL – no data available Treatment preference - no data available Adverse effects - no data available
Notes	
Risk of bias	
Bias	Authors' judgement Support for judgement
Random sequence generation (selection bias)	Unclear risk Described as randomised; other information not available
Allocation concealment (selection bias)	High risk "One person, who knew the randomly assigned sequence, opened the valve and regulated the gas flow as requested by another technician, who walked behind the patient recording the SaO ₂ measured by pulse oximetry (SpO ₂) values. Both this technician and the patient were blind about which gas was added."
Blinding of participants and personnel (performance bias) All outcomes	Low risk As above, and "Two indistinguishable cylinders located at the middle of the corridor, one with compressed air (CA) and one with oxygen, were connected by a Y-piece to a 15-m tube ending in a nasal cannula."
Blinding of outcome assessment (detection bias) All outcomes	Low risk See above.
Incomplete outcome data (attrition bias) All outcomes	Low risk All participants completed the study.

Jolly 2001a (Continued)

Selective reporting (re-reporting bias)	Low risk	According to the aims, adequately reported variables
Other bias	Unclear risk	It is unclear whether or not there was risk of other bias due to the lack of detail in the methods.

Jolly 2001b

Methods	Randomised, double-blinded (participants, investigators), cross-over trial
Participants	<p>Inclusion criteria: patients with COPD (ATS grade II/III) with ≥ 30 days of clinical stability; participating in respiratory rehab programme; $FEV_1 < 55\%$ and/or FEV_1 ratio $< 50\%$, resting $PaO_2 > 60$ mmHg</p> <p>Exclusion criteria: peripheral vascular disease; cardiac failure; active CAD</p> <p>Setting: laboratory</p> <p>11 participants classified as "desaturators," defined by a fall in $SpO_2 \geq 5\%$ to $< 90\%$ during the 6MWT on air</p> <p>10 male and 1 female</p> <p>Mean age: 67 (SEM 2)</p> <p>Mean FEV_1 (L): 0.9 (SEM 0.8)</p> <p>Mean FVC (% predicted): 68 (SEM 8)</p> <p>Mean TLC (L): 7.1 (SEM 0.6)</p> <p>Mean RV (L): 4.2 (SEM 0.45)</p> <p>Mean baseline SaO_2 (%): 95 (SE 0.3)</p> <p>Mean baseline PaO_2 (kPa): 9.9 (SE 0.3)</p> <p>Mean baseline $PaCO_2$ (kPa): 5.5 (SE 0.2)</p> <p>Mean breathlessness at baseline: 0.7 (SD 1.0)</p>
Interventions	Room air vs compressed air (3, 6, 9, 12 L/min) vs oxygen (3, 6, 9, 12 L/min) during 6MWT - amount of oxygen increased on the basis of desaturation during exercise. Two indistinguishable cylinders located at the middle of the corridor - 1 with compressed air (CA) and 1 with oxygen - were connected by a Y-piece to a 15-m tube, ending in a nasal cannula.
Outcomes	<p>Breathlessness measured by modified Borg (mBorg 0-10, 0 "none" to 10 "maximal")</p> <p>HRQOL - no data available</p> <p>Treatment preference - no data available</p> <p>Adverse effects - no data available</p>

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Described as randomised; other information not available
Allocation concealment (selection bias)	High risk	"One person, who knew the randomly assigned sequence, opened the valve and regulated the gas flow as requested by another technician, who walked behind the patient recording the SaO_2 measured by pulse oximetry (SpO_2)"

Jolly 2001b (Continued)

		values. Both this technician and the patient were blind about which gas was added."
Blinding of participants and personnel (performance bias) All outcomes	Low risk	As above, and "Two indistinguishable cylinders located at the middle of the corridor, one with compressed air (CA) and one with oxygen, were connected by a Y-piece to a 15-m tube ending in a nasal cannula."
Blinding of outcome assessment (detection bias) All outcomes	Low risk	As above
Incomplete outcome data (attrition bias) All outcomes	Low risk	All participants completed the study.
Selective reporting (reporting bias)	Low risk	According to the aims, adequately reported variables
Other bias	Unclear risk	It is unclear whether or not there was risk of other bias due to the lack of detail in the methods.

Killen 2000

Methods	Randomised, single-blind (participants), cross-over trial
Participants	<p>Inclusion criteria: COPD predominantly related to smoking; considered for symptomatic oxygen therapy; had stairs at home and found that ascending these produced breathlessness; desaturation to below 90% on ascent of 22 steps</p> <p>Exclusion criteria: history of ischaemic heart disease, left ventricular failure or other cause of reduced mobility such as severe arthritis; already on long-term oxygen or fulfilling the criteria for long-term oxygen therapy</p> <p>Setting: non-laboratory</p> <p>18 participants 8 male and 10 female Mean age: 67.5 (IQR 60.5 to 74.3)</p> <p>Median FEV₁ (L): 0.53 (IQR 0.45 to 0.76) Median DLCO (% predicted): 44 (IQR 28 to 64) Mean baseline SaO₂ (%): median 94 (IQR 91 to 95) Mean baseline PaO₂ (kPa): no data available</p> <p>Mean baseline PaCO₂ (kPa): no data available Mean breathlessness at baseline: no data available</p>
Interventions	Oxygen (2 L/min) vs compressed air 5 minutes before and/or 5 minutes after ascending 22 steps. During the 5 minutes before and after these ascents. they breathed from a cylinder of compressed air or oxygen, delivered at 2 L/min via face mask.
Outcomes	Breathlessness measured by 100-mm VAS ("not at all breathless" to "extremely breathless") HRQOL - no data available Treatment preference - no data available Adverse effects - no data available

Killen 2000 (Continued)

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	"The order of the ascents was determined by randomisation within a Latin square."
Allocation concealment (selection bias)	Unclear risk	Information on concealment of allocation not available
Blinding of participants and personnel (performance bias) All outcomes	High risk	"During the five minutes before and after these ascents they breathed from a cylinder of either compressed air or oxygen, delivered at 2 l/min via a face mask, in a single blind manner."
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	As above
Incomplete outcome data (attrition bias) All outcomes	Low risk	All participants completed
Selective reporting (reporting bias)	Low risk	According to the aims, adequately reported variables
Other bias	Unclear risk	It is unclear whether or not there was risk of other bias due to the lack of detail in the methods

Knebel 2000

Methods	Randomised, double-blinded (participants, investigators), cross-over trial
Participants	<p>Inclusion criteria: adults with obstructive lung disease due to AAT deficiency; FEV₁ < 70% of predicted; FEV₁/FVC < 0.70</p> <p>Exclusion criteria: FEV₁ < 1 L; hospitalisation in preceding 3 weeks; conditions prohibiting or limiting exercise; current use of oxygen; inability to understand English</p> <p>Setting: laboratory</p> <p>33 participants (31 completed)</p> <p>22 male and 13 female</p> <p>Mean age: 47 (SD 7)</p> <p>Mean FEV₁ (% predicted): 48 (SD 13)</p> <p>Mean TLC (% predicted): 105 (SD 14)</p> <p>Mean baseline SaO₂ (%): 97.1% (SD 1.7)</p> <p>Mean baseline PaO₂ (kPa): no data available</p> <p>Mean baseline PaCO₂ (kPa): no data available</p> <p>Mean breathlessness at baseline: 0.5 (SD 0.9)</p>

Knebel 2000 (Continued)

Interventions	Oxygen (4 L/min) vs compressed air (4 L/min) during 6MWT. Both gases were delivered by nasal cannula, using the minimum dose shown to improve exercise performance, 4 L/min.
Outcomes	Breathlessness measured by 10-cm horizontal VAS (0 = "No shortness of breath" and 10 = "Shortness of breath as bad as it can be") HRQOL – no data available Treatment preference - no data available Adverse effects - number of dropouts

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	"A table of random numbers identified the order of administration."
Allocation concealment (selection bias)	Unclear risk	Information not available
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"The tanks were covered so neither the patient nor the researcher knew which gas was being used."
Blinding of outcome assessment (detection bias) All outcomes	Low risk	See above.
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	"Two patients were unable to complete all of the walks because of unrelated problems." Data on remaining participants were analysed in the study.
Selective reporting (reporting bias)	Low risk	According to the aims, adequately reported variables
Other bias	Unclear risk	It is unclear whether or not there was risk of other bias due to the lack of detail in the methods.

Kurihara 1989

Methods	Randomised, single-blind, cross-over trial
Participants	Inclusion criteria: "COPD" Exclusion criteria: none mentioned (Cochrane translation) Setting: laboratory 14 participants 11 male and 3 female Mean age: 62 (SD 10.2) Mean FEV ₁ (L): 0.67 (SD 0.23) Mean FVC (% predicted): 58.3 (SD 6.2) Mean baseline SaO ₂ (%): no data available

Kurihara 1989 (Continued)

Mean baseline PaO₂ (kPa): 9.2 (SD 1.2)

Mean baseline PaCO₂ (kPa): no data available

Mean breathlessness at baseline: no data available

Interventions	Breathlessness by modified Borg scale and distance walked on treadmill
Outcomes	Breathlessness measured by modified Borg (mBorg, 1 "none" to 10 "extremely severe") HRQOL – no data available Treatment preference - no data available Adverse effects - no data available
Notes	Worked from translation sheet from Japanese as opposed to from the entire article

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Described as randomised; no other information available
Allocation concealment (selection bias)	Unclear risk	Information not available
Blinding of participants and personnel (performance bias) All outcomes	High risk	Single-blind study; no other information available
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	See above.
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Information not available
Selective reporting (reporting bias)	Unclear risk	Information not available
Other bias	Unclear risk	It is unclear whether or not there was risk of other bias due to the lack of detail in the methods.

Laude 2006

Methods	Randomised, controlled, double-blinded (participants, investigators), cross-over trial
Participants	<p>Inclusion criteria: COPD confirmed by FEV₁ < 80% predicted, FEV₁/FVC < 70%, a change in FEV₁ < 200 mL or < 15% of baseline, 15 minutes after inhaled salbutamol, dyspnoea on exertion defined by a Borg score after exercise ≥ 3 and freedom from exacerbations in the 6 weeks preceding the study</p> <p>Exclusion criteria: required domiciliary oxygen therapy; had resting hypercapnia, anaemia or a significant cardiac diagnosis</p> <p>Setting: laboratory</p> <p>Method of recruitment of participants: outpatients recruited from respiratory clinics at the 3 hospitals</p>

Laude 2006 (Continued)

82 participants (76 participants analysed in air group, 78 participants in oxygen group)

Gender not specified

Mean age: 69.7 (range 46 to 84)

Mean FEV₁ (L): 1.1 (SD 0.4)

Mean FVC (L): 2.6 (SD, 0.8)

Mean baseline SaO₂ (%): 94 (SD 2.3)

Mean baseline PaO₂ (kPa): no data available

Mean baseline PaCO₂ (kPa): no data available

Mean breathlessness at baseline: 1.8 (SD 1.1)

Interventions	Oxygen (0.28) vs medical air during treadmill exercise, delivered through a non-rebreathing mask and demand valve system connected to a portable cylinder
Outcomes	Breathlessness measured by modified Borg (mBorg 0-10, 0 "none" to 10 "maximal") and 100-mm VAS HRQOL – no data available Treatment preference - no data available Adverse effects - number of dropouts

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	"Independent randomisation of the gas mixtures was done using a Latin square design."
Allocation concealment (selection bias)	Unclear risk	Information not available
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"In all tests, the investigator carried the gas cylinder walking beside the patient and gave no encouragement. Patients were instructed not to speak while breathing the gas mixtures and for 2 min afterwards to avoid unblinding...would have un-blinded both patient and investigator." "Cylinders were coded as to content to maintain blinding."
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	As above
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Intention-to-treat population reported; specific details of how missing data were handled in the analysis not available
Selective reporting (reporting bias)	Low risk	Important outcomes reported
Other bias	Unclear risk	Unclear how participants were selected, data reported only for participants completing the study (7 withdrawals in total)

Leach 1992

Methods	Randomised, double-blinded (participants, investigators), cross-over trial
---------	--

Leach 1992 (Continued)

Participants	<p>Inclusion criteria: severely reduced exercise tolerance secondary to chronic respiratory disease; no previous experience of exercise testing</p> <p>Exclusion criteria: angina pectoris; impaired left ventricle heart function; locomotor disability</p> <p>Setting: laboratory</p> <p>20 participants Gender not specified Mean age: 63.4 Mean FEV₁ (L): 0.74 (0.3) Mean FVC (L): 1.94 (0.5)</p> <p>Mean baseline SaO₂ (%): no data available Mean baseline PaO₂ (kPa): 8.7 (SD 2.3)</p> <p>Mean baseline PaCO₂ (kPa): 5.6 (SD 1.2) Mean breathlessness at baseline: no data available</p>
Interventions	Oxygen (2, 4 or 6 L/min) vs compressed air (4 L/min) delivered through face mask during 6MWT and endurance shuttle walk test (walk as far as possible and stop when unable to go farther)
Outcomes	Breathlessness measured by 10-cm VAS ("not at all breathless" to "extremely breathless") HRQOL – no data available Treatment preference - no data available Adverse effects - no data available
Notes	
Risk of bias	
Bias	Authors' judgement Support for judgement
Random sequence generation (selection bias)	Unclear risk "The order of the four tests in which the gas was carried by the patient was randomised on each day."
Allocation concealment (selection bias)	Unclear risk Information not available
Blinding of participants and personnel (performance bias) All outcomes	High risk "The subject and the investigator were blinded to the flow rate and type of gas supplied, although in practice the investigator was frequently able to determine those patients having oxygen from the oxygen saturation shown by ear oximetry."
Blinding of outcome assessment (detection bias) All outcomes	High risk See above.
Incomplete outcome data (attrition bias) All outcomes	Low risk All participants completed
Selective reporting (reporting bias)	Low risk Adequately reported variables
Other bias	Unclear risk It is unclear whether or not there was risk of other bias due to the lack of detail in the methods

Lewis 2003

Methods	Randomised, single-blinded (participants), placebo-controlled, cross-over trial
Participants	<p>Inclusion criteria: moderate to severe COPD according to BTS criteria; significant self-reported breathlessness; on optimal treatment; no exacerbation of disease for > 4 weeks before study, with exacerbation defined as "a deterioration in respiratory symptoms requiring treatment with corticosteroids or antibiotics or both"</p> <p>Exclusion criteria: significant limiting or unstable comorbidities</p> <p>Setting: laboratory</p> <p>Method of recruitment of participants: Participants were identified through respiratory outpatient services at Green Lane Hospital, Auckland, New Zealand.</p> <p>22 participants (18 completed) 16 male and 2 female Mean age: 68.7 (SD 10.1) Mean FEV₁ (L): 0.91 (SD 0.4) Mean baseline SaO₂ (%): 94 (SD 1.6) Mean baseline PaO₂ (kPa): no data available</p> <p>Mean baseline PaCO₂ (kPa): no data available Mean breathlessness at baseline: 0.4 (SD 0.5)</p>
Interventions	Oxygen (2 L/min) vs air (2 L/min) before 6MWT, delivered via nasal cannulae
Outcomes	Breathlessness measured by modified Borg at end of exercise (mBorg 0-10, 0 "none" to 10 "maximal") HRQOL - no data available Treatment preference - no data available Adverse effects - number of dropouts
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Described as randomised; other information not available
Allocation concealment (selection bias)	Unclear risk	Information not available
Blinding of participants and personnel (performance bias) All outcomes	High risk	"...identical cylinders in a single-blind fashion. Due to the need for monitoring of oxygen saturation by the physiotherapist, double-blinding was not possible."
Blinding of outcome assessment (detection bias) All outcomes	High risk	As above
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Data reported only for participants completing the study (4 withdrawals)

Lewis 2003 (Continued)

Selective reporting (reporting bias)	Low risk	Important outcomes were reported.
Other bias	Unclear risk	It is unclear whether or not there was risk of other bias due to the lack of detail in the methods.

Maltais 2001

Methods	Randomised, double-blinded (participants, investigators), cross-over trial
Participants	<p>Inclusion criteria: "moderate to severe" COPD with diagnosis based on previous or current smoking history and PFTs (including spirometry, lung volume and CO diffusing capacity); "stable disease"</p> <p>Exclusion criteria: clinical cardiovascular, neurological or any condition that could alter the capacity to perform an exercise test according to medical history, physical exam, resting and exercise electrocardiogram and chest x-ray</p> <p>Setting: laboratory</p> <p>14 participants Gender not specified Mean age: 63 (SEM 3) Mean FEV₁ (L): 1.04 (SEM 0.1) Mean FVC (L): 2.64 (SEM 0.2)</p> <p>Mean baseline SaO₂ (%): no data available Mean baseline PaO₂ (kPa): 11.3 (SEM 0.5)</p> <p>Mean baseline PaCO₂ (kPa): 4.9 (SEM 0.3) Mean breathlessness at baseline: no data available</p>
Interventions	Room air vs oxygen (FiO ₂ 0.75) delivered through a mouthpiece during exercise testing
Outcomes	<p>Breathlessness measured by modified Borg (mBorg 0-10, 0 "none" to 10 "maximal")</p> <p>HRQOL – no data available</p> <p>Treatment preference - no data available</p> <p>Adverse effects - no data available</p>

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	"Patients performed two exercise tests in a random order"; other information not available
Allocation concealment (selection bias)	Unclear risk	Information not available
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"Patients and the physician supervising the exercise tests were blinded as to which inspiratory gas was used."
Blinding of outcome assessment (detection bias)	Low risk	See above.

Maltais 2001 (Continued)

All outcomes

Incomplete outcome data (attrition bias) All outcomes	Low risk	All participants completed the study.
Selective reporting (reporting bias)	Low risk	Adequately reported variables
Other bias	Unclear risk	It is unclear whether or not there was risk of other bias due to the lack of detail in the methods.

McDonald 1995

Methods	Randomised, double-blinded (participants, investigators), cross-over trial
Participants	<p>Inclusion criteria: stable severe COPD; resting PaO₂ > 60 mmHg; exertional breathlessness sufficient to interfere with daily activities; non-smoker; no exacerbations in preceding 3 months; maximal bronchodilator and/or corticosteroid therapy</p> <p>Exclusion criteria: symptomatic cardiac dysfunction; angina pectoris; locomotor disability</p> <p>Setting: non-laboratory</p> <p>33 participants (26 completed) 24 male and 2 female Mean age: 73 (SD 6) Mean FEV₁ (L): 0.9 (SD 0.4) Mean DLCO (mL/min/mmHg): 10.6 (SD 2.4) Mean baseline SaO₂ (%): 94 (SD 2.1) Mean baseline PaO₂ (kPa): 9.2 (SD 1.1)</p> <p>Mean baseline PaCO₂ (kPa): 5.5 (SD 0.4) Mean breathlessness at baseline: no data available</p>
Interventions	Oxygen vs compressed air delivered via nasal cannula at 4 L/min over long-term (successive 6-week periods during which participants were instructed to use portable gas cylinder during "any activity that would normally induce dyspnoea). Participants used portable gas cylinders fitted with a demand gas delivery system at a flow rate of 4 L/min intranasally.
Outcomes	Breathlessness measured by modified Borg at end of exercise (mBorg 0-10, 0 "none" to 10 "maximal") HRQOL – CRQ subdomains Treatment preference - no data available Adverse effects - number of dropouts
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Described as randomised; other information not available
Allocation concealment (selection bias)	Unclear risk	Information not available

McDonald 1995 (Continued)

Blinding of participants and personnel (performance bias) All outcomes	Low risk	"All gases were delivered from camouflaged cylinders, and both the patients and the investigator performing the tests were blinded to their contents."
Blinding of outcome assessment (detection bias) All outcomes	Low risk	As above
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Data reported only for participants completing the study (7 withdrawals)
Selective reporting (reporting bias)	Unclear risk	According to the aims, adequately reported variables
Other bias	Unclear risk	It is unclear whether or not there was risk of other bias due to the lack of detail in the methods.

McKeon 1988a

Methods	Randomised, double-blind (participant and investigator), cross-over trial
Participants	<p>Inclusion criteria: COPD with "significant disability with exertional breathlessness despite treatment with inhaled and oral bronchodilators"; FEV₁/FVC < 0.60 and TLC > 80% predicted with no significant change after bronchodilator; "stable condition"</p> <p>Exclusion criteria: not studied during exercise with history of angina, recent myocardial infarction, exercise-induced syncope or left ventricular failure</p> <p>Setting: laboratory</p> <p>Method of recruitment of participants: consecutive patients with COPD referred for assessment of need for oxygen therapy (Newcastle, Australia)</p> <p>20 participants 13 male and 7 female Mean age: 63.2 (SD 10) Mean FEV₁ (L): 0.79 (SD 0.3) Mean FVC (L): 2.30 (SD 0.7) Mean TLC (% predicted): 122 (SD 24) Mean RV (% predicted): 206 (SD 60) Mean DLCO (% predicted): 55 (SD 32)</p> <p>Mean baseline SaO₂ (%): 90 (SD 3) Mean baseline PaO₂ (kPa): 7.7 (SD 1.2)</p> <p>Mean baseline PaCO₂ (kPa): 5.9 (SD 1.2) Mean breathlessness at baseline: no data available</p>
Interventions	Compressed air vs oxygen via nasal cannula at 2.5 L/min during treadmill test
Outcomes	Breathlessness measured by 300-mm VAS HRQOL – no data available Treatment preference - no data available Adverse effects - no data available

McKeon 1988a (Continued)

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Described as randomised; information not available
Allocation concealment (selection bias)	Unclear risk	Information not available
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"Neither the patient nor the operator knew whether compressed air or supplemental oxygen had been given. Patients were told that both cylinders contained oxygen, but in different concentrations."
Blinding of outcome assessment (detection bias) All outcomes	Low risk	See above.
Incomplete outcome data (attrition bias) All outcomes	Low risk	All participants completed the study.
Selective reporting (reporting bias)	Low risk	According to the aims, adequately reported variables
Other bias	Unclear risk	It is unclear whether or not there was risk of other bias due to the lack of detail in the methods.

McKeon 1988b

Methods	Randomised, controlled, double-blind (participant and investigator), cross-over trial
Participants	<p>Inclusion criteria: COPD with "significant disability with exertional breathlessness despite treatment with inhaled and oral bronchodilators"; "stable condition" at the time of the study</p> <p>Exclusion criteria: not stated</p> <p>Setting: laboratory</p> <p>21 participants 11 women and 10 men Mean age: 62 (SD 9) Mean FEV₁ (L): 0.77 (SD 0.4) Mean FEV₁ (% predicted): 29 (SD 13) Mean FVC (L): 2.00 (SD 0.9) Mean FVC (% predicted): 58 (SD 20) Mean RV (L): 3.53 (SD 0.9) Mean TLC (L): 5.97 (SD 1.3)</p> <p>Mean baseline PaO₂ (kPa): 8.9 (SD 1.5) Mean breathlessness at baseline: no data available</p>
Interventions	Oxygen vs air delivered via nasal cannula at 4 L/min during treadmill test

McKeon 1988b (Continued)

Outcomes	Breathlessness measured by 300-mm VAS HRQOL – no data available Treatment preference - no data available Adverse effects - no data available
----------	---

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Described as randomised; no other information available
Allocation concealment (selection bias)	Unclear risk	Cylinders prepared by technician not involved in the study
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Identical cylinders used in the study
Blinding of outcome assessment (detection bias) All outcomes	Low risk	See above.
Incomplete outcome data (attrition bias) All outcomes	Low risk	All participants completed the study.
Selective reporting (reporting bias)	Low risk	According to the aims, adequately reported variables
Other bias	Unclear risk	It is unclear whether or not there was risk of other bias due to the lack of detail in the methods.

Miki 2012

Methods	Randomised, controlled, single-blinded (participants), cross-over trial
---------	---

Participants	<p>Inclusion criteria: COPD not stated as criterion, but study included patients with $FEV_1/FVC < 0.7$; clinical stability defined as no respiratory infection for ≥ 4 weeks before the exercise test; ability to tolerate exercise test for ≥ 4 minutes; and exercise limitation due primarily to exertional breathlessness during CPET (breathing discomfort alone or in conjunction with leg discomfort was the primary reason for stopping exercise)</p> <p>Exclusion criteria: significant pathology including neuromuscular, cardiac and/or peripheral vascular disease, malignancy or anaemia; qualified for LTOT before the exercise test</p> <p>Setting: laboratory (Japan)</p> <p>Method of recruitment of participants: no data</p> <p>35 participants</p> <p>All male</p>
--------------	--

Miki 2012 (Continued)

Mean age: 70.4 (SD 5.7)

Mean FEV₁ (L): 1.13 (SD 0.5)

Mean FEV₁ (% predicted): 43.1 (SD 17.4)

FEV₁/FVC: 0.46 (SD 13.5)

Mean baseline SaO₂ (%): no data available

Mean baseline PaO₂ (kPa): oxygen: 12.5 (SD 1.7); air: 10.2 (SD 1.3)

Mean baseline PaCO₂ (kPa): oxygen: 5.3 (SD 0.7); air: 5.1 (SD 0.7)

Mean breathlessness at baseline: oxygen: 0.1 (SD 0.2); air: 0.1 (SD 0.4)

Interventions	Oxygen (FiO ₂ 0.24) vs compressed air during 2 symptom-limited incremental exercise tests. Breathed through a mask attached to a low-resistance, 2-way, non-rebreathing valve from gas cylinders through 200-L Douglas bag
Outcomes	Breathlessness measured by modified Borg at iso-time (mBorg 0-10, 0 "none" to 10 "maximal") HRQOL - no data available Treatment preference - no data available Adverse effects - no data available

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Described as randomised; no other information available
Allocation concealment (selection bias)	Unclear risk	Information not available
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Single-blinded (participants): "Patients were blinded as to which oxygen concentration they were breathing."
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Information not available
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	All participants completed the study.
Selective reporting (reporting bias)	Low risk	Important outcomes were reported.
Other bias	Unclear risk	Unclear how participants were selected; blinding not optimal

Moore 2009

Methods	Randomised, controlled, double-blinded (participants and study personnel), cross-over trial
Participants	<p>Inclusion criteria: patients with a clinical diagnosis of COPD attending the respiratory laboratory for routine breathing tests</p> <p>Exclusion criteria: use of short-term bronchodilators within 4 hours; receipt of supplemental oxygen within 20 minutes</p> <p>Setting: tertiary hospital, not stated if single-centre, Melbourne, Australia</p> <p>Method of recruitment of participants: Patients with a clinical diagnosis of COPD attending the respiratory laboratory for routine breathing tests were invited to participate in the study.</p> <p>55 participants (51 completed)</p> <p>Male 40</p> <p>Mean age: 72.6 (SD 9.7)</p> <p>Mean FEV₁ (L): 1.40 (SD 0.78)</p> <p>Mean FEV₁ (% predicted): 54.7 (SD 24.9)</p> <p>Mean baseline SaO₂ (%): 95 (SD 3.2)</p> <p>Mean baseline PaO₂ (kPa): no data available</p> <p>Mean baseline PaCO₂ (kPa): no data available</p> <p>Mean breathlessness at baseline: no data available</p>
Interventions	Oxygen (44%) vs medical air via mouthpiece. Breathing circuit included a 100-L Douglas bag, a Hans Rudolf wide bore, non-rebreathing valve.
Outcomes	Breathlessness - modified Borg HRQOL – no data available Treatment preference - no data available Adverse effects - no data available

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Described as randomised; other information not available
Allocation concealment (selection bias)	Unclear risk	As above
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Double-blinded. "The study gases were delivered to the Douglas bag from cylinders of identical appearance."
Blinding of outcome assessment (detection bias) All outcomes	Low risk	As above
Incomplete outcome data (attrition bias)	Low risk	Data reported only for participants completing the study (1 withdrawal)

Moore 2009 (Continued)

All outcomes

Selective reporting (reporting bias)	Low risk	Important outcomes were reported.
Other bias	Unclear risk	It is unclear whether or not there was risk of other bias due to the lack of detail in the methods.

Moore 2011

Methods	Randomised, controlled, double-blinded (participants and study personnel), parallel-group trial
Participants	<p>Inclusion criteria: clinically stable ex-smokers with COPD on optimal medical treatment; $\text{PaO}_2 > 7.3$ kPa at rest breathing room air; and moderate to severe exertional breathlessness (mMRC Dyspnoea Scale grade > 2)</p> <p>Exclusion criteria: patients with $\text{PaCO}_2 > 6.0$ kPa undergoing repeat arterial blood gas analysis after breathing oxygen at 6 L/min, at rest, for 30 minutes; excluded if PaCO_2 increased by more than 0.7 kPa</p> <p>Setting: tertiary hospital, not stated if single centre, Melbourne, Australia</p> <p>Method of recruitment of participants: not described.</p> <p>143 participants (139 completed)</p> <p>Male 99 (69%), female 44</p> <p>Mean age: 71.8 (SD 9.8)</p> <p>Mean FEV_1 (L): 1.16 (SD 0.51)</p> <p>Mean baseline SaO_2 (%): no data available</p> <p>Mean baseline PaO_2 (kPa): oxygen: 9.7 (SD 8.2); air: 9.3 (SD 1.1)</p> <p>Mean baseline PaCO_2 (kPa): oxygen: 5.4 (SD 0.7); air: 5.4 (SD 0.6)</p> <p>Mean breathlessness at baseline: oxygen: 17.6 (SD 5.2); air: 17.5 (SD 4.9)</p>
Interventions	Domiciliary oxygen vs air at 6 L/min with 2-week run-in; 12 weeks ambulatory treatment. Delivered through nasal prongs by gas cylinders
Outcomes	Breathlessness measured by CRQ dyspnoea domain HRQOL – CRQ total score, CRQ subdomains Treatment preference - no data available Adverse effects - number of dropouts
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"Assigned them randomly to cylinder air or cylinder oxygen using a computer-generated sequence and concealed allocation"
Allocation concealment (selection bias)	Low risk	As above

Moore 2011 (Continued)

Blinding of participants and personnel (performance bias) All outcomes	Low risk	"Study personnel and participants were blinded to group allocation. Cylinders were of identical appearance."
Blinding of outcome assessment (detection bias) All outcomes	Low risk	As above
Incomplete outcome data (attrition bias) All outcomes	Low risk	Data reported only for participants completing the study. The number of dropouts was equal in both groups (4 dropouts in total).
Selective reporting (reporting bias)	Low risk	According to the aims, adequately reported variables.
Other bias	Unclear risk	It is unclear whether or not there was risk of other bias due to the lack of detail in the methods.

Nandi 2003

Methods	Randomised, controlled, double-blinded (participants, investigators), cross-over trial	
Participants	<p>Inclusion criteria: FEV₁ < 60% predicted with < 15% reversibility to inhaled salbutamol; smoking history > 20 pack-years; exertional desaturation ≥ 4% on pulse oximetry during submaximal exertion (corridor walking)</p> <p>Exclusion criteria: any other complicating medical condition</p> <p>Setting: laboratory</p> <p>34 participants 18 male and 16 female Mean age: 68 (SD 6.0) Mean FEV₁ (L): 0.88 (SD 0.3) Mean baseline SaO₂ (%): 92 (SD 5.2) Mean baseline PaO₂ (kPa): 7.7 (SD 1.5)</p> <p>Mean baseline PaCO₂ (kPa): no data available Mean breathlessness at baseline: no data available</p>	
Interventions	Oxygen (28% at 4 L/min) vs compressed air (4 L/min) before exercise. Gas was supplied via constant performance masks, which supply 28% oxygen at the mouth over a wide range of flow rates and breathing patterns.	
Outcomes	Breathlessness measured by 100-mm VAS ("not breathless at all" to "the most breathless I have ever been") HRQOL – no data available Treatment preference - no data available Adverse effects - no data available	
Notes		
Risk of bias		
Bias	Authors' judgement	Support for judgement

Nandi 2003 (Continued)

Random sequence generation (selection bias)	Unclear risk	Described as randomised; information not available
Allocation concealment (selection bias)	Unclear risk	Information not available
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"...neither the patient nor the test supervisor was aware of the gas mixture being used, or of oxygen saturation levels which were recorded by another observer."
Blinding of outcome assessment (detection bias) All outcomes	Low risk	See above.
Incomplete outcome data (attrition bias) All outcomes	Low risk	All participants completed the study.
Selective reporting (reporting bias)	Low risk	According to the aims, adequately reported variables
Other bias	Unclear risk	It is unclear whether or not there was risk of other bias due to the lack of detail in the methods.

Nonoyama 2007

Methods	Randomised, controlled, double-blind (participants and study personnel), N-of-1 cross-over trial
Participants	<p>Inclusion criteria: patients with a diagnosis of COPD (5) with dyspnoea limiting daily activities, and with desaturation $\leq 88\%$ for 2 continuous minutes during a room-air 6-minute-walk test</p> <p>Exclusion criteria: 18 years or younger; those who met criteria for mortality reduction with long-term oxygen; those who received oxygen for palliative care or isolated nocturnal hypoxaemia; those unable to complete the questionnaires or provide informed consent</p> <p>Setting: home setting, Canada, outpatient</p> <p>Method of recruitment of participants: Participants were recruited from local respiratory rehabilitation programmes, respirology offices and home oxygen companies. The study co-ordinator (MLN) was responsible for recruitment and enrolment.</p> <p>38 participants (27 completed)</p> <p>Male 17 (63%)</p> <p>Mean age: 69 (SD 10)</p> <p>Lung function: no data available</p> <p>Mean baseline SaO₂ (%): no data available</p> <p>Mean baseline PaO₂ (kPa): no data available</p> <p>Mean baseline PaO₂ (kPa): no data available</p> <p>Mean breathlessness at baseline: CRQ dyspnoea score 3.7 (SD 1.1)</p>
Interventions	Ambulatory oxygen (1-3 L/min) vs medical air (2 L/min) at exertion delivered through nasal prongs by gas cylinders. Participants undertook 3 pairs of 2-week treatment periods. Range of each period 9 to 21

Nonoyama 2007 (Continued)

days. Threshold for an adequate number of activities was set at 2 activities (chosen by participant), for a minimum duration of 1 hour a day

Outcomes	Breathlessness measured by modified Borg (mBorg 0-10, 0 "none" to 10 "maximal") and CRQ dyspnoea domain 1-7 p HRQOL – CRQ subdomains, SGRQ total score Treatment preference - no data available Adverse effects - number of dropouts
----------	---

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"Random allocation was centralized; method of randomization was computer generated."
Allocation concealment (selection bias)	Low risk	"Allocation was concealed and the order was randomly assigned to one of the eight possible orders of the pairs of placebo and active periods."
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"Oxygen and placebo gas in identical E-sized and D-sized cylinders, marked only with a code number indicating their contents." "Patients and the outcome assessors were blind to the gas mixture provided. This was achieved by an independent person ordering, receiving and preparing the appropriate cylinders and concentrator (oxygen or placebo)."
Blinding of outcome assessment (detection bias) All outcomes	Low risk	See above.
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Data reported only for participants completing the study. High proportion of dropouts (11 dropouts in total)
Selective reporting (reporting bias)	Low risk	According to the aims, adequately reported variables
Other bias	Unclear risk	It is unclear whether or not there was risk of other bias due to the lack of detail in the methods.

O'Donnell 1997

Methods	Randomised, placebo-controlled, double-blinded (participant, investigators), cross-over trial
Participants	<p>Inclusion criteria: advanced chronic airway limitation ($FEV_1 < 60\%$ predicted); mild hypoxaemia; did not meet criteria for LTOT; referred to an exercise programme because sedentary; poor exercise tolerance; severe activity-related breathlessness</p> <p>Exclusion criteria: clinical evidence of significant cardiovascular disease, other pulmonary disease (including cor pulmonale) or other disorders that could contribute to breathlessness or exercise limitation</p> <p>Method for recruiting participants: Patients were referred to an exercise programme because they were sedentary/had poor exercise tolerance/experienced severe activity-related breathlessness.</p> <p>Setting: laboratory (Ontario, Canada)</p>

O'Donnell 1997 (Continued)

11 participants
 7 male and 4 female
 Mean age: 68 (SEM 2)
 Mean FEV₁ (% predicted): 0.97 (SEM 0.13)
 Mean FVC (% predicted): 2.3 (SEM 0.3)
 Mean TLC (% predicted): 7.0 (SEM 0.5)
 Mean RV (% predicted): 4.4 (SEM 0.3)
 Mean DLCO (mL/min/mmHg): 8.8 (SEM 1.1)

 Mean baseline SaO₂ (%): no data available
 Mean baseline PaO₂ (kPa): 9.9 (SEM 0.4)

 Mean baseline PaCO₂ (kPa): 5.5 (SEM 0.3)
 Mean breathlessness at baseline: 5.1 (SD 0.3)

Interventions	Oxygen (FiO ₂ 0.60) vs room air (FiO ₂ 0.21) on endurance cycle exercise test. Participants breathed through a mouthpiece and a low-resistance 2-way nonrebreathing valve with the nose clipped.
Outcomes	Breathlessness measured by modified Borg (mBorg 0-10, 0 "none" to 10 "maximal") HRQOL – no data available Treatment preference - no data available Adverse effects - no data available
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Described as randomised; no other information available
Allocation concealment (selection bias)	Unclear risk	Information not available
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"Subjects were blinded with respect to the oxygen concentration being breathed, as was the investigator evaluating subjective responses and performing the data analysis; Identical breathing apparatus used in the study."
Blinding of outcome assessment (detection bias) All outcomes	Low risk	As above
Incomplete outcome data (attrition bias) All outcomes	Low risk	All participants completed the study.
Selective reporting (reporting bias)	Low risk	According to the aims, adequately reported variables
Other bias	Unclear risk	It is unclear whether or not there was risk of other bias due to the lack of detail in the methods.

O'Driscoll 2011

Methods	Randomised, controlled, single-blind (participants), cross-over trial
Participants	<p>Inclusion criteria: patients over 50 years of age; smoking history > 20 pack-years; severe COPD with FEV₁ < 40% of predicted; oxygen saturation at rest ≥ 93% breathing room air at time of recruitment to study and on arrival on study day; breathless during modest exercise such as climbing 1 or 2 flights of stairs; stable for 4 weeks (not requiring oral steroids or antibiotics or both); able to give informed consent; able to undertake a simple step test on an 18-cm exercise step</p> <p>Exclusion criteria: unable or unwilling to give informed consent; unable or unwilling to undertake step test for any reason; complicating comorbid conditions (e.g. arthritis of knees) that might interfere with exercise test or with the conduct of the trial; and already using oxygen at home</p> <p>Setting: laboratory</p> <p>Method of recruitment of participants: Participants were recruited from the chest clinic or pulmonary rehabilitation service at Salford Royal University Hospital (UK).</p> <p>39 participants (34 completed)</p> <p>Male 24 (71%)</p> <p>Mean age 69 (SD 10)</p> <p>Mean FEV₁ (L) 0.80 (SD 0.2)</p> <p>Mean FEV₁ (% predicted) 31.4 (SD 5.5)</p> <p>Mean baseline PaO₂ (kPa): no data available</p> <p>Mean baseline PaCO₂ (kPa): no data available</p> <p>Mean breathlessness at baseline: 1.5 (SD 1.1)</p>
Interventions	Oxygen (4 L/min) vs air (4 L/min) via face mask after exercise (step test)
Outcomes	Breathlessness measured by modified Borg after step exercise test (mBorg 0-10, 0 "none" to 10 "maximal") HRQOL – no data available Treatment preference - categorical, at study end Adverse effects - no data available
Notes	
Risk of bias	
Bias	Authors' judgement Support for judgement
Random sequence generation (selection bias)	Low risk "Patients received one of the following interventions in random order based on the patient drawing four pieces of folded paper sequentially from an opaque container prior to the commencement of the exercise task."
Allocation concealment (selection bias)	Low risk See above.
Blinding of participants and personnel (performance bias) All outcomes	High risk Single-blind. "The oxygen and air treatments were single blind. To achieve blinding for these two interventions, the investigator connected a concealed air or oxygen cylinder delivering a gas flow of 4 litres per minute to the same face mask for each patient."
Blinding of outcome assessment (detection bias)	Unclear risk "Breathlessness on a Borg scale reported by participants; unclear whether they were aware of saturation during, at the end of exercise and during the re-

O'Driscoll 2011 (Continued)

All outcomes		covery period. The SpO ₂ and pulse rate were recorded in the patient's trial record sheet each minute during the exercise test, at the end of exercise and every minute during recovery until 5 minutes after finishing the exercise task."
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Data reported only for participants completing the study. 5 dropouts in total
Selective reporting (reporting bias)	Low risk	According to the aims, adequately reported variables
Other bias	Unclear risk	It is unclear whether or not there was risk of other bias due to the lack of detail in the methods.

Oliveira 2012a

Methods	Randomised, controlled, double-blind (participants and study personnel), cross-over trial
Participants	<p>Inclusion criteria: stable, non-hypercapnic (PaCO₂ < 45 mmHg at rest) COPD; > 20 pack-years of smoking; resting saturation > 90% and PaO₂ > 8 kPa; no evidence of polycythemia or cor pulmonale. No participant had qualified previously for ambulatory O₂ therapy.</p> <p>Exclusion criteria: recent exacerbation (within 1 month); treatment with oral corticosteroids; previous or current heart disease</p> <p>Method of recruitment of participants: no data</p> <p>Setting: laboratory (UK)</p> <p>8 participants with COPD classified as "desaturators" with a drop in SaO₂ and end-exercise values < 88%</p> <p>All male</p> <p>Mean age: 66.7 (SD 7.9)</p> <p>Mean FEV₁ (L): 1.31 (SD 0.48)</p> <p>Mean FEV₁ (% predicted): 45.1 (SD 12.9)</p> <p>Mean baseline SaO₂ (%): no data available</p> <p>Mean baseline PaO₂ (kPa): 8.5 (SD 1.1)</p> <p>Mean baseline PaCO₂ (kPa): 5.2 (SD 0.6)</p> <p>Mean breathlessness at baseline: no data available</p>
Interventions	Hyperoxia (FiO ₂ 0.40) vs normoxia (FiO ₂ 0.21) during incremental cycle tests; 2 tests ≥ 2 days apart. Gas delivered through a facial mask connected to a low-resistance Douglas bag
Outcomes	<p>Breathlessness measured by modified Borg at peak exercise (mBorg 0-10, 0 "none" to 10 "maximal")</p> <p>HRQOL – no data available</p> <p>Treatment preference - no data available</p> <p>Adverse effects - no data available</p>
Notes	

Risk of bias

Oliveira 2012a (Continued)

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	"Test sequence was randomised"; other Information not available
Allocation concealment (selection bias)	Unclear risk	Information not available
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"Both patients and accompanying physician were blinded to the mixture being breathed."
Blinding of outcome assessment (detection bias) All outcomes	Low risk	See above.
Incomplete outcome data (attrition bias) All outcomes	Low risk	All participants
Selective reporting (reporting bias)	Low risk	According to the aims, adequately reported variables
Other bias	Low risk	We judged that this trial appeared to be free of other sources of bias.

Oliveira 2012b

Methods	Randomised, controlled, double-blind (participants and study personnel), cross-over trial
Participants	<p>Inclusion criteria: stable, non-hypercapnic ($\text{PaCO}_2 < 45$ mmHg at rest) COPD; > 20 pack-years of smoking; resting saturation $> 90\%$ and $\text{PaO}_2 > 8$ kPa; no evidence of polycythemia or cor pulmonale. No participant had qualified previously for ambulatory O_2 therapy.</p> <p>Exclusion criteria: recent exacerbation (within 1 month), treatment with oral corticosteroids; previous or current heart disease</p> <p>Method of recruitment of participants: no data</p> <p>Setting: laboratory (UK)</p> <p>12 participants classified as "non-desaturators"</p> <p>All male</p> <p>Mean age: 60.3 (SD 5.9)</p> <p>Mean FEV_1 (L): 1.56 (SD 0.41)</p> <p>Mean FEV_1 (% predicted): 48.6 (SD 10.8)</p> <p>Mean baseline SaO_2 (%): no data available</p> <p>Mean baseline PaO_2 (kPa): 10.0 (SD 1.3)</p> <p>Mean baseline PaCO_2 (kPa): 4.7 (SD 0.6)</p> <p>Mean breathlessness at baseline: no data available</p>

Oliveira 2012b (Continued)

Interventions	Hyperoxia (FiO ₂ 0.40) vs normoxia (FiO ₂ 0.21) during incremental cycle tests; 2 tests ≥ 2 days apart. Gas delivered through facial mask connected to a low-resistance Douglas bag
Outcomes	Breathlessness measured by modified Borg at peak exercise (mBorg 0-10, 0 "none" to 10 "maximal") HRQOL – no data available Treatment preference - no data available Adverse effects - no data available
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	"Test sequence was randomised"; other Information not available
Allocation concealment (selection bias)	Unclear risk	Information not available
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"Both patients and accompanying physician were blinded to the mixture being breathed."
Blinding of outcome assessment (detection bias) All outcomes	Low risk	See above.
Incomplete outcome data (attrition bias) All outcomes	Low risk	All participants completed the study.
Selective reporting (reporting bias)	Low risk	According to the aims, adequately reported variables
Other bias	Low risk	We judged that this trial appeared to be free of other sources of bias.

Ringbaek 2013

Methods	Randomised, controlled, single-blinded (participants), parallel-group trial
Participants	<p>Inclusion criteria: stable COPD (FEV₁ < 80% of predicted and FEV₁/FVC < 0.70); breathlessness with MRC score 3–5; motivated for PR; oxygen saturation at rest > 90%; desaturation > 4% and < 90% during ISWT or ESWT</p> <p>Exclusion criteria: LTOT; significant musculoskeletal, cardiac or cognitive problems</p> <p>Method of recruitment of participants: Of 391 participants starting PR at the centre between March 2007 and March 2011, 163 eligible patients were encouraged to participate in this trial.</p> <p>Setting: ambulatory rehabilitation training (Denmark)</p> <p>45 participants</p> <p>Male 53.3%</p>

Ringbaek 2013 (Continued)

Mean age: 69.0 (SD 8.7)

Mean FEV₁ (% predicted): 31.8 (SD 13.8)

Mean baseline SaO₂ (%): 94 (SD 2.0)

Mean baseline PaO₂ (kPa): no data available

Mean baseline PaCO₂ (kPa): no data available

Mean breathlessness at baseline, MRC (min-max): oxygen 4.7 (3-5); controls 4.3 (3-5)

Interventions	Ambulatory oxygen (2 L/min) vs control (no ambulatory oxygen); oxygen delivered through a 2.3-kg portable oxygen concentrator. Participants were asked to use oxygen during supervised and unsupervised exercise (≥ 30 min/d) for up to 33 weeks.
Outcomes	Breathlessness measured by modified Borg after 7 weeks of training (mBorg 0-10, 0 "none" to 10 "maximal")
	HRQL – SGRQ total score
	Treatment preference - no data available
	Adverse effects - COPD exacerbations, all hospital admissions, deaths at study end

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"Were randomised to either ambulatory oxygen (AO) (n=22) or to a control group (n=23) in blocks of 6, using sealed envelopes." Other information not available
Allocation concealment (selection bias)	Low risk	Sealed envelopes
Blinding of participants and personnel (performance bias) All outcomes	High risk	"without use of a sham concentrator"
Blinding of outcome assessment (detection bias) All outcomes	High risk	See above.
Incomplete outcome data (attrition bias) All outcomes	High risk	Data reported only for participants completing the study. High and unequal proportion of dropouts (7 missing for intervention; only 1 for comparison). "Baseline and week 7 only in the analysis"
Selective reporting (reporting bias)	Low risk	Have protocol and IPD data
Other bias	Low risk	We judged that this trial appeared to be free of other sources of bias.

Rooyackers 1997

Methods	Randomised, controlled, unblinded, parallel-group trial
---------	---

Rooyackers 1997 (Continued)

Participants

Inclusion criteria: hypoxaemia ($\text{SaO}_2 < 90\%$) at maximal exercise; increase in alveolar-arterial difference in oxygen tension ≥ 2 kPa from rest to maximal incremental exercise test; former smoker

Exclusion criteria: resting $\text{PaO}_2 < 64$ mmHg; mean nocturnal $\text{SaO}_2 < 90\%$; mean pulmonary artery pressure > 25 mmHg measured at rest by Doppler echocardiography; neuromuscular or cardiovascular disease; any medication change during the study

Setting: laboratory

Method for recruitment of participants: Patients were referred to the hospital for PR (Dekkerswald, Netherlands).

24 participants in total

12 participants allocated to 10 weeks of general exercise training breathing room air; "air group"

12 participants allocated to supplemental oxygen (4 L/min) during general exercise training for 10 weeks; "oxygen group"

10 male and 2 female in each group

See [Rooyackers 1997a](#); [Rooyackers 1997b](#) for baseline characteristics and resting pulmonary function data.

Interventions

See [Rooyackers 1997a](#); [Rooyackers 1997b](#).

Outcomes

HRQOL – CRQ total score (higher = better QoL), CRQ subdomains

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	No information available
Allocation concealment (selection bias)	Unclear risk	"The patients were randomly allocated either to general exercise training while breathing room air or while breathing at a flow rate of 4 L·min ⁻¹ through a dual-prong nasal cannula." No other information available
Blinding of participants and personnel (performance bias) All outcomes	High risk	While performing the tests: "During oxygen breathing, it was not possible to measure $\dot{V}\text{O}_2$, and blood gas analysis was not performed."
Blinding of outcome assessment (detection bias) All outcomes	High risk	As above
Incomplete outcome data (attrition bias) All outcomes	Low risk	All participants completed the study.
Selective reporting (reporting bias)	Low risk	According to the aims, adequately reported variables
Other bias	Unclear risk	It is unclear whether or not there was risk of other bias due to the lack of detail in the methods.

Rooyackers 1997a

Methods	Randomised, controlled, unblinded, parallel-group trial
Participants	<p>Inclusion criteria: hypoxaemia ($\text{SaO}_2 < 90\%$) at maximal exercise; increase in alveolar-arterial difference in oxygen tension ≥ 2 kPa from rest to maximal incremental exercise test; former smoker</p> <p>Exclusion criteria: resting $\text{PaO}_2 < 64$ mmHg; mean nocturnal $\text{SaO}_2 < 90\%$, mean pulmonary artery pressure > 25 mmHg measured at rest by Doppler echocardiography; neuromuscular or cardiovascular disease; any medication change during the study</p> <p>Setting: laboratory</p> <p>Method for recruitment of participants: Patients were referred to the hospital for PR (Dekkerswald, Netherlands).</p> <p>12 participants allocated to 10 weeks of general exercise training breathing room air; "air group" 10 male and 2 female Mean age: 59 (SD 13) Mean FEV_1 (L): 1.2 (SD 0.5) Mean TLC (% predicted): 114 (SD 20) Mean baseline SaO_2 (%): no data available Mean baseline PaO_2 (kPa): 10.5 (SD 1.1)</p> <p>Mean baseline PaCO_2 (kPa): 5.0 (SD 0.8) Mean breathlessness at baseline: no data available</p>
Interventions	Room air (RA) vs oxygen (4 L/min) during maximal incremental cycle exercise test, single-stage exercise test and 6MWT
Outcomes	<p>Breathlessness measured by modified Borg (mBorg 0-10, 0 "none" to 10 "maximal")</p> <p>HRQOL – CRQ total score (higher = better QoL), CRQ subdomains</p> <p>Treatment preference - no data available</p> <p>Adverse effects - no data available</p>
Notes	In both groups, the work rate was not allowed to exceed the level at which SaO_2 fell below 90%.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	No information available
Allocation concealment (selection bias)	Unclear risk	"The patients were randomly allocated either to general exercise training while breathing room air or while breathing supplemental oxygen at a flow rate of 4 L·min ⁻¹ through a dual-prong nasal cannula." No other information available
Blinding of participants and personnel (performance bias) All outcomes	High risk	While performing the tests: "During oxygen breathing, it was not possible to measure $\dot{V}\text{O}_2$, and blood gas analysis was not performed."
Blinding of outcome assessment (detection bias) All outcomes	High risk	As above

Rooyackers 1997a (Continued)

Incomplete outcome data (attrition bias) All outcomes	Low risk	All participants completed the study.
Selective reporting (reporting bias)	Low risk	According to the aims, adequately reported variables
Other bias	Unclear risk	It is unclear whether or not there was risk of other bias due to the lack of detail in the methods.

Rooyackers 1997b

Methods	Randomised, controlled, unblinded, parallel-group trial	
Participants	<p>Inclusion criteria: hypoxaemia (SaO₂ < 90%) at maximal exercise and increase in alveolar-arterial difference in oxygen tension ≥ 2 kPa from rest to maximal exercise during maximal incremental exercise; former smoker; no medication changes during the study</p> <p>Exclusion criteria: resting PaO₂ < 64 mmHg, mean nocturnal SaO₂ < 90%, mean pulmonary artery pressure > 25 mmHg measured at rest by Doppler echocardiography; neuromuscular or cardiovascular disorder</p> <p>Setting: laboratory</p> <p>Method for recruitment of participants: Patients were referred to the hospital for PR (Dekkerswald, Netherlands).</p> <p>12 participants allocated to supplemental oxygen (4 L/min) during general exercise training for 10 weeks; "oxygen group" 10 male and 2 female Mean age: 63 (SD 5) Mean FEV₁ (L): 0.9 (SD 0.3) Mean TLC (% predicted): 109 (SD 17)</p> <p>Mean baseline SaO₂ (%): no data available Mean baseline PaO₂ (kPa): 10.2 (SD 1.6)</p> <p>Mean baseline PaCO₂ (kPa): 5.1 (SD 1.1) Mean breathlessness at baseline: no data available</p>	
Interventions	Oxygen (4 L/min) vs room air (RA) during maximal incremental cycle exercise test, single-stage exercise test and 6MWT	
Outcomes	Breathlessness measured by modified Borg (mBorg 0-10, 0 "none" to 10 "maximal") HRQOL – CRQ total score (higher = better QoL), CRQ subdomains Treatment preference - no data available Adverse effects - no data available	
Notes		
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	No information available

Rooyackers 1997b (Continued)

Allocation concealment (selection bias)	Unclear risk	"The patients were randomly allocated either to general exercise training while breathing room air or while breathing supplemental oxygen at a flow rate of 4 L·min ⁻¹ through a dual-prong nasal cannula." No other information available
Blinding of participants and personnel (performance bias) All outcomes	High risk	While performing the tests: "During oxygen breathing, it was not possible to measure V'O ₂ , and blood gas analysis was not performed."
Blinding of outcome assessment (detection bias) All outcomes	High risk	As above
Incomplete outcome data (attrition bias) All outcomes	Low risk	All participants completed the study.
Selective reporting (reporting bias)	Low risk	According to the aims, adequately reported variables
Other bias	Unclear risk	It is unclear whether or not there was risk of other bias due to the lack of detail in the methods.

Scorsone 2010

Methods	Randomised, controlled, double-blind (participants and study personnel), parallel-group trial
Participants	<p>Inclusion criteria: clinical diagnosis of COPD and airflow obstruction confirmed by spirometry; although not stated as criteria, all participants had a history of smoking and were under regular bronchodilator treatment</p> <p>Exclusion criteria: not stated</p> <p>Method of recruitment of participants: not stated</p> <p>Setting: pulmonary rehab programme (Italy)</p> <p>20 participants</p> <p>Male 14 (70%)</p> <p>Mean age: air 68 (SD 7); oxygen 67 (SD 9)</p> <p>Mean FEV₁/FVC: air 47 (SD 12); oxygen 51 (SD 8)</p> <p>Mean FEV₁ (% predicted): air 50 (SD 12); oxygen 47 (SD 10)</p> <p>Mean baseline SaO₂ (%): no data available</p> <p>Mean baseline PaO₂ (kPa): oxygen 9.9 (SD 1.0); air 10.2 (SD 1.2)</p> <p>Mean baseline PaCO₂ (kPa): oxygen 5.3 (SD 0.4); air 5.1 (SD 0.4)</p> <p>Mean breathlessness at baseline: oxygen 7 (SD 3); air 7 (SD 3)</p>
Interventions	Supplemental oxygen (FiO ₂ 0.4) vs air delivered by mouthpiece from a Douglas bag; an 8-week, 3 times a week training programme conducted on a calibrated cycle ergometer

Scorsone 2010 (Continued)

Outcomes	Breathlessness measured by modified Borg at isotime of exertion (mBorg 0-10, 0 "none" to 10 "maximal") HRQOL – no data available Treatment preference - no data available Adverse effects - no data available
----------	--

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Described as randomised; other information not available
Allocation concealment (selection bias)	Unclear risk	Information not available
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"PaO ₂ saturation was continuously monitored by a technician blind to the design of the study." ".. according to a randomised, double-blind design in which subjects breathed humidified air, supplemental O ₂ 40%, or heliox 60:40 at random through a mouthpiece from a Douglas bag"
Blinding of outcome assessment (detection bias) All outcomes	Low risk	See above.
Incomplete outcome data (attrition bias) All outcomes	Low risk	All participants completed the study.
Selective reporting (reporting bias)	Low risk	According to the aims, adequately reported variables
Other bias	Unclear risk	It is unclear whether or not there was risk of other bias due to the lack of detail in the methods.

Somfay 2001

Methods	Randomised, controlled, single-blind (participants), cross-over trial
Participants	<p>Inclusion criteria: severe COPD (FEV₁ < 40% predicted); no more than mildly hypoxaemic (O₂ sat at rest > 92% and during exercise > 88%); none had previously qualified for home oxygen</p> <p>Exclusion criteria: clinically manifest cor pulmonale; severe cardiovascular comorbidity or other disease that might contribute to breathlessness or exercise limitation</p> <p>10 participants</p> <p>Setting: laboratory 6 male and 4 female</p> <p>Mean age: 67 (SD 7) Mean FEV₁ (L): 0.92 (SD 0.43) Mean FVC (% predicted): 76 (SD 15)</p>

Somfay 2001 (Continued)

Mean TLC (L): 7.3 (SD 1.5)
Mean RV (L): 4.3 (SD 1.3)
Mean baseline SaO₂ (%): 96 (SD 0.8)
Mean baseline PaO₂ (kPa): no data available

Mean baseline PaCO₂ (kPa): no data available
Mean breathlessness at baseline: no data available

Interventions	Oxygen (30%, 50%, 75% or 100%) vs compressed air during constant work rate test at 75% of peak work rate. Compressed air and oxygen were blended from gas cylinders into a 200-L meteorological balloon to be inspired during exercise. Participants breathed through a mouthpiece attached to a low-resistance, 2-way, nonrebreathing valve.
Outcomes	Breathlessness measured by modified Borg (mBorg 0-10, 0 "none" to 10 "maximal") HRQOL - no data available Treatment preference - no data available Adverse effects - no data available

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Described as randomised; no other information available
Allocation concealment (selection bias)	Unclear risk	Information not available
Blinding of participants and personnel (performance bias) All outcomes	High risk	Single-blind study; "Subjects were blinded with respect to the oxygen concentration they breathed."
Blinding of outcome assessment (detection bias) All outcomes	High risk	As above
Incomplete outcome data (attrition bias) All outcomes	Low risk	All participants completed the study.
Selective reporting (reporting bias)	Low risk	Adequately reported variables
Other bias	Unclear risk	It is unclear whether or not there was risk of other bias due to the lack of detail in the methods.

Spielmanns 2014

Methods	Randomised, controlled, parallel-group trial
Participants	Inclusion criteria: diagnosis of COPD and FEV ₁ /FVC < 0.7 after bronchodilation; FEV ₁ < 80% of predicted; stable clinical condition for ≥ 4 weeks; normoxaemia (SaO ₂ > 90%) at rest and during exercise without oxygen supply

Spielmanns 2014 (Continued)

Exclusion criteria: comorbidity that could interfere with training, such as ischaemic cardiac disease, musculoskeletal problems and psychological disorders

Method of recruitment of participants: invited to take part when referred by general practitioners to the St Remigius Hospital, in Leverkusen, Germany, from September 2009 to March 2011

Setting: 1 outpatient pulmonary rehabilitation programme (St Remigius Hospital, Leverkusen, Germany)

85 participants with COPD (39 withdrawals; 18 oxygen group; 21 medical air)

No data on sex

Mean age: oxygen 65 (SD 8.7); air 64 (SD 8.4)

Mean FEV₁ (L): oxygen 1.2 (SD 0.5); air 1.5 (SD 0.4)

Mean FEV₁ (% predicted): oxygen 44 (SD 10); air 43 (SD 12)

Mean baseline SaO₂ (%): > 90% according to inclusion criteria

Mean baseline PaO₂ (kPa): no data available

Mean baseline PaCO₂ (kPa): no data available

Mean breathlessness at baseline: no data available

Interventions	Oxygen (4 L/min) vs compressed air (4 L/min) during training cycle tests delivered via nasal cannula. Duration of treatment: training sessions throughout 24 weeks; 3 sessions/wk
Outcomes	Breathlessness - no data available HRQL - SF-36; subscores with range from 0 to 100; measured at baseline and after 12 weeks Treatment preference - no data available Adverse effects - number of dropouts
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"The randomization scheme was generated using the website Randomization.com using a computer-generated list of random numbers to assign subjects to either the oxygen or compressed air group."
Allocation concealment (selection bias)	Low risk	As above
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not sufficient information. "Tests were conducted by an unblinded coach; the subjects and investigator were not aware of the group to which they belonged until the end of all experiments."
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	As above
Incomplete outcome data (attrition bias) All outcomes	High risk	Data reported only for participants completing the study. High proportion of dropouts (39 in total; 18 from oxygen group; 21 from medical air)

Spielmanns 2014 (Continued)

Selective reporting (re-reporting bias)	Low risk	Adequately reported variables
Other bias	Low risk	We judged that this trial appeared to be free of other sources of bias.

Swinburn 1984

Methods	Randomised, single-blinded (participants), cross-over trial
Participants	Inclusion criteria: advanced obstructive airways disease but in "stable clinical state" Exclusion criteria: not stated Setting: laboratory 5 participants Mean age: 65 (range 53-72) Mean FEV ₁ (L): 0.8 (SD 0.2) Mean FVC (L): 1.8 (SD 0.4) Mean baseline SaO ₂ (%): 93.2 (SD 0.8) Mean baseline PaO ₂ (kPa): no data available Mean baseline PaCO ₂ (kPa): no data available Mean breathlessness at baseline: no data available
Interventions	Room air (RA) vs oxygen (60%) delivered through a mouthpiece during incremental cycle exercise test
Outcomes	Breathlessness measured by 10-cm VAS scale ("minimum" to "maximum") HRQOL - no data available Treatment preference - no data available Adverse effects - no data available

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Described as randomised; other information not available
Allocation concealment (selection bias)	Unclear risk	Information not available
Blinding of participants and personnel (performance bias) All outcomes	High risk	Single-blind. "The patients were informed that the purpose of the study was to investigate the effects of different gas mixtures on exercise performance."
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Information not available

Swinburn 1984 (Continued)

Incomplete outcome data (attrition bias) All outcomes	Low risk	All participants completed the study.
Selective reporting (reporting bias)	Low risk	Adequately reported outcomes
Other bias	Unclear risk	It is unclear whether or not there was risk of other bias due to the lack of detail in the methods.

Voduc 2010

Methods	Randomised, controlled, double-blind (participants and study personnel), cross-over trial
Participants	<p>Inclusion criteria: clinically stable COPD; accepted into the pulmonary rehabilitation programme; clinical diagnosis of COPD; FEV₁/FVC < 0.7; FEV₁ < 70% of predicted</p> <p>Exclusion criteria: hypoxia (SaO₂ < 89%) at rest or during ambulation; non-pulmonary conditions that could affect exercise performance (e.g. angina, congestive heart failure, severe musculoskeletal problems) or lung diseases other than COPD (e.g. asthma, bronchiectasis); and recent COPD exacerbations or changes in regular medications within the past 2 months</p> <p>Method of recruitment of participants: All patients were initially referred for pulmonary rehabilitation for improvement of exertional dyspnoea that was not satisfactorily managed by pharmacological therapy alone.</p> <p>Setting: 1 rehabilitation programme (Ottawa, Ontario, Canada)</p> <p>24 participants (16 completed)</p> <p>Male 14 (70%)</p> <p>Mean age: 65.9 (SD 6.6)</p> <p>Mean FEV₁/FVC: 41.5 (SD 13.8)</p> <p>Mean baseline SaO₂ (%): 97 (SD 1.9)</p> <p>Mean baseline PaO₂ (kPa): no data available</p> <p>Mean baseline PaCO₂ (kPa): no data available</p> <p>Mean breathlessness at baseline: no data available</p>
Interventions	Oxygen (FiO ₂ 0.5) vs room air delivered through a mask connected to a Douglas bag during 2 symptom-limited constant-load exercise tests
Outcomes	<p>Breathlessness measured by modified Borg at isotime (mBorg 0-10, 0 "none" to 10 "maximal")</p> <p>HRQOL – no data available</p> <p>Treatment preference - no data available</p> <p>Adverse effects - number of dropouts</p>

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Voduc 2010 (Continued)

Random sequence generation (selection bias)	Unclear risk	"The order of testing was randomised."
Allocation concealment (selection bias)	Unclear risk	No information provided
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"Both the patient and the supervising physician were blinded to the nature of the inhaled gas."
Blinding of outcome assessment (detection bias) All outcomes	Low risk	See above.
Incomplete outcome data (attrition bias) All outcomes	High risk	Included only participants who completed the whole 12-week training period. In total, 8 dropouts. They could have had a different acute response to oxygen.
Selective reporting (reporting bias)	Low risk	Adequately reported variables
Other bias	Unclear risk	It is unclear whether or not there was risk of other bias due to the lack of detail in the methods.

Wadell 2001

Methods	Randomised, controlled, single-blind (participants), parallel-group trial
Participants	<p>Inclusion criteria: age < 75 years; stopped smoking ≥ 6 months before entering the study; PaO₂ > 60 mmHg at rest; hypoxaemia (SaO₂ ≤ 92%) during exercise (6MWT test); FEV₁ < 70% of predicted; no infection in the 3 weeks preceding study enrolment; no change in medical treatment in the month preceding enrolment</p> <p>Exclusion criteria: past or present major illness, such as cardiac, orthopaedic or neurological disease that might have interfered with exercise performance</p> <p>Setting: laboratory</p> <p>Method of recruiting participants: recruited from previously diagnosed outpatients who had been treated at the Department of Respiratory Medicine and Allergy at the University Hospital in Umea, Sweden</p> <p>22 participants (20 completed) 10 in air group and 10 in oxygen group Median age: air 69 (60 to 72); oxygen 65 (52 to 73) Median FEV₁ (% predicted): air 51.6 (24 to 66); oxygen 39.3 (23 to 59) Median baseline SaO₂ (%): air 95 (91 to 97); oxygen 95 (93-97) Median baseline PaO₂ (kPa): air 9.3 (7.9 to 11.4); oxygen 9.5 (8.6 to 11.6)</p> <p>Median baseline PaCO₂ (kPa): 9.3 (7.9 to 11.4); oxygen 5.4 (4.5 to 5.8) Median breathlessness at baseline: 1.5 (0 to 3)</p>
Interventions	Oxygen (5 L/min) vs air (5 L/min) delivered through a dual-prong nasal cannula during 6MWT on treadmill - baseline effect of 2 interventions, as complete study involved training over an 8-week period
Outcomes	Breathlessness measured by modified Borg (mBorg 0-10, 0 "none" to 10 "maximal")

Wadell 2001 (Continued)

HRQOL – no data available
Treatment preference - no data available
Adverse effects - no data available

Notes The treadmill was stopped if participants rated 7/17 (respectively) or more on Borg scales and/or if participants' SaO₂ fell below 90%.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"The included patients were randomly allocated (randomization by blocks, men and women were randomised separately) to train either with air (AG) or with oxygen (OG)."
Allocation concealment (selection bias)	Unclear risk	Not enough information available to determine how order of treatment group assignment was concealed from investigators
Blinding of participants and personnel (performance bias) All outcomes	High risk	"Single-blind study"; "SaO ₂ and heart rate were continuously monitored with a pulse oximeter using a finger probe or a forehead probe."
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Outcomes self-reported by participant
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Data reported only for participants completing the study (2 withdrawals)
Selective reporting (reporting bias)	Low risk	Adequately reported variables
Other bias	Low risk	We judged that this trial appeared to be free of other sources of bias.

Woodcock 1981

Methods	Randomised, double-blind (participants and study personnel), cross-over trial
Participants	<p>Inclusion criteria: fixed airways obstruction; "moderate or severe breathlessness on exertion" (method of defining not stated); normal or low PaO₂</p> <p>Exclusion criteria: none stated</p> <p>Setting: laboratory</p> <p>10 participants 9 male and 1 female Mean age: 62 (range 43-70) Mean FEV₁ (L): 0.71 (SD 0.29) Mean FVC (L): 2.65 (SD 1.04) Mean baseline SaO₂ (%): no data available Mean baseline PaO₂ (kPa): 9.6 (SD 1.5)</p> <p>Mean baseline PaCO₂ (kPa): 4.6 (SD 0.6)</p>

Woodcock 1981 (Continued)

Mean breathlessness at baseline: 4 (SD 0.94)

Interventions	Oxygen (described as 100% delivered at 4 L/min) vs compressed air (4 L/min) during treadmill test and 6MWT
Outcomes	Breathlessness measured by 10-cm VAS HRQOL – no data available Treatment preference - no data available Adverse effects - no data available
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Described as randomised; other information not available
Allocation concealment (selection bias)	Unclear risk	Information not available
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"Compressed air versus oxygen delivered via coded unmarked cylinders"
Blinding of outcome assessment (detection bias) All outcomes	Low risk	See above.
Incomplete outcome data (attrition bias) All outcomes	Low risk	All participants completed the study.
Selective reporting (reporting bias)	Low risk	Adequately reported outcomes
Other bias	Low risk	We judged that this trial appeared to be free of other sources of bias.

6MWT: six-minute walk test; AAT: alpha-1 antitrypsin; ABG: arterial blood gas; ATS: American Thoracic Society; BTS: British Thoracic Society; CA: compressed air; CAD: coronary artery disease; CO: carbon monoxide; COPD: chronic obstructive pulmonary disease; CPET: cardiopulmonary exercise testing; CRQ: Chronic Respiratory Questionnaire; DLCO: diffusing capacity of the lung for carbon monoxide; ESWT: endurance shuttle walk test; FEV₁: forced expiratory volume in one second; FiO₂: fraction of inspired oxygen; FVC: forced vital capacity; HR: heart rate; HRQOL: health-related quality of life; IPD: individual patient data; IQR: interquartile range; ISWT: incremental shuttle walk test; LTOT: long-term oxygen relief therapy; PaCO₂: partial pressure of carbon dioxide in arterial blood; PaO₂: partial pressure of oxygen in arterial blood; PCO₂: partial pressure of carbon dioxide; PFT: pulmonary function test; PO₂: partial pressure of oxygen; PR: pulmonary rehabilitation; RA: room air; RV: right ventricle; SaO₂: oxygen saturation; SBOT: short-burst oxygen therapy; SD: standard deviation; SEM: standard error of the mean; SF-36: Short Form 36; SGRQ: St George's Respiratory Quotient; SMD: standard mean difference; SpO₂: oxygen saturation; TLC: total lung capacity; TR: tricuspid regurgitation; VAS: visual analogue scale; VC: vital capacity.

Characteristics of excluded studies [ordered by study ID]

Study	Reason for exclusion
Alexandre 2010	Other - abstract, no suitable outcome

Study	Reason for exclusion
Arnold 2008	Other - abstract
Balkissoon 2006	Other - review of another manuscript
Bjørgen 2009	No suitable outcome
Bradley 1978	No suitable outcome
Bye 1985	No suitable outcome
Campbell 2013	Patients already receiving home oxygen
Caspersen 2012	Other - abstract
Criner 1987	No suitable outcome
Cuvelier 2002	Patients already receiving home oxygen
Edvardsen 2007	Patients already receiving home oxygen
Evans 1986	No suitable outcome
Fraser 2013	Patients already receiving home oxygen
Fujimoto 2002	No suitable outcome
Garrod 2000	Patients already receiving home oxygen
Gosselin 2004	No suitable outcome
Heinzelmann 2011	Other - abstract
Helgerud 2010	No suitable outcome
King 1973	Mean PaO ₂ < 55 mmHg
King 2012	Other - abstract
Lane 1987	Not a randomised controlled trial
Leggett 1977	Mean PaO ₂ < 55 mmHg No suitable outcome
Lellouche 2011	Other - abstract
Light 1989	No suitable outcome
Liss 1988	Patients already receiving home oxygen
Lock 1992	Mean PaO ₂ < 55 mmHg No placebo or control arm
Maldonado 2014	Mean PaO ₂ < 55 mmHg; study performed at Bogota altitude

Study	Reason for exclusion
Mannix 1992	No suitable outcome
Marques-Magallanes 1998	Mean PaO ₂ < 55 mmHg No suitable outcome
Matsuzawa	Japanese with no capacity for translation
Nasilowski 2008	Patients already receiving home oxygen
Nguyen 2008	Intervention not oxygen vs medical air
Nosedá 1997	Intervention not oxygen vs medical air
O'Donnell 2001	Patients already receiving home oxygen
O'Driscoll 2003	Other - editorial
O'Driscoll 2007	No suitable outcome
O'Neill 2006	No suitable outcome
Ouyang 2006	Intervention not oxygen vs medical air
Palwai 2010	Patients already receiving home oxygen
Peters 2006	Intervention not oxygen vs medical air
Raimondi 1970	No suitable outcome
Roberts 1996	Mean PaO ₂ < 55 mmHg Patients already receiving home oxygen
Sandland 2008a	Patients hypoxic at rest
Sandland 2008b	Patients already receiving home oxygen or PRN oxygen No dyspnoea outcome
Sforza 2010	Other - abstract
Stein 1982	No suitable outcome
Stevenson 2004	No suitable outcome
Swinburn 1991	Mean PaO ₂ < 55 mmHg
Vyas 1971	No suitable outcome
Waterhouse 1983	No suitable outcome
Wedzicha 2006	Other - editorial
Womble 2010	Other - abstract
Womble 2012	No suitable outcome

PaO₂: partial pressure of oxygen in arterial blood; PRN: pro re nata, or 'as needed'.

DATA AND ANALYSES

Comparison 1. Oxygen versus air

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Breathlessness - all trials	32		Std. Mean Difference (Random, 95% CI)	-0.31 [-0.43, -0.20]
2 Breathlessness - subgroup analysis - short-burst oxygen vs not	32		SMD (Random, 95% CI)	Subtotals only
2.1 Studies using short-burst oxygen	4		SMD (Random, 95% CI)	-0.03 [-0.28, 0.22]
2.2 Studies not using short-burst oxygen	28		SMD (Random, 95% CI)	-0.36 [-0.48, -0.24]
3 Breathlessness - subgroup analysis - exertional desaturation vs not	31		SMD (Random, 95% CI)	Subtotals only
3.1 Studies with exertional desaturation	16		SMD (Random, 95% CI)	-0.28 [-0.39, -0.16]
3.2 Studies with no exertional desaturation	15		SMD (Random, 95% CI)	-0.47 [-0.69, -0.24]
4 Breathlessness - subgroup analysis - mean PaO ₂ < 9.3 kPa vs higher	32		SMD (Random, 95% CI)	Subtotals only
4.1 Studies with mean PaO ₂ < 9.3 kPa	7		SMD (Random, 95% CI)	-0.28 [-0.48, -0.07]
4.2 Studies with mean PaO ₂ ≥ 9.3 kPa	25		SMD (Random, 95% CI)	-0.33 [-0.47, -0.20]
5 Breathlessness - subgroup analysis - measured during exercise test vs not	32		SMD (Random, 95% CI)	-0.31 [-0.43, -0.20]
5.1 Studies measuring during exercise test	30		SMD (Random, 95% CI)	-0.34 [-0.46, -0.22]
5.2 Studies not measuring during exercise test	2		SMD (Random, 95% CI)	-0.13 [-0.37, 0.11]
6 Breathlessness - subgroup analysis - laboratory vs other	32		SMD (Random, 95% CI)	-0.31 [-0.43, -0.20]
6.1 Studies in laboratory setting	25		SMD (Random, 95% CI)	-0.37 [-0.52, -0.22]
6.2 Studies in non-laboratory setting	7		SMD (Random, 95% CI)	-0.23 [-0.36, -0.09]
7 Breathlessness - subgroup analysis - short-term vs long-term (training) effect of oxygen	32		Std. Mean Difference (Random, 95% CI)	-0.31 [-0.43, -0.20]
7.1 Studies of short-term effect of oxygen	29		Std. Mean Difference (Random, 95% CI)	-0.34 [-0.46, -0.22]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
7.2 Studies of long-term effect of oxygen	3		Std. Mean Difference (Random, 95% CI)	-0.09 [-0.37, 0.19]
8 Breathlessness - subgroup analysis - mean oxygen dose > 2 L/min vs lower	30		Std. Mean Difference (Random, 95% CI)	-0.31 [-0.43, -0.18]
8.1 Studies with a mean oxygen dose > 2 L/min	26		Std. Mean Difference (Random, 95% CI)	-0.35 [-0.49, -0.21]
8.2 Studies with a mean oxygen dose ≤ 2 L/min	5		Std. Mean Difference (Random, 95% CI)	-0.19 [-0.39, 0.01]
9 Breathlessness - sensitivity analysis - excluding measurements at peak exertion	26		Std. Mean Difference (Random, 95% CI)	-0.37 [-0.50, -0.24]
10 Breathlessness - sensitivity analysis - excluding studies with high risk of bias	25		SMD (Random, 95% CI)	-0.30 [-0.41, -0.20]
11 Breathlessness - sensitivity analysis - excluding outliers	26		Std. Mean Difference (Random, 95% CI)	-0.27 [-0.35, -0.18]
12 Breathlessness - sensitivity analysis - post hoc - excluding short-burst and outliers	22		SMD (Random, 95% CI)	-0.30 [-0.39, -0.20]
13 Health-related quality of life - all trials	5		Std. Mean Difference (Random, 95% CI)	0.12 [-0.04, 0.28]

Analysis 1.1. Comparison 1 Oxygen versus air, Outcome 1 Breathlessness - all trials.

Study or subgroup	Oxygen N	Air N	Std. Mean Difference (SE)	Std. Mean Difference IV, Random, 95% CI	Weight	Std. Mean Difference IV, Random, 95% CI
Abernethy 2010	0	0	-0.1 (0.172)		5.84%	-0.08[-0.41,0.26]
Bruni 2012a	0	0	-1.4 (0.48)		1.26%	-1.4[-2.34,-0.46]
Bruni 2012b	0	0	-1.8 (0.7)		0.63%	-1.77[-3.14,-0.4]
Davidson 1988	0	0	-0.2 (0.35)		2.18%	-0.22[-0.91,0.47]
Dean 1992	0	0	-1.4 (0.56)		0.95%	-1.4[-2.5,-0.3]
Eaton 2002	0	0	-0.4 (0.15)		6.71%	-0.42[-0.71,-0.13]
Emtner 2003a	0	0	-0.3 (0.38)		1.9%	-0.33[-1.07,0.41]
Emtner 2003b	0	0	-0.5 (0.4)		1.74%	-0.5[-1.28,0.28]
Eves 2006	0	0	-0.2 (0.45)		1.42%	-0.17[-1.05,0.71]
Jolly 2001a	0	0	-1.3 (0.64)		0.74%	-1.31[-2.56,-0.06]
Jolly 2001b	0	0	-1.2 (0.56)		0.95%	-1.22[-2.32,-0.12]
Killen 2000	0	0	-0.2 (0.34)		2.29%	-0.25[-0.92,0.42]
Knebel 2000	0	0	-0.1 (0.26)		3.46%	-0.13[-0.64,0.38]
Kurihara 1989	0	0	-0.6 (0.24)		3.88%	-0.58[-1.05,-0.11]
Laude 2006	0	0	-0.4 (0.16)		6.3%	-0.44[-0.75,-0.13]
Lewis 2003	0	0	-0.1 (0.2)		4.92%	-0.1[-0.49,0.29]
McDonald 1995	0	0	-0.4 (0.29)		2.94%	-0.4[-0.97,0.17]

Favours oxygen

-1 -0.5 0 0.5 1

Favours air

Analysis 1.2. Comparison 1 Oxygen versus air, Outcome 2 Breathlessness - subgroup analysis - short-burst oxygen vs not.

Analysis 1.3. Comparison 1 Oxygen versus air, Outcome 3 Breathlessness - subgroup analysis - exertional desaturation vs not.

Analysis 1.4. Comparison 1 Oxygen versus air, Outcome 4 Breathlessness - subgroup analysis - mean PaO₂ < 9.3 kPa vs higher.

Analysis 1.5. Comparison 1 Oxygen versus air, Outcome 5 Breathlessness - subgroup analysis - measured during exercise test vs not.

Analysis 1.6. Comparison 1 Oxygen versus air, Outcome 6 Breathlessness - subgroup analysis - laboratory vs other.

Analysis 1.7. Comparison 1 Oxygen versus air, Outcome 7 Breathlessness - subgroup analysis - short-term vs long-term (training) effect of oxygen.

Analysis 1.8. Comparison 1 Oxygen versus air, Outcome 8 Breathlessness - subgroup analysis - mean oxygen dose > 2 L/min vs lower.

Analysis 1.9. Comparison 1 Oxygen versus air, Outcome 9 Breathlessness - sensitivity analysis - excluding measurements at peak exertion.

Analysis 1.10. Comparison 1 Oxygen versus air, Outcome 10 Breathlessness - sensitivity analysis - excluding studies with high risk of bias.

Analysis 1.11. Comparison 1 Oxygen versus air, Outcome 11 Breathlessness - sensitivity analysis - excluding outliers.

Study or subgroup	Oxygen	Air	Std. Mean Difference	Std. Mean Difference	Weight	Std. Mean Difference
	N	N	(SE)	IV, Random, 95% CI		IV, Random, 95% CI
Test for overall effect: $Z=5.91(P<0.0001)$						
Favours oxygen -1 -0.5 0 0.5 1 Favours air						

Analysis 1.12. Comparison 1 Oxygen versus air, Outcome 12 Breathlessness - sensitivity analysis - post hoc - excluding short-burst and outliers.

Study or subgroup	Oxygen	Air	SMD	Std. Mean Difference	Weight	Std. Mean Difference
	N	N	(SE)	IV, Random, 95% CI		IV, Random, 95% CI
Abernethy 2010	0	0	-0.1 (0.172)		7.74%	-0.08[-0.41,0.26]
Davidson 1988	1	1	-0.2 (0.35)		1.87%	-0.22[-0.91,0.47]
Eaton 2002	1	1	-0.4 (0.15)		10.21%	-0.42[-0.71,-0.13]
Emtner 2003a	1	1	-0.3 (0.38)		1.59%	-0.33[-1.07,0.41]
Emtner 2003b	1	1	-0.5 (0.4)		1.44%	-0.5[-1.28,0.28]
Eves 2006	1	1	-0.2 (0.45)		1.13%	-0.17[-1.05,0.71]
Knebel 2000	1	1	-0.1 (0.26)		3.4%	-0.13[-0.64,0.38]
Kurihara 1989	1	1	-0.6 (0.24)		3.99%	-0.58[-1.05,-0.11]
Laude 2006	1	1	-0.4 (0.16)		8.97%	-0.44[-0.75,-0.13]
McDonald 1995	1	1	-0.4 (0.29)		2.73%	-0.4[-0.97,0.17]
Miki 2012	0	0	-0.4 (0.18)		7.09%	-0.45[-0.8,-0.1]
Moore 2011	0	0	-0.2 (0.17)		7.93%	-0.18[-0.52,0.15]
Nonoyama 2007	0	0	-0.2 (0.12)		15.95%	-0.22[-0.46,0.02]
O'Donnell 1997	1	1	-0.4 (0.16)		8.97%	-0.44[-0.75,-0.13]
Oliveira 2012a	0	0	0.1 (0.29)		2.73%	0.05[-0.52,0.62]
Oliveira 2012b	0	0	0.1 (0.35)		1.87%	0.11[-0.58,0.8]
Ringbaek 2013	0	0	0.2 (0.336)		2.04%	0.22[-0.44,0.87]
Rooyackers 1997a	1	1	-0.3 (0.41)		1.37%	-0.26[-1.06,0.54]
Rooyackers 1997b	1	1	-0.3 (0.42)		1.3%	-0.33[-1.15,0.49]
Scorsone 2010	0	0	0 (0.447)		1.15%	0[-0.88,0.88]
Voduc 2010	0	0	-0.3 (0.26)		3.4%	-0.28[-0.79,0.23]
Woodcock 1981	1	1	-0.7 (0.27)		3.15%	-0.71[-1.24,-0.18]
Total (95% CI)					100%	-0.3[-0.39,-0.2]
Heterogeneity: $\tau^2=0$; $\chi^2=15.71$, $df=21(P=0.79)$; $I^2=0\%$						
Test for overall effect: $Z=6.22(P<0.0001)$						
Favours oxygen -4 -2 0 2 4 Favours air						

Analysis 1.13. Comparison 1 Oxygen versus air, Outcome 13 Health-related quality of life - all trials.

Study or subgroup	Oxygen	Air	Std. Mean Difference	Std. Mean Difference	Weight	Std. Mean Difference
	N	N	(SE)	IV, Random, 95% CI		IV, Random, 95% CI
Eaton 2002	0	0	0.2 (0.12)		46.78%	0.22[-0.02,0.46]
Moore 2011	0	0	0.1 (0.17)		23.34%	0.06[-0.27,0.4]
Nonoyama 2007	0	0	-0.1 (0.19)		18.66%	-0.09[-0.46,0.28]
Rooyackers 1997	0	0	0.1 (0.409)		4.03%	0.12[-0.68,0.92]
Spielmanns 2014	0	0	0.2 (0.306)		7.19%	0.15[-0.45,0.75]
Favours Oxygen -2 -1 0 1 2 Favours Air

ADDITIONAL TABLES

Table 1. Characteristics of included trials

Study	Design	Setting	Blinding	O ₂ delivery	O ₂ dose	Baseline PaO ₂ (kPa)	Baseline SaO ₂ (%)	Baseline dyspnoea	Breathlessness outcome measure	HRQOL outcome measure
Abernethy 2010	Parallel	Non-laboratory	Double-blind	NC	2 L/min	10.0 (SD 1.5)	NA	4.8 (SD 2.1)	NRS	10-cm VAS
Bruni 2012a	Cross-over	Laboratory	Double-blind	Mouth-piece	50%	9.5 (SD 1.2)	NA	NA	Modified Borg	-
Bruni 2012b	Cross-over	Laboratory	Double-blind	Mouth-piece	50%	10.3 (SD 0.9)	NA	NA	Modified Borg	-
Davidson 1988	Cross-over	Laboratory	Double-blind	NC or valve	4 L/min	8.6 (SE 0.3)	NA	NA	10-cm VAS	-
Dean 1992	Cross-over	Laboratory	Double-blind	Mouth-piece	40%	9.5 (SE 0.3)	NA	NA	Modified Borg	-
Dyer 2012	Parallel	Laboratory	Single-blind	NC	2-6 L/min	NA	94 (SD 2)	3 (SD 1)	CRQ dyspnoea	CRQ sub-domains
Eaton 2002	Cross-over	Non-laboratory	Double-blind	NC	4 L/min	9.2 (SD 1.0)	94 (SD 1.9)	0.7 (SD 1.0)	Modified Borg	CRQ total CRQ sub-domains SF-36
Eaton 2006	Parallel	Non-laboratory	Double-blind	NC	2 L/min	Oxygen: 9.6 (SD 1.3) Air: 10.1 (SD 1.7)	Oxygen: 95 (SD 1.9) Air: 95 (SD 1.6)	Oxygen: 17.8 (SD 5.0) Air: 17.5 (SD 4.2)	CRQ dyspnoea	CRQ total CRQ sub-domains SF-36
Emtner 2003a	Parallel	Laboratory	Double-blind	Mouth-piece	30%	9.8 (SD 0.8)	NA	5.8 (SD 1.8)	Modified Borg	CRQ total CRQ sub-domains SF-36

Table 1. Characteristics of included trials (Continued)

Emtner 2003b	Parallel	Laboratory	Dou- ble-blind	Mouth- piece	30%	10.0 (SD 1.2)	NA	6.3 (2.5)	Modified Borg	CRQ total CRQ sub- domains SF-36
Eves 2006	Crossover	Laboratory	Dou- ble-blind	Mouth- piece	40%	9.1 (SD 0.9)	NA	NA	Modified Borg	-
Haidl 2004	Parallel	Non-labora- tory	Unblinded	NC	2 L/min	Oxygen: 9.0 (SD 0.9) Controls: 8.7 (SD 0.8)	NA	Oxygen: 5.0 (SD 2.1) Controls: 5.0 (SD 1.5)	Modified Borg	-
Ishimine 1995	Crossover	Laboratory	Dou- ble-blind	Unknown	3 L/min	10.1 (SD 1.1)	NA	NA	Dyspnoea ques- tionnaire	-
Jolly 2001a	Crossover	Laboratory	Dou- ble-blind	NC	3 L/min	10.5 (SE 0.4)	95.8 (SE 0.46)	0.56 (SE 0.34)	Modified Borg	-
Jolly 2001b	Crossover	Laboratory	Dou- ble-blind	NC	3 L/min	9.9 (SE 0.3)	94.7 (SE 0.27)	1.27 (SE 0.43)	Modified Borg	-
Killen 2000	Crossover	Non-labora- tory	Single-blind	Mask	2 L/min	NA	Median 94 (IQR 91, 95)	NA	100-mm VAS	-
Knebel 2000	Crossover	Laboratory	Dou- ble-blind	NC	4 L/min	NA	97.1 (SD 1.7) (range 92-100)	0.5 (SD 0.9)	10-cm VAS	-
Kurihara 1989	Crossover	Laboratory	Single-blind	NC	3 L/min	9.2 (SD 1.2)	NA	NA	Modified Borg	-
Laude 2006	Crossover	Laboratory	Unclear	Mask/ valve	28%	NA	93.9 (SD 2.3)	VAS 24.2 (19.0) Borg 1.8 (1.1)	100-mm VAS modified Borg	-

Table 1. Characteristics of included trials (Continued)

Leach 1992	Crossover	Laboratory	Single-blind	Mask	2 L/min	8.7 (SD 2.3)	NA	NA	10-cm VAS	-
Lewis 2003	Crossover	Laboratory	Single-blind	NC	2 L/min	NA	94.4 (1.6)	0.4 (0.5)	Modified Borg	-
Maltais 2001	Crossover	Laboratory	Dou- ble-blind	Mouth- piece	75%	11.3 (SEM 0.5)	NA	NA	Modified Borg	-
McDonald 1995	Crossover	Non-labora- tory	Dou- ble-blind	NC	4 L/min	9.2 (SD 1.1) (range 7.7-10.9)	94 (SD 2.1)	NA	Modified Borg	CRQ subdo- mains
McKeon 1988a	Crossover	Laboratory	Dou- ble-blind	NC	2.5 L/min	7.7 (SD 1.2) (range 5.7-10.9)	90 (SD 3) (range 84-96)	NA	300-mm VAS	-
McKeon 1988b	Crossover	Laboratory	Dou- ble-blind	NC	4 L/min	8.9 (SD 1.5)	NA	NA	300-mm VAS	-
Miki 2012	Crossover	Laboratory	Single-blind	Mask/ valve	24%	10.1 (SD 1.3)	NA	Oxygen: 0.1 (SD 0.2) Air: 0.1 (SD 0.4)	Modified Borg	-
Moore 2009	Crossover	Laboratory	Dou- ble-blind	Mouth- piece	44%	NA	95 (SD 3.2)	NA	Modified Borg	-
Moore 2011	Parallel	Non-labora- tory	Dou- ble-blind	NC	6 L/min	9.5 (SD 1.1)	NA	Oxygen: 17.6 (SD 5.2) Air: 17.5 (SD 4.9)	CRQ dyspnoea	CRQ total CRQ sub- domains
Nandi 2003	Crossover	Laboratory	Dou- ble-blind	Mask	4 L/min	7.7 (SD 1.5)	91.9 (SD 5.2) (range 76 to 97)	NA	100-mm VAS	-
Nonoyama 2007	Crossover	Non-labora- tory	Dou- ble-blind	NC	1-3 L/min	NA	NA	3.7 (SD 1.1)	Modified Borg	CRQ sub- domains SQRQ to- tal

Table 1. Characteristics of included trials (Continued)

O'Donnell 1997	Crossover	Laboratory	Dou- ble-blind	Mouth- piece	60%	9.9 (SEM 0.4)	NA	5.1 (SD 0.3) ^a	Modified Borg	-
O'Driscoll 2011	Crossover	Laboratory	Single-blind	Mask	4 L/min	NA	95.0 (SD, 1.3)	1.5 (SD 1.1)	Modified Borg	-
Oliveira 2012a	Crossover	Laboratory	Dou- ble-blind	Mask	40%	8.5 (SD 1.1)	NA	NA	Modified Borg	-
Oliveira 2012b	Crossover	Laboratory	Dou- ble-blind	Mask	40%	10.0 (SD 1.3)	NA	NA	Modified Borg	-
Ringbaek 2013	Parallel	Non-labora- tory	Single-blind	NC	2 L/min	NA	93.6 (SD 2.0)	5.3 (SD 1.8)	Modified Borg	SQRQ to- tal
Rooyackers 1997a	Parallel	Laboratory	Unclear	NC	4 L/min	10.2 (SD 1.2)	NA	NA	Modified Borg	CRQ total CRQ sub- domains
Rooyackers 1997b	Parallel	Laboratory	Unclear	NC	4 L/min	9.5 (SD 2.0)	NA	NA	Modified Borg	CRQ total CRQ sub- domains
Scorsone 2010	Parallel	Laboratory	Dou- ble-blind	Mouth- piece	40%	Oxygen: 9.9 (SD 1.0) Air: 10.2 (SD 1.2)	NA	7 (SD 3)	Modified Borg	
Somfay 2001	Cross-over	Laboratory	Single-blind	Mouth- piece	30%	NA	95.7 (0.8)	NA	Modified Borg	-
Spielmanns 2014	Parallel	Laboratory	Unclear	NC	4 L/min	NA	> 90%	NA	-	SF-36 total
Swinburn 1984	Cross-over	Laboratory	Single-blind	Mouth- piece	60%	NA	93.2 (SD 0.8)	NA	10-cm VAS	-
Voduc 2010	Cross-over	Laboratory	Dou- ble-blind	Mask	50%	NA	97.1 (SD 1.9)	NA	Modified Borg	-
Wadell 2001	Parallel	Laboratory	Single-blind	NC	5 L/min	Median 9.3	Median 94.6	Median 1.5	Modified Borg	-

Table 1. Characteristics of included trials (Continued)

						(range 7.9-11.4)	(range 90.7-97.2)	(range 0-3)		
Woodcock 1981	Cross-over	Laboratory	Dou- ble-blind	NC	4 L/min	9.6 (SD 1.5)	NA	4 (SD 0.94) ^b	10-cm VAS	-

CRQ: Chronic Respiratory Questionnaire; NA: not available; NC: nasal cannula; SD: standard deviation; SE: standard error; SF-36: Short Form-36 questionnaire; VAS: visual analogue scale.

Data presented as mean (standard deviation (SD)) unless otherwise specified.

^aAs measured by dyspnoea index.

^bAs measured by MRC dyspnoea grade.

APPENDICES

Appendix 1. Sources and search methods for the Cochrane Airways Group Specialised Register (CAGR)

Electronic searches: core databases

Database	Frequency of search
CENTRAL (the Cochrane Library)	Monthly
MEDLINE (Ovid)	Weekly
Embase (Ovid)	Weekly
PsycINFO (Ovid)	Monthly
CINAHL (EBSCO)	Monthly
AMED (EBSCO)	Monthly

Handsearches: core respiratory conference abstracts

Conference	Years searched
American Academy of Allergy, Asthma and Immunology (AAAAI)	2001 onwards
American Thoracic Society (ATS)	2001 onwards
Asia Pacific Society of Respiriology (APSR)	2004 onwards
British Thoracic Society Winter Meeting (BTS)	2000 onwards
Chest Meeting	2003 onwards
European Respiratory Society (ERS)	1992, 1994, 2000 onwards
International Primary Care Respiratory Group Congress (IPCRG)	2002 onwards
Thoracic Society of Australia and New Zealand (TSANZ)	1999 onwards

MEDLINE search strategy used to identify trials for the CAGR

COPD search

1. Lung Diseases, Obstructive/
2. exp Pulmonary Disease, Chronic Obstructive/
3. emphysema\$.mp.
4. (chronic\$ adj3 bronchiti\$).mp.
5. (obstruct\$ adj3 (pulmonary or lung\$ or airway\$ or airflow\$ or bronch\$ or respirat\$)).mp.

6. COPD.mp.

7. COAD.mp.

8. COBD.mp.

9. AECB.mp.

10. or/1-9

Filter to identify RCTs

1. exp "clinical trial [publication type]"/

2. (randomized or randomised).ab,ti.

3. placebo.ab,ti.

4. dt.fs.

5. randomly.ab,ti.

6. trial.ab,ti.

7. groups.ab,ti.

8. or/1-7

9. Animals/

10. Humans/

11. 9 not (9 and 10)

12. 8 not 11

The MEDLINE strategy and RCT filter are adapted to identify trials in other electronic databases

Appendix 2. Airways Group Specialised Register search strategy

#1 MeSH DESCRIPTOR Pulmonary Disease, Chronic Obstructive Explode All

#2 MeSH DESCRIPTOR Bronchitis, Chronic

#3 (obstruct*) near3 (pulmonary or lung* or airway* or airflow* or bronch* or respirat*)

#4 COPD:MISC1

#5 (COPD OR COAD OR COBD):TI,AB,KW

#6 #1 OR #2 OR #3 OR #4 OR #5

#7 MeSH DESCRIPTOR Oxygen Inhalation Therapy

#8 MeSH DESCRIPTOR Oxygen

#9 oxygen*

#10 O2:ti,ab

#11 LTOT:ti,ab

#12 inhalation* NEXT therap*

#13 #7 or #8 or #9 or #10 or #11 or #12

#14 #6 and #13

Appendix 3. CENTRAL search strategy

Search 1: COPD + oxygen

#1 MESH DESCRIPTOR Pulmonary Disease, Chronic Obstructive EXPLODE ALL TREES

#2 ((obstruct*) near3 (pulmonary or lung* or airway* or airflow* or bronch* or respirat*)):TI,AB,KY

#3 (COPD OR COAD OR COBD):TI,AB,KY

#4 #1 OR #2 OR #3

#5 MESH DESCRIPTOR Oxygen Inhalation Therapy EXPLODE ALL TREES

#6 MESH DESCRIPTOR Oxygen EXPLODE ALL TREES

#7 oxygen*:TI,AB,KY

#8 O2:TI,AB

#9 LTOT:TI,AB,KY

#10 (inhalation* NEXT therap*):TI,AB,KY

#11 #5 OR #6 OR #7 OR #8 OR #9 OR #10

#12 #4 AND #11

Search 2: Dyspnoea + oxygen

#1 MESH DESCRIPTOR Dyspnea EXPLODE ALL TREES

#2 (dyspnoea* or dyspnoea*):TI,AB,KY

#3 breathless*:TI,AB,KY

#4 ((shortness* or difficult*) NEAR2 (breath*)):TI,AB,KY

#5 #1 OR #2 OR #3 OR #4

#6 MESH DESCRIPTOR Oxygen EXPLODE ALL TREES

#7 MESH DESCRIPTOR Oxygen Inhalation Therapy EXPLODE ALL TREES

#8 LTOT:TI,AB,KY

#9 ((inhalation* NEXT therap*)):TI,AB,KY

#10 ((oxygen*) NEAR (therap* or palliative* or inhal* or long-term*)):TI,AB,KY

#11

#6 OR #7 OR #8 OR #9 OR #10

#12

#5 AND #11

Appendix 4. MEDLINE search strategy

1. exp Dyspnea/
2. (dyspnoea\$ or dyspnoea\$).ti,ab.
3. breathless\$.ti,ab.
4. ((shortness\$ or difficult\$) adj2 breath\$).ti,ab.
5. or/1-4

6. Oxygen/ad, tu [Administration & Dosage, Therapeutic Use]
7. exp Oxygen Inhalation Therapy/
8. LTOT.ti,ab.
9. (inhalation\$ adj3 therap\$).ti,ab.
10. (oxygen\$ adj3 (therap\$ or palliative\$ or inhal\$ or long-term\$)).ti,ab.
11. or/6-10
12. 5 and 11
13. (controlled clinical trial or randomized controlled trial).pt.
14. (randomized or randomised).ab,ti.
15. placebo.ab,ti.
16. dt.fs.
17. randomly.ab,ti.
18. trial.ab,ti.
19. groups.ab,ti.
20. or/13-19
21. Animals/
22. Humans/
23. 21 not (21 and 22)
24. 20 not 23
25. 12 and 24

Appendix 5. Embase search strategy

1. exp dyspnoea/
2. (dyspnoea\$ or dyspnoea\$).ti,ab.
3. breathless\$.ti,ab.
4. ((shortness\$ or difficult\$) adj2 breath\$).ti,ab.
5. or/1-4
6. oxygen/ad, cm, dt, ih [Drug Administration, Drug Comparison, Drug Therapy, Inhalational Drug Administration]
7. oxygen therapy/
8. LTOT.ti,ab.
9. (inhalation\$ adj3 therap\$).ti,ab.
10. (oxygen\$ adj3 (therap\$ or palliative\$ or inhal\$ or long-term\$)).ti,ab.
11. or/6-10
12. 5 and 11
13. Randomized Controlled Trial/
14. randomization/

15. controlled clinical trial/
16. Double Blind Procedure/
17. Single Blind Procedure/
18. Crossover Procedure/
19. (clinical\$ adj3 trial\$).tw.
20. ((singl\$ or doubl\$ or trebl\$ or tripl\$) adj3 (mask\$ or blind\$ or method\$)).tw.
21. exp Placebo/
22. placebo\$.ti,ab.
23. random\$.ti,ab.
24. ((control\$ or prospectiv\$) adj3 (trial\$ or method\$ or stud\$)).tw.
25. (crossover\$ or cross-over\$).ti,ab.
26. or/13-25
27. exp animals/ or exp invertebrate/ or animal experiment/ or animal model/ or animal tissue/ or animal cell/ or nonhuman/
28. human/ or normal human/ or human cell/
29. 27 and 28
30. 27 not 29
31. 26 not 30
32. 12 and 31

WHAT'S NEW

Date	Event	Description
14 August 2017	Amended	Errors in three treatment effects in the abstract corrected.

HISTORY

Protocol first published: Issue 2, 2007

Review first published: Issue 6, 2011

Date	Event	Description
12 July 2016	New citation required and conclusions have changed	This review update added 14 studies including 493 participants to the review. With the addition of these studies, the precision of the main analyses has been improved. Several new subanalyses have been performed, including level of desaturation during exercise, setting (laboratory or daily life), short-term versus long-term effects, type of measurement (iso-time or maximal exertion) and oxygen dose. We graded the quality of evidence and were able to meta-analyse data for health-related quality of life.
12 July 2016	New search has been performed	The systematic search, review and meta-analysis have been updated to 12 July 2016.

CONTRIBUTIONS OF AUTHORS

ME designed the first draft of the review update, participated in data collection and management and wrote the first draft of the review update, together with ZA.

ZA collected the review data and wrote the first draft of the review update, together with ME.

ABH contributed to the analysis plan and performed the statistical analyses.

DC contributed to the conception, design and analysis plan for this review.

AA conceived of, designed and co-ordinated the original review published in 2011 and contributed to the design of this review update.

All review authors contributed to generation of the review manuscript.

DECLARATIONS OF INTEREST

ME, ZA, ABH and DC declare no conflicts of interest relevant to this work.

AA is the Chief Medical Officer and Chief Scientific Officer at Flatiron Health, a cancer-focused health technology company in New York, New York, USA. Collaborators and end-users of the dataset and Flatiron's research & analytics services include academic researchers, pharmaceutical manufacturers and government users, including the US FDA. Investors in Flatiron include three pharmaceutical manufacturers (Roche, Amgen, Celgene), but these venture investments do not relate to any drug products. These companies have no specific data rights and no say in the day-to-day research activities at Flatiron.

SOURCES OF SUPPORT

Internal sources

- The authors declare that no such funding was received for this systematic review, Other.

External sources

- The Heart-Lung Foundation, Sweden.
- Wera and Emil Cornell Foundation, Sweden.
- The Swedish Society of Medicine, Sweden.
- The Scientific Committee of Blekinge County Council, Sweden.

DIFFERENCES BETWEEN PROTOCOL AND REVIEW

After statistical review, we included only post scores and not change scores in the meta-analyses, so as not to compare SMDs on the same scale. We also added a post hoc sensitivity analysis of breathlessness that excluded outliers and short-burst oxygen therapy.

INDEX TERMS

Medical Subject Headings (MeSH)

Dyspnea [etiology] [*therapy]; Exercise; Health Status; Home Care Services; Oxygen Inhalation Therapy [*methods]; Pulmonary Disease, Chronic Obstructive [complications] [*therapy]; Quality of Life; Randomized Controlled Trials as Topic

MeSH check words

Humans