Cochrane
Library

Cochrane Database of Systematic Reviews

Acupuncture for stroke rehabilitation (Review)

Yang A, Wu HM, Tang JL, Xu L, Yang M, Liu GJ

Yang A, Wu HM, Tang JL, Xu L, Yang M, Liu GJ.
Acupuncture for stroke rehabilitation.
Cochrane Database of Systematic Reviews 2016, Issue 8. Art. No.: CD004131.
DOI: [10.1002/14651858.CD004131.pub3](https://doi.org/10.1002/14651858.CD004131.pub3).

www.cochranelibrary.com

Acupuncture for stroke rehabilitation (Review)

Copyright © 2016 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

WILEY

TABLE OF CONTENTS

ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
SUMMARY OF FINDINGS	3
BACKGROUND	7
OBJECTIVES	7
METHODS	7
RESULTS	10
Figure 1.	11
Figure 2.	13
Figure 3.	14
Figure 4.	18
DISCUSSION	19
AUTHORS' CONCLUSIONS	20
ACKNOWLEDGEMENTS	20
REFERENCES	21
CHARACTERISTICS OF STUDIES	45
DATA AND ANALYSES	108
Analysis 1.1. Comparison 1 Acupuncture plus baseline treatment versus sham acupuncture plus baseline treatment, Outcome 1 Improvement of motor function.	109
Analysis 1.2. Comparison 1 Acupuncture plus baseline treatment versus sham acupuncture plus baseline treatment, Outcome 2 Improvement of cognitive function.	110
Analysis 1.3. Comparison 1 Acupuncture plus baseline treatment versus sham acupuncture plus baseline treatment, Outcome 3 Improvement of Health-Related Quality of Life (MOS SF-36).	110
Analysis 1.4. Comparison 1 Acupuncture plus baseline treatment versus sham acupuncture plus baseline treatment, Outcome 4 Improvement of Stroke-Specific Quality of Life(SS QOL).	110
Analysis 2.1. Comparison 2 Acupuncture plus baseline treatment versus baseline treatment alone, Outcome 1 Improvement of dependency at the end of treatment.	113
Analysis 2.2. Comparison 2 Acupuncture plus baseline treatment versus baseline treatment alone, Outcome 2 Improvement of dependency at the end of follow-up.	114
Analysis 2.3. Comparison 2 Acupuncture plus baseline treatment versus baseline treatment alone, Outcome 3 Improvement of global neurological deficit at the end of treatment.	114
Analysis 2.4. Comparison 2 Acupuncture plus baseline treatment versus baseline treatment alone, Outcome 4 Improvement of global neurological deficit at the end of follow-up.	114
Analysis 2.5. Comparison 2 Acupuncture plus baseline treatment versus baseline treatment alone, Outcome 5 Improvement of global neurological deficit at the end of treatment.	115
Analysis 2.6. Comparison 2 Acupuncture plus baseline treatment versus baseline treatment alone, Outcome 6 Improvement of motor function at the end of treatment.	115
Analysis 2.7. Comparison 2 Acupuncture plus baseline treatment versus baseline treatment alone, Outcome 7 Improvement of general motor function at the end of follow-up (Fugl-Meyer Assessment).	116
Analysis 2.8. Comparison 2 Acupuncture plus baseline treatment versus baseline treatment alone, Outcome 8 Improvement of motor function at the end of treatment (Fugl-Meyer Assessment).	116
Analysis 2.9. Comparison 2 Acupuncture plus baseline treatment versus baseline treatment alone, Outcome 9 Improvement of cognitive function at the end of treatment.	116
Analysis 2.10. Comparison 2 Acupuncture plus baseline treatment versus baseline treatment alone, Outcome 10 Improvement of cognitive function at the end of follow-up (MMSE).	117
Analysis 2.11. Comparison 2 Acupuncture plus baseline treatment versus baseline treatment alone, Outcome 11 Improvement of cognitive function at the end of treatment.	117
Analysis 2.12. Comparison 2 Acupuncture plus baseline treatment versus baseline treatment alone, Outcome 12 Improvement of depression at the end of treatment.	118
Analysis 2.13. Comparison 2 Acupuncture plus baseline treatment versus baseline treatment alone, Outcome 13 Improvement of depression at the end of treatment (HAMD).	118
Analysis 2.14. Comparison 2 Acupuncture plus baseline treatment versus baseline treatment alone, Outcome 14 Improvement of swallowing function at the end of treatment (Water drinking test).	119

Analysis 2.15. Comparison 2 Acupuncture plus baseline treatment versus baseline treatment alone, Outcome 15 Improvement of swallowing function at the end of treatment (Water drinking test).	119
Analysis 2.16. Comparison 2 Acupuncture plus baseline treatment versus baseline treatment alone, Outcome 16 Improvement of pain at the end of treatment (Visual Analogue Scale).	119
Analysis 2.17. Comparison 2 Acupuncture plus baseline treatment versus baseline treatment alone, Outcome 17 Improvement of sleep quality at the end of treatment (Rhone Planck Sleepiness Scale).	119
Analysis 2.18. Comparison 2 Acupuncture plus baseline treatment versus baseline treatment alone, Outcome 18 Improvement of spasticity at the end of treatment (Modified Ashworth Spasticity Rating Scale).	120
Analysis 2.19. Comparison 2 Acupuncture plus baseline treatment versus baseline treatment alone, Outcome 19 Improvement of quality of life at the end of treatment (MOS SF-36).	120
APPENDICES	120
WHAT'S NEW	128
HISTORY	129
CONTRIBUTIONS OF AUTHORS	129
DECLARATIONS OF INTEREST	129
SOURCES OF SUPPORT	129
INDEX TERMS	129

[Intervention Review]

Acupuncture for stroke rehabilitation

Ai Yang¹, Hong Mei Wu¹, Jin-Ling Tang², Li Xu³, Ming Yang¹, Guan J Liu⁴

¹Center of Geriatrics and Gerontology, West China Hospital, Sichuan University, Chengdu, China. ²Division of Epidemiology, The Jockey Club School of Public Health and Primary Care, The Chinese University of Hong Kong, Hong Kong SAR, China. ³Center of Health Care and Management, Hospital of Chengdu Office of People's Government of Tibetan Autonomous Region, Chengdu, China. ⁴Chinese Cochrane Centre, Chinese Evidence-Based Medicine Centre, West China Hospital, Sichuan University, Chengdu, China

Contact: Hong Mei Wu, Center of Geriatrics and Gerontology, West China Hospital, Sichuan University, 37 Guo Xue street, Chengdu, China. drwhm@163.com, drwhm@126.com.

Editorial group: Cochrane Stroke Group.

Publication status and date: New search for studies and content updated (no change to conclusions), published in Issue 8, 2016.

Citation: Yang A, Wu HM, Tang JL, Xu L, Yang M, Liu GJ. Acupuncture for stroke rehabilitation. *Cochrane Database of Systematic Reviews* 2016, Issue 8. Art. No.: CD004131. DOI: [10.1002/14651858.CD004131.pub3](https://doi.org/10.1002/14651858.CD004131.pub3).

Copyright © 2016 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

Stroke is the second most common cause of death in the world and in China it has now become the main cause of death. It is also a main cause of adult disability and dependency. Acupuncture for stroke has been used in China for hundreds of years and is increasingly practiced in some Western countries. This is an update of the Cochrane review originally published in 2006 .

Objectives

To determine the efficacy and safety of acupuncture therapy in people with subacute and chronic stroke. We intended to test the following hypotheses: 1) acupuncture can reduce the risk of death or dependency in people with subacute and chronic stroke at the end of treatment and at follow-up; 2) acupuncture can improve neurological deficit and quality of life after treatment and at the end of follow-up; 3) acupuncture can reduce the number of people requiring institutional care; and 4) acupuncture is not associated with any intolerable adverse effects.

Search methods

We searched the Cochrane Stroke Group Trials Register (June 2015), the Cochrane Central Register of Controlled Trials (CENTRAL; *Cochrane Library* 2015, Issue 7), MEDLINE (1966 to July 2015, Ovid), EMBASE (1980 to July 2015, Ovid), CINAHL (1982 to July 2015, EBSCO), and AMED (1985 to July 2015, Ovid). We also searched the following four Chinese medical databases: China Biological Medicine Database (July 2015); Chinese Science and Technique Journals Database (July 2015); China National Infrastructure (July 2015), and Wan Fang database (July 2015).

Selection criteria

Truly randomised unconfounded clinical trials among people with ischaemic or haemorrhagic stroke, in the subacute or chronic stage, comparing acupuncture involving needling with placebo acupuncture, sham acupuncture, or no acupuncture.

Data collection and analysis

Two review authors independently selected trials for inclusion, assessed quality, extracted and cross-checked the data.

Main results

We included 31 trials with a total of 2257 participants in the subacute or chronic stages of stroke. The methodological quality of most of the included trials was not high. The quality of evidence for the main outcomes was low or very low based on the assessment by the system of Grades of Recommendation, Assessment, Development and Evaluation (GRADE).

Acupuncture for stroke rehabilitation (Review)

Copyright © 2016 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

Two trials compared real acupuncture plus baseline treatment with sham acupuncture plus baseline treatment. There was no evidence of differences in the changes of motor function and quality of life between real acupuncture and sham acupuncture for people with stroke in the convalescent stage.

Twenty-nine trials compared acupuncture plus baseline treatment versus baseline treatment alone. Compared with no acupuncture, for people with stroke in the convalescent phase, acupuncture had beneficial effects on the improvement of dependency (activity of daily living) measured by Barthel Index (nine trials, 616 participants; mean difference (MD) 9.19, 95% confidence interval (CI) 4.34 to 14.05; GRADE very low), global neurological deficiency (seven trials, 543 participants; odds ratio (OR) 3.89, 95% CI 1.78 to 8.49; GRADE low), and specific neurological impairments including motor function measured by Fugl-Meyer Assessment (four trials, 245 participants; MD 6.16, 95% CI 4.20 to 8.11; GRADE low), cognitive function measured by the Mini-Mental State Examination (five trials, 278 participants; MD 2.54, 95% CI 0.03 to 5.05; GRADE very low), depression measured by the Hamilton Depression Scale (six trials, 552 participants; MD -2.58, 95% CI -3.28 to -1.87; GRADE very low), swallowing function measured by drinking test (two trials, 200 participants; MD -1.11, 95% CI -2.08 to -0.14; GRADE very low), and pain measured by the Visual Analogue Scale (two trials, 118 participants; MD -2.88, 95% CI -3.68 to -2.09; GRADE low). Sickness caused by acupuncture and intolerance of pain at acupoints were reported in a few participants with stroke in the acupuncture groups. No data on death, the proportion of people requiring institutional care or requiring extensive family support, and all-cause mortality were available in all included trials.

Authors' conclusions

From the available evidence, acupuncture may have beneficial effects on improving dependency, global neurological deficiency, and some specific neurological impairments for people with stroke in the convalescent stage, with no obvious serious adverse events. However, most included trials were of inadequate quality and size. There is, therefore, inadequate evidence to draw any conclusions about its routine use. Rigorously designed, randomised, multi-centre, large sample trials of acupuncture for stroke are needed to further assess its effects.

PLAIN LANGUAGE SUMMARY

Acupuncture for stroke rehabilitation

Review question

Acupuncture is a treatment based on ancient Chinese medicine in which fine needles or pressure is applied at certain sites in the body for therapeutic purposes. We wanted to know whether acupuncture is effective in improving the recovery of daily activities, movement, and quality of life in people who had experienced a stroke more than one month previously.

Background

Stroke is a major cause of death in the world and can also cause severe disability. Acupuncture is a relatively simple, inexpensive and safe treatment that has been used in China for hundreds of years and is increasingly practiced in some Western countries. However, it remains uncertain whether the existing evidence is sufficiently reliable to recommend the routine use of acupuncture.

Study characteristics

We identified 31 studies to July 2015 for inclusion in the review. These included a total of 2257 participants who had had a stroke more than one month previously. They all investigated acupuncture aimed at promoting recovery compared with no acupuncture or sham acupuncture. Outcomes included measures of daily activities (activities of daily living), neurological function, movement, cognition, depression, swallowing, pain, and quality of life. Most of the studies (29/31) were conducted in China; the studies varied considerably with respect to the time of stroke, specific techniques used, and the frequency of acupuncture.

Key results

We found some evidence that acupuncture improved activities of daily living and a number of aspects of neurological function. However, these conclusions were based on studies with low quality evidence. No serious side effects were reported and there was no information on the effects of acupuncture on death or the need for institutional care.

Quality of the evidence

It proved difficult to reliably determine the quality of the evidence because of poor reporting of study characteristics. Therefore, we have described most conclusions as having low or very low quality evidence.

SUMMARY OF FINDINGS

Summary of findings for the main comparison.

Acupuncture + baseline treatment versus baseline treatment alone

Patient or population: adults with stroke

Settings: inpatients

Intervention: Acupuncture + baseline treatment

Comparison: baseline treatment

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	Nº of participants (studies)	Quality of the evidence (GRADE)	Comments
	Risk with baseline treatment	Risk with Acupuncture + baseline treatment				
Improvement of dependency at the end of treatment assessed with Barthel Index	The mean improvement of dependency at the end of treatment was 0	The mean improvement of dependency at the end of treatment in the intervention group was 9.19 undefined more (4.34 more to 14.05 more)	-	616 (9 RCTs)	⊕⊕⊕⊕ VERY LOW ^{1,2}	Substantial heterogeneity in results. Most studies were at high or unclear risk of bias. All of the studies were carried out in China
Improvement of global neurological deficit at the end of treatment assessed with Modified Edinburgh and Scandinavian Stroke Scale	The mean improvement of global neurological deficit at the end of treatment was 0	The mean improvement of global neurological deficit at the end of treatment in the intervention group was 2.39 undefined fewer (3.34 fewer to 1.43 fewer)	-	240 (4 RCTs)	⊕⊕⊕⊕ LOW ¹	Most studies were at high or unclear risk of bias. All of the studies were carried out in China
Improvement of global neurological deficit at the end of treatment	Study population		OR 3.89 (1.78 to 8.49)	543 (7 RCTs)	⊕⊕⊕⊕ LOW ¹	Most studies were at high or unclear risk of bias. All of the studies were carried out in China
	674 per 1000	890 per 1000 (787 to 946)				
	Moderate					
	733 per 1000	914 per 1000				

	(830 to 959)					
Improvement of motor function at the end of treatment - upper and lower extremities motor function (FMA) assessed with Fugl-Meyer Assessment	The mean improvement of motor function at the end of treatment - upper and lower extremities motor function was 0	The mean improvement of motor function at the end of treatment - upper and lower extremities motor function in the intervention group was 6.16 undefined more (4.2 more to 8.11 more)	-	245 (4 RCTs)	⊕⊕⊕⊕ LOW ¹	Most studies were at high or unclear risk of bias. All of the studies were carried out in China
Improvement of motor function at the end of treatment - general motor function assessed with Motor assessment scale	The mean improvement of motor function at the end of treatment - general motor function was 0	The mean improvement of motor function at the end of treatment - general motor function in the intervention group was 4.53 undefined more (2.99 more to 6.07 more)	-	60 (1 RCT)	⊕⊕⊕⊕ LOW ¹	
Improvement of general motor function at the end of follow up assessed with Fugl-Meyer Assessment follow-up: mean 3 months	The mean improvement of general motor function at the end of follow-up was 0	The mean improvement of general motor function at the end of follow-up in the intervention group was 7.59 more (0.98 more to 14.2 more)	-	(1 RCT)	⊕⊕⊕⊕ MODERATE ³	
Improvement of motor function at the end of treatment assessed with Fugl-Meyer Assessment	Study population		OR 2.41 (0.98 to 5.96)	125 (2 RCTs)	⊕⊕⊕⊕ LOW ¹	
	710 per 1000	855 per 1000 (705 to 936)				
	Moderate					
	720 per 1000	861 per 1000 (716 to 939)				
Improvement of cognitive function at the end of treatment assessed with Mini-mental state examination	The mean improvement of cognitive function at the end of treatment was 0	The mean improvement of cognitive function at the end of treatment in the intervention group was 2.54 undefined more (0.03 more to 5.05 more)	-	278 (5 RCTs)	⊕⊕⊕⊕ VERY LOW ^{1,2}	Substantial heterogeneity in results. Most studies were at high or unclear risk of bias. All of the studies were carried out in China

Improvement of cognitive function at the end of follow-up assessed with Mini-mental state examination follow-up: 1 month	The mean improvement of cognitive function at the end of follow up was 0	The mean improvement of cognitive function at the end of follow-up in the intervention group was 3.47 undefined more (2.43 more to 4.51 more)	-	71 (1 RCT)	⊕⊕⊕⊕ LOW ¹	
Improvement of cognitive function at the end of treatment assessed with: Mini-mental state examination	Study population		OR 3.82 (1.89 to 7.72)	166 (3 RCTs)	⊕⊕⊕⊕ LOW ¹	
	512 per 1000	800 per 1000 (665 to 890)				
	Moderate					
	533 per 1000	814 per 1000 (684 to 898)				
Improvement of depression at the end of treatment assessed with Hamilton Depression Scale	The mean improvement of depression at the end of treatment was 0	The mean improvement of depression at the end of treatment in the intervention group was 2.58 undefined fewer (3.28 fewer to 1.87 fewer)	-	552 (6 RCTs)	⊕⊕⊕⊕ VERY LOW ^{1,2}	Substantial heterogeneity in results. Most studies were at high or unclear risk of bias. All of the studies were carried out in China
Improvement of depression at the end of treatment assessed with Hamilton Depression Scale	Study population		OR 2.03 (1.10 to 3.72)	342 (4 RCTs)	⊕⊕⊕⊕ LOW ¹	Most studies were at high or unclear risk of bias. All of the studies were carried out in China
	784 per 1000	880 per 1000 (799 to 931)				
	Moderate					
	807 per 1000	894 per 1000 (821 to 939)				
Improvement of swallowing function at the end of treatment	The mean improvement of swallowing function at the end of treatment was 0	The mean improvement of swallowing function at the end of treatment in the intervention group was 1.11 undefined fewer (2.08 fewer to 0.14 fewer)	-	200 (2 RCTs)	⊕⊕⊕⊕ VERY LOW ^{1,2}	

Improvement of pain at the end of treatment assessed with Visual Analogue Scale	The mean improvement of pain at the end of treatment was 0	The mean improvement of pain at the end of treatment in the intervention group was 2.88 undefined fewer (3.68 fewer to 2.09 fewer)	-	118 (2 RCTs)	⊕⊕⊕⊖ LOW ¹
Improvement of sleep quality at the end of treatment assessed with Rhone Planck Sleepiness Scale	The mean improvement of sleep quality at the end of treatment was 0	The mean improvement of sleep quality at the end of treatment in the intervention group was 1.09 undefined fewer (2.37 fewer to 0.19 more)	-	60 (1 RCT)	⊕⊕⊕⊖ MODERATE ³
Improvement of spasticity at the end of treatment assessed with Modified Ashworth Spasticity Rating Scale	The mean improvement of spasticity at the end of treatment was 0	The mean improvement of spasticity at the end of treatment in the intervention group was 0.4 undefined fewer (0.64 fewer to 0.16 fewer)	-	60 (1 RCT)	⊕⊕⊕⊖ LOW ¹
Improvement of quality of life at the end of treatment assessed with MOS SF-36	The mean improvement of quality of life was 0	The mean improvement of quality of life in the intervention group was 2.73 undefined more (0.54 fewer to 6 more)	-	71 (1 RCT)	⊕⊕⊕⊖ LOW ¹

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: Confidence interval; **RR:** Risk ratio; **OR:** Odds ratio;

GRADE Working Group grades of evidence

High quality: We are very confident that the true effect lies close to that of the estimate of the effect

Moderate quality: We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different

Low quality: Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect

Very low quality: We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect

¹ Downgraded by two levels for very serious risk of bias (none of the trials used adequate allocation concealment, nor blinding of participants or researchers. Also, most of them were at risk of attrition bias).

² Downgraded by one level for serious inconsistency (due to substantial heterogeneity, $I^2 = 57\%$).

³ Downgraded by one level for this trial did not use blinding of participants or researchers.

BACKGROUND

Description of the condition

Stroke is the second leading cause of death in the world (GBD2013 2015). In the USA, stroke accounted for about one of every 19 deaths in 2010. On average, someone dies of stroke every four minutes (Go 2014). In 2013, stroke has become the leading cause of death in cities and rural areas in China (GBD2013 2015). Although age-standardised rates of stroke mortality have decreased worldwide in the past two decades, the absolute numbers of people with first stroke (16.9 million), stroke survivors (33 million), stroke-related deaths (5.9 million), and disability-adjusted life-years (DALYs) lost (102 million) in 2010 were still high and had significantly increased since 1990 (68%, 84%, 26%, and 12% increase, respectively), imposing a great burden on families and communities in low-income and middle-income countries (Feigin 2014). Despite considerable research efforts on multiple treatment modalities, there is still no single rehabilitation intervention demonstrated unequivocally to aid recovery. This reality drives people to search for other modalities of treatment in an attempt to further improve the outcome of stroke rehabilitation, such as acupuncture and Chinese herbal medicine.

Description of the intervention

Acupuncture is one of the main modalities of treatment in traditional Chinese medicine and can be traced back more than 3000 years in China (Wu 1996). Being a relatively simple, inexpensive, and safe treatment compared with other conventional interventions, acupuncture has been well accepted by Chinese patients and is widely used to improve motor, sensation, speech, and other neurological functions in people with stroke. As a therapeutic intervention, acupuncture is also increasingly practiced in some Western countries (Hegyi 2012; Johansson 1993; NIH 1998; Salom-Moreno 2014; Schaechter 2007; Wayne 2005).

How the intervention might work

Many studies in animals and humans have demonstrated that acupuncture can cause multiple biological responses, including circulatory and biochemical effects. These responses can occur locally or close to the site of application, or at a distance. They are mediated mainly by sensory neurons to many structures within the central nervous system. This can lead to activation of pathways affecting various physiological systems in the brain as well as in the periphery (Jansen 1989; Johansson 1993; Magnusson 1994; Sun 2001; Wang 2001). In summary, possible mechanisms of the effects of acupuncture on neurological conditions include stimulation of neuronal cell proliferation (Cheng 2008), facilitation of neural plasticity (Ren 2008), reduction of the post-ischaemic inflammatory reaction (Liu 2009a), and prevention of neuronal apoptosis (Zhang 2008a).

Why it is important to do this review

There are a large number of studies of the clinical efficacy of acupuncture in stroke rehabilitation published in the world, especially in China, but not all demonstrate a beneficial effect on stroke rehabilitation (Hu 1993; Johansson 2001; Sun 2001; Sze 2002; Zhan 2014; Zhang 1996; Zhang 2015). Many systematic reviews of trials of acupuncture in stroke rehabilitation have been conducted (Ernst 1996; Hopwood 1996; Kong 2010; Li 2014; Park 2001; Smith 2002; Sze 2002; Wu 2010a). These reviews, however,

included mainly trials with stroke patients in either the acute, or subacute, and chronic stages. Some reviews only focused on specific types of stroke (Liu 2005; Wang 2012a; Zheng 2011a), or needling (Dong 2013; Kim 2010; Zhou 2013), or neurologic deficits (Lim 2015; Long 2012; Liu 2014a; Park 2014; Qi 2009; Wong 2012; Zhang 2009a; Zhang 2012a; Zhang 2014a; Zhu 2011a). Knowing that a large number of clinical studies had been completed since this review was originally published in 2006 (Wu 2006), we aimed to conduct an up-to-date systematic review of publications regardless of subtype of stroke, or needling, or neurologic deficits.

The aim of this review was to systematically analyse all the randomised controlled trials of acupuncture for subacute and chronic stroke to provide the best available evidence to inform clinical practice and further research planning on stroke treatment.

OBJECTIVES

To determine the efficacy and safety of acupuncture therapy in people with subacute and chronic stroke. We intended to test the following hypotheses: 1) acupuncture can reduce the risk of death or dependency in people with subacute and chronic stroke at the end of treatment and at follow-up; 2) acupuncture can improve neurological deficit and quality of life after treatment and at the end of follow-up; 3) acupuncture can reduce the number of people requiring institutional care; and 4) acupuncture is not associated with any intolerable adverse effects.

METHODS

Criteria for considering studies for this review

Types of studies

In the review, we included randomised controlled clinical trials (RCTs) comparing acupuncture with at least one control group that used placebo, sham treatment, or conventional treatment in people with subacute (one to three months since onset) or chronic stroke (over three months since onset). We excluded trials using quasi-randomisation or the allocation of participants using alternation, case record numbers, dates of birth, day of the week, or controlled trials using any other non-random allocation methods.

Types of participants

Trials involving participants of any age or sex with ischaemic or haemorrhagic stroke in the subacute (one to three months since onset) or chronic phases (over three months since onset) were eligible. Stroke must have been diagnosed according to the World Health Organization definition (rapidly developed clinical signs of focal (or global) disturbances of cerebral function, lasting more than 24 hours or leading to death, with no other apparent cause than of vascular origin (Asplund 1988)), or confirmed by computerised tomography (CT), or magnetic resonance imaging (MRI). We did not include trials of participants with subarachnoid haemorrhage or subdural haematoma. We also excluded trials that included people in the acute phase of stroke (within one month since onset).

Types of interventions

We included trials evaluating acupuncture therapy that involved needling after stroke onset at the subacute or chronic phases, regardless of times of treatment or length of treatment period. We included either traditional acupuncture, in which the needles

were inserted in classical meridian points, or contemporary acupuncture, in which the needles were inserted in non-meridian or trigger points, regardless of the source of stimulation (for example, hand or electrical stimulation). We excluded trials in which the acupuncture treatment did not involve needling, such as acupressure or laser acupuncture.

The control interventions were placebo acupuncture, sham acupuncture, or other conventional treatment. Placebo acupuncture refers to a needle attached to the skin surface (not penetrating the skin but at the same acupoints) (Van Tulder 2000). Sham acupuncture refers to:

1. a needle placed in an area close to but not in the acupuncture points (Van Tulder 2000);
2. subliminal skin electrostimulation via electrodes attached to the skin (SCSSS 1999).

The comparisons we investigated were:

1. acupuncture only compared with placebo or sham treatment;
2. acupuncture in addition to baseline medication or treatment compared with placebo or sham treatment in addition to baseline medication or treatment;
3. acupuncture in addition to baseline medication or treatment compared with baseline medication or treatment alone.

We excluded trials that compared different forms of acupuncture only and we also excluded trials reporting only physiological or laboratory parameters.

Types of outcome measures

We included trials that used at least one of the following outcome measures.

Primary outcomes

Death or dependency at the end of follow-up (at least three months or longer after stroke onset). We defined dependency as dependent on others in activities of daily living, based on the correlated definition of the Barthel scores (Activities of Daily Living, ADL) as a score of less than 60 or an Oxford handicap grade 3 to 6 (Sulter 1999), or the trialists' own definition.

Secondary outcomes

1. The proportion of people requiring institutional care or requiring extensive family support at the end of follow-up (at three months or longer after stroke onset). Family care is the main form of care for severely dependent people in developing countries.
2. Changes of neurological deficit after acupuncture treatment and at the end of follow-up (at three months or longer after stroke onset). The measures could focus on specific impairment (for example, Motricity Index, or Motor Assessment Scale, which assess only motor function), or global neurological deficit (for example, the National Institute of Health Stroke Scale, European Stroke Scale, the Scandinavian Stroke Scale) or two kinds of Chinese Stroke Recovery Scales, which involve motor, sensory and other impaired neurological functions. The Chinese Stroke Recovery Scale 1 (CSRS 1) refers to "the Revised Diagnostic Criteria of Acute Cerebral Infarction" formulated by the second National Academic Symposium on Cerebrovascular Diseases of

the Chinese Medical Association in 1986, which is similar to the Revised Scandinavian Stroke Scale (RSSS). The Chinese Stroke Recovery Scale 2 (CSRS 2) refers to "the Chinese Stroke Recovery Scale based on principles of traditional Chinese medicine".

3. Death from any causes during the entire treatment and follow-up period.
4. Quality of life (QOL) at the end of follow-up (at three months or longer after stroke onset). This could be measured by the Nottingham Health Profiles or Spitzer Quality of Life Index.
5. Possible adverse events including dizziness, difficulty in tolerating electrostimulation, infection, puncture of a lung, heart tamponades, spinal cord injury, disrupted pacemaker function; and presumed to be caused by acupuncture or electrostimulation. We evaluated the number of participants developing at least one severe adverse event listed above.

Search methods for identification of studies

See the 'Specialized register' section in the [Cochrane Stroke Group](#) module. We searched for all trials in all languages and arranged translation of relevant papers where necessary.

Electronic searches

Original searches (2005)

We initially obtained relevant trials from the following sources with no language restriction.

1. Cochrane Stroke Group Trials Register (November 2005).
2. Cochrane Central Register of Controlled Trials (CENTRAL) (Cochrane Library 2005, Issue 3).
3. MEDLINE (1966 to November 2005) combined with the Cochrane highly sensitive search strategy for identifying RCTs in MEDLINE (Dickersin 1994).
4. EMBASE (1980 to November 2005). Modified MEDLINE search and combined with the Cochrane highly sensitive search strategy for identifying RCTs in EMBASE (Lefebvre 1996).
5. Chinese Stroke Trials Register (November 2005).
6. Chinese Acupuncture Trials Register (November 2005).
7. Trials Register of the Cochrane Complementary Medicine Field (November 2005).
8. CINAHL (1982 to November 2005).
9. AMED (the Allied and Complementary Medicine Database, 1985 to November 2005).
10. Chinese Biological Medicine Database (CBM-disc, 1979 to November 2005).
11. National Center for Complementary and Alternative Medicine Register (http://nccam.nih.gov/clinical_trials/) and National Institute of Health Clinical Trials Database (<http://clinicaltrials.gov>) (searched November 2005).

Review update (2015)

We searched CENTRAL and the Cochrane Stroke Group Trials Register as above for new trials. We obtained relevant new trials from the following sources with no language restriction in the updated review.

1. Cochrane Stroke Group Trials Register (June 2015; [Appendix 1](#)).
2. Cochrane Central Register of Controlled Trials (CENTRAL; 2015, Issue 7) ([Appendix 2](#)).

3. MEDLINE (July 2015; Ovid; [Appendix 3](#)).
4. EMBASE (July 2015; Ovid; [Appendix 4](#)).
5. CINAHL (July 2015; EBSCO; [Appendix 5](#)).
6. AMED (the Allied and Complementary Medicine Database, July 2015; Ovid; [Appendix 6](#)).
7. Chinese Biological Medicine Database (July 2015; [Appendix 7](#)).
8. Chinese Science and Technique Journals Database (to July 2015; [Appendix 8](#)).
9. China National Infrastructure (to July 2015; [Appendix 9](#)).
10. WanFang database (to July 2015; [Appendix 10](#)).

Searching other resources

Original searches (2005)

1. We handsearched four Chinese journals relevant to acupuncture (from 1980 to November 2005):
 - a. *Acupuncture Research*;
 - b. *Chinese Acupuncture and Moxibustion*;
 - c. *Journal of Clinical Acupuncture and Moxibustion*;
 - d. *Shanghai Journal of Acupuncture and Moxibustion*.
2. We checked the reference lists of all relevant papers identified, including two systematic reviews ([Park 2001](#); [Sze 2002](#)), for further published and unpublished trials.

Review update (2015)

We searched the reference lists of all relevant papers identified.

Data collection and analysis

Selection of studies

Two review authors (AY, LX) independently checked the titles and abstracts of trials for inclusion based on the selection criteria outlined above. We retrieved the full text of the article if there was any doubt whether the article should be excluded or not. In cases of disagreement between the two review authors, a third member of the stroke research group (JLT or HMW) reviewed the information to decide on inclusion or exclusion of an article.

Data extraction and management

Two review authors (AY, L) independently extracted information on participants, methods, interventions, outcomes, and results by using a self-developed data extraction form. We translated studies not in English or Chinese before assessment. Where more than one publication for a study existed, we grouped reports together and we used the publication with the most complete data. Where relevant outcomes were only published in earlier versions, we used these data. We resolved disagreements by involving a third review author (JLT or HMW) or through discussion.

Assessment of risk of bias in included studies

We assessed the following items using the Cochrane risk of bias assessment tool ([Higgins 2011a](#); [Appendix 11](#)).

1. Was there adequate sequence generation?
2. Was allocation adequately concealed?
3. Was knowledge of the allocated interventions adequately prevented during the study?
4. Were incomplete outcome data adequately addressed?
5. Are reports of the study free of suggestion of selective outcome reporting?
6. Was the study apparently free of other problems that could put it at a risk of bias?

Two review authors (AY, LX) independently assessed risk of bias; any disagreements were resolved by a third review author (JLT or HMW).

Measures of treatment effect

For dichotomous outcomes (e.g. death or dependency, adverse effects), we expressed the results as odds ratios (ORs) with 95% confidence intervals (CIs). For continuous outcomes (e.g. quality of life), we used the mean difference (MD), or the standardised mean difference (SMD) if different scales were used.

Assessment of heterogeneity

We tested heterogeneity between trial results using a standard Chi² test on N-1 degrees of freedom, with a threshold value of P < 0.1, and with the I² test ([Higgins 2003](#)). I² values of 25%, 50% and 75% correspond to low, medium and high levels of heterogeneity.

Assessment of reporting biases

If we identified a sufficient number of studies, we planned to examine for potential publication bias using a funnel plot ([Sterne 2011](#)).

Data synthesis

We pooled data using the random-effects model but we also used the fixed-effect model to ensure robustness of the model chosen and susceptibility to outliers.

Subgroup analysis and investigation of heterogeneity

We planned to undertake subgroup analyses to explore possible sources of heterogeneity (e.g. participants, interventions, and study quality). Heterogeneity among participants could be related to the type and severity of stroke. Post-stroke neurological recovery is known to be obvious within three months, especially post-stroke motor recovery, and is unlikely beyond six months, so heterogeneity in treatments could be related to times to start of treatment. Mixing patients with interval from stroke onset of less than three months, three to six months, and more than six months in one sample would make the assessment of the efficacy of an intervention methodologically unsound. If appropriate data were available, we planned a subgroup analysis to compare participants with different times to start of treatment (within three months, three to six months, and after six months) using the method outlined by [Deeks 2001](#).

Sensitivity analysis

If appropriate data were available, we planned a sensitivity analysis to assess the effects of including only those trials:

1. that were double blind;
2. with adequate concealment of randomisation;
3. published in a language other than Chinese.

RESULTS

Description of studies

Results of the search

2006 version

For the 2006 version of this review, we identified 6402 potentially relevant articles, retrieved 50 full-text articles, and included five trials with 368 participants (Dai 1997; Li 1997a; Lun 1999; Naeser 1992; Wang 2001).

2015 update

For the update of this review, we identified 5874 potentially relevant articles from January 2005 to July 2015. Of these, we retrieved 371 full-text articles for further assessment and included 26 trials with 1889 participants (Bao 2012; Chou 2009; Gao 2014a; Guo 2011; Guo 2012; Huang 2008a; Ke 2015; Li 2010a; Li 2011a; Li 2013a; Liu 2013a; Sun 2013a; Sun 2015; Wang 2011a; Wang 2012; Wu 2008; Wu 2011a; Wu 2013a; Xu 2013; Yao 2014; Zhan 2014; Zhang 2013a; Zhang 2015; Zheng 2014; Zhou 2014; Zhu 2007). There are four relevant ongoing studies (Fu 2011; Liu 2013b; Xie 2006; Zhong 2010; [Characteristics of ongoing studies](#)). In total, we have included 31 trials in this updated review. Results of the search are displayed in [Figure 1](#).

Figure 1. Study flow diagram.

Figure 1. (Continued)

treatment versus baseline treatment
--

Included studies

A total of 31 trials were eligible for inclusion in this systematic review. For details of each included trial please see the [Characteristics of included studies](#) table.

Participants

Of the 31 included trials, 29 were conducted in China, one in the UK, and one in the USA.

The age of participants ranged from 24 to 95 years. More men than women were included in 23 trials (between 47% and 75% men) (Bao 2012; Dai 1997; Guo 2011; Guo 2012; Huang 2008a; Ke 2015; Li 2010a; Li 2011a; Liu 2013a; Lun 1999; Sun 2013a; Wang 2001; Wang 2011a; Wang 2012; Wu 2008; Wu 2011a; Yao 2014; Zhan 2014; Zhang 2013a; Zhang 2015; Zheng 2014; Zhou 2014; Zhu 2007). Three trials did not describe the gender of the participants (Li 1997a; Li 2013a; Naeser 1992). The range of time from stroke onset was from one month to 8.5 years.

There were five trials involving participants with interval from stroke onset between one to three months (Huang 2008a; Li 2013a; Naeser 1992; Wu 2013a; Zhou 2014), 10 trials more than three months (Dai 1997; Bao 2012; Chou 2009; Gao 2014a; Guo 2012; Li 2011a; Liu 2013a; Sun 2013a; Sun 2015; Zhang 2013a), and 16 trials including participants with interval from stroke onset of either less than three months or more than three months (Guo 2011; Ke 2015; Li 1997a; Li 2010a; Lun 1999; Wang 2001; Wang 2011a; Wang 2012; Wu 2008; Wu 2011a; Xu 2013; Yao 2014; Zhan 2014; Zhang 2015; Zheng 2014; Zhu 2007).

Seven trials included participants with ischaemic stroke only (Dai 1997; Bao 2012; Huang 2008a; Naeser 1992; Wu 2011a; Zhang 2015; Zhou 2014). Five trials did not describe the type of stroke (Guo 2012; Li 2011a; Sun 2015; Xu 2013; Zhan 2014). All other trials included participants with ischaemic and haemorrhagic stroke (Chou 2009; Gao 2014a; Guo 2011; Ke 2015; Li 1997a; Li 2010a; Li 2013a; Liu 2013a; Lun 1999; Sun 2013a; Wang 2001; Wang 2011a; Wang 2012; Wu 2008; Wu 2013a; Yao 2014; Zhang 2013a; Zheng 2014; Zhu 2007). All of the included trials used CT or MRI to confirm the diagnosis of stroke. The severity on entry was mild to severe in two trials (Dai 1997; Naeser 1992), without a definition of severity or not stated in the remaining 29 trials (Bao 2012; Chou 2009; Gao 2014a; Guo 2011; Guo 2012; Huang 2008a; Ke 2015; Li 1997a; Li 2010a; Li 2011a; Li 2013a; Liu 2013a; Lun 1999; Sun 2013a; Sun 2015; Wang 2001; Wang 2011a; Wang 2012; Wu 2008; Wu 2011a; Wu 2013a; Xu 2013; Yao 2014; Zhan 2014; Zhang 2013a; Zhang 2015; Zheng 2014; Zhou 2014; Zhu 2007).

Interventions

Two trials compared real acupuncture plus baseline treatment with sham acupuncture plus baseline treatment (Chou 2009; Naeser 1992); the remaining 29 trials compared acupuncture plus baseline medication or treatment with baseline medication or treatment alone. None of the trials compared acupuncture only with placebo or sham treatment.

Among the included trials there were 10 three-armed trials (Dai 1997; Guo 2011; Huang 2008a; Li 1997a; Li 2013a; Sun 2015; Wang 2001; Wu 2013a; Zhang 2013a; Zhang 2015) and one four-armed trial (Wu 2008) comparing acupuncture plus baseline medication or treatment with baseline medication treatment alone, one kind of acupuncture with another kind of acupuncture, or acupuncture only with other treatment. In this review, the baseline medication or treatment included Western medicine (WM), traditional Chinese medicine (TCM), non-pharmacological therapy, or a combination. WM included aspirin and other conventional drug therapies.

With one exception (Chou 2009), none of the other 30 included trials reported the acupuncture rationale or acupuncturists' background, including duration of relevant training, length of clinical experience and expertise in the specific condition. The acupuncture interventions used varied considerably across trials. Nineteen trials used only manual stimulation (Bao 2012; Dai 1997; Gao 2014a; Guo 2012; Huang 2008a; Ke 2015; Li 1997a; Li 2011a; Li 2013a; Lun 1999; Wang 2001; Wang 2011a; Wang 2012; Wu 2008; Wu 2013a; Xu 2013; Zhan 2014; Zhang 2015; Zheng 2014), four used only electrical stimulation (Chou 2009; Naeser 1992; Sun 2015; Wu 2011a), and eight used the combination of manual and electrical stimulation (Guo 2011; Li 2010a; Liu 2013a; Sun 2013a; Yao 2014; Zhang 2013a; Zhou 2014; Zhu 2007). Acupuncture point prescriptions were not consistent, with 15 trials involving either scalp or body acupoints (Chou 2009; Guo 2012; Huang 2008a; Ke 2015; Li 2011a; Li 2013a; Liu 2013a; Lun 1999; Wang 2001; Wang 2011a; Wang 2012; Wu 2013a; Zhan 2014; Zhang 2013a; Zhang 2015), and 16 trials using both body and scalp acupoints (Bao 2012; Dai 1997; Gao 2014a; Guo 2011; Li 1997a; Li 2010a; Naeser 1992; Sun 2013a; Sun 2015; Wu 2008; Wu 2011a; Xu 2013; Yao 2014; Zheng 2014; Zhou 2014; Zhu 2007). Numbers of points used ranged from one to 27 points in all included trials. The needle retention time was 15 to 40 minutes in all included trials. Thirteen trials reported the achievement of 'deqi', an irradiating feeling said to indicate effective needling (Bao 2012; Chou 2009; Dai 1997; Li 2010a; Li 2011a; Liu 2013a; Lun 1999; Sun 2013a; Sun 2015; Wang 2001; Yao 2014; Zheng 2014; Zhu 2007). Information on needle type was available in 13 trials (Bao 2012; Chou 2009; Dai 1997; Li 2010a; Lun 1999; Liu 2013a; Naeser 1992; Sun 2013a; Sun 2015; Wu 2011a; Yao 2014; Zheng 2014; Zhu 2007). The length of treatment period ranged from one to 24 weeks with the number of treatment sessions varying from six to 120 sessions and the frequency of treatment from five sessions per week to two sessions per day.

Outcomes

The most commonly reported outcomes were dependency, global neurological deficit, and specific neurological impairment after acupuncture treatment. Thirteen trials evaluated the effect of acupuncture on dependency (activities of daily living) at the end of acupuncture treatment. The measures employed included the Barthel Index (BI) or modified Barthel Index (MBI) in 11 trials (Bao 2012; Huang 2008a; Ke 2015; Li 2010a; Wang 2012; Wu 2011a; Yao 2014; Zhan 2014; Zhang 2015; Zheng 2014; Zhou 2014) and the Physical Self-maintenance Scale (PSMS) combined with

Instrumental Activities of Daily Living Scale (IADL) in two trials (Li 2011a; Sun 2013a). Twelve trials measured the global neurological deficit score or the proportion of participants with an improvement of global neurological deficit at the end of acupuncture treatment (Dai 1997; Gao 2014a; Guo 2011; Huang 2008a; Li 1997a; Lun 1999; Sun 2013a; Wang 2001; Yao 2014; Zhang 2015; Zhou 2014; Zhu 2007). The measures employed included NIHSS, CSRS1 (e.g. MESSS) and CSRS2 (TCM). Six trials evaluated the effect of acupuncture on motor function measured with Fugl-Meyer scale (FMA) and Motor assessment scale after treatment (Li 2013a; Wang 2011a; Wang 2012; Wu 2011a; Yao 2014; Zhou 2014). Eighteen trials evaluated the effect of acupuncture on specific neurological impairments (e.g. cognitive function; swallowing function; depression) at the end of treatment (Bao 2012; Gao 2014a; Guo 2011; Guo 2012; Li 2010a; Li 2011a; Liu 2013a; Sun 2013a; Sun 2015; Wang 2011a; Wu 2008; Wu 2011a; Wu 2013a; Xu 2013; Yao 2014; Zhang 2013a; Zheng 2014; Zhou 2014). Two trials reported quality of life measured with the Medical Outcomes Study 36-Item Short-Form Health Survey (MOS SF-36) (Chou 2009; Sun 2013a). Only five trials reported information on adverse events (Li 2010a; Sun 2013a; Sun 2015; Zhang 2013a; Zhou 2014). None of the 31 included trials provided any information on death, proportion of participants requiring institutional care or extensive family support after acupuncture treatment or at the end of follow-up.

Excluded studies

Of the 12,276 citations identified from English and Chinese databases, we excluded 11,855 citations during the initial screening of titles and abstracts. The main reasons for these exclusions were as follows.

1. Not stroke.
2. Studies not RCT.
3. Stroke duration since onset less than 30 days.
4. Acupuncture not involving needling: such as acupressure; laser acupuncture.
5. Studies comparing different kinds of acupuncture therapies.
6. Review articles.

We retrieved a total of 421 potentially eligible studies for screening of the full-text papers.

Of the 421 potentially eligible studies, we excluded 390 studies for the following reasons.

1. Questionable randomisation.
2. Comparing two different methods of acupuncture.
3. Stroke duration since onset less than 30 days.
4. Duplicates.
5. Others: abstract, review.

For details please see the [Characteristics of excluded studies](#) table.

Risk of bias in included studies

Please see [Characteristics of included studies](#) for details of the methodological quality, and [Figure 2](#) and [Figure 3](#) for summaries of the risk of bias findings in all included trials.

Figure 2. Risk of bias graph: review authors' judgements about each risk of bias item presented as percentages across all included studies.

Figure 3. Risk of bias summary: review authors' judgements about each risk of bias item for each included study.

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding (performance bias and detection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias
Bao 2012	+	?	?	?	?	?	?	?
Chou 2009	?	?	?	?	?	+	?	?
Dai 1997	?	?	?	?	?	?	?	?
Gao 2014a	+	?	?	?	?	+	?	?
Guo 2011	?	?	?	?	?	?	?	?
Guo 2012	?	?	?	?	?	?	?	?
Huang 2008a	+	?	?	?	?	?	?	?
Ke 2015	?	?	?	?	?	?	?	?
Li 1997a	?	?	?	?	?	?	?	?
Li 2010a	+	?	?	?	?	+	?	?
Li 2011a	?	?	?	?	?	?	?	?
Li 2013a	?	?	?	?	?	?	?	?
Liu 2013a	?	?	?	?	?	?	?	?
Lun 1999	?	?	?	?	?	?	?	?
Naeser 1992	?	?	?	+	?	?	?	?
Sun 2013a	+	?	?	?	?	+	?	?
Sun 2015	+	?	?	?	?	?	?	?
Wang 2001	?	?	?	?	?	?	?	?
Wang 2011a	+	?	?	?	?	?	?	?
Wang 2012	+	?	?	?	?	?	?	?

Figure 3. (Continued)

Wang 2012	+	?	?	?	?	?	?	?
Wu 2008	?	?	?	?	?	?	?	?
Wu 2011a	+	?	?	?	?	?	?	?
Wu 2013a	+	?	?	?	?	?	?	?
Xu 2013	?	?	?	?	?	?	?	?
Yao 2014	+	?	?	?	?	?	?	?
Zhan 2014	+	?	?	?	?	+	?	?
Zhang 2013a	+	+	-	-	?	?	?	?
Zhang 2015	+	?	?	?	?	?	?	?
Zheng 2014	+	?	?	?	?	+	?	?
Zhou 2014	+	+	+	?	+	+	?	?
Zhu 2007	?	?	?	?	?	?	?	?

Allocation

Random sequence generation

Thirteen trials randomly assigned participants to groups by using random number tables (Bao 2012; Huang 2008a; Gao 2014a; Li 2010a; Sun 2013a; Wang 2011a; Wang 2012; Wu 2013a; Yao 2014; Wu 2011a; Zhan 2014; Zhang 2015; Zheng 2014). The remaining 18 trials did not report their methods of random sequence generation.

Allocation concealment

Of the 31 included trials, only three trials reported adequate allocation concealment by using sealed envelopes (Sun 2015; Zhang 2013a; Zhou 2014).

Blinding

Zhang 2013a reported that participants, their physicians, and outcome assessors were not blinded. Zhou 2014 reported that the investigators were blinded but the participants were not. Naeser 1992 reported that the participants were blinded without describing the method in detail. No information on blinding was available in the remaining 28 trials.

Incomplete outcome data

Seven trials reported withdrawals, but the results were not analysed on an intention-to-treat basis (Chou 2009; Gao 2014a; Li 2010a; Sun 2013a; Zhan 2014; Zheng 2014; Zhou 2014). There was no statement on dropouts or withdrawals in any of the remaining 24 trials. For each of these 24 trials, the number of participants randomised was the same as participants analysed, so it appears that there were no exclusions from the trials after randomisation. We concluded that, although the results appeared to be analysed by intention-to-treat, we cannot be certain about this.

Selective reporting

The included trials in this review did not report some clinically important outcomes, such as death, requiring Institutional care, and all-cause mortality. Of the 31 included studies, only five studies reported adverse events (Li 2010a; Sun 2013a; Sun 2015; Zhang 2013a; Zhou 2014), therefore we assumed that this may have constituted some degree of reporting bias.

Other potential sources of bias

There was insufficient information reported to determine if there were any other potential sources of bias.

Effects of interventions

See: [Summary of findings for the main comparison](#)

Acupuncture plus baseline treatment versus sham acupuncture plus baseline treatment

Two trials compared real acupuncture plus baseline treatment with sham acupuncture plus baseline treatment (Chou 2009; Naeser 1992).

Changes of specific neurological impairments after acupuncture treatment and at the end of follow-up

One trial with 16 participants evaluated the effect of acupuncture on the improvement of motor function after treatment by using a categorical approach rather than continuous scales (Naeser 1992). It showed that real acupuncture was not superior to sham acupuncture in the improvement of motor function for participants with stroke in the convalescent stage (OR 9.00, 95% CI 0.40 to 203.30; Analysis 1.1), but the confidence interval was very wide and included clinically significant effects in both directions.

Another trial with 33 participants evaluated the effect of acupuncture on the change of cognitive function after treatment

(Chou 2009). There were significant improvements in orientation (MD 4.21, 95% CI 1.78 to 6.64), perception (MD 5.32, 95% CI 0.93 to 9.71), and praxis (MD 3.80, 95% CI 2.12 to 5.48) among participants in the real acupuncture group compared with the sham acupuncture group, but acupuncture was not superior to control in the improvement of visuomotor organisation (MD 2.76, 95% CI -0.58 to 6.10), thinking operation (MD 0.12 95% CI -0.64 to 0.88), and memory (MD 0.35, 95% CI -0.43 to 1.13) (Analysis 1.2).

Quality of life at the end of follow-up

One trial with 33 participants evaluated health-related quality of life measured by means of self-report using the Medical Outcomes Study 36-Item Short-Form Health Survey (SF-36) and Stroke-Specific Quality of Life Scale (SS-QOL) (Chou 2009). It showed that real acupuncture was not superior to sham acupuncture in the improvement of health-related quality of life measured by the SF-36 (physical component summary: MD -2.66, 95% CI -10.57 to 5.25; mental component summary: MD 11.52, 95% CI -1.76 to 24.80; Analysis 1.3), and the SS-QOL (language: MD 13.34, 95% CI -3.02 to 29.70; personality: MD 3.10, 95% CI -11.73 to 17.93; Analysis 1.4) from baseline to week eight among participants with stroke in the convalescent stage.

No data on death or dependency, the proportion of people requiring institutional care or requiring extensive family support, death from any cause and possible adverse events were available in these two trials (Chou 2009; Naeser 1992).

Acupuncture plus baseline treatment versus baseline treatment alone

Twenty-nine trials compared acupuncture plus baseline treatment with baseline treatment alone (Bao 2012; Dai 1997; Gao 2014a; Guo 2011; Guo 2012; Huang 2008a; Ke 2015; Li 1997a; Li 2010a; Li 2011a; Li 2013a; Liu 2013a; Lun 1999; Sun 2013a; Sun 2015; Wang 2001; Wang 2011a; Wang 2012; Wu 2008; Wu 2011a; Wu 2013a; Xu 2013; Yao 2014; Zhan 2014; Zhang 2013a; Zhang 2015; Zheng 2014; Zhou 2014; Zhu 2007).

Death or dependency at the end of treatment and follow-up

No data on death at the end of treatment or follow-up were available in any of the included trials.

Eleven trials with a total of 956 participants measured improvement of dependency after acupuncture treatment by using continuous scales only (Barthel Index (BI), Modified Barthel Index (MBI) and Activity of Daily Living Scale (ADL)) (Bao 2012; Huang 2008a; Ke 2015; Li 2010a; Li 2011a; Wang 2012; Wu 2011a; Yao 2014; Zhan 2014; Zhang 2015; Zheng 2014). There was significant improvement in dependency measured by the BI, the MBI, and the ADL Scale among participants in the acupuncture group compared with the control group (BI: MD 9.19, 95% CI 4.34 to 14.05; participants = 616; studies = 9; $I^2 = 95\%$; MBI: MD 3.44, 95% CI 0.30 to 6.58; participants = 60; studies = 1; ADL: MD 7.80, 95% CI 6.04 to 9.56; participants = 62; studies = 1; Analysis 2.1). Heterogeneity was significant and may be attributable to differences in severity of stroke, times of evaluation from stroke onset, and types of stroke.

One trial with 147 participants evaluated the effect of acupuncture on the improvement of dependency measured by the BI at the end of three months' follow-up (Zhou 2014). There was a beneficial effect of acupuncture on the improvement of dependency for

participants with stroke in the convalescent stage (MD 7.49, 95% CI 1.79 to 13.19; Analysis 2.2). One trial with 71 participants evaluated the effect of acupuncture on the improvement of dependency measured by the ADL scale at the end of one month's follow-up (Sun 2013a). It also showed a beneficial effect of acupuncture on the improvement of dependency (MD 3.83, 95% CI 2.67 to 4.99; Analysis 2.2).

Proportion requiring institutional care or requiring extensive family support at the end of follow-up

No data on the proportion of participants requiring institutional care or extensive family support at the end of follow-up were available in these 29 trials.

Changes of global neurological deficit after acupuncture treatment and at the end of follow-up

Eight trials measured global neurological deficit at the end of treatment and follow-up by using continuous scales. Of these, four trials evaluated the effect of acupuncture on global neurological function measured by the Modified Edinburgh and Scandinavian Stroke Scale (Gao 2014a; Huang 2008a; Zhang 2015; Zhu 2007). There was a significant decrease in the neurological deficit score among participants in the acupuncture group compared with the control group (MD -2.39, 95% CI -3.34 to -1.43; participants = 240; studies = 4; $I^2 = 23\%$; Analysis 2.3). Two trials evaluated the effect of acupuncture on global neurological function measured by the Neurological Function Deficit Scale (NFDS) (Guo 2011; Yao 2014). There was no significant improvement in global neurological function among participants in acupuncture group compared with control group (MD -1.02, 95% CI -5.80 to 3.76; participants = 123; studies = 2; $I^2 = 19\%$; Analysis 2.3). One trial evaluated the effect of acupuncture on the improvement of global neurological function measured by the NFDS at the end of one month follow-up (Sun 2013a). It showed a beneficial effect of acupuncture on the improvement of global neurological function (MD -6.15, 95% CI -7.09 to -5.21; Analysis 2.4). One trial evaluated the effect of acupuncture on the improvement of global neurological function by using the National Institute of Health Stroke Scale (NIHSS) at the end of three months' follow-up (Zhou 2014). There was no significant improvement in global neurological function among participants in acupuncture group compared with control group (MD -0.83, 95% CI -1.94 to 0.28; Analysis 2.4).

Seven trials with a total of 543 participants measured improvement of global neurological deficit after acupuncture treatment by using categorical approaches only (CSRS 1 and CSRS 2) rather than continuous scales, for example changes of global neurological deficit score (Dai 1997; Huang 2008a; Li 1997a; Lun 1999; Wang 2001; Yao 2014; Zhang 2015). There was significant heterogeneity among the seven trials ($I^2 = 55\%$), which was possibly due to differences in outcome measurements used, times of evaluation from stroke onset, and types of stroke. This means that the overall estimate of treatment effect is not reliable. Hence, the apparent improvement of global neurological deficit in the acupuncture group compared with the control group should be interpreted with caution (OR 3.89, 95% CI 1.78 to 8.49; participants = 543; studies = 7; $I^2 = 55\%$; Analysis 2.5).

Changes of specific neurological impairments after acupuncture treatment and at the end of follow-up

Of the 31 included trials, 20 measured specific neurological impairment at the end of treatment and follow-up, mainly including motor function, cognitive function, speech function, depression, swallowing function, pain, sleep, and spasticity.

Motor function

Four trials used the Fugl-Meyer Assessment (FMA) to measure motor function in the upper and lower extremities at the end of treatment (Li 2013a; Wang 2011a; Wu 2011a; Yao 2014). Acupuncture was superior to no acupuncture in the improvement of motor function in the upper and lower extremities (MD 6.16, 95% CI 4.20 to 8.11; participants = 245; studies = 4; $I^2 = 28\%$; Analysis 2.6). One trial with 60 participants evaluated the effect of acupuncture on the improvement of general motor function measured by the Motor Assessment Scale at the end of treatment (Wang 2012). There was a significant difference between the two groups (MD 4.53, 95% CI 2.99 to 6.07; Analysis 2.6). One trial with 147 participants evaluated the effect of acupuncture on the improvement of general motor function measured by the FMA at the end of three months' follow-up (Zhou 2014). There was a beneficial effect of acupuncture on the improvement of general motor function for participants with stroke in the convalescent stage (MD 7.59, 95% CI 0.98 to 14.20; Analysis 2.7).

Two trials measured improvement of motor function after acupuncture treatment by using categorical approaches (FMA) (Li 2013a; Wang 2011a). There was no significant improvement in motor function among participants in the acupuncture group compared with the control group (OR 2.41, 95% CI 0.98 to 5.96; Analysis 2.8).

Cognitive function

Five trials evaluated the effect of acupuncture on the improvement of cognitive function measured by the Mini-Mental State Examination (MMSE) at the end of treatment (Bao 2012; Li 2010a; Li 2011a; Liu 2013a; Zheng 2014). Acupuncture was superior to no acupuncture in the improvement of cognitive function for participants with stroke in the convalescent stage (MD 2.54, 95% CI 0.03 to 5.05; participants = 278; studies = 5; $I^2 = 98\%$; Analysis 2.9). The significant heterogeneity between the trials was possibly due to differences in times of evaluation from stroke onset, severity on entry, needling details, and outcome measurements used. Two trials measured this outcome by using the Montreal Cognitive Assessment Scale (MoCA) at the end of treatment (Bao 2012; Zheng 2014). There was also a beneficial effect of acupuncture on the improvement of cognitive function for participants with stroke in the convalescent stage (MD 1.34, 95% CI 0.76 to 1.92; participants = 120; studies = 2; $I^2 = 0\%$; Analysis 2.9). The difference in one trial in which cognitive function was evaluated by the Revised Hasegawa Dementia Scale (HDS-R) was significant too (Li 2010a) (MD 1.26, 95% CI 0.29 to 2.23; participants = 46; studies = 1; Analysis 2.9). One trial evaluated the effect of acupuncture on the improvement of cognitive function measured by MMSE at the end of one month's follow-up (Sun 2013a). There was a significant difference between the two groups (MD 3.47, 95% CI 2.43 to 4.51; Analysis 2.10).

Three trials measured improvement of cognitive function after acupuncture treatment by using categorical approaches (MMSE) (Bao 2012; Li 2010a; Zheng 2014). There was no significant

heterogeneity among them. It showed that acupuncture was beneficial for the improvement of cognitive function in participants with stroke in the convalescent stage (OR 3.82, 95% CI 1.89 to 7.72; participants = 166; studies = 3; $I^2 = 0\%$; Analysis 2.11). One trial with 46 participants evaluated this outcome measured by categorical approaches (HDS-R). It also showed that acupuncture was beneficial for the improvement of cognitive function in participants with stroke in the convalescent stage (OR 4.02, 95% CI 1.12 to 14.46; Analysis 2.11).

Depression

Six trials evaluated the effect of acupuncture on the improvement of depression measured by the Hamilton Depression Scale (HAMD) at the end of treatment (Guo 2011; Gao 2014a; Sun 2015; Wu 2008; Zhang 2013a; Zhou 2014). There was a beneficial effect of acupuncture on the improvement of depression for participants with stroke in the convalescent stage (MD -2.58, 95% CI -3.28 to -1.87; participants = 552; studies = 6; $I^2 = 71\%$; Analysis 2.12). One trial measured this outcome by using the Symptoms of Traditional Chinese Medicine (TCM) depression scale (Gao 2014a). There was also a beneficial effect of acupuncture on the improvement of depression for participants with stroke in the convalescent stage (MD -1.57, 95% CI -2.96 to -0.18; Analysis 2.12).

Four trials measured improvement of depression after acupuncture treatment by using categorical approaches (HAMD) (Gao 2014a; Sun 2015; Wu 2008; Zhang 2013a). There was no significant heterogeneity among them. It also showed that acupuncture was beneficial for the improvement of depression in participants with stroke in the convalescent stage (OR 2.03, 95% CI 1.10 to 3.72; participants = 342; studies = 4; $I^2 = 0\%$; Analysis 2.13).

Swallowing function

Three trials evaluated the effect of acupuncture on the improvement of swallowing function at the end of treatment (Guo 2012; Wu 2013a; Xu 2013). Acupuncture was superior to no acupuncture in the improvement of dysphagia measured by a drinking test in two trials (Wu 2013a; Xu 2013) (MD -1.11, 95% CI -2.08 to -0.14; participants = 200; studies = 2; $I^2 = 96\%$; Analysis 2.14). The difference was also significant in another trial in which dysphagia was evaluated categorically by a drinking test (Guo 2012) (OR 95.29, 95% CI 10.93 to 830.86; Analysis 2.15).

Pain

Two trials evaluated the effect of acupuncture on the improvement of pain measured by the Visual Analogue Scale (VAS) (Wang 2011a; Yao 2014). There was a significant decrease in scores among participants in the acupuncture group compared with the control group (MD -2.88, 95% CI -3.68 to -2.09; participants = 118; studies = 2; $I^2 = 0\%$; Analysis 2.16). It showed that acupuncture was superior to no acupuncture in the improvement of pain.

Sleep

One trial evaluated the effect of acupuncture on the improvement of sleep quality measured by the Rhone Planck Sleepiness Scale (Zhang 2013a). There was no significant improvement in sleep quality among participants in the acupuncture group compared with the control group (MD -1.09, 95% CI -2.37 to 0.19; Analysis 2.17).

Spasticity

One trial evaluated the effect of acupuncture on the improvement of spasticity measured by the Modified Ashworth Spasticity Rating Scale (Wu 2011a). There was a beneficial effect of acupuncture on the improvement of spasticity for participants with convalescent stroke (MD -0.40, 95% CI -0.64 to -0.16; participants = 60; studies = 1; Analysis 2.18).

Death from any cause during the whole treatment and follow-up period

No data on death from any cause during the period of treatment and follow-up were available in these 29 trials.

Quality of life at the end of follow-up

One trial reported no significant change in health-related quality of life measured by means of self-report using the Medical Outcomes Study 36-Item Short-Form Health Survey (SF-36) from baseline to one month after treatment among participants in the acupuncture group compared with the control group (Sun 2013a) (MD 2.73, 95% CI -0.54 to 6.00; participants = 71; studies = 1; Analysis 2.19).

Possible adverse events

Adverse events were reported in five trials (Li 2010a; Sun 2013a; Sun 2015; Zhang 2013a; Zhou 2014). Of these, three trials reported

no obvious adverse events related to acupuncture (Li 2010a; Sun 2015; Zhang 2013a). Zhou 2014 reported bleeding, haematoma, and pain at the acupoint in the acupuncture group and Sun 2013a found itchiness of the skin at the acupoint in one participant in the acupuncture group. No other serious adverse events were reported.

Subgroup analysis

We were unable to perform pre-determined subgroup analyses based on time of starting acupuncture and stroke severity because most trials did not provide information for the specified outcomes.

Sensitivity analysis

We were unable to perform the predetermined sensitivity analyses for trials with adequate concealment of randomisation and blinding, and that were published in languages other than Chinese because most trials did not provide the necessary data.

Publication bias

Nine trials comparing acupuncture plus baseline treatment with baseline treatment alone evaluated the effect of acupuncture on the improvement of dependency measured by the Barthel Index. Based on these nine trials, we produced a funnel plot to check for publication bias in this systematic review. We found that the funnel plot was asymmetric (Figure 4).

Figure 4. Funnel plot of comparison 2, Acupuncture + baseline treatment versus baseline treatment alone, outcome: 2.1 Improvement of dependency at the end of treatment.

Although funnel plot asymmetry has long been equated with publication bias, the funnel plot should be seen as a generic means of displaying small-study effects. Small-study effects may be due to factors other than publication bias, such as poor methodological quality leading to spuriously inflated effects in smaller studies, true heterogeneity, artefactual, and chance (Sterne 2011).

DISCUSSION

Summary of main results

We included 31 trials in this review, with a total of 2257 participants in the subacute or chronic stage of stroke. Two trials were conducted outside China (Chou 2009; Naeser 1992) and the remaining 29 trials were conducted in China. The present review on acupuncture for stroke is also not representative of different racial groups.

The results revealed the following.

1. Compared with no acupuncture, for people with stroke in the subacute or chronic stage, acupuncture may have a beneficial effect on improving dependency (activities of daily living); global neurological deficiency; and specific neurological impairments including motor function, cognitive function, depression, swallowing function, pain, and spasticity.
2. Acupuncture was not superior to sham acupuncture or no acupuncture on improving the quality of life in people with stroke in the subacute or chronic stage.
3. There were no serious adverse events reported in people with stroke in the convalescent stage using acupuncture.
4. Currently there is no evidence from the included RCTs for the effect of acupuncture on death, requiring institutional care, and all-cause mortality in people with stroke in the subacute or chronic stage.

Overall completeness and applicability of evidence

In this systematic review, a substantial number of the included studies were conducted in China and were published in Chinese. Our electronic searching successfully identified studies for which an abstract was available in Chinese, as well as a number of studies based on English titles. However, we believe it is likely that we will not have identified all relevant Chinese trials, in particular those for which only English titles were available and those not published in journals included in the electronic databases that we searched. The asymmetric funnel plot further proved that there was existing publication bias in this systematic review (Figure 4).

Many of the relevant trials that we included were published only as brief reports. This was frequently the case for studies published in Chinese, for which published versions were often less than two pages long. Although we contacted study authors, when possible, to confirm study eligibility, we did not have the time or resources to contact all study authors for further information on trial design or study results. Thus, in general the completeness of study information is low, resulting in a high number of studies for which risk of bias is classed as 'unclear'.

Characteristics of participants

The studies included in this review were predominantly conducted in China, so the review is not representative of different racial groups. The type and severity of stroke may alter the effects

of acupuncture on people with stroke. From the available information, it was not possible to perform pre-specified subgroup analyses comparing people with different severities of stroke, and different times to start of treatment after stroke. This was due firstly, to the limited number of studies for the specified outcomes, secondly, to only some of the included trials reporting data on severity, and thirdly, to most trials not clearly defining the interval between stroke onset and the start of the intervention treatment, and including participants with a mix of different intervals from stroke onset to start of intervention treatment.

Properties of interventions

The quality of acupuncture treatment is closely related to its effectiveness. Misleading results may have occurred if the treatment schedules were inadequate or administered by unskilled practitioners. However, information on the experience and training of the acupuncturists who gave the treatments was available in only one trial (Chou 2009). Furthermore, the acupuncture techniques, the number of acupoints, the number and duration of sessions, and the duration of the intervention period varied across trials. Some trials reported that the acupuncture points, the number of sessions, and the duration of treatment were individualised according to the practical conditions in each stroke patient. From the scarce description of treatment in all trials, it is difficult to evaluate if the acupuncture treatment was valid or not. The consensus of an international group of experienced acupuncturists and researchers was that clinical trials of acupuncture must use an optimal form of treatment, defined by examining standard texts and by surveying and consulting experts, and must be reported by using Revised Standards for Reporting Interventions in Clinical Trials of Acupuncture (STRICIA) (Hugh 2010).

Outcomes measures

Efficacy

Because post-stroke neurological recovery is known to be obvious within three months, especially post-stroke motor recovery, and unlikely beyond six months, mixing people with interval from stroke onset of less than three months, three to six months, and more than six months in one sample would make the assessment of the efficacy of an intervention methodologically unsound. From the available information, it was not possible to perform pre-specified subgroup analyses comparing people with different times to start of treatment (within three months, three to six months, and after six months) for each outcome. This was due to the limited amount of data, the varied outcome measurements, or both, in this systematic review.

The long-term goal of treatment for stroke is to reduce mortality and disability, and ultimately to prolong survival and improve quality of life. Data available from the included trials were mainly secondary outcomes. There was a lack of data from RCTs on clinically relevant outcomes at long-term follow up, such as mortality and quality of life. The lack of reliable evidence on primary outcomes meant that we were unable to draw conclusions about the influence of acupuncture on stroke.

Adverse events

Of the 31 included trials with a total of 2257 participants in the convalescent phase of stroke, only one participant was reported to have suffered from bleeding and three participants were reported

to have suffered from intolerance to pain at acupoints. There were no serious adverse events reported. From the available evidence, it appears that acupuncture is a safe treatment when used for people with stroke. However, it should be noted that this result was based on insufficient information from five trials with a small number of participants (Li 2010a; Sun 2013a; Sun 2015; Zhang 2013a; Zhou 2014). Most trials in this systematic review did not report whether any adverse events relevant to acupuncture were apparent in their participants. The reasons for insufficient reporting of adverse events were possibly as follows. Firstly, Chinese practitioners perceive acupuncture as free of side effects, secondly, because these are rare events, data from RCTs will almost never be sufficient to prove or disprove a causal relationship between a complementary and alternative medicine (CAM) therapy and a rare adverse event, and thirdly, study authors reported positive effects without reporting adverse events as well. However, some studies demonstrated that serious adverse events have been associated with acupuncture, such as infections (HIV, hepatitis, subacute bacterial endocarditis) caused by non-sterile needles, or complications (pneumothorax, cardiac tamponade) caused by tissue trauma, but the incidence of adverse events was unknown (Ernst 1997).

Quality of the evidence

The current available evidence shows that acupuncture may have beneficial effects on the improvement of dependency, global neurological deficiency, and specific neurological deficiency in some dimensions. Unfortunately due to low methodological quality with regard to methods of randomisation, allocation concealment and blinding of assessment of most included trials, and probable publication bias, there is currently insufficient evidence to support the routine use of acupuncture for people with stroke in the convalescent stage. The general low methodological quality of the included trials of acupuncture for stroke prohibited meaningful sensitivity analyses to illuminate the robustness of the results of the review to the exclusion of those trials with inadequate methodology. It was also not possible to perform sensitivity analysis to confirm the robustness of the results of the review to the exclusion of Chinese trials, because only two non-Chinese trials were included. However, this systematic review provides comprehensive and updated information on the effects of acupuncture for stroke in the convalescent stage for clinical practice. Therefore, further research on acupuncture for stroke is worthwhile in future.

Potential biases in the review process

The major limitations of this review are related to the weakness inherent in the available published literature on acupuncture for stroke. While most studies reported global and specific neurological deficit results, the measurement of the outcomes varied considerably across trials. The combined results from studies with such wide variation were unreliable. Furthermore, many studies did not provide detailed information on the severity of the disease, the exact time of starting the acupuncture treatment, the modalities of acupuncture technique

and acupuncturist's background, and possible adverse effects relevant to acupuncture treatment. This brings the generalisability of the results into question.

Another limitation of this systematic review is that publication bias might be present, as indicated by the asymmetric funnel plot for the effect of acupuncture on the improvement of dependency measured by the Barthel Index. Although we undertook extensive literature searches, we still could not exclude the possibility that studies with negative findings remain unpublished.

AUTHORS' CONCLUSIONS

Implications for practice

Although acupuncture may have positive effects in stroke rehabilitation and there were no reported serious adverse events, the small number of low quality studies and the probability of publication bias means that there was insufficient evidence to support the routine use of acupuncture for people with subacute or chronic stroke.

Implications for research

The widespread use of acupuncture, the promising results with less severe side effects, lower cost, and the insufficient quality of the available trials warrant further research. Large sham or placebo-controlled trials are needed to confirm or refute the available evidence. The following features should be addressed in further studies.

1. Detailed reporting of the generation of the allocation sequence and allocation concealment.
2. Application and clear description of blinding.
3. Use of placebo or sham acupuncture as the control.
4. Clear definition of the modality of acupuncture, and acupuncture technique based on evidence or a consensus of experts (STRICTA).
5. Use of standard validated outcome measures.
6. Reporting of clinically important outcome measures at long-term follow-up, such as mortality and quality of life.
7. Adverse events critically assessed by standardised monitoring or an effective self-report system. Attention should be paid to rare, severe adverse events relevant to acupuncture.
8. The study should be reported according to the STRICTA criteria (Hugh 2010) in conjunction with the CONSORT statement (Schulz 2010).

ACKNOWLEDGEMENTS

We thank Mrs Hazel Fraser for providing us with relevant trials and systematic reviews from the Cochrane Stroke Group Trials Register and Mrs Brenda Thomas for her help with developing the search strategy and helpful comments. We express our gratitude to Dr Valentina Assi, Dr Bo Wu, lead editor Prof Peter Langhorne, Julie Gildie, and Tam Watson for their very helpful comments.

REFERENCES

References to studies included in this review

Bao 2012 {published data only}

Bao Y. Clinical study on mild cognitive impairment after stroke treated with cord-supplementing and brain strengthening acupuncture [益髓健脑针刺法治疗脑梗死后轻度认知功能障碍的临床研究]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10228-1012467639.htm> (accessed 8 August 2015) 2012.

Chou 2009 {published data only (unpublished sought but not used)}

Chou P, Chu H, Lin J. Effects of electroacupuncture treatment on impaired cognition and quality of life in Taiwanese stroke patients. *Journal of Alternative & Complementary Medicine* 2009;**15**(10):1067-73.

Dai 1997 {published data only}

Dai CY, Lu F. Efficacy observation on the treatment of sequelae of cerebral infarction in 46 patients by integrating Chinese with modern medicine. *Practical Journal of Integrating Chinese with Modern Medicine* 1997;**10**(1):438.

Gao 2014a {published data only}

Gao X. Clinical observation on treating poststroke depression by scalp acupuncture and back shu points [针刺头穴结合五脏背俞穴治疗中风后抑郁的临床疗效观察]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10228-1014250907.htm> (accessed 8 August 2015) 2014.

Guo 2011 {published data only}

Guo AS, Li AH, Chen X, Cai JY, Wu QF. Effects of electric-acupuncture and fluoxetine on depression and neurological function of post-stroke depression patients [电针联合氟西汀对卒中后抑郁症患者抑郁状态及神经功能的影响]. *Shandong Medical Journal* 2011;**51**(25):9-11.

Guo 2012 {published data only}

Guo LJ, Yang GF. Clinical research of acupuncture treatment combined with popsicle stimulus in the treatment of dysphagia after apoplexy [针刺配合冰棒刺激治疗中风后吞咽困难的临床研究]. *China Journal of Chinese Medicine* 2012;**27**(7):918-9.

Huang 2008a {published data only}

Huang GF, Zhang HX, Zhang TF. Effects of scalp acupuncture combined with rehabilitation therapy on motor function in the hemiplegic patient of cerebral thrombosis at convalescent period [头针结合康复治疗对脑血栓形成恢复期偏瘫患者运动功能的影响]. *Chinese Acupuncture & Moxibustion* 2008;**28**(8):573-5.

Ke 2015 {published data only}

Ke J, Wang JH, Li CE. Acupuncture by linguiba method combined with rehabilitation for 40 patients after stroke [针刺灵龟八法配合康复锻炼治疗中风40例]. *Henan Traditional Chinese Medicine* 2015;**35**(1):153-5.

Li 1997a {published data only}

Li X. Efficacy analysis of rehabilitation in the treatment of hemiplegia in 112 stroke patients [112例中风偏瘫患者分组治

疗的康复疗效分析]. *Chinese Journal of Rehabilitative Theory and Practice* 1997;**3**(1):22-4.

Li 2010a {published data only}

Li W. Efficacy observation on the effects of acupuncture in the treatment of mild cognitive impairment after stroke [针刺对脑卒中后轻度认知功能障碍的疗效观察]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10026-2010145185.htm> (accessed 8 August 2015) 2010.

Li 2011a {published data only}

Li M. Clinical research on the treatment of cognitive impairment after stroke by pericardium acupuncture [针刺手厥阴心包经特定穴治疗脑卒中后认知障碍的临床研究]. Available at: <http://cdmd.cnki.com.cn/article/cdmd-10228-1011222586.htm> (accessed 8 August 2015) 2011.

Li 2013a {published data only}

Li XZ, Liu YX. Observation on the effects of acupuncture and rehabilitation in the treatment of shoulder-hand syndrome due to stroke [针刺结合康复训练分期治疗肩手综合征的观察]. *Chinese Journal of Gerontology* 2013;**33**(21):5247-8.

Liu 2013a {published data only}

Liu J, Feng XD. Clinical observation of treating cognitive impairment after stroke by electroacupuncture at Baihui and Shenting with cognitive rehabilitation training [电针百会、神庭穴配合康复训练治疗脑卒中后认知障碍临床研究]. *China Journal of Chinese Medicine* 2013;**28**(4):608-10.

Lun 1999 {published data only}

Lun X, Peng ZF, Peng SZ. Clinical observation on the treatment of sequelae of stroke by needling temporal three points [颞三针为主治疗中风后遗症的临床观察]. *Journal of Clinical Acupuncture and Moxibustion* 1999;**15**(4):8-9.

Naeser 1992 {published data only}

Naeser MA, Alexander MP, Stiassny-Eder D, Galler V, Hobbs J, Bachman D. Real versus sham acupuncture in the treatment of paralysis in acute stroke patients: a CT scan lesion site study. *Journal of Neurorehabilitation* 1992;**6**(4):163-73.

Sun 2013a {published data only}

Sun YT. Clinical study of post-stroke vascular cognitive impairment treatment with galvano-acupuncture and cognitive training [电针结合认知训练治疗梗死后血管性认知障碍的临床研究]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10572-1013226538.htm> (accessed 8 August 2015) 2013.

Sun 2015 {published data only}

Sun YT, Bao YH, Wang SL, Chu JM, Li LP. Efficacy observation on the treatment of post-stroke depression by acupuncture at the acupoints based on ziwuliuzhu and Prozac [子午流注取穴针刺配合百忧解治疗脑卒中后抑郁疗效观察]. *Chinese Acupuncture & Moxibustion* 2015;**35**(2):119-22.

Wang 2001 {published data only}

Wang ZY. Compared analyses among Chinese medicinal herbs, acupuncture and combination of them in the treatment of

hemiplegia in stroke patients [中药、针刺、针药并举对脑卒中偏瘫康复的疗效对比分析]. *Modern Rehabilitation* 2001;**5**(6):130.

Wang 2011a {published data only}

Wang X, Wang WQ. Observation on the treatment of fire needle therapy in 43 patients with shoulder-hand syndrome after stroke [火针疗法治疗肩手综合征43例]. *Guangming Journal of Chinese Medicine* 2011;**26**(4):754-6.

Wang 2012 {published data only}

Wang J, Cui X, Ni HH, Huang CS, Zhou CX, Wu Y, et al. Observation on the effects of interactive scalp acupuncture on motor function in patients with stroke [互动式头针治疗对脑卒中患者运动功能的影响]. *Chinese Journal of Rehabilitation Medicine* 2012;**27**(10):941-3.

Wu 2008 {published data only}

Wu XM, Wang B. Efficacy observation on the treatment of depression caused by stroke in 40 patients by acupuncture and fluoxetine [针刺配合百忧解口服治疗脑卒中后抑郁障碍40例疗效观察]. *Chinese Journal of Difficult and Complicated Cases* 2008;**7**(6):357-8.

Wu 2011a {published data only}

Wu W. Efficacy observation on the treatment of post-stroke strephenopodia by acupuncture and rehabilitation training [针刺配合康复训练治疗中风偏瘫侧足内翻疗效观察]. *Shanghai Journal of Acupuncture - Moxibustion* 2011;**30**(5):321-3.

Wu 2013a {published data only}

Wu YL, Wang L, Tuo S, Yu X, Wang Q. Clinical study on the effects of acupuncture kinesitherapy for dysphagia caused by pseudobulbar paralysis after stroke [互动式针刺法对卒中后假性延髓麻痹吞咽困难的疗效]. *Chinese Journal of Rehabilitation Medicine* 2013;**28**(8):739-42,757.

Xu 2013 {published data only}

Xu JY, Zhou ZL, Wu J. Clinical observation on the treatment of post-stroke dysphagia by Tiaoshen Tongluo Acupuncture combined with Tongue 3-needle and acupuncturing Double Yifeng Acupoints [调神通络针法结合舌三针及针刺双侧翳风穴治疗脑卒中后吞咽困难临床观察]. *Journal of Zhejiang University of Traditional Chinese Medicine* 2013;**37**(9):1117-8, 1132.

Yao 2014 {published data only}

Yao WJ. Clinical research on the effects of huici acupuncture combined with rehabilitation in the treatment of upper extremity dysfunction due to stroke [恢刺法结合康复训练对脑卒中后上肢功能障碍影响的临床研究]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10315-1014409892.htm> (accessed 8 August 2015) 2014.

Zhan 2014 {published data only}

Zhan LX. Acupuncture and moxibustion in combination with rehabilitation in the treatment of senile cerebral apoplexy hemiplegia: random parallel control study [针灸联合康复训练治疗老年脑卒中偏瘫随机平行对照研究]. *Journal of Practical Traditional Chinese Internal Medicine* 2014;**28**(1):134-6.

Zhang 2013a {published data only}

Zhang ZR. Clinical research on treating post-stroke depression by electric acupuncture shenwuxing [电针神五星治疗中风后抑郁的临床研究]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10633-1014171987.htm> (accessed 8 August 2015) 2013.

Zhang 2015 {published data only}

Zhan WY. Clinical research on the effects of scalp acupuncture combined with rehabilitation in the treatment of hemiplegia due to cerebral thrombosis [头针联合康复治疗脑血栓形成偏瘫患者的临床研究]. *Journal of Yanan University (Medical Sciences)* 2015;**13**(1):35-7.

Zheng 2014 {published data only}

Zheng Q. Clinical study on the treatment of mild cognitive impairment after stroke by yishentiaodu acupuncture [益肾调督针法治疗脑卒中后轻度认知功能障碍的临床研究]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10228-1014250904.htm> (accessed 8 August 2015) 2014.

Zhou 2014 {published data only}

Zhou YF. Acupuncture for poststroke depression: a single-center randomized controlled trial [针刺治疗缺血性脑卒中后抑郁的随机对照试验]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10344-1015546952.htm> (accessed 8 August 2015) 2014.

Zhu 2007 {published data only}

Zhu XJ, Gao WB, Yang XY. Clinical observation on the effects of acupuncture combined with rehabilitative exercises in the treatment of shoulder subluxation after stroke [针刺结合康复训练治疗中风后肩关节半脱位的临床观察]. *Journal of Beijing University of Traditional Chinese Medicine* 2007;**14**(3):23-5.

References to studies excluded from this review

Bai 2007 {published data only}

Bai L, Jin GY. Observation on the effects of acupuncture combined with rehabilitation on the neurologic function after stroke [针刺联合现代康复治疗脑梗死神经功能缺损的临床研究]. *Chinese Archives of Traditional Chinese Medicine* 2007;**25**(4):838-9.

Bai 2011 {published data only}

Bai ZQ. Observation on the effects of acupuncture on aphasia following ischemic stroke [针刺治疗脑梗塞失语症疗效的观察]. *Chinese Journal of Ethnomedicine and Ethnopharmacy* 2011;**20**(3):125.

Bai 2013 {published data only}

Bai YL, Li L, Hu YS. Prospective, randomized controlled trial of physiotherapy and acupuncture on motor function and daily activities in patients with ischemic stroke. *Journal of Alternative and Complementary Medicine* 2013;**19**:684-9.

Bao 2008 {published data only}

Bao J, Pan XH. Clinical observation of acupuncture combined with rehabilitation in the treatment of dysphagia after stroke [针刺加康复训练对脑卒中吞咽障碍的疗效观察]. *Journal of Hebei Medical University* 2008;**29**(6):876-7.

Bao 2010 {published data only}

Bao ZY, Kang P, Zhu HQ, Yang CY. Efficacy observation on the treatment of pseudobulbar palsy due to stroke by auditory evoked potential acupuncture [听觉诱发电位对针刺治疗中风后假性球麻痹疗效评定]. *Chinese Journal of Misdiagnosis* 2010;**10**(36):8857-8.

Bao 2012a {published data only}

Bao YH, Wang YW, Chu JM, Zhu GX, Wang CM, Hou HM. Observation of the curative effect of electrical acupuncture combined with rehabilitation on shoulder pain following hemiplegia [电针结合康复对中风后偏瘫肩痛患者疼痛及上肢运动功能改善作用的临床观察]. *Chinese Archives of Traditional Chinese Medicine* 2012;**19**(1):59-60.

Cai 2010 {published data only}

Cai SH, Yan LH, Nie B. Clinical observation on the treatment of apoplexy hemiplegia by electrical acupuncture at the back shu points and the front mu points [俞募穴电针治疗中风偏瘫临床观察]. *Clinical Journal of Traditional Chinese Medicine* 2010;**22**(2):124-6.

Calabro 2011 {published data only}

Calabro S. Scalp acupuncture for stroke. *Acupuncture Today* 2011; Vol. 12:37.

Cao 2010a {published data only}

Cao RL, Yang GQ, Huang LP, Zhang ZC. Acupuncture at tongue tri-point and ren ying point for swallowing dysfunction after stroke [舌三针及人迎穴为主治疗脑卒中后吞咽功能障碍]. *Shanxi Journal of Traditional Chinese Medicine* 2010;**31**(1):89.

Cao 2010b {published data only}

Cao XQ. Acupuncture combined with dialectical treatment for 54 cases with pseudobulbar palsy after stroke [辨证论治配合针刺治疗假性球麻痹54例]. *Chinese Journal of Traditional Chinese Medicine* 2010;**22**(8):718.

Chang 2010 {published data only}

Chang JL, Gao Y, Li SL, Wang J, Cao KG, Ma B. Effect of acupuncture and speech rehabilitation on motor aphasia after stroke [针刺配合语言康复治疗脑卒中后运动性失语]. *Chinese Journal of Rehabilitation Theory and Practice* 2010;**16**(1):58-9.

Chao 2009 {published data only}

Chao WB. Clinical study on treatment of aphasia due to apoplexy by acupuncture [针灸治疗中风失语症临床研究]. *Chinese Journal of Basic Medicine in Traditional Chinese Medicine* 2009;**15**(8):619,625.

Chau 2009 {published data only}

Chau AC, Cheung RT, Jiang X. Acupuncture of motor-implicated acupoints on subacute stroke patients: an fMRI evaluation study. *Medical acupuncture* 2009;**21**(4):233-41.

Che 2002 {published data only}

Che JL. Clinical observation on the treatment of sequelae of stroke in 40 patients with relatively long duration by needling abdominal points. *Jiangshu Traditional Chinese Medicine* 2002;**23**(1):31-2.

Chen 2000 {published data only}

Chen ZF, Jiang GH, Lai XS. Clinical observation on the effects of electronic acupuncture treating vascular dementia [电针治疗血管性痴呆临床疗效观察]. *Journal of Clinical Acupuncture and Moxibustion* 2000;**16**(12):18-20.

Chen 2007 {published data only}

Chen W, Gu HW, Liu XQ, Yu Q, Liu SH, Liu HL. Observation on the effects of neurological function in patients with ischemic stroke by needling Zusanli and Xuanzhong points [针刺足三里、悬钟对缺血性脑卒中神经功能的影响及其临床意义]. *Chinese Journal of Rehabilitation Medicine* 2007;**22**(12):1110-1.

Chen 2008a {published data only}

Chen AL, Li XP, Yu JL, Shi WH, Zhou J, Wang W. Effect of nerve trunk acupuncture combined with rehabilitation on hand function of stroke patients [针刺神经干及康复训练对脑卒中患者手功能的影响]. *Chinese Journal of Rehabilitation Medicine* 2008;**23**(6):416.

Chen 2008b {published data only}

Chen M. Results of acupuncture on the dysphagia due to cerebral stroke [针刺治疗对脑卒中吞咽障碍康复的效果]. *China Tropical Medicine* 2008;**8**(12):2177, 2194.

Chen 2009 {published data only}

Chen YQ. Clinical study with sEMG on efficacy of acupuncture in combination with rehabilitation in relieving the foot drop of the patients with stroke [针灸配合康复治疗对中风偏瘫后足下垂疗效的表面肌电研究]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10315-2009252155.htm> (accessed 8 August 2015) 2009.

Chen 2010a {published data only}

Chen X, Chen YF, Mao M, Mou X, Rao P, Bai Y. Clinical research on the effects of acupuncture therapy combined with contemporary rehabilitation techniques for hemiplegia and spasticity caused by apoplexy [针刺结合现代康复技术治疗中风偏瘫痉挛状态的临床研究]. *Journal of Chengdu University of TCM* 2010;**33**(1):27-9.

Chen 2010b {published data only}

Chen YY. Clinical observation on the effects of acupuncture combined with rehabilitation for patients with limb spasticity after stroke [针刺配合康复疗法对中风病肢体痉挛患者的临床观察]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10572-2010126355.htm> (accessed 8 August 2015) 2010.

Chen 2012a {published data only}

Chen HY. Clinical observation on Quanjing acupuncture therapy combined with rehabilitation training for treatment of hemiplegia and spasticity caused by apoplexy [全经针刺法结合康复训练治疗恢复期中风偏瘫痉挛状态的临床观察]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10541-1012401427.htm> (accessed 8 August 2015) 2012.

Chen 2012b {published data only}

Chen YH, Jiang TZ, Zhang B, Huang W, Liu N, Cao SJ. Clinical study on acupuncture in the treatment of swallowing dysfunction from apoplexy [针刺治疗脑梗死后吞咽障碍

临床研究]. *Clinical Journal of Traditional Chinese Medicine* 2012;**24**(11):1067-9.

Chen 2013 {published data only}

Chen LF. Acupuncture for young ischemic stroke: a controlled randomized trial. ChiCTR Chinese Clinical Trial Register (<http://www.chictr.org/>) (ChiCTR-TRC-13003317) (accessed 8 August 2015) 2013.

Chen 2014 {published data only}

Chen CY, Ge LB, Xu MS. Clinical study on the evaluation of acupuncture in treating ischemic stroke of recovery stage mostly based on static equilibrium [以静态平衡为主评估针刺治疗缺血性中风恢复期临床疗效]. *Shanghai Journal of Acupuncture and Moxibustion* 2014;**33**(4):291-5.

Cheng 2006 {published data only}

Cheng Y, Tang QS. Clinical random control study of acupuncture on treating stagnation of liver-qi with deficiency of the spleen syndrome of gerontic depressive disorder after stroke [针刺治疗老年期中风后郁证肝郁脾虚候的随机对照临床研究]. *Journal of TCM Brain Diseases* 2006;**2**(2):129-32.

Cheng 2007 {published data only}

Cheng Y, Tang QS. Abdominal acupuncture in treating liver-qi stagnation and spleen deficiency in the elderly with post-stroke depression: A randomized and controlled observation [腹针治疗老年期脑卒中后郁证肝郁脾虚候的随机对照观察]. *Journal of Clinical Rehabilitative Tissue Engineering Research* 2007;**11**(39):7791-4.

Cheng 2011 {published data only}

Cheng XK, Wang ZM, Sun L. Post-stroke hand dysfunction treated with acupuncture at Zhongzhu (TE 3) and Waiguan (TE 5) [针刺中渚、外关治疗脑卒中后手功能障碍]. *Chinese Acupuncture & Moxibustion* 2011;**31**(2):117-20.

Chow 2006 {published data only}

Chow ESL, Lau PMY, Chau RMW. Effectiveness of acupuncture treatment in stroke rehabilitation. *Neurorehabilitation and Neural Repair* 2006;**20**:209.

Chu 2007 {published data only}

Chu YJ, Wang CY, Zhang Y. Clinical observation of acupuncture for 72 cases with depression after stroke [针刺治疗脑卒中后抑郁症72例临床观察]. *Chinese Journal of Gerontology* 2007;**27**(9):1720-1.

Chu 2009 {published data only}

Chu GX, Yi XQ. Clinical observation on the treatment of spastic paralysis by electrical acupuncture combined with rehabilitation [电针结合康复治疗中风痉挛性瘫痪的临床观察]. *Hubei Journal of Traditional Chinese Medicine* 2009;**31**(8):13-4.

Cui 1992 {published data only}

Cui YY, Zhao JH, Huang WC, Wang AP. Observation on the effects of acupuncture in the treatment of compulsive crying and laughing due to multiple cerebral infarction [针刺治疗多发性脑梗塞性强哭强笑症的疗效观察]. *Journal of Acupuncture and Moxibustion* 1992;**1**:19-20.

Ding 2000 {published data only}

Ding BY, Cui YJ. Observation on the effects of acupuncture and cupping in the treatment of arthritic spasm in 52 stroke patients. *Shanghai Journal of Acupuncture and Moxibustion* 2000;**19**(3):28.

Dong 2007a {published data only}

Dong G. The optimized scheme of acupuncture treatment on symptom based stroke sequelae and clinical assessment on generic technology. ChiCTR Chinese Clinical Trial Register (<http://www.chictr.org/Site/English/Index.htm>) (ChiCTR-TRC-08000209) (accessed 8 August 2015) 2007.

Dong 2007b {published data only}

Dong JP, Sun WY, Wang S, Wu ZQ, Liu F. Clinical observation on the effects of head point-through-point electroacupuncture for treatment of poststroke depression [头部电针透穴治疗脑卒中后抑郁症临床观察]. *Chinese Acupuncture & Moxibustion* 2007;**27**(4):241-4.

Dong 2008a {published data only}

Dong G. Clinical curative effect and safe research on treatment of haemorrhagic stroke with stage acupuncture. ChiCTR Chinese Clinical Trial Register (<http://www.chictr.org/>) (ChiCTR-TRC-08000225) (accessed 8 August 2015) 2008.

Dong 2008b {published data only}

Dong JL, Xi GM, Zhou SH. Study of acupuncture Lianquan point combined with Fengchi point for 32 patients with pseudobulbar palsy after stroke [风池配合廉泉穴针灸治疗脑卒中后假性延髓性麻痹32例]. *Chinese Journal of Practical Nervous Diseases* 2008;**11**(1):38-9.

Dong 2009 {published data only}

Dong Y, Liu G, Wang T. Tongdu Kaiyin acupuncture for 30 patients with aphasia after stroke [通督开暗针法治疗脑卒中失语症30例]. *Journal of Anhui Traditional Chinese Medical College* 2009;**28**(1):34-6.

Du 2013 {published data only}

Du GY. Observation on the effects of acupuncture for 100 cases with postapoplectic pseudobulbar palsy [针灸治疗中风后假性球麻痹100例]. *Journal of Medical Theory and Practice* 2013;**25**(1):40-1.

Duan 2010 {published data only}

Duan YW, Sun XJ, Feng X, Xu JX. Observation of 40 patients with postapoplectic pseudobulbar palsy treated by acupuncture [针刺治疗脑卒中后假性球麻痹40例]. *Yunnan Journal of Traditional Chinese Medicine and Materia Medica* 2010;**31**(10):45-6.

Duan 2014 {published data only}

Duan JY, Ding BY, Zong L. Clinical observation of acupuncture at the thirteen ghost Aacupoints for post-stroke anxiety and depression [针刺十三鬼穴为主治疗脑卒中焦虑抑郁共病临床观察]. *Shanghai Journal of Acupuncture and Moxibustion* 2014;**33**(6):536-8.

Fang 2009 {published data only}

Fang GF, Liu JD. Effect of acupuncture combined with function training for post-stroke dysphagia [针刺配合功能训练治疗脑卒中后吞咽障碍疗效分析]. *Modern Medicine Journal of China* 2009;**11**(11):54-7.

Feng 1996a {published data only}

Feng SL, Guo ZQ, Xie GR. Observation on the clinical effects of hemiplegia from apoplexy treated by needling Feng Fu, Ya Men points. *Journal of Hunan College of Traditional Chinese Medicine* 1996;**16**(3):61-4.

Feng 1996b {published data only}

Feng SL, Guo ZQ, Xie GR. Clinical observation and experimental studies on the treatment of cerebral thrombus by needling "Feng Fu, Ya Men" points. *Journal of Guangzhou University of Traditional Chinese Medicine* 1996;**13**(2):20-3.

Feng 2011 {published data only}

Feng X. Acupuncture for 68 cases with supranuclear paralysis caused by stroke [针刺翳风穴配合咽后壁点刺治疗假性延髓麻痹68例]. *Guangming Journal of Traditional Chinese Medicine* 2011;**26**(5):1007.

Fink 2004 {published data only}

Fink M, Rollnik JD, Bijak M, Borstdt C, Duper J, Guergueltcheva V, et al. Needle acupuncture in chronic poststroke leg spasticity. *Archives of Physical Medicine and Rehabilitation* 2004;**85**(4):667-72.

Fu 2010 {published data only}

Fu WB, Guo Y, Chen XK. Comprehensive therapeutic protocol of electroacupuncture combined with Chinese herbs and rehabilitation training for treatment of cerebral infarction: a multi-center randomized controlled trial [电针联合中药、康复训练综合方案治疗脑梗死：多中心随机对照研究]. *Chinese acupuncture & moxibustion* 2010;**30**(1):6-9.

Fu 2013 {published data only}

Fu LX. Acupuncture treatment of ischemic stroke basal ganglia aphasia clinical curative effect evaluation: a randomized controlled study. ChiCTR Chinese Clinical Trial Register (<http://www.chictr.org/>) (ChiCTR-TRC-12003168) (accessed 8 August 2015) 2013.

Fu 2014 {published data only}

Fu ZY, Li XR, Wu QM, Li DD, Liu WA. Clinical observation on acupuncture and Chinese medicine combined with stroke unit for treatment of ischemic stroke at convalescence stage [针刺和中药结合卒中单元治疗缺血性脑卒中恢复期患者的临床研究]. *Chinese Journal of Integrative Medicine on Cardio/Cerebrovascular Disease* 2014;**12**(1):48-51.

Gao 2001 {published data only}

Gao HY, Yan LF, Liu BB. Clinical study on treatment of senile vascular dementia by acupuncture. *Journal of Traditional Chinese Medicine* 2001;**21**(2):103-9.

Gao 2008 {published data only}

Gao CY. Clinical observation on the effects of rehabilitation combined with acupuncture in the treatment of dysphagia after

stroke [脑卒中后吞咽困难康复训练加针刺治疗疗效观察]. *Chinese Journal of Gerontology* 2008;**28**(12):2486-7.

Gao 2011 {published data only}

Gao GY. Clinical research on the treatment of hemiplegia after stroke by scalp acupuncture and Brunnstrom technique [头针配合Brunnstrom促通技术治疗缺血性脑卒中后偏瘫患者的临床研究]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10507-1012253202.htm> (accessed 8 August 2015) 2011.

Gao 2014b {published data only}

Gao ZQ. Research on the treatment of hemiplegia after stroke by optimized techniques of acupuncture and moxibustion [针灸康复优化技术治疗中风后偏瘫的研究]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10369-1014334635.htm> (accessed 8 August 2015) 2014.

Gong 2008 {published data only}

Gong H. The clinical investigation of ultrasound, laser therapy and acupuncture for patients with cerebral infarction [超声、激光加针灸治疗脑梗塞的临床研究]. Available at: <http://cdmd.cnki.com.cn/article/cdmd-10422-2008189373.htm> (accessed 8 August 2015) 2008.

Gu 2007 {published data only}

Gu HW, Ma WP, Chen W, Li QS, Liu XQ, Liu SH. Treatment of ischemic stroke by acupuncture at Zusanli and Xuanzhong [针刺足三里和悬钟治疗缺血性脑卒中]. *Journal of Clinical Rehabilitation Tissue Engineering Research* 2007;**11**(25):4931-3.

Gu 2009 {published data only}

Gu YC. Clinical research on the effects of acupuncture at Jinjinyue point combined with speech therapy for broca aphasia after stroke [针刺金津玉液结合语言康复训练治疗中风后运动性失语的临床研究]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10393-2009201641.htm> (accessed 8 August 2015) 2009.

Guan 2009a {published data only}

Guan CX, Guo GH, Yu L. Observation on the effects of acupuncture plus deglutition training in the treatment of dysphagia caused by stroke [针刺配合吞咽训练治疗脑卒中后吞咽障碍的疗效观察]. *Journal of Medical Forum* 2009;**30**(4):19-23.

Guan 2009b {published data only}

Guan CY. Observation on therapeutic effect of Youxuan acupuncture and rehabilitation exercise in the treatment of hemiplegia after apoplexy [针刺优选法结合现代康复治疗脑卒中后偏瘫临床疗效观察]. Available at: <http://cdmd.cnki.com.cn/article/cdmd-10760-2009119611.htm> (accessed 8 August 2015) 2009.

Guan 2013 {published data only}

Guan C. Clinical research on the effects of acupuncture at heel vessel points and rehabilitation training in the treatment of strephenopodia after cerebral infarction [针刺跷脉穴结合康复训练治疗脑梗死后足内翻的临床研究]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10541-1013183193.htm> (accessed 8 August 2015) 2013.

Guo 1999 {published data only}

Guo XY, Wei YF. Clinical observation on the effects of acupuncture in the treatment of muscular spasm in 124 stroke patients. *Chinese Acupuncture and Moxibustion* 1999;**9**:553.

Guo 2006a {published data only}

Guo XL. Effects of scalp acupuncture combined with function training on the gait of stroke patients [头针结合康复训练对脑卒中后患者偏瘫步态的影响]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10393-2006162983.htm> (accessed 8 August 2015) 2006.

Guo 2006b {published data only}

Guo Y. Therapeutic effect observation of acupoint blocking on shoulder joint subluxation of post-apoplexy [穴位封闭治疗中风后肩关节半脱位疗效观察]. *Shanxi Journal of Traditional Chinese Medicine* 2006;**22**(3):39-40.

Guo 2013 {published data only}

Guo QS. Clinical observation on the treatment of stroke hemiplegia by acupuncture combined with Chinese medicine [针刺联合中药治疗中风偏瘫临床观察]. *Journal of Liaoning University of TCM* 2013;**15**(1):187-9.

Han 2008 {published data only}

Han P. Effects of scalp acupuncture and occupation therapy on motor functions of the upper limb in patients with stroke [头针结合作业疗法对脑卒中患者上肢运动功能及生活自理能力影响的临床对照研究]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10393-2008207891.htm> (accessed 8 August 2015) 2008.

Han 2010 {published data only}

Han XY, Liu Y. Clinical observation of acupuncture combined with rehabilitation in the treatment of dysphagia after stroke [针刺结合康复护理治疗脑卒中后吞咽困难的临床研究]. *China New Technologies and Products*. 2010;**4**:15.

Han 2011 {published data only}

Han DX, Zhuang LX, Zhang Y. Evaluation on efficacy of Jin's "Sanzhen" therapy combined with rehabilitation training for hemiplegia of stroke patients by Fugl-Meyer scale [用Fugl-Meyer量表评价靳三针结合康复训练对脑梗死偏瘫的疗效]. *Acupuncture research* 2011;**36**(3):209-14.

Han 2012 {published data only}

Han N. Acupuncture combined with rehabilitation training for swallowing dysfunction following stroke [针刺配合康复训练治疗中风后吞咽功能障碍]. *Journal of Community Medicine* 2012;**10**(9):32-3.

Han 2014 {published data only}

Han J, Wu Y. Post-stroke patients with dysarthria treated by acupuncture at neck points, tongue acupuncture plus speech rehabilitation training: a randomized parallel controlled study [项针、舌针联合言语康复训练治疗脑卒中后构音障碍随机平行对照研究]. *Journal of Practical Traditional Chinese Internal Medicine* 2014;**28**(5):135-7.

Hang 2014 {published data only}

Hang FJ, Fu XY, Wang TT, Dai DC, Tang CL, He XG. Effects of oral placement therapy combined with acupuncture of specific points on dysphagia after stroke [口腔定位疗法联合特定穴位针刺对脑卒中后吞咽障碍的影响]. *Chinese Journal of Rehabilitation Medicine* 2014;**29**(12):1129-33.

He 2008 {published data only}

He YJ. Clinical study on treating urinary incontinence after stroke by acupuncture [针灸治疗脑卒中后尿失禁的临床研究]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10572-2008100424.htm> (accessed 8 August 2015) 2008.

Hegyi 2012 {published data only}

Hegyi G, Szigeti G. Rehabilitation of stroke patients using Yamamoto New Scalp Acupuncture: a pilot study. *Journal of Alternative & Complementary Medicine* 2012;**18**(10):971-7.

Hong 2013 {published data only}

Hong YB, Cheng HY, Wang GL, Xia SW, Liu H, Liu HL. Clinical research of special acupuncture therapy by TCM differentiation on improving quality of life in patients of ischemic stroke [辨证应用特色针刺疗法对缺血性脑卒中患者生活质量的改善]. *Journal of Medical Research* 2013;**42**(2):41-5.

Hou 1998 {published data only}

Hou AL, Wang L, Pu Y. Clinical observation of treatment of multi-infarctional dementia by herbs, oxygen and acupuncture. *Shanghai Journal of Acupuncture and Moxibustion* 1998;**17**(2):12-3.

Hsing 2012 {published data only}

Hsing W, Imamura M, Weaver K, Fregni F, Azevedo Neto RS. Clinical effects of scalp electrical acupuncture in stroke: a sham-controlled randomized clinical trial. *Journal of Alternative & Complementary Medicine* 2012;**18**(4):341-6.

Hu 2011 {published data only}

Hu JK, Hu KM, Hu T. Clinical research of acupuncture combined with speech training for spasmodic dysphonia after apoplexy [针刺结合语言训练治疗脑卒中痉挛型构音障碍的临床研究]. *Journal of Sichuan of Traditional Chinese Medicine* 2011;**29**(2):117-9.

Hu 2012 {published data only}

Hu SL. Balanced muscular tension needling method with Shenqifutan capsule treatment of cerebral infarction recovering from clinical observation [张力平衡针法配合参芪复瘫胶囊治疗脑梗塞恢复期的临床观察]. Available at: <http://cdmd.cnki.com.cn/article/cdmd-10541-1012401619.htm> (accessed 8 August 2015) 2012.

Huang 2002 {published data only}

Huang HS, Huang X. Effects of "Chuan Xiong Qin and Huang Qi" on the treatment of upper and lower limbs pain due to stroke in 50 patients. *Research of Traditional Chinese Medicine* 2002;**18**(1):9-10.

Huang 2008b {published data only}

Huang F, Liu Y, Zhou FX. Effects of acupuncture on vascular cognitive impairment after stroke [针刺对脑梗死后血管性认知障碍的影响]. *Guangdong Medical Journal* 2008;**29**(11):1918-20.

Huang 2008c {published data only}

Huang F. Clinical observation on puncturing mainly back-shu points for treating blood-stasis blocking collateral type of ischemic apoplexy [针刺背俞穴为主治疗气虚血瘀型缺血性中风的临床观察]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10572-2008100417.htm> (accessed 8 August 2015) 2008.

Huang 2011 {published data only}

Huang ZJ, Chen JX, Chang ZY. Clinic analysis of half-length-flabbiness after hemiplegia stroke treated by acupuncture combined with rehabilitation training [针灸配合康复训练治疗脑卒中后半身不遂的临床研究]. *China Journal of Chinese Medicine* 2011;**26**(9):1149-50.

Huang 2012 {published data only}

Huang GW. Clinical study of acupuncture treatment for fecal and urine incontinence after stroke [针刺治疗中风后大小便失禁的临床研究]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10315-1012402005.htm> (accessed 8 August 2015) 2012.

Huang 2014 {published data only}

Huang LL. Randomized controlled clinical study on Shenque Fuxi Bagua acupuncture therapy in treating constipation after stroke [神阙伏羲八卦针刺疗法治疗中风后便秘的随机对照临床研究]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10633-1015543271.htm> (accessed 8 August 2015) 2014.

Jia 2012 {published data only}

Jia CJ, Ni GX, Tan H, Zhang X. Clinical observation on the effects of acupuncture combined with rehabilitation in the treatment of stage-IV shoulder-hand syndrome from apoplexy [康复训练结合针灸治疗脑卒中后肩手综合征IV期临床疗效观察]. *Journal of Changchun University of Traditional Chinese Medicine* 2012;**28**(4):711-2.

Jiang 1998 {published data only}

Jiang GH, Li YH, Yang WH. Clinical observation on treatment of sequelae of stroke with CT localization acupuncture. *Shanghai Journal of Acupuncture and Moxibustion* 1998;**17**(2):6-7.

Jiang 2000 {published data only}

Jiang ZY, Li CD. Needling paravertebral points in treatment of post-stroke thalamic pain. *International Journal of Clinical Acupuncture* 2000;**11**(1):7-10.

Jiang 2006 {published data only}

Jiang H, Wen XY, Shi TH, Zhang SM. Efficacy observation on the treatment of depression due to lacunar infarction by electroacupuncture [电针刺激治疗腔隙性脑梗死后抑郁状态的疗效观察]. *Chinese Journal of Physical Medicine and Rehabilitation* 2006;**28**(12):807-10.

Jiang 2009 {published data only}

Jiang D, Zhang L, Li B. Acupuncture for 27 cases with dysphagia after stroke [针刺治疗假性延髓麻痹吞咽障碍27例]. *Chinese Journal of Clinical Healthcare* 2009;**12**(5):518-9.

Jiang 2010 {published data only}

Jiang Y, Zhu JG, Xiong J. Clinical observation of consciousness restoring obstruction clearing needling technique in the treatment of subacute cerebral hemorrhage [醒脑开窍针刺法治疗亚急性期脑出血的临床观察]. *Chinese Journal of Ethnomedicine and Ethnopharmacy* 2010;**19**(18):89-90.

Jiang 2011a {published data only}

Jiang L, Zhao J. Acupuncture combined with drugs in the treatment of post-stroke depression [针灸结合药物治疗脑卒中后抑郁的临床研究]. *Journal of Clinical Acupuncture and Moxibustion* 2011;**27**(11):15-7.

Jiang 2011b {published data only}

Jiang YJ. Electroacupuncture DU20 and DU24 treatment on cognitive impairment after stroke [电针百会、神庭对脑卒中后认知功能障碍的影响]. Available at: <http://cdmd.cnki.com.cn/article/cdmd-10393-1011220783.htm> (accessed 8 August 2015) 2011.

Jiang 2012 {published data only}

Jiang L. Clinical research on acupuncture treatment for pseudobulbar paralysis after stroke [针刺治疗脑卒中后假性延髓麻痹的临床研究]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10228-1012468023.htm> (accessed 8 August 2015) 2012.

Jiang 2013 {published data only}

Jiang P, Zheng ZY. Clinical observation on the treatment of hemiplegia spasticity by acupuncture antagonist muscle corresponding acupoints and Bobath therapy [针刺拮抗肌相应穴位结合Bobath疗法治疗偏瘫痉挛状态的临床观察]. *Journal of Clinical Acupuncture and Moxibustion* 2013;**29**(1):18-20.

Jin 1993 {published data only}

Jin R, Lai XS, Li YH, Liang RA. Clinical observation on the treatment of sequelae of stroke with three temporal needling. *Chinese Acupuncture and Moxibustion*. 1993;**2**:11-2.

Jin 2010a {published data only}

Jin CX. Clinical research of acupuncture combined with rehabilitation for upper limb spasticity induced by stroke [针刺结合康复治疗中风偏瘫上肢痉挛状态的临床研究]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-87801-2010263051.htm> (accessed 8 August 2015) 2010.

Jin 2010b {published data only}

Jin RJ, Zhu TM, Wang Q. Effects of electric acupuncture at acupoint of antagonistic muscles combined with facilitation techniques on motor function and activity of daily living in patients with hemiplegia caused by cerebral infarction [电针拮抗肌腧穴复合易化技术对脑梗塞后偏瘫患者运动功能及日常生活能力的影响]. *Journal of Chengdu University of Traditional Chinese Medicine* 2010;**33**(3):9-12.

Kang 2011 {published data only}

Kang JJ. Clinical study on the effects of electroacupuncture on GV20 and EX-HN1 on stroke patients with cognitive impairment [电针百会及四神聪治疗脑卒中认知障碍患者的临床疗效观察]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10393-1011220779.htm> (accessed 8 August 2015) 2011.

Kang 2013 {published data only}

Kang GF, Chen JH, Li TL, Wu QM, Yang RF. Efficacy of Shen-qi Fu-tan capsule and tension balance acupuncture on cerebral infarction at convalescence: a clinical observation of 30 cases [参芪复瘫胶囊联合张力平衡针法治疗脑梗死恢复期30例临床观察]. *Guiding Journal of Traditional Chinese Medicine and Pharmacy* 2013;**19**(12):14-6.

Kjendahl 1997 {published data only}

Kjendahl A, Sallstrom S, Osten PE, Stanghelle JK. A one year follow-up study on the effects of acupuncture in the treatment of stroke patients in the subacute stage: a randomized, controlled study. *Clinical Rehabilitation* 1997;**11**:192-200.

Lai 1997a {published data only}

Lai XS. Observation on the effects of acupuncture in the treatment of senile vascular dementia. *Chinese Acupuncture and Moxibustion*. 1997;**4**:201-2.

Lai 1997b {published data only}

Lai XS, Zhang JW, Muo FZ, Jiang GH, Li JQ, Zheng ZC. Clinical observation on the effects of acupuncture in the treatment of vascular dementia. *Chinese Acupuncture and Moxibustion*. 1997;**12**:713-6.

Lai 1998 {published data only}

Lai XS, Muo FZ, Jiang GH, Li JQ, Chen JL, Kuang ZY. Observation of clinical effect of acupuncture on vascular dementia and its influence on superoxide dismutase, lipid peroxide and nitric oxide. *Chinese Journal of Integrated Traditional and Western Medicine* 1998;**18**(11):648-51.

Lai 2004 {published data only}

Lai XS, Liu JY, Jiang GM. Effect of combination of acupuncture treatment at Liench' tian and plum blossom needle treatment at the tip of tongue on aphasia after stroke and its significance on hemorheology. *Chinese Journal of Clinical Rehabilitation* 2004;**8**(19):3818-20.

Lai 2011 {published data only}

Lai J. Clinical research of acupuncture combined with rehabilitation therapy for dysarthria after stroke [针刺结合康复治疗脑卒中后构音障碍的临床研究]. Available at: <http://cdmd.cnki.com.cn/article/cdmd-10633-1011155206.htm> (accessed 8 August 2015) 2011.

Lang 2013 {published data only}

Lang JY, Zhuang LX, He J, Jia C, Zhou SH, Ke LP. Randomized controlled study on Jin's three needle therapy on spastic hemiplegia after ischemic stroke ["靳三针"疗法治疗缺血性中风后痉挛性偏瘫随机对照研究]. *Shanghai Journal of Acupuncture and Moxibustion* 2013;**32**(6):440-3.

Lao 2013 {published data only}

Lao LZ. Clinical research on motility dysarthria after stroke treated with acupuncture combining with rehabilitation training [针刺结合康复训练治疗脑卒中后运动性构音障碍的临床研究]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10572-1013229348.htm> (accessed 8 August 2015) 2013.

Lee 2007 {published data only}

Lee SW, Yun JM, Son JW, Kang BG, Park SM, Yun H J. The effect of electroacupuncture on upper-extremity spasticity of stroke patients. *Journal of Korean Oriental Medicine* 2007;**28**:492-501.

Lee 2009 {published data only}

Lee SY, Baek YH, Park SU. Intradermal acupuncture on Shen-Men and Nei-Kuan acupoints improves insomnia in stroke patients by reducing the sympathetic nervous activity: a randomized clinical trial. *American Journal of Chinese Medicine* 2009;**37**(6):1013-21.

Lei 2013 {published data only}

Lei YQ, Li RL, Lin QF, Han ZY, Wu CH, Huang XH. Zengshi Taichi acupuncture for 65 patients with pseudobulbar palsy after stroke [曾氏太极针法治疗脑卒中后假性球麻痹65例]. *Liaoning Journal of Traditional Chinese Medicine* 2013;**40**(4):780-1.

Li 1993 {published data only}

Li YH, Jin R. Three forms of acupuncture in the treatment of sequelae of cerebrovascular accident. *Chinese Acupuncture and Moxibustion* 1993;**4**:35-8.

Li 1994a {published data only}

Li CD, Huang Y, Li YK, Hu KM, Jiang ZY. Treating post-stroke depression with "mind-refreshing antidepressive" acupuncture therapy: a clinical study of 21 cases. *International Journal of Clinical Acupuncture* 1994;**5**(4):389-93.

Li 1994b {published data only}

Li YH, Jin R. Clinical study in the treatment of sequelae of cerebrovascular accident with needling temporal points. *Acupuncture Research* 1994;**19**(2):4-6.

Li 1997b {published data only}

Li YH, Yang WH, Zhuang LX, Zheng L. Clinical study of acupuncture in the treatment of vascular dementia. *Acupuncture Research* 1997;**3**(1):22-4.

Li 2006a {published data only}

Li CY, Fan YZ, Xue WH, Tian Y, Yan L, Ni AH. Clinical study on Baihui(GV 20) acupuncture in treating cerebral infarction [针刺百会穴治疗脑梗塞的临床研究]. *Shanghai Journal of Traditional Chinese Medicine* 2006;**40**(6):19-20.

Li 2006b {published data only}

Li GC, Zhao X. Clinical observation on treatment of apoplectic aphasia by acupuncture plus rehabilitation [针刺配合康复治疗脑卒中失语的临床观察]. *Shanghai Journal of Acupuncture and Moxibustion* 2006;**25**(9):13-4.

Li 2006c {published data only}

Li J, Wang Y, Liu B. Observation on the curative effects of acupuncture combined with rehabilitation on postapoplectic shoulder pain [针灸结合康复手法治疗脑卒中后肩痛的疗效观察]. *Chinese Journal of Trauma and Disability Medicine* 2006;**14**(1):37-8.

Li 2006d {published data only}

Li T, Tian GJ, Zhang L, Wei GW, Ma Q, Han JX. Clinical study of acupuncture on the neurological function in postoperative patients with extensive cerebral infarction [针刺对大面积脑梗塞术后病人神经功能的影响]. *Shanghai Journal of Acupuncture and Moxibustion* 2006;**25**(9):3-5.

Li 2006e {published data only}

Li T, Zhao JG, Tian GJ, Zhang L, Liu SJ. Clinical observation on the effects of acupuncture on neurological function in patient after operation of hypertensive cerebral hemorrhage [针刺对高血压脑出血术后病人神经功能的影响]. *Chinese Acupuncture & Moxibustion* 2006;**26**(4):247-9.

Li 2008a {published data only}

Li JA, Li JL. Clinical observation on the effects of acupuncture in the treatment of dysphagia induced by ischemic stroke [针刺治疗缺血性中风后吞咽困难的临床观察]. *Journal of Beijing University of Traditional Chinese Medicine* 2008;**15**(4):17-9.

Li 2008b {published data only}

Li P, Sun GF, Yao GM, Yan RY, Xia MH, Zhang WJ. Clinical observations on the treatment of dysphagia induced by pseudobulbar palsy by acupuncture plus rehabilitation therapy [针刺联合康复治疗脑卒中后吞咽障碍的临床观察]. *Shanghai Journal of Acupuncture and Moxibustion* 2008;**27**(5):10-1.

Li 2008c {published data only}

Li SW, Zhang ZX. Efficacy observation on the rehabilitation of stroke by acupuncture [针刺在脑梗死康复治疗中的应用价值观察]. *Journal of Zhejiang University of Traditional Chinese Medicine* 2008;**32**(4):514-5.

Li 2008d {published data only}

Li T. Clinical observation of acupuncture combined with language recovery training for 32 patients with motor aphasia after apoplexy [针刺配合语言训练治疗中风后运动性失语32例临床观察]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10199-2009021115.htm> (accessed 8 August 2015) 2008.

Li 2009a {published data only}

Li F. Clinical study on the treatment of post-stroke depression by acupuncture combined with fluoxetine [针刺结合百忧解治疗中风后抑郁症的临床观察]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10507-2009158081.htm> (accessed 8 August 2015) 2009.

Li 2009b {published data only}

Li Y, Zhang J, Chen YJ, Pan FQ, Xiao ZX, Tian JY. Research on the effects of neuro-facilitation technique combined with acupuncture on activities of daily living in stroke patients [神经促通技术联合针刺对脑卒中偏瘫患者日常生活活动能力的影响]. *Chinese Journal of Rehabilitation Medicine* 2009;**24**(12):1106-8.

Li 2010b {published data only}

Li CW. Effects of acupuncture on swallowing function in patients with stroke [针刺对脑卒中患者吞咽功能的影响]. *Journal of Emergency Traditional Chinese Medicine* 2010;**19**(9):1609.

Li 2010c {published data only}

Li HY, Li LF, Li Y. Clinical research on the effects of acupuncture combined with rehabilitation in the treatment of dysphagia after stroke [针刺联合康复训练对脑卒中后假性球麻痹吞咽困难干预作用的临床研究]. *Guangming Journal of Chinese Medicine* 2010;**25**(5):830-1.

Li 2011b {published data only}

Li PF, Pan YX, Wang Y, Sun PY, Pu F, Zhang YL. Clinical research on the effects of acupuncture combined with rehabilitation training in the treatment of dysphagia after stroke [针刺配合吞咽康复训练治疗脑卒中后吞咽障碍的临床研究]. *Journal of Clinical Acupuncture and Moxibustion* 2011;**27**(9):1-5.

Li 2011c {published data only}

Li WL. Clinical study on the effects of electroacupuncture combined with rehabilitation therapy for hemiplegia after ischemic stroke [电针结合康复训练治疗缺血性中风后偏瘫的临床研究]. *Jiangsu Journal of Traditional Chinese Medicine* 2011;**43**(9):68-9.

Li 2011d {published data only}

Li Y. Acupuncture treatment for motor aphasia after stroke [针刺治疗中风后运动性失语症]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10507-1012252976.htm> (accessed 8 August 2015) 2011.

Li 2012a {published data only}

Li D. Efficacy observation on treating swallow dysfunction after stroke by acupuncture combined with deglutition training [针灸配合吞咽训练对脑卒中后吞咽障碍的疗效观察]. *Clinical Journal of Chinese Medicine* 2012;**4**(23):38-9.

Li 2012b {published data only}

Li FF. Clinical effect of early rehabilitation combined with acupuncture therapy on stroke patients with dysphagia [早期康复结合针刺治疗脑卒中吞咽障碍临床疗效观察]. *Journal of New Chinese Medicine* 2012;**44**(6):147-8.

Li 2012c {published data only}

Li F, Tang JY, Cheng HL, Hong W, Yang J. Rehabilitation effect evaluation of tri-tongue-acus and cold stimulus add tongue muscle training for stroke patients with swallowing disorder [针刺加冷刺激配合舌肌训练对脑卒中吞咽障碍患者的康复疗效评价]. *China Journal of Traditional Chinese Medicine and Pharmacy* 2012;**27**(9):2471-4.

Li 2012d {published data only}

Li G, Zhang Y, Luo HQ. Clinical observation on the curative effect of acupuncture combined with estazolam for postapoplectic insomnia [针刺联合艾司唑仑治疗脑卒中后失眠的疗效观察]. *Medical Journal of National Defending Forces in Southwest China* 2012;**22**(6):641-2.

Li 2012e {published data only}

Li J. Clinical observation of acupuncture combined with rehabilitation for 35 patients with sequela of apoplexy [针刺结合康复治疗中风后遗症35例临床观察]. *Chinese Journal of Integrative Medicine On Cardio-/Cerebrovascular Disease* 2012;**10**(6):697-8.

Li 2012f {published data only}

Li Q. Clinical research of treating dysphagia after stroke by acupuncture [针刺治疗脑卒中后吞咽障碍的临床研究]. *Clinical Journal of Chinese Medicine* 2012;**4**(12):37-8.

Li 2012g {published data only}

Li SG, Zhou QS. Clinical research of acupuncture combined with massage for hemiplegia after stroke [针刺配合推拿治疗中风后偏瘫的临床研究]. *Modern Journal of Integrated Traditional Chinese and Western Medicine* 2012;**21**(23):2554-5.

Li 2013b {published data only}

Li F, Cai RL, Zhai L. Impacts of acupuncture at Jing-well points on the differentiated meridians and temple-three-needle therapy on P300 of patients with early vascular cognitive impairment [辨经刺并配合颞三针对早期血管性认知障碍患者P300的影响]. *Chinese Acupuncture & Moxibustion* 2013;**33**(9):784-8.

Li 2013c {published data only}

Li LH. Clinical observation on Quanjing acupuncture therapy combined with rehabilitation training treatment of convalescent the stroke hemiplegia movement dysfunction [全经针刺法结合康复训练治疗恢复期中风偏瘫运动功能障碍的疗效观察]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10541-1013183196.htm> (accessed 8 August 2015) 2013.

Li 2014 {published data only}

Li YG, Liu Q. Effect of acupuncture combined with neuromuscular electrical stimulation for postapoplectic dysphagia [针刺配合神经肌肉电刺激治疗对脑卒中后吞咽功能障碍的影响]. *Journal of Ningxia Medical University* 2014;**36**(4):450-2.

Li 2015 {published data only}

Li J, Hu KM, Yang L, Qu Y, Luo L, Li DH. Clinical observation on 90 cases of dysarthria after stroke treated with acupuncture and speech rehabilitation [针刺结合言语康复治疗脑卒中后构音障碍90例临床观察]. *Journal of Traditional Chinese Medicine* 2015;**56**(4):300-3, 310.

Liang 1993 {published data only}

Liang R. Clinical observation and experimental studies on the treatment of sequelae of stroke by needling temporal points. *International Journal of Clinical Acupuncture* 1993;**4**(1):19-26.

Liang 2010 {published data only}

Liang JL. Acupuncture combined with pulse electric acupuncture for 60 patients with aphasia after apoplexy [针刺加脉冲电针治疗中风后失语60例]. *Guangxi Journal of Traditional Chinese Medicine* 2010;**33**(2):30-1.

Liang 2012 {published data only}

Liang K. Clinical research on the effects of acupuncture in the treatment of pseudobulbar palsy due to cerebral infarction [针刺治疗脑梗死后假性延髓麻痹的临床疗效研究]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10228-1012467877.htm> (accessed 8 August 2015) 2012.

Liao 2006a {published data only}

Liao HW. Clinical observations on the efficacy of occupational therapy plus acupuncture for treating reflex sympathetic dystrophy [作业疗法合针灸治疗肩手综合征45例疗效观察]. *Shanghai Journal of Acupuncture and Moxibustion* 2006;**25**(3):9-10.

Liao 2006b {published data only}

Liao HW. Application of scalp acupuncture combined with language training in treatment of aphasia following stroke [头针结合语言治疗在卒中后失语症中的应用]. *Chinese Journal of Rehabilitation Medicine* 2006;**10**(11):115-6.

Liao 2013 {published data only}

Liao XD. Clinical observation of fire-needle therapy combined with Bobath therapy on spastic hemiplegia after stroke [火针点刺结合Bobath疗法治疗中风痉挛性偏瘫临床疗效观察]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10572-1013225433.htm> (accessed 8 August 2015) 2013.

Liao 2014 {published data only}

Liao SQ, Jiang Z, Zhuo LP, Huang SE. Clinical research on the effects of acupuncture at "Quchi" and "Waiguan" acupoints on hand function recovery after stroke [针刺"曲池" "外关"穴位促进脑卒中手功能恢复临床研究]. *Asia-Pacific Traditional Medicine* 2014;**10**(22):60-2.

Lim 2014 {published data only}

Lim CS, Park SK, Sun SH, Lee KH. Research on Korean Pharmacopuncture in South Korea since 2007. *Journal of Pharmacopuncture* 2014;**17**(4):15-21.

Lin 2010a {published data only}

Lin H, Ding XJ, Fu B. Effect of acupuncture combined with medications on postapoplectic cognitive impairment [针灸配合药物治疗对脑卒中后认知障碍的影响]. *Modern Journal of Integrated Traditional Chinese and Western Medicine* 2010;**19**(1):36-7.

Lin 2010b {published data only}

Lin ZL, Zhen J, Li FQ. Effects of acupuncture on recovery of depression after stroke [针刺对脑卒中后抑郁康复疗效的影响]. *China Prac Med* 2010;**5**(8):59-60.

Lin 2012 {published data only}

Lin JW. A clinical research on the effects of acupuncture at the Sun 13 ghost-points for 60 cases with post-stroke depression [针刺孙氏十三鬼穴治疗卒中后抑郁疗效观察]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10572-1012396179.htm> (accessed 8 August 2015) 2012.

Lin 2013 {published data only}

Lin ZX. Effectiveness of acupuncture for the treatment of post-stroke dysphagia: a pragmatic trial comparing real acupuncture treatment with waiting list. ChiCTR Chinese Clinical Trial Register (<http://www.chictr.org/>) (ChiCTR-TRC-12002621) (accessed 8 August 2015) 2013.

Liu 1998 {published data only}

Liu J, Peng XH, Lin DD, Li CD, Jiang ZY, Zeng L, et al. Clinical study of electrical scalp acupuncture in the treatment of

vascular dementia. *Chinese Acupuncture and Moxibustion* 1998;**4**:197-200.

Liu 2004a {published data only}

Liu GF, Zhou ZX. Effect of neuromuscular electrical stimulation in various modes on the rehabilitation of upper limb function in patients with hemiplegia. *Chinese Journal of Clinical Rehabilitation* 2004;**8**(22):4424-5.

Liu 2004b {published data only}

Liu HG, Huang SJ, Zhang HL. Effect of Jiawei simiao yongan tang combined with electroacupuncture in the recovery of motor function in patients with post-stroke sequelae. *Chinese Journal of Clinical Rehabilitation* 2004;**8**(10):1916-7.

Liu 2006 {published data only}

Liu LA, Guo RY, Pang SM, Li XZ, Dai SQ, Li XY. Clinical research of acupuncture combined with speech training for early phase of aphasia after stroke [针刺结合语言训练治疗中风早期失语症的临床研究]. *Chinese Archives of Traditional Chinese Medicine* 2006;**24**(12):2256-9.

Liu 2008a {published data only}

Liu J. Clinical observation on treatment of spasticity poststroke by electric acupuncture with motion point and yang meridian [电针治疗中风后痉挛状态的临床研究]. Available at: <http://cdmd.cnki.com.cn/article/cdmd-87801-2008138509.htm> (accessed 8 August 2015) 2008.

Liu 2008b {published data only}

Liu W, Mukherjee M, Sun C, Liu H, McPeak LK. Electroacupuncture may help motor recovery in chronic stroke survivors: a pilot study. *Journal of rehabilitation research and development* 2008;**45**(4):587-95.

Liu 2009 {published data only}

Liu CS. Clinical research of acupuncture combined with neuromuscular facilitation treatment for dysphagia after stroke [针刺复合易化技术治疗脑卒中后吞咽障碍的临床研究]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10633-2010077373.htm> (accessed 8 August 2015) 2009.

Liu 2010a {published data only}

Liu G, Lv CS, Yuan LX. Study on electroacupuncture improving the single-foot supporting phase rate of stroke patients [电针改善中风患者单足支撑相比例的研究]. *Chinese Acupuncture & Moxibustion* 2010;**30**(6):446-8.

Liu 2010b {published data only}

Liu WA, Wu QM, Li XR, Li DD, Fu L, Yi XC. Observation on the efficacy of combined treatment of stroke hemiplegia with scalp electroacupuncture and stroke unit [头电针配合卒中单元综合疗法治疗脑卒中偏瘫疗效观察]. *Shanghai Journal of Acupuncture and Moxibustion* 2010;**29**(3):149-51.

Liu 2011a {published data only}

Liu FC. Clinical observation on the effects of scalp acupuncture plus motor therapy in the treatment of post-stroke depression [头针配合运动疗法治疗脑卒中后抑郁疗效观察]. *Shanghai Journal of Acupuncture and Moxibustion* 2011;**30**(11):733-4.

Liu 2011b {published data only}

Liu Y. Observation on the treatment of hemiplegia due to stroke in 37 patients by Buyang Huanwu soup and acupuncture [补阳还五汤结合针刺治疗中风偏瘫37例临床观察]. *Guiding Journal of Traditional Chinese Medicine and Pharmacy* 2011;**17**(12):57-9.

Liu 2012a {published data only}

Liu HR, Li YF, Wu HQ, Xu LM, Xin JW, Jiang YN. Observation on the effects of acupuncture combined with swallowing & speech therapeutic instrument for dysphagia in patients after stroke [针灸结合吞咽言语诊治仪治疗脑卒中后吞咽障碍]. *China Pratical Medicine* 2012;**7**(24):49-50.

Liu 2012b {published data only}

Liu SD. Curative effect analysis of speech training and acupuncture therapy for dysarthria caused by stroke [言语训练结合针刺治疗脑卒中构音障碍的临床研究]. *Internal Medicine of China* 2012;**7**(4):336-8.

Liu 2012c {published data only}

Liu Y, Xue Z, Wu L. Clinical observation on 60 cases of hemiplegia treated with acupuncture. *International Journal of Clinical Acupuncture* 2012;**21**:16-8.

Liu 2013c {published data only}

Liu X, Wang PQ. Random parallel control study on the effects of acupuncture and rehabilitation combined with western medicine in the treatment of QDBS [针刺与康复训练联合西药治疗气虚血瘀型中风恢复期随机平行对照研究]. *Journal of Practical Traditional Chinese Internal Medicine* 2013;**27**(12):68-70.

Liu 2013d {published data only}

Liu XP. A random parallel control study on the effects of acupuncture combined with western medicine in the treatment of stroke [针灸联合西药治疗中风随机平行对照研究]. *Journal of Practical Traditional Chinese Internal Medicine* 2013;**27**(9):71-2.

Liu 2013e {published data only}

Liu Y, Liu L, Wang XM. Electroacupuncture at points Baliao and Huiyang (BL35) for post-stroke detrusor overactivity. *Neural Regeneration Research* 2013;**8**(18):1663-72.

Liu 2013f {published data only}

Liu ZF, Zhang JQ, Shao SJ, Wang SF. Therapeutic evaluation of post-stroke depression treated by acupuncture [针刺治疗卒中后抑郁的疗效评价]. *Journal of Sichuan of Traditional Chinese Medicine* 2013;**31**(2):111-3.

Liu 2014b {published data only}

Liu L. Effect of elongated needling therapy combined with deglutition training/rehabilitation training in treatment of post-stroke dysphagia [芒针针刺配合康复训练治疗脑卒中吞咽障碍疗效观察]. *Liaoning Journal of Traditional Chinese Medicine* 2014;**41**(11):2442-4.

Liu 2015 {published data only}

Liu LQ, Li HL, Xu JM. Clinical study of electroacupuncture on memory dysfunction after stroke [电针治疗脑卒中后记忆功

能障碍临床研究]. *Journal of Emergency in Traditional Chinese Medicine* 2015;**24**(5):775-7.

Long 2013 {published data only}

Long AG. Observation of clinical curative effect of acupuncture in patients with cerebral infarction [脑梗死患者针灸治疗的临床疗效观察]. *Chinese Journal of Medicine Guide* 2013;**15**(10):1725-6.

Lu 2010 {published data only}

Lu M, Meng L, Peng J. Therapeutic effects of neuromuscular electrical stimulation and electroacupuncture for dysphagia post stroke [神经肌肉电刺激疗法与电针治疗脑卒中后吞咽障碍的疗效对比研究]. *Chinese Journal of Rehabilitation Medicine* 2010;**25**(2):135-8.

Lu 2011a {published data only}

Lu FJ. Clinical observation on the effects of the combination of acupuncture at tendon junction and Bobath therapy for spastic paralysis after stroke [针刺经筋结点结合Bobath疗法治疗中风后痉挛性瘫痪的临床疗效观察]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10228-1011222551.htm> (accessed 8 August 2015) 2011.

Lu 2011b {published data only}

Lu XF. Clinical observation on acupuncture at the eight-confluent acupoints for hemiplegia after cerebral infarction [针刺八脉交会穴为主治疗脑梗死后偏瘫临床观察]. *Shanghai Journal of Acupuncture and Moxibustion* 2011;**30**(12):828-9.

Luo 2007 {published data only}

Luo WP. Clinical research of acupuncture combined with speech training on Broca aphasia following ischemic apoplexy [针刺配合语言训练治疗脑梗塞致运动性失语的临床研究]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10572-2007130686.htm> (accessed 8 August 2015) 2007.

Luo 2008 {published data only}

Luo WP, Tan JL, Huang HY. Clinical observation on treatment of cerebral infarction-induced Broca aphasia by Tiaoshen Fuyin acupuncture therapy combined with language training [调神复音针刺配合语言训练治疗脑梗死致运动性失语疗效观察]. *Chinese Acupuncture & Moxibustion* 2008;**28**(3):171-5.

Luo 2012 {published data only}

Luo YH. Effect of acupuncture treatment on neurological deficit scores of the hemiplegic stroke patients [针灸治疗对脑卒中偏瘫患者神经功能缺损评分的影响]. *Chinese Community Doctors* 2012;**14**(36):184-5.

Lv 2009 {published data only}

Lv H, Liu H, Ni JN. Electroacupuncture in treatment of postapoplectic swallowing dysfunction: a randomized controlled study [电针治疗卒中后吞咽困难的随机对照研究]. *Journal of China-Japan Friendship Hospital* 2009;**23**(4):228-9.

Lv 2014 {published data only}

Lv ST. Observation on the effects of acupuncture combined with rehabilitation therapy for spasticity after stroke [针刺配合康复治疗中风后肌张力增高的临床观察]. Available at:

<http://cdmd.cnki.com.cn/Article/CDMD-10162-1015546410.htm> (accessed 8 August 2015) 2014.

Ma 2011a {published data only}

Ma JQ, Zheng J, Zhu LW, Yang XL, Zhang XH. Effects of acupuncture combined with rehabilitation training on swallow dysfunction after stroke [针刺结合康复训练对脑卒中后吞咽功能障碍的影响]. *Journal of Clinical Acupuncture and Moxibustion* 2011;**27**(2):15-7.

Ma 2011b {published data only}

Ma XM, Li SZ, Cai NS, Yu HB, Zhang ZL. Research on the effects of quality of life in patients with stroke by tiaorentongdu acupuncture [调任通督针刺法改善脑卒中患者生存质量的研究]. *Shandong Journal of Traditional Chinese Medicine* 2011;**30**(6):396-8.

Ma 2013 {published data only}

Ma JL. Acupuncture treatment of post-stroke hemiplegic shoulder pain: a random parallel control study [针灸治疗中风后偏瘫肩痛随机平行对照研究]. *Journal of Practical Traditional Chinese Internal Medicine* 2013;**27**(6):122-3.

Mou 2010 {published data only}

Mou J, Fu LX, Lu YM, Ren YL, He JY, Qi YZ. Clinical observation on the Xinnao Kaiqiao acupuncture plus language rehabilitation training for motor aphasia caused by cerebral infarction: a report of 30 cases [醒脑开窍针刺法配合语言康复治疗脑梗死致运动性失语30例临床观察]. *Journal of Traditional Chinese Medicine* 2010;**51**(5):428-31.

Mu 2007 {published data only}

Mu JP, Zhu ZM, Cheng JM, Zhao DG, Yang DR, Ao JB. The clinical observation on the acupuncture therapy enlivening brain, dredging orifices cooperating with promoting treatment for hemiplegia after stroke [醒脑开窍针刺法配合促通技术治疗脑卒中偏瘫临床观察]. *Journal of Emergency in Traditional Chinese Medicine* 2007;**16**(5):520-1.

Mukherjee 2006 {published data only}

Mukherjee M, McPeak L, Redford J, Sun C, Liu W. The effect of electro-acupuncture on upper limb spasticity in chronic stroke survivors. *Stroke* 2006;**37**:631.

Muo 2001 {published data only}

Muo FZ, Li JQ, Chu L, Lai XS, Zhang JX, Liu SH. The influence of electroacupuncture on intelligence structure in patients with vascular dementia. *Shanghai Journal of Acupuncture and Moxibustion* 2001;**20**(6):6-8.

Ni 2011 {published data only}

Ni HH, Hu YS, Cui X, Wu Y, Huang DQ, Qu PY. Clinical observation on acupuncture combined with rehabilitation training for 20 patients with postapoplectic upper extremity spasticity [针刺结合康复训练治疗脑卒中后上肢痉挛20例临床观察]. *Journal of New Chinese Medicine* 2011;**43**(1):99-101.

Nie 2011 {published data only}

Nie WB. Clinical observation on the treatment of motor aphasia following stroke by Yinao Tiaoshen acupuncture [针灸益脑

调神法治疗脑卒中运动性失语疗效观察]. *Beijing Journal of Traditional Chinese Medicine* 2011;**30**(12):938-9.

Nie 2013 {published data only}

Nie RR, Huang CH. Post-stroke depression treated with acupuncture and moxibustion: an evaluation of therapeutic effect and safety [针灸治疗脑卒中后抑郁症疗效与安全性评价]. *Chinese Acupuncture & Moxibustion* 2013;**33**(6):490-4.

Pan 2013 {published data only}

Pan MZ. Clinical observation of acupuncture therapy on vascular dementia after stroke [针刺头穴为主治疗脑卒中后血管性痴呆临床疗效研究]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10315-1014128145.htm> (accessed 8 August 2015) 2013.

Pang 2010 {published data only}

Pang P, Lai X, Huang Y, Wang S, Wu J. The thoughts on treating post stroke limb spasticity with the four seas acupoints. *International Journal of Clinical Acupuncture* 2010;**19**(2):45-6.

Peng 2007a {published data only}

Peng XJ. Clinic observation of acupuncture combined with rehabilitation training for 30 cases with hemiplegia after stroke [针灸结合康复训练治疗中风偏瘫30例]. *China Naturopathy* 2007;**15**(5):53-4.

Peng 2009 {published data only}

Peng HY, Tan JL, Wang BG, Zhao MH. A clinical study of therapy of electroacupuncture at the temple-three-needle on post-stroke depression. *New Journal of Traditional Chinese Medicine* 2009;**10**:89-90.

Peng 2015 {published data only}

Peng HY. Clinical study on acupuncture combined with recovery massaging for patients with ischemic stroke [针刺结合康复按摩治疗缺血性中风的临床研究]. *Lishizhen Medicine and Materia Medica Research* 2015;**26**(3):664-5.

Qi 2000 {published data only}

Qi GS, Zhao XJ. Clinical observation on the treatment of stroke in rehabilitative stage with "Qing Nao Yi Shui" acupuncture therapy. *Journal of Tianjin College of Traditional Chinese Medicine* 2000;**19**(2):23-5.

Qi 2013 {published data only}

Qi M. Clinical research on the effects of acupuncture in treating 128 patients with dysphagia after stroke [针灸治疗128例卒中后吞咽障碍的临床研究]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10026-1013203840.htm> (accessed 8 August 2015) 2013.

Qi 2013a {published data only}

Qi M, Zhang GM. Efficacy study of acupuncture and moxibustion on dysphagia after stroke [针灸治疗卒中后吞咽障碍疗效研究]. *Beijing Journal of Traditional Chinese Medicine* 2013;**32**(7):512-4.

Qiao 2012 {published data only}

Qiao HF, Zhang QJ, Yuan HF, Zhang N, Wu ZH. Efficacy observation of acupuncture combined with paroxetine in

the treatment of depression after stroke. Available at: <http://cpfd.cnki.com.cn/Article/CPFDTOTAL-ZGKF201207001119.htm> (accessed 8 August 2015). 2012.

Qiu 2011 {published data only}

Qiu TT. A clinical study of acupuncture combined with therapeutic exercise in the treatment of shoulder-hand syndrome after stroke [针刺结合运动疗法治疗脑卒中后肩手综合征的临床研究]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10228-1011222549.htm> (accessed 8 August 2015) 2011.

Qu 1991 {published data only}

Qu RM, Qu XM. Observation on the treatment of sequelae of apoplexy in 105 patients with needling Du Mai points. *Jiang Xi Traditional Chinese Medicine* 1991;**22**(5):32-4.

Ran 2013 {published data only}

Ran XQ, Wu S, Ye DP, Min J. Effects of acupuncture combined with rehabilitation training on lower extremity blood flow on limb function with spastic paralysis after stroke [针刺结合康复训练对中风痉挛性偏瘫患者肢体功能及下肢血流的影响]. *Hubei Journal of Traditional Chinese Medicine* 2013;**35**(10):20-1.

Ren 2009 {published data only}

Ren YF. Clinical research on acupuncture in combination with rehabilitation for postapoplectic spasm [针刺结合康复治疗中风后痉挛的临床研究]. *Guangming Journal of Chinese Medicine* 2009;**24**(7):1313-5.

Ren 2010 {published data only}

Ren XM, Wang T, Wang CM, Yang GF. Clinical study on cerebral infarction treated with Danhong Tongjing Huoluo decoction combined acupuncture [丹红通经活络汤联合针刺治疗脑梗死临床研究]. *World Journal of Integrated Traditional and Western Medicine* 2010;**5**(10):886-8.

Ren 2012 {published data only}

Ren Z, Li XR, Li DD, Yan J, Tan T, Ye HM. Randomized controlled clinical study of the treatment of ischemic stroke (convalescence) with stroke unit combined with the acupuncture [针刺结合卒中单元治疗缺血性脑卒中(恢复期)随机对照临床研究]. *Journal of Practical Traditional Chinese Internal Medicine* 2012;**26**(7):4-6.

Sallstrom 1996 {published data only}

Sallstrom S, Kjendahl A, Osten PE, Stanghelle JH, Borchgrevink CF. Acupuncture in the treatment of stroke patients in the subacute stage: a randomized, controlled study. *Complementary Therapies in Medicine* 1996;**4**:193-7.

Salom-Moreno 2014 {published data only}

Salom-Moreno J, Sánchez-Mila Z, Ortega-Santiago R, Palacios-Ceña M, Truylol-Domínguez S, Fernández-de-las-Peñas C. Changes in spasticity, widespread pressure pain sensitivity, and baropodometry after the application of dry needling in patients who have had a stroke: a randomised controlled trial. *Journal of Manipulative and Physiological Therapeutics* 2014;**37**(8):569-79.

Schaechter 2007 {published data only}

Schaechter JD, Connell BD, Stason WB, Kaptchuk J, Krebs DE, Macklin EA, et al. Correlated change in upper limb function and motor cortex activation after verum and sham acupuncture in patients with chronic stroke. *Journal of Alternative and Complementary Medicine* 2007;**13**(5):527-32.

Seo 2013 {published data only}

Seo YR, Jung WS, Park SU, Moon SK, Park JM, Park JY. The effect of ouhyul herbal acupuncture point injections on shoulder pain after stroke. *Evidence-Based Complementary and Alternative Medicine* 2013;**2013**(2):1-5.

Shang 2008 {published data only}

Shang YJ, Ma CC, Cai YY, Wang DS, Kong LL. Clinical study on acupuncture combined with rehabilitation therapy for treatment of poststroke shoulder-hand syndrome [针刺结合康复治疗中风后肩-手综合征]. *Chinese Acupuncture & Moxibustion* 2008;**28**(5):331-3.

Shao 2012 {published data only}

Shao B. Clinical research on acupuncture in root and end and appearance and substance of kidney meridian associated with language training for Broca aphasia following the stroke [针刺"肾经标本根结"结合语言训练治疗中风后运动性失语的临床研究]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10393-1012510352.htm> (accessed 8 August 2015) 2012.

Shen 2008 {published data only}

Shen HJ, Kang XB, Xie YC, Yang H. Clinical observation of electroacupuncture for 40 cases with dysphagia after stroke [神经肌肉电刺激治疗脑卒中后吞咽困难40例]. *Shanxi Journal of Traditional Chinese Medicine* 2008;**29**(6):709-10.

Shen 2009 {published data only}

Shen SJ. Observation on the effects of acupuncture combined with rehabilitation training for sequela of apoplexy [针灸结合康复治疗中风后遗症的疗效观察]. *Chinese Journal of Ethnomedicine and Ethnopharmacy* 2009;**18**(20):147-8.

Sheng 2011 {published data only}

Sheng GB, Lu FJ, Liu CY, Wang W. The combination of acupuncture at meridian sinew nodal point and rehabilitation for spastic paralysis after stroke [针刺经筋结点结合康复治疗中风后痉挛性瘫痪]. *Journal of Clinical Acupuncture and Moxibustion* 2011;**27**(5):25-6.

Sheng 2013 {published data only}

Sheng GB, Gao C, Yu JK, Gao Y, Liu G. Clinical observation on the effects of electrical scalp acupuncture in the treatment of depression due to stroke [头部电针疗法治疗中风后抑郁的临床疗效观察]. *Information on Traditional Chinese Medicine* 2013;**30**(3):125-6.

Shi 2007 {published data only}

Shi HF, Zheng XH. Effect of scalp acupuncture drawing method on motor function in hemiplegic patients following cerebral thrombosis at convalescent period [头皮针抽提法对脑血栓形成后恢复期偏瘫患者运动功能的影响]. *Integrated Traditional and Western Medicine* 2007;**27**(4):361-3.

Shi 2007a {published data only}

Shi X. Research on the specific effects and the key influential factors of acupoints: research based on the treatment of brain resuscitating acupuncture for cerebral infarction. ChiCTR Chinese Clinical Trial Register (<http://www.chictr.org/>) (ChiCTR-TRC-07000036) (accessed 8 August 2015) 2007.

Song 2010 {published data only}

Song QM. Clinical research on post-stroke vascular cognitive impairment treatment with needling therapy [针刺治疗中风血管性认知障碍的临床观察]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10572-2010125609.htm> (accessed 8 August 2015) 2010.

Sui 2001 {published data only}

Sui MH. Clinical observation on the therapeutic effect of acupuncture of hand-Shaoyang Meridian acupoints for treatment of upper-limb spasm in stroke patients. *Acupuncture Research* 2001;**26**(2):131-3.

Sun 2008 {published data only}

Sun J, Mi Z, Wang H, Xu D, Chen H. Study on therapeutic effect of acupuncture on dysphagia after stroke. *Journal of Rehabilitation Medicine*. 2008:169.

Sun 2010 {published data only}

Sun QR. Clinical research on plum-blossom needle and scalp acupuncture combined with language rehabilitation training for treatment of ischemic aphasia from apoplexy [头针梅花针叩刺结合语言康复训练治疗缺血性中风失语临床研究]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-11919-2010248108.htm> (accessed 8 August 2015) 2010.

Sun 2011 {published data only}

Sun RL. Clinical observation of acupuncture combined with function training for 26 cases with dysphagia after stroke [针刺结合功能训练治疗脑卒中后吞咽困难26例临床观察]. *Journal of Shanxi College of Traditional Chinese Medicine* 2011;**12**(3):39-40.

Sun 2013b {published data only}

Sun YG. Acupuncture combined with rehabilitation training for 50 patients with postapoplectic swallowing dysfunction [针刺配合康复治疗脑卒中后吞咽障碍50例]. *Henan Traditional Chinese Medicine* 2013;**33**(7):1135-6.

Sun 2014 {published data only}

Sun JW. Research of acupuncture treatment technique of post-stroke dysphagia [针灸治疗中风后吞咽障碍技术的研究]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10026-1014242212.htm> (accessed 8 August 2015) 2014.

Suo 2008 {published data only}

Suo DM, Yu T, Liu L, Shi JW, Ding XR, Fan XN. Therapeutic effect of acupuncture for dysphagia after stroke [针刺对中风后吞咽障碍的影响]. *Henan Traditional Chinese Medicine* 2008;**28**(5):57-8.

Tang 2005 {published data only}

Tang XJ, She QQ, Wang YH, Zhang ZL. Efficacy analysis on prognosis of small amounts of thalamic hemorrhage by

acupuncture at early phase [早期针刺治疗对小量丘脑出血预后影响]. *Sichuan Medical Journal* 2005;**26**(11):51-2.

Tang 2012 {published data only}

Tang X, Tang CL, Xu FM, Xie HW, Li LM, Song YE. Effect of scalp acupuncture combined with body acupuncture on limb function in subacute stroke patients [头针联合体针对亚急性期脑卒中患者肢体功能的影响]. *Acupuncture research* 2012;**37**(6):488-92.

Tang 2013 {published data only}

Tang X, Huang L, Xie HW, Xu FM, Li LM, Song YE. Clinical observation on the effects of yin and yang acupuncture on subacute stroke [阴阳经联合针刺治疗亚急性期脑卒中的临床疗效]. *Chinese Journal of Gerontology* 2013;**33**(10):4668-70.

Tang 2013a {published data only}

Tang Yao. A random parallel control study on the effects of acupuncture combined with western medicine for early ischemic stroke hemiplegia [针灸联合西药治疗缺血性中风早期偏瘫随机平行对照研究]. *Journal of Practical Traditional Chinese Internal Medicine* 2013;**27**(4):140-1.

Tian 2014 {published data only}

Tian FL, Li Q, Liu GR, Zheng DS, Chen JM. Effects of yin-yang meridians acupuncture with respiratory reinforcing and reducing manipulation on lower limbs balance function in stroke patients [阴阳对刺呼吸补泻法对脑卒中患者下肢平衡功能的影响]. *Chinese Acupuncture & Moxibustion* 2014;**34**(11):1047-50.

Tong 1997 {published data only}

Tong LG, Li ZG, Ren YX, Yang J, Meng XY, Ma WM, et al. Body quick-needling plus scalp acupuncture in treating apoplexy and its sequelae. *International Journal of Clinical Acupuncture* 1997;**9**(2):133-9.

Tong 2013 {published data only}

Tong S, Su L, Lu HB, Liu JQ. Observation on the efficacy of acupuncture at key acupoints combined with rehabilitation therapy for spasmodic hemiplegia after cerebral infarction [关键点针刺法配合康复治疗脑梗死后痉挛期偏瘫疗效观察]. *Chinese Acupuncture & Moxibustion* 2013;**33**(5):399-402.

Tong 2014 {published data only}

Tong S, Lv HB, Liu JQ. Clinical observation on the curative effect of acupuncture in combination with deglutition training for postapoplectic (pharynx period) swallowing dysfunction [针刺结合吞咽功能训练治疗脑卒中后(咽期)吞咽障碍临床疗效观察]. *Acta Chinese Medicine and Pharmacology* 2014;**42**(2):129-31.

Wan 2013 {published data only}

Wan WR, Wang TL, Cheng SL, Zhao YL, Zhang W, Wu QY, et al. Post-stroke shoulder-hand syndrome treated with acupuncture and rehabilitation: a randomized controlled trial [针刺结合康复治疗中风后肩手综合征: 随机对照研究]. *Chinese Acupuncture & Moxibustion* 2013;**33**(11):970-4.

Wang 2004 {published data only}

Wang MH, Liang B, Wang M, Chen LQ, Wang YS, Li ZY, et al. Clinical study on the treatment of apoplectic hemiplegia by acupuncture plus rehabilitation training. *Shanghai Journal of Acupuncture and Moxibustion* 2004;**23**(4):7-9.

Wang 2006 {published data only}

Wang YH, Ying SG. Observations on the curative effects of acupuncture in the treatment of postapoplectic pseudobulbar paralysis [针刺治疗中风后假性球麻痹疗效观察]. *Shanghai Journal of Acupuncture and Moxibustion* 2006;**25**(6):15-6.

Wang 2007 {published data only}

Wang XY. Effects of the improved acupoints and rehabilitation exercise on locomotor ability of the upper limbs and ability of daily life in the patient of cerebral infarction [改良穴位配合康复训练对脑梗死患者上肢活动能力及日常生活能力的影响]. *Chinese Acupuncture & Moxibustion* 2007;**27**(3):179-81.

Wang 2008 {published data only}

Wang YJ. Clinical research on "Yiqi huoxue, only strengthen the pain" acupuncture combined with rehabilitation training treating shoulder-hand syndrome caused by stroke ["益气活血, 通络止痛"针法联合康复治疗脑卒中后肩手综合征的临床研究]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10162-2009031775.htm> (accessed 8 August 2015) 2008.

Wang 2008a {published data only}

Wang FY, Wang YQ, Li SH, Jv YB. Clinical observation on the treatment of post-stroke hiccup by acupuncture cooperated with compound of xuanfudaizhe soup [针灸配合旋覆代赭汤治疗中风后呃逆的临床观察]. *Chinese Archives of Traditional Chinese Medicine* 2008;**26**(6):1340-1.

Wang 2008b {published data only}

Wang B. Effect of acupuncture combined with cognitive therapy on cerebral-apoplexy-induced function impairment [针刺结合认知训练对老年脑卒中患者认知障碍及残疾状况的影响]. Available at: <http://cdmd.cnki.com.cn/article/cdmd-10507-2008118250.htm> (accessed 8 August 2015) 2008.

Wang 2009 {published data only}

Wang JJ, Li YX. Clinical effect observation of ice stimulation combined with acupuncture for dysphagia after stroke [冷刺激联合干扰电治疗脑卒中后吞咽困难的效果观察]. *Nursing and Rehabilitation* 2009;**8**(11):940-1.

Wang 2010a {published data only}

Wang LC, Wang ZY, Li BD, Ge JY. Study on electro-acupuncture on points combined with rehabilitation training for treatment of balance function disorder in cerebral infarction [电针背俞穴联合康复训练治疗脑梗死病人平衡功能障碍的对照研究]. *Chinese Journal of Integrative Medicine On Cardio-/Cerebrovascular Disease* 2010;**8**(11):1322-3.

Wang 2010b {published data only}

Wang TL. Effects of acupuncture combined with rehabilitation therapy on shoulder-hand syndrome after apoplexy [针刺结合康复治疗中风后肩手综合征的疗效观察]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10393-2010251306.htm> (accessed 8 August 2015) 2010.

Wang 2011b {published data only}

Wang HQ, Gao XY, Zhou YL. Clinical research on electroacupuncture of ren ying for dysphagia due to stroke [电针入迎穴治疗中风吞咽困难的临床研究]. *China Journal of Chinese Medicine* 2011;**26**(4):501-2.

Wang 2011c {published data only}

Wang JH. Effect on the motor function of stroke patients by combination of needling at back-shu point and trunk exercise [针刺背俞穴结合躯干训练对中风偏瘫患者运动功能的影响]. *Chinese Journal of Integrated Traditional and Western Medicine* 2011;**31**(6):745-6.

Wang 2011d {published data only}

Wang L, Liu SM, Liu M, Li BJ, Hui ZL. Research on the effects of acupuncture, psychological intervention and speech rehabilitation training for postapoplectic aphasia and depression/anxiety [针灸及心理干预配合语言康复训练治疗卒中后语言障碍伴抑郁/焦虑的研究]. *Journal of New Chinese Medicine* 2011;**43**(12):90-2.

Wang 2011e {published data only}

Wang L, Liu SM, Liu M, Li BJ, Hui ZL, Gao X. Post-stroke speech disorder treated with acupuncture and psychological intervention combined with rehabilitation training: a randomized controlled trial [针刺及心理干预配合康复训练治疗卒中后言语障碍: 随机对照研究]. *Chinese Acupuncture & Moxibustion* 2011;**31**(6):481-6.

Wang 2011f {published data only}

Wang LC, Liu HY, Wang QH, Yang YG. Clinical research on acupuncture at Neiguan point combined with rehabilitation training for treatment of postapoplectic hand cramps [针刺内关穴联合康复训练治疗中风后手部痉挛临床研究]. *Chinese Journal of Integrative Medicine On Cardio-/Cerebrovascular Disease* 2011;**9**(6):689-90.

Wang 2011g {published data only}

Wang M, Wang JH, Hua QH. Clinical observation on anandia after stroke which was treated by acupuncture [针刺治疗中风后运动性失语的临床观察]. *Journal of Sichuan of Traditional Chinese Medicine* 2011;**29**(7):116-8.

Wang 2011h {published data only}

Wang S. Clinical study of scalp acupuncture and body acupuncture combined with language rehabilitation training for aphasia after stroke [头针体针配合语言康复训练治疗中风后失语症的临床研究]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10199-1011224431.htm> (accessed 8 August 2015) 2011.

Wang 2011i {published data only}

Wang T. Clinical observation of Xinnao Shutong pill combined with acupuncture for dysphagia after stroke [心脑舒通联合针灸治疗缺血性中风后吞咽困难]. *Chinese Journal of Integrative Medicine on Cardio/Cerebrovascular Disease* 2011;**9**(7):809-10.

Wang 2011j {published data only}

Wang YK. Clinical observation on the effects of acupuncture combined with Chinese medicine in the treatment of ischemic stroke with Qi deficiency and blood stasis [针刺联合中药治

疗缺血性中风恢复期气虚血瘀证的临床观察]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10162-1011202597.htm> (accessed 8 August 2015) 2011.

Wang 2013 {published data only}

Wang WF. Clinical experience on acupuncture in the treatment of stroke [针灸治疗脑卒中的临床体会]. *Modern Diagnosis and Treatment* 2013;**24**(4):795, 814.

Wang 2014a {published data only}

Wang YJ, He J, Sun CC, Zhang YM, Du JG, Zhao JG. Clinical study on transcutaneous acupoint electrical stimulation and acupuncture on postapoplectic depression [经皮穴位电刺激与针刺治疗卒中后抑郁的临床研究]. *Clinical Journal of Rehabilitation Medicine* 2014;**29**(8):751-3.

Wang 2014b {published data only}

Wang ZY, Ye W, Yang LZ, Ma DY, Xu Q. Effects of acupuncture on recovery of neurological function in patients with hypertensive cerebral hemorrhage [针刺对高血压脑出血患者神经功能恢复情况的作用]. *Information on Traditional Chinese Medicine* 2014;**31**(4):142-3.

Wei 2005 {published data only}

Wei LL. Effect of Shuitu acupoint injection with stellate ganglion block on swallow dysfunction after stroke. *Chinese Journal of Clinical Rehabilitation* 2005;**9**(9):106-7.

Wei 2009 {published data only}

Wei CL. Clinical observation on the effects of acupuncture combined with Buyanghuanwu decoction treatment on Qi deficiency and blood stasis hemiplegia after ischemic stroke [针刺结合补阳还五汤治疗缺血性中风气虚血瘀型的临床疗效观察]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10315-2009252060.htm> (accessed 8 August 2015) 2009.

Wenli 2007 {published data only}

Zhang WL, Feng LH, Jia XH, Feng JP, Xu Y. Clinical observation on therapeutic effects of the point-penetrating method in acupuncture treatment of spastic hemiparalysis due to cerebrovascular disorders. *Journal of Traditional Chinese Medicine* 2007; Vol. 27, issue 3:170-2.

Wong 2013 {published data only}

Wong GK. Can acupuncture benefit putaminal hemorrhagic stroke patients? A randomized controlled clinical trial. *Stroke* 2013;**44**:AWP319.

Wu 1999 {published data only}

Wu D. Needling eye points combined with electrical scalp points in the treatment of urinary incontinence due to cerebrovascular accident in 40 patients. *Journal of Clinical Acupuncture and Moxibustion* 1999;**15**(9):33-4.

Wu 2009a {published data only}

Wu P, Liang FR, Hu KM, Tian WW, Yuan XL, Huang YL. Clinical research of acupuncture combined with facilitation technique for limb spasm after apoplexy hemiplegia - a report of 160 cases [针刺复合易化技术治疗脑卒中后偏瘫肢体痉挛的临床研究]. *Journal of Chengdu University of TCM* 2009;**32**(1):11-3.

Wu 2009b {published data only}

Wu XM, Yu YM. Clinical research on the effects of acupuncture of backshu point and jiaji point in the treatment of spastic paralysis due to stroke [针刺背俞穴、夹脊穴治疗脑卒中后痉挛性瘫痪的临床研究]. *Beijing Journal of Traditional Chinese Medicine* 2009;**28**(7):494-6.

Wu 2010 {published data only}

Wu XL, An ZH, Li YQ, Liu AG, Li XD. Effects of acupuncture on the rehabilitation of neurological function in the 157 hemiplegia patients due to cerebral infarction [针刺对157例不同分型脑梗死(偏瘫)患者神经运动功能康复的影响]. *Journal of Traditional Chinese Medicine* 2010;**51**(10):909-11.

Wu 2011 {published data only}

Wu P, Liang RF, Li Y, Yang L, Huang YL, Li AH. Clinical observation on acupuncture plus rehabilitation training for dysphagia after stroke: a multicenter randomized controlled trial [针刺联合康复训练治疗脑卒中后吞咽障碍多中心随机对照临床观察]. *Journal of Traditional Chinese Medicine* 2011;**52**(1):45-8.

Wu 2012 {published data only}

Wu JY, Huang RD, Ge LB. Clinical study on acupuncture for ischemic stroke sequelae [针刺治疗缺血性中风后遗症临床研究]. *Shanghai Journal of Acupuncture and Moxibustion* 2012;**31**(2):85-7.

Wu 2012a {published data only}

Wu HK, Tan F, Wan SY, Wang JL, Huang T, Ding DQ. Clinical study of curative effect of electro-acupuncture combined with functional electric stimulation on the treatment of spastic dysarthria in acute cerebral infarction [电针联合功能性电刺激早期干预对脑梗死后痉挛性构音障碍的疗效研究]. *Chinese Journal of Rehabilitation Medicine* 2012;**27**(12):1116-9.

Wu 2013b {published data only}

Wu XL, Lu BQ, Li YQ, Xu JH, Wang LY, Liu AG. Effects of acupuncture on disease severity score and activity of daily living scales in patients with partial anterior circulation infarction [针刺对部分前循环脑梗死患者疾病严重程度及日常生活能力的影响]. *Journal of Traditional Chinese Medicine* 2013;**54**(7):1210-3.

Wu 2014a {published data only}

Wu XL, Lu BQ, Li YQ, Xu JH, Zhang YH. Effects of acupuncture on motor function in patients with partial anterior circulation infarction [针刺对部分前循环老年脑梗死患者运动功能的康复效果]. *Chinese Journal of Gerontology* 2014;**34**(3):1463-5.

Wu 2014b {published data only}

Wu ZJ, Hu KM, Guo YG, Tu YM, Zhang HY, Wang Y. Acupuncture combined with speech rehabilitation training for post-stroke spasmodic dysphonia: a multicenter randomized controlled trial [针刺结合言语康复训练治疗脑卒中后痉挛性构音障碍:多中心随机对照研究]. *World Journal of Acupuncture-Moxibustion* 2014;**24**(4):12-6.

Xia 2008 {published data only}

Xia K, Yang SR, Dang DH, Sun Q. Ear acupuncture combined with Xinnaokaigiao recipe for dysphagia caused by cerebral hemorrhage [耳针配合醒脑开窍治疗脑出血外伤术后吞咽

困难的临床研究]. *Chinese Journal of Integrative Medicine on Cardio/Cerebrovascular Diseases* 2008;**6**(8):994-5.

Xia 2010 {published data only}

Xia WG, Zhu CJ, Zhu SQ, Tang ZP, Wang Y, Hua Q. Combination of feeding-swallowing training and acupuncture: an effective rehabilitation method for dysphagia after stroke [脑卒中后吞咽障碍综合康复治疗的评价]. *Acta Medicinæ Universitatis Scientiæ et Technologiæ Huazhong* 2010;**39**(5):614-9.

Xiao 1996 {published data only}

Xiao F, Liu W. Observation on treatment of dysphasia due to cerebrovascular accident with electroacupuncture at shengen point. *Journal of Chinese Medicine* 1996;**50**:27-9.

Xiao 2006 {published data only}

Xiao QY. Clinical observation on the effects of acupuncture combined with catgut-embedding for hemiplegic shoulder pain after stroke [针刺配合埋线治疗中风后偏瘫肩痛的临床疗效观察]. *Chinese Medicine Modern Distance Education of China* 2006;**4**(12):31-3.

Xiao 2008 {published data only}

Xiao AH, Liu YP, Huang SJ, Liu B. Influence of meridian exercises and activating collaterals oil knocking stimulation acupoints on rehabilitation effect of cerebral apoplexy patients [经络操加活络油叩击刺激穴位对脑卒中康复疗效的影响]. *Chinese Nursing Research* 2008;**22**(4):888-9.

Xiao 2011 {published data only}

Xiao D. Clinical study of bowels-unblocking and toxin-purging therapy combined with acupuncture for patients of post-stroke consciousness disturbance with phlegm-heat stagnation in Fu-organs [通腑泄毒法加针刺治疗痰热腑实型脑卒中意识障碍的临床研究]. *Journal of New Chinese Medicine* 2011;**43**(5):14-6.

Xie 2010 {published data only}

Xie Q. A clinical study of acupuncture combined with rehabilitation training in the treatment of shoulder-hand syndrome after stroke [针刺结合康复训练治疗脑卒中肩手综合征的临床研究]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10572-2010125364.htm> (accessed 8 August 2015) 2010.

Xie 2012 {published data only}

Xie XM, An JM, Huang LN, Yang YK. Effects of acupuncture on motor dysfunction of patients caused by ischemic brain stroke: a randomized and controlled study [针刺对缺血性中风患者运动功能障碍影响的随机对照研究]. *Journal of Clinical Acupuncture and Moxibustion* 2012;**28**(12):15-8.

Xie 2013 {published data only}

Xie HY, Lin HX, Lin GH. Clinical research of fire-needle combined with Huoluo Guben decoction for 44 patients with hemiplegia caused by stroke [火针配合活络固本汤治疗中风后偏瘫44例]. *World Science and Technology/Modernization of Traditional Chinese Medicine and Materia Medica* 2013;**15**(2):281-5.

Xing 2007 {published data only}

Xing J, Wang YJ, Li YR. Clinical study on acupuncture combined with hyperbaric oxygenation for improving balance function of

cerebral infarction [针刺并高压氧改善脑梗死平衡功能的临床观察]. *Chinese Acupuncture & Moxibustion* 2007;**27**(1):12-4.

Xu 2009a {published data only}

Xu HY, Zhang H. Clinical observation on the effects of acupuncture combined with pelvic floor muscle training for uracratia after stroke [针灸配合盆底肌锻炼治疗脑卒中尿失禁临床观察]. *Lishizhen Medicine and Materia Medica Research* 2009;**20**(1):225-6.

Xu 2009b {published data only}

Xu Z. Clinical study on the treatment of sequela of apoplexy by acupuncture [针灸治疗中风后遗症临床疗效观察]. *Chinese Journal of Modern Drug Application* 2009;**3**(24):132-3.

Xu 2010 {published data only}

Xu SF, Zhuang LX, Jia C, Chen XH, Wu SP, Jiang GM, et al. Effect of "Jin three-needle therapy" on motor function in patients of hemiplegia after stroke: a multi-center randomized controlled study ["靳三针"疗法与康复治疗改善脑卒中偏瘫患者肢体功能的疗效对比观察: 多中心随机对照]. *Chinese Acupuncture & Moxibustion* 2009;**29**(9):689-94.

Xu 2013a {published data only}

Xu RQ. Clinical study of acupuncture treatment for cerebral infarction in terms of heart-gall bladder theory [从心胆论治针灸治疗脑梗死的临床研究]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10572-1013226466.htm> (accessed 8 August 2015) 2013.

Xu 2014 {published data only}

Xu G. A random parallel control study of acupuncture combined fluoxetine treatment for post-stroke depression [针刺联合氟西汀治疗中风后抑郁症随机平行对照研究]. *Journal of Practical Traditional Chinese Internal Medicine* 2014;**28**(11):158-60.

Yan 2010 {published data only}

Yan J, Zhong XQ, Wang QZ, Meng J, Jiang SY, Wen XQ. Clinical observation on the effects of acupuncture combined with rehabilitation training for dysphagia after apoplexy [针刺结合康复训练治疗脑卒中后吞咽困难疗效观察]. *Modern Journal of Integrated Traditional Chinese and Western Medicine* 2010;**19**(31):3368-9.

Yan 2011 {published data only}

Yan XR. Different acupuncture and moxibustion methods in the treatment of post-stroke hiccup [不同针灸方法治疗中风后呃逆的临床观察]. *Journal of Clinical Acupuncture and Moxibustion* 2011;**27**(11):43-5.

Yang 2006 {published data only}

Yang SL. Effects of needling combined with cognitive function training on the cognitive function of stroke patients [针刺结合认知功能训练对脑卒中患者认知功能的影响]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10393-2006162984.htm> (accessed 8 August 2015) 2006.

Yang 2007a {published data only}

Yang D, Xu L, Zhang XR. Clinic observation of integrated acupuncture and Chinese medicine for 50 cases with hemiplegia after stroke [针药结合治疗脑卒中偏瘫

50例]. *Shanxi Journal of Traditional Chinese Medicine* 2007;**28**(8):1060-1.

Yang 2007b {published data only}

Yang DF, Lin XD, Yu ZM, Wu ZB, Wu XF, Lin JF. Effect of acupuncture combined with Bobath therapy on spasticity and motor function suffered from postapoplectic hemiplegia [组合针刺结合Bobath疗法对脑卒中偏瘫痉挛状态及运动功能的影响]. *Liaoning Journal of Traditional Chinese Medicine* 2007;**34**(2):215-6.

Yang 2007c {published data only}

Yang Y. Clinical observation on the effects of spastic paralysis after apoplexy with electric acupuncture combined with rehabilitation training [电针配合康复训练治疗脑梗死偏瘫肢体痉挛状态的临床观察]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10228-2007120088.htm> (accessed 8 August 2015) 2007.

Yang 2008 {published data only}

Yang SM. Acupuncture combined with low frequency impulse electrotherapy for 36 cases with dysphagia after stroke [针刺配合低频脉冲电治疗脑卒中后吞咽困难36例]. *Zhejiang Journal of Traditional Chinese Medicine* 2008;**43**(6):348.

Yang 2009 {published data only}

Yang ZH, Yu K, Li CY, Hou QF, Liu XQ. Application value of acupuncture in the rehabilitative treatment of patients with cerebral infarction [针灸在脑梗死康复治疗中的应用价值观察]. *Journal of Military Surgeon in Southwest China* 2009;**11**(3):434-5.

Yang 2010 {published data only}

Yang H, Wang J. Clinical observation on the effects of acupuncture treatment for 29 cases with dysphagia after stroke [针刺治疗中风后吞咽障碍29例]. *Journal of Practical Traditional Chinese Internal Medicine* 2010;**24**(2):93-4.

Yang 2011a {published data only}

Yang HF, Li XX, Wang T. The videofluoroscopic study on electroacupuncture in combination with rehabilitation in treatment of postapoplectic swallowing dysfunction [电针联合康复训练治疗脑卒中后吞咽障碍的电视荧光透视研究]. *Journal of Guangzhou University of TCM* 2011;**28**(3):239-42.

Yang 2011b {published data only}

Yang J. The clinical study on cognitive impairment after stroke by using the treatment of electroacupuncture given at DU20 and GB20 [电针百会、风池干预脑卒中后认知功能障碍的临床研究]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10393-1011220797.htm> (accessed 8 August 2015) 2011.

Yang 2011c {published data only}

Yang J. Study on acupuncture therapy of dredging governor vessel for regulating mentality in post ischemic stroke hemiplegic patients [通督调神针刺法治疗缺血性中风偏瘫的临床研究]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10572-1011132621.htm> (accessed 8 August 2015) 2011.

Yang 2012 {published data only}

Yang CD, Yang ZJ, Li JJ. Clinical observation of a combination of multiple acupuncture technique for hemiplegia after stroke [多种针灸手法联合治疗中风偏瘫的疗效观察]. *China Practical Medicine* 2012;**7**(33):239-40.

Yang 2014 {published data only}

Yang XH, Liu B, Ouyang BS. Effect of acupuncture combined with rehabilitative training on neural functional recovery of stroke patients during recovery phase: A randomized controlled trial [针刺治疗结合康复训练对恢复期卒中患者神经功能恢复的影响:随机对照研究]. *World Journal of Acupuncture Moxibustion*. 2014;**24**(4):17-23.

Yang 2014a {published data only}

Yang XH, Liu B, Ouyang BS. Effect of acupuncture combined with rehabilitative training on neural functional recovery of stroke patients during recovery phase: a randomized controlled trial. *World Journal of Acupuncture-Moxibustion* 2014;**24**(4):17-23.

Yang 2015 {published data only}

Yang HL, Zhang B, Liu T, Zheng JG. Efficacy of acupuncture in combination with medicine for mild cognitive impairment after cerebral infarction: a randomized controlled trial. *World Journal of Acupuncture-Moxibustion* 2015;**25**(1):7-12.

Yao 2006 {published data only}

Yao FD. Observation on the effects of acupuncture combined with language rehabilitation training for 22 patients with aphasia after stroke [针灸配合语言康复训练治疗脑卒中失语22例效果观察]. *Journal of Guangdong Medical College* 2006;**24**(3):290-1.

Yao 2013a {published data only}

Yao QF. Clinical observation on upper limb spastic hemiplegia following cerebral infarction treated by antagonistic acupuncture combined with rehabilitation [拮抗针法配合康复治疗脑梗死后上肢痉挛性偏瘫的临床疗效观察]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10228-1013249894.htm> (accessed 8 August 2015). 黑龙江中医药大学, 2013.

Yao 2013b {published data only}

Yao YF, Gao X, Jing R, Li SQ, Ba HZ. Efficacy observation on the treatment of poststroke hemiplegia in 30 patients by consciousness restoring obstruction clearing needling technique [醒脑开窍针刺法结合康复治疗脑卒中偏瘫患者30例]. *Shanxi Journal of Traditional Chinese Medicine* 2013;**34**(8):1047-8.

Yin 2013a {published data only}

Yin LL. A clinical study on acupuncture combined with rehabilitation therapy for 57 cases of swallowing disorders after stroke [针刺配合康复治疗中风后吞咽障碍57例临床研究]. *Journal of Traditional Chinese Medicine* 2013;**54**(9):766-8.

Yin 2013b {published data only}

Yin ZK. The clinical observation on treatment of dysphagia caused by pseudobulbar palsy after stroke with acupuncture Fengchi Fengfu mainly acupuncture point [针刺风池、风府为主穴治疗脑卒中后假性延髓麻痹吞咽困难临床

观察]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10228-1013249935.htm> (accessed 8 August 2015) 2013.

Yin 2014 {published data only}

Yin JC. Clinical observation on Quanjing acupuncture therapy combined with rehabilitation training for the treatment of shoulder hand syndrome after stroke [全经针刺结合康复训练治疗中风后肩手综合征的临床观察]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10541-1014241244.htm> (accessed 8 August 2015) 2014.

You 2011 {published data only}

You Y. A clinical study of electroacupuncture combined with rehabilitation training in the treatment of shoulder-hand syndrome of post-stroke [电针配合康复训练治疗脑卒中后肩手综合征的临床研究]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10441-1011204989.htm> (accessed 8 August 2015) 2011.

You 2014 {published data only}

You Y. Efficacy observation of disperse-dense wave electrical acupuncture combined with occupational therapy in the treatment of shoulder-hand syndrome due to stroke [电针疏密波配合作业疗法治疗脑卒中后肩手综合征的疗效观察]. *Chinese Journal of Convalescent Medicine* 2014;**23**(6):509-11.

Yu 2002 {published data only}

Yu MZ, Yang GZ, Zhan KR, Fan YY, Peng MF. Clinical observation on the effects of electrical stimulation at nervous trunk points in the treatment of sequelae of stroke. *Chinese Journal of Basic Medicine in Traditional Chinese Medicine* 2002;**8**(3):55-6.

Yu 2007 {published data only}

Yu XG, Sun SL, Cheng YH. Clinical observation on treatment of mild cognitive impairment after cerebral infarction with acupuncture dumaijingxue mainly point [以针刺督脉经穴为主治疗缺血性脑卒中后轻度认知功能障碍的临床观察]. *Chinese Journal of Integrative Medicine On Cardio-/Cerebrovascular Disease* 2007;**5**(12):1171-3.

Yu 2009 {published data only}

Yu JY, Guo XF. Observation on the effects of acupuncture in combination with rehabilitation in the treatment of spasticity following stroke [电针结合康复训练治疗脑卒中痉挛状态的疗效观察]. *Journal of External Therapy of TCM* 2009;**18**(4):42-3.

Yu 2011 {published data only}

Yu XM. Clinical observation on electric acupuncture combined with rehabilitation training for 50 patients with neurogenic bladder following stroke [电针结合康复训练治疗卒中后神经源性膀胱50例临床观察]. Available at: <http://cpfd.cnki.com.cn/Article/CPFDTOTAL-ZGKF201108001153.htm> (accessed 8 August 2015). 2011:176.

Yuan 2010 {published data only}

Yuan LX. The research on acupuncture treatment technology of dysphagia after stroke. ChiCTR Chinese Clinical Trial Register (<http://www.chictr.org/>) (ChiCTR-TRC-12002695) (accessed 8 August 2015) 2010.

Zeng 2010 {published data only}

Zeng XQ, Teng DS, Lin X, Yang T, Wang XF. Observations on the efficacy of acupuncture as main treatment for post-stroke dysphagia [针刺为主治疗中风后吞咽障碍疗效观察]. *Shanghai Journal of Acupuncture and Moxibustion* 2010;**29**(7):431-2.

Zhang 1988 {published data only}

Zhang ZJ. Observation on the effects of acupuncture in the treatment of post stroke aphasia. *Journal of Traditional Chinese Medicine* 1988;**5**:33-4.

Zhang 1997 {published data only}

Zhang SF, Ye XR, Dan QH, Zhang WF, Cui Y. The influence of acupuncture on SOD, CAT and serum intraproteinase fluorescence in stroke patients. *Chinese Acupuncture and Moxibustion* 1997;**9**:517-9.

Zhang 2002a {published data only}

Zhang L, Sheng L. Clinical observation on the effects of magnetic acupuncture in the treatment of sequelae of stroke. *Chinese Journal of Information in Traditional Chinese Medicine* 2002;**9**(1):63.

Zhang 2002b {published data only}

Zhang ZZ, Ma FJ, Ma Y. Observation on the effects of acupuncture in the treatment of urinary retention due to cerebral infarction in 36 patients [针刺对脑梗塞所致尿潴留36例疗效观察]. *Hei Long Jiang Medicine and Pharmacy* 2002;**25**(3):71.

Zhang 2007a {published data only}

Zhang HM. Clinical research of treatment of apoplectic aphemia with multi-needle puncture of scalp-points in combination with visual-listening-speech training [头穴丛刺结合言语康复治疗脑卒中运动性失语临床研究]. *Acupuncture research* 2007;**32**(3):190-4.

Zhang 2007b {published data only}

Zhang XT, Fan RX. Clinical observation on the effects of acupuncture combined with buyang huanwu decoction in the treatment of sequela of cerebral infarction [补阳还五汤配合针灸治疗脑梗塞后遗症疗效观察]. *Jilin Journal of Traditional Chinese Medicine* 2007;**27**(1):31-2.

Zhang 2008 {published data only}

Zhang WM, Zhen P. Acupuncture combined with swallowing therapy instrument for 82 cases with dysphagia after stroke [针刺结合吞咽言语诊治仪治疗中风后吞咽障碍82例]. *Jilin Journal of Traditional Chinese Medicine* 2008;**28**(11):819-20, 825.

Zhang 2009b {published data only}

Zhang NX, Huang TQ, Liu GZ, Hou YQ, Liu SQ, Luo JQ, et al. Effect of acupuncture and rehabilitation training on Barthel index in early-stage of stroke cases. *Journal of Acupuncture and Tuina Science* 2009;**7**(3):143-6.

Zhang 2009c {published data only}

Zhang W. Clinical research of acupuncture for deglutition handicap caused by supranuclear paralysis after cerebral infarction [针刺治疗脑梗死后假性延髓麻痹所致吞咽障碍

的临床研究]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10162-2009261860.htm> (accessed 8 August 2015) 2009.

Zhang 2010a {published data only}

Zhang CH. Clinical research of spastic paralysis after stroke by acupuncture. ChiCTR Chinese Clinical Trial Register (<http://www.chictr.org/>) (ChiCTR-TCH-11001547) (accessed 8 August 2015) 2010.

Zhang 2010b {published data only}

Zhang N. Clinical research of acupuncture combined with rehabilitation training for dysphagia recovery [针刺联用康复训练治疗中风病（血瘀证）恢复期吞咽障碍的临床研究]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10199-2010255182.htm> (accessed 8 August 2015) 2010.

Zhang 2010c {published data only}

Zhang NX, Liu GZ, Yao QH, Li WJ, Huang Y, Wang AM, et al. Effects of warming-reinforcing acupuncture combined with rehabilitation training on the early motor function of hemiparalysis patients caused by ischemic brain stroke: a randomized controlled study [热补针法结合康复训练对缺血性卒中偏瘫患者早期运动功能的影响:随机对照研究]. *Chinese Acupuncture & Moxibustion* 2010;**30**(6):441-5.

Zhang 2010d {published data only}

Zhang NX, Liu GZ, Yao QH, Li WJ, Huang Y, Wang AM. Effects of warming-reinforcing acupuncture combined with rehabilitation training on the early motor function of hemiparalysis patients caused by ischemic brain stroke: a randomized and controlled study. *Chinese Acupuncture & Moxibustion* 2010;**30**(6):441-5.

Zhang 2010e {published data only}

Zhang SH. Acupuncture combined with Jieyudan recipe for 33 cases with dysphagia after stroke [针灸配合解语丹加减治疗中风后吞咽困难33例]. *Zhejiang Journal of Traditional Chinese Medicine* 2010;**45**(8):509.

Zhang 2010f {published data only}

Zhang ZG. Acupuncture for postoperative intracranial hematoma after hypertensive intracerebral hemorrhage [针刺对高血压脑出血术后患者的临床疗效研究]. Available at: <http://cdmd.cnki.com.cn/article/cdmd-10441-1011205188.htm> (accessed 8 August 2015) 2010.

Zhang 2011a {published data only}

Zhang HM, Tang Q. Rehabilitation evaluation on post-stroke abnormal movement pattern prevented and treated with acupuncture and rehabilitation [针刺加康复防治卒中异常运动模式的康复学评定]. *Chinese acupuncture & moxibustion* 2011;**31**(6):487-92.

Zhang 2011b {published data only}

Zhang JH, Li JY, Zhao Y, Fan GQ, Wang JQ. Clinical observation of dysphagia after stroke by Tongnao Huoluo acupuncture therapy with swallowing training [通脑活络针刺法配合吞咽训练治疗脑卒中后吞咽障碍临床观察]. *Journal of Liaoning University of TCM* 2011;**13**(12):188-9.

Zhang 2011c {published data only}

Zhang WM. Post-apoplexy spasm treated by acupuncture on antagonistic muscles combined with modern rehabilitation training therapy [拮抗肌侧取穴针刺结合现代康复训练疗法治疗中风后痉挛]. *World Journal of Integrated Traditional and Western Medicine* 2011;**6**(4):291-3, 301.

Zhang 2011d {published data only}

Zhang XH, Feng GX, Wen N. Clinical study of warm-needling therapy on stroke patients with urinary retention [温针灸治疗中风后尿潴留患者的临床观察]. *Journal of TCM Univ. of Human* 2011;**31**(3):72-4.

Zhang 2011e {published data only}

Zhang XW. Clinical research on the effects of electroacupuncture and exercise therapy in the treatment of SHS [电针结合康复训练治疗中风后肩手综合征的疗效观察]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10572-1011135406.htm> (accessed 8 August 2015) 2011.

Zhang 2012b {published data only}

Zhang F. Study on acupuncture therapy of dredging governor vessel for regulating mentality in post ischemic stroke aphasia patients [通督调神针刺法治疗缺血性中风失语症临床研究]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10572-1012397626.htm> (accessed 8 August 2015) 2012.

Zhang 2012c {published data only}

Zhang JW, Hu JD, MPOVA, LOLA. Clinical study on acupuncture for stroke in Congo [针刺在刚果金治疗脑中风的临床研究]. *World Journal of Acupuncture-Moxibustion* 2012;**22**(1):35-8.

Zhang 2012d {published data only}

Zhang YS, Yu ZP, Zhang YL, Zhao XH, Lin CR. Therapeutic observation on acupuncture and rehabilitation in treating motor aphasia after cerebral apoplexy [针刺配合康复训练治疗脑卒中后运动性失语疗效观察]. *Shanghai Journal of Acupuncture and Moxibustion* 2012;**31**(11):781-3.

Zhang 2012e {published data only}

Zhang ZS, Qin YM, Li YH. Clinical observation of warming-promotion method of acupuncture combined with Buyanghuanwu decoction on ischemic stroke with qi-deficiency and blood-stasis syndrome [温通针法联合补阳还五汤治疗气虚血瘀型缺血性卒中临床观察]. *Hebei Journal of Traditional Chinese Medicine* 2012;**34**(11):1609-11.

Zhang 2013b {published data only}

Zhang W. Early intervention and prevention with acupuncture therapy on the post stroke depression: A randomized controlled clinical trial. ChiCTR Chinese Clinical Trial Register (<http://www.chictr.org/>) (ChiCTR-TRC-120034890) (accessed 8 August 2015) 2013.

Zhang 2014b {published data only}

Zhang H. Impact on rehabilitation effect of post-stroke abnormal movement pattern prevented and treated with multi-needle puncture of scalp-points. *Journal of alternative and complementary medicine* 2014;**20**(5):A47-8.

Zhang 2014c {published data only}

Zhang Y. A random parallel control study on the effects of acupuncture combined with fluoxetine tablet for depression after stroke [针灸联合氟西汀分散片治疗脑卒中后抑郁症随机平行对照研究]. *Journal of Practical Traditional Chinese Internal Medicine* 2014;**28**(2):123-4.

Zhang 2014d {published data only}

Zhang Y, Chen WP, Wang CY, Xie T, Wang M, Xia YJ. Modified acupuncture combined with rehabilitation therapy in treatment of limb spasticity in stroke patients [针灸解痉调气法结合康复治疗脑卒中痉挛期患者临床评价]. *Zhejiang Journal of Integrated Traditional and Western Medicine* 2014;**24**(4):290-2.

Zhao 2008 {published data only}

Zhao YY. Acupuncture for 82 cases with dysphagia after stroke ["醒脑开窍"针刺法治疗假球麻痹82例]. *Journal of Liaonin University of Traditional Chinese Medicine* 2008;**10**(7):91.

Zhao 2009 {published data only}

Zhao Y, Wang Q. Clinical observation of acupuncture for 60 cases of dysphagia after ischemic stroke [深刺治疗缺血性脑卒中后吞咽困难60例]. *Jilin Journal of Traditional Chinese Medicine* 2009;**29**(6):503-4.

Zhao 2010 {published data only}

Zhao GP, Sun J. Observation on the effects of acupuncture combined with language rehabilitation training for 174 patients with aphasia following stroke [针灸配合语言康复训练治疗脑卒中失语174例效果观察]. *Chinese Journal of Trauma and Disability Medicine* 2010;**18**(1):87-8.

Zhao 2011a {published data only}

Zhao NN. Clinical research of fire-needle combined with rehabilitation training on spastic hemiplegia after stroke [火针配合康复训练对脑卒中后痉挛性偏瘫的临床观察]. Available at: <http://cdmd.cnki.com.cn/article/cdmd-10572-1011135337.htm> (accessed 8 August 2015) 2011.

Zhao 2011b {published data only}

Zhao SJ. Clinical research on the effects of acupuncture therapy for the ischemic stroke [针刺治疗缺血性脑卒中的临床研究]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10162-1011201326.htm> (accessed 8 August 2015) 2011.

Zhao 2014a {published data only}

Zhao YH, Sun ZR, Zhang DQ, Li HN, Zhong JB, Liu J. Clinical observation on the electrical acupuncture with intermediate frequency combined with rehabilitation treating ischemic stroke [电针中频配合康复治疗缺血性脑卒中的临床观察]. *Chinese Journal of Integrative Medicine On Cardio/Cerebrovascular Disease* 2014;**12**(5):581-3.

Zhao 2014b {published data only}

Zhao XW, Chen XW, Wei CF, Tian QT, Yuan W. The effects of early intervention of rehabilitation training and acupuncture on stroke patients with neurological impairment and activities of daily living [康复训练结合针刺早期干预对脑卒中患者神经功能缺损及日常生活活动能力的作用]. *Medical Journal of Industrial Enterprise* 2014;**27**(1):541-3.

Zheng 1996 {published data only}

Zheng ZC. Needling scalp and lumbo-sacral points in the treatment of sequelae of stroke in 634 patients. *Shanghai Journal of Acupuncture and Moxibustion* 1996;**15**(5):15-6.

Zheng 2000 {published data only}

Zheng L, Zhuang LX, Li YH. Observations on the curative effect of acupuncture and herbs on vascular dementia. *Shanghai Journal of Acupuncture and Moxibustion* 2000;**19**(3):8-10.

Zheng 2011b {published data only}

Zheng QH, Yu B, Li Y, Yang ZQ, Yu SY. Clinical study of acupuncture with language training for aphasia after stroke [针刺配合语言训练治疗脑卒中后失语症临床研究]. *Journal of Liaoning University of TCM* 2011;**13**(1):105-7.

Zheng 2011c {published data only}

Zheng WX. Clinical Research of Poststroke Anxiety Disorder Treated by Acupuncture Combined with Deanxit [针灸配合黛力新治疗脑卒中后焦虑症的临床研究]. *Clinical Journal of Chinese Medicine* 2011;**26**(4):468-70.

Zhong 2014 {published data only}

Zhong CX. The clinical research of feasibility of "Xingnao Kaqiao" acupuncture therapy on preventing depression after stroke ["醒脑开窍"针法预防中风后抑郁的临床观察]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10315-1014409916.htm> (accessed 8 August 2015) 2014.

Zhou 1995 {published data only}

Zhou JZ, Zhang CS, Li L, Zhou SW, Shi XM. Clinical study in the treatment of sequelae of apoplexy with "Xing Nao Kai Qiao" acupuncture therapy. *Chinese Acupuncture and Moxibustion*. 1995;**3**:6-8.

Zhou 2008 {published data only}

Zhou GY. Observation on the effects of electroacupuncture and rehabilitation for 83 cases with stroke [电针和康复治疗脑卒中83例疗效观察]. *Chinese Journal of Information on Traditional Chinese Medicine* 2008;**15**(5):77.

Zhu 2010a {published data only}

Zhu T. Clinical research on the treatment of dysphagia after stroke by traditional Chinese medicine combined with acupuncture [中药加针刺治疗中风后吞咽障碍的临床研究]. *Guangming Journal of Chinese Medicine* 2010;**25**(11):2074-5.

Zhu 2010b {published data only}

Zhu T. Clinical observation of acupuncture in the treatment of dysphagia after stroke [针刺治疗卒中后吞咽功能障碍临床疗效观察]. *Chinese Journal of Practical Nervous Diseases* 2010;**13**(17):65-6.

Zhu 2011 {published data only}

Zhu F, Zhao J, Ding Y. Clinical observation on treating post-stroke depression with both acupuncture and medicine. *International Journal of Clinical Acupuncture* 2011;**20**:101-3.

Zhu 2012a {published data only}

Zhu ST. Clinical research on electro-acupuncture at swallow hole for dysphagia induced by true bulbar paralysis after

stroke [电针吞咽穴治疗脑梗死后真性延髓麻痹吞咽障碍的临床观察]. Available at: <http://cdmd.cnki.com.cn/Article/CDMD-10228-1012467791.htm> (accessed 8 August 2015) 2012.

Zhu 2012b {published data only}

Zhu ZQ, Xu GG, Guo YL, Ma B. Efficacy of stroke unit combined with acupuncture therapy on dysphagia in stroke patients [卒中单元模式下联合针刺治疗脑卒中吞咽障碍疗效观察]. *Chinese Medical Journal* 2012;**23**(2):144-6,149.

Zhuang 2012 {published data only}

Zhuang LX, Xu SF, D'Amado CR, Jia C, He J, Han DX, et al. An effectiveness study comparing acupuncture, physiotherapy, and their combination in poststroke rehabilitation: a multicentered, randomized, controlled clinical trial. *Alternative Therapies in Health and Medicine* 2012;**18**(3):8-14.

References to ongoing studies

Fu 2011 {published data only}

Fu Q. Randomised, controlled Phase III Trial to evaluate the efficacy and safety of comprehensive acupuncture treatment programs for poststroke motor rehabilitation. ChiCTR Chinese Clinical Trial Register (<http://www.chictr.org/>) (ChiCTR-TRC-11001347) (accessed 8 August 2015) 2011.

Liu 2013b {published data only}

Liu T. Acupuncture based on five elements body characteristics in the treatment of patients with post-stroke depression. ChiCTR Chinese Clinical Trial Register (<http://www.chictr.org/>) (ChiCTR-TRC-13003434) (accessed 8 August 2015) 2013.

Xie 2006 {published data only}

Xie Y. Randomized controlled study on the acupuncture for dysphagia in convalescence phase of apoplexy. ChiCTR Chinese Clinical Trial Register (<http://www.chictr.org/>) (ChiCTR-TRC-07000010) (accessed 8 August 2015) 2006.

Zhong 2010 {published data only}

Zhong H. The research of linguistic functional recovery mechanism based on fMRI after electroacupuncture at acupoints Tongli (HT5) and Xuanzhong (GB39) curing basal ganglia aphasia after stroke. ChiCTR Chinese Clinical Trial Register (<http://www.chictr.org/>) (ChiCTR-TRC-11001410) (accessed 8 August 2015) 2010.

Additional references

Asplund 1988

Asplund K, Tuomilehto J, Stegmayr B, Wester PO, Tunstall-Pedoe H. Diagnostic criteria and quality control of the registration of stroke events in the MONICA project. *Acta Medica Scandinavica Supplementum* 1988;**728**:26-39.

Cheng 2008

Cheng H, Yu J, Jiang Z. Acupuncture improves cognitive deficits and regulates the brain cell proliferation of SAMP8 mice. *Neuroscience Letters* 2008;**432**:111-6.

Deeks 2001

Deeks JJ, Altman DG, Bradburn MJ. Statistical methods for examining heterogeneity and combining results from several studies in meta analysis. In: Egger M, Davey Smith G, Altman DG editor(s). *Systematic Reviews in Health Care: Meta-analysis in Context*. 2nd Edition. London: BMJ Publication Group, 2001.

Dickersin 1994

Dickersin K, Scherer R, Lefebvre C. Identifying relevant studies for systematic reviews. *BMJ* 1994;**309**(6964):1286-91. [MEDLINE: 7718048]

Dong 2013

Dong QJ, Yang ZX. Systematic review of resuscitation needling technique for wind stroke. *Chinese Acupuncture & Moxibustion* 2013;**33**(5):475-80.

Ernst 1996

Ernst E, White AR. Acupuncture as an adjuvant therapy in stroke rehabilitation?. *Wien Medizinische Wochenschrift* 1996;**21**(22):556-8.

Ernst 1997

Ernst E, White A. Life-threatening adverse reactions after acupuncture? A systematic review. *Pain* 1997;**71**:123-6.

Feigin 2014

Feigin VL, Forouzanfar MH, Krishnamurthi R, Mensah GA, Connor M, Bennett DA, et al. Global and regional burden of stroke during 1990–2010: findings from the Global Burden of Disease Study 2010. *Lancet* 2014;**383**:245-54.

GBD2013 2015

GBD 2013 Mortality and Causes of Death Collaborators. Global, regional, and national age–sex specific all-cause and cause-specific mortality for 240 causes of death, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 2015;**385**(9963):117-71.

Go 2014

Go AS, Mozaffarian D, Roger VL, Benjamin EJ, Berry JD, Blaha MJ, et al. Heart Disease and Stroke Statistics—2014 Update. A report from the American Heart Association. *Circulation* 2014;**128**:00-00.

Higgins 2003

Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ* 2003;**327**(7414):557-60. [MEDLINE: 12958120]

Higgins 2011a

Higgins JPT, Altman DG, Sterne JAC (editors). Chapter 8: Assessing risk of bias in included studies. In: Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available from www.cochrane-handbook.org (accessed 10 August 2015).

Hopwood 1996

Hopwood V. Acupuncture in stroke recovery: a literature review. *Complementary Therapies in Medicine* 1996;**4**:258-63.

Hu 1993

Hu HH, Chung C, Liu TJ, Chen RC, Chen CH, Chou P, et al. A randomized controlled trial on the treatment for acute partial ischemic stroke with acupuncture. *Neuroepidemiology* 1993;**12**:106-13.

Hugh 2010

MacPherson H, Altman DG, Hammerschlag R, Youping L, Taixiang W, White A, et al. Revised STAndards for Reporting Interventions in Clinical Trials of Acupuncture (STRICTA): extending the CONSORT Statement. *Journal of Evidence-Based Medicine* 2010;**3**:140-55.

Jansen 1989

Jansen G, Lundeborg T, Kjartansson S, Samuelson UE. Acupuncture and sensory neuropeptides increase cutaneous blood flow in rats. *Neuroscience Letters* 1989;**97**:305-9.

Johansson 1993

Johansson K, Lindgren I, Widner H, Wiklund I, Johansson BB, Johansson K, et al. Can sensory stimulation improve the functional outcome in stroke patients?. *Neurology* 1993;**43**:2189-92.

Johansson 2001

Johansson BB, Haker E, Arbin MV. Acupuncture and transcutaneous nerve stimulation in stroke rehabilitation: a randomized, controlled trial. *Stroke* 2001;**32**:707-13.

Kim 2010

Kim MK, Choi TY, Lee MS, Lee H, Han CH. Contralateral acupuncture versus ipsilateral acupuncture in the rehabilitation of post-stroke hemiplegic patients: a systematic review. *BMC Complementary and Alternative Medicine* 2010;**10**(10):41-2.

Kong 2010

Kong JC, Lee MS, Shin BC, Song YS, Ernst E. Acupuncture for functional recovery after stroke: a systematic review of sham-controlled randomized clinical trials. *Canadian Medical Association Journal* 2010;**182**(16):1723-29.

Lefebvre 1996

Lefebvre C, McDonald S. Development of a sensitive search strategy for reports of randomized controlled trials in EMBASE. Fourth International Cochrane Colloquium; 1996 Oct 20-24; Adelaide (Australia). 1996.

Li 2014

Li L, Zhang H, Meng SQ, Qian HZ. An updated meta-analysis of the efficacy and safety of acupuncture treatment for cerebral infarction. *PLoS ONE* 2014;**9**(12):e114057.

Lim 2015

Lim SM, Yoo JH, Lee EJ, Kim HJ, Shin SW, Han GJ, et al. Acupuncture for spasticity after stroke: a systematic review and meta-analysis of randomized controlled trials. *Evidence-Based Complementary and Alternative Medicine* 2015;**2015**:870398. [DOI: [10.1155/2015/870398](https://doi.org/10.1155/2015/870398)]

Liu 2005

Zhang SH, Liu M, Asplund K, Li L. Acupuncture for acute stroke. *Cochrane Database of Systematic Reviews* 2005, Issue 2. [DOI: [10.1002/14651858.CD003317.pub2](https://doi.org/10.1002/14651858.CD003317.pub2)]

Liu 2009a

Liu Z, Fang JQ, Zeng C. Development of studies on the underlying mechanism of acupuncture intervention in reducing post-ischemic inflammatory reaction and the related new research thought. *Zhen Ci Yan Jiu* 2009;**34**:61-6.

Liu 2014a

Liu F, Li ZM, Jiang YJ, Chen LD. A meta-analysis of acupuncture use in the treatment of cognitive impairment after stroke. *Journal of Alternative and Complementary Medicine* 2014;**20**(7):535-44.

Long 2012

Long YB, Wu XP. A meta-analysis of the efficacy of acupuncture in treating dysphagia in patients with a stroke. *Acupuncture in Medicine* 2012;**30**:291-7.

Magnusson 1994

Magnusson M, Johansson K, Johansson BB. Sensory stimulation promotes normalization of postural control after stroke. *Stroke* 1994;**25**:1176-80.

NIH 1998

NIH Consensus Conference. Acupuncture. *JAMA* 1998;**280**:1518-24.

Park 2001

Park J, Hopwood V, White AR, Ernst E. Effectiveness of acupuncture for stroke: a systematic review. *Journal of Neurology* 2001;**248**:558-63.

Park 2014

Park SW, Yi SH, Lee JA, Hwang PW, Yoo HC, Kang KS, et al. Acupuncture for the treatment of spasticity after stroke: a meta-analysis of randomized controlled trials. *Journal of Alternative and Complementary Medicine* 2014;**20**(9):672-82.

Qi 2009

Qi YZ, Fu LX, Xiong J, Wang ZL, Mou J, Lu YM. Systematic evaluation of acupuncture for treatment of post-stroke spastic paralysis. *Chinese Acupuncture & Moxibustion* 2009;**129**(8):683-8.

Ren 2008

Ren L, Zhang WA, Fang NY. The influence of electro-acupuncture on neural plasticity in acute cerebral infarction. *Neurological Research* 2008;**30**:985-9.

Schulz 2010

Schulz KF, Altman DG, Moher D, for the CONSORT Group. CONSORT 2010 Statement: updated guidelines for reporting parallel group randomised trials. *BMJ* 2010;**340**:c332.

SCSSS 1999

Swedish Collaboration on Sensory Stimulation in Stroke. Sensory stimulation after stroke: a randomized controlled trial. *Cerebrovascular Diseases* 1999;**9 Suppl 1**:28.

Smith 2002

Smith LA, Moore OA, McQuay HJ, Moore A. Assessing the evidence of effectiveness of acupuncture for stroke rehabilitation: stepped assessment of likelihood of bias. Available at: www.jr2.ox.ac.uk/bandolier/booth/alternat/ACstroke.html. (accessed 10 August 2015) 2002.

Sterne 2011

Sterne JAC, Egger M, Moher D (editors). Chapter 10: Addressing reporting biases. In: Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Intervention*. Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available from www.cochrane-handbook.org. (accessed 10 August 2015).

Sulter 1999

Sulter G, Steen C, Keyser JD. Use of the Barthel Index and Modified Rankin Scale in acute stroke trials. *Stroke* 1999;**30**:1538-41.

Sun 2001

Sun HL, Li XM. Clinical study on treatment of cerebral apoplexy with penetration needling of scalp acupoints. *Chinese Acupuncture and Moxibustion* 2001;**21**(5):275-8.

Sze 2002

Sze FK, Wong E, Or KKH, Lau J, Woo J. Does acupuncture have additional value to standard poststroke motor rehabilitation?. *Stroke* 2002;**33**:186-94.

Van Tulder 2000

Van Tulder MW, Cherkin DC, Berman B, Lao L, Koes BW. Acupuncture for low back pain. *Cochrane Database of Systematic Reviews* 2002, Issue 2. [DOI: [10.1002/14651858.CD001351](https://doi.org/10.1002/14651858.CD001351)]

Wang 2012a

Wang Y, Shen JG, Wang XM, Fu DL, Chen CY, Lu LY, et al. Scalp acupuncture for acute ischemic stroke: a meta-analysis of randomized controlled trials. *Evidence-Based Complementary and Alternative Medicine* 2012;**2012**(6):1325.

Wayne 2005

Wayne PM, Krebs DE, Eric A, Macklin EA, Schnyer R, Kaptchuk TJ, et al. Acupuncture for upper-extremity rehabilitation in chronic stroke: a randomized sham-controlled study. *Archives of Physical Medicine and Rehabilitation* 2005;**86**(12):2248-55.

Wong 2012

Wong ISY, Ng KF, Tsang HWH. Acupuncture for dysphagia following stroke: a systematic review. *European Journal of Integrative Medicine* 2012;**4**:e141-e150.

Wu 1996

Wu JN. A short history of acupuncture. *Journal of Alternative Complementary Medicine* 1996;**2**:19-21.

Wu 2006

Wu HM, Tang JL, Lin XP, Lau J, Leung PC, Woo J, et al. Acupuncture for stroke rehabilitation. *Cochrane*

Database of Systematic Reviews 2006, Issue 3. [DOI:
10.1002/14651858.CD004131.pub2]

Wu 2010a

Wu P, Mills E, Moher D, Seely D. Acupuncture in poststroke rehabilitation: a systematic review and meta-analysis of randomized trials. *Stroke* 2010;**41**:e171-e179.

Zhang 1996

Zhang X, Yuan Y, Kuang P. The changes of vasoactive intestinal peptide, somato-statin and pancreatic polypeptide in blood and CSF of acute cerebral infarction patients and the effect of acupuncture on them. *Chen Tzu Yen Chiu* 1996;**21**(4):10-6.

Zhang 2008a

Zhang HZ, Zhang LX, She YF. Effect of electroacupuncture on apoptosis of hippocampus tissue in mice with vascular dementia. *Zhen Ci Yan Jiu* 2008;**33**:377-81.

Zhang 2009a

Zhang JB, Ren L, Sun Y. Meta-analysis on acupuncture for treatment of depression in patients of poststroke. *Chinese Acupuncture & Moxibustion* 2009;**29**(7):599-602.

Zhang 2012a

Zhang GC, Fu WB, Liu JH, Zhu XP, Liang ZH, Huang YF, et al. Meta analysis of the curative effect of acupuncture on post-stroke depression. *Traditional Chinese Medicine* 2012;**32**(1):6-11.

Zhang 2014a

Zhang JP, Chen J, Chen JQ, Li XH, Lai XY, Zhang SQ, et al. Early filiform needle acupuncture for poststroke depression: a meta-analysis of 17 randomized controlled clinical trials. *Neural Regeneration Research* 2014;**9**(7):773-84.

Zheng 2011a

Zheng GQ, Zhao ZM, Wang Y, Gu Y, Li Y, Chen XM, et al. Meta-analysis of scalp acupuncture for acute hypertensive intracerebral hemorrhage. *Journal of Alternative and Complementary Medicine* 2011;**17**(4):293-9.

Zhou 2013

Zhou JW, Li J, Zhao JJ, Xie HJ, Wang M. Meta analysis on ischemic stroke treated with scalp acupuncture. *World Journal of Acupuncture-Moxibustion* 2013;**23**(2):41-7.

Zhu 2011a

Zhu LL, Wang WX, Guo XG. Acupuncture for hiccups after stroke: a systematic review. *Chinese Journal of Evidence-based Medicine* 2011;**11**(3):325-8.

CHARACTERISTICS OF STUDIES

Characteristics of included studies [ordered by study ID]

Bao 2012

Methods	<p>RCT</p> <p>Method of randomisation: random number table</p> <p>Blinding: not stated</p> <p>Adverse effects: not stated</p> <p>ITT analysis: not stated</p> <p>Losses to FU: not stated</p>
Participants	<p>Country: China</p> <p>Number of participants included: 60 (30/30)</p> <p>Demographics: aged 50-73 years, 67% male</p> <p>Type of stroke: ischaemic only</p> <p>Diagnosis: WHO definition and all confirmed by CT</p> <p>Severity on entry: not stated</p> <p>Time from stroke onset: 3-12 months</p> <p>Setting: inpatient and outpatient</p> <p>Comparability: no significant difference in age or time post onset</p>
Interventions	<p>Comparison: acupuncture + WM versus WM</p> <p>Acupuncture treatment</p> <ul style="list-style-type: none"> Acupuncture rationale: not stated

Bao 2012 (Continued)

- Needling details
 - Points used: both body and scalp acupoints. Numbers of points used: 4 scalp acupoints and 3 body acupoints
 - Depths of insertion: not stated
 - Deqi elicited: yes
 - Needle stimulation: manual
 - Needle retention time: 50 minutes
 - Needle type: hua tuo brand
- Treatment regimen
 - Number of treatment sessions: 56 sessions
 - Frequency of treatment: 7 sessions/week
 - Total course: 8 weeks
- Practitioner background: not stated
- Co-intervention: WM

Control interventions: WM

Outcomes	<ul style="list-style-type: none"> • Improvement of cognitive function (MMSE, MoCA) • Improvement of independence (Barthel index) • FU: 8 weeks
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomisation by using random number table
Allocation concealment (selection bias)	Unclear risk	Information on allocation concealment was not reported
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Information on attrition was not reported
Selective reporting (reporting bias)	Unclear risk	Free of selective reporting bias was assessed as 'unclear' due to some clinically important outcomes unstated, such as quality of life, mortality and adverse events
Other bias	Unclear risk	No information provided

Chou 2009

Methods	<p>RCT</p> <p>Method of randomisation: not stated</p> <p>Blinding: not stated</p> <p>Adverse effects: not stated</p> <p>ITT analysis: not stated</p> <p>Losses to FU: 5</p>
Participants	<p>Country: UK</p> <p>Number of participants included: 33 (17/16)</p> <p>Demographics: aged 59-90 years, 48% male</p> <p>Type of stroke: both ischaemic and haemorrhagic strokes</p> <p>Diagnosis: CT/MRI</p> <p>Severity on entry: unclear</p> <p>Time from stroke onset: 13-33 months</p> <p>Setting: inpatient</p> <p>Comparability: comorbidity and past history similar</p>
Interventions	<p>Comparison: real acupuncture + PT versus sham acupuncture + PT</p> <p>Acupuncture treatment</p> <ul style="list-style-type: none"> • Acupuncture rationale: stated • Needling details <ul style="list-style-type: none"> ◦ Points used: body acupoints ◦ Numbers of points used: 2 acupoints ◦ Depths of insertion: 0.03 inch ◦ Deqi elicited: yes ◦ Needle stimulation: electrical ◦ Needle retention time: 20 minutes ◦ Needle type: the needles were 1 inch long • Treatment regimen <ul style="list-style-type: none"> ◦ Number of treatment sessions: 16 sessions ◦ Frequency of treatment: 2/week ◦ Total course: 8 weeks • Practitioner background: has been practicing Traditional Chinese Medicine for many years after graduating from medical school (China Medical University, Taichung, Taiwan) • Co-intervention: PT <p>Control interventions: sham acupuncture + PT</p>
Outcomes	<ul style="list-style-type: none"> • Improvement of cognitive function (LOTCA-G) • Quality of life (SF-36, SS-QOL) • FU: 8 weeks
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	The method of random sequence generation was not reported

Chou 2009 (Continued)

Allocation concealment (selection bias)	Unclear risk	Information on allocation concealment was not reported
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Incomplete outcome data (attrition bias) All outcomes	Low risk	Treatment: 3 participants could not finish the treatment protocol Control: 2 participants decided not to receive rehabilitation
Selective reporting (reporting bias)	Unclear risk	Free of selective reporting bias was assessed as 'unclear' due to some clinically important outcomes unstated, such as quality of life, mortality and adverse events
Other bias	Unclear risk	No information provided

Dai 1997

Methods	<p>RCT Method of randomisation: not stated Blinding: not stated</p> <p>Adverse effects: not stated ITT analysis: not stated Losses to FU: not stated</p>
Participants	<p>Country: China Number of participants included: 136 (46/45/45) Demographics: aged 48-86 years, 75% male Type of stroke: ischaemic only Diagnosis: WHO definition and all confirmed by CT Severity on entry: mild to severe Time from stroke onset: 3-14 months Setting: unclear Comparability: unclear</p>
Interventions	<p>3 arms:</p> <ul style="list-style-type: none"> • acupuncture + WM • acupuncture only • WM only <p>Comparison eligible: acupuncture + WM versus WM only</p> <p>Acupuncture treatment</p> <ul style="list-style-type: none"> • Acupuncture rationale: not stated

Acupuncture for stroke rehabilitation (Review)

Dai 1997 (Continued)

- Needling details:
 - points used: both body and scalp (three temporal points) acupoints
 - Numbers of points used: 5-11 points
 - Depths of insertion: 1.5-2.0 inches
 - Deqi elicited: yes
 - Needle stimulation: manual
 - Needle retention time: 30 minutes
 - Needle type: stainless steel, gauge 26-30#, length 2.0-2.6 inches
- Treatment regimen:
 - Number of treatment sessions: 30 sessions
 - Frequency of treatment: 1 session/day
 - Total course: 30 days
- Practitioner background: not stated
- Co-intervention:
 - aspirin 25 mg qd orally

Control interventions: WM: aspirin 25 mg qd orally

Outcomes	Number of participants with improvement in global neurological deficit (CSRS 1 score decrease > 18%) at the end of treatment FU: 30 days
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	The method of random sequence generation was not reported
Allocation concealment (selection bias)	Unclear risk	Information on allocation concealment was not reported
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Information on attrition was not reported
Selective reporting (reporting bias)	Unclear risk	Free of selective reporting bias was assessed as 'unclear' due to some clinically important outcomes unstated, such as quality of life, mortality and adverse events
Other bias	Unclear risk	No information provided

Gao 2014a

Methods	<p>RCT</p> <p>Method of randomisation: random number table</p> <p>Blinding: not stated</p> <p>Adverse effects: not stated</p> <p>ITT analysis: not stated</p> <p>Losses to FU: none</p>
Participants	<p>Country: China</p> <p>Number of patients included: 60 (30/30)</p> <p>Demographics: 50% male</p> <p>Type of stroke: both ischaemic and haemorrhagic strokes</p> <p>Diagnosis: CT/MRI</p> <p>Severity on entry: not stated</p> <p>Time from stroke onset: 3-6 months</p> <p>Setting: outpatient</p> <p>Comparability: no significant difference in age or time post onset</p>
Interventions	<p>Comparison: acupuncture + WM versus WM</p> <p>Acupuncture treatment</p> <ul style="list-style-type: none"> • Acupuncture rationale: not stated • Needling details <ul style="list-style-type: none"> ◦ Points used: both body and scalp acupoints ◦ Numbers of points used: 1 scalp acupoint and 5 body acupoints ◦ Depths of insertion: 0.66-1.05 inches ◦ Deqi elicited: not stated ◦ Needle stimulation: manual ◦ Needle retention time: 30 minutes ◦ Needle type: hua tuo brand, length 1.57 inches • Treatment regimen <ul style="list-style-type: none"> ◦ Number of treatment sessions: 24 sessions ◦ Frequency of treatment: 6 sessions/week ◦ Total course: 4 weeks • Practitioner background: not stated • Co-intervention: WM <p>Control interventions: WM</p>
Outcomes	<ul style="list-style-type: none"> • Improvement of depression (HAMD) • Improvement of neurological function (MESS and SSS) • TCM syndrome scoring criteria
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomisation by using random number table
Allocation concealment (selection bias)	Unclear risk	Information on the allocation concealment was not reported

Gao 2014a (Continued)

Blinding (performance bias and detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Incomplete outcome data (attrition bias) All outcomes	Low risk	No participants lost to follow-up
Selective reporting (reporting bias)	Unclear risk	Free of selective reporting bias was assessed as 'unclear' due to some clinically important outcomes unstated, such as quality of life, mortality and adverse events
Other bias	Unclear risk	No information provided

Guo 2011

Methods	<p>RCT Method of randomisation: not stated Blinding: not stated</p> <p>Adverse effects: not stated ITT analysis: not stated Losses to FU: not stated</p>
Participants	<p>Country: China Number of participants included: 95 (32/31/32) Demographics: aged 40-65 years, 61% male Type of stroke: both ischaemic and haemorrhagic strokes Diagnosis: CT/MRI Severity on entry: not stated Time from stroke onset: 1-6 months Setting: inpatient Comparability: no significant difference in age or time post onset</p>
Interventions	<p>3 arms:</p> <ul style="list-style-type: none"> • acupuncture only • WM only • acupuncture + WM <p>Comparison eligible: acupuncture + WM versus WM only</p> <p>Acupuncture treatment</p> <ul style="list-style-type: none"> • Acupuncture rationale: not stated

Guo 2011 (Continued)

- Needling details
 - Points used: both body and scalp acupoints
 - Numbers of points used: 4 scalp acupoints and 2 body acupoints
 - Depths of insertion: not stated
 - Deqi elicited: not stated
 - Needle stimulation: manual and electrical
 - Needle retention time: 30 minutes
 - Needle type: not stated
- Treatment regimen
 - Number of treatment sessions: 30 sessions
 - Frequency of treatment: 5 sessions/week
 - Total course: 6 weeks
- Practitioner background: not stated
- Co-intervention: WM

Control interventions: WM

Outcomes	<ul style="list-style-type: none"> • Improvement of depression (HAMD) • Improvement of neurological function (Clinical Neurological Function Defect Scale) • FU: 6 weeks
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	The method of random sequence generation was not reported
Allocation concealment (selection bias)	Unclear risk	Information on allocation concealment was not reported
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Information on attrition was not reported
Selective reporting (reporting bias)	Unclear risk	Free of selective reporting bias was assessed as 'unclear' due to some clinically important outcomes unstated, such as quality of life, mortality and adverse events
Other bias	Unclear risk	No information provided

Guo 2012

Methods	<p>RCT</p> <p>Method of randomisation: not stated</p> <p>Blinding: not stated</p> <p>Adverse effects: not stated</p> <p>ITT analysis: not stated</p> <p>Losses to FU: not stated</p>
Participants	<p>Country: China</p> <p>Number of participants included: 60 (30/30)</p> <p>Demographics: aged 38 to 73 years, 68% male</p> <p>Type of stroke: unclear</p> <p>Diagnosis: unclear</p> <p>Severity on entry: not stated</p> <p>Time from stroke onset: 3 months to 5 years</p> <p>Setting: inpatient</p> <p>Comparability: comorbidity and past history similar</p>
Interventions	<p>Comparison: acupuncture + OT versus OT</p> <p>Acupuncture treatment</p> <ul style="list-style-type: none"> • Acupuncture rationale: not stated • Needling details <ul style="list-style-type: none"> ◦ Points used: body acupoints ◦ Numbers of points used: 1 acupoint ◦ Depths of insertion: 0.28-0.55 inch ◦ Deqi elicited: not stated ◦ Needle stimulation: manual ◦ Needle retention time: not stated ◦ Needle type: the needles were 1.57 inches long • Treatment regimen <ul style="list-style-type: none"> ◦ Number of treatment sessions: 6-12 sessions ◦ Frequency of treatment: 6/week ◦ Total course: 1-2 weeks • Practitioner background: not stated • Co-intervention: OT <p>Control interventions: OT</p>
Outcomes	Number of participants with improvement in swallowing function (water drinking test)
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	The method of random sequence generation was not reported
Allocation concealment (selection bias)	Unclear risk	Information on allocation concealment was not reported
Blinding (performance bias and detection bias)	Unclear risk	Information on blinding was not reported

Acupuncture for stroke rehabilitation (Review)

Guo 2012 (Continued)

All outcomes

Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Information on attrition was not reported
Selective reporting (reporting bias)	Unclear risk	Free of selective reporting bias was assessed as 'unclear' due to some clinically important outcomes unstated, such as quality of life, mortality and adverse events
Other bias	Unclear risk	No information provided

Huang 2008a

Methods	<p>RCT</p> <p>Method of randomisation: random number table</p> <p>Blinding: not stated</p> <p>Adverse effects: not stated</p> <p>ITT analysis: not stated</p> <p>Losses to FU: not stated</p>
Participants	<p>Country: China</p> <p>Number of participants included: 90 (30/30/30)</p> <p>Demographics: 67% male</p> <p>Type of stroke: ischaemic only</p> <p>Diagnosis: CT/MRI</p> <p>Severity on entry: not stated</p> <p>Time from stroke onset: 1-3 months</p> <p>Setting: inpatient</p> <p>Comparability: no significant difference in age or time post onset</p>
Interventions	<p>3 arms:</p> <ul style="list-style-type: none"> • acupuncture only • PT only • acupuncture + PT <p>Comparison eligible: acupuncture + PT versus PT only</p> <p>Acupuncture treatment</p> <ul style="list-style-type: none"> • Acupuncture rationale: not stated

Huang 2008a (Continued)

- Needling details
 - Points used: scalp acupoints
 - Numbers of points used: 3 points
 - Depths of insertion: 0.98-1.38 inches
 - Deqi elicited: not stated
 - Needle stimulation: manual
 - Needle retention time: 30 minutes
 - Needle type: the needles were 1.57 inches long
- Treatment regimen
 - Number of treatment sessions: 24 sessions
 - Frequency of treatment: 6 sessions/week
 - Total course: 4 weeks
- Practitioner background: not stated
- Co-intervention: PT

Control interventions: PT

Outcomes	<ul style="list-style-type: none"> • Neurological function (MESS and SSS) • Independence (Barthel Index)
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomisation by using random number table
Allocation concealment (selection bias)	Unclear risk	Information on allocation concealment was not reported
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	No information provided
Selective reporting (reporting bias)	Unclear risk	Free of selective reporting bias was assessed as 'unclear' due to some clinically important outcomes unstated, such as quality of life, mortality and adverse events
Other bias	Unclear risk	No information provided

Ke 2015

Methods	<p>RCT</p> <p>Method of randomisation: not stated</p> <p>Blinding: not stated</p> <p>Adverse effects: not stated</p> <p>ITT analysis: not stated</p> <p>Losses to FU: not stated</p>
Participants	<p>Country: China</p> <p>Number of participants included: 80 (40/40)</p> <p>Demographics: aged 61-83 years, 55% male</p> <p>Type of stroke: both ischaemic and haemorrhagic strokes</p> <p>Diagnosis: WHO definition</p> <p>Severity on entry: not stated</p> <p>Time from stroke onset: 2-18 months</p> <p>Setting: inpatient</p> <p>Comparability: no significant difference in age or time post onset</p>
Interventions	<p>Comparison: acupuncture + PT versus PT</p> <p>Acupuncture treatment</p> <ul style="list-style-type: none"> • Acupuncture rationale: not stated • Needling details <ul style="list-style-type: none"> ◦ Points used: body acupoints ◦ Numbers of points used: 27 body acupoints ◦ Depths of insertion: 0.66-5.25 inches ◦ Deqi elicited: not stated ◦ Needle stimulation: manual ◦ Needle retention time: 30-40 minutes ◦ Needle type: not stated • Treatment regimen <ul style="list-style-type: none"> ◦ Number of treatment sessions: 30 sessions ◦ Frequency of treatment: 7 sessions/week ◦ Total course: 30 days • Practitioner background: not stated • Co-intervention: PT <p>Control interventions: PT</p>
Outcomes	<ul style="list-style-type: none"> • Improvement of syndrome (TCM syndrome scoring criteria) • Improvement of independence (Barthel Index) • FU: 30 days
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	The method of random sequence generation was not reported
Allocation concealment (selection bias)	Unclear risk	Information on allocation concealment was not reported

Ke 2015 (Continued)

Blinding (performance bias and detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Information on attrition was not reported
Selective reporting (reporting bias)	Unclear risk	Free of selective reporting bias was assessed as 'unclear' due to some clinically important outcomes unstated, such as quality of life, mortality and adverse events
Other bias	Unclear risk	No information provided

Li 1997a

Methods	<p>RCT Method of randomisation: not stated Blinding: not stated</p> <p>Adverse effects: not stated ITT analysis: not stated Losses to FU: not stated</p>
Participants	<p>Country: China Number of participants included: 112 (42/20/50) Demographics: aged 24-76 years Type of stroke: both ischaemic and haemorrhagic strokes Diagnosis: WHO definition and all confirmed by CT or MRI Severity on entry: not stated Time from stroke onset: 1 month to 8.5 years Setting: inpatients Comparability: comorbidity and past history similar</p>
Interventions	<p>3 arms:</p> <ul style="list-style-type: none"> • acupuncture + PT and OT • acupuncture only • PT and OT <p>Comparison eligible: acupuncture + PT and OT versus PT and OT</p> <p>Acupuncture treatment</p> <ul style="list-style-type: none"> • Acupuncture rationale: not stated

Li 1997a (Continued)

- Needling details
 - Points used: both body and scalp acupoints
 - Numbers of points used: 10-12 points
 - Depths of insertion: not stated
 - Deqi elicited: unclear
 - Needle stimulation: manual
 - Needle retention time: 30 minutes
 - Needle type: not stated
- Treatment regimen
 - Number of treatment sessions: 72 sessions
 - Frequency of treatment: 6 sessions/week
 - Total course: 3 months
- Practitioner background: not stated
- Co-intervention: baseline medication, PT and OT

Control interventions: baseline medication plus PT and OT

Outcomes	Number of participants with improvement in global neurological deficit (CSRS 1 score) at the end of treatment FU: 3 months
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	The method of random sequence generation was not reported
Allocation concealment (selection bias)	Unclear risk	Information on allocation concealment was not reported
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Information on attrition was not reported
Selective reporting (reporting bias)	Unclear risk	Free of selective reporting bias was assessed as 'unclear' due to some clinically important outcomes unstated, such as quality of life, mortality and adverse events
Other bias	Unclear risk	No information provided

Li 2010a

Methods	<p>RCT</p> <p>Method of randomisation: random number table</p> <p>Blinding: not stated</p> <p>Adverse effects: none</p> <p>ITT analysis: not stated</p> <p>Losses to FU: 4</p>
Participants	<p>Country: China</p> <p>Number of participants included: 46 (24/22)</p> <p>Demographics: aged 47-79 years, 59% male</p> <p>Type of stroke: both ischaemic and haemorrhagic strokes</p> <p>Diagnosis: CT/MRI</p> <p>Severity on entry: not stated</p> <p>Time from stroke onset: 1-36 months</p> <p>Setting: outpatient</p> <p>Comparability: no significant difference in age or time post onset</p>
Interventions	<p>Comparison: acupuncture + WM versus WM</p> <p>Acupuncture treatment</p> <ul style="list-style-type: none"> • Acupuncture rationale: not stated • Needling details <ul style="list-style-type: none"> ◦ Points used: both body and scalp acupoints ◦ Numbers of points used: 5 scalp acupoints and 6 body acupoints ◦ Depths of insertion: 0.66-1.5 inches ◦ Deqi elicited: yes ◦ Needle stimulation: manual and electrical ◦ Needle retention time: 30 minutes ◦ Needle type: gauge 30#, length 1.97 inches • Treatment regimen <ul style="list-style-type: none"> ◦ Number of treatment sessions: 60 sessions ◦ Frequency of treatment: 5 sessions/week ◦ Total course: 12 weeks • Practitioner background: not stated • Co-intervention: WM <p>Control interventions: WM</p>
Outcomes	<ul style="list-style-type: none"> • Improvement of cognitive function (MMSE HDS-R) • Improvement of independence (Barthel Index) • Adverse events • FU: 12 weeks
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomisation by using random number table

Li 2010a (Continued)

Allocation concealment (selection bias)	Unclear risk	Information on allocation concealment was not reported
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Incomplete outcome data (attrition bias) All outcomes	Low risk	Treatment: 1 participant could not finish the treatment protocol Control: 3 participants could not finish the control protocol
Selective reporting (reporting bias)	Unclear risk	Some patient-related outcomes were not reported, such as information on quality of life or all-cause mortality
Other bias	Unclear risk	No information provided

Li 2011a

Methods	RCT Method of randomisation: not stated Blinding: not stated Adverse effects: not stated ITT analysis: not stated Losses to FU: not stated
Participants	Country: China Number of participants included: 62 (31/31) Demographics: 54% male Type of stroke: unclear Diagnosis: CT/MRI Severity on entry: not stated Time from stroke onset: 6-20 months Setting: outpatient Comparability: no significant difference in age or time post onset
Interventions	Comparison: acupuncture + WM versus WM Acupuncture treatment • Acupuncture rationale: not stated

Li 2011a (Continued)

- Needling details
 - Points used: body acupoints
 - Numbers of points used: 7 body acupoints
 - Depths of insertion: 0.39-1.05 inches
 - Deqi elicited: not stated
 - Needle stimulation: manual
 - Needle retention time: not stated
 - Needle type: stainless steel needle gauge 28#, length 1.31 inches
- Treatment regimen
 - Number of treatment sessions: 28 sessions
 - Frequency of treatment: 7 sessions/week
 - Total course: 4 weeks
- Practitioner background: not stated
- Co-intervention: WM

Control interventions: WM

Outcomes	<ul style="list-style-type: none"> • Improvement of cognitive function (MMSE) • Improvement of independence (ADL) • FU: 4 weeks
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	The method of random sequence generation was not reported
Allocation concealment (selection bias)	Unclear risk	Information on allocation concealment was not reported
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Information on attrition was not reported
Selective reporting (reporting bias)	Unclear risk	Free of selective reporting bias was assessed as 'unclear' due to some clinically important outcomes unstated, such as quality of life, mortality and adverse events
Other bias	Unclear risk	No information provided

Li 2013a

Methods	<p>RCT</p> <p>Method of randomisation: not stated</p> <p>Blinding: not stated</p> <p>Adverse effects: not stated</p> <p>ITT analysis: not stated</p> <p>Losses to FU: not stated</p>
Participants	<p>Country: China</p> <p>Number of participants included: 60 (20/20/20)</p> <p>Demographics: aged 40-75 years</p> <p>Type of stroke: both ischaemic and haemorrhagic strokes</p> <p>Diagnosis: CT/MRI</p> <p>Severity on entry: not stated</p> <p>Time from stroke onset: 1-3 months</p> <p>Setting: inpatient</p> <p>Comparability: no significant difference in age or time post onset</p>
Interventions	<p>3 arms:</p> <ul style="list-style-type: none"> • acupuncture only • PT only • acupuncture + PT <p>Comparison eligible: acupuncture + PT versus PT only</p> <p>Acupuncture treatment</p> <ul style="list-style-type: none"> • Acupuncture rationale: not stated • Needling details <ul style="list-style-type: none"> ◦ Points used: body acupoints ◦ Numbers of points used: 12 points ◦ Depths of insertion: not stated ◦ Deqi elicited: not stated ◦ Needle stimulation: manual ◦ Needle retention time: 30 minute ◦ Needle type: not stated • Treatment regimen <ul style="list-style-type: none"> ◦ Number of treatment sessions: not stated ◦ Frequency of treatment: 7 sessions/week ◦ Total course: not stated • Practitioner background: not stated • Co-intervention: PT <p>Control interventions: PT</p>
Outcomes	<ul style="list-style-type: none"> • Improvement of motor function (Fugl-Meyer Assessment) • Improvement of pain (Visual Analogue Score) • Improvement of independence (ADL) • FU: not stated
Notes	-

Risk of bias

Li 2013a (Continued)

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	The method of random sequence generation was not reported
Allocation concealment (selection bias)	Unclear risk	Information on allocation concealment was not reported
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Information on attrition was not reported
Selective reporting (reporting bias)	Unclear risk	Free of selective reporting bias was assessed as 'unclear' due to some clinically important outcomes unstated, such as quality of life, mortality and adverse events
Other bias	Unclear risk	No information provided

Liu 2013a

Methods	<p>RCT Method of randomisation: not stated Blinding: not stated</p> <p>Adverse effects: not stated ITT analysis: not stated Losses to FU: not stated</p>
Participants	<p>Country: China Number of participants included: 50 (25/25) Demographics: aged 35-74 years, 68% male Type of stroke: both ischaemic and haemorrhagic strokes Diagnosis: WHO definition and all confirmed by CT Severity on entry: not stated Time from stroke onset: 3-12 months Setting: inpatient Comparability: no significant difference in age, comorbidity, or time post onset</p>
Interventions	<p>Comparison: acupuncture + daily OT versus daily OT</p> <p>Acupuncture treatment</p> <ul style="list-style-type: none"> Acupuncture rationale: not stated

Liu 2013a (Continued)

- Needling details
 - Points used: scalp acupoints
 - Numbers of points used: 2 scalp acupoints
 - Depths of insertion: 1.05-1.31/0.66 inches
 - Deqi elicited: yes
 - Needle stimulation: manual and electrical stimulation
 - Needle retention time: 90 minutes
 - Needle type: hua tuo brand, length 0.98 inches
- Treatment regimen
 - Number of treatment sessions: 24 sessions
 - Frequency of treatment: 6 sessions/week
 - Total course: 4 weeks
- Practitioner background: not stated
- Co-intervention: OT

Control interventions: OT

Outcomes	<ul style="list-style-type: none"> • Improvement of cognitive function (MMSE) • Independence (ADL) • FU: 24 days
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	The method of random sequence generation was not reported
Allocation concealment (selection bias)	Unclear risk	Information on allocation concealment was not reported
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Information on attrition was not reported
Selective reporting (reporting bias)	Unclear risk	Free of selective reporting bias was assessed as 'unclear' due to some clinically important outcomes unstated, such as quality of life, mortality and adverse events
Other bias	Unclear risk	No information provided

Lun 1999

Methods	<p>RCT</p> <p>Method of randomisation: not stated</p> <p>Blinding: not stated</p> <p>Adverse effects: not stated</p> <p>ITT analysis: not stated</p> <p>Losses to FU: not stated</p>
Participants	<p>Country: China</p> <p>Number of participants included: 109 (61/48)</p> <p>Demographics: aged 35-75 years, 60% male</p> <p>Type of stroke: both ischaemic and haemorrhagic strokes</p> <p>Diagnosis: WHO definition and all confirmed by CT before entry</p> <p>Severity on entry: not stated</p> <p>Time from stroke onset: 2 months to 5 years</p> <p>Setting: unclear</p> <p>Comparability: unclear</p>
Interventions	<p>Comparison: acupuncture + TCM versus TCM</p> <p>Acupuncture treatment</p> <ul style="list-style-type: none"> • Acupuncture rationale: not stated • Needling details <ul style="list-style-type: none"> ◦ Points used: scalp acupoints ◦ Numbers of points used: 5 points ◦ Depths of insertion: 2-2.6 inches ◦ Deqi elicited: yes ◦ Needle stimulation: manual ◦ Needle retention time: 30 minute ◦ Needle type: stainless steel needle, gauge 28-30#, length 2.6 inches • Treatment regimen <ul style="list-style-type: none"> ◦ Number of treatment sessions: 30 sessions ◦ Frequency of treatment: 1/day ◦ Total course: 45 days • Practitioner background: not stated • Co-intervention: baseline TCM <p>Control interventions: TCM</p>
Outcomes	<p>Number of participants with improvement in global neurological deficit (CSRS 2) at the end of treatment</p> <p>FU: 45 days</p>
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	The method of random sequence generation was not reported
Allocation concealment (selection bias)	Unclear risk	Information on allocation concealment was not reported

Lun 1999 (Continued)

Blinding (performance bias and detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Information on attrition was not reported
Selective reporting (reporting bias)	Unclear risk	Free of selective reporting bias was assessed as 'unclear' due to some clinically important outcomes unstated, such as quality of life, mortality and adverse events
Other bias	Unclear risk	No information provided

Naeser 1992

Methods	<p>RCT Method of randomisation: not stated Blinding: participants blinded</p> <p>Adverse effects: not stated ITT analysis: not stated Losses to FU: not stated</p>
Participants	<p>Country: USA Number of participants included: 16 (10/6) Demographics: aged 44-74 years Type of stroke: ischaemic only Diagnosis: all confirmed by CT Severity on entry: moderate Time from stroke onset: 1-3 months Setting: inpatients Comparability: no significant difference in age or time post onset</p>
Interventions	<p>Comparison: real acupuncture + daily PT versus sham acupuncture + daily PT</p> <p>Acupuncture treatment</p> <ul style="list-style-type: none"> • Acupuncture rationale: not stated • Needling details <ul style="list-style-type: none"> ◦ Points used: both body and scalp acupoints ◦ Numbers of points used: 11 body acupoints; numbers of scalp acupoints unclear ◦ Depths of insertion: unclear ◦ Deqi elicited: unclear ◦ Needle stimulation: electrical stimulation with frequency 1-2 Hz and amplitude unknown ◦ Needle retention time: 20 minutes ◦ Needle type: gauge 34#, length and material unknown

Naeser 1992 (Continued)

- Treatment regimen
 - Number of treatment sessions: 20 sessions
 - Frequency of treatment: 5 sessions/week
 - Total course: 4 weeks
- Practitioner background: not stated
- Co-intervention: PT

Control interventions

- Sham
- Acupuncture + PT

Outcomes	Number of participants with improvement in motor function (BMIT) within 5 days after completing treatment FU: 35 days
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	The method of random sequence generation was not reported
Allocation concealment (selection bias)	Unclear risk	Information on allocation concealment was not reported
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: "The sham acupuncture was done only on the non-paralyzed upper extremity and lower extremity. The patients were told that in China acupuncture is used on the non-paralysed side to treat the paralysed side. They were further told that the stimulation was low level, and they would not feel anything"
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Information on attrition was not reported
Selective reporting (reporting bias)	Unclear risk	Free of selective reporting bias was assessed as 'unclear' due to some clinically important outcomes unstated, such as quality of life, mortality and adverse events
Other bias	Unclear risk	No information provided

Sun 2013a

Methods	RCT Method of randomisation: random number table Blinding: not stated
---------	---

Acupuncture for stroke rehabilitation (Review)

Sun 2013a (Continued)

	Adverse effects: 1 ITT analysis: not stated Losses to FU: 7
Participants	Country: China Number of participants included: 71 (36/35) Demographics: aged 51-72 years, 56% male Type of stroke: both ischaemic and haemorrhagic strokes Diagnosis: WHO definition Severity on entry: not stated Time from stroke onset: 3-9 months Setting: inpatient Comparability: no significant difference in age or time post onset
Interventions	Comparison: acupuncture + OT + WM versus OT + WM Acupuncture treatment <ul style="list-style-type: none"> • Acupuncture rationale: not stated • Needling details <ul style="list-style-type: none"> ◦ Points used: both body and scalp acupoints ◦ Numbers of points used: 7 scalp acupoints and 5 body acupoints ◦ Depths of insertion: 0.26-1.97 inches ◦ Deqi elicited: yes ◦ Needle stimulation: manual and electrical ◦ Needle retention time: 30 minutes ◦ Needle type: hua tuo brand, length 1.31-1.97 inches • Treatment regimen <ul style="list-style-type: none"> ◦ Number of treatment sessions: 45 sessions ◦ Frequency of treatment: 1 session/day ◦ Total course: 45 days • Practitioner background: not stated • Co-intervention: OT + WM Control interventions: OT + WM
Outcomes	<ul style="list-style-type: none"> • Improvement of symptoms of turbid phlegm obstructing orifices • Improvement of cognitive function (MMSE) • Improvement of independence (ADL) • Improvement of neurological function (NDFS) • Improvement of quality of life (MOS, SF-36) • Adverse event: itchiness of the skin at the acupoint in one participant in the acupuncture group • FU: 45 days
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomisation by using random number table
Allocation concealment (selection bias)	Unclear risk	Information on allocation concealment was not reported

Sun 2013a (Continued)

Blinding (performance bias and detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Incomplete outcome data (attrition bias) All outcomes	Low risk	4 participants could not finish the treatment protocol, 3 participants lost to follow-up
Selective reporting (reporting bias)	Unclear risk	Some patient-related outcomes were not reported, such as information on quality of life or all-cause mortality
Other bias	Unclear risk	No information provided

Sun 2015

Methods	<p>RCT</p> <p>Method of randomisation: randomisation by using envelopes</p> <p>Blinding: not stated</p> <p>Adverse effects: none</p> <p>ITT analysis: not stated</p> <p>Losses to FU: not stated</p>
Participants	<p>Country: China</p> <p>Number of participants included: 93 (31/31/31)</p> <p>Demographics: aged 60-95 years, 47% male</p> <p>Type of stroke: unclear</p> <p>Diagnosis: WHO definition</p> <p>Severity on entry: not stated</p> <p>Time from stroke onset: 9-46 months</p> <p>Setting: inpatient and outpatient</p> <p>Comparability: no significant difference in age, comorbidity, or time post onset</p>
Interventions	<p>3 arms:</p> <ul style="list-style-type: none"> • acupuncture only • WM only • acupuncture + WM <p>Comparison eligible: acupuncture + WM versus WM only</p> <p>Acupuncture treatment</p> <ul style="list-style-type: none"> • Acupuncture rationale: not stated

Sun 2015 (Continued)

- Needling details
 - Points used: both body and scalp acupoints
 - Numbers of points used: 6 body acupoints, 3 scalp acupoints
 - Depths of insertion: not stated
 - Deqi elicited: yes
 - Needle stimulation: manual
 - Needle retention time: 30 minutes
 - Needle type: hua tuo brand, length 1.57 inches
- Treatment regimen
 - Number of treatment sessions: 120 sessions
 - Frequency of treatment: 5 sessions/week
 - Total course: 24 weeks
- Practitioner background: not stated
- Co-intervention: WM

Control interventions: WM

Outcomes	<ul style="list-style-type: none"> • Improvement of depression (HAMD) • Adverse event
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomisation by using envelopes assorted in a random manner
Allocation concealment (selection bias)	Unclear risk	Information on allocation concealment was not reported
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Information on attrition was not reported
Selective reporting (reporting bias)	Unclear risk	Some patient-related outcomes were not reported, such as information on quality of life or all-cause mortality
Other bias	Unclear risk	No information provided

Wang 2001

Methods	<p>RCT</p> <p>Method of randomisation: not stated</p> <p>Blinding: not stated</p> <p>Adverse effects: not stated</p> <p>ITT analysis: not stated</p> <p>Losses to FU: not stated</p>
Participants	<p>Country: China</p> <p>Number of participants included: 90 (34/30/26)</p> <p>Demographics: aged 39-75 years, 56% male</p> <p>Type of stroke: both ischaemic and haemorrhagic strokes</p> <p>Diagnosis: WHO definition and all confirmed by CT</p> <p>Severity on entry: not stated</p> <p>Time from stroke onset: 2 months to 5 years</p> <p>Setting: unclear</p> <p>Comparability: unclear</p>
Interventions	<p>3 arms:</p> <ul style="list-style-type: none"> • acupuncture + TCM • acupuncture only • TCM only <p>Comparison eligible: acupuncture + TCM versus TCM only</p> <p>Acupuncture treatment</p> <ul style="list-style-type: none"> • Acupuncture rationale: not stated • Needling details <ul style="list-style-type: none"> ◦ Points used: body acupoints ◦ Numbers of points used: 6-10 points ◦ Depths of insertion: not stated ◦ Deqi elicited: yes ◦ Needle stimulation: manual ◦ Needle retention time: 30 minutes ◦ Needle type: not stated • Treatment regimen <ul style="list-style-type: none"> ◦ Number of treatment sessions: 60 sessions ◦ Frequency of treatment: 10 sessions/13 days ◦ Total course: 78 days • Practitioner background: not stated • Co-intervention: TCM <p>Control interventions: TCM</p>
Outcomes	<ul style="list-style-type: none"> • Number of participants with improvement in global neurological deficit (CSRS 2) at the end of treatment • FU: 78 days
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Wang 2001 (Continued)

Random sequence generation (selection bias)	Unclear risk	The method of random sequence generation was not reported
Allocation concealment (selection bias)	Unclear risk	Information on allocation concealment was not reported
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Information on attrition was not reported
Selective reporting (reporting bias)	Unclear risk	Free of selective reporting bias was assessed as 'unclear' due to some clinically important outcomes unstated, such as quality of life, mortality and adverse events
Other bias	Unclear risk	No information provided

Wang 2011a

Methods	<p>RCT Method of randomisation: random number table Blinding: not stated</p> <p>Adverse effects: not stated ITT analysis: not stated Losses to FU: not stated</p>
Participants	<p>Country: China Number of participants included: 85 (43/42) Demographics: aged 40-80 years, 65% male Type of stroke: both ischaemic and haemorrhagic strokes Diagnosis: CT/MRI Severity on entry: not stated Time from stroke onset: 33-192 days Setting: inpatient and outpatient Comparability: no significant difference in age, comorbidity, or time post onset</p>
Interventions	<p>Comparison: acupuncture + PT + OT versus PT + OT</p> <p>Acupuncture treatment</p> <ul style="list-style-type: none"> Acupuncture rationale: not stated

Wang 2011a (Continued)

- Needling details
 - Points used: body acupoints
 - Numbers of points used: 10 acupoints
 - Depths of insertion: 0.08-0.2 inches
 - Deqi elicited: not stated
 - Needle stimulation: manual
 - Needle retention time: not stated
 - Needle type: not stated
- Treatment regimen
 - Number of treatment sessions: 10 sessions
 - Frequency of treatment: 3-4 sessions/week
 - Total course: 3 weeks
- Practitioner background: not stated
- Co-intervention: PT + OT

Control interventions: PT + OT

Outcomes	<ul style="list-style-type: none"> • Improvement of motor function (Fugl-Meyer Assessment) • Improvement of pain (Visual Analogue Scale) • FU: 3 weeks
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomisation by using random number table
Allocation concealment (selection bias)	Unclear risk	Information on allocation concealment was not reported
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Information on attrition was not reported
Selective reporting (reporting bias)	Unclear risk	Free of selective reporting bias was assessed as 'unclear' due to some clinically important outcomes unstated, such as quality of life, mortality and adverse events
Other bias	Unclear risk	No information provided

Wang 2012

Methods	<p>RCT</p> <p>Method of randomisation: random number table</p> <p>Blinding: not stated</p> <p>Adverse effects: not stated</p> <p>ITT analysis: not stated</p> <p>Losses to FU: not stated</p>
Participants	<p>Country: China</p> <p>Number of participants included: 60 (30/30)</p> <p>Demographics: aged 43-77 years, 65% male</p> <p>Type of stroke: both ischaemic and haemorrhagic strokes</p> <p>Diagnosis: WHO definition and all confirmed by CT/MRI</p> <p>Severity on entry: not stated</p> <p>Time from stroke onset: 1-6 months</p> <p>Setting: inpatient</p> <p>Comparability: no significant difference in age or time post onset</p>
Interventions	<p>Comparison: acupuncture + PT versus PT</p> <p>Acupuncture treatment</p> <ul style="list-style-type: none"> • Acupuncture rationale: not stated • Needling details <ul style="list-style-type: none"> ◦ Points used: scalp acupoints ◦ Numbers of points used: 4 points ◦ Depths of insertion: 0.98-1.38 inches ◦ Deqi elicited: not stated ◦ Needle stimulation: manual ◦ Needle retention time: 30 minutes ◦ Needle type: stainless steel needle, gauge 30#, length 1.97 inches • Treatment regimen <ul style="list-style-type: none"> ◦ Number of treatment sessions: 45 sessions ◦ Frequency of treatment: 15 sessions/month ◦ Total course: 3 months • Practitioner background: not stated • Co-intervention: PT <p>Control interventions: PT</p>
Outcomes	<ul style="list-style-type: none"> • Improvement of motor function (Motor Assessment Scale) • Improvement of independence (modified Barthel Index) • FU: 3 months
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomisation by using random number table
Allocation concealment (selection bias)	Unclear risk	Information on allocation concealment was not reported

Wang 2012 (Continued)

Blinding (performance bias and detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Information on attrition was not reported
Selective reporting (reporting bias)	Unclear risk	Free of selective reporting bias was assessed as 'unclear' due to some clinically important outcomes unstated, such as quality of life, mortality and adverse events
Other bias	Unclear risk	No information provided

Wu 2008

Methods	<p>RCT Method of randomisation: not stated Blinding: not stated</p> <p>Adverse effects: not stated ITT analysis: not stated Losses to FU: not stated</p>
Participants	<p>Country: China Number of participants included: 160 (40/40/40/40) Demographics: aged 40-69 years, 63% male Type of stroke: both ischaemic and haemorrhagic strokes Diagnosis: CT/MRI Severity on entry: not stated Time from stroke onset: 1-6 months Setting: inpatient and outpatient Comparability: no significant difference in age, comorbidity, or time post onset</p>
Interventions	<p>4 arms:</p> <ul style="list-style-type: none"> PT + OT acupuncture + PT + OT WM + PT + OT acupuncture + WM + PT + OT <p>Comparisons eligible:</p> <ul style="list-style-type: none"> acupuncture + PT + OT versus PT + OT acupuncture + WM + PT + OT versus WM + PT + OT <p>Acupuncture treatment</p>

Wu 2008 (Continued)

- Acupuncture rationale: not stated
- Needling details
 - Points used: both body and scalp acupoints
 - Numbers of points used: 2 scalp acupoints and 2 body acupoints
 - Depths of insertion: not stated
 - Deqi elicited: not stated
 - Needle stimulation: manual
 - Needle retention time: not stated
 - Needle type: not stated
- Treatment regimen
 - Number of treatment sessions: 56 sessions
 - Frequency of treatment: 7 sessions/week
 - Total course: 8 weeks
- Practitioner background: not stated
- Co-intervention: PT + OT/WM + PT + OT

Control interventions: PT + OT/WM + PT + OT

Outcomes	<ul style="list-style-type: none"> • Improvement of depression (HAMD)
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	The method of random sequence generation was not reported
Allocation concealment (selection bias)	Unclear risk	Information on allocation concealment was not reported
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Information on attrition was not reported
Selective reporting (reporting bias)	Unclear risk	Free of selective reporting bias was assessed as 'unclear' due to some clinically important outcomes unstated, such as quality of life, mortality and adverse events
Other bias	Unclear risk	No information provided

Wu 2011a

Methods	<p>RCT</p> <p>Method of randomisation: randomisation by using random number table</p> <p>Blinding: not stated</p> <p>Adverse effects: not stated</p> <p>ITT analysis: not stated</p> <p>Losses to FU: not stated</p>
Participants	<p>Country: China</p> <p>Number of participants included: 60 (30/30)</p> <p>Demographics: aged 41-76 years, 55% male</p> <p>Type of stroke: ischaemic only</p> <p>Diagnosis: confirmed by CT/MRI</p> <p>Severity on entry: not stated</p> <p>Time from stroke onset: 1 month to 8 years</p> <p>Setting: inpatient</p> <p>Comparability: no significant difference in age or time post onset</p>
Interventions	<p>Comparison: acupuncture + daily PT versus daily PT</p> <p>Acupuncture treatment</p> <ul style="list-style-type: none"> • Acupuncture rationale: not stated • Needling details <ul style="list-style-type: none"> ◦ Points used: both body and scalp acupoints ◦ Numbers of points used: 5 scalp acupoints and 16 body acupoints ◦ Depths of insertion: not stated ◦ Deqi elicited: yes ◦ Needle stimulation: electrical ◦ Needle retention time: 30 minutes ◦ Needle type: hua tuo brand, length 1.57 inches • Treatment regimen <ul style="list-style-type: none"> ◦ Number of treatment sessions: 84 sessions ◦ Frequency of treatment: 7 sessions/week ◦ Total course: 12 weeks • Practitioner background: not stated • Co-intervention: PT <p>Control interventions: PT</p>
Outcomes	<ul style="list-style-type: none"> • Improvement of spasticity (Ashworth) • Improvement of motor function (Fugl-Meyer Assessment) • Improvement of independence (Barthel Index) • FU: 12 weeks
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomisation by using random number table
Allocation concealment (selection bias)	Unclear risk	Information on allocation concealment was not reported

Acupuncture for stroke rehabilitation (Review)

Wu 2011a (Continued)

Blinding (performance bias and detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Information on attrition was not reported
Selective reporting (reporting bias)	Unclear risk	Free of selective reporting bias was assessed as 'unclear' due to some clinically important outcomes unstated, such as quality of life, mortality and adverse events
Other bias	Unclear risk	No information provided

Wu 2013a

Methods	<p>RCT Method of randomisation: random number table Blinding: not stated</p> <p>Adverse effects: not stated ITT analysis: not stated Losses to FU: not stated</p>
Participants	<p>Country: China Number of participants included: 90 (30/30/30) Demographics: 45-80 years, 57% male Type of stroke: both ischaemic and haemorrhagic strokes Diagnosis: WHO definition and all confirmed by CT/MRI Severity on entry: not stated Time from stroke onset: 1-3 months Setting: inpatient Comparability: no significant difference in age or time post onset</p>
Interventions	<p>3 arms:</p> <ul style="list-style-type: none"> • routine acupuncture + OT + WM • acupuncture kinesitherapy simultaneously + OT + WM • OT + WM <p>Comparisons eligible:</p> <ul style="list-style-type: none"> • routine acupuncture + OT + WM versus OT + WM • acupuncture kinesitherapy simultaneously + OT + WM versus OT + WM <p>Acupuncture treatment</p> <ul style="list-style-type: none"> • Acupuncture rationale: not stated

Wu 2013a (Continued)

- Needling details
 - Points used: scalp acupoints
 - Numbers of points used: 7 points
 - Depths of insertion: 0.66-1.57 inches
 - Deqi elicited: not stated
 - Needle stimulation: manual
 - Needle retention time: 30 minutes
 - Needle type: length 1.57 inches
- Treatment regimen
 - Number of treatment sessions: 30 sessions
 - Frequency of treatment: 5 sessions/week
 - Total course: 6 weeks
- Practitioner background: not stated
- Co-intervention: OT + WM

Control interventions: OT + WM

Outcomes	<ul style="list-style-type: none"> • Improvement of swallowing function (Water drinking test) • FU: 6 weeks
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomisation by using random number table
Allocation concealment (selection bias)	Unclear risk	Information on allocation concealment was not reported
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Information on attrition was not reported
Selective reporting (reporting bias)	Unclear risk	Free of selective reporting bias was assessed as 'unclear' due to some clinically important outcomes unstated, such as quality of life, mortality and adverse events
Other bias	Unclear risk	No information provided

Xu 2013

Methods	<p>RCT</p> <p>Method of randomisation: not stated</p> <p>Blinding: not stated</p> <p>Adverse effects: not stated</p> <p>ITT analysis: not stated</p> <p>Losses to FU: not stated</p>
Participants	<p>Country: China</p> <p>Number of participants included: 140 (70/70)</p> <p>Demographics: 45-75 years, 47% male</p> <p>Type of stroke: unclear</p> <p>Diagnosis: WHO definition and all confirmed by CT/MRI</p> <p>Severity on entry: not stated</p> <p>Time from stroke onset: 1-12 months</p> <p>Setting: inpatient</p> <p>Comparability: no significant difference in age or time post onset</p>
Interventions	<p>Comparison: acupuncture + WM versus WM</p> <p>Acupuncture treatment</p> <ul style="list-style-type: none"> • Acupuncture rationale: not stated • Needling details <ul style="list-style-type: none"> ◦ Points used: both body and scalp acupoints ◦ Numbers of points used: 4 scalp acupoints and 16 body acupoints ◦ Depths of insertion: 1.18-1.97 inches ◦ Deqi elicited: yes ◦ Needle stimulation: manual ◦ Needle retention time: 30 minutes ◦ Needle type: stainless steel needle, length 1.97 inches • Treatment regimen <ul style="list-style-type: none"> ◦ Number of treatment sessions: 80 sessions ◦ Frequency of treatment: 2 sessions/day ◦ Total course: 40 days • Practitioner background: not stated • Co-intervention: WM <p>Control interventions: WM</p>
Outcomes	<ul style="list-style-type: none"> • Improvement of swallowing function (Water drinking test) • FU: 40 days
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	The method of random sequence generation was not reported
Allocation concealment (selection bias)	Unclear risk	Information on allocation concealment was not reported

Xu 2013 (Continued)

Blinding (performance bias and detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Information on attrition was not reported
Selective reporting (reporting bias)	Unclear risk	Free of selective reporting bias was assessed as 'unclear' due to some clinically important outcomes unstated, such as quality of life, mortality and adverse events
Other bias	Unclear risk	No information provided

Yao 2014

Methods	<p>RCT Method of randomisation: random number table Blinding: not stated</p> <p>Adverse effects: not stated ITT analysis: not stated Losses to FU: not stated</p>
Participants	<p>Country: China Number of participants included: 90 (30/30/30) Demographics: 54% male Type of stroke: both ischaemic and haemorrhagic strokes Diagnosis: WHO definition and all confirmed by CT/MRI Severity on entry: not stated Time from stroke onset: 1-12 months Setting: inpatient Comparability: no significant difference in age or time post onset</p>
Interventions	<p>Comparison eligible:</p> <ul style="list-style-type: none"> • routine acupuncture + PT versus PT only <p>Acupuncture treatment</p> <ul style="list-style-type: none"> • Acupuncture rationale: not stated

Yao 2014 (Continued)

- Needling details
 - Points used: both body and scalp acupoints
 - Numbers of points used: 15 points
 - Depths of insertion: not stated
 - Deqi elicited: yes
 - Needle stimulation: manual and electrical
 - Needle retention time: 30 minutes
 - Needle type: stainless steel needle, length 1.57 inches
- Treatment regimen
 - Number of treatment sessions: 24 sessions
 - Frequency of treatment: 3 sessions/week
 - Total course: 8 weeks
- Practitioner background: not stated
- Co-intervention: PT

Control interventions: PT

Outcomes	<ul style="list-style-type: none"> • Improvement of motor function (Fugl-Meyer Motor Assessment) • Improvement of neurological function (Neurological Deficit Score) • Improvement of independence (Barthel Index) • Improvement of pain (Visual Analogue Score) • FU: 8 weeks
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomisation by using random number table
Allocation concealment (selection bias)	Unclear risk	Information on allocation concealment was not reported
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Information on attrition was not reported
Selective reporting (reporting bias)	Unclear risk	Free of selective reporting bias was assessed as 'unclear' due to some clinically important outcomes unstated, such as quality of life, mortality and adverse events

Yao 2014 (Continued)

Other bias	Unclear risk	No information provided
------------	--------------	-------------------------

Zhan 2014

Methods	<p>RCT</p> <p>Method of randomisation: random number table</p> <p>Blinding: not stated</p> <p>Adverse effects: not stated</p> <p>ITT analysis: not stated</p> <p>Losses to FU: none</p>
Participants	<p>Country: China</p> <p>Number of participants included: 130 (65/65)</p> <p>Demographics: aged 45-75 years, 59% male</p> <p>Type of stroke: unclear</p> <p>Diagnosis: WHO definition</p> <p>Severity on entry: not stated</p> <p>Time from stroke onset: 2-11 months</p> <p>Setting: inpatient and outpatient</p> <p>Comparability: no significant difference in age or time post onset</p>
Interventions	<p>Comparison: acupuncture + PT versus PT</p> <p>Acupuncture treatment</p> <ul style="list-style-type: none"> Acupuncture rationale: not stated Needling details <ul style="list-style-type: none"> Points used: body acupoints Numbers of points used: 8 points Depths of insertion: not stated Deqi elicited: not stated Needle stimulation: manual Needle retention time: 15-20 minutes Needle type: not stated Treatment regimen <ul style="list-style-type: none"> Number of treatment sessions: 6 sessions Frequency of treatment: 6 sessions/week Total course: 1 week Practitioner background: not stated Co-intervention: PT <p>Control interventions: PT</p>
Outcomes	<ul style="list-style-type: none"> Improvement of independence (Barthel Index) FU: 1 week
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomisation by using random number table

Zhan 2014 (Continued)

Allocation concealment (selection bias)	Unclear risk	Information on allocation concealment was not reported
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Incomplete outcome data (attrition bias) All outcomes	Low risk	No participants withdrew or were lost to follow-up
Selective reporting (reporting bias)	Unclear risk	Free of selective reporting bias was assessed as 'unclear' due to some clinically important outcomes unstated, such as quality of life, mortality and adverse events
Other bias	Unclear risk	No information provided

Zhang 2013a

Methods	<p>RCT Method of randomisation: closed envelopes Blinding: Investigators, participants, not blinded Adverse effects: not stated</p> <p>ITT analysis: none Losses to FU: not stated</p>
Participants	<p>Country: China Number of participants included: 90 (30/30/30) Demographics: 43-75 years, 54% male Type of stroke: both ischaemic and haemorrhagic strokes Diagnosis: WHO definition and all confirmed by CT/MRI Severity on entry: not stated Time from stroke onset: 6-19 months Setting: inpatient Comparability: no significant difference in age or time post onset</p>
Interventions	<p>3 arms:</p> <ul style="list-style-type: none"> • acupuncture only • WM only • acupuncture + WM <p>Comparison eligible: acupuncture + WM versus WM</p> <p>Acupuncture treatment</p> <ul style="list-style-type: none"> • Acupuncture rationale: not stated

Zhang 2013a (Continued)

- Needling details
 - Points used: scalp acupoints
 - Numbers of points used: 4 acupoints
 - Depths of insertion: 0.66-1.05 inches
 - Deqi elicited: yes
 - Needle stimulation: manual and electrical
 - Needle retention time: 30 minutes
 - Needle type: hua tuo brand, length 0.98 inches
- Treatment regimen
 - Number of treatment sessions: 30 sessions
 - Frequency of treatment: 5 sessions/week
 - Total course: 6 weeks
- Practitioner background: not stated
- Co-intervention: WM

Control interventions: WM

Outcomes	<ul style="list-style-type: none"> • Improvement of depression (Hamilton Depression Scale) • Improvement of sleep (Rhone Planck Sleepiness Scale) • Adverse event • FU: 6 weeks
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomisation by using closed envelopes
Allocation concealment (selection bias)	Low risk	Randomisation by using closed envelopes
Blinding (performance bias and detection bias) All outcomes	High risk	Quote: "unable to blind to participants and operators"
Blinding of participants and personnel (performance bias) All outcomes	High risk	Quote: "unable to blind to participants and operators"
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Details not given
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Information on attrition was not reported
Selective reporting (reporting bias)	Unclear risk	Some patient-related outcomes were not reported, such as information on quality of life or all-cause mortality
Other bias	Unclear risk	No information provided

Zhang 2015

Methods	<p>RCT</p> <p>Method of randomisation: random number table</p> <p>Blinding: not stated</p> <p>Adverse effects: not stated</p> <p>ITT analysis: not stated</p> <p>Losses to FU: not stated</p>
Participants	<p>Country: China</p> <p>Number of participants included: 90 (30/30/30)</p> <p>Demographics: 66% male</p> <p>Type of stroke: ischaemic only</p> <p>Diagnosis: CT/MRI</p> <p>Severity on entry: not stated</p> <p>Time from stroke onset: 2-6 months</p> <p>Setting: inpatient</p> <p>Comparability: no significant difference in age or time post onset</p>
Interventions	<p>3 arms:</p> <ul style="list-style-type: none"> • acupuncture only • PT only • acupuncture + PT <p>Comparison eligible: acupuncture + PT versus PT only</p> <p>Acupuncture treatment</p> <ul style="list-style-type: none"> • Acupuncture rationale: not stated • Needling details <ul style="list-style-type: none"> ◦ Points used: scalp acupoints ◦ Numbers of points used: 3 points ◦ Depths of insertion: 0.98-1.18 inches ◦ Deqi elicited: not stated ◦ Needle stimulation: manual ◦ Needle retention time: 30 minutes ◦ Needle type: length 1.57 inches • Treatment regimen <ul style="list-style-type: none"> ◦ Number of treatment sessions: 24 sessions ◦ Frequency of treatment: 6 sessions/week ◦ Total course: 4 weeks • Practitioner background: not stated • Co-intervention: PT <p>Control interventions: PT</p>
Outcomes	<ul style="list-style-type: none"> • Improvement of neurological function (Modified Edinburgh And Scandinavia Stroke Scale) • Improvement of independence (Barthel Index) • FU: 4 weeks
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Zhang 2015 (Continued)

Random sequence generation (selection bias)	Low risk	Randomisation by using random number table
Allocation concealment (selection bias)	Unclear risk	Information on allocation concealment was not reported
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Information on attrition was not reported
Selective reporting (reporting bias)	Unclear risk	Free of selective reporting bias was assessed as 'unclear' due to some clinically important outcomes unstated, such as quality of life, mortality and adverse events
Other bias	Unclear risk	No information provided

Zheng 2014

Methods	<p>RCT Method of randomisation: randomisation by using random number table Blinding: not stated</p> <p>Adverse effects: not stated ITT analysis: not stated Losses to FU: none</p>
Participants	<p>Country: China Number of participants included: 60 (30/30) Demographics: aged 51-79 years, 60% male Type of stroke: both ischaemic and haemorrhagic strokes Diagnosis: WHO definition Severity on entry: not stated Time from stroke onset: 1-6 months Setting: inpatient Comparability: no significant difference in age or time post onset</p>
Interventions	<p>Comparison: acupuncture + WM versus WM</p> <p>Acupuncture treatment</p> <ul style="list-style-type: none"> Acupuncture rationale: not stated

Zheng 2014 (Continued)

- Needling details
 - Points used: both body and scalp acupoints
 - Numbers of points used: 2 scalp acupoints and 5 body acupoints
 - Depths of insertion: 0.39-1.97 inches
 - Deqi elicited: yes
 - Needle stimulation: manual
 - Needle retention time: 30 minutes
 - Needle type: hua tuo brand, length 1.57 inches
- Treatment regimen
 - Number of treatment sessions: 48 sessions
 - Frequency of treatment: 6 sessions/week
 - Total course: 8 weeks
- Practitioner background: not stated
- Co-intervention: WM

Control interventions: WM

Outcomes	<ul style="list-style-type: none"> • Improvement of cognitive function (MMSE, MoCA) • Improvement of independence (Barthel Index) • FU: 8 weeks
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomisation by using random number table
Allocation concealment (selection bias)	Unclear risk	Information on allocation concealment was not reported
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "No participants lost to follow-up"
Selective reporting (reporting bias)	Unclear risk	Free of selective reporting bias was assessed as 'unclear' due to some clinically important outcomes unstated, such as quality of life, mortality and adverse events
Other bias	Unclear risk	No information provided

Zhou 2014

Methods	<p>RCT</p> <p>Method of randomisation: closed envelopes</p> <p>Blinding: Investigators but not participants blinded</p> <p>Adverse effects: 20</p> <p>ITT analysis: not stated</p> <p>Losses to FU: none</p>
Participants	<p>Country: China</p> <p>Number of participants included: 147 (75/72)</p> <p>Demographics: aged 35-80 years, 54% male</p> <p>Type of stroke: ischaemic only</p> <p>Diagnosis: WHO definition and all confirmed by CT/MRI</p> <p>Severity on entry: not stated</p> <p>Time from stroke onset: 1-3 months</p> <p>Setting: inpatient</p> <p>Comparability: no significant difference in age or time post onset</p>
Interventions	<p>Comparison: acupuncture + daily PT versus daily PT</p> <p>Acupuncture treatment</p> <ul style="list-style-type: none"> • Acupuncture rationale: not stated • Needling details <ul style="list-style-type: none"> ◦ Points used: both body and scalp acupoints ◦ Numbers of points used: 8 scalp acupoints and 9 body acupoints ◦ Depths of insertion: 1.18-1.38 inches ◦ Deqi elicited: not stated ◦ Needle stimulation: manual and electrical ◦ Needle retention time: 6-8 hours/30 minutes ◦ Needle type: hua tuo brand, length 1.57 inches • Treatment regimen <ul style="list-style-type: none"> ◦ Number of treatment sessions: 40 sessions ◦ Frequency of treatment: 5 sessions/week ◦ Total course: 8 weeks • Practitioner background: not stated • Co-intervention: PT <p>Control interventions: PT</p>
Outcomes	<ul style="list-style-type: none"> • Improvement of depression (HAMD) • Improvement of neurological function (NIHSS) • Improvement of independence (Barthel Index) • Improvement of motor function (Fugl-Meyer scale) • Adverse event: bleeding, haematoma, and pain
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomisation by using closed envelopes

Zhou 2014 (Continued)

Allocation concealment (selection bias)	Low risk	Randomisation by using closed envelopes
Blinding (performance bias and detection bias) All outcomes	Low risk	Investigators blinded
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Details not given
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Outcome assessors blinded
Incomplete outcome data (attrition bias) All outcomes	Low risk	Treatment: no participants lost to follow-up Control: 3 participants lost to follow-up
Selective reporting (reporting bias)	Unclear risk	Some patient-related outcomes were not reported, such as information on quality of life or all-cause mortality
Other bias	Unclear risk	No information provided

Zhu 2007

Methods	RCT Method of randomisation: not stated Blinding: not stated ITT analysis: not stated Losses to FU: not stated
Participants	Country: China Number of participants included: 60 (30/30) Demographics: aged 45-69 years, 53% male Type of stroke: both ischaemic and haemorrhagic strokes Diagnosis: WHO definition Severity on entry: not stated Time from stroke onset: 1-6 months Setting: inpatient Comparability: no significant difference in age, comorbidity, or time post onset
Interventions	Comparison: acupuncture + PT versus PT Acupuncture treatment <ul style="list-style-type: none"> Acupuncture rationale: not stated Needling details <ul style="list-style-type: none"> Points used: both body and scalp acupoints Numbers of points used: 2 scalp acupoints and 4 body acupoints Depths of insertion: 1.57 inches Deqi elicited: yes Needle stimulation: manual and electrical stimulation Needle retention time: 30 minutes Needle type: length 1.57 inches

Zhu 2007 (Continued)

- Treatment regimen
 - Number of treatment sessions: 24 sessions
 - Frequency of treatment: 6 sessions/week
 - Total course: 4 weeks
- Practitioner background: not stated
- Co-intervention: PT

Control interventions: PT

Outcomes	<ul style="list-style-type: none"> • Improvement of neurological function (Modified Edinburgh Stroke Scale and Scandinavian Stroke Scale)
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	The method of random sequence generation was not reported
Allocation concealment (selection bias)	Unclear risk	Information on allocation concealment was not reported
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Information on blinding was not reported
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Information on blinding was not reported
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Information on attrition was not reported
Selective reporting (reporting bias)	Unclear risk	Free of selective reporting bias was assessed as 'unclear' due to some clinically important outcomes unstated, such as quality of life, mortality and adverse events
Other bias	Unclear risk	No information provided

BMIT: Boston Motor Inventory Test

CSRS 1: Chinese Stroke Recovery Scale based on the revised diagnostic criteria of acute cerebral infarction formulated by the second National Academic Symposium on Cerebrovascular Diseases of the Chinese Medical Association in 1986, which is similar to the Revised Scandinavian Stroke Scale (RSSS)

CSRS 2: Chinese Stroke Recovery Scale based on principles of traditional Chinese medicine

CT: computerised tomography

FU: follow-up

HAMD: Hamilton Depression Scale

ITT: intention-to-treat

LOTCA: Loewenstein Occupational Therapy Cognitive Assessment

MESS: modified Edinburgh Stroke Scale

Acupuncture for stroke rehabilitation (Review)

Copyright © 2016 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

MMSE: Mini Mental State Examination
MoCA: Montreal Cognitive Assessment Scale
MRI: magnetic resonance imaging
OT: occupational therapy
PT: physical therapy
qd: once per day
RCT: randomised controlled trial
SF-36: 36-Item Short Form Health Survey
SS-QOL: Stroke Specific Quality of Life Scale
SSS: Scandinavian Stroke Scale
TCM: Traditional Chinese Medicine
WHO: World Health Organization
WM: Western medicine

Characteristics of excluded studies *[ordered by study ID]*

Study	Reason for exclusion
Bai 2007	The trial included people with stroke less than 1 month since onset
Bai 2011	The trial included people with stroke less than 1 month since onset
Bai 2013	The trial included people with stroke less than 1 month since onset although the median time to post-stroke was more than 1 month
Bao 2008	The duration of stroke since onset was not stated in the trial
Bao 2010	The duration of stroke since onset was not clear
Bao 2012a	The trial included people with stroke less than 1 month since onset
Cai 2010	The trial included people with stroke less than 1 month since onset
Calabro 2011	Abstract only; clarification of randomisation and intervention sought but not obtained
Cao 2010a	The trial included people with stroke less than 1 month since onset
Cao 2010b	Not RCT
Chang 2010	The trial included people with stroke less than 1 month since onset
Chao 2009	The trial included people with stroke less than one month since onset
Chau 2009	It was not possible to include data from this trial in the analysis. The trial aimed to assess effects of 2 methods of acupuncture on subacute or chronic stroke. Acupoints were different between the 2 groups
Che 2002	The trial aimed to assess effects of 2 methods of acupuncture on subacute or chronic stroke. Acupoints were different between the 2 groups
Chen 2000	Confounded; the trial aimed to assess effects of acupuncture only compared with drug therapy (such as WM or TCM)
Chen 2007	The trial included people with stroke less than 1 month since onset
Chen 2008a	The duration of stroke since onset was not clear

Study	Reason for exclusion
Chen 2008b	Stroke population not randomised to treatment: quasi-randomisation of participants (based on administrative procedure) to 1 of 2 groups
Chen 2009	The trial included people with stroke less than 1 month since onset
Chen 2010a	The trial included people with stroke less than 1 month since onset
Chen 2010b	The duration of stroke since onset was not clear
Chen 2012a	The trial included people with stroke less than 1 month since onset
Chen 2012b	The trial included people with stroke less than 1 month since onset
Chen 2013	The trial included people with acute stroke (less than 1 month since onset)
Chen 2014	The trial included people with stroke less than 1 month since onset
Cheng 2006	The duration of stroke since onset was not stated in the trial
Cheng 2007	The duration of stroke since onset was not stated in the trial
Cheng 2011	It was not possible to include data from this trial in the analysis. Stroke population not randomised to treatment: quasi-randomisation of participants (based on administrative procedure) to 1 of 2 groups
Chow 2006	Abstract only; clarification of randomisation and intervention sought but not obtained
Chu 2007	The trial included people with stroke less than 1 month since onset
Chu 2009	The trial included people with stroke less than 1 month since onset
Cui 1992	It was not possible to include data from this trial in the analysis; this confounded trial aimed to assess effects of acupuncture only compared with drug therapy (such as WM or TCM)
Ding 2000	It was not possible to include data from this trial in the analysis. The trial aimed to assess effects of 2 methods of acupuncture on subacute or chronic stroke. Acupoints were different between the 2 groups
Dong 2007a	The trial included people with acute stroke (less than 1 month since onset)
Dong 2007b	The duration of stroke since onset was not clear
Dong 2008a	The trial included people with acute stroke (less than 1 month since onset)
Dong 2008b	The trial included people with acute stroke (less than 1 month since onset)
Dong 2009	The trial included people with stroke less than 1 month since onset
Du 2013	The duration of stroke since onset was not stated in the trial
Duan 2010	The trial included people with stroke less than 1 month since onset
Duan 2014	The duration of stroke since onset was not clear
Fang 2009	The duration of stroke since onset was not stated in the trial

Study	Reason for exclusion
Feng 1996a	It was not possible to include data from this trial in the analysis. The trial aimed to assess effects of 2 methods of acupuncture on subacute or chronic stroke. Acupoints were different between the 2 groups
Feng 1996b	It was not possible to include data from this trial in the analysis. The trial aimed to assess effects of 2 methods of acupuncture on subacute or chronic stroke. Acupoints were different between the 2 groups
Feng 2011	The trial included people with stroke less than 1 month since onset
Fink 2004	It was not possible to include data from this trial in the analysis. MAS scores (as a clinical measure of spasticity) were assessed before and after the treatment period but mean change of neurological score after the treatment period was not available
Fu 2010	The trial included people with stroke less than 1 month since onset
Fu 2013	The trial included people with stroke less than 1 month since onset
Fu 2014	The trial included people with stroke less than 1 month since onset
Gao 2001	It was not possible to include data from this trial in the analysis. This confounded trial aimed to assess effects of acupuncture only compared with drug therapy (such as WM or TCM)
Gao 2008	The duration of stroke since onset was not stated in the trial
Gao 2011	The duration of stroke since onset was not clear
Gao 2014b	The trial included people with stroke less than 1 month since onset
Gong 2008	The trial included people with stroke less than 1 month since onset
Gu 2007	The trial included people with acute stroke (less than 1 month since onset)
Gu 2009	The trial included people with stroke less than 1 month since onset
Guan 2009a	It was not possible to include data from this trial in the analysis. Stroke population not randomised to treatment: quasi-randomisation of participants (based on administrative procedure) to 1 of 2 groups
Guan 2009b	Quasi-randomised study
Guan 2013	The trial included people with stroke less than 1 month since onset
Guo 1999	The trial aimed to assess effects of acupuncture only compared with PT and OT
Guo 2006a	The trial included people with stroke less than 1 month since onset
Guo 2006b	The trial included people with stroke less than 1 month since onset
Guo 2013	The trial included people with stroke less than 1 month since onset
Han 2008	The trial included people with stroke less than 1 month since onset
Han 2010	The trial included people with stroke less than 1 month since onset

Study	Reason for exclusion
Han 2011	The trial included people with stroke less than 1 month since onset
Han 2012	The duration of stroke since onset was not stated in the trial
Han 2014	Quasi-randomised study
Hang 2014	The trial included people with stroke less than 1 month since onset
He 2008	The trial included people with stroke less than 1 month since onset
Hegyi 2012	The trial included people with stroke less than 1 month since onset
Hong 2013	The duration of stroke since onset was not clear
Hou 1998	The trial aimed to assess the effects of the combination of acupuncture, oxygen and herbs compared with acupuncture or oxygen respectively
Hsing 2012	The trial aimed to assess effects of electrical acupuncture compared with routine acupuncture
Hu 2011	The duration of stroke since onset was not stated in the trial
Hu 2012	The trial included people with stroke less than 1 month since onset
Huang 2002	It was not possible to include data from this trial in the analysis; this confounded trial aimed to assess effects of acupuncture only compared with drug therapy (such as WM or TCM)
Huang 2008b	The duration of stroke since onset was not clear
Huang 2008c	The duration of stroke since onset was not clear
Huang 2011	The trial included people with stroke less than 1 month since onset
Huang 2012	The trial included people with stroke less than 1 month since onset
Huang 2014	The duration of stroke since onset was not clear
Jia 2012	The trial included people with stroke less than 1 month since onset
Jiang 1998	It was not possible to include data from this trial in the analysis. The trial aimed to assess effects of 2 methods of acupuncture on subacute or chronic stroke. Acupoints were different between the 2 groups
Jiang 2000	It was not possible to include data from this trial in the analysis; this confounded trial aimed to assess effects of acupuncture only compared with drug therapy (such as WM or TCM)
Jiang 2006	The trial included people with stroke less than 1 month since onset
Jiang 2009	The trial included people with stroke less than 1 month since onset
Jiang 2010	The duration of stroke since onset was not stated in the trial
Jiang 2011a	The duration of stroke since onset was not clear
Jiang 2011b	The trial included people with stroke less than 1 month since onset

Study	Reason for exclusion
Jiang 2012	The trial included people with stroke less than 1 month since onset
Jiang 2013	The trial included people with stroke less than 1 month since onset
Jin 1993	It was not possible to include data from this trial in the analysis. The trial aimed to assess effects of 2 methods of acupuncture on subacute or chronic stroke. Acupoints were different between the 2 groups
Jin 2010a	Quasi-randomised study
Jin 2010b	The duration of stroke since onset was not stated in the trial
Kang 2011	The trial included people with stroke less than 1 month since onset
Kang 2013	The trial included people with stroke less than 1 month since onset
Kjendahl 1997	The trial included people with stroke less than 1 month since onset although the median time to post-stroke was more than 1 month
Lai 1997a	It was not possible to include data from this trial in the analysis; this confounded trial aimed to assess effects of acupuncture only compared with drug therapy (such as WM or TCM)
Lai 1997b	The trial aimed to assess the effects of electrical acupuncture compared with manual acupuncture. Acupoints were the same between the 2 groups
Lai 1998	The trial aimed to assess the effects of electrical acupuncture compared with manual acupuncture. Acupoints were the same between 2 groups
Lai 2004	It was not possible to include data from this trial in the analysis. The trial aimed to assess effects of 2 methods of acupuncture on subacute or chronic stroke. Acupoints were different between the 2 groups
Lai 2011	The duration of stroke since onset was not clear
Lang 2013	The trial included people with stroke less than 1 month since onset
Lao 2013	The trial included people with stroke less than 1 month since onset
Lee 2007	The full-text with English or Chinese language was not available
Lee 2009	The duration of stroke since onset was not stated in the trial
Lei 2013	The trial included people with stroke less than 1 month since onset
Li 1993	It was not possible to include data from this trial in the analysis. The trial aimed to assess effects of 2 methods of acupuncture on subacute or chronic stroke. Acupoints were different between the 2 groups
Li 1994a	It was not possible to include data from this trial in the analysis. The trial aimed to assess effects of 2 methods of acupuncture on subacute or chronic stroke. Acupoints were different between the 2 groups
Li 1994b	It was not possible to include data from this trial in the analysis. The trial aimed to assess effects of 2 methods of acupuncture on subacute or chronic stroke. Acupoints were different between the 2 groups

Study	Reason for exclusion
Li 1997b	No outcome measures of interest were available in the trial, which aimed to assess the effects of acupuncture on the change of intellectual disturbance in people with vascular dementia caused by cerebrovascular disease
Li 2006a	The trial included people with stroke less than 1 month since onset
Li 2006b	The trial included people with stroke less than 1 month since onset
Li 2006c	The trial included people with stroke less than 1 month since onset
Li 2006d	The trial included people with stroke less than 1 month since onset
Li 2006e	The trial included people with stroke less than 1 month since onset
Li 2008a	The trial included people with acute stroke (less than 1 month since onset)
Li 2008b	The duration of stroke since onset was not clear
Li 2008c	The trial included people with stroke less than 1 month since onset
Li 2008d	The trial included people with stroke less than 1 month since onset
Li 2009a	The trial included people with stroke less than 1 month since onset
Li 2009b	The trial included people with stroke less than 1 month since onset
Li 2010b	The trial included people with stroke less than 1 month since onset
Li 2010c	The trial aimed to assess the effects of the combination of acupuncture and PT compared with PT. The duration of stroke since onset was not stated in the trial
Li 2011b	The trial included people with stroke less than 1 month since onset
Li 2011c	The trial included people with stroke less than 1 month since onset
Li 2011d	The trial included people with acute stroke (less than 1 month since onset)
Li 2012a	The duration of stroke since onset was not stated in the trial
Li 2012b	The duration of stroke since onset was not stated in the trial
Li 2012c	The trial included people with stroke less than 1 month since onset
Li 2012d	Quasi-randomised study
Li 2012e	The duration of stroke since onset was not clear
Li 2012f	The trial included people with stroke less than 1 month since onset although the median time to post-stroke was more than 1 month
Li 2012g	The duration of stroke since onset was not stated in the trial
Li 2013b	The trial included people with stroke less than 1 month since onset. Stroke population not randomised to treatment: quasi-randomisation of participants (based on administrative procedure) to 1 of 2 groups

Study	Reason for exclusion
Li 2013c	The trial included people with stroke less than 1 month since onset
Li 2014	The trial included people with stroke less than 1 month since onset
Li 2015	The trial included people with stroke less than 1 month since onset
Liang 1993	It was not possible to include data from this trial in the analysis. The trial aimed to assess effects of 2 methods of acupuncture on subacute or chronic stroke. Acupoints were different between the 2 groups
Liang 2010	Quasi-randomised study
Liang 2012	The trial included people with stroke less than 1 month since onset
Liao 2006a	The duration of stroke since onset was not clear
Liao 2006b	The duration of stroke since onset was not stated in the trial
Liao 2013	The trial included people with stroke less than 1 month since onset
Liao 2014	The duration of stroke since onset was not clear
Lim 2014	Review
Lin 2010a	The trial included people with stroke less than 1 month since onset
Lin 2010b	The duration of stroke since onset was not stated in the trial
Lin 2012	The duration of stroke since onset was not clear
Lin 2013	The trial included people with stroke less than 1 month since onset
Liu 1998	It was not possible to include data from this trial in the analysis. ADL was assessed before and after the treatment period but the number of participants who were independent after the treatment period was not available. The trial primarily aimed to assess the effects of acupuncture on the intellectual disturbance in people with vascular dementia due to stroke
Liu 2004a	The trial included people with stroke less than 1 month since onset (that is, acute stroke)
Liu 2004b	Confounded (acupuncture plus 1 type of Chinese herbs versus another type of Chinese herbs)
Liu 2006	The trial included people with acute stroke (less than 1 month since onset)
Liu 2008a	The trial included people with stroke less than 1 month since onset
Liu 2008b	Non-randomised controlled study
Liu 2009	The trial included people with stroke less than 1 month since onset
Liu 2010a	The trial included people with stroke less than 1 month since onset
Liu 2010b	The trial included people with stroke less than 1 month since onset
Liu 2011a	The duration of stroke since onset was not stated in the trial

Study	Reason for exclusion
Liu 2011b	The trial included people with stroke less than 1 month since onset
Liu 2012a	The trial included people with stroke less than 1 month since onset
Liu 2012b	The duration of stroke since onset was not clear
Liu 2012c	The trial included people with stroke less than 1 month since onset
Liu 2013c	The trial included people with stroke less than 1 month since onset
Liu 2013d	The duration of stroke since onset was not clear
Liu 2013e	The duration of stroke since onset was not stated in the trial
Liu 2013f	The duration of stroke since onset was not stated in the trial
Liu 2014b	The trial included people with stroke less than 1 month since onset
Liu 2015	The duration of stroke since onset was not clear
Long 2013	The duration of stroke since onset was not stated in the trial
Lu 2010	The trial included people with stroke less than 1 month since onset
Lu 2011a	The trial included people with stroke less than 1 month since onset
Lu 2011b	The duration of stroke since onset was not stated in the trial
Luo 2007	The trial included people with stroke less than 1 month since onset
Luo 2008	The trial included people with stroke less than 1 month since onset
Luo 2012	The duration of stroke since onset was not stated in the trial
Lv 2009	The trial included people with stroke less than 1 month since onset
Lv 2014	The trial included people with stroke less than 1 month since onset
Ma 2011a	The trial included people with acute stroke (less than 1 month since onset)
Ma 2011b	The trial included people with stroke less than 1 month since onset
Ma 2013	The trial included people with stroke less than 1 month since onset
Mou 2010	The trial included people with stroke less than 1 month since onset
Mu 2007	The duration of stroke since onset was not stated in the trial
Mukherjee 2006	Abstract only; clarification of randomisation and intervention sought but not obtained
Muo 2001	It was not possible to include data from this trial in the analysis. This confounded trial aimed to assess effects of acupuncture only compared with drug therapy (such as WM or TCM)
Ni 2011	The trial included people with stroke less than 1 month since onset

Study	Reason for exclusion
Nie 2011	The trial included people with stroke less than 1 month since onset
Nie 2013	The duration of stroke since onset was not stated in the trial
Pan 2013	The duration of stroke since onset was not stated in the trial
Pang 2010	Not RCT
Peng 2007a	The trial aimed to assess the effects of the combination of acupuncture, PT and WM compared with WM. The duration of stroke since onset was not stated in the trial
Peng 2009	The trial included people with stroke less than 1 month since onset
Peng 2015	The duration of stroke since onset was not stated in the trial
Qi 2000	It was not possible to include data from this trial in the analysis. The trial aimed to assess effects of 2 methods of acupuncture on subacute or chronic stroke. Acupoints were different between the 2 groups
Qi 2013	The trial included people with stroke less than 1 month since onset
Qi 2013a	The trial included people with stroke less than 1 month since onset
Qiao 2012	Abstract only; clarification of randomisation and intervention sought but not obtained
Qiu 2011	Quasi-randomised study
Qu 1991	It was not possible to include data from this trial in the analysis. The trial aimed to assess effects of 2 methods of acupuncture on subacute or chronic stroke. Acupoints were different between the 2 groups
Ran 2013	The trial included people with stroke less than 1 month since onset
Ren 2009	The duration of stroke since onset was not stated in the trial
Ren 2010	The trial included people with acute stroke (less than 1 month since onset)
Ren 2012	The trial included people with stroke less than 1 month since onset
Sallstrom 1996	The median time to post-stroke was more than 1 month, but the trial included people with acute stroke (that is, less than 1 month)
Salom-Moreno 2014	The duration of stroke since onset was not stated in the trial
Schaechter 2007	The duration of stroke since onset was not stated in the trial
Seo 2013	The trial aimed to assess the effects of ouhyul herbal acupuncture point injection compared with normal saline acupuncture point injection
Shang 2008	Quasi-randomised study
Shao 2012	The duration of stroke since onset was not clear
Shen 2008	The trial included people with stroke less than 1 month since onset

Study	Reason for exclusion
Shen 2009	The trial included people with stroke less than 1 month since onset
Sheng 2011	The trial included people with stroke less than 1 month since onset
Sheng 2013	The duration of stroke since onset was not clear
Shi 2007	The trial aimed to assess the effects of combination of acupuncture, PT and WM compared with WM
Shi 2007a	The trial included people with stroke less than 1 month since onset
Song 2010	Not stroke
Sui 2001	It was not possible to include data from this trial in the analysis. The trial aimed to assess effects of 2 methods of acupuncture on subacute or chronic stroke. Acupoints were different between the 2 groups
Sun 2008	Abstract only; clarification of randomisation and intervention sought but not obtained
Sun 2010	The duration of stroke since onset was not clear
Sun 2011	Not RCT
Sun 2013b	The duration of stroke since onset was not stated in the trial
Sun 2014	The trial included people with stroke less than 1 month since onset
Suo 2008	The trial aimed to assess the effects of combination of sham acupuncture and PT compared with acupuncture only
Tang 2005	The trial included people with stroke less than 1 month since onset
Tang 2012	The trial included people with stroke less than 1 month since onset
Tang 2013	The trial included people with stroke less than 1 month since onset
Tang 2013a	The trial aimed to assess the effects of combination of acupuncture and WM compared with combination of WM and PT
Tian 2014	The trial included people with stroke less than 1 month since onset
Tong 1997	It was not possible to include data from this trial in the analysis. The trial aimed to assess effects of 2 methods of acupuncture on subacute or chronic stroke. Acupoints were different between the 2 groups
Tong 2013	The trial included people with stroke less than 1 month since onset
Tong 2014	The trial included people with stroke less than 1 month since onset
Wan 2013	The duration of stroke since onset was not clear in the trial
Wang 2004	Randomisation was not stated; the trial included people with acute stroke (less than 1 month since onset)
Wang 2006	The trial included people with stroke less than 1 month since onset

Study	Reason for exclusion
Wang 2007	The trial included people with stroke less than 1 month since onset
Wang 2008	The trial included people with stroke less than 1 month since onset
Wang 2008a	The trial included people with stroke less than 1 month since onset
Wang 2008b	The trial included people with acute stroke (less than 1 month since onset)
Wang 2009	Quasi-randomised study
Wang 2010a	The duration of stroke since onset was not clear
Wang 2010b	The duration of stroke since onset was not clear
Wang 2011b	The trial included people with stroke less than 1 month since onset
Wang 2011c	The trial included people with stroke less than 1 month since onset
Wang 2011d	The duration of stroke since onset was not clear
Wang 2011e	The trial included people with stroke less than 1 month since onset
Wang 2011f	The duration of stroke since onset was not clear
Wang 2011g	The trial included people with stroke less than 1 month since onset
Wang 2011h	The trial included people with stroke less than 1 month since onset
Wang 2011i	The trial included people with stroke less than 1 month since onset
Wang 2011j	The trial included people with stroke less than 1 month since onset
Wang 2013	The duration of stroke since onset was not stated in the trial
Wang 2014a	The trial aimed to assess the effects of acupuncture compared with transcutaneous electrical acupoint stimulation
Wang 2014b	The duration of stroke since onset was not stated in the trial
Wei 2005	The trial included people with ischaemic stroke less than 1 month since onset (that is, acute stroke). Data for participants with haemorrhagic stroke more than 150 days were not separated from data for participants with ischaemic stroke
Wei 2009	The trial included people with stroke less than 1 month since onset
Wenli 2007	The trial included people with stroke less than 1 month since onset
Wong 2013	Abstract only; clarification of randomisation and intervention sought but not obtained
Wu 1999	Data on pre-planned outcome measures of interest were not available from the trial, which aimed to assess the effects of acupuncture on urinary incontinence in people with stroke
Wu 2009a	The trial included people with stroke less than 1 month since onset
Wu 2009b	The trial aimed to assess the effects of three different forms of acupuncture

Study	Reason for exclusion
Wu 2010	The trial included people with acute stroke (less than 1 month since onset)
Wu 2011	The trial included people with stroke less than 1 month since onset
Wu 2012	The duration of stroke since onset was not clear
Wu 2012a	The trial included people with acute stroke (less than 1 month since onset)
Wu 2013b	The trial included people with acute stroke (less than 1 month since onset)
Wu 2014a	The trial included people with acute stroke (less than 1 month since onset)
Wu 2014b	The duration of stroke since onset was not clear
Xia 2008	The trial included people with stroke less than 1 month since onset
Xia 2010	The trial included people with stroke less than 1 month since onset
Xiao 1996	It was not possible to include data from this trial in the analysis. The trial aimed to assess effects of 2 methods of acupuncture on subacute or chronic stroke. Acupoints were different between the 2 groups
Xiao 2006	The trial included people with stroke less than 1 month since onset
Xiao 2008	The trial included people with acute stroke (less than 1 month since onset)
Xiao 2011	The trial included people with acute stroke (less than 1 month since onset)
Xie 2010	The trial included people with stroke less than 1 month since onset
Xie 2012	The trial included people with stroke less than 1 month since onset
Xie 2013	Quasi-randomised study
Xing 2007	The trial included people with stroke less than 1 month since onset
Xu 2009a	The duration of stroke since onset was not stated in the trial
Xu 2009b	The duration of stroke since onset was not stated in the trial
Xu 2010	The trial included people with acute stroke (less than 1 month since onset)
Xu 2013a	The trial included people with stroke less than 1 month since onset
Xu 2014	The duration of stroke since onset was not clear
Yan 2010	The duration of stroke since onset was not clear
Yan 2011	The trial included people with stroke less than 1 month since onset
Yang 2006	The duration of stroke since onset was not clear
Yang 2007a	Quasi-randomised study
Yang 2007b	The trial included people with stroke less than 1 month since onset

Study	Reason for exclusion
Yang 2007c	The trial included people with stroke less than 1 month since onset
Yang 2008	The trial aimed to assess the effects of electrical acupuncture compared with routine acupuncture
Yang 2009	The duration of stroke since onset was not clear
Yang 2010	The trial included people with stroke less than 1 month since onset
Yang 2011a	The trial included people with stroke less than 1 month since onset
Yang 2011b	The duration of stroke since onset was not clear
Yang 2011c	The duration of stroke since onset was not clear
Yang 2012	The trial included people with stroke less than 1 month since onset
Yang 2014	The duration of stroke since onset was not clear
Yang 2014a	The duration of stroke since onset was not clear
Yang 2015	The trial included people with stroke less than 1 month since onset
Yao 2006	The duration of stroke since onset was not stated in the trial
Yao 2013a	The trial included people with stroke less than 1 month since onset
Yao 2013b	The trial included people with acute stroke (less than 1 month since onset)
Yin 2013a	The trial included people with acute stroke (less than 1 month since onset)
Yin 2013b	The trial included people with stroke less than 1 month since onset
Yin 2014	The duration of stroke since onset was not clear
You 2011	The trial included people with stroke less than 1 month since onset
You 2014	The trial included people with stroke less than 1 month since onset
Yu 2002	It was not possible to include data from this trial in the analysis. The trial aimed to assess effects of 2 methods of acupuncture on subacute or chronic stroke. Acupoints were different between the 2 groups. The trial was also confounded; it aimed to assess effects of acupuncture only compared with drug therapy (such as WM or TCM)
Yu 2007	The trial included people with stroke less than 1 month since onset
Yu 2009	The trial included people with stroke less than 1 month since onset
Yu 2011	Abstract only; clarification of randomisation and intervention sought but not obtained
Yuan 2010	The trial included people with stroke less than 1 month since onset
Zeng 2010	The trial included people with stroke less than 1 month since onset
Zhang 1988	It was not possible to include data from this trial in the analysis. This confounded trial aimed to assess effects of acupuncture only compared with drug therapy (such as WM or TCM)

Study	Reason for exclusion
Zhang 1997	It was not possible to include data from this trial in the analysis. This confounded trial aimed to assess effects of acupuncture only compared with drug therapy (such as WM or TCM)
Zhang 2002a	The trial aimed to assess the effects of magnetic acupuncture compared with routine acupuncture. Acupoints were similar between the 2 groups
Zhang 2002b	The duration of stroke since onset was not stated in the trial; it was not possible to include data from this trial in the analysis. This confounded trial aimed to assess effects of acupuncture only compared with drug therapy (such as WM or TCM)
Zhang 2007a	The trial included people with stroke less than 1 month since onset
Zhang 2007b	The trial aimed to assess the effects of the combination of acupuncture, buyang huanwu decoction and routine treatment compared with routine treatment
Zhang 2008	The trial included people with stroke less than 1 month since onset
Zhang 2009b	The trial included people with stroke less than 1 month since onset
Zhang 2009c	The duration of stroke since onset was not stated in the trial
Zhang 2010a	The trial aimed to assess the effects of basic therapy, rehabilitation, acupuncture, TCM, and massage therapy compared with basic treatment and rehabilitation treatment
Zhang 2010b	The trial included people with acute stroke (less than 1 month since onset)
Zhang 2010c	The trial included people with acute stroke (less than 1 month since onset)
Zhang 2010d	The trial included people with stroke less than 1 month since onset
Zhang 2010e	The duration of stroke since onset was not clear
Zhang 2010f	The trial included people with acute stroke (less than 1 month since onset)
Zhang 2011a	The trial included people with stroke less than 1 month since onset
Zhang 2011b	The duration of stroke since onset was not stated in the trial
Zhang 2011c	The duration of stroke since onset was not stated in the trial
Zhang 2011d	The trial included people with stroke less than 1 month since onset
Zhang 2011e	The duration of stroke since onset was not clear
Zhang 2012b	The trial included people with stroke less than 1 month since onset
Zhang 2012c	The trial included people with stroke less than 1 month since onset
Zhang 2012d	The trial included people with stroke less than 1 month since onset
Zhang 2012e	The trial included people with stroke less than 1 month since onset
Zhang 2013b	The trial included people with stroke less than 1 month since onset
Zhang 2014b	Abstract only; clarification of randomisation and intervention sought but not obtained

Study	Reason for exclusion
Zhang 2014c	The duration of stroke since onset was not clear
Zhang 2014d	The trial included people with stroke less than 1 month since onset
Zhao 2008	The duration of stroke since onset was not stated in the trial
Zhao 2009	The trial aimed to assess the effects of deep acupuncture compared with routine acupuncture. Acupoints were the same between 2 groups
Zhao 2010	The duration of stroke since onset was not stated in the trial
Zhao 2011a	The duration of stroke since onset was not clear
Zhao 2011b	The trial included people with acute stroke (less than 1 month since onset)
Zhao 2014a	The trial included people with stroke less than 1 month since onset
Zhao 2014b	The duration of stroke since onset was not stated in the trial
Zheng 1996	The trial aimed to assess the effects of combination of body and scalp acupuncture compared with body acupuncture only or scalp acupuncture only
Zheng 2000	Confounded (acupuncture plus Chinese herbs versus WM only)
Zheng 2011b	The trial aimed to assess the effects of different forms of acupuncture
Zheng 2011c	The trial included people with stroke less than 1 month since onset
Zhong 2014	The trial included people with stroke less than 1 month since onset
Zhou 1995	It was not possible to include data from this trial in the analysis. The trial aimed to assess effects of 2 methods of acupuncture on subacute or chronic stroke. Acupoints were different between the 2 groups
Zhou 2008	The trial aimed to assess the effects of combination of acupuncture, PT and WM compared with WM. The duration of stroke since onset was not stated in the trial
Zhu 2010a	The duration of stroke since onset was not stated in the trial
Zhu 2010b	The duration of stroke since onset was not stated in the trial
Zhu 2011	Abstract only; clarification of randomisation and intervention sought but not obtained
Zhu 2012a	The trial included people with stroke less than 1 month since onset
Zhu 2012b	The duration of stroke since onset was not stated in the trial
Zhuang 2012	The trial included people with stroke less than 1 month since onset

ADL: activities of daily living

MAS: Modified Ashworth Scale

OT: occupational therapy

PT: physical therapy

TCM: Traditional Chinese Medicine

WM: Western medicine

Characteristics of ongoing studies [ordered by study ID]

Fu 2011

Trial name or title	Randomised, controlled Phase III trial to evaluate the efficacy and safety of comprehensive acupuncture treatment programmes for post-stroke motor rehabilitation
Methods	Randomised parallel controlled trial
Participants	People with haemorrhagic or ischaemic stroke
Interventions	Acupuncture + rehabilitation training
Outcomes	<ul style="list-style-type: none"> • BOSS • Fugl-Meyer Motor Assessment
Starting date	1 January 2011
Contact information	Shanghai Municipal Health Bureau
Notes	Recruiting status : completed

Liu 2013b

Trial name or title	Acupuncture based on five elements body characteristics in the treatment of people with post-stroke depression
Methods	Randomised parallel controlled trial
Participants	People with post-stroke depression
Interventions	<ul style="list-style-type: none"> • Five pedestrian body dialectical acupuncture • Conventional acupuncture
Outcomes	<ul style="list-style-type: none"> • HAMD • ADL • Neurologic function score
Starting date	1 July 2013
Contact information	The First Affiliated Hospital of Guangxi Traditional Chinese Medicine University
Notes	Recruiting status

Xie 2006

Trial name or title	Randomised controlled study on the acupuncture for dysphagia in convalescence phase of apoplexy
Methods	Parallel RCT
Participants	People with dysphagia in the convalescent phase of apoplexy

Xie 2006 (Continued)

Interventions	Acupuncture with swallowing training
Outcomes	<ul style="list-style-type: none"> • Mortality • Improvement • Neurological function scale • Length of hospitalisation • Nutrition status • Quality of life • TCM outcomes • Rate of Lung Infection • Loss of follow-up, and withdraw • Adverse events
Starting date	1 October 2006
Contact information	Department of Acupuncture, Huguosi TCM Hospital attached to the Beijing University of TCM
Notes	Date last refreshed: 7 February 2015

Zhong 2010

Trial name or title	The research of linguistic functional recovery mechanism based on fMRI after electroacupuncture at acupoints Tongli (HT5) and Xuanzhong (GB39) curing basal ganglia aphasia after stroke
Methods	Randomised parallel controlled trial
Participants	People who have explicit syndrome of aphasia, presenting within ischaemic stroke only in basal ganglia
Interventions	Electroacupuncture
Outcomes	<ul style="list-style-type: none"> • Linguistic functional evaluation after 30 days' and 60 days' therapy • fMRI after 30 days' and 60 days' therapy
Starting date	1 January 2010
Contact information	State Administration of Traditional Chinese Medicine
Notes	Recruiting status : completed

ADL: activities of daily living

BOSS: Burden of Stroke Scale

fMRI: functional magnetic resonance imaging

HAMD: Hamilton Depression Scale

TCM: traditional Chinese medicine

DATA AND ANALYSES

Comparison 1. Acupuncture plus baseline treatment versus sham acupuncture plus baseline treatment

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Improvement of motor function	1		Odds Ratio (M-H, Fixed, 95% CI)	Totals not selected
2 Improvement of cognitive function	1		Mean Difference (IV, Fixed, 95% CI)	Totals not selected
2.1 Orientation	1		Mean Difference (IV, Fixed, 95% CI)	0.0 [0.0, 0.0]
2.2 Perception	1		Mean Difference (IV, Fixed, 95% CI)	0.0 [0.0, 0.0]
2.3 Praxis	1		Mean Difference (IV, Fixed, 95% CI)	0.0 [0.0, 0.0]
2.4 Visuomotor organisation	1		Mean Difference (IV, Fixed, 95% CI)	0.0 [0.0, 0.0]
2.5 Thinking operation	1		Mean Difference (IV, Fixed, 95% CI)	0.0 [0.0, 0.0]
2.6 Memory	1		Mean Difference (IV, Fixed, 95% CI)	0.0 [0.0, 0.0]
2.7 Attention	1		Mean Difference (IV, Fixed, 95% CI)	0.0 [0.0, 0.0]
3 Improvement of Health-Related Quality of Life (MOS SF-36)	1		Mean Difference (IV, Fixed, 95% CI)	Totals not selected
3.1 Physical component summary	1		Mean Difference (IV, Fixed, 95% CI)	0.0 [0.0, 0.0]
3.2 Mental component summary	1		Mean Difference (IV, Fixed, 95% CI)	0.0 [0.0, 0.0]
4 Improvement of Stroke-Specific Quality of Life(SS QOL)	1		Mean Difference (IV, Fixed, 95% CI)	Totals not selected
4.1 Language	1		Mean Difference (IV, Fixed, 95% CI)	0.0 [0.0, 0.0]
4.2 Personality	1		Mean Difference (IV, Fixed, 95% CI)	0.0 [0.0, 0.0]

Analysis 1.1. Comparison 1 Acupuncture plus baseline treatment versus sham acupuncture plus baseline treatment, Outcome 1 Improvement of motor function.

Study or subgroup	Acupuncture n/N	Control n/N	Odds Ratio M-H, Fixed, 95% CI	Odds Ratio M-H, Fixed, 95% CI
Naeser 1992	4/10	0/6		9[0.4,203.3]

Analysis 1.2. Comparison 1 Acupuncture plus baseline treatment versus sham acupuncture plus baseline treatment, Outcome 2 Improvement of cognitive function.

Analysis 1.3. Comparison 1 Acupuncture plus baseline treatment versus sham acupuncture plus baseline treatment, Outcome 3 Improvement of Health-Related Quality of Life (MOS SF-36).

Analysis 1.4. Comparison 1 Acupuncture plus baseline treatment versus sham acupuncture plus baseline treatment, Outcome 4 Improvement of Stroke-Specific Quality of Life(SS QOL).

Comparison 2. Acupuncture plus baseline treatment versus baseline treatment alone

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Improvement of dependency at the end of treatment	11		Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Barthel Index	9	616	Mean Difference (IV, Random, 95% CI)	9.19 [4.34, 14.05]
1.2 Modified Barthel Index	1	60	Mean Difference (IV, Random, 95% CI)	3.44 [0.30, 6.58]
1.3 ADL	1	62	Mean Difference (IV, Random, 95% CI)	7.80 [6.04, 9.56]
2 Improvement of dependency at the end of follow-up	2		Mean Difference (IV, Random, 95% CI)	Subtotals only
2.1 Barthel index	1	147	Mean Difference (IV, Random, 95% CI)	7.49 [1.79, 13.19]
2.2 ADL	1	71	Mean Difference (IV, Random, 95% CI)	3.83 [2.67, 4.99]
3 Improvement of global neurological deficit at the end of treatment	6		Mean Difference (IV, Random, 95% CI)	Subtotals only
3.1 Modified Edinburgh and Scandinavian Stroke Scale	4	240	Mean Difference (IV, Random, 95% CI)	-2.39 [-3.34, -1.43]
3.2 Neurological function deficit scale	2	123	Mean Difference (IV, Random, 95% CI)	-1.02 [-5.80, 3.76]
4 Improvement of global neurological deficit at the end of follow-up	2		Mean Difference (IV, Random, 95% CI)	Subtotals only
4.1 Neurological function deficit scale	1	71	Mean Difference (IV, Random, 95% CI)	-6.15 [-7.09, -5.21]
4.2 NIHSS	1	147	Mean Difference (IV, Random, 95% CI)	-0.83 [-1.94, 0.28]
5 Improvement of global neurological deficit at the end of treatment	7	543	Odds Ratio (M-H, Random, 95% CI)	3.89 [1.78, 8.49]
6 Improvement of motor function at the end of treatment	5		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
6.1 Upper and lower extremities motor function(Fugl-Meyer Assessment)	4	245	Mean Difference (IV, Fixed, 95% CI)	6.16 [4.20, 8.11]
6.2 General motor function (Motor assessment scale)	1	60	Mean Difference (IV, Fixed, 95% CI)	4.53 [2.99, 6.07]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
7 Improvement of general motor function at the end of follow-up (Fugl-Meyer Assessment)	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
8 Improvement of motor function at the end of treatment (Fugl-Meyer Assessment)	2	125	Odds Ratio (M-H, Random, 95% CI)	2.41 [0.98, 5.96]
9 Improvement of cognitive function at the end of treatment	5		Mean Difference (IV, Random, 95% CI)	Subtotals only
9.1 MMSE	5	278	Mean Difference (IV, Random, 95% CI)	2.54 [0.03, 5.05]
9.2 MoCA	2	120	Mean Difference (IV, Random, 95% CI)	1.34 [0.76, 1.92]
9.3 HDS-R	1	46	Mean Difference (IV, Random, 95% CI)	1.26 [0.29, 2.23]
10 Improvement of cognitive function at the end of follow-up (MMSE)	1	71	Mean Difference (IV, Fixed, 95% CI)	3.47 [2.43, 4.51]
11 Improvement of cognitive function at the end of treatment	3		Odds Ratio (M-H, Random, 95% CI)	Subtotals only
11.1 MMSE	3	166	Odds Ratio (M-H, Random, 95% CI)	3.82 [1.89, 7.72]
11.2 HDS-R	1	46	Odds Ratio (M-H, Random, 95% CI)	4.02 [1.12, 14.46]
12 Improvement of depression at the end of treatment	6		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
12.1 HAMD	6	552	Mean Difference (IV, Fixed, 95% CI)	-2.58 [-3.28, -1.87]
12.2 Symptoms of TCM depression scale	1	60	Mean Difference (IV, Fixed, 95% CI)	-1.57 [-2.96, -0.18]
13 Improvement of depression at the end of treatment (HAMD)	4	342	Odds Ratio (M-H, Random, 95% CI)	2.03 [1.10, 3.72]
14 Improvement of swallowing function at the end of treatment (Water drinking test)	2	200	Mean Difference (IV, Random, 95% CI)	-1.11 [-2.08, -0.14]
15 Improvement of swallowing function at the end of treatment (Water drinking test)	1	60	Odds Ratio (M-H, Random, 95% CI)	95.29 [10.93, 830.86]
16 Improvement of pain at the end of treatment (Visual Analogue Scale)	2	118	Mean Difference (IV, Fixed, 95% CI)	-2.88 [-3.68, -2.09]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
17 Improvement of sleep quality at the end of treatment (Rhone Planck Sleepiness Scale)	1	60	Mean Difference (IV, Fixed, 95% CI)	-1.09 [-2.37, 0.19]
18 Improvement of spasticity at the end of treatment (Modified Ashworth Spasticity Rating Scale)	1	60	Mean Difference (IV, Fixed, 95% CI)	-0.4 [-0.64, -0.16]
19 Improvement of quality of life at the end of treatment (MOS SF-36)	1	71	Mean Difference (IV, Fixed, 95% CI)	2.73 [-0.54, 6.00]

Analysis 2.1. Comparison 2 Acupuncture plus baseline treatment versus baseline treatment alone, Outcome 1 Improvement of dependency at the end of treatment.

Analysis 2.2. Comparison 2 Acupuncture plus baseline treatment versus baseline treatment alone, Outcome 2 Improvement of dependency at the end of follow-up.

Analysis 2.3. Comparison 2 Acupuncture plus baseline treatment versus baseline treatment alone, Outcome 3 Improvement of global neurological deficit at the end of treatment.

Analysis 2.4. Comparison 2 Acupuncture plus baseline treatment versus baseline treatment alone, Outcome 4 Improvement of global neurological deficit at the end of follow-up.

Analysis 2.5. Comparison 2 Acupuncture plus baseline treatment versus baseline treatment alone, Outcome 5 Improvement of global neurological deficit at the end of treatment.

Analysis 2.6. Comparison 2 Acupuncture plus baseline treatment versus baseline treatment alone, Outcome 6 Improvement of motor function at the end of treatment.

Analysis 2.7. Comparison 2 Acupuncture plus baseline treatment versus baseline treatment alone, Outcome 7 Improvment of general motor function at the end of follow-up (Fugl-Meyer Assessment).

Analysis 2.8. Comparison 2 Acupuncture plus baseline treatment versus baseline treatment alone, Outcome 8 Improvement of motor function at the end of treatment (Fugl-Meyer Assessment).

Analysis 2.9. Comparison 2 Acupuncture plus baseline treatment versus baseline treatment alone, Outcome 9 Improvement of cognitive function at the end of treatment.

Analysis 2.10. Comparison 2 Acupuncture plus baseline treatment versus baseline treatment alone, Outcome 10 Improvement of cognitive function at the end of follow-up (MMSE).

Analysis 2.11. Comparison 2 Acupuncture plus baseline treatment versus baseline treatment alone, Outcome 11 Improvement of cognitive function at the end of treatment.

Analysis 2.12. Comparison 2 Acupuncture plus baseline treatment versus baseline treatment alone, Outcome 12 Improvement of depression at the end of treatment.

Analysis 2.13. Comparison 2 Acupuncture plus baseline treatment versus baseline treatment alone, Outcome 13 Improvement of depression at the end of treatment (HAMD).

Analysis 2.14. Comparison 2 Acupuncture plus baseline treatment versus baseline treatment alone, Outcome 14 Improvement of swallowing function at the end of treatment (Water drinking test).

Analysis 2.15. Comparison 2 Acupuncture plus baseline treatment versus baseline treatment alone, Outcome 15 Improvement of swallowing function at the end of treatment (Water drinking test).

Analysis 2.16. Comparison 2 Acupuncture plus baseline treatment versus baseline treatment alone, Outcome 16 Improvement of pain at the end of treatment (Visual Analogue Scale).

Analysis 2.17. Comparison 2 Acupuncture plus baseline treatment versus baseline treatment alone, Outcome 17 Improvement of sleep quality at the end of treatment (Rhone Planck Sleepiness Scale).

Analysis 2.18. Comparison 2 Acupuncture plus baseline treatment versus baseline treatment alone, Outcome 18 Improvement of spasticity at the end of treatment (Modified Ashworth Spasticity Rating Scale).

Analysis 2.19. Comparison 2 Acupuncture plus baseline treatment versus baseline treatment alone, Outcome 19 Improvement of quality of life at the end of treatment (MOS SF-36).

APPENDICES

Appendix 1. Cochrane Stroke Group Trials Register search strategy

Stage: Late treatment (> 30 days)

Disease: Not specified

Condition: Not specified

Intervention type: Complementary medical therapy

Intervention code: Acupuncture

Appendix 2. CENTRAL search strategy

The Cochrane Library, June 2015

#1 [mh "cerebrovascular disorders"] or [mh "basal ganglia cerebrovascular disease"] or [mh "brain ischemia"] or [mh "carotid artery diseases"] or [mh "cerebral small vessel diseases"] or [mh "intracranial arterial diseases"] or [mh "intracranial embolism and thrombosis"]

or [mh "intracranial hemorrhages"] or [mh ^stroke] or [mh "brain infarction"] or [mh ^"stroke, lacunar"] or [mh ^"vasospasm, intracranial"] or [mh ^"vertebral artery dissection"]

#2 (stroke or poststroke or "post-stroke" or cerebrovasc* or brain next vasc* or cerebral next vasc* or cva* or apoplex* or SAH):ti,ab

#3 ((brain* or cerebr* or cerebell* or intracran* or intracerebral) near/5 (isch*emi* or infarct* or thrombo* or emboli* or occlus*)):ti,ab

#4 ((brain* or cerebr* or cerebell* or intracerebral or intracranial or subarachnoid) near/5 (haemorrhage* or hemorrhage* or haematoma* or hematoma* or bleed*)):ti,ab

#5 [mh ^hemiplegia] or [mh ^paresis]

#6 (hemipleg* or hemipar* or paresis or paretic):ti,ab

#7 [mh ^"brain injuries"] or [mh ^"brain injury, chronic"]

#8 #1 or #2 or #3 or #4 or #5 or #6 or #7

#9 [mh ^acupuncture] or [mh ^"acupuncture therapy"] or [mh ^"acupuncture analgesia"] or [mh ^"acupuncture, ear"] or [mh ^electroacupuncture] or [mh ^meridians] or [mh ^"acupuncture points"] or [mh ^"trigger points"]

#10 (acupuncture* or electroacupuncture or "electro-acupuncture" or acupoint* or meridians or needling):ti,ab

#11 ((meridian or non-meridian or trigger) near/10 point*):ti,ab

#12 #9 or #10 or #11

#13 #8 and #12

Appendix 3. MEDLINE search strategy

MEDLINE (Ovid) November 2005 to July 2015

1. cerebrovascular disorders/ or exp basal ganglia cerebrovascular disease/ or exp brain ischemia/ or exp carotid artery diseases/ or exp cerebral small vessel diseases/ or exp intracranial arterial diseases/ or exp "intracranial embolism and thrombosis"/ or exp intracranial hemorrhages/ or stroke/ or exp brain infarction/ or stroke, lacunar/ or vasospasm, intracranial/ or vertebral artery dissection/

2. (stroke or poststroke or post-stroke or cerebrovasc\$ or brain vasc\$ or cerebral vasc\$ or cva\$ or apoplex\$ or SAH).tw.

3. ((brain\$ or cerebr\$ or cerebell\$ or intracran\$ or intracerebral) adj5 (isch?emi\$ or infarct\$ or thrombo\$ or emboli\$ or occlus\$)).tw.

4. ((brain\$ or cerebr\$ or cerebell\$ or intracerebral or intracranial or subarachnoid) adj5 (haemorrhage\$ or hemorrhage\$ or haematoma\$ or hematoma\$ or bleed\$)).tw.

5. hemiplegia/ or exp paresis/

6. (hemipleg\$ or hemipar\$ or paresis or paretic).tw.

7. brain injuries/ or brain injury, chronic/

8. or/1-7

9. acupuncture/ or acupuncture therapy/ or acupuncture analgesia/ or acupuncture, ear/ or electroacupuncture/ or meridians/ or acupuncture points/ or trigger points/

10. (acupuncture\$ or electroacupuncture or electro-acupuncture or acupoint\$ or meridians or needling).tw.

11. ((meridian or non-meridian or trigger) adj10 point\$).tw.

12. 9 or 10 or 11

13. 8 and 12

14. Randomized Controlled Trials as Topic/

15. random allocation/

16. Controlled Clinical Trials as Topic/

17. control groups/
18. clinical trials as topic/ or clinical trials, phase i as topic/ or clinical trials, phase ii as topic/ or clinical trials, phase iii as topic/ or clinical trials, phase iv as topic/
19. double-blind method/
20. single-blind method/
21. Placebos/
22. placebo effect/
23. cross-over studies/
24. randomized controlled trial.pt.
25. controlled clinical trial.pt.
26. (clinical trial or clinical trial phase i or clinical trial phase ii or clinical trial phase iii or clinical trial phase iv).pt.
27. (random\$ or RCT or RCTs).tw.
28. (controlled adj5 (trial\$ or stud\$)).tw.
29. (clinical\$ adj5 trial\$).tw.
30. ((control or treatment or experiment\$ or intervention) adj5 (group\$ or subject\$ or patient\$)).tw.
31. (quasi-random\$ or quasi random\$ or pseudo-random\$ or pseudo random\$).tw.
32. ((control or experiment\$ or conservative) adj5 (treatment or therapy or procedure or manage\$)).tw.
33. ((singl\$ or doubl\$ or tripl\$ or trebl\$) adj5 (blind\$ or mask\$)).tw.
34. (cross-over or cross over or crossover).tw.
35. (placebo\$ or sham).tw.
36. trial.ti.
37. (assign\$ or allocat\$).tw.
38. controls.tw.
39. or/14-38
40. 13 and 39
41. exp animals/ not humans/
42. 40 not 41

Appendix 4. EMBASE search strategy

EMBASE (Ovid) November 2005 to July 2015

1. cerebrovascular disease/ or exp basal ganglion hemorrhage/ or exp brain hematoma/ or exp brain hemorrhage/ or exp brain infarction/ or exp brain ischemia/ or exp carotid artery disease/ or cerebral artery disease/ or exp cerebrovascular accident/ or exp intracranial aneurysm/ or exp occlusive cerebrovascular disease/ or stroke unit/ or stroke patient/
2. (stroke or poststroke or post-stroke or cerebrovasc\$ or brain vasc\$ or cerebral vasc\$ or cva\$ or apoplex\$ or SAH).tw.
3. ((brain\$ or cerebr\$ or cerebell\$ or intracran\$ or intracerebral) adj5 (isch?emi\$ or infarct\$ or thrombo\$ or emboli\$ or occlus\$)).tw.
4. ((brain\$ or cerebr\$ or cerebell\$ or intracerebral or intracranial or subarachnoid) adj5 (haemorrhage\$ or hemorrhage\$ or haematoma\$ or hematoma\$ or bleed\$)).tw.
5. hemiparesis/ or hemiplegia/ or paresis/

6. (hemipleg\$ or hemipar\$ or paresis or paretic).tw.
7. brain injury/ or acquired brain injury/
8. or/1-7
9. acupuncture/ or acupuncture analgesia/ or electroacupuncture/ or acupuncture needle/
10. (acupuncture\$ or electroacupuncture or electro-acupuncture or acupoint\$ or meridians or needling).tw.
11. ((meridian or non-meridian or trigger) adj10 point\$).tw.
12. 9 or 10 or 11
13. 8 and 12
14. Randomized Controlled Trial/ or "randomized controlled trial (topic)"/
15. Randomization/
16. Controlled clinical trial/ or "controlled clinical trial (topic)"/
17. control group/ or controlled study/
18. clinical trial/ or "clinical trial (topic)"/ or phase 1 clinical trial/ or phase 2 clinical trial/ or phase 3 clinical trial/ or phase 4 clinical trial/
19. Crossover Procedure/
20. Double Blind Procedure/
21. Single Blind Procedure/ or triple blind procedure/
22. placebo/ or placebo effect/
23. (random\$ or RCT or RCTs).tw.
24. (controlled adj5 (trial\$ or stud\$)).tw.
25. (clinical\$ adj5 trial\$).tw.
26. ((control or treatment or experiment\$ or intervention) adj5 (group\$ or subject\$ or patient\$)).tw.
27. (quasi-random\$ or quasi random\$ or pseudo-random\$ or pseudo random\$).tw.
28. ((control or experiment\$ or conservative) adj5 (treatment or therapy or procedure or manage\$)).tw.
29. ((singl\$ or doubl\$ or tripl\$ or trebl\$) adj5 (blind\$ or mask\$)).tw.
30. (cross-over or cross over or crossover).tw.
31. (placebo\$ or sham).tw.
32. trial.ti.
33. (assign\$ or allocat\$).tw.
34. controls.tw.
35. or/14-34
36. 13 and 35
37. (exp animals/ or exp invertebrate/ or animal experiment/ or animal model/ or animal tissue/ or animal cell/ or nonhuman/) not (human/ or normal human/ or human cell/)
38. 36 not 37

Appendix 5. CINAHL search strategy

CINAHL (EBSCO) November 2005 to July 2015

S1 -(MH "Cerebrovascular Disorders") OR (MH "Basal Ganglia Cerebrovascular Disease+") OR (MH "Carotid Artery Diseases+") OR (MH "Cerebral Ischemia+") OR (MH "Cerebral Vasospasm") OR (MH "Intracranial Arterial Diseases+") OR (MH "Intracranial Embolism and Thrombosis") OR (MH "Intracranial Hemorrhage+") OR (MH "Stroke") OR (MH "Vertebral Artery Dissections")

S2 -(MH "Stroke Patients") OR (MH "Stroke Units")

S3 -TI (stroke* or poststroke or apoplex* or cerebral vasc* or brain vasc* or cerebrovasc* or cva* or SAH) or AB (stroke* or poststroke or apoplex* or cerebral vasc* or brain vasc* or cerebrovasc* or cva* or SAH)

S4 -TI (brain or cerebr* or cerebell* or intracran* or intracerebral) or AB (brain or cerebr* or cerebell* or intracran* or intracerebral)

S5 -TI (ischemi* or ischaemi* or infarct* or thrombo* or emboli* or occlus*) or AB (ischemi* or ischaemi* or infarct* or thrombo* or emboli* or occlus*)

S6 -S4 and S5

S7 -TI (brain* or cerebr* or cerebell* or intracerebral or intracran* or subarachnoid) or AB (brain* or cerebr* or cerebell* or intracerebral or intracran* or subarachnoid)

S8 -TI (haemorrhage* or hemorrhage* or haematoma* or hematoma* or bleed*) or AB (haemorrhage* or hemorrhage* or haematoma* or hematoma* or bleed*)

S9 -S7 and S8

S10 -(MH "Hemiplegia")

S11 -TI (hemipleg* or hemipar* or paresis or paretic) or AB (hemipleg* or hemipar* or paresis or paretic)

S12 -(MH "Brain Injuries")

S13 -S1 OR S2 OR S3 OR S6 OR S9 OR S10 OR S11 OR S12

S14 -(MH "Acupuncture") OR (MH "Acupuncture Analgesia") OR (MH "Acupuncture Anesthesia") OR (MH "Acupuncture, Ear") OR (MH "Electroacupuncture") OR (MH "Meridians") OR (MH "Acupuncture Points") OR (MH "Acupuncturists") OR (MH "Trigger Point")

S15 -TI (acupuncture* or electroacupuncture or electro-acupuncture or acupoint* or meridians or needling) OR AB (acupuncture* or electroacupuncture or electro-acupuncture or acupoint* or meridians or needling)

S16 -TI ((meridian or non-meridian or trigger) N10 point*) or AB ((meridian or non-meridian or trigger) N10 point*)

S17 -S14 OR S15 OR S16

S18 -S13 AND S17

Appendix 6. AMED search strategy

AMED (Ovid) November 2005 to July 2015

1. cerebrovascular disorders/ or cerebral hemorrhage/ or cerebral infarction/ or cerebral ischemia/ or cerebrovascular accident/ or stroke/

2. (stroke or poststroke or post-stroke or cerebrovasc\$ or brain vasc\$ or cerebral vasc\$ or cva\$ or apoplex\$ or SAH).tw.

3. ((brain\$ or cerebr\$ or cerebell\$ or intracran\$ or intracerebral) adj5 (isch?emi\$ or infarct\$ or thrombo\$ or emboli\$ or occlus\$)).tw.

4. ((brain\$ or cerebr\$ or cerebell\$ or intracerebral or intracranial or subarachnoid) adj5 (haemorrhage\$ or hemorrhage\$ or haematoma\$ or hematoma\$ or bleed\$)).tw.

5. hemiplegia/

6. (hemipleg\$ or hemipar\$ or paresis or paretic).tw.

7. brain injuries/

8. or/1-7

9. acupuncture/ or acupuncture therapy/ or acupoints/ or neiguan/ or acupuncture analgesia/ or ear acupuncture/ or electroacupuncture/ or meridians/ or needling/ or scalp acupuncture/

10. (acupuncture\$ or electroacupuncture or electro-acupuncture or acupoint\$ or meridians or needling).tw.

11. ((meridian or non-meridian or trigger) adj10 point\$).tw.

12. 9 or 10 or 11

13. 8 and 12

Appendix 7. CBM search strategy

#1 中风

#2 卒中

#3 脑血管

#4 脑*塞

#5 脑*血

#6 脑*栓

#7 蛛网膜下腔出血

#8 主题词="中风/全部副主题/全部树"

#9 主题词="脑血管意外/全部副主题/全部树"

#10 主题词="垂体卒中/全部副主题"

#11 主题词="梗塞, 大脑中动脉/全部副主题"

#12 主题词="梗塞, 大脑前动脉/全部副主题"

#13 主题词="梗塞, 大脑后动脉/全部副主题"

#14 主题词="中风后遗症/全部副主题"

#15 主题词="中风先兆症/全部副主题"

#16 主题词="蛛网膜下腔出血/全部副主题/全部树"

#17 #1 or #2 or #3 or #4 or #5 or #6 or #8 or #9 or #10 or #11 or #12 or #13 or #14 or #15

#18 随机

#19 盲法

#20 单盲

#21 双盲

#22 三盲

#23 安慰剂

#24 主题词="随机对照试验 [文献类型]"

#25 主题词="随机分配"

#26 主题词="随机对照试验/全部副主题"

#27 #18 or #19 or #20 or #21 or #22 or #23 or #24 or #25 or #26

#28 针

#29 电针
#30 长针
#31 芒针
#32 皮下针
#33 火针
#34 头针
#35 手捻针
#36 巨针
#37 头皮针
#38 体针
#39 温针
#40 透刺
#41 巨刺
#42 针法
#43 刺法
#44 眼针
#45 磁极针
#46 毫针
#47 缪刺
#48 皮内针
#49 鑱针
#50 园针
#51 鍤针
#52 锋针
#53 铍针
#54 圆利针
#55 大针

#56 #28 or #29 or #30 or #31 or #32 or #33 or #34 or #35 or #36 or #37 or #38 or #39 or #40 or #41 or #42 or #43 or #44 or #45 or #46 or #47 or #48 or #49 or #50 or #51 or #52 or #52 or #53 or #54 or #55

Appendix 8. CNKI search strategy

(主题=中风 OR 主题=卒中 OR 主题=脑血管 OR 主题=脑梗塞 OR 主题=脑梗死 OR 主题=脑血栓 OR 主题=脑栓塞 OR 主题=脑出血 OR 主题=脑溢血 OR 主题=蛛网膜下腔出血) AND (主题=随机 OR 主题=盲法 OR 主题=单盲 OR 主题=双盲 OR 主题=三盲 OR 主题=安慰剂) AND (主题=针灸 OR 主题=针刺 OR 主题=电针 OR 主题=芒针 OR 主题=皮下针 OR 主题=火针 OR 主题=头针 OR 主题=手捻针 OR 主题=针刀 OR 主题=长针 OR 主题=巨针 OR 主题=头皮针 OR 主题=体针 OR 主题=温针 OR 主题=透刺 OR 主题=巨刺 OR 主题=针法 OR 主题=刺法 OR 主题=眼针 OR 主题=磁极针 OR 主题=毫针 OR 主题=缪刺 OR 主题=皮内针 OR 主题=鑱针 OR 主题=园针 OR 主题=鍤针 OR 主题=锋针 OR 主题=铍针 OR 主题=圆利针 OR 主题=大针)

Appendix 9. VIP search strategy

(U=中风+U=卒中+U=脑血管+U=脑梗塞+U=脑梗死+U=脑血栓+U=脑栓塞+U=脑出血+U=脑溢血+U=蛛网膜下腔出血)*(U=随机+U=盲法+U=安慰剂)*(U=针灸+U=针刺+U=电针+U=针法+U=刺法+U=磁极针)*全部期刊*年=2005-2015

Appendix 10. Wanfang search strategy

((中风或卒中或脑血管*或脑梗*或脑*栓或脑栓*或脑*血或蛛网膜下腔出血) and (((("随机") or ("盲法")) or ("单盲")) or ("双盲")) or ("安慰剂"))) and (((("针灸") or ("针法")) or ("针刺")) or ("电针")) and ("刺法"))

Appendix 11. Risk of bias assessment tool

Potential source of bias	Assessment criteria
Random sequence generation Selection bias (biased allocation to interventions) due to inadequate generation of a randomised sequence	<p><i>Low risk of bias:</i> Random number table; computer random number generator; coin tossing; shuffling cards or envelopes; throwing dice; drawing of lots; minimization (minimization may be implemented without a random element, and this is considered to be equivalent to being random).</p> <p><i>High risk of bias:</i> Sequence generated by odd or even date of birth; date (or day) of admission; sequence generated by hospital or clinic record number; allocation by judgement of the clinician; by preference of the participant; based on the results of a laboratory test or a series of tests; by availability of the intervention.</p> <p><i>Unclear:</i> Insufficient information about the sequence generation process to permit judgement.</p>
Allocation concealment Selection bias (biased allocation to interventions) due to inadequate concealment of allocations prior to assignment	<p><i>Low risk of bias:</i> Randomisation method described that would not allow investigator/participant to know or influence intervention group before eligible participant entered in the study (e.g. central allocation, including telephone, web-based, and pharmacy-controlled, randomisation; sequentially numbered drug containers of identical appearance; sequentially numbered, opaque, sealed envelopes).</p> <p><i>High risk of bias:</i> Using an open random allocation schedule (e.g. a list of random numbers); assignment envelopes were used without appropriate safeguards (e.g. if envelopes were unsealed or non-opaque or not sequentially numbered); alternation or rotation; date of birth; case record number; any other explicitly unconcealed procedure.</p> <p><i>Unclear:</i> Randomisation stated but no information on method used is available.</p>
Blinding of participants and personnel Performance bias due to knowledge of the allocated interventions by participants and personnel during the study	<p><i>Low risk of bias:</i> No blinding or incomplete blinding, but the review authors judge that the outcome is not likely to be influenced by lack of blinding; blinding of participants and key study personnel ensured, and unlikely that the blinding could have been broken.</p> <p><i>High risk of bias:</i> No blinding or incomplete blinding, and the outcome is likely to be influenced by lack of blinding; blinding of key study participants and personnel attempted, but likely that the blinding could have been broken, and the outcome is likely to be influenced by lack of blinding.</p> <p><i>Unclear:</i> Insufficient information to permit judgement</p>
Blinding of outcome assessment Detection bias due to knowledge of the allocated interventions by outcome assessors.	<p><i>Low risk of bias:</i> No blinding of outcome assessment, but the review authors judge that the outcome measurement is not likely to be influenced by lack of blinding; blinding of outcome assessment ensured, and unlikely that the blinding could have been broken.</p> <p><i>High risk of bias:</i> No blinding of outcome assessment, and the outcome measurement is likely to be influenced by lack of blinding; blinding of outcome assessment, but likely that the blinding could have been broken, and the outcome measurement is likely to be influenced by lack of blinding.</p> <p><i>Unclear:</i> Insufficient information to permit judgement</p>

(Continued)

Incomplete outcome data

Attrition bias due to amount, nature or handling of incomplete outcome data.

Low risk of bias: No missing outcome data; reasons for missing outcome data unlikely to be related to true outcome (for survival data, censoring unlikely to be introducing bias); missing outcome data balanced in numbers across intervention groups, with similar reasons for missing data across groups; for dichotomous outcome data, the proportion of missing outcomes compared with observed event risk not enough to have a clinically relevant impact on the intervention effect estimate; for continuous outcome data, plausible effect size (difference in means or standardized difference in means) among missing outcomes not enough to have a clinically relevant impact on observed effect size; missing data have been imputed using appropriate methods.

High risk of bias: Reason for missing outcome data likely to be related to true outcome, with either imbalance in numbers or reasons for missing data across intervention groups; for dichotomous outcome data, the proportion of missing outcomes compared with observed event risk enough to induce clinically relevant bias in intervention effect estimate; for continuous outcome data, plausible effect size (difference in means or standardized difference in means) among missing outcomes enough to induce clinically relevant bias in observed effect size; 'as-treated' analysis done with substantial departure of the intervention received from that assigned at randomisation; potentially inappropriate application of simple imputation.

Unclear: Insufficient information to permit judgement

Selective reporting

Reporting bias due to selective outcome reporting

Low risk of bias: The study protocol is available and all of the study's pre-specified (primary and secondary) outcomes that are of interest in the review have been reported in the pre-specified way; the study protocol is not available but it is clear that the published reports include all expected outcomes, including those that were pre-specified (convincing text of this nature may be uncommon).

High risk of bias: Not all of the study's pre-specified primary outcomes have been reported; one or more primary outcomes is reported using measurements, analysis methods or subsets of the data (e.g. subscales) that were not pre-specified; one or more reported primary outcomes were not pre-specified (unless clear justification for their reporting is provided, such as an unexpected adverse effect); one or more outcomes of interest in the review are reported incompletely so that they cannot be entered in a meta-analysis; the study report fails to include results for a key outcome that would be expected to have been reported for such a study.

Unclear: Insufficient information to permit judgement

Other bias

Bias due to problems not covered elsewhere in the table

Low risk of bias: The study appears to be free of other sources of bias.

High risk of bias: Had a potential source of bias related to the specific study design used; stopped early due to some data-dependent process (including a formal-stopping rule); had extreme baseline imbalance; has been claimed to have been fraudulent; had some other problem.

Unclear: Insufficient information to assess whether an important risk of bias exists; insufficient rationale or evidence that an identified problem will introduce bias.

WHAT'S NEW

Date	Event	Description
12 January 2016	New citation required but conclusions have not changed	Compared with the 2006 version of this review, a substantial amount of new information has been included, but there is no change to the main conclusions in this updated review.
12 January 2016	New search has been performed	We have updated the searches to July 2015. We included 26 new trials (1889 participants) in this version. There are now 31 trials, with 2257 participants, included in this updated review.

HISTORY

Protocol first published: Issue 2, 2003

Review first published: Issue 3, 2006

Date	Event	Description
20 October 2008	Amended	Contact details updated
15 July 2008	Amended	Converted to new review format.

CONTRIBUTIONS OF AUTHORS

Wu HM: developing the search strategy, assessment of studies, data extraction, data entry, data analysis, writing of protocol and review.

Yang A: developing the search strategy, assessment of studies, data extraction, data entry, data analysis, writing of review.

Tang JL: data analysis, resolution of disagreements, writing protocol and review.

Xu Li: assessment of studies, data extraction, data entry, suggestions and corrections.

Yang M: data extraction, suggestions and corrections.

Liu GJ: data analysis, suggestions and corrections.

DECLARATIONS OF INTEREST

Ai Yang: none known.

Hong Mei Wu: none known.

Jin-Ling Tang: none known.

Li Xu: none known.

Ming Yang: none known.

Guan J Liu: none known.

SOURCES OF SUPPORT

Internal sources

- Hong Kong Branch of Chinese Cochrane Center, China.
- Department of Community & Family Medicine, Chinese University of Hong Kong, China.
- Chinese Cochrane Center, West China Hospital, Si Chuan University, China.

External sources

- Hong Kong Croucher Foundation, China.

INDEX TERMS

Medical Subject Headings (MeSH)

*Acupuncture Therapy; *Stroke Rehabilitation; Activities of Daily Living; Convalescence; Randomized Controlled Trials as Topic

MeSH check words

Humans