Cochrane
Library

Cochrane Database of Systematic Reviews

Medical treatment for early fetal death (less than 24 weeks) (Review)

Neilson JP, Hickey M, Vazquez JC

Neilson JP, Hickey M, Vazquez JC.

Medical treatment for early fetal death (less than 24 weeks).

Cochrane Database of Systematic Reviews 2006, Issue 3. Art. No.: CD002253.

DOI: 10.1002/14651858.CD002253.pub3.

www.cochranelibrary.com

TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
BACKGROUND	2
OBJECTIVES	3
METHODS	3
RESULTS	4
DISCUSSION	6
AUTHORS' CONCLUSIONS	6
ACKNOWLEDGEMENTS	6
REFERENCES	7
CHARACTERISTICS OF STUDIES	17
DATA AND ANALYSES	36
Analysis 1.1. Comparison 1 Vaginal misoprostol versus placebo, Outcome 1 Complete miscarriage < 24 hours after treatment.	41
Analysis 1.2. Comparison 1 Vaginal misoprostol versus placebo, Outcome 2 Complete miscarriage < 48 hours.	41
Analysis 1.3. Comparison 1 Vaginal misoprostol versus placebo, Outcome 3 Complete miscarriage without ERPC day 7.	42
Analysis 1.4. Comparison 1 Vaginal misoprostol versus placebo, Outcome 4 Uterine curettage.	43
Analysis 1.5. Comparison 1 Vaginal misoprostol versus placebo, Outcome 5 Opiates for pain relief.	43
Analysis 1.6. Comparison 1 Vaginal misoprostol versus placebo, Outcome 6 Blood transfusion.	44
Analysis 1.7. Comparison 1 Vaginal misoprostol versus placebo, Outcome 7 Haemoglobin difference > 10 g/L.	44
Analysis 1.8. Comparison 1 Vaginal misoprostol versus placebo, Outcome 8 Nausea.	45
Analysis 1.9. Comparison 1 Vaginal misoprostol versus placebo, Outcome 9 Diarrhoea.	45
Analysis 1.10. Comparison 1 Vaginal misoprostol versus placebo, Outcome 10 Fever.	46
Analysis 1.11. Comparison 1 Vaginal misoprostol versus placebo, Outcome 11 Uterine perforation.	46
Analysis 1.12. Comparison 1 Vaginal misoprostol versus placebo, Outcome 12 Vaginal bleeding 2 weeks after treatment.	47
Analysis 1.13. Comparison 1 Vaginal misoprostol versus placebo, Outcome 13 Satisfaction with treatment.	47
Analysis 2.1. Comparison 2 Vaginal misoprostol versus surgical evacuation of uterus, Outcome 1 Surgical evacuation of uterus.	48
Analysis 2.2. Comparison 2 Vaginal misoprostol versus surgical evacuation of uterus, Outcome 2 Post-treatment haematocrit (%).	48
Analysis 2.3. Comparison 2 Vaginal misoprostol versus surgical evacuation of uterus, Outcome 3 Nausea.	49
Analysis 2.4. Comparison 2 Vaginal misoprostol versus surgical evacuation of uterus, Outcome 4 Pain relief.	50
Analysis 2.5. Comparison 2 Vaginal misoprostol versus surgical evacuation of uterus, Outcome 5 Diarrhoea.	50
Analysis 2.6. Comparison 2 Vaginal misoprostol versus surgical evacuation of uterus, Outcome 6 Uterine perforation.	51
Analysis 2.7. Comparison 2 Vaginal misoprostol versus surgical evacuation of uterus, Outcome 7 Asherman syndrome.	51
Analysis 3.1. Comparison 3 Vaginal misoprostol versus vaginal gemeprost, Outcome 1 Miscarriage < 24 hours.	52
Analysis 3.2. Comparison 3 Vaginal misoprostol versus vaginal gemeprost, Outcome 2 Temperature > 38 degrees C.	52
Analysis 3.3. Comparison 3 Vaginal misoprostol versus vaginal gemeprost, Outcome 3 Vomiting.	53
Analysis 3.4. Comparison 3 Vaginal misoprostol versus vaginal gemeprost, Outcome 4 Diarrhoea.	54
Analysis 3.5. Comparison 3 Vaginal misoprostol versus vaginal gemeprost, Outcome 5 Opiate analgesia.	54
Analysis 4.1. Comparison 4 Vaginal misoprostol versus vaginal prostaglandin E1/2, Outcome 1 Surgical evacuation.	55
Analysis 4.2. Comparison 4 Vaginal misoprostol versus vaginal prostaglandin E1/2, Outcome 2 Blood transfusion.	55
Analysis 4.3. Comparison 4 Vaginal misoprostol versus vaginal prostaglandin E1/2, Outcome 3 Hospital stay (days).	56
Analysis 4.4. Comparison 4 Vaginal misoprostol versus vaginal prostaglandin E1/2, Outcome 4 Complete miscarriage.	56
Analysis 4.5. Comparison 4 Vaginal misoprostol versus vaginal prostaglandin E1/2, Outcome 5 Nausea.	57
Analysis 5.1. Comparison 5 Vaginal misoprostol lower versus higher dose regimens, Outcome 1 Miscarriage.	57
Analysis 5.2. Comparison 5 Vaginal misoprostol lower versus higher dose regimens, Outcome 2 Fever.	58
Analysis 5.3. Comparison 5 Vaginal misoprostol lower versus higher dose regimens, Outcome 3 Nausea.	59
Analysis 5.4. Comparison 5 Vaginal misoprostol lower versus higher dose regimens, Outcome 4 Diarrhoea.	59
Analysis 6.1. Comparison 6 Vaginal misoprostol wet versus dry vaginal preparations, Outcome 1 Miscarriage < 3 days.	60

Analysis 6.2. Comparison 6 Vaginal misoprostol wet versus dry vaginal preparations, Outcome 2 Miscarriage < 8 days.	60
Analysis 6.3. Comparison 6 Vaginal misoprostol wet versus dry vaginal preparations, Outcome 3 Miscarriage < 15 days.	61
Analysis 6.4. Comparison 6 Vaginal misoprostol wet versus dry vaginal preparations, Outcome 4 Miscarriage < 30 days.	61
Analysis 6.5. Comparison 6 Vaginal misoprostol wet versus dry vaginal preparations, Outcome 5 Diarrhoea < 48 hours after treatment.	62
Analysis 6.6. Comparison 6 Vaginal misoprostol wet versus dry vaginal preparations, Outcome 6 Chills < 48 hours of treatment.	62
Analysis 6.7. Comparison 6 Vaginal misoprostol wet versus dry vaginal preparations, Outcome 7 Vomiting < 48 hours of treatment.	63
Analysis 6.8. Comparison 6 Vaginal misoprostol wet versus dry vaginal preparations, Outcome 8 Would wish/probably wish same treatment in future nonviable pregnancy.	63
Analysis 7.1. Comparison 7 Vaginal misoprostol + methotrexate versus vaginal misoprostol alone, Outcome 1 Miscarriage not complete.	64
Analysis 7.4. Comparison 7 Vaginal misoprostol + methotrexate versus vaginal misoprostol alone, Outcome 4 Additional surgical evacuation.	64
Analysis 7.5. Comparison 7 Vaginal misoprostol + methotrexate versus vaginal misoprostol alone, Outcome 5 Haemorrhage.	65
Analysis 7.6. Comparison 7 Vaginal misoprostol + methotrexate versus vaginal misoprostol alone, Outcome 6 Pain relief.	65
Analysis 8.1. Comparison 8 Vaginal misoprostol plus laminaria tents versus vaginal misoprostol alone, Outcome 1 Miscarriage < 24 hours.	66
Analysis 8.2. Comparison 8 Vaginal misoprostol plus laminaria tents versus vaginal misoprostol alone, Outcome 2 Miscarriage < 48 hours.	66
Analysis 9.1. Comparison 9 Oral misoprostol versus vaginal misoprostol, Outcome 1 Complete miscarriage.	67
Analysis 9.2. Comparison 9 Oral misoprostol versus vaginal misoprostol, Outcome 2 Vomiting.	68
Analysis 9.9. Comparison 9 Oral misoprostol versus vaginal misoprostol, Outcome 9 Nausea.	68
Analysis 9.10. Comparison 9 Oral misoprostol versus vaginal misoprostol, Outcome 10 Diarrhoea.	69
Analysis 9.12. Comparison 9 Oral misoprostol versus vaginal misoprostol, Outcome 12 Pain (visual analogue scale).	69
Analysis 9.13. Comparison 9 Oral misoprostol versus vaginal misoprostol, Outcome 13 Fever.	70
Analysis 9.14. Comparison 9 Oral misoprostol versus vaginal misoprostol, Outcome 14 Women's satisfaction with treatment.	70
Analysis 9.15. Comparison 9 Oral misoprostol versus vaginal misoprostol, Outcome 15 Time to delivery (hours).	71
Analysis 9.16. Comparison 9 Oral misoprostol versus vaginal misoprostol, Outcome 16 Oxytocin infusion.	71
Analysis 9.17. Comparison 9 Oral misoprostol versus vaginal misoprostol, Outcome 17 Manual removal of placenta.	72
Analysis 10.1. Comparison 10 Oral misoprostol + mifepristone versus expectant management, Outcome 1 Empty uterine cavity at day 5.	72
Analysis 10.2. Comparison 10 Oral misoprostol + mifepristone versus expectant management, Outcome 2 Urgent surgical evacuation for bleeding.	73
Analysis 10.3. Comparison 10 Oral misoprostol + mifepristone versus expectant management, Outcome 3 Pelvic inflammatory disease.	73
Analysis 10.4. Comparison 10 Oral misoprostol + mifepristone versus expectant management, Outcome 4 Pain (visual analogue scale day 5).	74
Analysis 10.5. Comparison 10 Oral misoprostol + mifepristone versus expectant management, Outcome 5 Sick leave (days).	74
Analysis 10.6. Comparison 10 Oral misoprostol + mifepristone versus expectant management, Outcome 6 Bleeding (days).	75
Analysis 10.7. Comparison 10 Oral misoprostol + mifepristone versus expectant management, Outcome 7 Satisfaction with treatment (visual analogue scale day 14).	75
Analysis 11.1. Comparison 11 Sublingual misoprostol versus vaginal misoprostol, Outcome 1 Complete miscarriage.	76
Analysis 11.2. Comparison 11 Sublingual misoprostol versus vaginal misoprostol, Outcome 2 Nausea.	76
Analysis 11.3. Comparison 11 Sublingual misoprostol versus vaginal misoprostol, Outcome 3 Vomiting.	77
Analysis 11.4. Comparison 11 Sublingual misoprostol versus vaginal misoprostol, Outcome 4 Diarrhoea.	77
Analysis 11.5. Comparison 11 Sublingual misoprostol versus vaginal misoprostol, Outcome 5 Haemoglobin day 43.	78
Analysis 11.6. Comparison 11 Sublingual misoprostol versus vaginal misoprostol, Outcome 6 Intolerable pain.	78

Analysis 11.7. Comparison 11 Sublingual misoprostol versus vaginal misoprostol, Outcome 7 Satisfied with treatment.	79
Analysis 12.1. Comparison 12 Mifepristone versus placebo, Outcome 1 Miscarriage < 48 hours.	79
Analysis 12.2. Comparison 12 Mifepristone versus placebo, Outcome 2 Miscarriage < 3 days.	80
Analysis 12.3. Comparison 12 Mifepristone versus placebo, Outcome 3 Miscarriage < 4 days.	80
Analysis 12.4. Comparison 12 Mifepristone versus placebo, Outcome 4 Miscarriage < 5 days.	81
Analysis 12.5. Comparison 12 Mifepristone versus placebo, Outcome 5 Vaginal bleeding before day 5.	81
Analysis 12.6. Comparison 12 Mifepristone versus placebo, Outcome 6 Pain before day 5.	82
Analysis 13.1. Comparison 13 Vaginal gemeprost versus surgical evacuation of uterus, Outcome 1 Surgical evacuation.	82
Analysis 13.2. Comparison 13 Vaginal gemeprost versus surgical evacuation of uterus, Outcome 2 Perforation of uterus.	83
Analysis 13.3. Comparison 13 Vaginal gemeprost versus surgical evacuation of uterus, Outcome 3 Nausea.	83
WHAT'S NEW	83
HISTORY	84
CONTRIBUTIONS OF AUTHORS	84
DECLARATIONS OF INTEREST	84
SOURCES OF SUPPORT	84
DIFFERENCES BETWEEN PROTOCOL AND REVIEW	85
INDEX TERMS	85

Medical treatment for early fetal death (less than 24 weeks)

James P Neilson¹, Martha Hickey², Juan C Vazquez³

¹Department of Women's and Children's Health, The University of Liverpool, Liverpool, UK. ²The University of Melbourne, The Royal Women's Hospital, Melbourne, Australia. ³Departamento de Salud Reproductiva, Instituto Nacional de Endocrinología (INEN), Habana, Cuba

Contact address: James P Neilson, Department of Women's and Children's Health, The University of Liverpool, First Floor, Liverpool Women's NHS Foundation Trust, Crown Street, Liverpool, L8 7SS, UK. jneilson@liverpool.ac.uk.

Editorial group: Cochrane Pregnancy and Childbirth Group.

Publication status and date: Edited (no change to conclusions), published in Issue 11, 2012.

Review content assessed as up-to-date: 24 April 2006.

Citation: Neilson JP, Hickey M, Vazquez JC. Medical treatment for early fetal death (less than 24 weeks). *Cochrane Database of Systematic Reviews* 2006, Issue 3. Art. No.: CD002253. DOI: 10.1002/14651858.CD002253.pub3.

Copyright © 2012 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

In most pregnancies that miscarry, arrest of embryonic or fetal development occurs some time (often weeks) before the miscarriage occurs. Ultrasound examination can reveal abnormal findings during this phase by demonstrating anembryonic pregnancies or embryonic or fetal death. Treatment before 14 weeks has traditionally been surgical but medical treatments may be effective, safe, and acceptable, as may be waiting for spontaneous miscarriage.

Objectives

To assess the effectiveness, safety and acceptability of any medical treatment for early pregnancy failure (anembryonic pregnancies or embryonic and fetal deaths before 24 weeks).

Search methods

We searched the Cochrane Pregnancy and Childbirth Group Trials Register (30 November 2005). We updated this search on 6 August 2012 and added the results to the awaiting classification section of the review.

Selection criteria

Randomised trials comparing medical treatment with another treatment (e.g. surgical evacuation), or placebo, or no treatment for early pregnancy failure. Quasi-random studies were excluded.

Data collection and analysis

Data were extracted unblinded.

Main results

Twenty four studies (1888 women) were included.

Vaginal misoprostol hastens miscarriage (complete or incomplete) when compared with placebo: e.g. miscarriage less than 24 hours (two trials, 138 women, relative risk (RR) 4.73, 95% confidence interval (CI) 2.70 to 8.28), with less need for uterine curettage (two trials, 104 women, RR 0.40, 95% CI 0.26 to 0.60) and no significant increase in nausea or diarrhoea. Lower-dose regimens of vaginal misoprostol tend to be less effective in producing miscarriage (three trials, 247 women, RR 0.85, 95% CI 0.72 to 1.00) with

similar incidence of nausea. There seems no clear advantage to administering a 'wet' preparation of vaginal misoprostol or of adding methotrexate, or of using laminaria tents after 14 weeks. Vaginal misoprostol is more effective than vaginal prostaglandin E in avoiding surgical evacuation. Oral misoprostol was less effective than vaginal misoprostol in producing complete miscarriage (two trials, 218 women, RR 0.90, 95% CI 0.82 to 0.99). Sublingual misoprostol had equivalent efficacy to vaginal misoprostol in inducing complete miscarriage but was associated with more frequent diarrhoea. The two trials of mifepristone treatment generated conflicting results. There was no statistically significant difference between vaginal misoprostol and gemeprost in the induction of miscarriage for fetal death after 13 weeks.

Authors' conclusions

Available evidence from randomised trials supports the use of vaginal misoprostol as a medical treatment to terminate non-viable pregnancies before 24 weeks. Further research is required to assess effectiveness and safety, optimal route of administration and dose. Conflicting findings about the value of mifepristone need to be resolved by additional study.

[Note: the 108 citations in the awaiting classification section of the review may alter the conclusions of the review once assessed.]

PLAIN LANGUAGE SUMMARY

Medical treatment for early fetal death (less than 24 weeks)

Medical treatments for inevitable miscarriage.

Pregnancies that miscarry can sometimes be identified earlier at an ultrasound scan if the loss is due to the baby having died or no baby having developed. In the past, treatment before 14 weeks has usually been by surgery (D&C) but drugs have now been developed which may be helpful, or waiting for the miscarriage to happen may be a better alternative. The review of trials assessed various potential drug treatments using different routes and different doses, compared with waiting for the miscarriage. This review identified 24 studies involving 1888 women of less than 24 weeks gestation, where the baby had died in the uterus or the baby had not formed in the uterus. Most studies were of good quality. Vaginal misoprostol brought forward the time of the miscarriage, but the studies were too small to adequately assess potential adverse effects, including future fertility. Oral misoprostol seemed less effective than the vaginal route, and women took more sick-leave with the oral drugs. Some women may wish to hasten an inevitable miscarriage, and others may not. It appears that both forms of care can be available to women. Women who are breastfeeding an older baby may prefer to wait rather than have drug treatment. Further research is needed on drug doses, routes of administration and potential adverse effects, including future fertility, and also on women's views of drug treatment, surgery and waiting for spontaneous miscarriage.

BACKGROUND

The incidence of clinically obvious miscarriage is considered to be between 10% and 15% of all pregnancies, although the real incidence may be considerably higher ([Grudzinskas 1995](#); [Howie 1995](#); [Simpson 1991](#)).

The widespread use of ultrasound in early pregnancy for either specific reasons (for example, vaginal bleeding) or as a routine examination ([Neilson 1998](#)) reveals 'non-viable pregnancies' destined inevitably to miscarry in due course. These are termed 'anembryonic pregnancies' (formerly called 'blighted ova') if no embryo has developed within the gestation sac, or 'missed abortions' if an embryo or fetus is present, but is dead.

The protocol for this review aimed to combine trials of medical treatments for both non-viable pregnancies and for incomplete miscarriage but on further reflection, this was illogical. Non-viable pregnancies contain viable trophoblast (placental) tissue, which produces hormones, which may in theory make these pregnancies more susceptible to anti-hormone therapy and more resistant to uterotonic (stimulating uterine contractions) therapy than pregnancies in which (incomplete) miscarriage has already taken place. This review will therefore focus exclusively on non-viable pregnancies, before miscarriage. Another review will assess trials of medical treatments after miscarriage has occurred ([Vazquez 2000](#)). A further review compares expectant management with surgical treatment for miscarriage ([Nanda 2002](#)).

Traditionally, early non-viable pregnancies (less than 14 weeks) have been terminated by surgical evacuation. Later pregnancies (14 to 24 weeks) have been ended by medical induction of miscarriage.

Various types of medical treatment could be suitable as alternatives to surgical treatment: misoprostol is a prostaglandin E1 analogue, marketed for the prevention and treatment of peptic ulcers. Recognized as a potent method for terminating unwanted viable pregnancies (Costa 1993; Norman 1991), it is cheap, stable at room temperature and has few systemic effects, although vomiting, diarrhoea, hypertension and even potential teratogenicity when misoprostol fails to induce abortion have been reported (Fonseca 1991). Misoprostol has been shown to be an effective myometrial stimulant of the pregnant uterus, selectively binding to EP-2/EP-3 prostanoid receptors (Senior 1993). It is rapidly absorbed orally and vaginally. Vaginally absorbed serum levels are more prolonged and vaginal misoprostol may have locally mediated effects (Zieman 1997).

Misoprostol could be especially useful in developing countries, where transport and storage facilities are inadequate and the availability of uterotonic agents and blood is limited. Its use in obstetrics and gynecology has been explored, especially to induce first and second trimester abortion (Ashok 1998; Bugalho 1996), for the induction of labour (Alfirevic 2001; Hofmeyr 2003) and for the prevention of postpartum haemorrhage (Gulmezoglu 2004) despite the fact that it has not been registered for such use.

Other uterotonic drugs that could have a role would include ergometrine, oxytocin, and prostaglandin F2alpha.

The progesterone antagonist, mifepristone, is of value in terminating early unwanted pregnancies and may be of use in non-viable pregnancies and spontaneous miscarriage (Baulieu 1986; Kovacs 1984), alone or in combination with prostaglandin (Cameron 1986).

Methotrexate may be helpful in the medical treatment of ectopic pregnancy and might therefore have a place in the treatment of intrauterine non-viable pregnancies as well. It has also been used for the early termination of unwanted pregnancy, followed by a uterotonic agent such as misoprostol.

Although clotting problems occasionally occur in women with prolonged retention of a dead fetus, this is rare and does not usually happen within the first month after fetal death. There are, therefore, not pressing medical reasons to terminate non-viable pregnancies. Although, anecdotally, many women favour early termination, so-called 'expectant management' (that is, awaiting spontaneous miscarriage) is a legitimate alternative and this policy should be considered in clinical care and in planning trials.

OBJECTIVES

To assess, from clinical trials, the effectiveness and safety of different medical treatments for the termination of non-viable pregnancies, with reference to death or serious complications, additional surgical evacuation, blood transfusion, haemorrhage, blood loss, anaemia, days of bleeding, pain relief, pelvic infection, cervical damage, duration of stay in hospital, psychological effects, subsequent fertility, women's satisfaction and costs.

METHODS

Criteria for considering studies for this review

Types of studies

Randomised clinical trials comparing a medical treatment with another treatment (for example, surgical evacuation), or placebo, or no treatment to terminate non-viable pregnancies; random allocation to treatment and comparison groups; reasonable measures to ensure allocation concealment; and violations of allocated management not sufficient to materially affect outcomes. Quasi-random studies were excluded.

Types of participants

Women with non-viable pregnancies (i.e. where the embryo or fetus had died in utero, and in whom miscarriage would have happened inevitably in due course) if less than 24 weeks estimated gestational age. Subgroup analyses to be performed, if possible, for women at less than 14 weeks, and those between 14 and 23 weeks estimated gestational age.

Types of interventions

Trials were considered if they compared medical treatment with other methods (for example, expectant management, placebo or any other intervention including surgical evacuation). Comparisons between different routes of administration of medical treatment (for example, oral versus vaginal), or between different drugs or doses of drug, or duration or timing of treatment, were also included if data existed.

Types of outcome measures

Trials were considered if any of the following outcomes were reported.

Primary outcomes

1. Complete miscarriage (i.e. no pregnancy tissues remaining in uterus - based on clinical findings at surgery and/or ultrasound examination after a specific period).
2. Death or serious complications (e.g. uterine rupture, uterine perforation, hysterectomy, organ failure, intensive care unit admission).

Secondary outcomes

1. Additional surgical evacuation
2. Blood transfusion
3. Haemorrhage
4. Blood loss
5. Anemia
6. Days of bleeding
7. Pain relief
8. Pelvic infection
9. Cervical damage
10. Digestive disorders (nausea or vomiting or diarrhoea)
11. Hypertensive disorders
12. Duration of stay in hospital
13. Psychological effects
14. Subsequent fertility
15. Woman's satisfaction/acceptability of method
16. Costs

Search methods for identification of studies

Electronic searches

We searched the Cochrane Pregnancy and Childbirth Group Trials Register by contacting the Trials Search Co-ordinator (30 November 2005). We updated this search on 6 August 2012 and added the results to Studies awaiting classification.

The Cochrane Pregnancy and Childbirth Group's Trials Register is maintained by the Trials Search Co-ordinator and contains trials identified from:

1. monthly searches of the Cochrane Central Register of Controlled Trials (CENTRAL);
2. weekly searches of MEDLINE;
3. weekly searches of EMBASE;
4. handsearches of 30 journals and the proceedings of major conferences;
5. weekly current awareness alerts for a further 44 journals plus monthly BioMed Central email alerts.

Details of the search strategies for CENTRAL, MEDLINE and EMBASE, the list of handsearched journals and conference proceedings, and the list of journals reviewed via the current awareness service can be found in the 'Specialized Register' section

within the editorial information about the [Cochrane Pregnancy and Childbirth Group](#).

Trials identified through the searching activities described above are each assigned to a review topic (or topics). The Trials Search Co-ordinator searches the register for each review using the topic list rather than keywords.

We did not apply any language restrictions.

Data collection and analysis

We assessed all potential trials for eligibility according to the criteria specified in the protocol. A single author extracted data from each publication and co-authors checked the data. We resolved any discrepancies by discussion. In addition to the main outcome measures listed above, information on the setting of the study (country, type of population, socioeconomic status), the method of randomisation, a detailed description of the regimen used (drug(s), route, dose, frequency), definitions of the outcomes (if provided), and whether or not clinicians and participants were 'blind' to treatment allocated, were all collected. An intention-to-treat analysis was performed where possible. Any information on completeness of follow up was collected as well.

Trials were assessed for methodological quality using the standard Cochrane criteria of adequacy of allocation concealment:

- (A) adequate;
- (B) unclear;
- (C) inadequate;
- (D) allocation concealment was not used.

We collected information on blinding of outcome assessment and loss to follow up.

Separate comparisons were made of different drug regimens, grouped where appropriate by number of doses given and the route of administration. Summary relative risks were calculated using a fixed-effect model (providing there was no significant heterogeneity between trials - defined as I^2 greater than 50%). Because of the small number of trials and comparisons, it was impossible to perform sensitivity analysis using trial quality (A versus B, C, D).

RESULTS

Description of studies

See: [Characteristics of included studies](#); [Characteristics of excluded studies](#); [Characteristics of ongoing studies](#).

This review has included 24 studies comparing vaginal misoprostol to expectant management (Bagratee 2004), placebo (Herabutya 1997; Kovavisarach 2002; Lister 2005; Wood 2002), surgical evacuation (Demetroulis 2001; Graziosi 2004; Muffley 2002), oral or

sublingual misoprostol (Creinin 1997; Ngoc 2004; Tang 2003), other types of vaginal or intracervical prostaglandin preparation (Al Inizi 2003; Eng 1997*; Fadalla 2004*; Kara 1999*); different doses (Heard 2002; Kovavisarach 2005; Niromanesh 2005*) and preparations (Gilles 2004) of vaginal misoprostol; the addition to vaginal misoprostol of methotrexate (Autry 1999) or laminaria tents (Jain 1996*); mifepristone versus placebo (Lelaidier 1993); mifepristone plus oral misoprostol versus expectant management (Nielsen 1999), and vaginal gemeprost versus surgical evacuation (Egarter 1995).

The Bagratee 2004 trial used a comparison of vaginal misoprostol versus placebo to explore comparisons with expectant management (up to seven days) and, therefore, differed in concept from the Herabutya 1997 and Wood 2002 studies in which early surgical intervention occurred after, respectively, 24 and 48 hours.

Five of the 24 included studies addressed medical treatment of non-viable pregnancies after 14 weeks (Eng 1997*; Fadalla 2004*; Jain 1996*; Kara 1999*; Niromanesh 2005*). These are labelled with an asterisk for ease of interpretation.

There are additional trials that have included data on women with both non-viable pregnancies and incomplete miscarriages (for example, Ngai 2001). If these can be separated by the researchers, these data may be included in the future.

(One hundred reports from an updated search on 6 August 2012 have been added to Studies awaiting classification.)

Risk of bias in included studies

Thirteen studies used robust methods of allocation concealment (Autry 1999; Bagratee 2004; Creinin 1997; Demetroulis 2001; Gilles 2004; Graziosi 2004; Kovavisarach 2005; Lelaidier 1993; Lister 2005; Muffley 2002; Ngoc 2004; Tang 2003; Wood 2002). Nine reports failed to describe the process of randomisation (Al Inizi 2003; Egarter 1995; Fadalla 2004*; Herabutya 1997; Jain 1996*; Kara 1999*; Kovavisarach 2002; Nielsen 1999; Niromanesh 2005*). One study has been reported only in abstract - without randomisation details (Heard 2002). In one study, allocation was based on picking an un-numbered envelope from a pack - a method that is recognised to be less robust (Eng 1997*). In most trials, analysis by intention-to-treat was performed.

It was not possible to blind the physicians to the method of treatment in some studies - if this involved surgical evacuation of the uterus, alternative routes of drug administration (oral versus vaginal) or a policy of expectant management. It is, however, possible to blind the evaluator who assessed complications during the follow-up visit but no study made mention of this.

There was variation between studies in the timing of scheduled follow-up visits.

Effects of interventions

Twenty four studies, with a total of 1888 women, were included. Nineteen of the studies addressed termination of non-viable pregnancies before 14 weeks.

Treatment with vaginal misoprostol hastens miscarriage (passage of products of conception, whether complete or incomplete) when compared with placebo: miscarriage less than 24 hours (two trials, 138 women, relative risk (RR) 4.73, 95% confidence interval (CI) 2.70 to 8.28); miscarriage less than 48 hours (two (other) trials, 84 women, RR 5.74, 95% CI 2.70 to 12.19); complete miscarriage without need for surgical intervention at seven days (one trial, 83 women, RR 2.99, 95% CI 1.80 to 4.99). There was less need for uterine curettage (two trials, 104 women, RR 0.40, 95% CI 0.26 to 0.60) and no statistically significant increase in adverse effects: nausea (two trials, 88 women, RR 1.38, 95% CI 0.43 to 4.40), diarrhoea (two trials, 88 women, RR 2.21, 95% CI 0.35 to 14.06). One study showed a reduction in costs associated with a strategy of starting treatment with misoprostol, compared to immediate curettage (mean difference EUR192, 95% CI 33 to 351), no obvious difference in subsequent fertility, and similar numbers of women (58%) who would choose the same treatment strategy in the future (Graziosi 2004); although more women who had complete miscarriage after misoprostol (76%) would choose this treatment than those who required subsequent curettage (38%). Consistent with these observations, treatment with vaginal misoprostol decreases the need for surgical evacuation of the uterus when compared with a policy of arranging immediate surgical evacuation (three trials, 254 women, RR 0.42, 95% CI 0.34 to 0.52) at a cost of more nausea (one trial, 154 women, RR 21.85, 95% CI 1.31 to 364.37) and diarrhoea (one trial, 154 women, RR 40.85, 95% CI 2.52 to 662.57).

Vaginal misoprostol has been administered in doses of 400 mcg, 600 mcg, and 800 mcg in trials: lower-dose regimens tend to be less effective in producing miscarriage (three trials, 247 women, RR 0.85, 95% CI 0.72 to 1.00) with similar incidence of nausea (two trials, 214 women, RR 0.67, 95% CI 0.31 to 1.41). There seems no clear advantage to administering a 'wet' preparation of vaginal misoprostol compared to a 'dry' preparation: miscarriage less than three days (one trial, 80 women, RR 1.14, 95% CI 0.85 to 1.54). Adding methotrexate treatment to vaginal misoprostol has not been demonstrated to be advantageous in the single small trial to address this: miscarriage not complete after treatment (21 women, RR 0.26, 95% CI 0.01 to 5.65). Nor are laminaria tests proven useful adjuncts to vaginal misoprostol during the second trimester: complete miscarriage less than 24 hours (one trial, 38 women, RR 0.90, 95% CI 0.65 to 1.25), less than 48 hours (one trial, 38 women, RR 1.07, 95% CI 0.88 to 1.29). Vaginal misoprostol was more effective than vaginal prostaglandin E in avoiding surgical evacuation (one trial, 80 women, RR 0.39, 95% CI 0.21 to 0.72) and effecting complete miscarriage in the second trimester (one trial, 65 women, RR 1.44, 95% CI 1.06 to 1.96).

Overall, oral misoprostol was found to be less effective than vaginal misoprostol in producing complete miscarriage (two trials, 218

women, RR 0.90, 95% CI 0.82 to 0.99) but this difference was seen only with the 400 mcg oral dose (one trial, 20 women, RR 0.29, 95% CI 0.10 to 0.79) and not the 800 mcg oral dose (one trial, 198 women, RR 0.96, 95% CI 0.88 to 1.05). There was less vomiting with the oral regimen (one trial, 190 women, RR 0.29, 95% CI 0.10 to 0.84) but similar incidence of diarrhoea (two trials, 210 women, RR 1.05, 95% CI 0.67 to 1.66). There were high (and similar) levels of satisfaction with treatment (one trial, 198 women, RR 0.96, 95% CI 0.86 to 1.06). Oral misoprostol was slower than vaginal misoprostol in effecting miscarriage in a single trial of women with second trimester fetal death (weighted mean difference 4.10 hours, 95% CI 2.64 to 5.56).

Sublingual misoprostol had equivalent efficacy to vaginal misoprostol in inducing complete miscarriage (one trial, 80 women, RR 1.00, 95% CI 0.85 to 1.18) but was associated with more frequent diarrhoea (RR 2.65, 95% CI 1.48 to 4.38) but not other side-effects.

The two trials of mifepristone treatment generated conflicting results. One study found mifepristone to be much more effective than placebo: miscarriage complete by day five after treatment (46 women, RR 9.50, 95% CI 2.49 to 36.19). The other study compared treatment with mifepristone plus oral misoprostol with a policy of expectant management (no treatment); there was no statistically significant difference in complete miscarriage by day five (122 women, RR 1.08, 95% CI 0.90 to 1.30).

There was no statistically significant difference between vaginal misoprostol and gemeprost in the induction of miscarriage less than 24 hours for fetal death after 13 weeks (one trial, 50 women, RR 1.24, 95% CI 0.90 to 1.70).

There were few reports of serious adverse effects in the reported trials, but one woman required a bowel resection after uterine perforation at evacuation of the uterus (Egarter 1995).

DISCUSSION

The large majority of included trials (21/24) addressed the use of misoprostol (mainly by vaginal administration). There is intense interest in the reproductive uses of misoprostol because it appears a potent method for pregnancy interruption as well as being cheap and stable at room temperature, and thus potentially especially useful in developing countries, where transport and storage facilities are inadequate and the availability of uterotonic agents and blood is limited. However, ultrasound imaging is needed to di-

agnose non-viable pregnancies and equipment is sparse in many developing countries. The reproductive use of misoprostol is considered in other Cochrane reviews, for indications that include termination of unwanted pregnancies (Kulier 2004; Say 2002), induction of labour (Alfirevic 2001; Hofmeyr 2003; Muzonzini 2004) and prevention and treatment of postpartum haemorrhage (Gulmezoglu 2004; Mousa 2003).

AUTHORS' CONCLUSIONS

Implications for practice

Available evidence from randomised trials supports the use of vaginal misoprostol as one possible option for the treatment of non-viable pregnancies before 24 weeks.

Implications for research

Ultrasound demonstration of early pregnancy failure before 14 weeks is a common problem that merits greater research effort than has occurred to date. Further research is required to assess the effectiveness, safety and side-effects of misoprostol, including optimal route of administration and dose. Conflicting findings about the value of mifepristone need to be resolved by additional study. Women's views about the acceptability of medical treatment, surgical treatment and expectant management should be integral to future research protocols, as should economic assessments. Long-term outcome, notably subsequent fertility, deserves further study in appropriately powered randomised controlled studies.

[Note: the 108 citations in the awaiting classification section of the review may alter the conclusions of the review once assessed.]

ACKNOWLEDGEMENTS

Thanks to Sonja Henderson and Lynn Hampson of the Cochrane Pregnancy and Childbirth Group, Liverpool Women's Hospital, Liverpool, UK.

As part of the pre-publication editorial process, this review has been commented on by two peers (an editor and referee who are external to the editorial team), one or more members of the Pregnancy and Childbirth Group's international panel of consumers and the Group's Statistical Adviser.

REFERENCES

References to studies included in this review

Al Inizi 2003 *{published data only}*

Al Inizi SA, Ezimokhai M. Vaginal misoprostol versus dinoprostone for the management of missed abortion. *International Journal of Gynecology & Obstetrics* 2003;**83**: 73–4.

Autry 1999 *{published data only}*

Autry A, Jacobson G, Sandhu R, Isbill K. Medical management of non-viable early first trimester pregnancy. *International Journal of Gynecology & Obstetrics* 1999;**67**(1): 9–13.

Bagratee 2004 *{published data only}*

Bagratee JS, Khullar V, Regan L, Moodley J, Kagoro H. A randomized controlled trial comparing medical and expectant management of first trimester miscarriage. *Human Reproduction* 2004;**19**:266–71.

Creinin 1997 *{published data only}*

Creinin MD, Moyer R, Guido R. Misoprostol for medical evacuation of early pregnancy failure. *Obstetrics & Gynecology* 1997;**89**:768–72.

Demetroulis 2001 *{published data only}*

* Demetroulis C, Saridogan E, Kunde D, Naftalin AA. A prospective randomized control trial comparing medical and surgical treatment for early pregnancy failure. *Human Reproduction* 2001;**16**:365–9.
Demetroulis C, Saridogan E, Kunde D, Naftalin AA. A prospective randomized control trial comparing medical and surgical treatment for early pregnancy failure. XVI FIGO World Congress of Obstetrics and Gynecology; 2000 Sept 3–8; Washington DC, USA. 2000.

Egarter 1995 *{published data only}*

Egarter C, Lederhilger J, Kurz C, Karas H, Reisenberger K. Gemeprost for first trimester missed abortion. *Archives of Gynecology and Obstetrics* 1995;**256**:29–32.

Eng 1997* *{published data only}*

Eng NS, Guan AC. Comparative study of intravaginal misoprostol with gemeprost as an abortifacient in second trimester missed abortion. *Australian and New Zealand Journal of Obstetrics and Gynaecology* 1997;**37**:331–4.

Fadalla 2004* *{published data only}*

Fadalla FA, Mirghani OA, Adam I. Oral misoprostol vs vaginal misoprostol for termination of pregnancy with intrauterine fetal demise in the second-trimester. *International Journal of Gynecology & Obstetrics* 2004;**86**: 52–3.

Gilles 2004 *{published data only}*

Barnhart KT, Bader T, Huang X, Frederick MM, Timbers KA, Zhang JJ. Hormone pattern after misoprostol administration for a nonviable first-trimester gestation. *Fertility and Sterility* 2004;**81**:1099–105.
Creinin MD, Harwood B, Guido RS, Fox MC, Zhang J, NICHD Early Pregnancy Failure Trial. Endometrial thickness after misoprostol use for early pregnancy failure.

International Journal of Gynecology & Obstetrics 2004;**86**: 22–6.

Davis AR, Robilotto CM, Westhoff CL, Forman S, Zhang J, NICHD Management of Early Pregnancy Failure Trial. Bleeding patterns after vaginal misoprostol for treatment of early pregnancy failure. *Human Reproduction* 2004;**19**: 1655–8.

Gilles J, Creinin MM, Barnhart KT, Westhoff C, Frederick MM, Zhang J, et al. Wet versus dry misoprostol application for treatment of early pregnancy failure. *Fertility and Sterility* 2002;**78**:S64–S65.

* Gilles JM, Creinin MD, Barnhart K, Westhoff C, Frederick MM, Zhang J. A randomized trial of saline solution-moistened misoprostol versus dry misoprostol for the first-trimester pregnancy failure. *American Journal of Obstetrics and Gynecology* 2004;**190**:389–94.

Graziosi 2004 *{published data only}*

Graziosi GCM, Bruinse HW, Reuwer PJH, Teteringen O, Mol BWJ. Fertility outcome after a randomized trial comparing curettage with misoprostol for treatment of early pregnancy failure. *Human Reproduction* 2005;**20**:1749–50.
Graziosi GCM, Bruinse HW, Reuwer PJH, van Kessel PH, Westerweel PE, Mol BW. Misoprostol versus curettage in women with early pregnancy failure: impact on women's health-related quality of life. A randomized controlled trial. *Human Reproduction* 2005;**20**:2340–7.

* Graziosi GCM, Mol BWJ, Reuwer PJH, Drogtrap A, Bruinse HW. Misoprostol versus curettage in women with early pregnancy failure after initial expectant management: a randomized trial. *Human Reproduction* 2004;**19**:1894–9.
Graziosi GCM, van der Steeg JW, Reuwer PHW, Drogtrap AP, Bruinse HW, Mol BWJ. Economic evaluation of misoprostol in the treatment of early pregnancy failure compared to curettage after an expectant management. *Human Reproduction* 2005;**20**:1067–71.

Heard 2002 *{published data only}*

Heard MJ, Stewart GM, Buster JE, Carson SA, Miller HJ. Outpatient management of missed abortion with vaginal misoprostol [abstract]. *Obstetrics & Gynecology* 2002;**99**(4 Suppl):20S.

Herabutya 1997 *{published data only}*

Herabutya Y, O-Prasertsawat P. Misoprostol in the management of missed abortion. *International Journal of Gynecology & Obstetrics* 1997;**56**:263–6.

Jain 1996* *{published data only}*

Jain JK, Mishell DR. A comparison of misoprostol with and without laminaria tents for induction of second-trimester abortion. *American Journal of Obstetrics and Gynecology* 1996;**175**:173–7.

Kara 1999* *{published data only}*

Kara M, Ozden S, Eroglu M, Cetin A, Arioglu P. Comparison of misoprostol and dinoprostol administration for the induction of labour in second trimester pregnancies

- in cases of intrauterine fetal loss. *Italian Journal of Gynecology and Obstetrics* 1999;**1**:13–6.
- Kovavisarach 2002** *{published data only}*
Kovavisarach E, Sathapanachai U. Intravaginal 400 micrograms misoprostol for pregnancy termination in cases of blighted ovum: a randomised controlled trial. *Australian and New Zealand Journal of Obstetrics and Gynaecology* 2002;**42**:161–3.
- Kovavisarach 2005** *{published data only}*
Kovavisarach E, Jamnansiri C. Intravaginal misoprostol 600mcg and 800mcg for the treatment of early pregnancy failure. *International Journal of Gynecology & Obstetrics* 2005;**90**:208–12.
- Lelaidier 1993** *{published data only}*
Lelaidier C, Baton-Saint-Mleux C, Fernandez H, Bourget P, Frydman R. Mifepristone (RU 486) induces embryo expulsion in first trimester non-developing pregnancies: a prospective randomized trial. *Human Reproduction* 1993;**8**:492–5.
- Lister 2005** *{published data only}*
Lister MS, Shaffer LET, Bell JG, Lutter KQ, Moorma KH. Randomized, double-blind, placebo-controlled trial of vaginal misoprostol for management of early pregnancy failures. *American Journal of Obstetrics and Gynecology* 2005;**193**:1338–43.
- Muffley 2002** *{published data only}*
Muffley PE, Stitely ML, Gherman RB. Early intrauterine pregnancy failure: a randomized trial of medical versus surgical treatment. *American Journal of Obstetrics and Gynecology* 2002;**187**:321–6.
- Ngoc 2004** *{published data only}*
Ngoc NTN, Blum J, Westheimer E, Quan TTV, Winikoff B. Medical treatment of missed abortion using misoprostol. *International Journal of Gynecology & Obstetrics* 2004;**87**:138–42.
- Nielsen 1999** *{published data only}*
Nielsen S, Hahlin M, Platz-Christensen J. Expectant management or pharmacological treatment for first trimester spontaneous abortion: a randomised trial. *Acta Obstetrica et Gynecologica Scandinavica* 1997;**76**(167:2):77.
* Nielsen S, Hahlin M, Platz-Christensen J. Randomised trial comparing expectant with medical management for first trimester miscarriages. *British Journal of Obstetrics and Gynaecology* 1999;**106**:804–7.
- Niromanesh 2005*** *{published data only}*
Niromanesh S, Hashemi-Feasharaki M, Mosavi-Jarrahi A. Second trimester abortion using intravaginal misoprostol. *International Journal of Gynecology & Obstetrics* 2005;**89**:276–7.
- Tang 2003** *{published data only}*
Tang OS, Lau WNT, Ng EHY, Lee SWH, Ho PC. A prospective randomized study to compare the use of repeated doses of vaginal with sublingual misoprostol in the management of first trimester silent miscarriage. *Human Reproduction* 2003;**18**:176–81.
- Wood 2002** *{published data only}*
Wood SL, Brain PH. Medical management of missed abortion: a randomized clinical trial. *Obstetrics & Gynecology* 2002;**99**:563–6.

References to studies excluded from this review

- Abdel Fattah 1997** *{published data only}*
Abdel Fattah IH. PGE1 analogue for the induction of midtrimester abortion in cases of intrauterine fetal death. *Acta Obstetrica et Gynecologica Scandinavica Supplement* 1997;**76**(167:2):26.
- Almog 2005** *{published data only}*
Almog B, Levin I, Winkler N, Fainaru O, Pauzner D, Lessing JB, et al. The contribution of laminaria placement for cervical ripening in second trimester termination of pregnancy induced by intra-amniotic injection of prostaglandin F2alpha followed by concentrated oxytocin infusion. *European Journal of Obstetrics & Gynecology and Reproductive Biology* 2005;**118**:32–5.
- Anderman 2000** *{published data only}*
Anderman S, Jaschevatzky OE, Ballas S. Comparison between a double balloon device and the foley catheter in extraamniotic prostaglandin F2a infusion for termination of midtrimester missed abortion. XVI FIGO World Congress of Obstetrics & Gynecology; 2000 Sept 3-8; Washington DC, USA. 2000:162.
- Avila-Vergara 1997** *{published data only}*
Avila-Vergara MA, Morgan-Ortiz F, Fragoza-Sosa O, Haro-Garcia L. Cervical labor induction with prostaglandin E2 in patients with fetal death [Maduracion cervical con prostaglandina E2 en pacientes con feto muerto]. *Ginecologia y Obstetricia de Mexico* 1997;**65**:155–8.
- Bebbington 2002** *{published data only}*
Bebbington MW, Kent N, Lim K, Gagnon A, Delisle MF, Tessier F, et al. A randomized controlled trial comparing two protocols for the use of misoprostol in midtrimester pregnancy termination. *American Journal of Obstetrics and Gynecology* 2002;**187**:853–7.
- Cabrol 1990** *{published data only}*
Cabrol D, Dubois C, Cronje H, Gonnet JM, Guillot M, Maria B, et al. Induction of labour with mifepristone (RU 486) in intrauterine fetal death. *American Journal of Obstetrics and Gynecology* 1990;**163**:540–2.
- Clevin 2001** *{published data only}*
Clevin L, Munk T, Hansen TR. Spontaneous abortion. Drug treatment versus surgery [Spontan abort. Medicinsk versus kirurgisk behandling]. *Ugeskrift for Laeger* 2001;**163**(15):2136–9.
- David 2003** *{published data only}*
David M, Chen FCK, Lichtenegger W. NO-donor nitroglycerin versus the prostaglandin gemeprost for cervical ripening in first trimester missed abortion. *International Journal of Gynecology & Obstetrics* 2003;**83**:71–2.
- Dickinson 1998** *{published data only}*
Dickinson JE, Godfrey M, Evans SF. Efficacy of intravaginal misoprostol in second-trimester pregnancy termination:

- a randomized controlled trial. *Journal of Maternal-Fetal Medicine* 1998;**7**:115–9.
- Dickinson 2002** *{published data only}*
Dickinson JE, Evans SF. The optimization of intravaginal misoprostol dosing schedules in second-trimester pregnancy termination. *American Journal of Obstetrics and Gynecology* 2002;**186**:470–4.
- Dickinson 2003** *{published data only}*
Dickinson JE, Evans SF. A comparison of oral misoprostol with vaginal misoprostol administration in second-trimester pregnancy termination for fetal abnormality. *Obstetrics & Gynecology* 2003;**101**:1294–9.
- Eppel 2005** *{published data only}*
Eppel W, Facchinetti F, Schleussner E, Piccinini F, Pizzi C, Gruber DM, et al. Second trimester abortion using isosorbide mononitrate in addition to gemeprost compared with gemeprost alone: a double blind randomized, placebo-controlled multicenter trial. *American Journal of Obstetrics and Gynecology* 2005;**192**:856–61.
- Feldman 2003** *{published data only}*
Feldman DM, Borgida AF, Rodis JF, Leo MV, Campbell WA. A randomized comparison of two regimens of misoprostol for second-trimester pregnancy termination. *American Journal of Obstetrics and Gynecology* 2003;**189**:710–3.
- Ghorab 1998** *{published data only}*
Ghorab MNM, El Helw BA. Second-trimester termination of pregnancy by extra-amniotic prostaglandin F2alpha or endocervical misoprostol. *Acta Obstetrica et Gynecologica Scandinavica* 1998;**77**:429–32.
- Gonzalez 2001** *{published data only}*
Gonzalez JA, Carlan SJ, Alverson MW. Outpatient second trimester pregnancy termination. *Contraception* 2001;**63**:89–93.
- Grimes 2004** *{published data only}*
Grimes DA, Smith MS, Witham AD. Mifepristone and misoprostol versus dilatation and evacuation for midtrimester abortion: a pilot randomised controlled trial. *BJOG: an international journal of obstetrics and gynaecology* 2004;**111**:148–53.
- Gronland 2002** *{published data only}*
Gronland A, Gronland L, Clevin L, Andersen B, Palmegren N, Lidegaard O. Management of missed abortion: comparison of medical treatment with either mifepristone + misoprostol or misoprostol alone with surgical evacuation. A multi-center trial in Copenhagen county, Denmark. *Acta Obstetrica et Gynecologica Scandinavica* 2002;**81**:1060–5.
- Hausler 1997** *{published data only}*
Hausler MCH, Koroschetz F, Tamussino K, Walcher W. Is a curettage after spontaneous abortion still relevant. A prospective randomised study [Ist eine Curettage nach Abortus completus noch zeitgemäß?. Eine prospektiv randomisierte Studie]. *Geburthshilfe und Frauenheilkunde* 1997;**57**:396–9.
- Herabutya 2005** *{published data only}*
Herabutya Y, Chanrachakul B, Punyavachira P. A randomised controlled trial of 6 and 12 hourly administration of vaginal misoprostol for second trimester pregnancy termination. *BJOG: an international journal of obstetrics and gynaecology* 2005;**112**:1297–301.
- Hidar 2001** *{published data only}*
Hidar S, Fekih M, Chaieb A, Bibi M, Mellouli R, Khairi H. Oxytocin and misoprostol administered intravaginally for termination of pregnancy at 13–29 weeks of amenorrhea. A prospective randomized trial [Apport de l'association d'ocytocine au misoprostol administre en intravaginal au cours des interruptions de grossesses entre 13 et 29 semaines d'amenorrhée. Essai clinique prospectif randomisé]. *Journal de Gynecologie, Obstetrique et Biologie de la Reproduction* 2001;**30**:439–43.
- Hill 1991** *{published data only}*
Hill NCW, Selinger M, Ferguson J, MacKenzie IZ. Management of intra-uterine fetal death with vaginal administration of gemeprost or prostaglandin E2: a random allocation controlled trial. *Journal of Obstetrics and Gynaecology* 1991;**11**:422–6.
- Hogg 2000** *{published data only}*
Hogg B, Owen J. Laminaria versus extraamniotic saline infusion (EASI) for cervical ripening and mid-trimester labor induction. *American Journal of Obstetrics and Gynecology* 2000;**182**(1 Pt 2):S135.
- Jain 1994** *{published data only}*
Jain JK, Mishell DR. A comparison of intravaginal misoprostol with prostaglandin E2 for termination of second-trimester pregnancy. *New England Journal of Medicine* 1994;**331**:290–3.
- Jain 1999** *{published data only}*
Jain JK, Kuo J, Mishell DR. A comparison of two dosing regimens of intravaginal misoprostol for second-trimester pregnancy termination. *Obstetrics & Gynecology* 1999;**93**:571–5.
- Johnson 1997** *{published data only}*
Johnson N, Priestnall M, Marsay T, Ballard P, Watters J. A randomised trial evaluating pain and bleeding after a first trimester miscarriage treated surgically or medically. *European Journal of Obstetrics & Gynecology and Reproductive Biology* 1997;**72**(2):213–5.
- Kanhai 1989** *{published data only}*
* Kanhai HHH, Keirse MJNC. Induction of labour after fetal death: a randomized controlled trial of two prostaglandin regimens. *British Journal of Obstetrics and Gynaecology* 1989;**96**:1400–4.
Kanhai HHH, Keirse MJNC. Intravenous administration of sulprostone for the induction of labour after fetal death: a randomised comparison of two dose schedules. Proceedings of 1st European Congress on Prostaglandins in Reproduction; 1988 July 6–9; Vienna, Austria. 1988:45.
Kanhai HHH, Keirse MJNC. Intravenous administration of sulprostone for the induction of labour after fetal death: a randomised comparison of two dose schedules. *World*

- Congress of Gynecology and Obstetrics; 1988 October 23-28; Brazil. 1988:201-2.
- Lippert 1978** *{published data only}*
Lippert TH, Luthi A. Induction of labour with prostaglandin E2 gel in cases of intrauterine fetal death. *Prostaglandins* 1978;**15**:533-42.
- Machtinger 2002** *{published data only}*
Machtinger R, Stockheim D, Goldenberg M, Soriano D, Atlas M, Seidman DS. A randomized prospective study of misoprostol alone or combined with mifepristone for treatment of first trimester spontaneous abortion. *Fertility and Sterility* 2002;**78**(3 Suppl 1):S64.
- Machtinger 2004** *{published data only}*
Machtinger R, Stockheim D, Shulman A, Dulitzki M, Schiff E, Seidman DS. A randomized prospective study comparing the effectiveness of four protocols for treatment of first trimester spontaneous abortion. *Fertility and Sterility* 2004;**82** Suppl 2:S80.
- Makhlouf 2003** *{published data only}*
Makhlouf AM, Al-Hussaini TK, Habib DM, Makarem MH. Second-trimester pregnancy termination: comparison of three different methods. *Journal of Obstetrics and Gynaecology* 2003;**23**:407-11.
- Martin 1965** *{published data only}*
Martin RH, Menzies DN. Oestrogen therapy in missed abortion and labour. *Journal of Obstetrics and Gynaecology of the British Commonwealth* 1965;**62**:256-8.
- Nakintu 2001** *{published data only}*
Nakintu N. A comparative study of vaginal misoprostol and intravenous oxytocin for induction of labour in women with intrauterine fetal death in Mulago Hospital, Uganda. *African Health Sciences* 2001;**1**:55-9.
- Ngai 2001** *{published data only}*
Ngai SW, Chan YM, Tang OS, Ho PC. Vaginal misoprostol as medical treatment for first trimester spontaneous miscarriage. *Human Reproduction* 2001;**16**(7):1493-6.
- Nuthalapaty 2004** *{published data only}*
Nuthalapaty F, Ramsey P, Biggio J, Owen J. Comparative efficacy of high dose vaginal misoprostol versus concentrated oxytocin + low dose vaginal misoprostol for mid-trimester labor induction. *American Journal of Obstetrics and Gynecology* 2004;**191**(6 Suppl 1):S73.
- Nuutila 1997** *{published data only}*
Nuutila M, Toivonen J, Ylikorkala O, Halmesmaki E. A comparison between two doses of intravaginal misoprostol and gemeprost for induction of second-trimester abortion. *Obstetrics & Gynecology* 1997;**90**:896-900.
- Owen 1999** *{published data only}*
Owen J, Hauth JC. Vaginal misoprostol vs concentrated oxytocin plus low-dose prostaglandin E2 for second trimester pregnancy termination. *Journal of Maternal-Fetal Medicine* 1999;**8**:48-50.
- Paraskevaides 1992** *{published data only}*
Paraskevaides E, Prendiville W, Stuart B, Scanail SN, Walsh D, McGuinness N, et al. Medical evacuation of first trimester (twelve weeks gestation) incomplete abortion and missed abortion. *Journal of Gynecologic Surgery* 1992;**8**:159-63.
- Perry 1999** *{published data only}*
Perry KG, Rinehart BK, Terrone DA, Martin RW, May WL, Roberts WE. Second-trimester uterine evacuation: a comparison of intra-amniotic (15S)-15-methyl-prostaglandin F2alpha and intravaginal misoprostol. *American Journal of Obstetrics and Gynecology* 1999;**181**:1057-61.
- Piotrowski 1979** *{published data only}*
Piotrowski J, Basta A, Klimczyk K, Malolepazy A, Dluzniewska M, Splawinski JA. Indomethacin increases abortifacient effect of PGE2 in man. *Prostaglandins* 1979;**17**:451-9.
- Pongsatha 2004** *{published data only}*
Pongsatha S, Tongsong T. Intravaginal misoprostol for pregnancy termination. *International Journal of Gynecology & Obstetrics* 2004;**87**:176-7.
- Ramsey 2004** *{published data only}*
Ramsey PS, Savage K, Lincoln T, Owen J. Vaginal misoprostol versus concentrated oxytocin and vaginal PGE2 for second-trimester labor induction. *Obstetrics & Gynecology* 2004;**104**:138-45.
- Roy 2003** *{published data only}*
Roy G, Ferreira E, Hudon L, Marquette G. The efficacy of oral versus vaginal misoprostol for second-trimester termination of pregnancy: a double blind, randomized placebo-controlled trial. *American Journal of Obstetrics and Gynecology* 2003;**189**(6):S70.
- Salamalekis 1990** *{published data only}*
Salamalekis E, Loghis C, Kassanos D, Traka A, Zourlas PA. Comparison of extra-amniotic prostaglandin F2alpha and dinoprostone use for labor induction after second trimester intrauterine fetal death. Proceedings of 12th European Congress of Perinatal Medicine; 1990; Lyon, France. 1990:228.
- Su 2005** *{published data only}*
Su LL, Biswas A, Choolani M, Kalaichelvan V, Singh K. A prospective randomized comparison of vaginal misoprostol versus intra-amniotic prostaglandins for midtrimester termination of pregnancy. *American Journal of Obstetrics and Gynecology* 2005;**193**:1410-4.
- Surita 1997** *{published data only}*
Surita FGC, Cecatti JG, Pinto e Silva JL. Misoprostol versus laminaria for cervical ripening in intrauterine fetal death. *Acta Obstetrica et Gynecologica Scandinavica Supplement* 1997;**76**(167:2):32.
- Thavarasah 1986** *{published data only}*
Thavarasah AS, Almohdzar SA. Prostaglandin (F2alpha) in missed abortion. Intravenous extra-amniotic and intramuscular administration - a randomized study. *Biological Research in Pregnancy* 1986;**7**:106-10.
- Toppozada 1994** *{published data only}*
Toppozada MK, Shaala SA, Anwar MY, Haiba NA, Abdrabbo S, El-Absy HM. Termination of pregnancy with

fetal death in the second and third trimesters - the double balloon versus extra-amniotic prostaglandin. *International Journal of Gynecology & Obstetrics* 1994;**45**:269–73.

Yapar 1996 {published data only}

Yapar EG, Senoz S, Urkutur M, Batioglu S, Gokmen O. Second trimester pregnancy termination including fetal death: comparison of five different methods. *European Journal of Obstetrics & Gynecology and Reproductive Biology* 1996;**69**:97–102.

Zhang 2005 {published data only}

* Zhang J, Gilles JM, Barnhart K, Creinin MD, Westhoff C, Frederick MM, et al. A comparison of medical management with misoprostol and surgical management for early pregnancy failure. *New England Journal of Medicine* 2005; **353**:761–9.
Zhang J, Gilles K, Barnhart K, Creinin M, Westhoff C, Frederick M. Medical management with misoprostol for early pregnancy failure: a multicenter, randomized equivalence trial. *Fertility and Sterility* 2004;**82** Suppl 2: S53–S54.

References to studies awaiting assessment

Aberdeen 1993 {published data only}

Henshaw RC, Hinshaw K, Smith NC, Templeton AA. The medical management of miscarriage. Fertility Society of Australia / Australian Gynaecological Endoscopy Society; 1995 November 19–25; Melbourne, Australia. 1995:FSA 75.
* Hinshaw K, Rispin N, Smith N, Templeton A. Medical versus surgical management in first trimester miscarriage: a prospective, pragmatic random allocation trial. *Journal of Obstetrics and Gynaecology* 1993;**13**:404–5.
Hinshaw K, Rispin R, Henshaw R, Smith N, Templeton A. Medical versus surgical uterine evacuation in first trimester miscarriage: a prospective, pragmatic randomised trial. 27th British Congress of Obstetrics and Gynaecology; 1995 July 4–7; Dublin. 1995:4.
Hughes J, Ryan M, Hinshaw K, Henshaw R, Rispin R, Templeton A. The costs of treating miscarriage: a comparison of medical and surgical management. *British Journal of Obstetrics and Gynaecology* 1996;**103**:1217–21.
Rispin R, Hinshaw K, Henshaw R, Smith N, Templeton A. New aspects of care in the management of miscarriage. Proceedings of Research in Midwifery Conference; 1993 September 14; Birmingham, UK. 1993.

Al-Bdour 2007 {published data only}

Al-Bdour AN, Akasheh H, Al-Jayousi T. Missed abortion: termination using single-dose versus two doses of vaginal misoprostol tablets. *Pakistan Journal of Medical Sciences* 2007;**23**(6):920–3.

Altaf 2006 {published data only}

Altaf F, Sultana N, Iqbal N. Therapeutic abortions; efficacy of intra-vaginal misoprostol in comparison to extra amniotically administered prostaglandin f2a. *Professional Medical Journal* 2006;**13**(3):417–22.

Anderson 2009 {published data only}

Anderson J, Gouk E, Young L, Turnbull L, Sayeed G, Elattar A, et al. A randomised controlled trial of oral versus vaginal misoprostol for medical management of early fetal demise. *International Journal of Gynecology & Obstetrics* 2009;**107**(Suppl 2):S533.

Ara 2009 {published data only}

Ara G, Nargis S, Khatun R, Saha A. Vaginal misoprostol as a medical management in early pregnancy loss. *International Journal of Gynecology & Obstetrics* 2009;**107**(Suppl 2): S533–4.

Arellano 2009 {published data only}

Arellano M, Durocher J, Leon W, Montesinos R, Pena M, Winikoff B. Introduction of misoprostol for incomplete abortion care in Latin America: evidence from Ecuador. *International Journal of Gynecology & Obstetrics* 2009;**107** (Suppl 2):S49.

Ayudhaya 2006 {published data only}

Ayudhaya OP, Herabutya Y, Chanrachakul B, Ayuthaya NI, O-Prasertsawat P. A comparison of the efficacy of sublingual and oral misoprostol 400 microgram in the management of early pregnancy failure: a randomized controlled trial. *Journal of the Medical Association of Thailand* 2006;**89** (Suppl 4):S5–S10.

Azra 2007 {published data only}

Azra B, Shakeel S, Nilofer M. A comparison of two protocols of intra vaginal misoprostol for second trimester medical termination of pregnancy. *Pakistan Armed Forces Medical Journal* 2007;**57**(1):61–5.

Bagratee 2009 {published data only}

Bagratee J, Regan L, Khullar V, Moodley J, Connolly C. Does the volume of retained products of conception and hormonal parameters influence the success of conservative methods of management of first trimester miscarriage?. *International Journal of Gynecology & Obstetrics* 2009;**107** (Suppl 2):S116.

Bangkok 2003 {published data only}

Chittacharoen A, Herabutya Y, Punyavachira P. A randomized trial of oral and vaginal misoprostol to manage delivery in cases of fetal death. *Obstetrics & Gynecology* 2003;**101**:70–3.

Bani-Irshaid 2006 {published data only}

Bani-Irshaid I, Athamneh TZ, Bani-Khaled D, Al-Momani M, Dahamsheh H. Termination of second and early third trimester pregnancy: comparison of 3 methods. *Eastern Mediterranean Health Journal* 2006;**12**(5):605–9.

Behrashi 2008 {published data only}

Behrashi M, Mahdian M. Vaginal versus oral misoprostol for second-trimester pregnancy termination: a randomized trial. *Pakistan Journal of Biological Sciences* 2008;**11**(21): 2505–8.

Behrashi 2010 {published data only}

Behrashi M. Comparison between the oral and vaginal misoprostol effects on pregnancy termination in second trimester. IRCT Iranian Registry of Clinical Trials (www.irct.ir) (accessed 6 December 2010).

Ben-Meir 2009 {published data only}

Ben-Meir A, Erez Y, Feigenberg T, Hamani Y, Laufer N, Rojansky N. Mifepristone followed by high-dose oxytocin drip for second-trimester abortion: a randomized, double-blind, placebo-controlled, pilot study. *Journal of Reproductive Medicine* 2009;**54**(8):511–6.

Betstadt 2008 {published data only}

Betstadt S. MiMi: a randomized trial of mifepristone and misoprostol for treatment of early pregnancy failure. <http://clinicaltrials.gov/ct2/show/NCT00468299> (accessed 20 February 2008).

Bique 2007 {published data only}

Bique C, Usta M, Debora B, Chong E, Westheimer E, Winikoff B. Comparison of misoprostol and manual vacuum aspiration for the treatment of incomplete abortion. *International Journal of Gynecology & Obstetrics* 2007;**98**(3): 222–6.

Biswas 2007 {published data only}

Biswas SC, Dey R, Jana R, Chattopadhyay N. Comparative study of intravaginal misoprostol and extra amniotic ethacridine lactate instillation for mid trimester pregnancy termination. *Journal of Obstetrics and Gynaecology of India* 2007;**57**(3):211–3.

Blohm 2005 {published data only}

Blohm F, Friden BE, Milsom I, Platz-Christensen JJ, Nielsen S. A randomised double blind trial comparing misoprostol or placebo in the management of early miscarriage. *BJOG: an international journal of obstetrics and gynaecology* 2005; **112**(8):1090–5.

Brouns 2010 {published data only}

Brouns JF, van Wely M, Burger MP, van Wijngaarden WJ. Comparison of two dose regimens of misoprostol for second-trimester pregnancy termination. *Contraception* 2010;**82**(3):266–75.

Cabrol 1990a {published data only}

Cabrol D, Dubois C, Cronje H, Gonnet JM, Guillot M, Maria B, et al. Induction of labor with mifepristone (RU 486) in intrauterine fetal death. *American Journal of Obstetrics and Gynecology* 1990;**163**:540–2.

Caliskan 2005 {published data only}

Caliskan E, Dilbaz S, Doger E, Ozeren S, Dilbaz B. Randomized comparison of 3 misoprostol protocols for abortion induction at 13–20 weeks of gestation. *Journal of Reproductive Medicine* 2005;**50**(3):173–80.

Caliskan 2009 {published data only}

Caliskan E, Doger E, Cakiroglu Y, Corakci A, Yucesoy I. Sublingual misoprostol 100 microgram versus 200 microgram for second trimester abortion: a randomised trial. *European Journal of Contraception & Reproductive Health Care* 2009;**14**(1):55–60.

Chen 2008 {published data only}

Chen BA, Reeves MF, Creinin MD, Gilles JM, Barnhart K, Westhoff C, et al. Misoprostol for treatment of early pregnancy failure in women with previous uterine surgery. *American Journal of Obstetrics & Gynecology* 2008;**198**(6): 626.e1–5.

Creinin 2006 {published data only}

Creinin MD, Huang X, Westhoff C, Barnhart K, Gilles JM, Zhang J, et al. Factors related to successful misoprostol treatment for early pregnancy failure. *Obstetrics & Gynecology* 2006;**107**(4):901–7.

Dabash 2009 {published data only}

Dabash R, Cherine M, Darwish E, Blum J, Hassanein N, Abdel Daiem T, et al. Bleeding following surgical (MVA) and medical (400 ug sublingual misoprostol) treatment of incomplete abortion. *International Journal of Gynecology & Obstetrics* 2009;**107**(Suppl 2):S150–1.

Dao 2007 {published data only}

Dao B, Blum J, Thieba B, Raghavan S, Ouedraogo M, Lankoande J, et al. Is misoprostol a safe, effective and acceptable alternative to manual vacuum aspiration for postabortion care? Results from a randomised trial in Burkina Faso, West Africa. *BJOG: an international journal of obstetrics and gynaecology* 2007;**114**(11):1368–75.

David 2005 {published data only}

David M, Chen FCK. Comparison of isosorbide mononitrate (Mono Mack) and misoprostol (Cytotec) for cervical ripening in the first trimester missed abortion. *Archives of Gynecology and Obstetrics* 2005;**273**(3):144–5.

Davis 2007 {published data only}

Davis AR, Hendlish SK, Westhoff C, Frederick MM, Zhang J, Gilles JM, et al. Bleeding patterns after misoprostol vs surgical treatment of early pregnancy failure: results from a randomized trial. *American Journal of Obstetrics and Gynecology* 2007;**196**(1):31.

Diop 2009 {published data only}

Diop A, Raghavan S, Rakotovo JP, Comendant R, Blumenthal PD, Winikoff B. Two routes of administration for misoprostol in the treatment of incomplete abortion: a randomized clinical trial. *Contraception* 2009;**79**(6): 456–62.

Elhassan 2008 {published data only}

Elhassan EM, Abubaker MS, Adam I. Sublingual compared with oral and vaginal misoprostol for termination of pregnancy with second-trimester fetal demise. *International Journal of Gynecology & Obstetrics* 2008;**100**(1):82–3.

Eslamian 2007 {published data only}

Eslamian L, Gosili R, Jamal A, Alyassin A. A prospective randomized controlled trial of two regimens of vaginal misoprostol in second trimester termination of pregnancy. *Acta Medica Iranica* 2007;**45**(6):497–500.

Fang 2009 {published data only}

Fang AH, Chen QF, Zheng W, Li YH, Chen RY. Termination of missed abortion in a combined procedure: a randomized controlled trial. *Journal of Reproduction and Contraception* 2009;**20**(1):45–9.

Fiala 2005 {published data only}

Fiala C, Swahn ML, Stephansson O, Gemzell-Danielsson K. The effect of non-steroidal anti-inflammatory drugs on medical abortion with mifepristone and misoprostol at 13–

- 22 weeks gestation. *Human Reproduction* 2005;**20**(11): 3072–7.
- Guix 2005** {published data only}
Guix C, Palacio M, Figueras F, Bennasar M, Zamora L, Coll O, et al. Efficacy of two regimens of misoprostol for early second-trimester pregnancy termination. *Fetal Diagnosis and Therapy* 2005;**20**(6):544–8.
- Halimi 2004** {published data only}
Halimi M. Therapeutic termination of second trimester pregnancy: a comparison of extra-amniotic foley's catheter balloon alone with the combined use of foley's catheter balloon and extra-amniotic instillation of prostaglandin f2-alpha. *Journal of Postgraduate Medical Institute* 2004;**18**(3): 408–18.
- Harwood 2008** {published data only}
Harwood B, Nansel T, for the NIOFCHandHDMofEPFT. Quality of life and acceptability of medical versus surgical management of early pregnancy failure. *BJOG: an international journal of obstetrics and gynaecology* 2008;**115**(4):501–8.
- Hassan 2007** {published data only}
Hassan FI, Mostapha MK, Sattar MA, Marouf E, Azim SA. Oral versus rectal route of misoprostol administration: a randomized controlled trial. *Middle East Fertility Society Journal* 2007;**12**(1):53–6.
- Herabutya 1997a** {published data only}
Herabutya Y, O-Prasertsawat P. A comparison of intravaginal misoprostol with intracervical prostaglandin E2 gel for the management of dead fetus in utero. *Thai Journal of Obstetrics and Gynaecology* 1997;**9**(2):95–8.
- Hidar 2005** {published data only}
Hidar S, Bouddebous M, Chaieb A, Jerbi M, Bibi M, Khairi H. Randomized controlled trial of vaginal misoprostol versus vaginal misoprostol and isosorbide dinitrate for termination of pregnancy at 13–29 weeks. *Archives of Gynecology and Obstetrics* 2005;**273**(3):157–60.
- Hughes 1996** {published data only}
Hughes J, Ryan M, Hinshaw K, Henshaw R, Rispin R, Templeton A. The costs of treating miscarriage: a comparison of medical and surgical management. *British Journal of Obstetrics and Gynaecology* 1996;**103**(12): 1217–21.
- Imran 2010** {published data only}
Imran F, Anser A, Danish N, Fatima N. Misoprostol for the purpose of mid-trimester termination of pregnancy: a comparative study with prostaglandin F2 alpha. *Journal of Ayub Medical College, Abbottabad: JAMC* 2010;**22**(4): 87–91.
- Islam 2006** {published data only}
Islam A, Abbasi AN, Sarwar I. Use of Foley's catheter and prostaglandin F-2 alpha in second trimester termination of pregnancy. *Journal of Ayub Medical College, Abbottabad* 2006;**18**(3):35–9.
- Jabir 2009** {published data only}
Jabir M, Smeets R. Comparison of oral and vaginal misoprostol for cervical ripening before evacuation of first trimester missed miscarriage. *International Journal of Gynecology & Obstetrics* 2009;**107**(Suppl 2):S209.
- Jabir 2009a** {published data only}
Jabir M, Smeets RI. Comparison of oral and vaginal misoprostol for cervical ripening before evacuation of first trimester missed miscarriage. *Saudi Medical Journal* 2009;**30**(1):82–7.
- Kamal 2005** {published data only}
Kamal R, Parveen F, Mazhar SB. Role of misoprostol in vaginal versus double oro-vaginal route for termination of pregnancy in mid trimester pregnancy. *Annals of Pakistan Institute of Medical Sciences* 2005;**1**(4):196–200.
- Kapp 2007** {published data only}
Kapp N, Todd CS, Yadgarova KT, Alibayeva G, Nazarova D, Loza O, et al. A randomized comparison of misoprostol to intrauterine instillation of hypertonic saline plus a prostaglandin F2alpha analogue for second-trimester induction termination in Uzbekistan. *Contraception* 2007;**76**(6):461–6.
- Kurshid 2010** {published data only}
Kurshid R, Ahmed A, Mir S, Ul Shamas I. To assess the efficacy of two regimens of misoprostol for second trimester pregnancy termination—a randomized comparison. Internet Journal of Gynecology and Obstetrics 2010; Vol. 14, issue 1.
- Kushwah 2009** {published data only}
Kushwah B, Singh A. Sublingual versus oral misoprostol for uterine evacuation following early pregnancy failure. *International Journal of Gynecology & Obstetrics* 2009;**106**(1):43–5.
- Kushwah 2011** {published data only}
Kushwah DS, Kushwah B, Salman MT, Verma VK. Acceptability and safety profile of oral and sublingual misoprostol for uterine evacuation following early fetal demise. *Indian Journal of Pharmacology* 2011;**43**(3): 306–10.
- Lahore 1999** {published data only}
Amjad T, Akhtar S. Termination of pregnancy with foetal death in second trimester: Foley's catheter versus extra amniotic prostaglandins. *Journal of College of Physicians & Surgeons Pakistan* 1999;**9**:403–5.
- Lughmani 2007** {published data only}
Lughmani ST. A comparison of intravaginal misoprostol with prostaglandin E2 for termination of 1st trimester pregnancy. Double blind randomized trial [abstract]. 31st British International Congress of Obstetrics and Gynaecology; 2007 July 4–6; London, UK. 2007:209.
- Lughmani 2008** {published data only}
Lughmani ST. A comparison of intravaginal misoprostol with prostaglandin E2 for termination of 1st trimester pregnancy. *BJOG: an international journal of obstetrics and gynaecology* 2008;**115**(s1):179.
- Mahjabeen, 2009** {published data only}
Mahjabeen, Khawaja NP, Rehman R. Comparison of oral versus vaginal misoprostol for mid-trimester pregnancy

- termination. *Journal of the College of Physicians & Surgeons Pakistan* 2009;**19**(6):359–62.
- Montesinos 2011** {published data only}
Montesinos R, Durocher J, Leon W, Arellano M, Pena M, Pinto E, et al. Oral misoprostol for the management of incomplete abortion in Ecuador. *International Journal of Gynecology & Obstetrics* 2011;**115**(2):135–9.
- Moran 2005** {published data only}
Moran T, Deutsch R. Methotrexate/misoprostol vs. a more standard approach for termination of pregnancies of undetermined location: a randomized, controlled trial. *Journal of Reproductive Medicine* 2005;**50**(10):784–92.
- Mostafa-Gharebaghi 2010** {published data only}
Mostafa-Gharebaghi P, Mansourfar M, Sadeghi-Bazargani H. Low dose vaginal misoprostol versus prostaglandin E2 suppository for early uterine evacuation: a randomized clinical trial. *Pakistan Journal of Biological Sciences* 2010;**13**(19):946–50.
- Mulayim 2009** {published data only}
Mulayim B, Celik NY, Onalan G, Zeyneloglu HB, Kescu E. Sublingual misoprostol after surgical management of early termination of pregnancy. *Fertility & Sterility* 2009;**92**(2):678–81.
- Nasreen 2009** {published data only}
Nasreen Z. What for early pregnancy failure manual vacuum aspiration (MVA) with small dose misoprostol or misoprostol alone?. *International Journal of Gynecology & Obstetrics* 2009;**107**(Suppl 2):S539–40.
- Nassar 2006** {published data only}
Nassar AH. A randomized trial of two regimens of misoprostol for second trimester termination for intrauterine fetal death (ongoing trial). *ClinicalTrials.gov* (<http://clinicaltrials.gov/>) (accessed 21 March 2006) 2006.
- Nguyen 2005** {published data only}
Nguyen TN, Blum J, Durocher J, Quan TT, Winikoff B. A randomized controlled study comparing 600 versus 1,200 micrograms oral misoprostol for medical management of incomplete abortion. *Contraception* 2005;**72**(6):438–42.
- Niinimäki 2006** {published data only}
Niinimäki M, Jouppila P, Martikainen H, Talvensaari-Mattila A. A randomized study comparing efficacy and patient satisfaction in medical or surgical treatment of miscarriage. *Fertility and Sterility* 2006;**86**(2):367–72.
- Niinimäki 2009** {published data only}
Niinimäki M, Karinen P, Hartikainen AL, Pouta A. Treating miscarriages: a randomised study of cost-effectiveness in medical or surgical choice. *BJOG: an international journal of obstetrics and gynaecology* 2009;**116**(7):984–90.
- Nor 2006** {published data only}
Nor Azlin MI, Abdullah HS, Zainul Rashid MR, Jamil MA. Misoprostol (alone) in second trimester terminations of pregnancy: as effective as Gemeprost?. *Journal of Obstetrics and Gynaecology* 2006;**26**(6):546–9.
- Nuthalapaty 2005** {published data only}
Nuthalapaty FS, Ramsey PS, Biggio JR, Owen J. High-dose vaginal misoprostol versus concentrated oxytocin plus low-dose vaginal misoprostol for midtrimester labor induction: a randomized trial. *American Journal of Obstetrics and Gynecology* 2005;**193**(3 Pt 2):1065–70.
- Paritakul 2010** {published data only}
Paritakul P, Phupong V. Comparative study between oral and sublingual 600 microg misoprostol for the treatment of incomplete abortion. *Journal of Obstetrics and Gynaecology Research* 2010;**36**(5):978–83.
- Petrou 2006** {published data only}
Petrou S, Trinder J, Brocklehurst P, Smith L. Economic evaluation of alternative management methods of first-trimester miscarriage based on results from the MIST trial. *BJOG: An International Journal of Obstetrics & Gynaecology* 2006;**113**(8):879–89.
- Prasartsakulchai 2004** {published data only}
Prasartsakulchai C, Tannirandorn Y. A comparison of vaginal misoprostol 800 microg versus 400 microg in early pregnancy failure: a randomized controlled trial. *Journal of the Medical Association of Thailand* 2004;**87** Suppl 3: S18–23.
- Ramadan 2009** {published data only}
Ramadan MC. Misoprostol versus MVA for incomplete abortion: results from a randomized controlled trial in Egypt. *International Journal of Gynecology & Obstetrics* 2009;**107**(Suppl 2):S68–9.
- Rausch 2012** {published data only}
Rausch M, Lorch S, Chung K, Frederick M, Zhang J, Barnhart K. A cost-effectiveness analysis of surgical versus medical management of early pregnancy loss. *Fertility and Sterility* 2012;**97**(2):355–60.e1.
- Reeves 2006** {published data only}
Reeves MF, Lohr PA, Harwood B, Creinin MD. Sonographic findings after misoprostol or vacuum aspiration for early pregnancy failure. *Contraception* 2006;**74**(2):182.
- Reeves 2008** {published data only}
Reeves MF, Lohr PA, Harwood BJ, Creinin MD. Ultrasonographic endometrial thickness after medical and surgical management of early pregnancy failure. *Obstetrics & Gynecology* 2008;**111**(1):106–12.
- Rita 2006** {published data only}
Rita Gupta S, Kumar S. A randomised comparison of oral and vaginal misoprostol for medical management of first trimester missed abortion. *JK Science* 2006;**8**(1):35–8.
- Rivero-Lopez 1998** {published data only}
Rivero-Lopez E, Marquez-Maraver F, Duenas-Diez JL, Cabezas-Sanchez B. Deferred miscarriage: effectiveness of intravaginal misoprostol versus laminaria alone [Aborto diferido: eficacia del misoprostol intravaginal versus la aplicacion de tallos de laminaria]. *Progresos de Obstetricia y Ginecologia* 1998;**41**:579–81.

Robledo 2007 {published data only}

Robledo C, Zhang J, Troendle J, Barnhart K, Creinin MD, Westhoff C, et al. Clinical indicators for success of misoprostol treatment after early pregnancy failure. *International Journal of Gynecology & Obstetrics* 2007;**99**(1): 46–51.

Rosenberg 2006 {published data only}

Rosenberg P. Expectant versus immediate medical management for the evacuation of the nonevolutive pregnancies before 13 GW (ongoing trial). ClinicalTrials.gov (<http://clinicaltrials.gov/>) (accessed 21 March 2006) 2006.

Ruangchainikhom 2006 {published data only}

Ruangchainikhom W, Phongphisanou E, Bhekasuta J, Sarapak S. Effectiveness of 400 or 600 micrograms of vaginal misoprostol for terminations of early pregnancies. *Journal of the Medical Association of Thailand* 2006;**89**(7): 928–33.

Saichua 2009 {published data only}

Saichua C, Phupong V. A randomized controlled trial comparing powdery sublingual misoprostol and sublingual misoprostol tablet for management of embryonic death or anembryonic pregnancy. *Archives of Gynecology and Obstetrics* 2009;**280**(3):431–5.

Sathapanachai 2000 {published data only}

Sathapanachai U. Intravaginal 400 micrograms misoprostol for pregnancy termination in cases of blighted ovum. *Thai Journal of Obstetrics and Gynaecology* 2000;**12**(4):363.

Shah 2010 {published data only}

Shah N, Azam SI, Khan NH. Sublingual versus vaginal misoprostol in the management of missed miscarriage. *JPMA - Journal of the Pakistan Medical Association* 2010;**60**(2):113–6.

Shaikh 2008 {published data only}

Shaikh ZAN. Comparison between misoprostol alone and misoprostol with manual vacuum aspiration for the treatment of missed and incomplete miscarriage. *BJOG: an international journal of obstetrics and gynaecology* 2008;**115**(s1):83.

Shelley 2005 {published data only}

Shelley JM, Healy D, Grover S. A randomised trial of surgical, medical and expectant management of first trimester spontaneous miscarriage. *Australian and New Zealand Journal of Obstetrics and Gynaecology* 2005;**45**: 122–7.

Shobeira 2007 {published data only}

Shobeira JM, Atashkhoii S. Second trimester pregnancy termination by intravaginal and parenteral form of prostaglandin E2 [abstract]. 31st British International Congress of Obstetrics and Gynaecology; 2007 July 4–6; London, UK. 2007:210.

Shokry 2009 {published data only}

Shokry M, Shahin AY, Fathalla MM, Shaaban OM. Oral misoprostol reduces vaginal bleeding following surgical evacuation for first trimester spontaneous abortion. *International Journal of Gynecology & Obstetrics* 2009;**107**(2):117–20.

Shwekerela 2007 {published data only}

Shwekerela B, Kalumuna R, Kipingili R, Mashaka N, Westheimer E, Clark W, et al. Misoprostol for treatment of incomplete abortion at the regional hospital level: results from Tanzania. *BJOG: an international journal of obstetrics and gynaecology* 2007;**114**(11):1363–7.

Smith 2006 {published data only}

Smith L. Extension to: randomised controlled trial of expectant, medical and surgical management of early miscarriage. Research Findings Register (www.refer.nhs.uk) (accessed 7 March 2006) 2006.

Smith 2006a {published data only}

Smith LF, Frost J, Levitas R, Bradley H, Garcia J. Women's experiences of three early miscarriage management options: a qualitative study. *British Journal of General Practice* 2006;**56**(524):198–205.

Smith 2009 {published data only}

Smith LF, Ewings PD, Quinlan C. Incidence of pregnancy after expectant, medical, or surgical management of spontaneous first trimester miscarriage: long term follow-up of miscarriage treatment (MIST) randomised controlled trial. *BMJ* 2009;**339**:b3827.

Srikhao 2005 {published data only}

Srikhao N, Tannirandorn Y. A comparison of vaginal misoprostol 800 microg versus 400 microg for anembryonic pregnancy: a randomized comparative trial. *Journal of the Medical Association of Thailand* 2005;**88**(Suppl 2):S41–7.

Sripramote 2000 {published data only}

Sripramote M, Chatsuphang W. A randomized comparison of oral and vaginal misoprostol for cervical priming before uterine curettage in the first trimester of pregnancy. *Vajira Medical Journal* 2000;**44**(3):207–15.

Stockheim 2006 {published data only}

Stockheim D, Machtinger R, Wiser A, Dulitzky M, Soriano D, Goldenberg M, et al. A randomized prospective study of misoprostol or mifepristone followed by misoprostol when needed for the treatment of women with early pregnancy failure. *Fertility and Sterility* 2006;**86**(4):956–60.

Suchonwanit 1999 {published data only}

Suchonwanit P. Comparative study between vaginal misoprostol 200 mg and 400 mg in first trimester intrauterine fetal death and anembryonic gestation. *Thai Journal of Obstetrics and Gynaecology* 1999;**11**(4):263.

Tam 2005 {published data only}

Tam WH, Tsui MH, Lok IH, Yip SK, Yuen PM, Chung TK. Long-term reproductive outcome subsequent to medical versus surgical treatment for miscarriage. *Human Reproduction* 2005;**20**(12):3355–9.

Tang 2006 {published data only}

Tang OS, Ong CY, Tse KY, Ng EH, Lee SW, Ho PC. A randomized trial to compare the use of sublingual misoprostol with or without an additional 1 week course for the management of first trimester silent miscarriage. *Human Reproduction* 2006;**21**(1):189–92.

Tanha 2010 {published data only}

Tanha FD. Comparison of the efficacy of two routes of misoprostol administration (sublingual and vaginal) for termination of second trimester pregnancy. IRCT Iranian Registry of Clinical Trials (www.irct.ir) (accessed 6 December 2010). IRCT138807272576N1 2010.

Tanha 2010a {published data only}

Tanha FD, Feizi M, Shariat M. Sublingual versus vaginal misoprostol for the management of missed abortion. *Journal of Obstetrics and Gynaecology Research* 2010;**36**(3): 525–32.

Taylor 2011 {published data only}

Taylor J, Diop A, Blum J, Dolo O, Winikoff B. Oral misoprostol as an alternative to surgical management for incomplete abortion in Ghana. *International Journal of Gynecology & Obstetrics* 2011;**112**(1):40–4.

Toptas 2011 {published data only}

Toptas T, Mendilcioglu I, Simsek M, Taskin O. Comparison of intravaginal misoprostol alone to combination of intravaginal misoprostol and extraamniotic Foley catheter for the second trimester of pregnancies. *American Journal of Obstetrics and Gynecology* 2011;**204**(1 Suppl):S126.

Torre 2012 {published data only}

Torre A, Huchon C, Bussieres L, Machevin E, Camus E, Fauconnier A. Immediate versus delayed medical treatment for first-trimester miscarriage: a randomized trial. *American Journal of Obstetrics & Gynecology* 2012;**206**(3):215.e1–6.

Trinder 2006 {published data only}

Trinder J, Brocklehurst P, Porter R, Read M, Vyas S, Smith L. Management of miscarriage: expectant, medical, or surgical? Results of randomised controlled trial (miscarriage treatment (MIST) trial). *BMJ* 2006;**332**(7552):1235–40.

Van Mensel 2009 {published data only}

Van Mensel K, Claerhout F, Debois P, Keirse MJNC, Hanssens M. A randomized controlled trial of misoprostol and sulprostone to end pregnancy after fetal death [Article ID 496320]. *Obstetrics and Gynecology International* 2009.

Yilmaz 2005 {published data only}

Yilmaz B, Kelekci S, Ertas IE, Kahyaoglu S, Ozel M, Sut N, et al. Misoprostol moistened with acetic acid or saline for second trimester pregnancy termination: a randomized prospective double-blind trial. *Human Reproduction* 2005;**20**(11):3067–71.

Yilmaz 2007 {published data only}

Yilmaz B, Kelekci S, Ertas IE, Ozel M, Sut N, Mollamahmutoglu L, et al. Randomized comparison of second trimester pregnancy termination utilizing saline moistened or dry misoprostol. *Archives of Gynecology and Obstetrics* 2007;**276**(5):511–6.

Zanganeh 2010 {published data only}

Zanganeh M. Comparing the effects of multiple doses of misoprostol with single dose of misoprostol plus oxytocin in induction of second trimester abortion. IRCT Iranian Registry of Clinical Trials (www.irct.ir) (accessed 6 December 2010).

Zhang 2000 {published data only}

Zhang C, Cheng W. A contrastive analysis of the efficacy of misoprostol and li fan nuo in intermediate term of pregnancy. *Journal of Wuban University of Science and Technology (Natural Science Edition)* 2000;**23**(4):409–11.

References to ongoing studies**Australia 2000 {published data only}**

Louey K. Misoprostol for the medical management of miscarriage (RCT). Personal communication 2000.

UK 1999 {published data only}

Trinder J. Miscarriage Treatment Study (MIST). National Research Register www.nrr.nhs.uk (accessed 1999).

Additional references**Alfirevic 2001**

Alfirevic Z. Oral misoprostol for induction of labour. *Cochrane Database of Systematic Reviews* 2001, Issue 2.

Ashok 1998

Ashok PW, Penney GC, Flett GM, Templeton A. An effective regimen for early medical abortion: a report of 2000 consecutive cases. *Human Reproduction* 1998;**13**: 2962–5.

Baulieu 1986

Baulieu E, Ulmann A. Antiprogestosterone activity of RU-486 and its contraceptive and other applications. *Human Reproduction* 1986;**1**:107–10.

Bugalho 1996

Bugalho A, Faundes A, Jamisse L, Usfa M, Maria E, Bique C. Evaluation of the effectiveness of vaginal misoprostol to induce first trimester abortion. *Contraception* 1996;**53**: 244–6.

Cameron 1986

Cameron IT, Michie AF, Baird DT. Therapeutic abortion in early pregnancy with antiprogesterone RU486 alone or in combination with prostaglandin analogue (gemeprost). *Contraception* 1986;**34**(5):459–68.

Costa 1993

Costa SH, Vessey MP. Misoprostol and illegal abortion in Rio de Janeiro, Brazil. *Lancet* 1993;**341**:1258–61.

Fonseca 1991

Fonseca W, Alencar AJC, Mota FSB, Coelho HLL. Misoprostol and congenital malformations. *Lancet* 1991;**338**:56.

Grudzinskas 1995

Grudzinskas JG. Endocrinological and metabolic assessment of early pregnancy. In: Chamberlain G editor (s). *Turnbull's obstetrics*. London: Pearson Professional Ltd, 1995:185–93.

Gulmezoglu 2004

Gulmezoglu AM, Forna F, Villar J, Hofmeyr GJ. Prostaglandins for prevention of postpartum haemorrhage. *Cochrane Database of Systematic Reviews* 2004, Issue 1.

Hofmeyr 2003

Hofmeyr GJ, Gulmezoglu AM. Vaginal misoprostol for cervical ripening and induction of labour. *Cochrane Database of Systematic Reviews* 2003, Issue 1.

Howie 1995

Howie PG. Abortion and ectopic pregnancy. In: Whitfield CR editor(s). *Dewhurst's Textbook of Obstetrics and Gynecology for Postgraduates*. Oxford: Blackwell Science Ltd, 1995:140–63.

Kovacs 1984

Kovacs L, Sas M, Resch BA, Ugocsai G, Swahn ML, Bygdeman, et al. Termination of very early pregnancy by RU 486—an antiprogesterone compound. *Contraception* 1984; **29**(5):399–410.

Kulier 2004

Kulier R, Gulmezoglu AM, Hofmeyr GJ, Cheng LN, Campana A. Medical methods for first trimester abortion. *Cochrane Database of Systematic Reviews* 2004, Issue 1.

Mousa 2003

Mousa HA, Alfirevic Z. Treatment for primary postpartum haemorrhage. *Cochrane Database of Systematic Reviews* 2003, Issue 1.

Muzonzini 2004

Muzonzini G, Hofmeyr GJ. Buccal or sublingual misoprostol for cervical ripening and induction of labour. *Cochrane Database of Systematic Reviews* 2004, Issue 4.

Nanda 2002

Nanda K, Peloggia A, Nanda G, Grimes D. Expectant care versus surgical treatment for miscarriage. *Cochrane Database of Systematic Reviews* 2002, Issue 1.

Neilson 1998

Neilson JP. Ultrasound for fetal assessment in early pregnancy. *Cochrane Database of Systematic Reviews* 1998, Issue 4.

Norman 1991

Norman JE, Thong KJ, Baird DT. Uterine contractility and induction of abortion in early pregnancy by misoprostol and mifepristone. *Lancet* 1991;**338**:1233–6.

Say 2002

Say L, Kulier R, Gulmezoglu AM, Campana A. Medical versus surgical methods for first trimester termination of pregnancy. *Cochrane Database of Systematic Reviews* 2002, Issue 4.

Senior 1993

Senior J, Marshall K, Sangha R, Clayton JK. In vitro characterisation of prostanoid receptors on human myometrium at term pregnancy. *British Journal of Pharmacology* 1993;**108**:501–6.

Simpson 1991

Simpson JL. Fetal wastage. In: Gabe SG, Simpson JL editor (s). *Obstetrics. Normal and problem pregnancies*. New York: Churchill Livingstone Inc, 1991:783–807.

Vazquez 2000

Vazquez JC, Hickey M, Neilson JP. Medical treatment for miscarriage. *Cochrane Database of Systematic Reviews* 2000, Issue 3.

Zieman 1997

Zieman M, Fong SK, Benowitz NL, Bankster D, Darney PD. Absorption kinetics of misoprostol with oral or vaginal administration. *Obstetrics & Gynecology* 1997;**90**:88–92.

* Indicates the major publication for the study

CHARACTERISTICS OF STUDIES

Characteristics of included studies [ordered by study ID]

Al Inizi 2003

Methods	'Random allocation'. Details unknown.	
Participants	60 women with early non-viable pregnancies diagnosed by ultrasound	
Interventions	Vaginal misoprostol 400 mcg repeated twice a day to maximum of 1600 mcg (n = 27) versus dinoprostone (PGE2) vaginal tablets repeated 6 hourly intervals to maximum of 36 mg (n = 33)	
Outcomes	Complete miscarriage/need for surgical evacuation.	
Notes	Authors contacted.	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	B - Unclear

Autry 1999

Methods	Randomisation using a random number tables. Allocation concealment was accomplished in sequentially numbered opaque sealed envelopes made available at the time of enrollment in the study. Intention to treat analysis	
Participants	21 women diagnosed with a non-viable first trimester intrauterine pregnancy up to 49 days gestation. Evidence of non-viability included one of the following findings on TVS: 1) mean gestational sac diameter greater than 18 mm and no embryonic pole; 2) embryonic pole 5-10 mm without cardiac activity; 3) intrauterine gestational sac with abnormal hCG titers. Others entry criteria: 1) 18 years of age or greater; 2) closed cervix on digital exam; 3) no known intolerance or allergy to misoprostol or MTX; 4) hemoglobin of 9 g/dl or greater; 5) platelet count of 100,000/mcl or greater; 6) no history of blood clotting disorders; 7) no active liver or renal disease; 8) ability and willingness to comply with visit schedule; 9) hCG less than 40 000 IU/l; and 10) easy access to a telephone and transportation	
Interventions	Combined group (n = 12): IM MTX 50 mg/m ² body surface area (day 1) followed two days later (day 3) by vaginal misoprostol 800 mcg (by vaginal placement of four 200 mcg-tablets of misoprostol). If the gestational sac was present vaginal misoprostol was repeated. Misoprostol only group (n = 9): four 200 mcg-tablets placed in the vagina on day 1. The remainder of the follow up was similar to that for combined group	
Outcomes	Successful complete abortion: MTX plus misoprostol 12/12 vs misoprostol only 8/9. No blood transfusion or antibiotics. Positive urine pregnancy test at the initial follow-up appointment: 2/9 vs 7/7. Pain relief: 4/12 vs 4/9	

Autry 1999 (Continued)

Notes	Wisconsin, Milwaukee, USA. All women received: 1) prescription for 10 tablets acetaminophen with codeine (300 mg/30 mg) and 8 tablets of ibuprofen (600 mg); 2) instruction sheet including phone number to contact physician 24 hours/day; and a diary sheet to record symptoms, side-effects, and pain medication use. Data about side-effects (headache, nausea and emesis) and women's satisfaction reported as no separate data. Authors conclude that both treatments are effective regimens for the complete evacuation of non-viable early first trimester pregnancy, and represent a reasonable alternative for women wishing to avoid surgery
-------	---

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

Bagratee 2004

Methods	Computer-generated random allocation of study number. Numbered envelopes containing misoprostol or placebo
Participants	104 women who attended Early Pregnancy Unit, St Mary's Hospital, with incomplete miscarriage or early pregnancy failure < 13 weeks
Interventions	600 mcg misoprostol (n = 52) or placebo [expectant management] (n = 52). Second dose next day unless complete miscarriage had occurred in meantime. Review day 7 and surgical evacuation if miscarriage not complete. Further review at day 14
Outcomes	Primary: complete miscarriage without need for ERPC by day 7. Secondary outcomes: clinical, side-effects, satisfaction and future choices
Notes	Primary outcome reported for both non-viable pregnancies and incomplete miscarriages, but not for secondary outcomes. These will be added if authors can provide data separately for non-viable pregnancies and incomplete miscarriages

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

Creinin 1997

Methods	Sealed, numbered, sequential envelopes containing instructions based on computer-generated random number table
Participants	20 women with non-viable pregnancies diagnosed by transvaginal ultrasound; < 9 weeks; closed cervix; no contra-indication to misoprostol; no heavy bleeding
Interventions	400 mcg misoprostol orally, repeated after 24 hours if the pregnancy had not been expelled (n = 12); vaginal misoprostol 800 mcg - repeated after 24 hours if necessary (as above) (n = 8). Surgical evacuation offered to women in both groups after 48 hours if treatment unsuccessful
Outcomes	Miscarriage; pain (visual analogue scale); side-effects.
Notes	Pilot study.

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

Demetroulis 2001

Methods	Randomisation by opening sealed opaque envelope containing computer generated allocation code number. No attempt at masking given the manifest differences between medical and surgical interventions
Participants	80 women with incomplete miscarriage or anembryonic pregnancy or missed miscarriage < 13 weeks, diagnosed by ultrasound. The data in this review are derived only from the subgroup with non-viable pregnancies (n = 50) and not those with incomplete miscarriages. Women were reviewed 8-10 hours after medical treatment; if they had empty uteruses on ultrasound examination they were discharged home; if not, surgical evacuation was arranged
Interventions	Vaginal misoprostol 800 mcg once only (n = 26) versus surgical evacuation of the uterus (n = 24)
Outcomes	Need for surgical evacuation, symptoms including pain and bleeding, 'satisfaction'
Notes	Authors contacted for information on outcomes according to indication for treatment. Only usable data currently available are on incidence of surgical evacuation

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

Egarter 1995

Methods	Women “randomly assigned”; no details.	
Participants	87 women in Austria with non-viable pregnancies between 8 and 12 weeks, diagnosed by ultrasound	
Interventions	Vaginal gemeprost 1 mg every 3 hours up to maximum of 3 mg daily for 2 days (n = 43) versus uterine curettage (n = 44)	
Outcomes	Need for surgical curettage. Adverse effects.	
Notes		
<i>Risk of bias</i>		
Bias	Authors’ judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	B - Unclear

Eng 1997*

Methods	Randomised by “blindly picking a sealed number from a box”. Treatment allocation was then based on whether the number was odd or even	
Participants	50 women with intrauterine fetal death at 13-26 weeks of pregnancy	
Interventions	Vaginal misoprostol 200 mcg 3-hourly up to a maximum dose of 1200 mcg (n = 25) versus vaginal gemeprost 1 mg 3-hourly up to a maximum dose of 5 mg (n = 25)	
Outcomes	Main outcome - “treatment failure” defined as failure to miscarry within 24 hours, or side-effects severe enough to preclude use of additional dose of drug	
Notes		
<i>Risk of bias</i>		
Bias	Authors’ judgement	Support for judgement
Allocation concealment (selection bias)	High risk	C - Inadequate

Fadalla 2004*

Methods	“Randomised”; no details.	
Participants	70 women in the Wad Medeni Teaching Hospital, Sudan, with fetal deaths between 13 and 28 weeks, diagnosed by ultrasound	

Fadalla 2004* (Continued)

Interventions	Oral misoprostol (n = 35) versus vaginal misoprostol (n = 35) - both administered as 100 mcg tablets 4-hourly until initiation of labour	
Outcomes	Time to delivery; oxytocin infusion; manual removal of placenta	
Notes		
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	B - Unclear

Gilles 2004

Methods	Random allocation by computer-automated telephone response system. Stratification by pregnancy type. Random permuted blocks of size 4 or 8	
Participants	80 women with anembryonic pregnancy < 46 mm sac diameter or embryonic/fetal death with crown-rump length < 41 mm. Four centres	
Interventions	“Wet misoprostol” 800 mcg + 2 ml saline vaginally (n = 41) versus “dry misoprostol” (as above without saline) (n = 39). Second dose given day 3 if no miscarriage	
Outcomes	Primary outcome: miscarriage without need for curettage before 30 days. Secondary outcomes: miscarriage < 3, < 8 and < 15 days; side-effects, women’s views	
Notes		
<i>Risk of bias</i>		
Bias	Authors’ judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

Graziosi 2004

Methods	Consent for study obtained at time of diagnosis of early pregnancy failure. Randomised after at least one week of expectant management. Computer programme with block randomisation sequence. Stratification by previous vaginal birth; Gestational age < or > 10 weeks; centre	
Participants	154 women with ultrasound-diagnosed early pregnancy failure - either anembryonic pregnancy or fetal death at 6-14 weeks. 6 centre study in Netherlands	

Graziosi 2004 (Continued)

Interventions	Vaginal misoprostol 800 mcg; repeated after 24 hours if ultrasound indicated remaining tissue in the uterus. Curettage after 3 days if miscarriage hadn't occurred or was incomplete (n = 79) or suction curettage within a week of randomisation (n = 75)	
Outcomes	Primary: complete evacuation. Secondary: side-effects, pain and need for analgesia, intensity/duration of bleeding	
Notes	Of 241 eligible women, 87 (36%) declined to participate and chose curettage	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

Heard 2002

Methods	“Randomized” - no details.	
Participants	33 women with “missed abortion”.	
Interventions	Vaginal misoprostol 400 mcg (n = 12) versus 800 mcg (n = 21)	
Outcomes	Only usable outcome reported in abstract was miscarriage.	
Notes	Abstract - no explanation for unbalanced numbers.	
<i>Risk of bias</i>		
Bias	Authors’ judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	B - Unclear

Herabutya 1997

Methods	"Random allocation" but method not discussed in paper.	
Participants	84 women with ultrasound confirmation of fetal death with uterine size < 14 weeks, no bleeding, and cervix closed	
Interventions	Misoprostol (200 micrograms vaginally) (n = 42) or vaginal placebo (n = 42) on admission to hospital	
Outcomes	Primary outcome was miscarriage within 24 hours of treatment. Some information available on complications	

Herabutya 1997 (Continued)

Notes	Much of the outcome data reported describes only the subgroups who did miscarry before surgical evacuation	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	B - Unclear

Jain 1996*

Methods	“Random number table”.	
Participants	70 women in Los Angeles, USA, with either fetal death (n = 40) or medical or genetic indications for termination of pregnancy (n = 30) at 12-22 weeks. Only data from pregnancies complicated by fetal death included here	
Interventions	Vaginal misoprostol 200 mcg 12-hourly plus laminaria tents (n = 20) versus vaginal misoprostol 200 mcg 12-hourly alone	
Outcomes	Miscarriage.	
Notes	Adverse effects are described for the groups as wholes, so are not included here. 2 women excluded from analyses - 1 protocol violation; 1 was found to have interstitial ectopic pregnancy	
<i>Risk of bias</i>		
Bias	Authors’ judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	B - Unclear

Kara 1999*

Methods	"Random allocation". No details.	
Participants	65 women in Istanbul, Turkey, with ultrasound-diagnosed fetal death in second trimester	
Interventions	Vaginal misoprostol 200 mcg (n = 32) versus intracervical dinoprostol 0.5 mg (n = 33). Intravenous oxytocin started after 6 hours if no 'effective contractions'	
Outcomes	Complete miscarriage. Adverse effects.	
Notes	Misoprostol dose reported as 200 mg. Assumed to be 200 mcg. Time to miscarriage not included as standard deviations seem incorrect	

<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	B - Unclear

Kovavisarach 2002

Methods	"Random allocation". Method not discussed.	
Participants	54 women with anembryonic pregnancies < 12 weeks diagnosed by TVS. Single centre study in Bangkok, Thailand	
Interventions	Vaginal misoprostol 400 mcg (n = 27) or placebo (n = 27). Reviewed after 24 hours and curettage offered if no or incomplete miscarriage. Further review after 1 week	
Outcomes	Primary: complete miscarriage within 24 hours of treatment.	
Notes		

<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	B - Unclear

Kovavisarach 2005

Methods	Random allocation using sealed, sequentially numbered envelopes, prepared using published table of random numbers	
Participants	114 women in Bangkok, Thailand, with non-viable pregnancies (anembryonic or fetal deaths) at < 12 weeks, diagnosed by TVS. Women with open cervixes were not eligible for recruitment	
Interventions	Vaginal misoprostol 600 mcg (n = 57) or 800 mcg (n = 57). If complete miscarriage not effected within 24 hours, or if clinical circumstances dictated (pain, bleeding), uterine curettage was performed	
Outcomes	Primary: complete miscarriage without need for uterine curettage within 24 hours. Secondary: adverse effects	
Notes		

<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement

Allocation concealment (selection bias)	Low risk	A - Adequate
---	----------	--------------

Lelaidier 1993

Methods	Drug or identical placebo supplied by pharmacy using randomisation list using permutation blocks of four
Participants	46 women with non-viable pregnancies diagnosed by ultrasound on two examinations separated by one week. < 14 weeks. No bleeding or pain
Interventions	Mifepristone 600 mg orally (n = 23) or placebo (n = 23). All women were reviewed after 5 days and if miscarriage had not occurred, surgical evacuation was performed that day
Outcomes	Primary outcome was expulsion of the pregnancy. Symptoms also recorded
Notes	Two women in the placebo group underwent surgical evacuation by private practitioners before 5th day review. Both were in the process of miscarriage and were classed as expulsion positive; no information available on symptoms

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

Lister 2005

Methods	Random allocation - blocked and stratified by physician office and by day of recruitment - day of diagnosis, or after day of diagnosis
Participants	34 women in Columbus, Ohio, USA, with early pregnancy failure (anembryonic pregnancies or early fetal deaths) diagnosed by TVS
Interventions	Vaginal misoprostol 800 mcg, repeated after 24 hours if sac still present on TVS (n = 18) or placebo (n = 16)
Outcomes	Primary: miscarriage complete at 48 hours.
Notes	Planned sample size 84 but trial stopped after interim analysis of first 36 women. Two women excluded from analysis - one protocol violation; one did not meet entry criteria. Two women did not come for assessment 2 weeks after initial treatment

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Lister 2005 (Continued)

Allocation concealment (selection bias)	Low risk	A - Adequate
---	----------	--------------

Muffley 2002

Methods	Computer-generated random number table with blocked permutations - group assignments recorded in sealed opaque numbered envelopes
Participants	50 women with non-viable pregnancies diagnosed by ultrasound (anembryonic or embryonic/fetal deaths) < 12 weeks. Exclusions: excessive bleeding, anaemia, unstable vital signs, coagulopathy, asthma or other contra-indication to prostaglandin treatment, infection, open cervix
Interventions	800 mcg misoprostol vaginally, repeated after 24 hours if ultrasound showed tissue still present in uterus; final review after further 24 hours - if tissue still present, surgical evacuation performed (n = 25). Suction curettage (n = 25)
Outcomes	Primary outcome: miscarriage.
Notes	Analysis by intention to treat. Details about nausea, vomiting, diarrhoea reported only for misoprostol group

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

Ngoc 2004

Methods	Randomised by opening sequentially numbered envelope - prepared by computer-generated code in blocks of 10
Participants	200 women in Ho Chi Minh City, Vietnam, with non-viable first trimester pregnancies (anembryonic or early fetal death) diagnosed by ultrasound; cervix closed
Interventions	Oral misoprostol 800 mcg (n = 100) versus vaginal misoprostol 800 mcg (n = 98). Women reviewed after 48 hours; if retained products present, they were given option of surgical evacuation or further review after another 5 days (when evacuation was performed if there were still products present)
Outcomes	Primary: complete miscarriage without need for surgical evacuation. Secondary: adverse effects
Notes	2 women lost to follow up.

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

Nielsen 1999

Methods	Randomisation method not discussed in paper.
Participants	122 women < 13 weeks with symptoms of threatened miscarriage (bleeding +/- pain), a closed cervix, and ultrasound demonstration of pregnancy non-viability (anembryonic pregnancy n = 44; embryonic/fetal death n = 46; 'complex mass with deformed gestational sac' n = 32). Surgical evacuation at day 5 if transvaginal ultrasound showed retained products > 15 mm diameter
Interventions	Mifepristone (400 mg orally) followed by oral misoprostol (400 micrograms) 48 hours later (n = 60) versus expectant management (n = 62)
Outcomes	Clinical events; routine transvaginal ultrasound at 5 days to identify retained products; visual analogue scale to assess pain at day 5; visual analogue scale to assess satisfaction at day 14
Notes	Seeking clarification from authors if "complex mass with deformed gestational sac" represents missed or incomplete miscarriage. Data included in meantime

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	B - Unclear

Niromanesh 2005*

Methods	Randomisation method not discussed in paper.
Participants	100 women in Tehran, Iran, with fetal deaths between 14 and 25 weeks
Interventions	Vaginal misoprostol: 400 mcg (n = 50) versus 600 mcg (n = 50) - both 12-hourly for 48 hours
Outcomes	Miscarriage; surgical evacuation; adverse effects.
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Allocation concealment (selection bias)	Unclear risk	B - Unclear
---	--------------	-------------

Tang 2003

Methods	Randomisation by "computer-generated random numbers".
Participants	80 women with non-viable pregnancies diagnosed by ultrasound < 13 weeks
Interventions	Group 1: 600 mcg misoprostol sublingually every 3 hours for maximum of 3 doses (n = 40); Group 2: 600 mcg misoprostol vaginally every 3 hours for maximum of 3 doses (n = 40). Women discharged home after completion of treatment and reassessed day 7 - when surgical evacuation performed if gestation sac still present, or retained products of conception plus heavy bleeding
Outcomes	Primary outcome: complete miscarriage (defined as no need for surgical evacuation up until return of menstruation)
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

Wood 2002

Methods	Computer-generated random number list in blocks. Pharmacy prepared numbered envelopes. Tablets not identical so placed by nurse in opaque vaginal introducer for physician to insert - to maintain allocation concealment
Participants	50 women with ultrasound diagnosed non-viable pregnancies. Gestational age 7-17 weeks but women not included if fetal size by ultrasound > 12 weeks equivalent. Also excluded from recruitment if experiencing uterine cramping or bleeding
Interventions	Misoprostol (800 mcg vaginally) (n = 25) or vaginal placebo (n = 25). If complete miscarriage not suspected after 24 hours, treatment was repeated. At 48 hours, if no miscarriage or miscarriage thought to be incomplete, uterine curettage was offered
Outcomes	Sample size based on reduction of uterine curettage from 50% to 10%. Women's satisfaction also assessed, but are not included in analyses as data not reported from control group
Notes	Analysis by intention to treat.

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

ERPC: evacuation of retained products of conception

hCG: human chorionic gonadotropin

IM: intramuscular

IU: international units

mcg: microgrammes

MTX: methotrexate

POCs: products of conception

TVS: transvaginal sonography

vs: versus

Characteristics of excluded studies [ordered by study ID]

Study	Reason for exclusion
Abdel Fattah 1997	Conference abstract. No information about gestational age but, given title, probably includes pregnancies > 24 weeks as well as < 24 weeks
Almog 2005	Termination of 'viable' pregnancies - mainly with fetal anomalies
Anderman 2000	Conference abstract. Includes pregnancies > 24 weeks as well as < 24 weeks
Avila-Vergara 1997	Intrauterine deaths mainly third trimester.
Bebbington 2002	Termination of viable pregnancies.
Cabrol 1990	Trial of mifepristone for induction of labour after intrauterine death - but mainly late second and third trimester pregnancies
Clevin 2001	Abstract in Danish. A prospective, randomised study carried out to clarify the effect of vaginal administration of a prostaglandin E1 analogue (gemeprost) versus surgical management (curettage) on miscarriages at up to twelve weeks of gestation. Three groups: 1 (n = 27), 2A (n = 17) and 2B (n = 17), allocated according to the endometrial thickness. The measured outcomes were reduction of endometrial thickness, duration of vaginal bleeding and pain, reported in a non-suitable format for analysis
David 2003	Randomised trial (details of randomisation unclear) of two methods to soften the cervix before surgical evacuation for early non-viable pregnancies. No usable clinical data, given short timescale between treatment and surgery

(Continued)

Dickinson 1998	Trial included women with fetal malformations and maternal indications for pregnancy termination between 14 and 28 weeks, as well as pregnancies with fetal death. Data will be included for the latter if these can be obtained from the authors
Dickinson 2002	Trial included women with fetal malformations and maternal indications for pregnancy termination between 14 and 30 weeks, as well as pregnancies with fetal death. Data will be included for the latter if these can be obtained from the authors
Dickinson 2003	Randomised trial comparing oral with vaginal administration of misoprostol to terminate pregnancies with fetal malformations - not non-viable pregnancies
Eppel 2005	Trial included women with fetal malformations and maternal indications for pregnancy termination between 14 and 23 weeks, as well as pregnancies with fetal death. Data will be included for the latter if these can be obtained from the authors
Feldman 2003	Trial included women with fetal malformations and maternal indications for pregnancy termination between 14 and 23 weeks, as well as pregnancies with fetal death. Data will be included for the latter if these can be obtained from the authors
Ghorab 1998	Trial included women with fetal malformations for pregnancy termination, as well as pregnancies with fetal death. Data will be included for the latter if these can be obtained from the authors
Gonzalez 2001	Trial included women with fetal malformations and maternal indications for pregnancy termination between 14 and 23 weeks, as well as pregnancies with fetal death. Data will be included for the latter if these can be obtained from the authors
Grimes 2004	Trial included women with other reasons for pregnancy termination, as well as pregnancies with fetal death. Data will be included for the latter if these can be obtained from the authors
Gronland 2002	Not a randomised trial. Three centre study of women with non-viable pregnancies comparing three treatment regimens: misoprostol, mifepristone + misoprostol, surgical evacuation - with treatment regimen changing at each hospital every four months
Hausler 1997	Prospective randomised controlled trial evaluating three interventions for complete spontaneous abortion. Diagnosis was based on positive pregnant test, vaginal bleeding and/or evacuation of tissue from the vagina, a closed uterine orifice with only slight bleeding on admission and a possible clear sonographic pregnancy diagnosis in the history. Interventions: A) n = 15 curettage; B) n = 20 only controlled and; C) n = 15 additionally treated for 10 days with an oral hormone intake of 2 mg norethisterone acetate and 0.01 mg ethinyl oestradiol 3 x day. Randomisation by sealed unmarked envelopes. 63 patients were included in the study and allocated randomly to each group. 13 women (20.6%) were excluded from the study after randomisation: 10 did not report for the planned follow-up control, one did not report for curettage, in one the height of the endometrium was > 8 mm and in one an ectopic pregnancy was diagnosed 6 days after the randomisation. The study only presents outcomes, in a non-suitable format, regarding hCG clearing time and duration of the secondary haemorrhage from the day of randomisation
Herabutya 2005	RCT of misoprostol for terminating viable pregnancies.

(Continued)

Hidar 2001	Trial included women with fetal malformations and maternal indications for pregnancy termination between 13 and 29 weeks, as well as pregnancies with fetal death. Data will be included for the latter if these can be obtained from the authors
Hill 1991	Trial includes fetal deaths in both second and third trimesters
Hogg 2000	Abstract. Trial included women with other reasons for pregnancy termination, as well as pregnancies with fetal death. Data will be included for the latter if these can be obtained from the authors
Jain 1994	Trial included women with fetal malformations and maternal indications for pregnancy termination between 12 and 22 weeks, as well as pregnancies with fetal death. Data will be included for the latter if these can be obtained from the authors
Jain 1999	Trial included women with fetal malformations and maternal indications for pregnancy termination between 12 and 22 weeks, as well as pregnancies with fetal death. Data will be included for the latter if these can be obtained from the authors
Johnson 1997	Randomised controlled trial evaluating pain and bleeding and comparing surgical to medical treatment. Surgical arm (n = 12) uterine curettage under general anesthesia. Medical arm (n = 17) include three different participant conditions and treatments: a) no treatment if women had a complete abortion and uterine cavity echo (nyometrium-myometrium) less than 1.5 mm; b) women with incomplete abortion : 1 mg pessary of Gemeprost (Cervagem, May and Baker) and remained in hospital for 4 hours or until the had passed POC; and c) women with intact gestational sac (but non-viable fetus) 200 mg RU 486 (mifepristone) and then allowed home, readmitted 36-48 hours later for 1 mg of vaginal Cervagem. Data from each subgroup in the medical arm are not separated. The sample size is too small to detect any difference among such number of groups
Kanhai 1989	Includes both second and third trimester fetal deaths.
Lippert 1978	Second and third trimester fetal deaths. Not obviously randomised
Machtlinger 2002	Abstract. Appears to include both non-viable pregnancies and miscarriages. Await full report
Machtlinger 2004	Abstract. Appears to include both non-viable pregnancies and miscarriages. Await full report
Makhlouf 2003	Not clear from paper if all pregnancies complicated by fetal death. Seeking clarification from authors
Martin 1965	Allocation based on alternation, not randomisation. Alternation violated
Nakintu 2001	Both second and third trimester fetal deaths. Seeking separate data from author
Ngai 2001	Includes data on women with both non-viable pregnancies and incomplete miscarriages. If these data can be separated by the researchers, these data may be included in the future
Nuthalapaty 2004	Abstract. Clinical indications for termination not described
Nuutila 1997	Trial included women with fetal malformations and maternal indications for pregnancy termination between 12 and 24 weeks, as well as pregnancies with fetal death. Data will be included for the latter if these can be obtained from the authors

(Continued)

Owen 1999	Trial included women with fetal malformations and maternal indications for pregnancy termination between 16 and 24 weeks, as well as pregnancies with fetal death. Data will be included for the latter if these can be obtained from the authors
Paraskevaides 1992	Small study of 16 women "randomised" to surgical evacuation or prostaglandin F2alpha or Trilostane treatment. No details about clinical presentation or ultrasound and clinical findings, but from abstract includes both women with non-viable pregnancies and incomplete miscarriage
Perry 1999	Excluded women with fetal deaths.
Piotrowski 1979	Not clear that this was a randomised trial.
Pongsatha 2004	Trial excluded women with fetal deaths.
Ramsey 2004	Trial included women with other reasons for pregnancy termination, as well as pregnancies with fetal death. Data will be included for the latter if these can be obtained from the authors
Roy 2003	Abstract. Not clear if fetal death included as indication for termination
Salamalekis 1990	Abstract only. Treatment allocation by alternation, not by randomisation
Su 2005	Termination of pregnancy for fetal anomalies, social reasons or maternal disease; not for non-viable pregnancies
Surita 1997	Abstract only. May include third trimester fetal deaths.
Thavarasah 1986	Unclear from paper but allocation may have been by alternation. Will seek clarification from authors
Toppozada 1994	Includes third trimester fetal deaths.
Yapar 1996	Includes indications for termination other than fetal death. High degree of protocol violation (60/400). Results not presented as intention-to-treat
Zhang 2005	Includes both non-viable pregnancies and miscarriages. Data will be included for the former if these can be obtained from the authors

hCG: human chorionic gonadotropin

POC: products of conception

RCT: randomised controlled trial

Characteristics of ongoing studies [ordered by study ID]

Australia 2000

Trial name or title	Misoprostol for the medical management of miscarriage (randomised controlled trial)
Methods	
Participants	Women with a spontaneous incomplete or missed abortion up to and including 12 weeks' gestation
Interventions	Drugs and medications; surgery.
Outcomes	Primary outcome measures: (1) need for evacuation of retained products of conception in theatre within 1 month of the intervention; (2) infection requiring antibiotics; (3) change in haemoglobin from pre-intervention to 72 hours and 4 weeks post-intervention. Secondary outcomes: (1) side-effects e.g. nausea, vomiting, diarrhoea, abdominal pain; (2) analgesia required in hospital - narcotics or oral; (3) antiemetics required in hospital; (4) duration of bleeding; (5) complication rates based on ethnic extraction.
Starting date	April 1999.
Contact information	Katharine Louey 7 Eagle 11 Bridge Rd Mackay Qld 4740 Australia Tel: + 61 7 49531817 e-mail: klouey@mrbean.net.au
Notes	

UK 1999

Trial name or title	Miscarriage Treatment Study (MIST).
Methods	
Participants	Women with confirmed missed and incomplete miscarriage attending early pregnancy clinics, with gestational dates less than 91 days (13 weeks), or uterine cavity volume being less than 91 days. Other inclusion criteria: women can understand English, willing to be randomised to any treatment group and there is uncertainty about the best management. Exclusion criteria: severe haemorrhage, severe pain, pyrexia above 37.5 C, severe asthma, blood dyscrasia, diabetes, current anticoagulants or corticosteroid therapy, twin or higher order pregnancy, use of prostaglandin contra-indicated, cannot understand written English, refuses written consent
Interventions	Expectant group: sent home with written advice and analgesia. Medical group: will be admitted (immediately if incomplete miscarriage and 24-48 hours after oral administration of mifepristone if missed miscarriage) and vaginal misoprostol will be administered. Surgical group: as is current practice

Outcomes	<p>Primary outcome measure:</p> <p>(1) gynecological infection within 14 days of the confirmation of miscarriage by TVS. Secondary outcome measures:</p> <p>(1) treatment with antibiotic (within 14 days and 6 weeks);</p> <p>(2) days of pain, days of vaginal bleeding, time off work, return to usual daily activities;</p> <p>(3) haemoglobin and packed cell volume (both at 10-14 days), unplanned surgical ERPC or other admission (within 14 days or 6 weeks);</p> <p>(4) complete evacuation of the uterus by TVS (at 10-14 days);</p> <p>(5) depression anxiety and general health (at 6 weeks).</p>
Starting date	
Contact information	<p>Miss Lohanna Trinder MIST Study Co-ordinating Center Dept of Women's Health Cotswold Center Southmead Hospital Bristol BS10 5NB UK Tel: 01179595186 FAX 01179595178 e-mail: jo.trinder@msn.com.uk</p>
Notes	

ERPC: evacuation of retained products of conception

TVS: transvaginal sonography

DATA AND ANALYSES

Comparison 1. Vaginal misoprostol versus placebo

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Complete miscarriage < 24 hours after treatment	2	138	Risk Ratio (M-H, Fixed, 95% CI)	4.73 [2.70, 8.28]
2 Complete miscarriage < 48 hours	2	84	Risk Ratio (M-H, Fixed, 95% CI)	5.74 [2.70, 12.19]
3 Complete miscarriage without ERPC day 7	1	83	Risk Ratio (M-H, Fixed, 95% CI)	2.99 [1.80, 4.99]
4 Uterine curettage	2	104	Risk Ratio (M-H, Fixed, 95% CI)	0.40 [0.26, 0.60]
5 Opiates for pain relief	1	84	Risk Ratio (M-H, Fixed, 95% CI)	5.0 [0.25, 101.11]
6 Blood transfusion	1	84	Risk Ratio (M-H, Fixed, 95% CI)	0.2 [0.01, 4.04]
7 Haemoglobin difference > 10 g/L	1	50	Risk Ratio (M-H, Fixed, 95% CI)	1.25 [0.38, 4.12]
8 Nausea	2	88	Risk Ratio (M-H, Fixed, 95% CI)	1.38 [0.43, 4.40]
9 Diarrhoea	2	88	Risk Ratio (M-H, Fixed, 95% CI)	2.21 [0.35, 14.06]
10 Fever	1	54	Risk Ratio (M-H, Fixed, 95% CI)	9.0 [0.51, 159.43]
11 Uterine perforation	1	84	Risk Ratio (M-H, Fixed, 95% CI)	0.33 [0.01, 7.96]
12 Vaginal bleeding 2 weeks after treatment	1	32	Risk Ratio (M-H, Fixed, 95% CI)	1.0 [0.41, 2.45]
13 Satisfaction with treatment	1	32	Risk Ratio (M-H, Fixed, 95% CI)	1.17 [0.83, 1.64]

Comparison 2. Vaginal misoprostol versus surgical evacuation of uterus

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Surgical evacuation of uterus	3	254	Risk Ratio (M-H, Fixed, 95% CI)	0.42 [0.34, 0.52]
2 Post-treatment haematocrit (%)	1	50	Mean Difference (IV, Fixed, 95% CI)	-1.40 [-3.51, 0.71]
3 Nausea	1	154	Risk Ratio (M-H, Fixed, 95% CI)	21.85 [1.31, 364.37]
4 Pain relief	1	154	Risk Ratio (M-H, Fixed, 95% CI)	1.42 [0.82, 2.46]
5 Diarrhoea	1	154	Risk Ratio (M-H, Fixed, 95% CI)	40.85 [2.52, 662.57]
6 Uterine perforation	1	154	Risk Ratio (M-H, Fixed, 95% CI)	0.32 [0.01, 7.65]
7 Asherman syndrome	1	154	Risk Ratio (M-H, Fixed, 95% CI)	0.32 [0.01, 7.65]

Comparison 3. Vaginal misoprostol versus vaginal gemeprost

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Miscarriage < 24 hours	1	50	Risk Ratio (M-H, Fixed, 95% CI)	1.24 [0.90, 1.70]
1.2 Gestation 13-24 weeks	1	50	Risk Ratio (M-H, Fixed, 95% CI)	1.24 [0.90, 1.70]
2 Temperature > 38 degrees C	1	50	Risk Ratio (M-H, Fixed, 95% CI)	1.5 [0.27, 8.22]
2.2 Gestation 13-24 weeks	1	50	Risk Ratio (M-H, Fixed, 95% CI)	1.5 [0.27, 8.22]
3 Vomiting	1	50	Risk Ratio (M-H, Fixed, 95% CI)	3.0 [0.13, 70.30]
3.2 Gestation 13-24 weeks	1	50	Risk Ratio (M-H, Fixed, 95% CI)	3.0 [0.13, 70.30]
4 Diarrhoea	1	50	Risk Ratio (M-H, Fixed, 95% CI)	0.14 [0.01, 2.63]
4.2 Gestation 13-24 weeks	1	50	Risk Ratio (M-H, Fixed, 95% CI)	0.14 [0.01, 2.63]
5 Opiate analgesia	1	50	Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
5.2 Gestation 13-24 weeks	1	50	Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]

Comparison 4. Vaginal misoprostol versus vaginal prostaglandin E1/2

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Surgical evacuation	1	60	Risk Ratio (M-H, Fixed, 95% CI)	0.39 [0.21, 0.72]
2 Blood transfusion	1	60	Risk Ratio (M-H, Fixed, 95% CI)	6.07 [0.30, 121.33]
3 Hospital stay (days)	1	60	Mean Difference (IV, Fixed, 95% CI)	-2.38 [-3.36, -1.40]
4 Complete miscarriage	1	65	Risk Ratio (M-H, Fixed, 95% CI)	1.44 [1.06, 1.96]
5 Nausea	1	65	Risk Ratio (M-H, Fixed, 95% CI)	1.03 [0.28, 3.78]

Comparison 5. Vaginal misoprostol lower versus higher dose regimens

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Miscarriage	3	247	Risk Ratio (M-H, Fixed, 95% CI)	0.85 [0.72, 1.00]
1.1 400 mcg versus 600 mcg	1	100	Risk Ratio (M-H, Fixed, 95% CI)	1.05 [0.87, 1.26]
1.2 400 mcg versus 800 mcg	1	33	Risk Ratio (M-H, Fixed, 95% CI)	0.77 [0.45, 1.31]
1.3 600 mcg versus 800 mcg	1	114	Risk Ratio (M-H, Fixed, 95% CI)	0.67 [0.48, 0.93]
2 Fever	2	214	Risk Ratio (M-H, Fixed, 95% CI)	0.73 [0.41, 1.30]
3 Nausea	2	214	Risk Ratio (M-H, Fixed, 95% CI)	0.67 [0.31, 1.41]
4 Diarrhoea	2	214	Risk Ratio (M-H, Fixed, 95% CI)	0.54 [0.15, 1.91]

Comparison 6. Vaginal misoprostol wet versus dry vaginal preparations

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Miscarriage < 3 days	1	80	Risk Ratio (M-H, Fixed, 95% CI)	1.14 [0.85, 1.54]
2 Miscarriage < 8 days	1	80	Risk Ratio (M-H, Fixed, 95% CI)	1.04 [0.84, 1.29]
3 Miscarriage < 15 days	1	80	Risk Ratio (M-H, Fixed, 95% CI)	0.92 [0.78, 1.10]
4 Miscarriage < 30 days	1	80	Risk Ratio (M-H, Fixed, 95% CI)	0.95 [0.79, 1.14]
5 Diarrhoea < 48 hours after treatment	1	77	Risk Ratio (M-H, Fixed, 95% CI)	1.75 [0.89, 3.42]
6 Chills < 48 hours of treatment	1	77	Risk Ratio (M-H, Fixed, 95% CI)	1.36 [0.94, 1.98]
7 Vomiting < 48 hours of treatment	1	77	Risk Ratio (M-H, Fixed, 95% CI)	0.93 [0.33, 2.62]
8 Would wish/probably wish same treatment in future nonviable pregnancy	1	73	Risk Ratio (M-H, Fixed, 95% CI)	1.18 [0.93, 1.49]

Comparison 7. Vaginal misoprostol + methotrexate versus vaginal misoprostol alone

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Miscarriage not complete	1	21	Risk Ratio (M-H, Fixed, 95% CI)	0.26 [0.01, 5.65]
4 Additional surgical evacuation	1	21	Risk Ratio (M-H, Fixed, 95% CI)	0.26 [0.01, 5.65]
5 Haemorrhage	1	21	Risk Ratio (M-H, Fixed, 95% CI)	2.31 [0.10, 50.85]
6 Pain relief	1	21	Risk Ratio (M-H, Fixed, 95% CI)	0.75 [0.25, 2.22]

Comparison 8. Vaginal misoprostol plus laminaria tents versus vaginal misoprostol alone

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Miscarriage < 24 hours	1	38	Risk Ratio (M-H, Fixed, 95% CI)	0.90 [0.65, 1.25]
2 Miscarriage < 48 hours	1	38	Risk Ratio (M-H, Fixed, 95% CI)	1.07 [0.88, 1.29]

Comparison 9. Oral misoprostol versus vaginal misoprostol

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Complete miscarriage	2	218	Risk Ratio (M-H, Fixed, 95% CI)	0.90 [0.82, 0.99]
1.1 400 mcg oral misoprostol versus 800 mcg vaginal misoprostol	1	20	Risk Ratio (M-H, Fixed, 95% CI)	0.29 [0.10, 0.79]
1.2 800 mcg oral misoprostol versus 800 mcg vaginal misoprostol	1	198	Risk Ratio (M-H, Fixed, 95% CI)	0.96 [0.88, 1.05]
2 Vomiting	1	190	Risk Ratio (M-H, Fixed, 95% CI)	0.29 [0.10, 0.84]
9 Nausea	1	20	Risk Ratio (M-H, Fixed, 95% CI)	0.80 [0.37, 1.74]
10 Diarrhoea	2	210	Risk Ratio (M-H, Fixed, 95% CI)	1.05 [0.67, 1.66]
12 Pain (visual analogue scale)	1	18	Mean Difference (IV, Fixed, 95% CI)	-1.90 [-4.82, 1.02]
13 Fever	1	190	Risk Ratio (M-H, Fixed, 95% CI)	1.0 [0.36, 2.74]
14 Women's satisfaction with treatment	1	198	Risk Ratio (M-H, Fixed, 95% CI)	0.96 [0.86, 1.06]
15 Time to delivery (hours)	1	70	Mean Difference (IV, Fixed, 95% CI)	4.1 [2.64, 5.56]
16 Oxytocin infusion	1	70	Risk Ratio (M-H, Fixed, 95% CI)	2.75 [0.97, 7.81]
17 Manual removal of placenta	1	70	Risk Ratio (M-H, Fixed, 95% CI)	4.5 [1.05, 19.35]

Comparison 10. Oral misoprostol + mifepristone versus expectant management

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Empty uterine cavity at day 5	1	122	Risk Ratio (M-H, Fixed, 95% CI)	1.08 [0.90, 1.30]
2 Urgent surgical evacuation for bleeding	1	122	Risk Ratio (M-H, Fixed, 95% CI)	0.34 [0.01, 8.29]
3 Pelvic inflammatory disease	1	122	Risk Ratio (M-H, Fixed, 95% CI)	0.52 [0.05, 5.55]
4 Pain (visual analogue scale day 5)	1	122	Mean Difference (IV, Fixed, 95% CI)	4.10 [-5.92, 14.12]
5 Sick leave (days)	1	122	Mean Difference (IV, Fixed, 95% CI)	1.8 [0.63, 2.97]
6 Bleeding (days)	1	122	Mean Difference (IV, Fixed, 95% CI)	0.70 [-0.43, 1.83]
7 Satisfaction with treatment (visual analogue scale day 14)	1	122	Mean Difference (IV, Fixed, 95% CI)	3.40 [-5.54, 12.34]

Comparison 11. Sublingual misoprostol versus vaginal misoprostol

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Complete miscarriage	1	80	Risk Ratio (M-H, Fixed, 95% CI)	1.0 [0.85, 1.18]
2 Nausea	1	80	Risk Ratio (M-H, Fixed, 95% CI)	1.2 [0.80, 1.79]
3 Vomiting	1	80	Risk Ratio (M-H, Fixed, 95% CI)	0.78 [0.32, 1.88]
4 Diarrhoea	1	80	Risk Ratio (M-H, Fixed, 95% CI)	2.55 [1.48, 4.38]
5 Haemoglobin day 43	1	80	Mean Difference (IV, Fixed, 95% CI)	0.10 [-0.38, 0.58]
6 Intolerable pain	1	80	Risk Ratio (M-H, Fixed, 95% CI)	0.75 [0.29, 1.97]
7 Satisfied with treatment	1	77	Risk Ratio (M-H, Fixed, 95% CI)	0.99 [0.79, 1.25]

Comparison 12. Mifepristone versus placebo

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Miscarriage < 48 hours	1	46	Risk Ratio (M-H, Fixed, 95% CI)	5.0 [0.25, 98.75]
2 Miscarriage < 3 days	1	46	Risk Ratio (M-H, Fixed, 95% CI)	19.0 [1.17, 308.40]
3 Miscarriage < 4 days	1	46	Risk Ratio (M-H, Fixed, 95% CI)	14.0 [2.00, 97.88]
4 Miscarriage < 5 days	1	46	Risk Ratio (M-H, Fixed, 95% CI)	9.5 [2.49, 36.19]
5 Vaginal bleeding before day 5	1	44	Risk Ratio (M-H, Fixed, 95% CI)	3.92 [1.89, 8.10]
6 Pain before day 5	1	44	Risk Ratio (M-H, Fixed, 95% CI)	2.19 [0.93, 5.17]

Comparison 13. Vaginal gemeprost versus surgical evacuation of uterus

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Surgical evacuation	1	87	Risk Ratio (M-H, Fixed, 95% CI)	0.24 [0.14, 0.41]
2 Perforation of uterus	1	87	Risk Ratio (M-H, Fixed, 95% CI)	0.20 [0.01, 4.14]
3 Nausea	1	87	Risk Ratio (M-H, Fixed, 95% CI)	1.79 [0.56, 5.68]

Analysis 1.1. Comparison 1 Vaginal misoprostol versus placebo, Outcome 1 Complete miscarriage < 24 hours after treatment.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 1 Vaginal misoprostol versus placebo

Outcome: 1 Complete miscarriage < 24 hours after treatment

Analysis 1.2. Comparison 1 Vaginal misoprostol versus placebo, Outcome 2 Complete miscarriage < 48 hours.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 1 Vaginal misoprostol versus placebo

Outcome: 2 Complete miscarriage < 48 hours

Analysis 1.3. Comparison 1 Vaginal misoprostol versus placebo, Outcome 3 Complete miscarriage without ERPC day 7.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 1 Vaginal misoprostol versus placebo

Outcome: 3 Complete miscarriage without ERPC day 7

Analysis 1.4. Comparison 1 Vaginal misoprostol versus placebo, Outcome 4 Uterine curettage.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 1 Vaginal misoprostol versus placebo

Outcome: 4 Uterine curettage

Analysis 1.5. Comparison 1 Vaginal misoprostol versus placebo, Outcome 5 Opiates for pain relief.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 1 Vaginal misoprostol versus placebo

Outcome: 5 Opiates for pain relief

Analysis 1.6. Comparison 1 Vaginal misoprostol versus placebo, Outcome 6 Blood transfusion.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 1 Vaginal misoprostol versus placebo

Outcome: 6 Blood transfusion

Analysis 1.7. Comparison 1 Vaginal misoprostol versus placebo, Outcome 7 Haemoglobin difference > 10 g/L.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 1 Vaginal misoprostol versus placebo

Outcome: 7 Haemoglobin difference > 10 g/L

Analysis 1.8. Comparison 1 Vaginal misoprostol versus placebo, Outcome 8 Nausea.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 1 Vaginal misoprostol versus placebo

Outcome: 8 Nausea

Analysis 1.9. Comparison 1 Vaginal misoprostol versus placebo, Outcome 9 Diarrhoea.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 1 Vaginal misoprostol versus placebo

Outcome: 9 Diarrhoea

Analysis 1.10. Comparison 1 Vaginal misoprostol versus placebo, Outcome 10 Fever.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 1 Vaginal misoprostol versus placebo

Outcome: 10 Fever

Analysis 1.11. Comparison 1 Vaginal misoprostol versus placebo, Outcome 11 Uterine perforation.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 1 Vaginal misoprostol versus placebo

Outcome: 11 Uterine perforation

Analysis 1.12. Comparison 1 Vaginal misoprostol versus placebo, Outcome 12 Vaginal bleeding 2 weeks after treatment.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 1 Vaginal misoprostol versus placebo

Outcome: 12 Vaginal bleeding 2 weeks after treatment

Analysis 1.13. Comparison 1 Vaginal misoprostol versus placebo, Outcome 13 Satisfaction with treatment.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 1 Vaginal misoprostol versus placebo

Outcome: 13 Satisfaction with treatment

Analysis 2.1. Comparison 2 Vaginal misoprostol versus surgical evacuation of uterus, Outcome 1 Surgical evacuation of uterus.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 2 Vaginal misoprostol versus surgical evacuation of uterus

Outcome: 1 Surgical evacuation of uterus

Analysis 2.2. Comparison 2 Vaginal misoprostol versus surgical evacuation of uterus, Outcome 2 Post-treatment haematocrit (%).

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 2 Vaginal misoprostol versus surgical evacuation of uterus

Outcome: 2 Post-treatment haematocrit (%)

Analysis 2.3. Comparison 2 Vaginal misoprostol versus surgical evacuation of uterus, Outcome 3 Nausea.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 2 Vaginal misoprostol versus surgical evacuation of uterus

Outcome: 3 Nausea

Analysis 2.4. Comparison 2 Vaginal misoprostol versus surgical evacuation of uterus, Outcome 4 Pain relief.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 2 Vaginal misoprostol versus surgical evacuation of uterus

Outcome: 4 Pain relief

Analysis 2.5. Comparison 2 Vaginal misoprostol versus surgical evacuation of uterus, Outcome 5 Diarrhoea.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 2 Vaginal misoprostol versus surgical evacuation of uterus

Outcome: 5 Diarrhoea

Analysis 2.6. Comparison 2 Vaginal misoprostol versus surgical evacuation of uterus, Outcome 6 Uterine perforation.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 2 Vaginal misoprostol versus surgical evacuation of uterus

Outcome: 6 Uterine perforation

Analysis 2.7. Comparison 2 Vaginal misoprostol versus surgical evacuation of uterus, Outcome 7 Asherman syndrome.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 2 Vaginal misoprostol versus surgical evacuation of uterus

Outcome: 7 Asherman syndrome

Analysis 3.1. Comparison 3 Vaginal misoprostol versus vaginal gemeprost, Outcome 1 Miscarriage < 24 hours.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 3 Vaginal misoprostol versus vaginal gemeprost

Outcome: 1 Miscarriage < 24 hours

Analysis 3.2. Comparison 3 Vaginal misoprostol versus vaginal gemeprost, Outcome 2 Temperature > 38 degrees C.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 3 Vaginal misoprostol versus vaginal gemeprost

Outcome: 2 Temperature > 38 degrees C

Analysis 3.3. Comparison 3 Vaginal misoprostol versus vaginal gemeprost, Outcome 3 Vomiting.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 3 Vaginal misoprostol versus vaginal gemeprost

Outcome: 3 Vomiting

Analysis 3.4. Comparison 3 Vaginal misoprostol versus vaginal gemeprost, Outcome 4 Diarrhoea.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 3 Vaginal misoprostol versus vaginal gemeprost

Outcome: 4 Diarrhoea

Analysis 3.5. Comparison 3 Vaginal misoprostol versus vaginal gemeprost, Outcome 5 Opiate analgesia.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 3 Vaginal misoprostol versus vaginal gemeprost

Outcome: 5 Opiate analgesia

Analysis 4.1. Comparison 4 Vaginal misoprostol versus vaginal prostaglandin E1/2, Outcome 1 Surgical evacuation.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 4 Vaginal misoprostol versus vaginal prostaglandin E1/2

Outcome: 1 Surgical evacuation

Analysis 4.2. Comparison 4 Vaginal misoprostol versus vaginal prostaglandin E1/2, Outcome 2 Blood transfusion.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 4 Vaginal misoprostol versus vaginal prostaglandin E1/2

Outcome: 2 Blood transfusion

Analysis 4.3. Comparison 4 Vaginal misoprostol versus vaginal prostaglandin E1/2, Outcome 3 Hospital stay (days).

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 4 Vaginal misoprostol versus vaginal prostaglandin E1/2

Outcome: 3 Hospital stay (days)

Analysis 4.4. Comparison 4 Vaginal misoprostol versus vaginal prostaglandin E1/2, Outcome 4 Complete miscarriage.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 4 Vaginal misoprostol versus vaginal prostaglandin E1/2

Outcome: 4 Complete miscarriage

Analysis 4.5. Comparison 4 Vaginal misoprostol versus vaginal prostaglandin E1/2, Outcome 5 Nausea.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 4 Vaginal misoprostol versus vaginal prostaglandin E1/2

Outcome: 5 Nausea

Analysis 5.1. Comparison 5 Vaginal misoprostol lower versus higher dose regimens, Outcome 1 Miscarriage.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 5 Vaginal misoprostol lower versus higher dose regimens

Outcome: 1 Miscarriage

(Continued ...)

Analysis 5.2. Comparison 5 Vaginal misoprostol lower versus higher dose regimens, Outcome 2 Fever.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 5 Vaginal misoprostol lower versus higher dose regimens

Outcome: 2 Fever

Analysis 5.3. Comparison 5 Vaginal misoprostol lower versus higher dose regimens, Outcome 3 Nausea.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 5 Vaginal misoprostol lower versus higher dose regimens

Outcome: 3 Nausea

Analysis 5.4. Comparison 5 Vaginal misoprostol lower versus higher dose regimens, Outcome 4 Diarrhoea.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 5 Vaginal misoprostol lower versus higher dose regimens

Outcome: 4 Diarrhoea

Analysis 6.1. Comparison 6 Vaginal misoprostol wet versus dry vaginal preparations, Outcome 1 Miscarriage < 3 days.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 6 Vaginal misoprostol wet versus dry vaginal preparations

Outcome: 1 Miscarriage < 3 days

Analysis 6.2. Comparison 6 Vaginal misoprostol wet versus dry vaginal preparations, Outcome 2 Miscarriage < 8 days.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 6 Vaginal misoprostol wet versus dry vaginal preparations

Outcome: 2 Miscarriage < 8 days

Analysis 6.3. Comparison 6 Vaginal misoprostol wet versus dry vaginal preparations, Outcome 3 Miscarriage < 15 days.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 6 Vaginal misoprostol wet versus dry vaginal preparations

Outcome: 3 Miscarriage < 15 days

Analysis 6.4. Comparison 6 Vaginal misoprostol wet versus dry vaginal preparations, Outcome 4 Miscarriage < 30 days.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 6 Vaginal misoprostol wet versus dry vaginal preparations

Outcome: 4 Miscarriage < 30 days

Analysis 6.5. Comparison 6 Vaginal misoprostol wet versus dry vaginal preparations, Outcome 5 Diarrhoea < 48 hours after treatment.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 6 Vaginal misoprostol wet versus dry vaginal preparations

Outcome: 5 Diarrhoea < 48 hours after treatment

Analysis 6.6. Comparison 6 Vaginal misoprostol wet versus dry vaginal preparations, Outcome 6 Chills < 48 hours of treatment.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 6 Vaginal misoprostol wet versus dry vaginal preparations

Outcome: 6 Chills < 48 hours of treatment

Analysis 6.7. Comparison 6 Vaginal misoprostol wet versus dry vaginal preparations, Outcome 7 Vomiting < 48 hours of treatment.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 6 Vaginal misoprostol wet versus dry vaginal preparations

Outcome: 7 Vomiting < 48 hours of treatment

Analysis 6.8. Comparison 6 Vaginal misoprostol wet versus dry vaginal preparations, Outcome 8 Would wish/probably wish same treatment in future nonviable pregnancy.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 6 Vaginal misoprostol wet versus dry vaginal preparations

Outcome: 8 Would wish/probably wish same treatment in future nonviable pregnancy

Analysis 7.1. Comparison 7 Vaginal misoprostol + methotrexate versus vaginal misoprostol alone, Outcome 1 Miscarriage not complete.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 7 Vaginal misoprostol + methotrexate versus vaginal misoprostol alone

Outcome: 1 Miscarriage not complete

Analysis 7.4. Comparison 7 Vaginal misoprostol + methotrexate versus vaginal misoprostol alone, Outcome 4 Additional surgical evacuation.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 7 Vaginal misoprostol + methotrexate versus vaginal misoprostol alone

Outcome: 4 Additional surgical evacuation

Analysis 7.5. Comparison 7 Vaginal misoprostol + methotrexate versus vaginal misoprostol alone, Outcome 5 Haemorrhage.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 7 Vaginal misoprostol + methotrexate versus vaginal misoprostol alone

Outcome: 5 Haemorrhage

Analysis 7.6. Comparison 7 Vaginal misoprostol + methotrexate versus vaginal misoprostol alone, Outcome 6 Pain relief.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 7 Vaginal misoprostol + methotrexate versus vaginal misoprostol alone

Outcome: 6 Pain relief

Analysis 8.1. Comparison 8 Vaginal misoprostol plus laminaria tents versus vaginal misoprostol alone, Outcome 1 Miscarriage < 24 hours.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 8 Vaginal misoprostol plus laminaria tents versus vaginal misoprostol alone

Outcome: 1 Miscarriage < 24 hours

Analysis 8.2. Comparison 8 Vaginal misoprostol plus laminaria tents versus vaginal misoprostol alone, Outcome 2 Miscarriage < 48 hours.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 8 Vaginal misoprostol plus laminaria tents versus vaginal misoprostol alone

Outcome: 2 Miscarriage < 48 hours

Analysis 9.1. Comparison 9 Oral misoprostol versus vaginal misoprostol, Outcome 1 Complete miscarriage.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 9 Oral misoprostol versus vaginal misoprostol

Outcome: 1 Complete miscarriage

Analysis 9.2. Comparison 9 Oral misoprostol versus vaginal misoprostol, Outcome 2 Vomiting.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 9 Oral misoprostol versus vaginal misoprostol

Outcome: 2 Vomiting

Analysis 9.9. Comparison 9 Oral misoprostol versus vaginal misoprostol, Outcome 9 Nausea.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 9 Oral misoprostol versus vaginal misoprostol

Outcome: 9 Nausea

Analysis 9.10. Comparison 9 Oral misoprostol versus vaginal misoprostol, Outcome 10 Diarrhoea.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 9 Oral misoprostol versus vaginal misoprostol

Outcome: 10 Diarrhoea

Analysis 9.12. Comparison 9 Oral misoprostol versus vaginal misoprostol, Outcome 12 Pain (visual analogue scale).

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 9 Oral misoprostol versus vaginal misoprostol

Outcome: 12 Pain (visual analogue scale)

Analysis 9.13. Comparison 9 Oral misoprostol versus vaginal misoprostol, Outcome 13 Fever.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 9 Oral misoprostol versus vaginal misoprostol

Outcome: 13 Fever

Analysis 9.14. Comparison 9 Oral misoprostol versus vaginal misoprostol, Outcome 14 Women's satisfaction with treatment.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 9 Oral misoprostol versus vaginal misoprostol

Outcome: 14 Women's satisfaction with treatment

Analysis 9.15. Comparison 9 Oral misoprostol versus vaginal misoprostol, Outcome 15 Time to delivery (hours).

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 9 Oral misoprostol versus vaginal misoprostol

Outcome: 15 Time to delivery (hours)

Analysis 9.16. Comparison 9 Oral misoprostol versus vaginal misoprostol, Outcome 16 Oxytocin infusion.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 9 Oral misoprostol versus vaginal misoprostol

Outcome: 16 Oxytocin infusion

Analysis 9.17. Comparison 9 Oral misoprostol versus vaginal misoprostol, Outcome 17 Manual removal of placenta.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 9 Oral misoprostol versus vaginal misoprostol

Outcome: 17 Manual removal of placenta

Analysis 10.1. Comparison 10 Oral misoprostol + mifepristone versus expectant management, Outcome 1 Empty uterine cavity at day 5.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 10 Oral misoprostol + mifepristone versus expectant management

Outcome: 1 Empty uterine cavity at day 5

Analysis 10.2. Comparison 10 Oral misoprostol + mifepristone versus expectant management, Outcome 2 Urgent surgical evacuation for bleeding.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 10 Oral misoprostol + mifepristone versus expectant management

Outcome: 2 Urgent surgical evacuation for bleeding

Analysis 10.3. Comparison 10 Oral misoprostol + mifepristone versus expectant management, Outcome 3 Pelvic inflammatory disease.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 10 Oral misoprostol + mifepristone versus expectant management

Outcome: 3 Pelvic inflammatory disease

Analysis 10.4. Comparison 10 Oral misoprostol + mifepristone versus expectant management, Outcome 4 Pain (visual analogue scale day 5).

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 10 Oral misoprostol + mifepristone versus expectant management

Outcome: 4 Pain (visual analogue scale day 5)

Analysis 10.5. Comparison 10 Oral misoprostol + mifepristone versus expectant management, Outcome 5 Sick leave (days).

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 10 Oral misoprostol + mifepristone versus expectant management

Outcome: 5 Sick leave (days)

Analysis 10.6. Comparison 10 Oral misoprostol + mifepristone versus expectant management, Outcome 6 Bleeding (days).

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 10 Oral misoprostol + mifepristone versus expectant management

Outcome: 6 Bleeding (days)

Analysis 10.7. Comparison 10 Oral misoprostol + mifepristone versus expectant management, Outcome 7 Satisfaction with treatment (visual analogue scale day 14).

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 10 Oral misoprostol + mifepristone versus expectant management

Outcome: 7 Satisfaction with treatment (visual analogue scale day 14)

Analysis 11.1. Comparison 11 Sublingual misoprostol versus vaginal misoprostol, Outcome 1 Complete miscarriage.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 11 Sublingual misoprostol versus vaginal misoprostol

Outcome: 1 Complete miscarriage

Analysis 11.2. Comparison 11 Sublingual misoprostol versus vaginal misoprostol, Outcome 2 Nausea.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 11 Sublingual misoprostol versus vaginal misoprostol

Outcome: 2 Nausea

Analysis 11.3. Comparison 11 Sublingual misoprostol versus vaginal misoprostol, Outcome 3 Vomiting.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 11 Sublingual misoprostol versus vaginal misoprostol

Outcome: 3 Vomiting

Analysis 11.4. Comparison 11 Sublingual misoprostol versus vaginal misoprostol, Outcome 4 Diarrhoea.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 11 Sublingual misoprostol versus vaginal misoprostol

Outcome: 4 Diarrhoea

Analysis 11.5. Comparison 11 Sublingual misoprostol versus vaginal misoprostol, Outcome 5 Haemoglobin day 43.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 11 Sublingual misoprostol versus vaginal misoprostol

Outcome: 5 Haemoglobin day 43

Analysis 11.6. Comparison 11 Sublingual misoprostol versus vaginal misoprostol, Outcome 6 Intolerable pain.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 11 Sublingual misoprostol versus vaginal misoprostol

Outcome: 6 Intolerable pain

Analysis 11.7. Comparison 11 Sublingual misoprostol versus vaginal misoprostol, Outcome 7 Satisfied with treatment.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 11 Sublingual misoprostol versus vaginal misoprostol

Outcome: 7 Satisfied with treatment

Analysis 12.1. Comparison 12 Mifepristone versus placebo, Outcome 1 Miscarriage < 48 hours.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 12 Mifepristone versus placebo

Outcome: 1 Miscarriage < 48 hours

Analysis 12.2. Comparison 12 Mifepristone versus placebo, Outcome 2 Miscarriage < 3 days.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 12 Mifepristone versus placebo

Outcome: 2 Miscarriage < 3 days

Analysis 12.3. Comparison 12 Mifepristone versus placebo, Outcome 3 Miscarriage < 4 days.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 12 Mifepristone versus placebo

Outcome: 3 Miscarriage < 4 days

Analysis 12.4. Comparison 12 Mifepristone versus placebo, Outcome 4 Miscarriage < 5 days.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 12 Mifepristone versus placebo

Outcome: 4 Miscarriage < 5 days

Analysis 12.5. Comparison 12 Mifepristone versus placebo, Outcome 5 Vaginal bleeding before day 5.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 12 Mifepristone versus placebo

Outcome: 5 Vaginal bleeding before day 5

Analysis 12.6. Comparison 12 Mifepristone versus placebo, Outcome 6 Pain before day 5.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 12 Mifepristone versus placebo

Outcome: 6 Pain before day 5

Analysis 13.1. Comparison 13 Vaginal gemeprost versus surgical evacuation of uterus, Outcome 1 Surgical evacuation.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 13 Vaginal gemeprost versus surgical evacuation of uterus

Outcome: 1 Surgical evacuation

Analysis 13.2. Comparison 13 Vaginal gemeprost versus surgical evacuation of uterus, Outcome 2 Perforation of uterus.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 13 Vaginal gemeprost versus surgical evacuation of uterus

Outcome: 2 Perforation of uterus

Analysis 13.3. Comparison 13 Vaginal gemeprost versus surgical evacuation of uterus, Outcome 3 Nausea.

Review: Medical treatment for early fetal death (less than 24 weeks)

Comparison: 13 Vaginal gemeprost versus surgical evacuation of uterus

Outcome: 3 Nausea

WHAT'S NEW

Last assessed as up-to-date: 24 April 2006.

Date	Event	Description
8 August 2012	Amended	Search updated. One hundred reports added to Studies awaiting classification

HISTORY

Protocol first published: Issue 3, 2000

Review first published: Issue 3, 2006

Date	Event	Description
18 September 2008	Amended	Converted to new review format.

CONTRIBUTIONS OF AUTHORS

Juan Vazquez: protocol development and revisions to first draft of review.

Martha Hickey: protocol development and revisions to the first draft of review.

Jim Neilson: supervision of protocol development; completion of first draft of review.

DECLARATIONS OF INTEREST

None known.

SOURCES OF SUPPORT

Internal sources

- America Arias Hospital, Havana, Cuba.
- The University of Liverpool, UK.

External sources

- HRP/WHO, Geneva, Switzerland.

DIFFERENCES BETWEEN PROTOCOL AND REVIEW

The protocol for this review aimed to include both trials for treatment of both ultrasound diagnosed non-viable pregnancies and incomplete miscarriage. For the reasons described in the review, two separate reviews now address these topics - thus, the change in title from 'Medical management for miscarriage'.

INDEX TERMS

Medical Subject Headings (MeSH)

*Fetal Death [diagnostic imaging]; Abortifacient Agents [*therapeutic use]; Abortion, Induced [*methods]; Administration, Intravaginal; Administration, Oral; Mifepristone [*therapeutic use]; Misoprostol [*therapeutic use]; Randomized Controlled Trials as Topic; Ultrasonography, Prenatal

MeSH check words

Female; Humans; Pregnancy