**Cochrane
Library**

Cochrane Database of Systematic Reviews

Computed tomography versus magnetic resonance imaging versus bone scintigraphy for clinically suspected scaphoid fractures in patients with negative plain radiographs (Review)

Mallee WH, Wang J, Poolman RW, Kloen P, Maas M, de Vet HCW, Doornberg JN

Mallee WH, Wang J, Poolman RW, Kloen P, Maas M, de Vet HCW, Doornberg JN.

Computed tomography versus magnetic resonance imaging versus bone scintigraphy for clinically suspected scaphoid fractures in patients with negative plain radiographs.

Cochrane Database of Systematic Reviews 2015, Issue 6. Art. No.: CD010023.

DOI: [10.1002/14651858.CD010023.pub2](https://doi.org/10.1002/14651858.CD010023.pub2).

www.cochranelibrary.com

Computed tomography versus magnetic resonance imaging versus bone scintigraphy for clinically suspected scaphoid fractures in patients with negative plain radiographs (Review)

Copyright © 2015 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

WILEY

TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
SUMMARY OF FINDINGS	4
BACKGROUND	7
OBJECTIVES	9
METHODS	9
RESULTS	11
Figure 1.	12
Figure 2.	14
Figure 3.	15
Figure 4.	16
Figure 5.	17
DISCUSSION	18
AUTHORS' CONCLUSIONS	19
ACKNOWLEDGEMENTS	19
REFERENCES	21
CHARACTERISTICS OF STUDIES	26
DATA	46
Test 1. CT.	46
Test 2. MRI.	46
Test 3. BS.	46
ADDITIONAL TABLES	46
APPENDICES	50
CONTRIBUTIONS OF AUTHORS	52
DECLARATIONS OF INTEREST	52
SOURCES OF SUPPORT	52
DIFFERENCES BETWEEN PROTOCOL AND REVIEW	52
INDEX TERMS	52

[Diagnostic Test Accuracy Review]

Computed tomography versus magnetic resonance imaging versus bone scintigraphy for clinically suspected scaphoid fractures in patients with negative plain radiographs

Wouter H Mallee¹, Junfeng Wang², Rudolf W Poolman³, Peter Kloen¹, Mario Maas⁴, Henrica CW de Vet⁵, Job N Doornberg¹

¹Department of Orthopaedic Surgery, Academic Medical Center, Amsterdam, Netherlands. ²Department of Clinical Epidemiology, Biostatistics and Bioinformatics, Academic Medical Center, Amsterdam, Netherlands. ³Department of Orthopaedic Surgery, Onze Lieve Vrouwe Gasthuis, Amsterdam, Netherlands. ⁴Department of Radiology, Academic Medical Center, Amsterdam, Netherlands. ⁵Department of Epidemiology and Biostatistics, EMGO Institute for Health and Care Research, VU University Medical Center, Amsterdam, Netherlands

Contact address: Wouter H Mallee, Department of Orthopaedic Surgery, Academic Medical Center, Secretariaat G4-Noord, Meibergdreef 9, Amsterdam, 1057 GB, Netherlands. woutermallee@hotmail.com.

Editorial group: Cochrane Bone, Joint and Muscle Trauma Group.

Publication status and date: New, published in Issue 6, 2015.

Citation: Mallee WH, Wang J, Poolman RW, Kloen P, Maas M, de Vet HCW, Doornberg JN. Computed tomography versus magnetic resonance imaging versus bone scintigraphy for clinically suspected scaphoid fractures in patients with negative plain radiographs. *Cochrane Database of Systematic Reviews* 2015, Issue 6. Art. No.: CD010023. DOI: [10.1002/14651858.CD010023.pub2](https://doi.org/10.1002/14651858.CD010023.pub2).

Copyright © 2015 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

In clinically suspected scaphoid fractures, early diagnosis reduces the risk of non-union and minimises loss in productivity resulting from unnecessary cast immobilisation. Since initial radiographs do not exclude the possibility of a fracture, additional imaging is needed. Computed tomography (CT), magnetic resonance imaging (MRI) and bone scintigraphy (BS) are widely used to establish a definitive diagnosis, but there is uncertainty about the most appropriate method.

Objectives

The primary aim of this study is to identify the most suitable diagnostic imaging strategy for identifying clinically suspected fractures of the scaphoid bone in patients with normal radiographs. Therefore we looked at the diagnostic performance characteristics of the most used imaging modalities for this purpose: computed tomography, magnetic resonance imaging and bone scintigraphy.

Search methods

In July 2012, we searched the Cochrane Register of Diagnostic Test Accuracy Studies, MEDLINE, EMBASE, the Database of Abstracts of Reviews of Effects, the Cochrane Central Register of Controlled Trials, the NHS Economic Evaluation Database. In September 2012, we searched MEDION, ARIF, Current Controlled Trials, the World Health Organization (WHO) International Clinical Trials Registry Platform, conference proceedings and reference lists of all articles.

Selection criteria

We included all prospective or retrospective studies involving a consecutive series of patients of all ages that evaluated the accuracy of BS, CT or MRI, or any combination of these, for diagnosing suspected scaphoid fractures. We considered the use of one or two index tests or six-week follow-up radiographs as adequate reference standards.

Data collection and analysis

Two review authors independently screened titles and abstracts and assessed full-text reports of potentially eligible studies. The same authors extracted data from full-text reports and assessed methodological quality using the QUADAS checklist. For each index test, estimates of sensitivity and specificity from each study were plotted in ROC space; and forest plots were constructed for visual examination of variation in test accuracy. We performed meta-analyses using the HSROC model to produce summary estimates of sensitivity and specificity.

Main results

We included 11 studies that looked at diagnostic accuracy of one or two index tests: four studies (277 suspected fractures) looked at CT, five studies (221 suspected fractures) looked at MRI and six studies (543 suspected fractures) looked at BS. Four of the studies made direct comparisons: two studies compared CT and MRI, one study compared CT and BS, and one study compared MRI and BS. Overall, the studies were of moderate to good quality, but relevant clinical information during evaluation of CT, MRI or BS was mostly unclear or unavailable.

As few studies made direct comparisons between tests with the same participants, our results are based on data from indirect comparisons, which means that these results are more susceptible to bias due to confounding. Nonetheless, the direct comparisons showed similar patterns of differences in sensitivity and specificity as for the pooled indirect comparisons.

Summary sensitivity and specificity of CT were 0.72 (95% confidence interval (CI) 0.36 to 0.92) and 0.99 (95% CI 0.71 to 1.00); for MRI, these were 0.88 (95% CI 0.64 to 0.97) and 1.00 (95% CI 0.38 to 1.00); for BS, these were 0.99 (95% CI 0.69 to 1.00) and 0.86 (95% CI 0.73 to 0.94). Indirect comparisons suggest that diagnostic accuracy of BS was significantly higher than CT and MRI; and CT and MRI have comparable diagnostic accuracy. The low prevalence of a true fracture among suspected fractures (median = 20%) means the lower specificity for BS is problematic. For example, in a cohort of 1000 patients, 112 will be over-treated when BS is used for diagnosis. If CT is used, only 8 will receive unnecessary treatment. In terms of missed fractures, BS will miss 2 fractures and CT will miss 56 fractures.

Authors' conclusions

Although quality of the included studies is moderate to good, findings are based on only 11 studies and the confidence intervals for the summary estimates are wide for all three tests. Well-designed direct comparison studies including CT, MRI and BS could give valuable additional information.

Bone scintigraphy is statistically the best diagnostic modality to establish a definitive diagnosis in clinically suspected fractures when radiographs appear normal. However, physicians must keep in mind that BS is more invasive than the other modalities, with safety issues due to level of radiation exposure, as well as diagnostic delay of at least 72 hours. The number of overtreated patients is substantially lower with CT and MRI.

Prior to performing comparative studies, there is a need to raise the initially detected prevalence of true fractures in order to reduce the effect of the relatively low specificity in daily practice. This can be achieved by improving clinical evaluation and initial radiographical assessment.

PLAIN LANGUAGE SUMMARY

Comparing different types of scan (CT, MRI, bone scan) for diagnosis of clinically suspected scaphoid fractures, when initial radiographs are negative

This summary of a Cochrane review presents what we know from research about the accuracy of imaging tests to detect true scaphoid fractures among suspected fractures.

When a patient presents to the emergency department with wrist injury and clinical signs of a scaphoid fracture, normal initial radiographs do not exclude a fracture. Approximately 20% of them do have a true scaphoid fracture and need additional imaging to establish a definitive diagnosis. Because of the low healing potential of the scaphoid bone, adequate diagnosis and treatment is vital to prevent complications such as non-union. If a patient is clinically suspected for a scaphoid fracture, their wrist will be immobilised in a cast until definitive diagnosis is obtained. This fear of under-treatment results in a large amount of over-treated wrist injuries. Computed tomography (CT), magnetic resonance imaging (MRI) and bone scintigraphy (BS; bone scan) are all imaging modalities that can be chosen at this stage. The aim of this systematic review was to establish which is the superior technique for identifying a true fracture and preventing unnecessary treatment. A high sensitivity reduces the risk of missing fractures; a low specificity increases the number of unnecessary treatments.

We conducted a thorough search of electronic databases, trial registers and conference proceedings up to July 2012. We included 11 studies in our analysis. The studies were moderate to good quality. Four studies (277 suspected fractures) looked at CT, five studies (221 suspected fractures) looked at MRI and six studies (543 suspected fractures) looked at BS. Four of these studies directly compared two modalities, such as both CT and MRI. When we compared the pooled data for the different imaging tests from all studies, we found that BS has the highest sensitivity, but specificity was lower than CT and MRI. All three imaging tests were found to be highly accurate for definitive diagnosis. CT and MRI were comparable in diagnostic accuracy (the correct diagnosis is made). Although BS had significantly better accuracy than

CT and MRI, it could lead to more people receiving unnecessary treatment. Moreover, BS is an invasive technique and is believed to be inappropriate for use in some populations, especially children.

Future studies should focus on improving clinical evaluation to raise the prevalence of true fractures. In addition, more direct comparison studies could add valuable data to determine which modality is superior in diagnosis of suspected scaphoid fractures.

SUMMARY OF FINDINGS

Summary of findings 1. Summary of findings: Diagnostic accuracy data

Comparing the diagnostic accuracies of computed tomography versus magnetic resonance imaging versus bone scintigraphy for clinically suspected scaphoid fractures in patients with negative plain radiographs

Patient population	Patients with a clinically suspected scaphoid fracture but normal radiographs after trauma of the wrist
Prior test-ing	Clinical evaluation
Setting	Emergency departments
Index tests	Computed tomography (CT), magnetic resonance imaging (MRI) and bone scintigraphy (BS)
Reference standard	Most studies used radiographs obtained after 6 weeks. Otherwise, 1 index test or 2 index tests with the same diagnosis (fracture or no fracture) were used
Target con-dition	Scaphoid fractures
Importance	Early definitive diagnosis of a scaphoid fracture ensures adequate treatment, prevents unnecessary immobilisation and minimises the risk of long-term complications (e.g. non-union)
Included studies	4 studies for CT; 5 studies for MRI; 6 studies for BS 2 studies compared CT and MRI; 1 study directly compared CT with BS; and 1 compared MRI with BS
Number of suspected fractures (patients) studied	277 (276 patients) for CT; 221 (221 patients) for MRI; 543 (542 patients) for BS
Quality concerns	Overall quality of the included studies was moderate to good. Of most concern was the lack of availability of relevant clinical information during evaluation of the images as this does not mimic daily practice. Five studies did not clearly describe fracture criteria for a positive test
Limitations	No study compared all three tests (CT, MRI and BS) in the same population Only four comparison studies were included. Current comparisons are based on indirect evidence with possible variations in confounding factors like pa-tient population and study characteristics Some studies were performed with only small cohorts

The confidence intervals for summary estimates are wide for all three tests

Test	Number of studies	Number of suspected fractures	Summary sensitivity (95% CI)	Summary specificity (95%)	Summary LR+ ¹ (95% CI)	Summary LR- ² (95% CI)	Consequences in a cohort of 1000 ³	
							Missed fractures	Overtreated
CT	4	277	0.72 (0.36 to 0.92)	0.99 (0.71 to 1.00)	119.98 (1.49 to 9655.66)	0.28 (0.10 to 0.85)	56	8
MRI	5	221	0.88 (0.64 to 0.97)	1.00 (0.38 to 1.00)	826.64 (0.51 to 1334596)	0.12 (0.03 to 0.42)	24	0
BS	6	543	0.99 (0.69 to 1.00)	0.86 (0.73 to 0.94)	7.35 (3.51 to 15.37)	0.01 (0.00 to 0.49)	2	112

Comparisons of the imaging tests

Comparison	Findings
CT, MRI and BS for diagnosis of clinically suspected scaphoid fractures	<ul style="list-style-type: none"> Overall diagnostic accuracy (DOR) of BS was significantly higher compared with CT ($\text{Chi}^2 = 50.3$, $\text{df} = 1$, $P < 0.01$) and MRI ($\text{Chi}^2 = 29.7$, $\text{df} = 1$, $P < 0.01$) CT and MRI were comparable in diagnostic accuracy ($\text{Chi}^2 = 1.9$, $\text{df} = 1$, $P = 0.17$) <p>The direct comparisons had similar patterns of differences in sensitivity and specificity as for the indirect comparisons.</p> <p>Given a median prevalence of 20%, 200 out of 1000 patients will have a scaphoid fracture.</p> <p>Of 200 cases, 56 will be missed if diagnosed using CT, 24 will be missed if diagnosed using MRI and 2 will be missed if diagnosed using BS.</p> <p>Of 800 patients without a scaphoid fracture, 8 will receive unnecessary treatment when CT is used for diagnosis, 0 when MRI is used for diagnosis and 112 if BS is used for diagnosis</p>

Conclusions:

The meta-analyses showed that DOR of BS is significantly better than CT ($P < 0.01$) and MRI ($P < 0.01$). This is based on a large difference in sensitivity. Conversely, specificities of CT and MRI are both higher than for BS. CT and MRI have comparable diagnostic accuracy. Direct comparisons showed similar patterns of differences in sensitivity and specificity. Reflecting the small number of studies, the confidence intervals for summary estimates are wide for all three tests.

There is a concern that the number of over-treated patients with BS is considerable, as well as the number of missed fractures on CT.

Quality of included studies was moderate to good, but there were only four direct comparison studies.

Well-designed studies directly comparing CT, MRI and BS could give valuable additional information.

1. LR+ Positive likelihood ratio
2. LR- Negative likelihood ratio
3. The median prevalence was 20%, calculated by using all studies. Missed fractures and over-treated patients were calculated using the median prevalence

BACKGROUND

Target condition being diagnosed

The scaphoid bone is one of the carpal wrist bones and is located in the proximal row. Its surface mainly consists of cartilage and it articulates with the distal radius, and with four other carpal bones: the lunate, trapezium, trapezoid and capitate. When flexing and extending the wrist, the scaphoid rotates forwards and backwards. The same movements can be found when twisting the wrist from the radial to the ulnar side. Owing to the scaphoid's anatomy, position and kinematics, it serves a key role in the function of the wrist.

Sustaining a fall on an outstretched hand (FOOSH) is the typical mechanism for fracturing the scaphoid. 'Axial fist' trauma, involving transmission of an external force through the second metacarpal when the fist is clenched, as when punching, is another, less common, cause. These types of trauma are most common in young and active males performing sports. Scaphoid fractures constitute approximately 2% to 3% of all fractures (Hove 1999). The scaphoid is the most commonly fractured carpal bone (Dennis 2011; Hove 1999; Van der Molen 1999; Van Onselen 2003).

One of the problems with fracturing the scaphoid is its low healing potential. The scaphoid's blood circulation mainly derives from small branches of the radial artery entering the bone from the distal part. The blood supply is fragile and can be interrupted when fractured (Gelberman 1986; Rhemrev 2011). If untreated, this can lead to non-union, with or without avascular necrosis, and finally carpal collapse and disability (Gelberman 1986; Merrell 2002). Early detection and adequate treatment can provide predictable and satisfactory rates of healing (Dias 2005). In contrast, delay of diagnosis and failure to recognise displacement are important risk factors for non-union of scaphoid wrist fractures (Adey 2007; Lozano-Calderon 2006).

When someone with a FOOSH or 'axial fist' trauma presents to the emergency department, certain clinical findings can lead to suspecting a scaphoid fracture. The most important physical examinations are pressing the anatomical snuffbox and applying longitudinal thumb compression (Pillai 2005; Rhemrev 2010a; Unay 2009). If either of these result in pain in the scaphoid area, radiographs of the wrist and the scaphoid are necessary. Usually x-rays are then obtained in four views: postero-anterior, true lateral, semipronated oblique, and posteroanterior with the wrist in ulnar deviation (Yin 2010). Most scaphoid fractures will be identified with this imaging technique, but up to 16% are missed on initial radiographs (Jenkins 2008; Mallee 2011). These missed fractures are also known as occult fractures. When clinical and radiographic findings do not match, we speak of a 'clinically suspected scaphoid fracture' and additional imaging (second-line imaging) is needed.

In cases of inadequate or delayed diagnosis, possible problems in union (bone healing) can lead to functional wrist problems (Merrell 2002; Rhemrev 2011). Therefore, despite the normal radiographs, current clinical practice is to immobilise the scaphoid in a cast or splint until further imaging is established. The fear of under-treatment results in over-treatment of five out of six patients (Mallee 2011; Rhemrev 2010b).

Difficulties in detecting occult scaphoid fractures have been addressed in many radiological studies, aiming at exploring

the value of novel imaging techniques or updates of already known techniques such as computed tomography (CT), magnetic resonance imaging (MRI), bone scintigraphy (BS) and ultrasound (US) (Breitenseher 1997; Roolker 1997; Senall 2004; Tiel-van Buul 1993). However, there is currently no consensus regarding which modality is best to detect an occult scaphoid fracture. Several worldwide and national studies showed considerable variation in the management of occult scaphoid fractures (Brookes-Fazakerley 2009; Groves 2006). This is partly attributed to the availability of the imaging tools and differences in costs, but also to the controversies regarding the best method to detect true scaphoid fractures. The international questionnaire-based survey of Groves 2006 revealed equivalent imaging strategies for suspected scaphoid fractures in only 6.7% of the, mainly university, hospitals. Groves 2006 reported that the most commonly used second-line imaging modality in Europe was CT, whereas it was BS in Australasia and MRI in North America. This variation shows that there is a lack of agreed standard diagnostic practice, which amplifies the need for this review. Furthermore, the increase in availability of CT scanning in emergency (radiology) departments and dedicated MRI equipment, such as tailored sequences and dedicated wrist coils, enables earlier use of these techniques in daily clinical practice. Yet clear evidence of optimal scaphoid conventional imaging protocols is lacking, especially concerning cost effectiveness and patient safety (radiation protection).

Besides detecting a fracture, the location of the scaphoid fracture is important too. The proximal pole of the scaphoid is prone to complications after fracture owing to its limited vascularity. It has been proposed that these fractures need to be treated operatively because cast immobilisation will not ensure adequate healing. This differs from undisplaced fractures through the waist of the scaphoid for which union rates of up to 95% have been reported after cast immobilisation (Geissler 2012).

In general, the key to evaluating the performance of a diagnostic test is an agreed-upon reference standard that is used to define the presence or absence of a disease. We know that an important caveat in the interpretation of studies of the diagnostic performance characteristics of various imaging modalities for triage of suspected scaphoid fractures is the lack of an agreed-upon reference standard for the diagnosis of a true fracture of the scaphoid. The most commonly applied test is the six-week follow-up set of radiographs. This is generally considered to be the most valid reference test (Mallee 2011). When we examine some of the prospective trials studying one or more index tests, lists of reference standards are often given. Other methods used are:

- if two of the index tests are positive (MRI, CT, BS), the diagnosis is a fracture;
- if two of the index tests are negative (MRI, CT, BS), the diagnosis is 'no fracture';
- clinical follow-up and radiographs after two weeks;
- clinical follow-up and MRI;
- single use of an index test (MRI, CT, BS);
- single use of clinical follow-up.

These methods are sometimes used in research as reference standards but some are considered suboptimal. These differences in approach hamper the interpretation of the scaphoid imaging literature because most of the results found are not checked with an optimal reference test. We consider the single use of an index test

(MRI, CT, BS) and the clinical follow-up with radiographs after two weeks as a 'suboptimal' reference test. The use of clinical follow-up alone is even more unsatisfactory as a reference standard.

Treatment of a non-displaced or minimally displaced fractured scaphoid can be operative or non-operative and is mainly based on the location of the fracture. The majority of the fractures are located in the waist of the scaphoid (Geissler 2012). Whereas waist and distal pole fractures seem to heal with acceptable rates with cast treatment, it is a fracture of the proximal pole that is prone to non-union. Therefore, these fractures are considered unstable and require operative treatment (Rettig 1999). The non-operative method is with use of a cast or splint that prevents the scaphoid's movements. Healing of a scaphoid fracture to union is a time-consuming process that results in the need for a long period of immobilisation, ranging from 6 to 12 weeks (Bond 2001; Dias 2005; Vinnars 2008). To avoid this burden, operative fixation with a headless compression screw can be performed (Fowler 2010). Surgical treatment is favourable in terms of time off work and functional outcome, but can lead to more (minor) complications (Buijze 2010).

Index test(s)

The tests evaluated in this review are multi-slice CT, MRI and BS.

CT creates axial images of the wrist that can be reconstructed in different planes, such as anatomical coronal and sagittal series. Several studies show preferable use of reconstructions in planes defined by the long axis of the scaphoid (Mallee 2011; Sanders 1988; Ty 2008). Image reconstruction in CT is a mathematical process that generates images from X-ray projection data acquired at many different angles around the patient. Image reconstruction has a fundamental impact on image quality and therefore on radiation dose. No literature could be found comparing different types of image reconstruction; we will therefore evaluate all types in this review.

MRI generates a strong magnetic field to align the hydrogen atoms in the body. This alignment is altered with use of radiofrequency pulses and can be detected to build the images. MRI was the first non-invasive method to create high-resolution images of the musculoskeletal system. In scaphoid injury, bone bruising or bone marrow oedema consists mainly of liquid with hydrogen atoms, and thus is well visualised. Cortical involvement of the fracture can, therefore, be less obvious. The exact value of bone marrow oedema in the clinical spectrum of scaphoid injury is unclear; as is its relationship to patient outcome.

BS is widely described for scaphoid disorders. After an intravenous injection with radioactive isotopes, the osteoblastic activity can be visualised. A gamma camera can detect the radiation emitted by the isotopes. Where there is a fracture, osteoblastic activity is high at the fracture site indicating the natural healing process of the bone. This activity is displayed as a dense spot in the bone. BS provides a radiation burden and is thus potentially harmful, especially to the younger age group.

When we consider the negative aspects of the additional imaging methods, we find that:

MRI:

- is known for its low availability and generally higher costs compared with CT;
- produces images in which bone bruising can be difficult to distinguish from a fracture (Mallee 2011). No clear criteria for a bruise or a fracture are established. When bone bruising is detected, the possibility of fracture development must be remembered (Thavarajah 2011); and thus follow-up is important.

CT:

- is one of the modalities that uses radiation. Although the dose of 0.03 mSv for imaging the wrist is very low (Biswas 2009), its use in the younger patient group is debatable.

BS:

- uses radiation. With 4 mSv, the dose is much higher than CT, but still only the same as two years of natural background radiation (Rhemrev 2010b). BS is not recommended for children;
- needs radioactive isotopes that must be injected intravenously, which makes BS the most invasive procedure of all;
- can only be performed with an interval of 72 hours after injury. This delay is needed to capture osteoblastic activity at the fracture site in all patients (McDougall 1989);
- in the lead author's hospital, the costs of BS are comparable with those for MRI.

Alternative test(s)

Ultrasound (US) can be used to diagnose suspected scaphoid fractures. The literature evaluating its performance characteristics is scarce and the latest review including US shows inferior results compared with MRI, CT or BS (Ring 2008). In addition, an international survey of imaging strategies among hospitals revealed no use of US for these injuries (Groves 2006). This review therefore does not consider US.

Another test, six-week follow-up radiographs, is extensively used in literature as a reference standard (Mallee 2011; Memarsadeghi 2006); but its accuracy is being questioned (Mallee 2011). One of the main disadvantages is the time interval before this test can be performed, given the need for immobilisation. The importance of immediate diagnosis rules out the use of the follow-up radiographs as an adequate diagnostic tool. Moreover, a positive CT, MRI or bone scan can be accompanied by normal x-rays after six weeks. These disadvantages make the quality and clinical applicability of this test questionable.

Rationale

In clinically suspected scaphoid fractures, early diagnosis reduces the risk of non-union and minimises any loss in productivity resulting from unnecessary cast immobilisation (Dorsay 2001). This means improvement of short-term management (avoid unnecessary immobilisation) and long-term outcome (risk of non-union, avascular necrosis). The value of an imaging tool with the highest accuracy is of great importance for both the patient and economically in terms of healthcare costs and productivity loss.

There are many controversies surrounding the choice of imaging modality; this is reflected in the considerable variation in practice (Groves 2006). All three imaging modalities (CT, MRI and BS) are

widely used and reviews of these have reported that all show high sensitivity and specificity rates ([Ring 2008](#); [Yin 2010](#)). The most recent review searched up to October 2008, but did not include non-English studies even though there were three potentially eligible reports in foreign languages ([Yin 2010](#)). Since 2000, several articles evaluating one or two tests have been published. Hence, an update of the evidence was warranted.

With this review, we evaluated the diagnostic performance characteristics of BS, MRI and CT with an updated search for diagnostic accuracy studies and the inclusion of non-English literature.

OBJECTIVES

The primary aim of this study is to identify the most suitable diagnostic imaging strategy for identifying clinically suspected fractures of the scaphoid bone in patients with normal radiographs. Therefore we looked at the diagnostic performance characteristics ([Appendix 1](#)) of the most used imaging modalities for this purpose: computed tomography, magnetic resonance imaging and bone scintigraphy.

Secondary objectives

To investigate which imaging technique is the best for determining the location of the fracture (proximal, waist or distal).

Investigation of sources of heterogeneity

We assessed the potential influence of sources of heterogeneity on the diagnostic accuracy of the tests, especially the type of reference standard and blinded evaluation of the reference test (if reported).

METHODS

Criteria for considering studies for this review

Types of studies

All prospective or retrospective studies involving a consecutive series of patients. We only included trials using reference standards that we considered optimal or adequate. Randomised controlled trials would have been included if these had been found.

Participants

People of all ages who presented at hospital or clinic within one week of trauma with a clinically suspected scaphoid fracture and negative post-trauma radiographs. Clinical suspicion of a scaphoid fracture is based on pain in the anatomical snuffbox or by longitudinal compression of the thumb, or both. The radiographs generally include two images of the wrist (postero-anterior and lateral views) and at least one of two additional scaphoid views.

Index tests

CT, MRI or BS, or a combination of two of these tests. Because the criteria for a fracture may differ (especially in MRI), we report all study characteristics, including 'fracture criteria', in [Characteristics of included studies](#).

Target conditions

Clinically suspected scaphoid fractures (which could be proximal, waist or distal) with negative plain radiographs.

Reference standards

Various reference standards were included.

1. A scaphoid plain radiograph series, conducted six to 14 weeks after the initial injury, consisting of the following four views: posteroanterior with the wrist in neutral position; lateral; semipronated oblique scaphoid; and radial oblique scaphoid. An abnormal lucent line within the scaphoid is considered evidence of a fracture.
2. The use of two index tests. If both are positive or negative, a final diagnosis is obtained.
3. In addition, clinical findings are often combined with an index test or repeated radiographs obtained after six weeks to formulate a reference standard.
4. The use of only one of the second-line modalities has been described; this is somewhat unsatisfactory because these diagnostic techniques are still under study.

We considered six-week follow-up radiographs (1) the most suitable reference standard. Next we considered the use of two index tests with the same outcome and one index test including clinical findings (2 and 3). Although we considered the fourth option to be suboptimal, it was included in the review.

We did not include studies using clinical findings only six to 14 weeks after trauma or the single use of one- to two-week follow-up radiographs as a reference standard as we consider these inadequate.

Search methods for identification of studies

Electronic searches

We searched the Cochrane Register of Diagnostic Test Accuracy Studies (July 2012), MEDLINE (1946 to July Week 1 2012) and EMBASE (1974 to 2012 Week 27). We also searched the Database of Abstracts of Reviews of Effects (*The Cochrane Library* 2012 Issue 7), [MEDION](#) (Meta-analyses van Diagnostisch Onderzoek) (September 15th 2012) and the [Aggressive Research Intelligence Facility](#) (ARIF) reviews database (15 September 2012) for relevant diagnostic reviews. In addition, we searched the Cochrane Central Register of Controlled Trials (*The Cochrane Library* 2012 Issue 7) and the NHS Economic Evaluation Database (*The Cochrane Library* 2012 Issue 7) for comparative and cost-effectiveness studies looking at different diagnostic modalities. We searched [Current Controlled Trials](#) (15 September 2012) and the [WHO International Clinical Trials Registry Platform](#) (15 September 2012) for ongoing studies.

We developed a sensitive search strategy for MEDLINE (Ovid Web), EMBASE (Ovid Web) and *The Cochrane Library* (Wiley Online Library) as recommended in Chapter 7 of the *Cochrane Handbook for Systematic Reviews of Diagnostic Test Accuracy* ([de Vet 2008](#)). The search strategies for all databases are shown in [Appendix 2](#).

There were no restrictions based on language or publication status.

Searching other resources

We checked the reference lists of all articles, including reviews, for relevant primary diagnostic studies and systematic or narrative reviews.

We handsearched the abstracts of the conference proceedings of two societies: the [American Society for Surgery of the Hand](#)

annual meetings (2000 to 2012); and the [American Academy of Orthopaedic Surgeons](#) annual meetings (2011 to 2013). If potentially eligible abstracts were found, we searched for the full reports.

We also contacted experts in the field and main investigators of relevant ongoing studies for additional information.

Data collection and analysis

Selection of studies

Two review authors (WHM and JND) independently screened the titles and abstracts of retrieved publications to identify potentially eligible studies for inclusion. WHM and JND assessed full-text reports of potentially eligible studies and independently determined study inclusion or exclusion. Any disagreement was either resolved by discussion; or, if necessary, by an arbiter (RWP). When WHM and JND were involved in one of the studies, two other authors (RWP and PK) were asked to assess eligibility. Only results of full reports were evaluated.

Data extraction and management

Two review authors (WHM and JND) independently extracted data from full-text reports. If studies had been published more than once, only data from the latest or most suitable report were included. (In cases of overlapping patient data, we only used the data once.) Any disagreement was discussed, either until consensus was achieved, or, if necessary, with an arbiter (RWP). When WHM and JND were involved in one of the included studies, two other authors were asked to extract data. Where necessary, we contacted study authors for additional information or data.

The following data were collected:

1. general information: title, journal, year, publication status, country of study, period of study, primary objective and study design (prospective versus retrospective and consecutive versus non-consecutive; randomised);
2. sample size (screened and included);
3. baseline characteristics: age, sex, side of injury, trauma mechanism, time of presentation, inclusion and exclusion criteria;
4. target condition, as reported;
5. index test: description of technique, criteria for a fracture, timing of test and expertise of the tester;
6. reference standard test: description of technique, criteria for a fracture, time from trauma to reference test and expertise of the tester;
7. sensitivity and specificity;
8. number of true positives (TP), true negatives (TN), false positives (FP), and false negatives (FN).

Assessment of methodological quality

Two review authors (WHM and JND) independently assessed the methodological quality of the included studies using a slightly modified version of the QUADAS checklist ([Whiting 2003](#)). Both review authors had prior knowledge of the methodological aspects of diagnostic accuracy studies. Where any disagreement on the quality assessment occurred, a third review author (RWP) was asked to arbitrate. When WHM and JND were involved in one

of the included studies, two other authors were asked to assess the methodological quality. We used the QUADAS checklist with previously set criteria specific to the review topic ([Table 1](#)).

To inform our assessment of overall methodological quality we established the following general 'rules'. We considered the methodological quality was 'excellent' if all QUADAS items were met; and 'good' if at least item 2 (acceptable reference standard?) was scored as 'yes', with the other items open for discussion between the two review authors (WHM and JND). We considered quality was 'moderate' if either item 1 (representative spectrum?) or item 2 was scored as 'unclear' or 'no'; again with the other items open for discussion. We considered quality was 'poor' if both items 1 and 2 were scored 'no'.

Statistical analysis and data synthesis

The main target was to identify the index test with the highest diagnostic accuracy for diagnosing suspected scaphoid fractures. With the outcomes of each primary study, we generated 2 x 2 tables (with TPs, TNs, FPs and FNs) for each diagnostic test according to the presence or absence of a true fracture. With these data, sensitivity and specificity fractions are presented. Where results were reported as 'inconclusive' (as in [Nielsen 1983](#)), we treated these as negative findings. If the data presented in trials had been uninterpretable in that 2 x 2 tables could not be generated, we planned to contact the original authors of the study for clarification, and otherwise present the data only descriptively.

The two main parameters of diagnostic test accuracy are sensitivity and specificity. As there is a trade-off between these parameters, they should not be analysed separately. For descriptive purposes, coupled forest plots are presented showing the pairs of sensitivities and specificities with 95% confidence intervals (CIs). Sensitivity and specificity are displayed in the receiver operating characteristic (ROC) space.

Diagnostic accuracy was first evaluated for each index test individually. For pooling sensitivities and specificities, we assume there is at least one common criterion for test positivity used across studies for a given test. Given the fact that different studies may have slightly different criteria for test positivity, and individual observers within a study may interpret the criteria a little differently, the bivariate random effects model was used to get the summary estimates of sensitivity and specificity. A separate model was fitted for each index test with bivariate approach except CT. For CT, the estimation from the bivariate model did not converge. This may be due to the small number of studies (four studies for CT) included in the meta-analysis. So we used the HSROC model as an alternative, which could give mathematical equivalent estimates of bivariate approach. Both models produced summary estimates of the mean sensitivity and specificity with corresponding 95% CIs. Summary estimates of sensitivity, specificity, positive likelihood ratio, negative likelihood ratio and their 95% CIs were calculated by using "estimate" command in SAS.

Pairwise comparisons between CT, MRI and BS were based on the overall performance, measured by diagnostic odds ratio (DOR). We added test type as covariate into the HSROC model and tested the statistical significance of the covariate effects on the test accuracy. The strategy of comparison was as follows: first, we had model (a) ([Table 2](#)), which included covariates for shape (beta), accuracy (alpha) and threshold (theta); then covariate for shape was dropped

and we got model (b), and a χ^2 test was performed on the change in the -2 log likelihood from model (a) to model (b). If the curves had different shapes, it indicated that the differences in test accuracy depended on threshold. Otherwise, we continued to drop the covariate for accuracy and got model (c), and then compared -2 log likelihood with model (b) using the χ^2 test. If the likelihood test showed a significant change from model (b) to model (c), then we can say there is a significant difference in the accuracy between the tests being compared.

Our second target was to identify the accuracy of fracture location detection (proximal, waist, distal). This was not done for the current version of the review. Should there be sufficient studies containing adequate information about fracture location in future, we plan to include only the fractured scaphoids and generate 2 x 2 tables for each diagnostic test. We plan to present sensitivity, specificity and predictive values and calculate these in the same way as our main target. We also intend to consider a second option, which is to keep the entire dataset (i.e. including people with no fracture), and compute the relative sensitivity and specificity for fractures in different locations; and thereby compare the accuracy to detect the presence of a fracture at each location.

Investigations of heterogeneity

Heterogeneity in diagnostic test accuracy reviews is expected. Aside from analyses in which the different index tests are presented as subgroups, none of the planned subgroup analyses to investigate heterogeneity were performed. Should there be sufficient data available in future, we will conduct subgroup analyses based on the assessment of methodological quality (yes versus no or unclear) from items 2 (acceptable reference standard?), 3 (acceptable delay between tests?), 4 (partial verification avoided?), 5 (differential verification avoided?) and 6 (incorporation avoided?) of the QUADAS criteria. Additionally, if there are sufficient studies, we will perform a meta-regression analysis. Characteristics of the index test, study population (adults/children), and judgements for the five QUADAS items will be added to the model as covariates, to analyse their influence on diagnostic accuracy. Heterogeneity will be judged on the scatter of points and from the prediction ellipse. This graphical information will also be used to decide about subgroups. We will present pooled estimates per clinical relevant subgroups. The possibilities of performing meta-regression analyses will depend on the number of studies available for a specific index test providing sufficient information.

Sensitivity analyses

During the review, a number of subjective choices were made with regard to eligibility, methodological quality and clinical similarity. The influence of these decisions on the outcome of the review

should ideally be explored in sensitivity analyses (e.g. QUADAS item 12 (clearly described fracture criteria for index test)), but this was not possible since there were too few studies for proper analyses.

Our planned sensitivity analysis based on indirect comparison versus direct comparison was also hindered because of the small numbers of studies making direct comparisons. In order to compare the accuracy of the index tests, two strategies could be applied. We could include all studies examining one or more index test or we could include only studies that presented a direct comparison between two or more index tests. Although the first analysis is based on all available data, the second analysis potentially gives more valid data for the comparison. These two strategies may lead to different conclusions, so, while we decided to include all studies, we also checked the results of the few direct comparison studies. If there had been sufficient data, we would also have examined whether the results of the meta-analyses would have changed if we had included only direct comparison studies.

RESULTS

Results of the search

For this search (main search date July 2012), we screened a total of 2900 records from the following databases: Cochrane Register of Diagnostic Test Accuracy Studies (14 records); MEDLINE (1226); EMBASE (1586), the Database of Abstracts of Reviews of Effects (2); MEDION (0); ARIF (3), the Cochrane Central Register of Controlled Trials (34); the NHS Economic Evaluation Database (8); the WHO International Clinical Trials Registry Platform (13) and Current Controlled Trials (14). We did not identify potentially eligible studies from other sources.

The search resulted in the identification of 64 potentially eligible articles, for which (where possible) full reports were obtained. Upon study selection, we included 11 studies (Beeres 2008; Breitenseher 1997 (published in five reports); De Zwart 2012 (published in three reports); Ilica 2011; Mallee 2011; Memarsadeghi 2006; Nielsen 1983; O'Carroll 1982; Stordahl 1984; Tiel-van Buul 1993 (published in two reports); Tiel-van Buul 1996); and 45 studies were excluded, one of which was published in two reports (Lepage 2004). There were no ongoing trials or studies awaiting classification. All studies were written in English. Five studies were conducted in The Netherlands, two in Austria and one in each of Turkey, Ireland, Norway and Denmark. All studies included patients that presented to the emergency department with clinical suspicion of a scaphoid fracture, but with normal initial radiographs.

A flow diagram summarising the study selection process is shown in Figure 1.

Figure 1. Study flow diagram

Figure 1. (Continued)

Included studies

The characteristics of the individual studies are reported in the [Characteristics of included studies](#).

Four studies evaluated CT ([De Zwart 2012](#); [Ilica 2011](#); [Mallee 2011](#); [Memarsadeghi 2006](#)); five studies evaluated MRI ([Beeres 2008](#); [Breitenseher 1997](#); [Mallee 2011](#); [Memarsadeghi 2006](#); [Tiel-van Buul 1996](#)); and six studies evaluated BS ([Beeres 2008](#); [De Zwart 2012](#); [Nielsen 1983](#); [O'Carroll 1982](#); [Stordahl 1984](#); [Tiel-van Buul 1993](#)). Of these studies, two compared CT with MRI ([Mallee 2011](#); [Memarsadeghi 2006](#)); one study compared BS with CT ([De Zwart 2012](#)); and one compared MRI with BS ([Beeres 2008](#)).

The main objective for all studies was the detection of a true scaphoid fracture among clinically suspected scaphoid fractures. A total of 717 patients with 719 clinically suspected scaphoid fractures were assessed. For CT, 276 patients with 277 suspected fractures provided data; 221 patients for MRI; and 542 patients with 543 suspected fractures for BS. The sample size ranged from 16 to 159, with a mean of 65 patients. The weighted mean age of the studies was 36.5 years (range 10 to 88 years). Five studies included children, one of which evaluated MRI ([Breitenseher 1997](#)); and the other four of which evaluated BS ([Nielsen 1983](#); [O'Carroll 1982](#); [Stordahl 1984](#); [Tiel-van Buul 1993](#)). The gender distribution was available for 10 studies, in which the proportion of men ranged from 49.7% ([De Zwart 2012](#)), to 100% ([Ilica 2011](#)).

Seven studies assessed patients within 72 hours of the patient injuring their wrist; four studies did not report the timing of presentation to the emergency department ([Breitenseher 1997](#); [Nielsen 1983](#); [O'Carroll 1982](#); [Stordahl 1984](#)). In seven studies, the index test was performed within 10 days of trauma ([Beeres 2008](#); [Breitenseher 1997](#); [De Zwart 2012](#); [Ilica 2011](#); [Mallee 2011](#); [Memarsadeghi 2006](#); [Nielsen 1983](#)). Tenderness in the anatomical snuffbox was clearly incorporated in clinical evaluation in six studies ([Beeres 2008](#); [Breitenseher 1997](#); [De Zwart 2012](#); [Ilica 2011](#); [Mallee 2011](#); [Tiel-van Buul 1993](#)). One study reported 'pain over the

scaphoid' as being clinically suspected ([Memarsadeghi 2006](#)). Four studies did not define the content of clinical evaluation ([Nielsen 1983](#); [O'Carroll 1982](#); [Stordahl 1984](#); [Tiel-van Buul 1996](#)). Images of BS were evaluated by a consultant clinical nuclear physician in four studies ([Beeres 2008](#); [De Zwart 2012](#); [Tiel-van Buul 1993](#); [Tiel-van Buul 1996](#)); three studies (two when BS was an index test, one when BS was a reference standard) did not provide the expertise of the observer(s) ([Nielsen 1983](#); [O'Carroll 1982](#); [Stordahl 1984](#)). For MRI and CT, evaluation was performed by at least one experienced radiologist.

This review focused on true scaphoid fractures among clinically suspected scaphoid fractures. In addition, all studies reported on the diagnosis of other wrist fractures (see [Characteristics of included studies](#)).

Excluded studies

We excluded 45 studies; the characteristics of these studies are presented in the [Characteristics of excluded studies](#). The most common reasons for exclusion were that no reference standard was used or that it was inadequate (21 studies), or that patients were included after a second clinical evaluation after one to two weeks (eight studies). Inadequate reference tests included repeating the radiographs after 10 days or using only clinical evaluation after one to two weeks. Some studies did not perform any other test besides initial clinical and radiographic evaluation.

Methodological quality of included studies

The included studies were diverse but all were of moderate to good quality ([Figure 2](#) and [Figure 3](#)). Five studies were considered 'good quality' ([Breitenseher 1997](#); [Mallee 2011](#); [Memarsadeghi 2006](#); [Nielsen 1983](#); [Tiel-van Buul 1993](#)); and six studies were considered 'moderate quality'. Of these, two studies had three items scored as low quality ([Beeres 2008](#); [De Zwart 2012](#)) and one study had five items scored as unclear and one item scored as low quality ([O'Carroll 1982](#)).

Figure 2. Methodological quality graph: review authors' judgements about each methodological quality item presented as percentages across all included studies

Figure 3. Methodological quality summary: review authors' judgements about each methodological quality item for each included study

	Representative spectrum?	Acceptable reference standard?	Acceptable delay between tests?	Partial verification avoided?	Differential verification avoided?	Incorporation avoided?	Index test results blinded?	Reference standard results blinded?	Relevant clinical information?	Uninterpretable results reported?	Withdrawals explained?	Fracture criteria defined?
Beeres 2008	+	+	+	+	-	-	+	+	-	+	+	?
Breitenseher 1997	?	+	+	+	+	+	+	+	?	+	+	+
De Zwart 2012	+	+	+	+	-	-	+	+	-	+	+	?
Ilica 2011	+	?	+	+	+	+	+	+	-	+	+	?
Mallee 2011	+	+	+	+	+	+	+	+	?	+	+	+
Memarsadeghi 2006	+	+	+	+	+	+	?	+	+	+	+	+
Nielsen 1983	?	+	+	+	+	+	?	+	-	+	+	+
O'Carroll 1982	?	+	?	+	+	+	?	?	?	+	+	-
Stordahl 1984	-	+	+	+	+	+	+	+	?	+	+	+
Tiel-van Buul 1993	+	+	+	+	+	+	+	+	?	+	+	+
Tiel-van Buul 1996	+	?	+	+	+	+	+	+	-	+	+	?

All studies recruited patients consecutively as per our inclusion criteria. A prospective study design was clearly reported in eight studies. In three studies this was unclear, but due to the use of a reference standard, we assumed these were prospective as well. In only one study (Stordahl 1984) was the spectrum of patients not clear; since the timing of presentation and precise aspects of clinical evaluation were not reported, we judged this study to be low quality for this item. Participants in nine studies

received an acceptable reference standard: seven studies used follow-up radiographs in four or more views after at least six weeks (Breitenseher 1997; Mallee 2011; Memarsadeghi 2006; Nielsen 1983; O'Carroll 1982; Stordahl 1984; Tiel-van Buul 1993); and two studies used a mixed reference standard (same outcome in two index tests or six-week follow-up radiographs) (Beeres 2008; De Zwart 2012). Two studies were judged at lower quality as they used suboptimal reference standards: one used MRI (Ilica 2011); and the

other used BS (Tiel-van Buul 1996). Because of the mixed use of at least one index test as a reference test, differential verification and incorporation bias could not be avoided in these two studies (Beeres 2008; De Zwart 2012). Only one study reported the use of clinically relevant information during evaluation of the images (Memarsadeghi 2006); five studies excluded this information intentionally (Beeres 2008; De Zwart 2012; Ilica 2011; Nielsen 1983; Tiel-van Buul 1996). The criteria for diagnosing a fracture was not defined in O'Carroll 1982 for BS; for CT in De Zwart 2012 and Ilica 2011; and for MRI in Beeres 2008 and Tiel-van Buul 1996. However, we rated the latter four studies as unclear for this item because of other information and that the evaluation of test results was performed by at least two observers.

Findings

Indirect comparisons

The forest plots of the diagnostic performance characteristics of CT, MRI and BS are presented in Figure 4. The median prevalence of a true scaphoid fracture among clinically suspected scaphoid fractures with normal radiographs is 20% (range 11% to 44%). For CT, sensitivity estimates ranged from 0.67 (95% CI 0.22 to 0.96) to 0.88 (95% CI 0.62 to 0.98) and specificity estimates from 0.96 (95% CI 0.82 to 1.00) to 1.00 (95% CI 0.81 to 1.00). For MRI, sensitivity estimates ranged from 0.67 (95% CI 0.22 to 0.96) to 1.00 (95% CI 0.72 to 1.00) and specificity estimates from 0.89 (95% CI 0.72 to 0.98) to 1.00 (95% CI 0.66 to 1.00). For BS, sensitivity estimates ranged from 0.95 (95% CI 0.75 to 1.00) to 1.00 (95% CI 0.54 to 1.00) and specificity from 0.52 (95% CI 0.41 to 0.63) to 1.00 (95% CI 0.82 to 1.00).

Figure 4. Forest plot of tests: 1 CT, 2 MRI, 3 BS

CT

MRI

BS

The study specific and pooled estimates and 95% confidence regions are displayed in a scatter plot for CT, MRI and BS (Figure 5). The pooled estimates for CT sensitivity and specificity were 0.72 (95% CI 0.36 to 0.92) and 0.99 (95% CI 0.71 to 1.00), respectively; the

pooled estimates for MRI sensitivity and specificity were 0.88 (95% CI 0.64 to 0.97) and 1.00 (95% CI 0.38 to 1.00), respectively; and the pooled estimates for BS sensitivity and specificity were 0.99 (95% CI 0.69 to 1.00) and 0.86 (95% CI 0.73 to 0.94), respectively.

Figure 5. Study specific and pooled estimates of test performance for CT, MRI and BS with 95% confidence regions

Pairwise comparisons were performed using HSROC model (see Statistical analysis and data synthesis section above). The -2 log likelihood of model (a), (b) and (c) of comparisons between each pair of tests are shown in [Table 2](#). By comparing -2 log likelihood between model (a) and model (b), we found that the differences in test accuracy do not depend on threshold (since CT, MRI and BS do not have thresholds), thus we could continue to compare the overall accuracy (DOR) between tests. When comparing the overall accuracy (comparing model (b) and model (c)) of these tests, significant differences were found in 'CT versus BS' ($\chi^2 = 50.3$, $df = 1$, P value < 0.01) and 'MRI versus BS' ($\chi^2 = 29.7$, $df = 1$, P value < 0.01), which indicates that the overall accuracy of BS is higher than CT and MRI; while no evidence was found for a difference in accuracy between CT and MRI ($\chi^2 = 1.9$, $df = 1$, P value = 0.17). These results may be explained by the findings from the summary estimates (see [Summary of findings 1](#)): BS has a slightly lower specificity but a much higher sensitivity than CT and MRI, which leads to higher DOR for BS.

Direct comparisons

The separate findings of the four studies providing direct comparisons between tests are shown in [Table 3](#). The direct comparisons showed similar patterns of differences in sensitivity and specificity as for the indirect comparisons.

The two studies directly comparing CT and MRI found comparable sensitivities and specificities for the two tests ([Mallee 2011](#); [Memarsadeghi 2006](#)), with neither trial finding statistically significant differences between tests (reported P values > 0.05). The study directly comparing CT with BS ([De Zwart 2012](#)) reported a lower sensitivity, which was not statistically significant (reported $P = 0.13$) and a higher specificity (reported $P = 0.02$) for CT, but no statistically significant difference in the percentage of "correct predictions (accuracy)" (reported $P = 0.63$). The study directly comparing MRI with BS ([Beeres 2008](#)), which found a lower sensitivity and higher specificity for MRI, reported no statistically significant difference in "the percentage of correct predictions with MRI and bone scintigraphy ($p = 0.388$)".

Secondary objectives

There was no information about the diagnostic accuracy of the tests for identifying the location of the fracture (proximal, waist, distal).

DISCUSSION

Summary of main results

Early diagnosis and treatment of patients with a clinically suspected scaphoid fracture minimises the risk of complications and prevents unnecessary cast immobilisation. If initial radiographs appear normal, approximately 20% will still have a true fracture. In clinical practice, a definitive diagnosis is established by using CT, MRI or BS. This systematic review summarised the evidence and compared the diagnostic accuracies of these three imaging modalities. Eleven studies, four which evaluated two index tests, were included in the comparison: four studies for CT, five studies for MRI and six for BS.

We found evidence that BS has a significantly higher diagnostic accuracy (DOR) than CT and MRI; which reflects the higher sensitivity for BS. The summary sensitivity and specificity of BS were 0.99 and 0.86, respectively. For CT, summary sensitivity and

specificity were 0.72 and 0.99. For MRI, summary sensitivity and specificity were 0.88 and 1.00. Specificities of CT and MRI were both higher than BS. The single studies that directly compared CT and BS and MRI and BS found a similar pattern of the differences in sensitivity and specificity; however, both studies reported a lack of significant difference in the percentage of correct predictions. No differences were found between the diagnostic accuracies of CT and MRI. This finding applied also to the data from the two studies directly comparing CT and MRI. A summary of all results is presented in [Summary of findings 1](#).

Quality assessment showed moderate quality (six studies) to good quality (five studies). All patients were consecutive cohorts and at least eight (though the methodology suggests all) studies were explicitly prospective research. 'Relevant clinical information' was often not available during evaluation of index tests and is therefore a possible risk of bias. This should be included in future studies as omitting it is not representative of clinical practice. The other 11 items were mainly scored as 'Yes', implying good quality.

We could not find any information on which imaging technique is best for determining the location of the fracture (proximal, waist or distal). Some articles presented the location of a scaphoid fracture when presenting results for an index test; however, diagnostic accuracy calculations were not performed. In scaphoid fractures, healing is believed to be more problematic when fractures occur in the proximal part since blood supply is interrupted.

Strengths and weaknesses of the review

The evidence provided by this review is based on a comprehensive and sensitive literature search with the aim of identifying all relevant studies. All major electronic databases were searched and articles were selected with clear inclusion and exclusion criteria. Only studies with consecutive series of patients were included, which mimics clinical practice.

Another strength of this review is the usage of a well-regarded assessment tool to evaluate the quality of included studies: QUADAS. This tool provides detailed evaluation of quality and enables a simple and clear presentation of the assessment ([Figure 2](#); [Figure 3](#)).

A key issue in diagnostic accuracy studies is the application of an adequate reference standard to test for true disease status. This issue is much debated in scaphoid literature and the lack of evidence and consensus on the right reference standard limits evaluation of diagnostic accuracy. Even though it is debated, follow-up radiographs at six weeks is generally considered to be the most suitable reference standard. The timing of visualisation of a lucent line on a radiograph is unknown but believed to be two to six weeks. This supports our choice to exclude reference standards that only consisted of repeated (radiographical) evaluation after one to two weeks, as this has been shown to be inadequate.

In our decision to pool data from studies, the similarity or equivalence in the criteria for test positivity is a critical issue. Thus the failure of some studies to report clear fracture criteria, which is a vital aspect for the interpretation of images, is clearly a problem. Where the criteria were not described, we considered the evaluation of images by two observers provided some assurance of an appropriate process. Since when reported, the criteria for CT, MRI and BS were sufficiently similar to merit pooling, we decided

that it was a reasonable assumption that similar criteria would have been applied in all studies. Clearly, more precise criteria would be desirable in all future studies.

Another weakness of the review is the lack of direct comparison studies that include all three index tests. In addition, only four direct comparison studies including two index tests were evaluated. This means that comparison of CT, MRI and BS is mainly based on studies testing diagnostic accuracy of only one index test, i.e. indirect evidence. Another limitation of the review is that the findings derive from only a few studies. Therefore, sensitivity analyses could not be performed and potential sources for heterogeneity could not be investigated formally.

Our secondary objective for the review, accuracy of determining the location of the fracture, could not be answered and is therefore a weakness of the review. To date, we know of no studies that present these results.

A key limitation is the date of the search, July 2012; however, we are not aware of any new studies or current research on this topic.

Applicability of findings to the review question

The quality of the included studies was moderate to good and the data from these suggest that BS is the most sensitive modality to use in diagnosis of suspected scaphoid fractures. Direct comparison studies were few as indeed were the numbers of indirect comparative studies for each test. The low number of included studies for data analyses lowers the precision of the data. There are several other aspects that also need attention or additional research in order to determine the most suitable diagnostic method. The low prevalence of true scaphoid fractures among suspected fractures must be emphasised. The relatively low specificity of BS means that the number of over-treated patients would be much higher than with CT or MRI.

The effect of the low prevalence (20%) of true fractures among suspected scaphoid fractures is clearer when we apply the diagnostic accuracies in a cohort of 1000 patients ([Summary of findings 1](#)). BS has a higher sensitivity and would lead to only 2 missed fractures in a cohort of 1000 patients, compared with 56 and 24 missed fractures by CT and MRI, respectively. The relatively low specificity of BS would result in unnecessary treatment of 112 patients, compared with only 8 over-treated patients when diagnosis is performed using CT and none when diagnosis is performed using MRI. Although we could not detect statistically significant differences between the specificities of all three modalities, the clinical impact of lower specificity combined with the low prevalence of a fracture is substantial. This shows the challenges in the diagnostic management of scaphoid fractures. A possible way to improve the diagnostic accuracy and lower the impact on clinical practice is by raising the prevalence of true fractures among suspected fractures. This can be achieved by improving clinical evaluation or initial radiographic assessment, or both.

An interesting finding was the number of other fractures reported by all three imaging modalities. This review is focused on the scaphoid, but carpal and distal radius fractures were frequently found. The clinical significance for detecting these fractures is unknown, but does emphasise the questionable accuracy of current initial diagnostic methods.

Moreover, BS is the most invasive method to use with the intravenous application of radioactive isotopes and, compared with CT, gives a much higher dose of radiation. Therefore, BS is generally not recommended for children. BS also requires a delay of at least 72 hours to capture the osteoblastic activity at the fracture site and is therefore not applicable for instant diagnosis. Therefore, while BS might be the imaging modality with the highest sensitivity, it may not be the most suitable in practice.

AUTHORS' CONCLUSIONS

Implications for practice

The diagnostic accuracy (DOR) of all three modalities studied in this review is considered good. However, we found evidence that BS has a significantly higher diagnostic accuracy than CT and MRI. In the meta-analysis, BS shows better sensitivity than CT and MRI. However, BS is also characterised by a lower specificity than either CT or MRI. The number of studies included is small and the confidence intervals for summary estimates are wide for all three tests. Even fewer studies directly compared index tests. This reduces the precision and generalisability of our results. The more invasive aspects of BS need also to be borne in mind. This test is less favourable compared with CT and MRI in terms of timing and safety due to a diagnostic delay of more than 72 hours and the intravenous administration of radioactive isotopes. It is debatable whether sensitivity or specificity is more important in this scenario. With the big impact of over-treatment due to the relatively low specificity and with the invasive character of BS in mind, we would not recommend performing BS. CT and MRI both have good and comparable diagnostic accuracies, as shown in both meta-analyses and direct comparative studies. Given the data do not discriminate between the use of these tests, either of these tests can be used where available.

Implications for research

Prospective studies, perhaps involving randomisation of diagnostic tests, with direct comparisons of CT and MRI in the same patient population would add valuable data. We question the need for further research evaluating BS because of its limited use and invasive character. It would be useful if such studies incorporated economic (direct and indirect costs) and patient-related outcome measures (e.g. Disabilities of the Arm Shoulder and Hand, Patient Related Wrist Evaluation). Given the debate on the current best available reference standard (six week radiographs), consideration should be given to the practicalities of a check radiological follow-up, perhaps at one year, to examine for missed fractures. Prior to these, studies looking at ways to improve initial diagnostic management are needed to increase the identification of true scaphoid fractures among clinically suspected fractures.

ACKNOWLEDGEMENTS

We would like to thank Helen Handoll, the editors of the Cochrane Diagnostic Test Accuracy Working Group (particularly Petra Macaskill (review) and Gianni Virgili (protocol)), Peter Amadio, Lindsey Elstub, Laura MacDonald and Steven Rhemrev for their constructive feedback, comments and support in drafting the protocol and review. We also thank Joanne Elliott and the librarian of the VUMC of Amsterdam, Marijke Mol, for their support in developing the search strategies and running searches.

This project was supported by the National Institute for Health Research (NIHR) via Cochrane Infrastructure funding to the Cochrane Bone, Joint and Muscle Trauma Group, and the Cochrane

Diagnostic Test Accuracy Working Group. The views and opinions expressed herein are those of the review authors and do not necessarily reflect those of the Systematic Reviews Programme, NIHR or the UK National Health Service or Department of Health.

REFERENCES

References to studies included in this review

Beeres 2008 {published data only}

Beeres FJ, Rhemrev SJ, den Hollander P, Kingma LM, Meylaerts SA, le Cessie S, et al. Early magnetic resonance imaging compared with bone scintigraphy in suspected scaphoid fractures. *Journal of Bone and Joint Surgery. British volume* 2008;**90**(9):1205-9. [PUBMED: 18757961]

Breitenseher 1997 {published data only}

Breitenseher. MRI in radiologically occult scaphoid fractures. Initial experiences with 1.0 Tesla (whole body-middle field equipment) versus 0.2 Tesla (dedicated low-field equipment) [MRT bei radiologisch okkulten Kahnbeinfrakturen. Erste Erfahrungen von 1.0 Tesla (Ganzkörper-Mittelfeldgerät) versus 0.2 Tesla (dediziertes Niederfeldgerät)]. *Der Radiologe* 1997;**37**(10):812-8.

* Breitenseher MJ, Metz VM, Gilula LA, Gaebler C, Kukla C, Fleischmann D, et al. Radiographically occult scaphoid fractures: value of MR imaging in detection. *Radiology* 1997;**203**(1):245-50. [PUBMED: 9122402]

Gaebler C, Kukla C, Breitenseher MJ, Trattig S, Mittlboeck M, Vecsei V. Magnetic resonance imaging of occult scaphoid fractures. *Journal of Trauma* 1996;**41**(1):73-6.

Kukla C, Gaebler C, Breitenseher MJ, Trattig S, Vecsei V. Occult fractures of the scaphoid: The diagnostic usefulness and indirect economic repercussions of radiography versus magnetic resonance scanning. *Journal of Hand Surgery - British volume* 1997;**22B**(6):810-3.

Kukla C, Gaebler C, Breitenseher MJ, Trattig S, Vecsei V. Prospective comparison of MRI vs. direct magnification radiography in occult fractures of the scaphoid bone [Prospektiver Vergleich MRT vs. direkte Vergrößerungsradiographie bei okkulten Frakturen des Os scaphoideum]. *Unfallchirurg* 1998;**101**(1):32-6.

De Zwart 2012 {published data only}

* De Zwart A, Rhemrev SJ, Kingma LM, Meylaerts SA, Arndt JW, Schipper IB, et al. Early CT compared with bone scintigraphy in suspected scaphoid fractures. *Clinical Nuclear Medicine* 2012; Vol. 37, issue 10:981. [PUBMED: 22955071]

Rhemrev SJ, De Zwart AD, Kingma LM, Meylaerts SAG, Arndt JW, Schipper IB, et al. Early computed tomography compared with bone scintigraphy in suspected scaphoid fractures. *Clinical Nuclear Medicine* 2010;**35**(12):931-4.

Rhemrev SJ, van Leerdam RH, Beeres FJP, Arndt J-W, Meylaerts SAG. Bone scintigraphy in patients with clinical suspicion of a scaphoid fracture. *European Journal of Emergency Medicine* 2010;**17**(2):124-5.

Ilica 2011 {published data only}

Ilica AT, Ozyurek S, Kose O, Durusu M. Diagnostic accuracy of multidetector computed tomography for patients with suspected scaphoid fractures and negative radiographic

examinations. *Japanese Journal of Radiology* 2011;**29**(2):98-103. [PUBMED: 21359934]

Mallee 2011 {published data only}

Mallee W, Doornberg JN, Ring D, van Dijk CN, Maas M, Goslings JC. Comparison of CT and MRI for diagnosis of suspected scaphoid fractures. *Journal of Bone and Joint Surgery. American Volume* 2011;**93**(1):20-8. [PUBMED: 21209265]

Memarsadeghi 2006 {published data only}

Memarsadeghi M, Breitenseher MJ, Schaefer-Prokop C, Weber M, Aldrian S, Gabler C, et al. Occult scaphoid fractures: comparison of multidetector CT and MR imaging--initial experience. *Radiology* 2006;**240**(1):169-76. [PUBMED: 16793977]

Nielsen 1983 {published data only}

Nielsen PT, Hedeboe J, Thommesen P. Bone scintigraphy in the evaluation of fracture of the carpal scaphoid bone. *Acta Orthopaedica Scandinavica* 1983;**54**(2):303-6. [PUBMED: 6221504]

O'Carroll 1982 {published data only}

O'Carroll PF, Doyle J, Duffy G. Radiography and scintigraphy in the diagnosis of carpal scaphoid fractures. *Irish Journal of Medical Science* 1982;**151**(7):211-3. [PUBMED: 7129827]

Stordahl 1984 {published data only}

Stordahl A, Schjoth A, Woxholt G, Fjermeros H. Bone scanning of fractures of the scaphoid. *Journal of Hand Surgery* 1984;**9**(2):189-90. [PUBMED: 6747427]

Tiel-van Buul 1993 {published data only}

* Tiel-van Buul MM, van Beek EJ, Broekhuizen AH, Bakker AJ, Bos KE, van Royen EA. Radiography and scintigraphy of suspected scaphoid fracture. A long-term study in 160 patients. *Journal of Bone and Joint Surgery. British volume* 1993;**75**(1):61-5. [PUBMED: 8421037]

Tiel-van Buul MMC, van Beek EJR, Dijkstra PF, Bakker AJ, Broekhuizen TH, van Royen EA. Significance of a hot spot on the bone scan after carpal injury: evaluation by computed tomography. *European Journal of Nuclear Medicine* 1993;**20**:159-64.

Tiel-van Buul 1996 {published data only}

Tiel-van Buul MM, Roolker W, Verbeeten BW, Broekhuizen AH. Magnetic resonance imaging versus bone scintigraphy in suspected scaphoid fracture. *European Journal of Nuclear Medicine* 1996;**23**(8):971-5. [PUBMED: 8753688]

References to studies excluded from this review

Adey 2007 {published data only}

Adey L, Souer JS, Lozano-Calderon S, Palmer W, Lee SG, Ring D. Computed tomography of suspected scaphoid fractures. *Journal of Hand Surgery* 2007;**32**(1):61-6. [PUBMED: 17218177]

Akdemir 2004 {published data only}

Akdemir UO, Atasever T, Sipahioğlu S, Turkolmez S, Kazimoglu C, Sener E. Value of bone scintigraphy in patients with carpal trauma. *Annals of Nuclear Medicine* 2004;**18**(6):495-9. [PUBMED: 15515749]

Asche 1982 {published data only}

Asche G. Value of bone scintigraphy in the recognition of scaphoid bone fracture difficult to detect roentgenologically [Der Wert der Knochenszintigraphie bei der Erkennung der röntgenologisch schwer sichtbaren Kahnbeinfraktur]. *Handchirurgie, Mikrochirurgie, Plastische Chirurgie* 1982;**14**(2):114-7. [PUBMED: 7182293]

Beeres 2005 {published data only}

Beeres FJ, Hogervorst M, den Hollander P, Rhemrev S. Outcome of routine bone scintigraphy in suspected scaphoid fractures. *Injury* 2005;**36**(10):1233-6. [PUBMED: 15979624]

Beeres 2007 {published data only}

Beeres FJ, Hogervorst M, Rhemrev SJ, den Hollander P, Jukema GN. A prospective comparison for suspected scaphoid fractures: bone scintigraphy versus clinical outcome. *Injury* 2007;**38**(7):769-74. [PUBMED: 17307177]

Breederveld 2004 {published data only}

Breederveld RS, Tuinebreijer WE. Investigation of computed tomographic scan concurrent criterion validity in doubtful scaphoid fracture of the wrist. *Journal of Trauma* 2004;**57**(4):851-4. [PUBMED: 15514541]

Bretlau 1999 {published data only}

Bretlau T, Christensen OM, Edstrom P, Thomsen HS, Lausten GS. Diagnosis of scaphoid fracture and dedicated extremity MRI. *Acta Orthopaedica Scandinavica* 1999;**70**(5):504-8. [PUBMED: 10622486]

Brismar 1988 {published data only}

Brismar J. Skeletal scintigraphy of the wrist in suggested scaphoid fracture. *Acta Radiologica* 1988;**29**(1):101-7. [PUBMED: 2964829]

Bury 2010 {published data only}

Bury R. Detection of occult wrist fractures using quantitative scintigraphy. *Nuclear Medicine Communications* 2010;**31**(2):178; author reply 178-9. [PUBMED: 20032769]

Cook 1997 {published data only}

Cook PA, Yu JS, Wiand W, Cook AJ 2nd, Coleman CR, Cook AJ. Suspected scaphoid fractures in skeletally immature patients: application of MRI. *Journal of Computer Assisted Tomography* 1997;**21**(4):511-5. [PUBMED: 9216753]

Coupland 1996 {published data only}

Coupland DB. Determining the presence of scaphoid fracture with a day 4 bone scan. *Clinical Journal of Sport Medicine* 1996;**6**(2):137. [PUBMED: 8673573]

Cruikshank 2007 {published data only}

Cruikshank J, Meakin A, Breadmore R, Mitchell D, Pincus S, Hughes T, et al. Early computerized tomography accurately

determines the presence or absence of scaphoid and other fractures. *Emergency Medicine Australasia* 2007;**19**(3):223-8. [PUBMED: 17564689]

Fowler 1998 {published data only}

Fowler C, Sullivan B, Williams LA, McCarthy G, Savage R, Palmer A. A comparison of bone scintigraphy and MRI in the early diagnosis of the occult scaphoid waist fracture. *Skeletal Radiology* 1998;**27**(12):683-7. [PUBMED: 9921930]

Fusetti 2008 {published data only}

Fusetti C, Garavaglia G. A prospective comparison for suspected scaphoid fractures: bone scintigraphy versus clinical outcome. *Injury* 2008; Vol. 39, issue 4:500-1. [PUBMED: 18314126]

Gäbler 2001 {published data only}

Gäbler C, Kukla C, Breitenseher MJ, Trattnig S, Vecsei V. Diagnosis of occult scaphoid fractures and other wrist injuries. Are repeated clinical examinations and plain radiographs still state of the art?. *Langenbeck's Archives of Surgery / Deutsche Gesellschaft für Chirurgie* 2001;**386**(2):150-4. [PUBMED: 11374049]

Ganel 1979 {published data only}

Ganel A, Engel J, Oster Z, Farine I. Bone scanning in the assessment of fractures of the scaphoid. *Journal of Hand Surgery* 1979;**4**(6):540-3. [PUBMED: 512312]

Geijer 2011 {published data only}

Geijer M, Borjesson AM, Gothlin JH. Clinical utility of tomosynthesis in suspected scaphoid fracture. A pilot study. *Skeletal Radiology* 2011;**40**(7):863-7. [PUBMED: 21057785]

Hobby 2001 {published data only}

Hobby JL, Tom BD, Bearcroft PW, Dixon AK. Magnetic resonance imaging of the wrist: diagnostic performance statistics. *Clinical Radiology* 2001;**56**(1):50-7. [PUBMED: 11162698]

Hunter 1997 {published data only}

Hunter JC, Escobedo EM, Wilson AJ, Hanel DP, Zink-Brody GC, Mann FA. MR imaging of clinically suspected scaphoid fractures. *AJR. American Journal of Roentgenology* 1997;**168**(5):1287-93. [PUBMED: 9129428]

Jenkins 2008 {published data only}

Jenkins PJ, Slade K, Huntley JS, Robinson CM. A comparative analysis of the accuracy, diagnostic uncertainty and cost of imaging modalities in suspected scaphoid fractures. *Injury* 2008;**39**(7):768-74. [PUBMED: 18541243]

Jonsson 1992 {published data only}

Jonsson K, Jonsson A, Sloth M, Kopylov P, Wingstrand H. CT of the wrist in suspected scaphoid fracture. *Acta Radiologica* 1992;**33**(5):500-1. [PUBMED: 1389664]

Keim 2003 {published data only}

Keim S, Mack M, Vogl T, Windolf J. Clinical impact of MRI in acute wrist injuries. Indication and therapeutic consequences [Klinische Relevanz der MRT bei akuten Handgelenkverletzungen. Indikation und therapeutische

Konsequenz]. *Der Unfallchirurg* 2003;**106**(2):127-35. [PUBMED: 12624687]

Klein 2002 {published data only}

Klein HM, Vrsalovic V, Balas R, Neugebauer F. Imaging diagnostics of the wrist: MRI and Arthrography/Arthro-CT [Bildgebende Diagnostik des Handgelenkes: MRT und Arthrographie/Arthro-CT]. *RoFo : Fortschritte auf dem Gebiete der Röntgenstrahlen und der Nuklearmedizin* 2002;**174**(2):177-82. [PUBMED: 11898079]

Kristensen 1983 {published data only}

Kristensen J, Carlsen O, Kirkeby HJ, Noreng M, Thousig OP. Bone scintigraphy in suspected fracture of the scaphoid bone [Knoglescintigrafi ved mistanke om fraktur af os scaphoideum]. *Ugeskrift for Laeger* 1983;**145**(34):2591-4. [PUBMED: 6612886]

Kumar 2005 {published data only}

Kumar S, O'Connor A, Despois M, Galloway H. Use of early magnetic resonance imaging in the diagnosis of occult scaphoid fractures: the CAST Study (Canberra Area Scaphoid Trial). *New Zealand Medical Journal* 2005;**118**(1209):U1296. [PUBMED: 15711629]

Kusano 2002 {published data only}

Kusano N, Churei Y, Shiraishi E, Kusano T. Diagnosis of occult carpal scaphoid fracture: a comparison of magnetic resonance imaging and computed tomography techniques. *Techniques in Hand & Upper Extremity Surgery* 2002;**6**(3):119-23. [PUBMED: 16520609]

Lanik 1986 {published data only}

Lanik R, Stanek O. Possibilities of scintigraphy in the early diagnosis of fracture of the scaphoid bone in the hand [Moznosti scintigrafie v casne diagnostice zlomenin clunkove kosti ruky]. *Ceskoslovenska Radiologie* 1986;**40**(5):315-22. [PUBMED: 3502957]

Lapa 2000 {published data only}

Lapa P, Ferrer-Antunes A, Lima J. Bone scintigraphy in the diagnosis of scaphoid fracture [Cintigrafia ossea no diagnostico de fractura do escafoide]. *Revista Espanola de Medicina Nuclear* 2000;**19**(7):504. [PUBMED: 11171509]

Lepage 2004 {published data only}

* Lepage D, Obert L, Garbuio P, Tropet Y, Paratte B, Runge M, et al. Contribution of quantitative radio-scintigraphy to diagnosis of wrist injuries undetected on plain films: a prospective study of 154 cases [Apport de la radioscintigraphie quantitative dans les traumatismes du poignet avec radiographies initiales normales: etude prospective de 154 cas]. *Revue de Chirurgie Orthopedique et Reparatrice de l'Appareil Moteur* 2004;**90**(6):542-9. [PUBMED: 15672921]

Vichard P, Garbuio P, Lepage D, Tropet Y. Occult fractures of the carpal navicular. Detection by quantitative radiosintigraphy. Social and medico-legal repercussions [Les fractures occultes du scaphoide carpien. Detection par la radiosintigraphie quantitative. Incidences sociales et medico-legales]. *Bulletin de l'Academie Nationale de Medecine* 2001;**185**(8):1399-1413.

Lindequist 1998 {published data only}

Lindequist S, Houshian S, Rosbach SB, Riegels-Nielsen P. MRI scanning in suspected fractures of the scaphoid bone [MR-scanning ved mistanke om fraktur af os scaphoideum]. *Ugeskrift for Laeger* 1998;**160**(51):7438-41. [PUBMED: 9889659]

Low 2005 {published data only}

Low G, Raby N. Can follow-up radiography for acute scaphoid fracture still be considered a valid investigation?. *Clinical Radiology* 2005;**60**(10):1106-10. [PUBMED: 16179171]

Lozano-Calderon 2006 {published data only}

Lozano-Calderon S, Blazar P, Zurakowski D, Lee SG, Ring D. Diagnosis of scaphoid fracture displacement with radiography and computed tomography. *Journal of Bone and Joint Surgery. American Volume* 2006;**88**(12):2695-703. [PUBMED: 17142420]

Moller 2004 {published data only}

Moller JM, Larsen L, Bovin J, Lausten GS, Hasselqvist M, Jensen CM, et al. MRI diagnosis of fracture of the scaphoid bone: impact of a new practice where the images are read by radiographers. *Academic Radiology* 2004;**11**(7):724-8. [PUBMED: 15217588]

Murphy 1995 {published data only}

Murphy DG, Eisenhauer MA, Powe J, Pavlofsky W. Can a day 4 bone scan accurately determine the presence or absence of scaphoid fracture?. *Annals of Emergency Medicine* 1995;**26**(4):434-8. [PUBMED: 7574124]

Querellou 2009 {published data only}

Querellou S, Moineau G, Le Duc-Pennec A, Guillo P, Turzo A, Cotonea Y, et al. Detection of occult wrist fractures by quantitative radiosintigraphy: a prospective study on selected patients. *Nuclear Medicine Communications* 2009;**30**(11):862-7. [PUBMED: 19668095]

Raby 2001 {published data only}

Raby N. Magnetic resonance imaging of suspected scaphoid fractures using a low field dedicated extremity MR system. *Clinical Radiology* 2001;**56**(4):316-20. [PUBMED: 11286584]

Schick 1999 {published data only}

Schick S, Trattinig S, Gabler C, Kukla C, Gahleitner A, Kainberger F, et al. Occult fractures of the wrist joint: high resolution image magnification roentgen versus MRI [Okkulte Handgelenksfrakturen: Feinfokusvergrößerungsrontgen versus MRT]. *RoFo : Fortschritte auf dem Gebiete der Röntgenstrahlen und der Nuklearmedizin* 1999;**170**(1):16-21. [PUBMED: 10071639]

Tennoe 2000 {published data only}

Tennoe B, Stiris MG. MR imaging in suspected acute trauma of wrist bones. *Acta Radiologica* 2000; Vol. 41, issue 3:303. [PUBMED: 10866091]

Thorpe 1996 {published data only}

Thorpe AP, Murray AD, Smith FW, Ferguson J. Clinically suspected scaphoid fracture: a comparison of magnetic resonance imaging and bone scintigraphy. *British Journal of Radiology* 1996;**69**(818):109-13. [PUBMED: 8785636]

Tiel-van Buul 1993b {published data only}

Tiel-van Buul MM, van Beek EJ, Borm JJ, Gubler FM, Broekhuizen AH, van Royen EA. The value of radiographs and bone scintigraphy in suspected scaphoid fracture. A statistical analysis. *Journal of Hand Surgery* 1993;**18**(3):403-6. [PUBMED: 8345279]

Tiel-van Buul 1995 {published data only}

Tiel-van Buul MM, Broekhuizen TH, van Beek EJ, Bossuyt PM. Diagnosis of scaphoid fractures: the role of nuclear medicine. *Journal of nuclear medicine : official publication, Society of Nuclear Medicine* 1995; Vol. 36, issue 9:1725-6. [PUBMED: 7658240]

Tiel-van Buul 1998 {published data only}

Tiel-van Buul MM, van Beek EJ. Value of MR imaging in the detection of occult scaphoid fractures. *Radiology* 1998;**206**(1):291-2. [PUBMED: 9423687]

Vrettos 1996 {published data only}

Vrettos BC, Adams BK, Knottenbelt JD, Lee A. Is there a place for radionuclide bone scintigraphy in the management of radiograph-negative scaphoid trauma?. *South African Medical Journal = Suid-Afrikaanse Tydskrif vir Geneeskunde* 1996;**86**(5):540-2. [PUBMED: 8711552]

Wilson 1986 {published data only}

Wilson AW, Kurer MH, Peggington JL, Grant DS, Kirk CC. Bone scintigraphy in the management of X-ray-negative potential scaphoid fractures. *Archives of Emergency Medicine* 1986;**3**(4):235-42. [PUBMED: 3801112]

You 2007 {published data only}

You JS, Chung SP, Chung HS, Park IC, Lee HS, Kim SH. The usefulness of CT for patients with carpal bone fractures in the emergency department. *Emergency Medicine Journal* 2007;**24**(4):248-50. [PUBMED: 17384376]

Additional references
Biswas 2009

Biswas D, Bible JE, Bohan M, Simpson AK, Whang PG, Grauer JN. Radiation exposure from musculoskeletal computerized tomographic scans. *Journal of Bone and Joint Surgery. American Volume* 2009;**91**(8):1882-9. [PUBMED: 19651945]

Bond 2001

Bond CD, Shin AY, McBride MT, Dao KD. Percutaneous screw fixation or cast immobilization for nondisplaced scaphoid fractures. *Journal of Bone and Joint Surgery. American Volume* 2001;**83**(4):483-8. [PUBMED: 11315775]

Brookes-Fazakerley 2009

Brookes-Fazakerley SD, Kumar AJ, Oakley J. Survey of the initial management and imaging protocols for occult scaphoid fractures in UK hospitals. *Skeletal Radiology* 2009;**38**(11):1045-8. [PUBMED: 19194702]

Buijze 2010

Buijze GA, Doornberg JN, Ham JS, Ring D, Bhandari M, Poolman RW. Surgical compared with conservative treatment for acute nondisplaced or minimally displaced scaphoid fractures: a systematic review and meta-analysis of randomized controlled trials. *Journal of Bone and Joint Surgery. American Volume* 2010;**92**(6):1534-44. [PUBMED: 20516332]

de Vet 2008

de Vet HCW, Eisinga A, Riphagen II, Aertgeerts B, Pewsner D. Chapter 7: Searching for studies. *Cochrane Handbook for Systematic Reviews of Diagnostic Test Accuracy Version 0.4* (updated September 2008). The Cochrane Collaboration, 2008.

Dennis 2011

Dennis HH, Sze AC, Murphy D. Prevalence of carpal fracture in Singapore. *Journal of Hand Surgery. American Volume* 2011;**36**(2):278-83. [PUBMED: 21276891]

Dias 2005

Dias JJ, Wildin CJ, Bhowal B, Thompson JR. Should acute scaphoid fractures be fixed? A randomized controlled trial. *Journal of Bone and Joint Surgery. American Volume* 2005;**87**(10):2160-8. [PUBMED: 16203878]

Dorsay 2001

Dorsay TA, Major NM, Helms CA. Cost-effectiveness of immediate MR imaging versus traditional follow-up for revealing radiographically occult scaphoid fractures. *AJR. American Journal of Roentgenology* 2001;**177**(6):1257-63. [PUBMED: 11717059]

Fowler 2010

Fowler JR, Ilyas AM. Headless compression screw fixation of scaphoid fractures. *Hand Clinics* 2010;**26**(3):351-61, vi. [PUBMED: 20670800]

Geissler 2012

Geissler WB, Adams JE, Bindra RR, Lanzinger WD, Slutsky DJ. Scaphoid fractures: what's hot, what's not. *Instructional Course Lectures* 2012;**61**:71-84. [PUBMED: 22301223]

Gelberman 1986

Gelberman RH, Gross MS. The vascularity of the wrist. Identification of arterial patterns at risk. *Clinical Orthopaedics and Related Research* 1986;**(202)**:40-9. [PUBMED: 3514029]

Groves 2006

Groves AM, Kayani I, Syed R, Hutton BF, Bearcroft PP, Dixon AK, et al. An international survey of hospital practice in the imaging of acute scaphoid trauma. *AJR. American Journal of Roentgenology* 2006;**187**(6):1453-6. [PUBMED: 17114536]

Hove 1999

Hove LM. Epidemiology of scaphoid fractures in Bergen, Norway. *Scandinavian Journal of Plastic and Reconstructive Surgery and Hand Surgery / Nordisk Plastikkirurgisk Forening [and] Nordisk Klubb for Handkirurgi* 1999;**33**(4):423-6. [PUBMED: 10614752]

McDougall 1989

McDougall IR, Rieser RP. Scintigraphic techniques in musculoskeletal trauma. *Radiologic Clinics of North America* 1989;**27**(5):1003-11. [PUBMED: 2672079]

Merrell 2002

Merrell GA, Wolfe SW, Slade JF 3rd. Treatment of scaphoid nonunions: quantitative meta-analysis of the literature. *Journal of Hand Surgery. American Volume* 2002;**27**(4):685-91. [PUBMED: 12132096]

Pillai 2005

Pillai A, Jain M. Management of clinical fractures of the scaphoid: results of an audit and literature review. *European Journal of Emergency Medicine* 2005;**12**(2):47-51. [PUBMED: 15756078]

Rettig 1999

Rettig ME, Raskin KB. Retrograde compression screw fixation of acute proximal pole scaphoid fractures. *Journal of Hand Surgery* 1999;**24**(6):1206-10. [PUBMED: 10584942]

Rhemrev 2010a

Rhemrev SJ, Beeres FJ, van Leerdam RH, Hogervorst M, Ring D. Clinical prediction rule for suspected scaphoid fractures a prospective cohort study. *Injury* 41;**10**:1026-30. [PUBMED: 20510414]

Rhemrev 2010b

Rhemrev SJ, de Zwart AD, Kingma LM, Meylaerts SA, Arndt JW, Schipper IB, et al. Early computed tomography compared with bone scintigraphy in suspected scaphoid fractures. *Clinical Nuclear Medicine* 2010;**35**(12):931-4. [PUBMED: 21206223]

Rhemrev 2011

Rhemrev SJ, Ootes D, Beeres FJ, Meylaerts SA, Schipper IB. Current methods of diagnosis and treatment of scaphoid fractures. *International Journal of Emergency Medicine* 2011;**4**:4. [PUBMED: 21408000]

Ring 2008

Ring D, Lozano-Calderon S. Imaging for suspected scaphoid fracture. *Journal of Hand Surgery. American Volume* 2008;**33**(6):954-7. [PUBMED: 18656772]

Roolker 1997

Roolker W, Tiel-van Buul MM, Ritt MJ, Verbeeten B Jr, Griffioen FM, Broekhuizen AH. Experimental evaluation of scaphoid X-series, carpal box radiographs, planar tomography, computed tomography, and magnetic resonance imaging in the diagnosis of scaphoid fracture. *Journal of Trauma* 1997;**42**(2):247-53. [PUBMED: 9042876]

Sanders 1988

Sanders WE. Evaluation of the humpback scaphoid by computed tomography in the longitudinal axial plane of the scaphoid. *Journal of Hand Surgery. American Volume* 1988;**13**(2):182-7. [PUBMED: 3351239]

Senall 2004

Senall JA, Failla JM, Bouffard JA, van Holsbeeck M. Ultrasound for the early diagnosis of clinically suspected scaphoid fracture. *Journal of Hand Surgery. American Volume* 2004;**29**(3):400-5. [PUBMED: 15140480]

Thavarajah 2011

Thavarajah D, Syed T, Shah Y, Wetherill M. Does scaphoid bone bruising lead to occult fracture? A prospective study of 50 patients. *Injury* 2011;**42**(11):1303-6. [PUBMED: 21497809]

Ty 2008

Ty JM, Lozano-Calderon S, Ring D. Computed tomography for triage of suspected scaphoid fractures. *Hand* 2008;**3**(2):155-8. [PUBMED: 18780093]

Unay 2009

Unay K, Gokcen B, Ozkan K, Poyanli O, Eceviz E. Examination tests predictive of bone injury in patients with clinically suspected occult scaphoid fracture. *Injury* 2009;**40**(12):1265-8. [PUBMED: 19524914]

Van der Molen 1999

Van der Molen AB, Groothoff JW, Visser GJ, Robinson PH, Eisma WH. Time off work due to scaphoid fractures and other carpal injuries in The Netherlands in the period 1990 to 1993. *Journal of Hand Surgery. British Volume* 1999;**24**(2):193-8. [PUBMED: 10372775]

Van Onselen 2003

Van Onselen EB, Karim RB, Hage JJ, Ritt MJ. Prevalence and distribution of hand fractures. *Journal of Hand Surgery. British Volume* 2003;**28**(5):491-5. [PUBMED: 12954264]

Vinnars 2008

Vinnars B, Pietreanu M, Bodestedt A, Ekenstam F, Gerdin B. Nonoperative compared with operative treatment of acute scaphoid fractures. A randomized clinical trial. *Journal of Bone and Joint Surgery. American Volume* 2008;**90**(6):1176-85. [PUBMED: 18519309]

Whiting 2003

Whiting P, Rutjes AW, Reitsma JB, Bossuyt PM, Kleijnen J. The development of QUADAS: a tool for the quality assessment of studies of diagnostic accuracy included in systematic reviews. *BMC Medical Research Methodology* 2003;**3**:25. [PUBMED: 14606960]

Yin 2010

Yin ZG, Zhang JB, Kan SL, Wang XG. Diagnosing suspected scaphoid fractures: a systematic review and meta-analysis. *Clinical Orthopaedics and Related Research* 2010;**468**(3):723-34. [PUBMED: 19756904]

* Indicates the major publication for the study

CHARACTERISTICS OF STUDIES

Characteristics of included studies [ordered by study ID]

Beeres 2008

Clinical features and settings	<p>Inclusion criteria: presentation to the ED within 48 hours after trauma. Clinically suspected for scaphoid fracture: tenderness ASB and painful ASB on longitudinal compression of thumb or index fingers. No fracture on initial radiographs in three views (PA, Lat, oblique with ulnar deviation)</p> <p>Exclusion criteria: polytrauma patients, patients under the age of 18 years, and those in whom MRI was contraindicated</p>
Participants	<p>Study location: The Hague, The Netherlands</p> <p>Study period: March 2004 to January 2007</p> <p>Participants enrolled: 100; sex: 50 men and 50 women; mean age 42 years (range 18 to 84)</p> <p>Participants included in analyses: 100</p>
Study design	Prospective, consecutive cohort
Target condition and reference standard(s)	<p>Target condition(s): true scaphoid fracture among clinically suspected scaphoid fractures</p> <p>Reference test: MRI & BS (-) = no fracture; MRI & BS (+) = fracture. If discrepancy between MRI & BS, 6-week follow-up radiographs including physical examination</p>
Index and comparator tests	<p>Index test(s): MRI and BS</p> <p>Magnetic resonance imaging</p> <p><i>Timing:</i> performed within 24 hours after presentation to the ED</p> <p><i>Description of imaging technique (including patient position):</i> 1.5 Tesla with flexible surface coil around the wrist. Patient in prone position with the injured hand extended forward, palm down and overhead. T1 and T2 images were obtained. A slice thickness of 3 mm and distance factor of 10% was used</p> <p><i>Interpretation:</i> images were evaluated by a trainee and a consultant radiologist</p> <p><i>Criteria for a positive test:</i> not given</p> <p>Bone scintigraphy</p> <p><i>Timing:</i> performed within 3 to 5 days after presentation to the ED</p> <p><i>Description of imaging technique (including patient position):</i> palmar and dorsal images of both wrists between 2.5 and 4 hours after injection of 500 MBq 99mTc-HDP</p> <p><i>Interpretation:</i> images were evaluated by a consultant clinical nuclear physician</p> <p><i>Criteria for a positive test:</i> osteoblast activity</p>
Follow-up	None reported after final reference standard test was performed
Notes	<p>Prevalence of scaphoid fracture: 20% (20/100)</p> <p>Reporting of other fractures: MRI reported 24 other fractures, BS reported 40 other fractures, reference standard reported 32 other fractures</p>

Table of Methodological Quality

Item	Authors' judgement	Description
------	--------------------	-------------

Beeres 2008 (Continued)

Representative spectrum? All tests	Yes	Consecutive series of patients with acute wrist injury, clinical signs for a scaphoid fracture and normal initial radiographs
Acceptable reference standard? All tests	Yes	Reference standard was adequate according to the requirements mentioned in our protocol: 2 index tests with the same outcome, or 6-week follow-up radiographs if there was a discrepancy
Acceptable delay between tests? All tests	Yes	Reference standard was either same outcome in 2 index tests or radiographs obtained after 6 weeks, which was an acceptable interval
Partial verification avoided? All tests	Yes	All patients received the intended reference standard
Differential verification avoided? All tests	No	Different reference standards were used. If there was discrepancy between test outcomes, 6-week follow-up radiographs were used
Incorporation avoided? All tests	No	Index tests were part of the reference standard
Index test results blinded? All tests	Yes	All images were separately evaluated in a blinded fashion
Reference standard results blinded? All tests	Yes	All images were separately evaluated in a blinded fashion
Relevant clinical information? All tests	No	Clinical information was not present during evaluation of the images
Uninterpretable results reported? All tests	Yes	All outcomes of index tests and reference standard were described; no uninterpretable test results were present
Withdrawals explained? All tests	Yes	All patients underwent the entire study protocol
Fracture criteria defined? All tests	Unclear	No fracture criteria for MRI or radiographs, however evaluation was performed by 2 observers. Criteria for BS were given

Breitenseher 1997

Clinical features and settings	<p>Inclusion criteria: presentation to the ED after acute trauma. Clinically suspected for scaphoid fracture: pain, swelling and tenderness of ASB during evaluation in the ED. No fracture on initial radiographs in 6 views (AP, 2 x Lat, PA with ulnar deviation, 2 x oblique)</p> <p>Exclusion criteria: fracture on initial radiographs</p>	
Participants	<p>Study location: Vienna, Austria</p> <p>Study period: January 1995 to March 1996</p> <p>Participants enrolled: 42: 23 men and 19 women. Mean age 30.5 (range 10 to 66)</p>	

Breitenseher 1997 (Continued)

Participants included in analyses: 42

Study design	Prospective, consecutive cohort
Target condition and reference standard(s)	<p>Target condition(s): true scaphoid fracture among clinically suspected scaphoid fractures</p> <p>Reference test: 6-week follow-up radiographs compared with initial radiographs</p>
Index and comparator tests	<p>Index test: MRI</p> <p>Magnetic resonance imaging:</p> <p><i>Timing:</i> performed within 7 days (mean 3.8 days) after trauma</p> <p><i>Description of imaging technique (including patient position):</i> 1 Tesla with circular surface coil. T1, STIR (both section thickness of 2.4 mm and intersection gap of 0.3 mm) and T2 (section thickness of 1.5 mm) images were performed. Patient position unknown</p> <p><i>Interpretation:</i> images were evaluated by two experienced radiologists</p> <p><i>Criteria for a positive test:</i> evidence of cortical or trabecular fracture line, diffuse area of bone marrow oedema or combination of all</p>
Follow-up	None reported after final reference standard test was performed
Notes	<p>Prevalence of scaphoid fracture: 33% (14/42)</p> <p>Reporting of other fractures: MRI reported 7 other fractures, reference standard reported none.</p>

Table of Methodological Quality

Item	Authors' judgement	Description
Representative spectrum? All tests	Unclear	Unclear what the delay between wrist trauma and presentation to the ED was
Acceptable reference standard? All tests	Yes	Reference standard was adequate according to the requirements mentioned in the protocol: 6-week follow-up radiographs
Acceptable delay between tests? All tests	Yes	6 weeks for radiographs is an acceptable delay
Partial verification avoided? All tests	Yes	All patients received the intended reference standard
Differential verification avoided? All tests	Yes	All patients received the same reference standard: 6-week follow-up radiographs
Incorporation avoided? All tests	Yes	The index test, MRI, was not part of the reference standard
Index test results blinded? All tests	Yes	All images were evaluated in a blinded fashion
Reference standard results blinded?	Yes	All images were evaluated in a blinded fashion

Breitenseher 1997 (Continued)

All tests

Relevant clinical information? All tests	Unclear	It was unclear if clinical information was present during evaluation of MRI
Uninterpretable results reported? All tests	Yes	All outcomes of index tests and reference standard were described, no uninterpretable test results were present
Withdrawals explained? All tests	Yes	Consecutive series without loss to follow-up. All patients included in the study underwent both MRI and radiographs
Fracture criteria defined? All tests	Yes	Clear fracture criteria for both radiographs and MRI were given

De Zwart 2012

Clinical features and settings	<p>Inclusion criteria: presentation to the ED within 48 hours after trauma. Clinically suspected for scaphoid fracture: tenderness ASB and painful ASB on longitudinal compression of thumb or index finger. No fracture on initial radiographs in 3 views (PA, Lat, oblique with ulnar deviation)</p> <p>Exclusion criteria: polytrauma patients, patients younger than 18 years and those with contraindications for bone scintigraphy or CT</p>	
Participants	<p>Study location: The Hague, The Netherlands</p> <p>Study period: November 2007 to January 2011</p> <p>Participants enrolled: 159: 79 men and 80 women. Mean age 41 (range 17 to 88)</p> <p>Participants included in analyses: 159</p>	
Study design	Prospective, consecutive cohort	
Target condition and reference standard(s)	<p>Target condition(s): true scaphoid fracture among clinically suspected scaphoid fractures</p> <p>Reference test: CT & BS (-) = no fracture; CT & BS (+) = fracture. If discrepancy between CT & BS: 6-week follow-up radiographs including physical examination</p>	
Index and comparator tests	<p>Index test: CT and BS</p> <p>Computed tomography</p> <p><i>Timing:</i> performed within 24 hours after presentation to the ED</p> <p><i>Description of imaging technique (including patient position):</i> prone position with the hand extended forward palm down over the patient's head with the wrist in neutral flexion and neutral radial-ulnar deviation. Scout images were obtained to ensure that the scanning plane corresponded with the scans that provided a lateral view of the scaphoid bone as defined by the central longitudinal axis of the scaphoid. Coronal plane images defined as images that provided a posteroanterior view of the scaphoid in the anatomic plane and in line with the axis of the scaphoid were obtained by supinating the forearm 90° keeping the wrist in a neutral position. Slice thickness was 0.625 mm with reconstructions every 0.4 mm. For multiplanar reformatted images, parameters were 2 mm slice thickness, 2 mm interval</p> <p><i>Interpretation:</i> images were evaluated by a resident and a consultant radiologist</p> <p><i>Criteria for a positive test:</i> none given</p>	

De Zwart 2012 (Continued)

Bone scintigraphy

Timing: performed within 3 to 5 days after presentation to the ED

Description of imaging technique (including patient position): palmar and dorsal images of both wrists were performed between 2.5 and 4 hours after the intravenous injection of 500 MBq of Tc-99m-HDP visualizing the osteoblastic activity with a planar collimator

Interpretation: images were evaluated by a consultant clinical nuclear physician

Criteria for a positive test: osteoblastic activity

Follow-up	None reported after final reference standard test was performed
Notes	<p>Prevalence of scaphoid fracture: 13% (20/159)</p> <p>Reporting of other fractures: CT reported 35 other fractures, BS reported 57 other fractures, reference standard unknown.</p>

Table of Methodological Quality

Item	Authors' judgement	Description
Representative spectrum? All tests	Yes	Consecutive series of patients with acute wrist injury, clinical signs for a scaphoid fracture and normal initial radiographs
Acceptable reference standard? All tests	Yes	Reference standard was adequate according to the requirements mentioned in the protocol: 2 index tests with the same outcome, or 6-week follow-up radiographs if there was a discrepancy
Acceptable delay between tests? All tests	Yes	Reference standard was either same outcome in 2 index tests or radiographs obtained after 6 weeks, which was an acceptable interval
Partial verification avoided? All tests	Yes	All patients received the intended reference standard
Differential verification avoided? All tests	No	Different reference standards were used. If there was discrepancy between test outcomes, 6-week follow-up radiographs were used
Incorporation avoided? All tests	No	Index tests were part of the reference standard
Index test results blinded? All tests	Yes	All images were separately evaluated in a blinded fashion
Reference standard results blinded? All tests	Yes	All images were separately evaluated in a blinded fashion
Relevant clinical information? All tests	No	Clinical information was not present during evaluation of the images. Observers were blinded from all other data.
Uninterpretable results reported? All tests	Yes	All outcomes of index tests and reference standard were described, no uninterpretable test results were present

De Zwart 2012 (Continued)

Withdrawals explained? All tests	Yes	Patients that did not have both CT and BS were reported including explanations
Fracture criteria defined? All tests	Unclear	No fracture criteria for either CT or radiographs were given, however evaluation was performed by 2 observers. Criteria for BS were given

Ilica 2011

Clinical features and settings	<p>Inclusion criteria: presentation to the ED within 72 hours after trauma. Clinically suspected for scaphoid fracture: tenderness ASB and scaphoid's tubercle. No fracture on initial radiographs in 3 views (PA with ulnar deviation, Lat, oblique with wrist in semi-supination).</p> <p>Exclusion criteria: Patients admitted more than 72 hours after the trauma or less than 18 years of age.</p>	
Participants	<p>Study location: Ankara, Turkey</p> <p>Study period: December 2007 to November 2008</p> <p>Participants enrolled: 54 (55 wrists): all men. Mean age 22 (range 20 to 40)</p> <p>Participants included in analyses: 55 (wrists)</p>	
Study design	Prospective, consecutive cohort	
Target condition and reference standard(s)	<p>Target condition(s): True scaphoid fracture among clinically suspected scaphoid fractures.</p> <p>Reference test: MRI</p>	
Index and comparator tests	<p>Index test: CT</p> <p>Computed tomography</p> <p><i>Timing:</i> performed within 7 days after trauma</p> <p><i>Description of imaging technique (including patient position):</i> 64-detector multislice system. Prone position with the hand above the head and the wrist placed flat. Acquisition with 0.6 mm detectors, slice width of 0.6 mm</p> <p><i>Interpretation:</i> images were evaluated by two experienced radiologists</p> <p><i>Criteria for a positive test:</i> none given</p> <p>Magnetic resonance imaging:</p> <p><i>Timing:</i> performed within 7 days after trauma</p> <p><i>Description of imaging technique (including patient position):</i> 1.5 Tesla with dedicated wrist coil. Prone position with the affected arm above the body. Coronal and axial T1 images (3 to 5 mm slice thickness with a 0.5 mm gap); coronal and axial fat-saturated proton density-weighted images with fat saturation (3 to 5 mm slice thickness with 0.5 to 1.0 mm gap) and coronal T2 images (3 mm slice thickness) were obtained.</p> <p><i>Interpretation:</i> 2 radiologists</p> <p><i>Criteria for a positive test:</i> evidence of a cortical fracture line, a trabecular fracture line, or a combination of these abnormalities</p>	
Follow-up	None reported after final reference standard test was performed	
Notes	Fracture prevalence: 29% (16/55)	

Ilica 2011 (Continued)

Reporting of other fractures: MRI reported 6 other fractures, CT reported 5 other fractures

Table of Methodological Quality

Item	Authors' judgement	Description
Representative spectrum? All tests	Yes	Consecutive series of patients with acute wrist injury, clinical signs for a scaphoid fracture and normal initial radiographs
Acceptable reference standard? All tests	Unclear	Reference standard was one index test: MRI
Acceptable delay between tests? All tests	Yes	Both CT and MRI were performed within 1 week after wrist injury
Partial verification avoided? All tests	Yes	All patients received the intended reference standard
Differential verification avoided? All tests	Yes	All patients received the same reference standard: MRI
Incorporation avoided? All tests	Yes	The index test, CT, was not part of the reference standard
Index test results blinded? All tests	Yes	All images were separately evaluated in a blinded fashion
Reference standard results blinded? All tests	Yes	All images were separately evaluated in a blinded fashion
Relevant clinical information? All tests	No	Clinical information was not present during evaluation of the images. Observers were blinded from all other data
Uninterpretable results reported? All tests	Yes	All outcomes of index tests and reference standard were described, no uninterpretable test results were present
Withdrawals explained? All tests	Yes	Patients who did not return for MRI were reported including explanations
Fracture criteria defined? All tests	Unclear	Fracture criteria given for initial radiographs and MRI. However, these were not given for CT although it was well defined and evaluation was performed by 2 observers

Mallee 2011

Clinical features and settings	Inclusion criteria: presentation to the ED within 24 hours after trauma. Clinically suspected for scaphoid fracture: fall on outstretched hand, tenderness ASB. No fracture on initial radiographs in four views (PA with ulnar deviation, 2x Lat, PA with angulated wrist)
--------------------------------	--

Mallee 2011 (Continued)

Exclusion criteria: age of less than 18 years; any concurrent distal ulnar, radial, or carpal fracture; previous scaphoid fracture; rheumatoid arthritis and cognitive dysfunction that would limit physical examination

Participants	<p>Study location: Amsterdam, The Netherlands</p> <p>Study period: April 2008 to October 2008</p> <p>Participants enrolled: 40: 25 men and 15 women. Mean age 44.3 years (range 20 to 84)</p> <p>Participants included in analyses: 34</p>
Study design	Prospective, consecutive cohort
Target condition and reference standard(s)	<p>Target condition(s): true scaphoid fracture among clinically suspected scaphoid fractures.</p> <p>Reference test: 6-week follow-up radiographs compared with initial radiographs</p>
Index and comparator tests	<p>Index tests: MRI and CT</p> <p>Magnetic resonance imaging</p> <p><i>Timing:</i> performed within 10 days (mean 3.6 days) after trauma</p> <p><i>Description of imaging technique (including patient position):</i> 1.0 Tesla with dedicated wrist coil. slice thickness of 3 mm and a 0.6 mm gap and included the following series: a localizer image, a coronal slice of a short tau inversion recovery (STIR) sequence, and a coronal slice of a spin-echo T1-weighted sequence, in coronal views. The patient was positioned supine, with the forearm and wrist alongside the body. The open MRI allowed for central placement of the hand relative to the magnetic field</p> <p><i>Interpretation:</i> evaluated by a consensus panel: a musculoskeletal radiologist, a trauma surgeon and an orthopaedic surgeon</p> <p><i>Criteria for a positive test:</i> presence of a cortical fracture line, a trabecular fracture line, or a combination of both. Extensive focal zone of oedema without a clear cortical fracture line, comparable with that seen with a stress fracture, was discussed to decide if the findings represented a fracture or not</p> <p>Computed tomography</p> <p><i>Timing:</i> performed within 10 days (mean 3.6 days) after trauma</p> <p><i>Description of imaging technique (including patient position):</i> multidetector 64-slice, high-resolution 0.5 mm slice section thickness. Prone position with the affected arm above the body and palm down. Reconstructions in planes defined by the long axis of the scaphoid were made</p> <p><i>Interpretation:</i> evaluated by a consensus panel: a musculoskeletal radiologist, a trauma surgeon and an orthopaedic surgeon</p> <p><i>Criteria for a positive test:</i> presence of a sharp lucent line within the trabecular bone pattern, a break in the continuity of the cortex, a sharp step in the cortex, or a dislocation of bone fragments</p>
Follow-up	None reported after final reference standard test was performed
Notes	<p>Fracture prevalence: 18% (6/34)</p> <p>Reporting of other fractures: MRI reported 12 other fractures, CT reported 15 other fractures, reference standard reported 4 other fractures.</p>

Table of Methodological Quality

Item	Authors' judgement	Description
------	--------------------	-------------

Mallee 2011 (Continued)

Representative spectrum? All tests	Yes	Consecutive series of patients with acute wrist injury, clinical signs for a scaphoid fracture and normal initial radiographs
Acceptable reference standard? All tests	Yes	Reference standard was adequate according to the requirements mentioned in the protocol: 6-week follow-up radiographs
Acceptable delay between tests? All tests	Yes	The interval between injury and reference test was acceptable: 6 weeks. CT and MRI were performed on the same day within 10 days
Partial verification avoided? All tests	Yes	All patients received the intended reference standard
Differential verification avoided? All tests	Yes	All patients received the same reference standard: 6-week follow-up radiographs
Incorporation avoided? All tests	Yes	Index tests, CT and MRI, were not part of the reference standard
Index test results blinded? All tests	Yes	All images were separately evaluated in a blinded fashion
Reference standard results blinded? All tests	Yes	All images were separately evaluated in a blinded fashion
Relevant clinical information? All tests	Unclear	It was not reported whether clinical information was present or not during evaluation of the images
Uninterpretable results reported? All tests	Yes	All outcomes of index tests and reference standard were described, no uninterpretable test results were present
Withdrawals explained? All tests	Yes	Patients who did not return for 6-week follow-up radiographs were reported including explanations
Fracture criteria defined? All tests	Yes	Fracture criteria for radiographs, CT and MRI were well defined

Memarsadeghi 2006

Clinical features and settings	Inclusion criteria: presentation to the ED within 72 hours. Clinically suspected for scaphoid fracture: pain over the scaphoid. No fracture on initial radiographs in 4 views (PA, Lat, oblique semipronated wrist, radial oblique scaphoid) Exclusion criteria: none reported	
Participants	Study location: Vienna, Austria Study period: June 2000 to July 2002 Participants enrolled: 29: 17 men and 12 women. Mean age 34 years (range 17 to 62)	

Memarsadeghi 2006 (Continued)

Participants included in analyses: 29

Study design	Prospective, consecutive cohort
Target condition and reference standard(s)	Target condition: True scaphoid fracture among clinically suspected scaphoid fractures Reference test: 6-week follow-up radiographs
Index and comparator tests	Index tests: MRI and CT Magnetic resonance imaging <i>Timing:</i> performed on the same day within 6 days (mean 4.1 days) after trauma <i>Description of imaging technique (including patient position):</i> 1.0 Tesla with appropriate receive-only surface coil. Coronal and transverse STIR images (2.4 mm section thickness), coronal and transverse T1 images (2.4 mm section thickness), and coronal three-dimensional T2 images (1.5 mm section thickness) were made. Prone position with affected arm above the body <i>Interpretation:</i> evaluated by a resident and an experienced musculoskeletal radiologist <i>Criteria for a positive test:</i> presence of a cortical fracture line, a trabecular fracture line, or a combination of both Computed tomography <i>Timing:</i> performed on the same day within 6 days (mean 4.1 days) after trauma <i>Description of imaging technique (including patient position):</i> four-detector with detector configuration of two sections at 0.5 mm section thickness. Multiplanar reformations of 0.7 mm thickness were calculated in coronal, sagittal, and transverse planes relative to the wrist. Prone position with affected arm above the body <i>Interpretation:</i> evaluated by a resident and an experienced musculoskeletal radiologist <i>Criteria for a positive test:</i> presence of a sharp lucent line within the trabecular bone pattern, a break in the continuity of the cortex, a sharp step in the cortex, or a dislocation of bone fragments
Follow-up	None reported after final reference standard test was performed
Notes	Fracture prevalence: 38% (11/29) Reporting of other fractures: MRI reported 11 other fractures, CT reported 5 other fractures, reference standard reported 13 other fractures

Table of Methodological Quality

Item	Authors' judgement	Description
Representative spectrum? All tests	Yes	Consecutive series of patients with acute wrist injury, clinical signs for a scaphoid fracture and normal initial radiographs
Acceptable reference standard? All tests	Yes	Reference standard was adequate according to the requirements mentioned in the protocol: 6-week follow-up radiographs
Acceptable delay between tests? All tests	Yes	The interval between injury and reference test was acceptable: 6 weeks. CT and MRI were performed on the same day within 6 days
Partial verification avoided?	Yes	All patients received the intended reference standard

Computed tomography versus magnetic resonance imaging versus bone scintigraphy for clinically suspected scaphoid fractures in patients with negative plain radiographs (Review)

35

Memarsadeghi 2006 (Continued)

All tests

Differential verification avoided? All tests	Yes	All patients received the same reference standard: 6-week follow-up radiographs
Incorporation avoided? All tests	Yes	Index tests, CT and MRI, were not part of the reference standard
Index test results blinded? All tests	Unclear	Not reported if observers were blinded for results of CT and MRI during evaluation of the 6-week follow-up radiographs
Reference standard results blinded? All tests	Yes	Evaluation of CT and MRI was performed without knowledge of the reference test results
Relevant clinical information? All tests	Yes	Site of clinical symptoms were known during evaluation of the CT and MRI
Uninterpretable results reported? All tests	Yes	All outcomes of index tests and reference standard were described; no uninterpretable test results were present
Withdrawals explained? All tests	Yes	All included patients were reported in the results without withdrawals
Fracture criteria defined? All tests	Yes	Fracture criteria for radiographs, CT and MRI were well defined

Nielsen 1983

Clinical features and settings	<p>Inclusion criteria: clinically suspected for scaphoid fracture. No fracture, or inconclusive, on initial radiographs in 4 views (2x unknown, 2x oblique). Timing of presentation to the ED not reported.</p> <p>Exclusion criteria: none reported.</p>
Participants	<p>Study location: Denmark, Randers.</p> <p>Study period: 1980</p> <p>Participants enrolled: 100 (101 wrists): 61 men and 39 women. Mean age 33 years (range 10 to 80)</p> <p>Participants included in analyses: 101 (wrists)</p>
Study design	Prospective, consecutive cohort
Target condition and reference standard(s)	<p>Target condition(s): true scaphoid fracture among clinically suspected scaphoid fractures.</p> <p>Reference test: 2-week and 2-month follow-up radiographs</p>
Index and comparator tests	<p>Index test: BS</p> <p>Bone scintigraphy</p> <p><i>Timing:</i> performed within 10 days after trauma.</p>

Nielsen 1983 (Continued)

Description of imaging technique (including patient position): scanner was equipped with a parallel hole collimator, 3 hours after intravenous administration of 99m-Tc-MDP

Interpretation: not reported who evaluated the images

Criteria for a positive test: "negative": homogeneous and symmetrical tracer uptake in both wrists; "positive": focal uptake radially or centrally in the wrist; "inconclusive": diffuse tracer uptake in the wrist or focal uptake in the ulnar part of the wrist

Follow-up	None reported after final reference standard test was performed
Notes	Fracture prevalence: 11% (11/101) Reporting of other fractures: BS reported 43 other fractures, reference standard reported 9 other fractures

Table of Methodological Quality

Item	Authors' judgement	Description
Representative spectrum? All tests	Unclear	Delay between wrist injury and presentation to the ED was not reported
Acceptable reference standard? All tests	Yes	Reference standard was adequate according to the requirements mentioned in the protocol: both 2-week and 2-month follow-up radiographs
Acceptable delay between tests? All tests	Yes	The interval between injury and reference test was acceptable: 2 months
Partial verification avoided? All tests	Yes	All patients received the intended reference standard
Differential verification avoided? All tests	Yes	All patients received the same reference standard: 2-week and 2-month follow-up radiographs
Incorporation avoided? All tests	Yes	Index test, BS, was not part of the reference standard
Index test results blinded? All tests	Unclear	It was not reported if BS results were present during evaluation of the follow-up radiographs
Reference standard results blinded? All tests	Yes	BS was already evaluated prior to the follow-up radiographs
Relevant clinical information? All tests	No	Clinical and radiographic information was not present during evaluation of BS
Uninterpretable results reported? All tests	Yes	All outcomes of index tests and reference standard were described; inconclusive test results were present and reported
Withdrawals explained? All tests	Yes	All included patients underwent the entire study protocol

Nielsen 1983 (Continued)

Fracture criteria defined? All tests	Yes	Fracture criteria for BS were defined
---	-----	---------------------------------------

O'Carroll 1982

Clinical features and settings	<p>Inclusion criteria: timing of presentation to the ED not reported. Clinically suspected for scaphoid fracture. No fracture on initial radiographs, number of views unknown (AP, Lat, oblique).</p> <p>Exclusion criteria: none reported.</p>
Participants	<p>Study location: Dublin, Ireland</p> <p>Study period: not reported</p> <p>Participants enrolled: 30: 21 men/9 women. Mean age 32 (range 11-72)</p> <p>Participants included in analyses: 30</p>
Study design	Prospective, consecutive cohort
Target condition and reference standard(s)	<p>Target condition(s): true scaphoid fracture among clinically suspected scaphoid fractures.</p> <p>Reference test: 2-week, 4-week and 6-week follow-up radiographs</p>
Index and comparator tests	<p>Index test: BS</p> <p>Bone scintigraphy</p> <p><i>Timing:</i> performed within 10 days (range 3 to 32 days) after trauma.</p> <p><i>Description of imaging technique (including patient position):</i> Tc-99m-HDP and a large field of view Gamma camera. Ventral views of both wrists were obtained 3 hours following intravenous injection of the radiopharmaceutical and 500,000 counts were obtained in each picture</p> <p><i>Interpretation:</i> not reported who evaluated the images</p> <p><i>Criteria for a positive test:</i> none reported</p>
Follow-up	None reported after final reference standard test was performed
Notes	<p>Fracture prevalence: 20% (6/30)</p> <p>Reporting of other fractures: none reported.</p>

Table of Methodological Quality

Item	Authors' judgement	Description
Representative spectrum? All tests	Unclear	Delay between wrist injury and presentation to ED was not reported
Acceptable reference standard? All tests	Yes	Reference standard was adequate according to the requirements mentioned in the protocol: 2-, 4-, 6-week follow-up radiographs
Acceptable delay between tests? All tests	Unclear	BS was performed up to 32 days after wrist injury

O'Carroll 1982 (Continued)

Partial verification avoided? All tests	Yes	All patients received the intended reference standard
Differential verification avoided? All tests	Yes	All patients received the same reference standard: 2-, 4-, 6-week follow-up radiographs
Incorporation avoided? All tests	Yes	Index test, BS, was not part of the reference standard
Index test results blinded? All tests	Unclear	Not reported if results of BS was present during evaluation of radiographs
Reference standard results blinded? All tests	Unclear	Not reported if reference test results were present during evaluation of BS
Relevant clinical information? All tests	Unclear	Not reported if clinical information was present during evaluation of the images
Uninterpretable results reported? All tests	Yes	All outcomes of index tests and reference standard were described; inconclusive test results were not present
Withdrawals explained? All tests	Yes	All included patients received both index test and reference standard; no withdrawals
Fracture criteria defined? All tests	No	Fracture criteria for BS were not well defined

Stordahl 1984

Clinical features and settings	<p>Inclusion criteria: timing of presentation to the ED not reported. Clinically suspected for scaphoid fracture. No fracture, or inconclusive, on initial radiographs in 4 views (PA, Lat, 2x oblique).</p> <p>Exclusion criteria: none reported.</p>
Participants	<p>Study location: Kristiansand, Norway</p> <p>Study period: not reported</p> <p>Participants enrolled: 30: 18 men and 12 women. Mean age 31 years (range 10 to 69)</p> <p>Participants included in analyses: 28 (2 patients were excluded because, upon "revision of the material", their fractures were "already evident on the initial x-rays")</p>
Study design	Not reported, consecutive
Target condition and reference standard(s)	<p>Target condition(s): true scaphoid fracture among clinically suspected scaphoid fractures</p> <p>Reference test: 2-week, 4-week and 6-week follow-up radiographs</p>
Index and comparator tests	<p>Index test: BS</p> <p>Bone scintigraphy</p>

Stordahl 1984 (Continued)

Timing: performed 2 weeks after trauma.

Description of imaging technique (including patient position): radionuclide imaging was performed three hours after the administration of Tc-99m-HDP. A Pho/Gamma 4 Camera with divergent low energy collimator, or pinhole collimator was used

Interpretation: two examiners of unknown expertise evaluated the images

Criteria for a positive test: increased radioactivity, focal or diffuse

Follow-up	None reported after final reference standard test was performed
Notes	Fracture prevalence: 32% (9/28) Reporting of other fractures: BS reported 3 other fractures,

Table of Methodological Quality

Item	Authors' judgement	Description
Representative spectrum? All tests	No	Delay between wrist injury and presentation was not reported and inconclusive radiographs were included
Acceptable reference standard? All tests	Yes	Reference standard was adequate according to the requirements mentioned in the protocol: 2-, 4-, 6-week follow-up radiographs
Acceptable delay between tests? All tests	Yes	The interval between injury and reference test was acceptable: 6 weeks. BS was performed within 2 weeks
Partial verification avoided? All tests	Yes	All patients received the intended reference standard
Differential verification avoided? All tests	Yes	All patients received the same reference standard: 2-, 4-, 6-week follow-up radiographs
Incorporation avoided? All tests	Yes	Index test, BS, was not part of the reference standard
Index test results blinded? All tests	Yes	All images were separately evaluated in a blinded fashion
Reference standard results blinded? All tests	Yes	All images were separately evaluated in a blinded fashion
Relevant clinical information? All tests	Unclear	Not well reported if clinical findings were present during evaluation of radiographs and BS
Uninterpretable results reported? All tests	Yes	All outcomes of index tests and reference standard were described; no uninterpretable test results were present
Withdrawals explained? All tests	Yes	All included patients received both index test and reference standard. 2 patients were excluded since they had fractures on initial radiographs

Stordahl 1984 (Continued)

Fracture criteria defined? All tests	Yes	Fracture criteria for BS were well defined
---	-----	--

Tiel-van Buul 1993

Clinical features and settings	Inclusion criteria: presentation to ED within 24 hours. Clinically suspected for scaphoid fracture: FOOSH, tenderness in ASB. No fracture initial radiographs in four views (PA with ulnar deviation, Lat, 2x oblique). Exclusion criteria: none reported.
--------------------------------	---

Participants	Study location: Amsterdam, The Netherlands Study period: September 1987 to September 1990 Participants enrolled: 160: 82 men and 78 women. Mean age 38.6 years (range 12 to 84) Participants included in analyses: 125
--------------	---

Study design	Not reported, consecutive
--------------	---------------------------

Target condition and reference standard(s)	Target condition(s): true scaphoid fracture among clinically suspected scaphoid fractures. Reference test: 2-week and 6-week follow-up radiographs
--	---

Index and comparator tests	Index test: BS Bone scintigraphy <i>Timing:</i> performed 3 to 34 days (mean 12.3) after injury <i>Description of imaging technique (including patient position):</i> after administration of 200 MBq Tc-99m-HDP, dynamic phase (after 2 to 5 minutes) and static phase (after 2 to 3 hours) were recorded. Anterior and lateral static views were obtained <i>Interpretation:</i> evaluated by an experienced nuclear physician. All radiographs were judged blindly by a panel consisting of a bone radiologist, a general radiologist and a senior traumatologist <i>Criteria for a positive test:</i> presence of focally increased activity in both dynamic and static phase
----------------------------	--

Follow-up	Additional follow-up after a minimum of 1 year
-----------	--

Notes	Fracture prevalence: 17% (21/125) Reporting of other fractures: BS reported 49 other fractures, reference standard reported 24 other fractures
-------	---

Table of Methodological Quality

Item	Authors' judgement	Description
Representative spectrum? All tests	Yes	Consecutive series of patients with acute wrist injury, clinical signs for a scaphoid fracture and normal initial radiographs
Acceptable reference standard? All tests	Yes	Reference standard was adequate according to the requirements mentioned in the protocol: 2- and 6-week follow-up radiographs

Tiel-van Buul 1993 (Continued)

Acceptable delay between tests? All tests	Yes	The interval between injury and reference test was acceptable: 6 weeks. BS was performed within 2 weeks
Partial verification avoided? All tests	Yes	All patients received the intended reference standard
Differential verification avoided? All tests	Yes	All patients received the same reference standard: 2- and 6-week follow-up radiographs
Incorporation avoided? All tests	Yes	Index test, BS, was not part of the reference standard
Index test results blinded? All tests	Yes	All images were separately evaluated in a blinded fashion
Reference standard results blinded? All tests	Yes	BS was performed and evaluated prior to the reference standard
Relevant clinical information? All tests	Unclear	Not reported if clinical information was present during evaluation of BS
Uninterpretable results reported? All tests	Yes	All outcomes of index tests and reference standard were described, no uninterpretable test results were present
Withdrawals explained? All tests	Yes	Withdrawals were reported including explanation
Fracture criteria defined? All tests	Yes	Fracture criteria for BS were well defined

Tiel-van Buul 1996

Clinical features and settings	<p>Inclusion criteria: presentation to ED within 72 hours after trauma. Clinically suspected for scaphoid fracture. No fracture initial radiographs in four views (PA, oblique with ulnar deviation, Lat, PA with tilted beam).</p> <p>Exclusion criteria: claustrophobia</p>
Participants	<p>Study location: Amsterdam, The Netherlands.</p> <p>Study period: not reported</p> <p>Participants enrolled: 16: 11 men and 5 women. Mean age 36 years (range 24 to 60)</p> <p>Participants included in analyses: 16</p>
Study design	Not reported, consecutive
Target condition and reference standard(s)	<p>Target condition(s): true scaphoid fracture among clinically suspected scaphoid fractures</p> <p>Reference test: BS performed in 3 to 14 days (mean 10)</p>

Tiel-van Buul 1996 (Continued)

Index and comparator
tests

Index test: MRI and BS

Magnetic resonance imaging
Timing: performed in 3 to 14 days (mean 10)

Description of imaging technique (including patient position): 1.5 Tesla with a circular surface coil. Prone position with arm extended overhead with wrist fixed in neutral position. T1 and T2 images were obtained with 3.0 mm slice thickness

Interpretation: images were evaluated by an experienced radiologist

Criteria for a positive test: none reported

Bone scintigraphy
Timing: performed in 3 to 14 days (mean 10)

Description of imaging technique (including patient position): after administration of 200 MBq Tc-99m-HDP, dynamic phase and static phase were recorded. Anterior and lateral static views were obtained

Interpretation: all carpal radiographs, MR images and bone scans were reviewed by a panel of the same radiologists and nuclear physician

Criteria for a positive test: presence of focally increased activity in both dynamic and static phase

Follow-up

None reported after final reference standard test was performed

Notes

Fracture prevalence: 44% (7/16)

Reporting of other fractures: BS reported one lunate fracture

Table of Methodological Quality

Item	Authors' judgement	Description
Representative spectrum? All tests	Yes	Consecutive series of patients with acute wrist injury, clinical signs for a scaphoid fracture and normal initial radiographs
Acceptable reference standard? All tests	Unclear	Reference standard was one index test: BS
Acceptable delay between tests? All tests	Yes	The interval between injury and reference test was acceptable: both tests were performed within 14 days
Partial verification avoided? All tests	Yes	All patients received the intended reference standard
Differential verification avoided? All tests	Yes	All patients received the same reference standard: BS
Incorporation avoided? All tests	Yes	Index test, MRI, was not part of the reference standard
Index test results blinded? All tests	Yes	All images were separately evaluated in a blinded fashion

Tiel-van Buul 1996 *(Continued)*

Reference standard results blinded? All tests	Yes	All images were separately evaluated in a blinded fashion
Relevant clinical information? All tests	No	BS was evaluated blinded from other data
Uninterpretable results reported? All tests	Yes	All outcomes of index tests and reference standard were described, no uninterpretable test results were present
Withdrawals explained? All tests	Yes	Withdrawals were reported including explanation
Fracture criteria defined? All tests	Unclear	Fracture criteria for BS were well defined but those for MRI were lacking

ASB = Anatomical snuffbox
PA = Posteroanterior
Lat = Lateral
ED = Emergency department
BS = Bone Scintigraphy
MRI = Magnetic Resonance Imaging
CT = Computed Tomography
FOOSH = Fall on outstretched hand

Characteristics of excluded studies *[ordered by study ID]*

Study	Reason for exclusion
Adey 2007	The series of patients was not consecutive
Akdemir 2004	Patients were only included if they were still clinically suspected to have a scaphoid fracture after 1 to 2 weeks
Asche 1982	Inadequate reference standard
Beeres 2005	No reference standard
Beeres 2007	Inadequate reference standard
Breederveld 2004	Patients were only included if they were still clinically suspected after 1 to 2 weeks
Bretlau 1999	Patients were only included if they were still clinically suspected after 1 to 2 weeks
Brismar 1988	Inadequate reference standard
Bury 2010	Letter to editor
Cook 1997	Inadequate patient selection since patients were not excluded when they had a fracture on initial radiographs
Coupland 1996	Inadequate reference standard

Study	Reason for exclusion
Cruickshank 2007	Inadequate reference standard
Fowler 1998	Patients were only included if they were still clinically suspected after 1 to 2 weeks
Fusetti 2008	Letter to editor
Ganel 1979	No reference standard
Geijer 2011	No index test
Gäbler 2001	Patients were only included if they were still clinically suspected after 1 to 2 weeks
Hobby 2001	Review
Hunter 1997	Inadequate reference standard
Jenkins 2008	No reference standard
Jonsson 1992	Patients were only included if they were still clinically suspected after 7 to 18 days
Keim 2003	No reference standard
Klein 2002	Inadequate patient selection since all patients with wrist pain were included; this study did not focus on the diagnostic test accuracy for clinically suspected scaphoid fractures
Kristensen 1983	Inadequate reference standard
Kumar 2005	No reference standard
Kusano 2002	No reference standard
Lanik 1986	Inadequate reference standard
Lapa 2000	Case report
Lepage 2004	No reference standard
Lindequist 1998	No reference standard
Low 2005	The series of patients was not consecutive
Lozano-Calderon 2006	Inadequate patient selection since a convenience sample was used
Moller 2004	No reference standard
Murphy 1995	Patients were only included if they were still clinically suspected after 1 to 2 weeks
Querellou 2009	Not focused on scaphoid fractures
Raby 2001	No reference standard
Schick 1999	Not focused on scaphoid fractures
Tenneo 2000	Letter to editor

Study	Reason for exclusion
Thorpe 1996	Patients were only included if they were still clinically suspected after 1 to 2 weeks
Tiel-van Buul 1993b	Inadequate reference standard
Tiel-van Buul 1995	Letter to editor
Tiel-van Buul 1998	Letter to editor
Vrettos 1996	Inadequate reference standard
Wilson 1986	Inadequate reference standard
You 2007	No reference standard

DATA

Presented below are all the data for all of the tests entered into the review.

Table Tests. Data tables by test

Test	No. of studies	No. of participants
1 CT	4	277
2 MRI	5	221
3 BS	6	543

Test 1. CT.

Test 2. MRI.

Test 3. BS.

ADDITIONAL TABLES

Table 1. QUADAS checklist and assessment criteria

	Item question	Item answer
1.	Was the spectrum of patients representative of the patients who will re-	Yes: 1) presentation to the emergency department within 72 hours; 2) all included patients were suspected of having a scaphoid fracture

Computed tomography versus magnetic resonance imaging versus bone scintigraphy for clinically suspected scaphoid fractures in patients with negative plain radiographs (Review)

Copyright © 2015 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

Table 1. QUADAS checklist and assessment criteria (Continued)

	ceive the test in practice? (representative spectrum)	with normal radiographs; 3) prospective study design; and 4) consecutive series Unclear: if insufficient information is presented on study design or inclusion criteria No: 1) patients presented after 72 hours; 2) retrospective study design; or 3) not a consecutive series of patients
2.	Is the reference standard likely to classify the target condition correctly? (acceptable reference standard)	Yes: 1) if reference standard is 6-week follow-up radiographs (this is the most commonly used reference standard); 2) if 2 index tests report the same outcome; or 3) if 1 index test is used as a reference standard combined with clinical evaluation Unclear: suboptimal would be if only 1 index test is used No: 1) if only clinical evaluation after 6 weeks is considered to be the reference standard; or 2) if only clinical evaluation or radiographs, or both, after 2 weeks is considered to be the reference standard; 3) if insufficient information is given
3.	Is the time period between reference standard and index test short enough to be reasonably sure that the target condition did not change between the 2 tests? (acceptable delay between tests)	Yes: if average interval between trauma and follow-up radiographs was 6 to 14 weeks. We will allow follow-up radiographs taken at least 2 weeks after trauma although this is considered to be a suboptimal reference standard No: if interval was not clearly reported or before 2 weeks or greater than 14 weeks after trauma
4.	Did the whole sample or a random selection of the sample receive verification using the intended reference standard? (partial verification avoided)	Yes: if all patients received both index test and reference standard. We will allow for a random selection Unclear: if insufficient information was available to judge this No: if some of the patients who received the index test did not receive verification of their true disease state, and the selection of patients to receive the reference standard was not random
5.	Did patients receive the same reference standard irrespective of the index test result? (differential verification avoided)	Yes: if all patients received the same reference standard, irrespective of the index test result Unclear: if it is unclear whether different reference standards were used No: if the outcome of the index test influenced the choice of reference standard
6.	Was the reference standard independent of the index test (i.e. the index test did not form part of the reference standard)? (incorporation avoided)	Yes: if index test was not part of the reference standard Unclear: unclear No: if index test was part of the reference standard
7.	Were the reference standard results interpreted without knowledge of the results of the index test? (index test results blinded)	Yes: if the evaluation was blinded from the index test results Unclear: if insufficient information was given on the blinded evaluation of the reference standard No: if the index test results were present during evaluation of the reference standard
8.	Were the index test results interpreted without knowledge of the results	Yes: if the evaluation of the index test results was blinded from the results of the reference standard

Table 1. QUADAS checklist and assessment criteria (Continued)

	of the reference standard? (reference standard results blinded)	Unclear: if insufficient information was given on the blinded evaluation of the index test No: if the results of the reference standard were present during evaluation of the index test
9.	Were the same clinical data available when test results were interpreted as would be available when the test is used in practice? (relevant clinical information)	Yes: if available clinical data during evaluation of the test are the same as in daily practice Unclear: if insufficient information is given on the available clinical data during evaluation of the test No: if the usual clinical data were not available during evaluation of the test
10.	Were uninterpretable/intermediate test results reported? (uninterpretable results reported?)	Yes: if the number of uninterpretable/intermediate test results is stated or if results match the number of initially included patients Unclear: if insufficient information to permit judgement No: if uninterpretable/intermediate test results are reported, without amount, or were excluded
11.	Were withdrawals from the study explained? (withdrawals explained)	Yes: if any withdrawals are stated and explained Unclear: if insufficient information to permit judgement No: if withdrawals are not mentioned or explained
12.	Did the study provide a clear definition of what was considered to be a 'positive' result?	Yes: if fracture criteria are well defined, even though they can differ between studies Unclear: if insufficient information but evaluation was performed by at least 2 observers No: if no fracture criteria are defined

Table 2. -2 Log Likelihood of models in each pairwise comparison

	-2 Log Likelihood		
	Model (a)	Model (b)	Model (c)
CT vs MRI	48.0	48.2	50.1
CT vs BS	63.8	65.0	115.3
MRI vs BS	72.3	72.6	102.3

Model (a) assumed different shape (beta), accuracy (alpha) and threshold effect (theta)

Model (b) assumed different shape (beta) and accuracy (alpha)

Model (c) assumed different shape (beta) only

Table 3. Direct comparisons (CT versus MRI; CT versus BS; MRI versus BS) for detection of scaphoid fractures

Study	Cases	Non-cases	Sensitivity (95% CI)	Specificity (95% CI)	Sensitivity (95% CI)	Specificity (95% CI)	Difference in sensitivity (95% CI) P value	Difference in specificity (95% CI) P value
CT			MRI					
Mallee 2011	6	28	0.67 (0.22 to 0.96)	0.96 (0.82 to 1.00)	0.67 (0.22 to 0.96)	0.89 (0.72 to 0.98)	0.00 (-0.53 to 0.53) P = 1.0	0.07 (-0.06 to 0.21) P = 0.611
Memarsadegh 2006	11	11	0.73 (0.39 to 0.94)	1.00 (0.81 to 1.00)	1.00 (0.72 to 1.00)	1.00 (0.81 to 1.00)	-0.27 (-0.54 to -0.01) P = 0.214	0.00 (Standard error is zero; P undefined)
CT			BS					
De Zwart 2012	20	139	0.70 (0.46 to 0.88)	0.99 (0.96 to 1.00)	0.95 (0.75 to 1.00)	0.94 (0.88 to 0.97)	-0.25 (-0.47 to -0.03) P = 0.092	0.05 (0.01 to 0.10) P = 0.019
MRI			BS					
Beeres 2008	20	80	0.80 (0.56 to 0.94)	1.00 (0.95 to 1.00)	1.00 (0.83 to 1.00)	0.90 (0.81 to 0.96)	-0.20 (-0.38 to -0.03) P = 0.106	0.10 (0.03 to 0.17) P = 0.007

Sensitivity, specificity and their CIs are recalculated with Review Manager
P values were based on the 2-sided Fisher's exact test

APPENDICES

Appendix 1. Diagnostic test performance characteristics

Sensitivity; specificity; accuracy

- Sensitivity: the proportion of patients who had a scaphoid fracture according to the reference standard and who were classified as having a positive test.
- Specificity: the proportion of patients who had no scaphoid fracture according to the reference standard and who were classified as having a negative test.
- Accuracy: the proportion of patients who were correctly classified by the test.

Positive predictive value (PPV); negative predictive value (NPV)

- PPV: the probability that a patient with a positive test has a scaphoid fracture.
- NPV: the probability that a patient with a negative test has no scaphoid fracture.

Appendix 2. Search strategies

MEDLINE (Ovid Web)

```

1 exp Magnetic Resonance Imaging/ (269663)
2 ((magnetic resonance or MR or NMR) adj2 (imag* or tomograph* or scan*)).tw. (149610)
3 (MRI or MRIs or NMRI).tw. (112187)
4 (diffusion weighted imag* or DWI or T2-weighted imag*).tw. (12506)
5 or/1-4 (321609)
6 exp Tomography, X-Ray Computed/ (267102)
7 (comput* adj3 tomograph*).tw. (154076)
8 (CT or CAT).tw. (246660)
9 micro-computed tomog*.tw. (1454)
10 or/6-9 (453788)
11 Radionuclide Imaging/ (24045)
12 (scintigra* or radioscintigra*).tw. (39873)
13 (bone adj3 scan).tw. (4431)
14 scintiscan*.tw. (1491)
15 or/11-14 (63106)
16 or/5,10,15 (746644)
17 Scaphoid bone/ or Wrist injuries/ or Wrist Joint/ (11542)
18 exp Fractures, Bone/ (129394)
19 and/17-18 (3618)
20 ((Scaphoid* or wrist or navicular) adj3 (fracture* or injur* or trauma)).tw. (2892)
21 or/19-20 (5363)
22 and/16,21 (761)
23 Diagnostic Imaging/ or Diagnosis, Differential/ or exp "Sensitivity and Specificity"/ or "Predictive Value of Tests"/ (723481)
24 scaphoid.mp. (3223)
25 and/23-24 (231)
26 Fractures, Bone/di, ra, ri [Diagnosis, Radiography, Radionuclide Imaging] (10875)
27 and/24,26 (586)
28 or/22,25,27 (1226)

```

EMBASE (Ovid Web)

```

1 exp Nuclear Magnetic Resonance Imaging/ (434436)
2 ((magnetic resonance or MR or NMR) adj2 (imag* or tomograph* or scan*)).tw. (188295)
3 (MRI or MRIs or NMRI).tw. (166239)
4 (diffusion weighted imag* or DWI or T2-weighted imag*).tw. (17259)
5 or/1-4 (476776)
6 exp Computer Assisted Tomography/ (512291)
7 (comput* adj3 tomograph*).tw. (193911)
8 (CT or CAT).tw. (336635)
9 micro-computed tomog*.tw. (1660)
10 or/6-9 (702231)
11 Bone Scintiscanning/ or Radiodiagnosis/ (50657)
12 (scintigra* or radioscintigra*).tw. (54696)

```

Computed tomography versus magnetic resonance imaging versus bone scintigraphy for clinically suspected scaphoid fractures in patients with negative plain radiographs (Review)

Copyright © 2015 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

13 (bone adj3 scan).tw. (6365)
14 scintiscan*.tw. (2105)
15 or/11-14 (103388)
16 or/5,10,15 (1102315)
17 Scaphoid Fracture/ (1339)
18 Scaphoid Bone/ or Wrist Injury/ or Wrist/ or Wrist Radiography/ (20278)
19 exp Fracture/ (183932)
20 and/18-19 (3626)
21 ((scaphoid* or wrist or navicular) adj3 (fracture* or injur* or trauma)).tw. (3638)
22 or/17,20-21 (6434)
23 and/16,22 (1208)
24 "Sensitivity and Specificity"/ or Diagnostic Imaging/ or Receiver Operating Characteristic/ or Diagnostic Accuracy/ or Diagnostic Test/ or Diagnostic Value/ or Diagnostic Procedure/ (557240)
25 scaphoid.mp. (4476)
26 and/24-25 (382)
27 Fracture/di [Diagnosis] (3601)
28 and/25,27 (112)
29 Scaphoid Fracture/di [Diagnosis] (497)
30 or/23,26,28-29 (1586)

Cochrane Database of Abstracts of Reviews of Effects, Cochrane Central Register of Controlled Trials, NHS Economic Evaluation Database (Wiley Online Library)

#1 MeSH descriptor Magnetic Resonance Imaging explode all trees 4459
#2 ((magnetic resonance or MR or NMR) NEAR/2 (imag* or tomograph* or scan*)):ti,ab,kw 5453
#3 (MRI or MRIs or NMRI):ti,ab,kw 2723
#4 (diffusion weighted imag* or DWI or T2-weighted imag*):ti,ab,kw 404
#5 (#1 OR #2 OR #3 OR #4) 6382
#6 MeSH descriptor Tomography, X-Ray Computed explode all trees 3159
#7 (comput* NEAR/3 tomograph*):ti,ab,kw 6556
#8 (CT or CAT):ti,ab,kw 19978
#9 (micro-computed tomog*):ti,ab,kw 12
#10 (#6 OR #7 OR #8 OR #9) 24123
#11 MeSH descriptor Radionuclide Imaging, this term only 213
#12 (scintigra* or radioscintigra*):ti,ab,kw 1550
#13 (bone NEAR/3 scan):ti,ab,kw 177
#14 (scintiscan*):ti,ab,kw 201
#15 (#11 OR #12 OR #13 OR #14) 1933
#16 (#5 OR #10 OR #15) 31013
#17 MeSH descriptor Scaphoid Bone, this term only 41
#18 MeSH descriptor Wrist Injuries, this term only 103
#19 MeSH descriptor Wrist Joint, this term only 164
#20 (#17 OR #18 OR #19) 284
#21 MeSH descriptor Fractures, Bone explode all trees 3390
#22 (#20 AND #21) 121
#23 (Scaphoid* or wrist or navicular) NEAR/3 (fracture* or injur* or trauma):ti,ab,kw 226
#24 (#22 OR #23) 243
#25 (#16 AND #24) 31
#26 MeSH descriptor Diagnostic Imaging, this term only 187
#27 MeSH descriptor Diagnosis, Differential, this term only 1330
#28 MeSH descriptor Sensitivity and Specificity explode all trees 13581
#29 MeSH descriptor Predictive Value of Tests explode all trees 5043
#30 (#26 OR #27 OR #28 OR #29) 14595
#31 (scaphoid) 100
#32 (#30 AND #31) 7
#33 MeSH descriptor Fractures, Bone, this term only with qualifiers: DI,RA,RI 116
#34 (#31 AND #33) 23
#35 (#25 OR #32 OR #34) 34
#36 (diagnos* and scaphoid and fractur*):ti,ab,kw in Trials 17
#37 (#35 OR #36) 2 (DARE), 34 (CENTRAL), 8 (NHS EED)

Other databases

We searched the following databases for the term "scaphoid":

MEDION (0 records), ARIF (3), WHO International Clinical Trials Registry Platform (13), and Current Controlled Trials (14)

CONTRIBUTIONS OF AUTHORS

All authors contributed to the development of the review and commented on and approved the final version. The guarantor of this review is Wouter Mallee.

DECLARATIONS OF INTEREST

Wouter H Mallee: was involved in the publication of a potentially eligible study. Therefore, this trial was assessed by other review authors.
Job N Doornberg: was involved in the publication of a potentially eligible study. Therefore, this trial was assessed by other review authors.
Rudolf W Poolman: Dr Poolman's institution was supported for several research activities by several implant and drug companies, none of which are related to this review (for example, Link, AMGEN, AstraTech).

Peter Kloen: none known

Mario Maas: has provided consultancy to several companies (Genzyme, Shire, Protalix, Actelion) in an unrelated area of "Gaucher Disease and Imaging".

Henrica CW de Vet: none known

J Wang: none known

SOURCES OF SUPPORT

Internal sources

- Academic Medical Center of Amsterdam, Netherlands.
- Onze Lieve Vrouwe Gasthuis Amsterdam, Netherlands.
- Vrije Universiteit Medical Center of Amsterdam, Netherlands.
- Massachusetts General Hospital, USA.

External sources

- No sources of support supplied

DIFFERENCES BETWEEN PROTOCOL AND REVIEW

To improve clarity, minor edits were made to Background and Methods.

In [Assessment of methodological quality](#), we added in a paragraph describing the key rules for rating overall methodological quality.

In [Data extraction and management](#), we clarified on our usage of overlapping data where studies had been published more than once. We clarified that in cases of overlapping patient data, we only used the data once.

To compare CT, MRI and BS, we used the HSROC model instead of the bivariate model. When we tried to use the bivariate model, the parameter estimates did not converge in SAS PROC NLMIXED. This might be caused by the small number of studies.

INDEX TERMS

Medical Subject Headings (MeSH)

*Magnetic Resonance Imaging; *Radionuclide Imaging; *Tomography, X-Ray Computed; Fractures, Bone [*diagnosis] [diagnostic imaging]; Prospective Studies; Retrospective Studies; Scaphoid Bone [diagnostic imaging] [*injuries]; Sensitivity and Specificity

MeSH check words

Humans