Published in final edited form as:

Org Lett. 2018 December 07; 20(23): 7400–7404. doi:10.1021/acs.orglett.8b03035.

Electron-Mediated Aminyl and Iminyl Radicals from C5 Azido-Modified Pyrimidine Nucleosides Augment Radiation Damage to Cancer Cells

Zhiwei Wen[†], Jufang Peng[†], Paloma R. Tuttle[†], Yaou Ren[†], Carol Garcia[†], Dipra Debnath[‡], Sunny Rishi[‡], Cameron Hanson[‡], Samuel Ward[‡], Anil Kumar[‡], Yanfeng Liu[§], Weixi Zhao[§], Peter M. Glazer[§], Yuan Liu[†], Michael D. Sevilla^{*‡}, Amitava Adhikary^{*‡}, and Stanislaw F. Wnuk^{*‡}

[†]Department of Chemistry and Biochemistry, Florida International University, Miami, Florida 33199, United States

[‡]Department of Chemistry, Oakland University, Rochester, Michigan 48309, United States

[§]Department of Therapeutic Radiology, Yale University School of Medicine, New Haven, Connecticut 06520, United States

Abstract

Two classes of azido-modified pyrimidine nucleosides were synthesized as potential radiosensitizers; one class is 5-azidomethyl-2'-deoxyuridine (AmdU) and cytidine (AmdC), while the second class is 5-(1-azidovinyl)-2'-deoxyuridine (AvdU) and cytidine (AvdC). The addition of radiation-produced electrons to C5-azido nucleosides leads to the formation of π -aminyl radicals followed by facile conversion to σ -iminy radicals either via a bimolecular reaction involving intermediate α -azidoalkyl radicals in AmdU/AmdC or by tautomerization in AvdU/AvdC. AmdU demonstrates effective radiosensitization in EMT6 tumor cells.

Graphical Abstract

*Corresponding Authors sevilla@oakland.edu., adhikary@oakland.edu., wnuk@fiu.edu.

ASSOCIATED CONTENT

* Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: [10.1021/acs.orglett.8b03035](https://doi.org/10.1021/acs.orglett.8b03035).

Experimental procedures, mechanistic and theoretical studies, and compound characterization data for all new compounds (PDF)

Notes

The authors declare no competing financial interest.

Pyrimidine nucleosides modified at C5 (e.g., 5-bromo-2'-deoxyuridine) are well-investigated as radiosensitizers in cancer therapy.¹ 5-(Phenylselenenyl)methyl-2'-deoxyuridine and 5-thiocyanato-2'-deoxyuridine were shown to induce reactive benzyl-type² or uracil-C5-thiyl radical³ for DNA interstrand cross-linking^{2,3} and apoptosis in cancer cells.⁴

3'-Azido-3'-deoxythymidine (AZT) has shown a significant radiosensitization in irradiated human colon cancer, larynx squamous carcinoma, and malignant glioma cells.⁵ Moreover, AZT has been employed as a radiation sensitizer in radiotherapy of tumors for HIV-positive patients.⁶ We found that the radiation-produced prehydrated electrons⁷ leads to the site-specific formation of a localized π -aminyl radical ($\text{RNH}^\bullet$) in azido-substituted nucleosides⁸ and sugars.^{9,10} Our ESR studies show that $\text{RNH}^\bullet$ formed from AZT undergoes bimolecular H atom abstraction from a C5-methyl group, generating a thymine allyl radical ($\text{dUCH}_2^\bullet$), or from a sugar moiety yielding $\text{C5}'^\bullet$.⁸ Moreover, $\text{RNH}^\bullet$ and modified base radicals can produce sugar radicals which lead to DNA-strand breaks^{1a} and eventually lead to induction of apoptosis.^{1c} We hypothesize that incorporation of azido-modified nucleosides into genomic DNA would augment radiation-induced damage in cells owing to the reactions of aminyl radicals under hypoxic conditions and, therefore, act as potential radiosensitizers.

Owing to the poor incorporation of AZT in genomic DNA,¹¹ in this work, we have employed 5-azidomethyl-2'-deoxyuridine (AmdU, 4), 5-azidomethyl-2'-deoxycytidine (AmdC, 6), 5-(1-azidovinyl)-2'-deoxyuridine (AvdU, 12), and 5-(1-azidovinyl)-2'-deoxycytidine (AvdC, 13) analogues^{12,13} (Schemes 1 and 2) to study the radiation-mediated formation of $\text{RNH}^\bullet$ and its subsequent reactions. Recently, AmdU has been metabolically incorporated into DNA in living cells for click labeling of DNA.¹⁴ Moreover, its 5'-triphosphate was found to be the substrate for DNA polymerases and PCR amplification.^{15,16} Herein, we report that π -aminyl $\text{RNH}^\bullet$ generated by radiation-produced electron addition to azido-nucleosides undergoes conversion to the thermodynamically more stable σ -iminyl radicals ($\text{R N}^\bullet$). We also show incorporation of AmdU and AmdC phosphates into DNA fragments by polymerase-catalyzed reactions and present evidence that these azidonucleosides can act as radio-sensitizers.

AmdU 4 was synthesized from thymidine 1 by sequential silylation, bromination with NBS, displacement of bromide with NaN_3 , and desilylation (45% overall, Scheme 1).^{14,17} Treatment of 3 with 2,4,6-triisopropylbenzenesulfonyl chloride (TIPBSCI) in the presence of

TEA/DMAP followed by displacement of the resulting aryl sulfate with NH_4OH and deprotection of 5 with TBAF provided AmdC 6 (57% overall). AmdU and AmdC were triphosphorylated¹⁸ to give AmdUTP 7 (76%) and AmdCTP 8 (23%).

Silver-catalyzed hydroazidation¹⁹ of the protected 5-ethynyl-2'-deoxyuridine²⁰ 9 with TMSN_3 regioselectively produced α -azidovinyl 11 (52%; Scheme 2).²¹ Deacetylation of 11 gave AvdU¹³ 12 (90%). Analogous hydroazidation of the protected 5-ethynyl-2'-deoxycytidine²² 10 followed by deacetylation provided AvdC 13 (42% overall). Conversion of uracil base in 11 to a cytosine counterpart by treatment²³ of 11 with TIPBSCl followed by ammonolysis also gave 13 (Supporting Information (SI)).

Radiation-produced electron addition to 4 allowed us to identify three radicals: (i) π -aminyl radical, $\pi\text{-U-5-CH}_2\text{-ND}^\bullet$; (ii) α -azidoalkyl radical, $\text{U-5-CH}^\bullet\text{-N}$; and (iii) σ -iminyl radical, $\text{U-5-CH N}^\bullet$ (Figures 1 and S1, Scheme 3). The center of the spectrum in Figure 1A (blue) does not show the doublet²⁴ expected for $\text{U}^{\bullet-}$, thereby indicating that the azide moiety of 4 scavenged nearly all radiation-produced electrons to form the π -aminyl $\text{RNH}^\bullet$ via a dissociative electron attachment. The initial electron attachment is followed by rapid protonation of the incipient nitrene anion radical (Scheme 3). The spectrum extends over 178.5 G, and its hyperfine structure shows line components from the hyperfine coupling (HFCC) due to single axially symmetric anisotropic nitrogen [$A_{\parallel} = \sim 42.5$ G, $A_{\perp} = \sim 0$] with $g_{\parallel}$ and $g_{\perp}$ values that are typical of aminyl radical nitrogen (Table S1). In $\text{U-5-CH}_2\text{-ND}^\bullet$, the radical-site p-orbital strongly couples with two $\beta\text{-CH}_2^-$ protons, generating the wide doublet (ca. 93.5 G). The simulated spectrum of $\text{U-5-CH}_2\text{-ND}^\bullet$ (Figure 1A, red; see the SI for simulation parameters) matches the line components of the experimental blue spectrum in Figure 1A.

On subsequent stepwise annealing at 140, 160, 165, and 170 K, new line components appear which are assigned to two new species (Figure 1B–E, blue). The first species initially observed in the center of spectrum 1C and resolved as a multiplet in spectrum 1D is assigned to a C-centered α -azidoalkyl radical ($\text{U-5-CH}^\bullet\text{-N}_3$) formed via H-abstraction by $\text{RNH}^\bullet$ from a proximate parent 4 (Scheme 3). In the spectrum in Figure 1C, the second species begins to form. The total hyperfine splitting decreases by ca. 8 G from that found in Figure 1A,B (i.e., the wings line components move in), and this new species becomes the only radical found in Figure 1E. The spectrum in Figure 1E is assigned to the σ -iminyl radical, $\text{U-5-CH N}^\bullet$ resulting from couplings of one anisotropic nitrogen and one β -proton. For assignments of radicals contributing to the spectra in Figure 1D,E and HFCC, see the SI (pages S6 and S7, Table S1, and Figures S1 and S2). These results show that the reactive $\pi\text{-U-5-CH}_2\text{-ND}^\bullet$ abstracts an H-atom from the parent 4 to form the azidoalkyl radical intermediate, $\text{U-5-CH}^\bullet\text{-N}_3$. Subsequently, $\text{U-5-CH}^\bullet\text{-N}_3$ promptly undergoes a unimolecular $\beta\text{-N}_2$ elimination from the azide group to produce the thermodynamically more stable $\sigma\text{-U-5-CH N}^\bullet$ (Scheme 3).

The $\pi\text{-C-5-CH-ND}^\bullet$ formation and its subsequent conversion to the σ -iminyl radical, $\text{C-5-CH N}^\bullet$, has also been observed upon radiation-produced electron attachment to cytosine counterpart AmdC 6 (Scheme S1, Figure S2, and Table S1). The α -azidoalkyl radicals generated from photolysis of α -azidoacetophenones²⁵ or γ -azidobutyrophenone²⁶ as well as

α -azido o-iodoanilides under tin-mediated radical reactions²⁷ are known to undergo similar conversion to the σ -iminy radical.

A radiation-produced electron addition to the vinyl azide **12** also generates π -RNH $\cdot$, which undergoes facile tautomerization to thermodynamically more stable σ -iminy radical (Figure 2 and Scheme 4). The 77 K ESR spectrum (black, Figure 2A) shows line components from single axially symmetric anisotropic nitrogen HFCCs and from two anisotropic protons of the CH₂ group in **12** (Table S1). We assign these to the π -aminyl radical, U-5-C(ND $\cdot$) CH₂ (Scheme 4). Formation of U-5-C(ND $\cdot$) CH₂ results from a dissociative electron attachment pathway similar to that shown in Scheme 3. The simulated red spectrum (see Table S1 for simulation parameters) matches line components of the black spectrum and supports our assignment (Figure 2A). HFCC values for the anisotropic nitrogen and CH₂ α -protons in U-5-C(ND $\cdot$) CH₂ and its optimized geometry are reported in Table S1 and Figure S5.

ESR spectra obtained upon progressive annealing (Figure 2B–D, black) show that height of the singlet at the center increases with concomitant decrease of the line components from the two anisotropic α CH₂ protons. The black spectrum (Figure 2D) is due solely to an axially symmetric anisotropic nitrogen (for HFCCs see Table S1) and is assigned to the σ -iminy radical, U-5-C(CH₃) N $\cdot$. The simulated spectrum (Figure 2D, red; Table S1) matches the line components of the black spectrum.²⁸ Nearly identical spectra were obtained from one-electron attachment to AvdC, proving that formation of the π -aminyl radical (C-5-C(ND $\cdot$) CH₂) and its facile tautomerization to the σ -iminy radical (C-5-C(CH₃) N $\cdot$) has a general character and occurs independent of the nucleobase.

Increasing the concentration (1 to 5 mg/mL) of **12** has no effect on the extent of this conversion as observed from spectra recorded under the same microwave power, modulation, and gain. From these results, we conclude that the conversion of π -RNH $\cdot$ to σ -R N $\cdot$ observed in **12** and **13** is by tautomerization; it occurs via facile intramolecular proton transfer from the aminyl group to the double-bonded CH₂ group in the π -RNH $\cdot$. Conversion of π -RNH $\cdot$ to σ -R N $\cdot$ has also been observed in one-electron oxidized 1-methylcytosine and its derivatives.²⁸

In contrast to our previous results in which π -RNH $\cdot$ produced from one-electron reduction of AZT and azidopen-toses^{8,9} undergoes H atom abstraction reactions, we find that π -RNH $\cdot$ from **4**, **6**, **12**, and **13** undergoes conversion to σ -R N $\cdot$. For **4** and **6**, the π -RNH $\cdot$ to σ -R N $\cdot$ conversion is bimolecular which, at first, involves H-abstraction to form an α -azidoalkyl radical followed by a unimolecular decomposition to σ R N $\cdot$ (Scheme 3). However, **12** and **13** undergo tautomerization of π -RNH $\cdot$ to σ -R N $\cdot$ (Scheme 4). Owing to the high free-radical scavenger concentrations in cells,^{1a} the bimolecular conversion of π -RNH $\cdot$ to σ -R N $\cdot$ from **4** and **6** is unlikely to take place. However, the tautomerization of π -RNH $\cdot$ to σ -R N $\cdot$ from **12** and **13** should occur even in cells. Further, reactivity of σ -R N $\cdot$ from **12** and **13** is far less than that of a π -RNH $\cdot$. Therefore, it is expected that the π -RNH $\cdot$ from **4** and **6** could augment radiation damage more effectively than the σ -R N $\cdot$ from **12** and **13**.

Incorporation of nucleosides into DNA is important for them to exhibit radiosensitization.^{1b–d,29} Thus, AmdUTP was successfully incorporated by the *E. coli* Klenow fragment of DNA polymerase I (pol I) and human repair DNA polymerase β (pol β) during DNA leading and lagging strand synthesis and BER (base excision repair) using an open template, onenucleotide gap substrates, and one-nucleotide substrate containing a 5'-THF (a tetrahydrofuran ring which mimics a sugar residue; Figure S6, Table S2; see the SI for more details). Incorporation of AmdUTP in the presence of dATP, dGTP, and dCTP showed that both polymerases readily inserted AmdUTP into dsDNA and extended the nucleotide during DNA replication and BER (Figure S7). AmdU incorporation can be ligated into duplex DNA during DNA replication and BER in the presence of LIG I (Figure S8). These results are consistent with a recent finding that AmdU was efficiently incorporated into newly synthesized DNA in human cancer cells.¹⁴ The AmdCTP 8 was also incorporated into DNA by pol β during DNA replication and BER (Table S2, template 2; Figure S9).

To test our hypothesis that azidonucleosides incorporated into DNA can act as radiosensitizers, we investigated the radiation response of EMT6 breast cancer cells to the presence of 100 μ M azido-modified nucleosides in both aerobic and hypoxic environments in terms of survival fraction (SF). For survival fraction measurements in aerobic cells, the cultures were treated with a 100 μ M azido compound or vehicle for 48 h. For the measurements in hypoxic cells, hypoxic conditions were applied for 4 h after a 44 h aerobic incubation with 100 μ M azido compounds or vehicles. To investigate the radiosensitizing effect, cells were irradiated with 7.5 Gy X-ray during the final few minutes of the 48 h incubation (Figure 3 and Table S3).

AmdU showed radiosensitization under both normoxic and hypoxic environments with sensitization enhancement ratios (SER) at 7.5 Gy X-ray ($SER_{SF7.5}$) of 4.57 and 4.10, respectively (Figure 3, Table S3). These results show that aminyl radicals generated in AmdU augments radiation damage to cells. In a hypoxic microenvironment, $RNH^{\bullet}$ formed from electron addition to AmdU can be involved in the H atom abstraction reactions^{1a,8,9} leading to lesions that can induce apoptosis of cancer cells.^{1c} On the other hand, in the aerobic cells, the aminyl radical generated from AmdU can react with oxygen to generate aminylperoxyl radical $RNHOO^{\bullet}$ and eventually lead to aminoxyl (nitroxyl) radicals $RNO^{\bullet}$,³⁰ which also can lead to DNA damage.³¹ Other azido nucleosides showed a lower radiosensitizing effect with $SER_{SF7.5}$ index of 1.35 for AmdC under hypoxic cells and 1.37 for AvdC under aerobic cells.

Reasons for the differences in radiosensitization between AmdU 4 and AmdC 6 are unclear. Possibilities include differences in drug uptake into cells, metabolic phosphorylation,³² and/or reactivity of the aminyl radical generated at the uracil and cytosine base. The higher radiosensitizing effect of AmdU compared with that of AvdU 12 and AvdC 13 could be explained by the higher reactivity of π - $RNH^{\bullet}$ from AmdU compared to that of σ -R $N^{\bullet}$ from AvdU and AvdC.

In conclusion, the pyrimidine nucleosides modified at the C5 position with azidomethyl or azidovinyl moieties have been designed as potential radiosensitizers. The 5'-phosphates of 5-azidomethyl analogues were incorporated into DNA fragments by polymerase-catalyzed

reactions. The π -aminy radicals generated from 4, 6, 12, and 13 undergo facile conversion to more stable σ -iminy radicals, either bimolecularly involving an α -azidoalkyl radical intermediate or by tautomerization. AmdU may act as effective radiosensitizer in EMT6 cancer cells in the presence or absence of oxygen. Since AmdU 4 has been used for DNA labeling in cells,^{14,33} 4 can serve a dual purpose of labeling tumor cells prior to, during, or after radiotherapy and may radiosensitize the tumor during radiotherapy.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENTS

The authors are thankful for financial support from the NIH (M.D.S., R01 CA045424; Y.L., ES023569; S.F.W., SC1CA138176). Z.W. is a recipient of the FIU Dissertation Year Fellowship. J.P. is grateful for support from the Oversea Scholarship Program of Shaanxi Normal University.

REFERENCES

- (1). (a) von Sonntag C Free-Radical-Induced DNA Damage and Its Repair; Springer-Verlag: Berlin, 2006. (b) Rak J; Chomicz L; Wicz J; Westphal K; Zdrowowicz M; Wityk P; Zyndul M; Makurat S; Golon L J. Phys. Chem. B 2015, 119, 8227. [PubMed: 26061614] (c) Greenberg MM Radiat. Phys. Chem 2016, 128, 82. (d) Schurmann R; Vogel S; Ebel K; Bald I Chem. - Eur. J 2018, 24, 1.
- (2). Hong IS; Ding H; Greenberg MM J. Am. Chem. Soc 2006, 128, 485. [PubMed: 16402835]
- (3). Zdrowowicz M; Chomicz L; Zyndul M; Wityk P; Rak J; Wiegand TJ; Hanson CG; Adhikary A; Sevilla MD Phys. Chem. Chem. Phys 2015, 17, 16907. [PubMed: 26059609]
- (4). Kim BM; Rode AB; Han EJ; Hong IS; Hong SH Apoptosis 2012, 17, 200. [PubMed: 22002103]
- (5). (a) Coucke PA; Cottin E; Decosterd LA Acta Oncol. 2007, 46, 612. [PubMed: 17562437] (b) Zhou F-X; Liao Z-K; Dai J; Xiong J; Xie C-H; Luo Z-G; Liu S-Q; Zhou Y-F Biochem. Biophys. Res. Commun 2007, 354, 351. [PubMed: 17223082] (c) Liao Z-KZ; F, X.; Luo Z-G; Zhang W-J; Jie X; Bao J; Han G; Zhang M-S; Xie C-H; Zhou Y-F Oncol. Rep 2008, 19, 281. [PubMed: 18097608]
- (6). Housri N; Yarchoan R; Kaushal A Cancer 2010, 116, 273. [PubMed: 20014399]
- (7). Ma J; Wang F; Denisov SA; Adhikary A; Mostafavi M Sci. Adv 2017, 3, e1701669 [PubMed: 29250599]
- (8). Adhikary A; Khanduri D; Pottiboyina V; Rice CT; Sevilla MD J. Phys. Chem. B 2010, 114, 9289. [PubMed: 20575557]
- (9). Mudgal M; Rishi S; Lumpuy DA; Curran KA; Verley KL; Sobczak AJ; Dang TP; Sulimoff N; Kumar A; Sevilla MD; Wnuk SF; Adhikary AJ Phys. Chem. B 2017, 121, 4968.
- (10). (a) For a photochemical generation and reactivity of nitrogen-centered purine nucleoside radicals, see: Kuttappan-Nair V; Samson-Thibault F; Wagner JR Chem. Res. Toxicol 2010, 23, 48. [PubMed: 20000474] (b) Zheng L; Lin L; Qu K; Adhikary A; Sevilla MD; Greenberg MM Org. Lett 2017, 19, 6444. [PubMed: 29125775] (c) Zheng L; Griesser M; Pratt DA; Greenberg MM J. Org. Chem 2017, 82, 3571. [PubMed: 28318253]
- (11). Chiu DT; Duesberg PH Genetica 1995, 95, 103. [PubMed: 7744255]
- (12). AmdU inhibits thymidylate synthetase and thymidine kinase activities: Balzarini J; De Clercq E; Mertes MP; Shugar D; Torrence PF Biochem. Pharmacol 1982, 31, 3673. [PubMed: 7181950]
- (13). AvdU inhibits the replication of HSV-1 and VZV and is cytostatic against HSV-1 and HSV-2 TK gene transfected FM3A tumor cells with its cytostatic effect enhanced 5-fold after short exposure to UV irradiation at 254 nm: Balzarini J; Andrei G; Kumar R; Knaus EE; Wiebe LI; De Clercq E FEBS Lett 1995, 373, 41. [PubMed: 7589430]
- (14). Neef AB; Luedtke NW ChemBioChem 2014, 15, 789. [PubMed: 24644275]

- (15). Ren X; El-Sagheer AH; Brown T *Analyst* 2015, 140, 2671. [PubMed: 25734317]
- (16). (a) Other azido-modified nucleosides have been applied for the metabolic labeling of DNA (ref 16a) and RNA (ref 16b) and for live cell fluorescence imaging (ref 16c): Kukwikila M; Gale N; El-Sagheer AH; Brown T; Tavassoli A *Nat. Chem* 2017, 9, 1089. [PubMed: 29064492]
(b) Nguyen K; Fazio M; Kubota M; Nainar S; Feng C; Li X; Atwood SX; Bredy TW; Spitaleri RC *J. Am. Chem. Soc* 2017, 139, 2148 Warminski M; Kowalska J; Jemielity J *Org. Lett* 2017, 19, 3624 Mamot, A; Sikorski PJ; Warminski M; Kowalska J; Jemielity J *Angew. Chem., Int. Ed* 2017, 56, 15628. (c) Zayas J; Annoual MD; Das JK; Felty Q; Gonzalez WG; Miksovska J; Sharifai N; Chiba A; Wnuk SF *Bioconjugate Chem.* 2015, 26, 1519.
- (17). Krim J; Taourirte M; Grünwald C; Krstic I; Engels J. *Synthesis* 2013, 45, 396.
- (18). Kovacs T; Otvos L *Tetrahedron Lett.* 1988, 29, 4525.
- (19). Liu Z; Liao P; Bi X *Org. Lett* 2014, 16, 3668. [PubMed: 24971828]
- (20). Liang Y; Suzol SH; Wen Z; Artiles AG; Mathivathanan L; Raptis RG; Wnuk SF *Org. Lett* 2016, 18, 1418. [PubMed: 26933954]
- (21). For the review on the reactivities of vinyl azides see: Hu B; DiMagno SG *Org. Biomol. Chem* 2015, 13, 3844. [PubMed: 25731154]
- (22). Suzol SH; Howlader AH; Wen Z; Ren Y; Laverde EE; Garcia C; Liu Y; Wnuk SF *ACS Omega* 2018, 3, 4276. [PubMed: 29732453]
- (23). Fauster K; Hartl M; Santner T; Aigner M; Kreutz C; Bister K; Ennifar E; Micura R *ACS Chem. Biol* 2012, 7, 581. [PubMed: 22273279]
- (24). (a) Bernhard WA *Wiley Ser. React. Intermed. Chem. Biol.* 2009, 2, 41. (b) Sagstuen E; Hole EO In *Electron Paramagnetic Resonance*; Brustolon M, Giamello E, Eds.; John Wiley & Sons, Inc., 2009; p 325. (c) Close D In *Radiation Induced Molecular Phenomena in Nucleic Acids*; Shukla M, Leszczynski J, Eds.; Springer, 2008; Vol. 5, p 493.
- (25). Mandel SM; Singh PND; Muthukrishnan S; Chang M; Krause JA; Gudmundsdottir AD *Org. Lett* 2006, 8, 4207. [PubMed: 16956188]
- (26). Muthukrishnan S; Sankaranarayanan J; Klima RF; Pace TCS; Bohne C; Gudmundsdottir AD *Org. Lett* 2009, 11, 2345. [PubMed: 19432450]
- (27). (a) Bencivenni G; Lanza T; Leardini R; Minozzi M; Nanni D; Spagnolo P; Zanardi G J. *Org. Chem* 2008, 73, 4721. [PubMed: 18471016] (b) Minozzi M; Nanni D; Spagnolo P *Chem. - Eur. J* 2009, 15, 7830. [PubMed: 19551769]
- (28). The axially symmetric anisotropic nitrogen HFCCs and the g values of U-5-C(CH₃) N[•] are found to be nearly identical to the reported values of σ -iminyl radicals from one-electron oxidized 1-methylcytosine and its derivatives: Adhikary A; Kumar A; Bishop CT; Wiegand TJ; Hindi RM; Adhikary A; Sevilla MD *J. Phys. Chem. B* 2015, 119, 11496. [PubMed: 26237072]
- (29). (a) Hou D; Greenberg MM *J. Org. Chem* 2014, 79, 1877. [PubMed: 24559326] (b) Chomicz L; Zdrowowicz M; Kasprzykowski F; Rak J; Buonaugurio A; Wang Y; Bowen KHJ. *J. Phys. Chem. Lett* 2013, 4, 2853.
- (30). Gottinger HA; Zubarev V. e.; Brede O *J. Chem. Soc., Perkin Trans. 2* 1997, 2, 2167.
- (31). (a) Hohman WF; Palcic B; Skarsgard LD *Int. J. Radiat. Biol. Relat. Stud. Phys., Chem. Med* 1976, 30, 247. [PubMed: 1086839] (b) Millar BC; Fielden EM; Smith CE *British J. Cancer. Suppl* 1978, 3, 73.
- (32). Jagiello K; Makurat S; Pere S; Rak J; Puzyn T *Struct. Chem* 2018, 29, 1367.
- (33). AmdU 5'-monophosphate prodrugs have been recently shown to enhance incorporation into the DNA of wild-type cells and animals: Tera M; Glasauer SMK; Luedtke NW *ChemBioChem* 2018, 19, 1939 [PubMed: 29953711]

Figure 1.

(A) ESR spectrum (blue) after radiation-produced one-electron addition to AmdU 4 (2.2 mg/mL) at 77 K (γ -irradiation, 500 Gy) in 7.5 M LiCl/D₂O in dark. Spectra B–E were obtained via stepwise annealing of the sample for 15 min at 140, 160, 165, and 170 K in the dark. All spectra were recorded at 77 K. The radiation produced background Cl₂^{•−} spectrum was subtracted from spectra A and B for clarity. Three reference triangles show Frey's salt resonances with central marker at $g = 2.0056$.

Figure 2.

(A) ESR spectrum (black) after radiation-produced one-electron addition to 12 (1 mg/mL) at 77 K (γ -irradiation, 500 Gy) in 7.5 M LiCl/D₂O in dark. (B–D) Spectra after stepwise annealing at 135, 150, and 165 K recorded at 77 K. The red spectra in (A) and (D) are the simulated spectra. The background $\text{Cl}_2^{\bullet-}$ spectrum has been subtracted from spectra A and B for clarity.

Figure 3. Radiosensitizing effect of 100 μ M azido-modified nucleosides on EMT6 cells: (A) normoxic and (B) hypoxic conditions

Scheme 1.

Synthesis of 5-Azidomethyl Pyrimidine Nucleosides and Their 5'-Triphosphates

Scheme 2.
 Synthesis of 5-Azidovinyl Pyrimidine Nucleosides

Scheme 3.

Formation of π -Aminyl Radical from AmdU and Its Bimolecular Conversion to the σ -Iminyl Radical

Scheme 4. Tautomerization of π Aminyl Radical, Generated from AvdU, to σ -Iminyl Radical