Published in final edited form as:

J Aging Phys Act. 2017 January ; 25(1): 116–122. doi:10.1123/japa.2016-0058.

Rest-Activity Patterns in Older Adults with Heart Failure and Healthy Older Adults

Daniel Liebzeit, RN-BSN [PhD student],

University of Wisconsin-Madison and William S. Middleton Memorial VA

Hospitaldaniel.liebzeit@wisc.edu. Phone: 715-701-0679.

Cynthia Phelan, PhD, RN [Nurse Scientist],

GRECC, William S. Middleton Memorial VA Hospital, Madison, WI. University of Wisconsin-

Madison School of Nursing, Madison, WI.

Chooza Moon, RN-MS [PhD candidate],

candidate, University of Wisconsin-Madison School of Nursing, Madison, WI. University of

Wisconsin-Madison and William S. Middleton Memorial VA Hospital

Roger Brown, PhD [Professor], and

Professor, University of Wisconsin-Madison School of Nursing, Madison, WI. University of

Wisconsin-Madison and William S. Middleton Memorial VA Hospital

Lisa Bratzke, PhD [Assistant Professor]

Assistant Professor, University of Wisconsin-Madison School of Nursing, Madison, WI. University

of Wisconsin-Madison and William S. Middleton Memorial VA Hospital

Abstract

The purpose of this investigation is to examine differences in rest-activity patterns and sleep characteristics in older adults with HF and healthy older adults. The sample included older adults with HF ($n = 20$) and a reference group of healthy older adults ($n = 20$). Traditional cosinor analysis was used to assess three parameters of rest-activity from wrist actigraphy data: amplitude (range of activity), mesor (mean activity), and acrophase (time of peak activity). Traditional sleep characteristics were also determined from actigraphy data: Total Sleep Time, Sleep Latency, Sleep efficiency, and Wake after Sleep Onset. The HF group demonstrated significantly lower mesor and amplitude than the reference group ($p < 0.01$). The HF group had significantly greater TST ($p < 0.01$), but the groups had similar SE, SL, and WASO. Despite similar sleep characteristics to healthy older adults, overall rest-activity patterns were significantly dampened in those with HF.

Keywords

heart failure; circadian rhythm; rest-activity patterns; sleep disturbance; geriatrics

Over 5 million people in the United States of America suffer from Heart Failure (HF) (“Division for Heart Disease and Stroke Prevention: Heart Failure Fact Sheet,” 2014). HF is commonly associated with activity intolerance (“Division for Heart Disease and Stroke Prevention: Heart Failure Fact Sheet,” 2014) and sleep disturbances (both subjective and objective) (“Division for Heart Disease and Stroke Prevention: Heart Failure Fact Sheet,”

2014; M. Y. Wang, Hung, & Tsai, 2011). Subjectively, individuals with HF report overall sleep duration similar to healthy adults (Redeker & Stein, 2006). However, between 50–96% of individuals with HF report poor sleep quality (Garcia et al., 2012; Moon, Phelan, Lauver, & Bratzke, 2015; Redeker & Stein, 2006; T. Wang, Lee, Tsay, & Tung, 2010), as compared to 16–51% of healthy adults (Kim et al., 2015; Redeker & Stein, 2006). Findings regarding daytime sleepiness and daytime dysfunction due to poor sleep in HF are variable. 7%–44% of those surveyed report excessive daytime sleepiness (Moon et al., 2015; Redeker et al., 2010; Redeker & Stein, 2006), and there are inconsistent reports of whether daytime dysfunction is more common in HF than healthy adults (Moon et al., 2015; Redeker & Stein, 2006).

Objective sleep methods (e.g., actigraphy) support the subjective findings of normal sleep duration and disturbed sleep (Redeker & Stein, 2006). Individuals with HF show more frequent awakenings during their sleep interval but a lower percentage of time awake, as compared to healthy adults (Redeker & Stein, 2006).

Disturbed sleep has been linked to increased risk of mortality in individuals with HF (Reinhard et al., 2013). Poor sleep quality in individuals with HF has also been associated with reduced cognitive performance, reduced quality of life, depression, and reduced functional independence (Garcia et al., 2012). Reports of daytime dysfunction in HF, due to sleep, have been associated with poorer cognitive function (Moon et al., 2015).

Rest-activity patterns can be observed as a biomarker of an individual's circadian rhythm. Circadian regulation, that is, consistently following a 24-hour rest-activity pattern, is associated with better health outcomes, including decreased mortality (Evans & Davidson, 2013). Dysregulated rest-activity is a manifestation of a dysregulated circadian system, making rest-activity patterns a commonly used measure of circadian rhythmicity (Gehrman et al., 2005). Rest-activity methods are less invasive and data are easier to collect than other measures of circadian rhythms, such as core body temperature or hormonal markers (i.e. melatonin) (Gehrman et al., 2005). When identified, dysregulated circadian rhythms are amenable to interventions (i.e. changes in the environment, biological processes, etc.) (Gehrman et al., 2005).

Rest-activity analysis goes beyond traditional sleep quality and sleep characteristics to examine rest and activity levels over a 24-hour day. Sleep characteristics include total sleep time (time in bed asleep), sleep latency (time in bed before falling asleep), wake after sleep onset (time spent awake after falling asleep), and sleep efficiency (percent of time spent in bed asleep). These sleep characteristics provide no new information about daytime rest and activity (see Figure 1). A study of 24-hour rest-activity patterns demonstrates whether an individual's circadian rhythm is synchronized or dysregulated. Rest-activity analysis reveals characteristics about an individual's nighttime activity, as well as its relationship to daytime activity. Rest-activity patterns have been studied extensively in older adults suffering from dementia, where disrupted nighttime sleep and extending sleep into daytime hours is common. Findings indicate impaired circadian rhythms and low levels of exposure to bright light result in loss of important environmental cues that regulate sleep (Gehrman et al., 2005). In Dementia, circadian interventions to improve nighttime sleep have been effective

(Deschenes & McCurry, 2009; Gehrman et al., 2005; Skjerve, Bjorvatn, & Holsten, 2004). A more thorough understanding of 24-hour rest-activity patterns may contribute interventions to improve both daytime functioning and nighttime sleep in HF.

Rest-activity patterns have not been examined in-depth in the HF population. A study of individuals suffering from HF with and without anergia (lack of energy) demonstrated very different rest-activity patterns (Maurer et al., 2009). Individuals with anergia and HF had a significantly smaller range of activity between the night and day than those without anergia (Maurer et al., 2009). No studies have compared rest-activity patterns of older adults suffering from HF to healthy older adults. This gap in rest-activity assessment is important since older adults with HF are reported to have disturbed sleep and lower daytime activity, and expected daytime sleepiness following disturbed sleep is not consistently reported. The purpose of this investigation is to: (1) examine differences in rest-activity patterns (e.g. mean activity, range of activity, and time of peak activity) between older adults with HF and healthy older adults, (2) further elucidate objective sleep characteristics (e.g. total sleep time, sleep efficiency, wake after sleep onset, and sleep latency) in older adults with HF, when compared to healthy older adults, and (3) explore factors that may explain group differences in rest-activity patterns.

Methods

Sample

Two samples were recruited for this study. Older adults (aged 58 and older) with HF were recruited from a Midwestern HF clinic (Bauer et al., 2012), as part of a larger HF study examining the psychometric properties of a battery of neuropsychological tests (Bauer & Pozehl, 2011). The inclusion and exclusion criteria for the parent HF study can be seen in Table 1 (Bauer & Pozehl, 2011). Individuals were included in the current analysis if they had at least five days of actigraphy data ($n = 20$).

A reference sample of healthy older adults with actigraphy data was obtained from the Midlife Development in the United States (MIDUS) II study. Comprehensive biological assessments were conducted on a subsample of the 1,255 MIDUS participants from 2004–2009 (Ryff, Seeman, & Weinstein, 2010). MIDUS participants, with at least 5 days of actigraphy data ($n = 440$), were selected to match (by age, gender, and race) the HF group. Individuals were excluded if they had a diagnosis of HF or depression. The group of participants selected for this study ($n = 20$) were matched within 0.80 years of age, on average, and were matched for race and gender to the older adults in the HF group. The MIDUS study design and methods are described in detail elsewhere (Ryff et al., 2010).

Study Design

All study procedures for this secondary analysis were reviewed and received IRB approval.

Measures

Demographic data were obtained via chart review and self-report in the parent studies.

Number of Comorbidities: A number of comorbidities score was derived by summing the number of comorbid health conditions identified (via chart abstraction for the HF group and self-report for the reference group). This method is consistent with a well-established severity of illness scale (DUSOI, Duke University) (Parkerson, Broadhead, & Tse, 1993; Parkerson, Schuntermann, & Sattler, 1998).

Depression Score: Depression was assessed using the Personal Health Questionnaire Depression Scale (PHQ-8)(Kroenke et al., 2009) (HF group) and the Center for Epidemiological Studies Depression Inventory (CES-D) (Radloff, 1977) (reference group). The PHQ-8 is a self-administered 8-item scale to diagnose and assess severity of depressive disorders in clinical studies (Kroenke et al., 2009). The measure has demonstrated validity in a large epidemiological population-based study (Kroenke et al., 2009). The CES-D inventory is a 20-item self-report scale designed to measure depressive symptomatology (Radloff, 1977). This measure emphasizes depressed mood or affect and has demonstrated validity in community-based older adult population (Radloff, 1977). Z-scores were created to compare the respective scales in the subsequent statistical analyses (Rubin, 2012).

Heart Failure Status: The following data were obtained via chart abstraction (HF group): duration of HF diagnosis, left ventricular ejection fraction (LVEF), total number of medications, New York Heart Association (NYHA) class, and etiology of HF (Bauer et al., 2012).

Rest-Activity Patterns: Wrist-actigraphy, using the Actiwatch® 64, was used to assess 24-hour rest-activity patterns (Ryff et al., 2010). All participants wore wrist-actigraphy for a continuous 7-day period (Bauer & Pozehl, 2011; Ryff et al., 2010). The reference group also completed sleep diaries during the period that the watch was worn, to track subjective sleep patterns, and the time in and out of bed was used to mark Rest and Exclusion Intervals in the Actiwatch® (Ryff et al., 2010). Traditional cosinor analysis was used to assess three parameters of rest and activity: amplitude, mesor, and acrophase (Gehrman et al., 2005). Amplitude is the difference between the maximum and minimum activity level; lower values indicate a weaker rhythm. Mesor is the 24-hour activity level; lower values indicate less overall activity. Acrophase is the time, during the 24-hour day, that activity peaks. The R-squared (RSQ) value indicates the reduction in squared error from using a model to summarize data compared to using the mean.

Sleep Characteristics: Wrist-actigraphy was also used to assess sleep characteristics (Ryff et al., 2010). This study explored standard Actiware® sleep characteristics: Total Sleep Time (TST), Sleep Latency (SL), Wake after Sleep Onset (WASO), and Sleep Efficiency (SE). TST is the time in bed asleep. SL is the time in bed before falling asleep. WASO is the time spent awake after falling asleep. SE is the percent of time spent in bed asleep (e.g., time asleep/time in bed).

Data Analysis

Actigraphy data were downloaded to Respironics Actiware® software Version 5.59. Sleep characteristics were obtained using the Actiware® software standard analysis. Rest-activity

data (activity each epoch), were exported from Actiware® software to Microsoft Excel. The activity data collected outside of the intended data collection interval (i.e. the time that the Actiwatch® was turned 'on' before the participant started wearing the device and after final removal) was excluded from data analysis. Rest-activity data collected in 15-second and 30-second epochs were converted to 1-minute epochs by averaging the levels of activity in each 1-minute interval. For the analysis, five full 24-hour days were randomly selected for each HF group participant, using simple random selection. The total epochs were comparable between the two groups. Data were exported into SAS for traditional cosinor analysis.

Statistical analyses were conducted using IBM® SPSS® Statistics Version 22. Paired samples t-tests were conducted to describe the sample and test for group differences in demographics and health characteristics. A Chi-square test of independence was used to test for group differences in categorical variables (e.g. education, gender, and race). Paired samples t-test was used to examine group differences in sleep characteristics and rest-activity patterns. Multiple regression was used to test predictors of the rest-activity patterns seen in these groups. Diagnosis of HF, depression scores, and number of comorbidities were included as predictors of each of the three rest-activity parameters in the multiple regression analysis. Depression has been associated with sleep and daytime function (Jausset et al., 2011) and increased number of comorbidities has been associated with lower physical activity in older adults (Beveridge et al., 2015). The presences of particular co-morbid conditions were not included as covariates due to the limited sample size. Counts of common comorbid conditions in the HF and reference groups are shown in Table 2. Age, gender, and race were not included because the groups were matched on these variables.

Results

Demographics

The demographic information for the two groups is displayed in Table 3. The HF group had significantly more comorbidities (mean = 5.25; SD = 2.67) than the reference group (mean = 1.90; SD = 1.74; $p < 0.01$). It was not feasible to determine the total number of medications for the reference group.

Rest-Activity Patterns

The HF group demonstrated significantly lower mesor (mean = 69.88; SD = 27.72) than the reference group (mean = 225.77; SD = 98.30; $p < 0.01$) (Table 4). Additionally, the HF group had significantly lower activity amplitude (mean = 58.58; SD = 27.96) than the reference group (mean = 192.01; SD = 85.87; $p < 0.01$). The HF group had a lower average RSQ value (mean = .4575; SD = .154) for the cosinor model than the reference group (mean = .6275; SD = .123; $p < 0.01$). No significant group differences were noted in acrophase (the peak of daytime activity).

In multiple regression analyses, HF diagnosis was the only significant predictor of lower mesor (95% Confidence Interval (CI) [91.12, 209.58]; $p < 0.01$) and amplitude (95% CI [76.59, 182.91]; $p < 0.01$), after controlling for depression score and number of

comorbidities (Table 5). Neither HF, depression, nor the total number of comorbidities were significant predictors of acrophase (Table 5).

Sleep Characteristics

The overall characteristics of sleep in those with HF were similar to the healthy controls (Table 6), except the HF group had significantly greater TST (mean = 470.06 minutes; SD = 67.08) than the reference group (mean = 394.89 minutes; SD = 75.05; $p < 0.01$). There were no significant differences in SL, SE, or WASO between the HF and reference groups.

Discussion

This study sought to: (1) examine differences in rest-activity patterns between older adults with HF and healthy older adults, (2) further elucidate objective sleep characteristics in older adults with HF, when compared to healthy older adults, and (3) explore factors that may explain group differences in rest-activity patterns. Our findings revealed significant differences in 24-hour mean activity levels and the range of activity in the older adults with HF, despite similar sleep characteristics among the groups.

The HF group had significantly dampened rest-activity pattern: the mean activity level (mesor) and range of activity during the 24 hours (amplitude) was lower than the reference group. The lower mesor suggests that older adults with HF are less active, overall, than the comparison group. Changes in rest-activity patterns have not yet been well documented, but a previous study has reported lower levels of daytime physical activity in individuals with HF (Oka, Stotts, Dae, Haskell, & Gortner, 1993). Higher rates of napping (planned and unplanned) and activity intolerance (“Division for Heart Disease and Stroke Prevention: Heart Failure Fact Sheet,” 2014) in older adults with HF may also explain these findings. Higher activity levels at night are consistent with previous reports of difficulty staying asleep and frequent interruptions in sleep (Redeker et al., 2010; Redeker & Stein, 2006).

The lower amplitude suggests that older adults with HF do not reach the high levels of daytime activity observed in healthy adults. Higher maximum levels of daily activity have been associated with fewer harmful events, including death, hospitalizations, emergency department visits, and intercurrent illnesses (Howell et al., 2010). An explanation for these findings could be the severity of HF observed in this study. The HF group in this study did have moderate HF, and, therefore, could have lower activity tolerance. Older adults with higher NYHA class in HF fatigue more easily and/or tolerate less activity.

The finding of lower amplitude in the HF group matches what was found previously in individuals suffering from HF and anergia (lack of energy) (Maurer et al., 2009). It is unclear whether anergia contributed to the findings in this study, because data about symptoms of anergia were not collected. Other symptoms not measured, such as pain or fatigue, could also contribute to some of the observed differences between the groups. The findings of lower mesor and amplitude match what has been found in a group of women with endometrial cancer, post-surgery, compared to women without a history of cancer (Rumble et al., 2015). This is the first study to compare the mesor and amplitude rest-activity patterns in HF to a group of healthy older adults.

In this investigation, we expected that the comorbid conditions found routinely in HF (Ho, Pinsky, Kannel, Levy, & Pitt, 1993) could explain the dampened rest-activity pattern; but it did not. The older adults with HF had significantly more comorbid conditions than the control, but the HF diagnosis persists as the only significant predictor of differences in mean activity level (mesor) and range of activity (amplitude). The differences in number of comorbidities between the HF and reference groups did not explain the differences observed in rest-activity patterns. While it may be feasible that certain comorbid conditions that occur with HF (i.e. diabetes and HF) may impact rest-activity patterns, we were not able to explore that assumption.

Certain medications common in older adults with HF, such as beta-blockers, may dampen activity tolerance, and, therefore, 24-hour activity levels. Potential issues with dosing or side effects of specific medications can affect daytime sleepiness or function. The multiplicative effect of many medications (i.e. polypharmacy) can further increase feelings of fatigue, and, thereby, decrease activity. It is important to understand how medication management in older adults with HF may impact rest-activity patterns. Differences in the databases did not allow for direct analysis of specific medication, but in post hoc analysis, we found the number of medications, but not LVEF or NYHA class, was related to differences in mesor and amplitude within the HF group.

In addition to disruptions in rest-activity patterns, the HF group had significantly lower R-squared value in the cosinor analysis, indicating lower circadian rhythmicity. In other words, the lower R-squared means that the older adults with HF conform less to the typical rise of activity in the day and lower activity at night (i.e. the expected circadian rhythm of activity). A more flat level of activity throughout the day and night would contribute to a low R-squared and circadian rhythmicity. The lower mesor and amplitude in the older adults with HF supports this finding of a more flat circadian rhythm. The lower rhythmicity in older adults with HF could reflect a problem of both nighttime and daytime dysfunction. Individuals with HF may exhibit frequent nighttime awakenings due to how well their HF is managed, leading to circadian rhythm disturbances. For example, variability in adherence to dietary sodium restrictions, nighttime cardiovascular-related fluctuations in fluid volume and adherence to medication regimes, and daytime napping can precipitate nighttime awakenings.

Unlike a previous study using wrist actigraphy, older adults with HF had significantly more total sleep time (TST) than the comparison group (Redeker & Stein, 2006). The comparison group in this study and the previous study have similar sleep duration, which indicates that the discrepancy is between the sleep duration of the two HF groups (Redeker & Stein, 2006). One possible explanation for this finding is that the previous study recruited individuals with more advanced HF (i.e. NYHA classification) and lower ejection fractions than what was observed in this study. The previous study had an average LVEF of 22.92 (SD = 8.99) (Redeker & Stein, 2006), compared to 30.75 (SD = 7.99) in this study. The previous study also had 4 (6.8%) individuals with NYHA class IV HF (Redeker & Stein, 2006), while this study had none. Results from our study suggest that older adults with HF may spend more time sleeping per day (total sleep time), without having any significant differences in sleep efficiency, wake after sleep onset, and time in bed until falling asleep (sleep latency).

Overall, the sleep characteristics were very similar to the group of healthy older adults. Since our results contradict what has been documented previously, confirmatory follow-up studies should be conducted.

This study has some limitations. The sample size is small and lacks diversity, so generalizability is decreased. The majority of the sample was white, male, with at least a high school education. Anthropometric data (e.g. BMI or waist circumference) for the participants were not available and could have explained some of the differences observed. The wrist-actigraph used in both groups could not indicate off-wrist status (i.e. whether the participant removed the Actiwatch® at any time during the intended data collection interval). Newer technology can address this in future studies.

Future studies are needed to examine how factors, such as increased activity or napping restrictions, may impact rest-activity patterns in older adults with HF. Additionally, further understanding of rest-activity patterns may reveal opportunities to improve both sleep and daytime function in this group. Increased activity during the day could improve sleep at night. Consolidation of daytime napping may help older adults with HF better consolidate their nighttime sleep. Use of daytime diaries or accelerometry may help to determine potential explanations of lower daytime activity and may provide a new perspective for interventions (i.e. increased activity or napping restrictions) to promote environmental synchrony and more normal circadian patterns of rest-activity. Finally, the adverse outcomes associated with abnormal patterns of rest-activity should be investigated in subsequent analysis; it would be important to focus on how dampened rest-activity patterns may specifically affect older adults suffering from HF.

This study is the first to explore rest-activity patterns among older adults with HF, compared to healthy older adults. This study contributes to current research on sleep in HF by revealing that, in spite of very similar objective sleep characteristics between older adults with HF and healthy older adults, the rest-activity patterns in older adults with HF are unique. Findings suggest that examination of factors influencing rest-activity patterns in HF may reveal opportunities to manage symptoms of daytime dysfunction.

Acknowledgments:

The MIDUS study was supported by a grant from the National Institute on Aging (P01-AG020166) to conduct a longitudinal follow-up of the MIDUS (Midlife in the U.S.) investigation. The original study was supported by the John D. and Catherine T. MacArthur Foundation Research Network on Successful Midlife Development. The MIDUS study was supported by the following grants: M01-RR023942 (Georgetown), M01-RR00865 (UCLA) from the General Clinical Research Centers Program and 1UL1RR025011 (UW) from the Clinical and Translational Science Award (CTSA) program of the National Center for Research Resources, National Institutes of Health. MIDUS research participants were admitted to or studied on the University of Wisconsin-Clinical and Translational Research Core (UW-CTRC). The described study was supported by the National Institute of Nursing Research of the National Institutes of Health under Award Number R00NR012773 (Bratzke) and in part by Career Development Award #IK2CX000535 from the U.S. Department of Veterans Affairs Clinical Science Research & Development Service (Phelan). The content is solely the responsibility of the authors and does not represent the official views of the National Institutes of Health, the U.S. Department of Veterans Affairs or the United States Government.

References

- Bauer L, & Pozehl B (2011). Measurement of cognitive function in chronic heart failure: a feasibility study. *Applied Nursing Research*, 24(4), 223–228. [PubMed: 20974084]

- Bauer L, Pozehl B, Hertzog M, Johnson J, Zimmerman L, & Filipi M (2012). A brief neuropsychological battery for use in the chronic heart failure population. *European Journal of Cardiovascular Nursing*, 11(2), 223–230. [PubMed: 21514892]
- Beveridge C, Knutson K, Spampinato L, Flores A, Meltzer DO, Van Cauter E, & Arora VM (2015). Daytime Physical Activity and Sleep in Hospitalized Older Adults: Association with Demographic Characteristics and Disease Severity. *Journal of the American Geriatrics Society*, 63(7), 1391–1400. doi:10.1111/jgs.13520 [PubMed: 26131982]
- Deschenes CL, & McCurry SM (2009). Current treatments for sleep disturbances in individuals with dementia. *Curr Psychiatry Rep*, 11(1), 20–26. [PubMed: 19187704]
- Division for Heart Disease and Stroke Prevention: Heart Failure Fact Sheet. (2014). Retrieved from http://www.cdc.gov/dhdsdp/data_statistics/fact_sheets/fs_heart_failure.htm
- Evans JA, & Davidson AJ (2013). Health consequences of circadian disruption in humans and animal models. *Prog Mol Biol Transl Sci*, 119, 283–323. doi:10.1016/b978-0-12-396971-2.00010-5 [PubMed: 23899601]
- Garcia S, Alosco ML, Spitznagel MB, Cohen R, Raz N, Sweet L, ... Gunstad J (2012). Poor sleep quality and reduced cognitive function in persons with heart failure *Int J Cardiol* (Vol. 156, pp. 248–249). Netherlands. [PubMed: 22360947]
- Gehrman P, Marler M, Martin JL, Shochat T, Corey-Bloom J, & Ancoli-Israel S (2005). The relationship between dementia severity and rest/activity circadian rhythms. *Neuropsychiatr Dis Treat*, 1(2), 155–163. [PubMed: 18568061]
- Ho KK, Pinsky JL, Kannel WB, Levy D, & Pitt B (1993). The epidemiology of heart failure: the Framingham Study. *Journal of the American College of Cardiology*, 22(4s1), A6–A13.
- Howell J, Strong BM, Weisenberg J, Kakade A, Gao Q, Cuddihy P, ... Maurer MS (2010). Maximum daily 6 minutes of activity: an index of functional capacity derived from actigraphy and its application to older adults with heart failure. *J Am Geriatr Soc*, 58(5), 931–936. doi:10.1111/j.1532-5415.2010.02805.x [PubMed: 20374397]
- Jausse I, Bouyer J, Ancelin ML, Akbaraly T, Peres K, Ritchie K, ... Dauvilliers Y (2011). Insomnia and daytime sleepiness are risk factors for depressive symptoms in the elderly. *Sleep*, 34(8), 1103–1110. doi:10.5665/sleep.1170 [PubMed: 21804672]
- Kim CW, Chang Y, Zhao D, Cainzos-Achirica M, Ryu S, Jung HS, ... Sung E (2015). Sleep Duration, Sleep Quality, and Markers of Subclinical Arterial Disease in Healthy Men and Women. *Arterioscler Thromb Vasc Biol*. doi:10.1161/atvbaha.115.306110
- Kroenke K, Strine TW, Spitzer RL, Williams JB, Berry JT, & Mokdad AH (2009). The PHQ-8 as a measure of current depression in the general population. *Journal of affective disorders*, 114(1), 163–173. [PubMed: 18752852]
- Maurer MS, Cuddihy P, Weisenberg J, Delisle S, Strong BM, Gao Q, ... Howell J (2009). The prevalence and impact of anergia (lack of energy) in subjects with heart failure and its associations with actigraphy. *J Card Fail*, 15(2), 145–151. doi:10.1016/j.cardfail.2008.10.021 [PubMed: 19254674]
- Moon C, Phelan CH, Lauver DR, & Bratzke LC (2015). Is sleep quality related to cognition in individuals with heart failure? *Heart Lung*, 44(3), 212–218. doi:10.1016/j.hrtlng.2015.02.005 [PubMed: 25796476]
- Oka RK, Stotts NA, Dae MW, Haskell WL, & Gortner SR (1993). Daily physical activity levels in congestive heart failure. *Am J Cardiol*, 71(11), 921–925. [PubMed: 8465782]
- Parkerson GR Jr., Broadhead WE, & Tse CK (1993). The Duke Severity of Illness Checklist (DUSOI) for measurement of severity and comorbidity. *J Clin Epidemiol*, 46(4), 379–393. [PubMed: 8483003]
- Parkerson GR Jr., Schuntermann MF, & Sattler LL (1998). [The Duke Severity of Illness Checklist]. *Rehabilitation (Stuttg)*, 37(1), suppl I–XV.
- Radloff LS (1977). The CES-D scale a self-report depression scale for research in the general population. *Applied psychological measurement*, 1(3), 385–401.
- Redeker NS, Jeon S, Muench U, Campbell D, Walsleben J, & Rapoport DM (2010). Insomnia symptoms and daytime function in stable heart failure. *Sleep*, 33(9), 1210–1216. [PubMed: 20857868]

- Redeker NS, & Stein S (2006). Characteristics of sleep in patients with stable heart failure versus a comparison group. *Heart Lung*, 35(4), 252–261. doi:10.1016/j.hrtlng.2005.10.007 [PubMed: 16863897]
- Reinhard W, Plappert N, Zeman F, Hengstenberg C, Riegger G, Novack V, ... Arzt M (2013). Prognostic impact of sleep duration and sleep efficiency on mortality in patients with chronic heart failure. *Sleep Med*, 14(6), 502–509. doi:10.1016/j.sleep.2012.12.014 [PubMed: 23628241]
- Rubin A (2012). *Statistics for evidence-based practice and evaluation*: Cengage Learning.
- Rumble ME, Rose SL, White KH, Moore AH, Gehrman P, Benca RM, & Costanzo ES (2015). Circadian actigraphic rest-activity rhythms following surgery for endometrial cancer: A prospective, longitudinal study. *Gynecol Oncol*, 137(3), 448–455. doi:10.1016/j.ygyno.2015.04.001 [PubMed: 25906914]
- Ryff CD, Seeman T, & Weinstein M (2010). National Survey of Midlife Development in the United States (MIDUS II): Biomarker project, 2004–2009.
- Skjerve A, Bjorvatn B, & Holsten F (2004). Light therapy for behavioural and psychological symptoms of dementia. *Int J Geriatr Psychiatry*, 19(6), 516–522. doi:10.1002/gps.1087 [PubMed: 15211528]
- Wang MY, Hung HL, & Tsai PS (2011). The sleep log and actigraphy: congruency of measurement results for heart failure patients. *J Nurs Res*, 19(3), 173–180. doi:10.1097/JNR.0b013e318229c42f [PubMed: 21857324]
- Wang T, Lee S, Tsay S, & Tung H (2010). Factors influencing heart failure patients' sleep quality. *Journal of Advanced Nursing*, 66(8), 1730–1740. doi:10.1111/j.1365-2648.2010.05342.x [PubMed: 20557385]

Figure 1. Circadian Rhythm Analysis versus Traditional Sleep Characteristics

Note. The shaded region represents the time of day that traditional sleep characteristics typically reflect. The red line of activity shows the information gathered from circadian rest-activity analysis, including three parameters: mesor, amplitude, and acrophase. These parameters are determined from activity levels throughout the entire day.

Table 1

Inclusion/Exclusion Criteria for Subjects with HF

Inclusion Criteria:	
>21 years of age	
English speaking and writing	
Duration of HF >6 months	
Stable medication regimen for past 4 weeks	
Exclusion Criteria:	
Current diagnosis of neurological illness or dementia (e.g. Alzheimer's dementia, Parkinson's, epilepsy)	
History of substance abuse/treatment	
PhQ-8 >10, indicating major depression	
History of stroke	
Hepatic insufficiency (AST/ALT more than twice normal limits)	
Severe renal failure (serum creatinine >2.5 mg/dl)	
Anemia	
Left ventricular assist device	
History of the following within 3 months of study enrollment:	
Acute myocardial infarction	
Unstable angina	
Coronary artery bypass graft surgery	
percutaneous transluminal coronary angioplasty	
Biventricular pacemaker insertion	
History of Hospitalization within past 4 weeks	

(Bauer & Pozehl, 2011) Note. ALT = alanine aminotransferase; AST = aspartate aminotransferase.

Table 2

Comorbidity Counts by Group

	HF group (<i>n</i> = 20)	Reference group (<i>n</i> = 20)
Comorbidity	<i>n</i> (%)	<i>n</i> (%)
Diabetes Mellitus	4 (20)	2 (10)
Cancer	9 (45)	3 (15)
Liver Disease	0 (0)	1 (5)
Hypertension	10 (50)	1 (5)
Arthritis	2 (10)	13 (65)

Table 3

Demographic and Health Characteristics by Group

	HF group (<i>n</i> = 20)	Reference group (<i>n</i> = 20)
Demographic	mean (SD)	mean (SD)
Age	74.40 (7.49)	74.00 (7.12)
Male, <i>n</i> (%)	12 (60)	12 (60)
White, <i>n</i> (%)	20 (100)	20 (100)
Education, <i>n</i> (%)		
<HS	1 (5)	3 (15)
HS or equiv.	9 (45)	10 (50)
College	6 (30)	3 (15)
Bachelor's Degree	3 (15)	2 (10)
Master's Degree	1 (5)	2 (10)
Health Characteristics	Mean (SD)	Mean (SD)
Depression	2.80 (3.09)	5.65 (5.01)
Comorbidities *	5.25 (2.67)	1.90 (1.74)
Duration of HF (months)	56.55 (46.68)	
LVEF	30.75 (7.99)	
Medications (#)	10.3 (3.61)	
NYHA Class 2, <i>n</i> (%)	8 (40)	
NYHA Class 3, <i>n</i> (%)	12 (60)	
Ischemic HF, <i>n</i> (%)	14 (70)	

Note.

* $p < 0.01$. Depression was assessed using the PHQ-8 for the HF group and CES-D for the reference group. The scores reflected in this table represent the raw depression scores, not the adjusted z-scores used for analysis.

Table 4

Rest-Activity Patterns of the HF and Reference Groups

	HF group (<i>n</i> = 20)	Reference group (<i>n</i> = 20)
Variable	mean (SD)	mean (SD)
Mesor *	69.88 (27.72)	225.77 (98.30)
Amplitude *	58.58 (27.96)	192.01 (85.87)
Acrophase	13.95 (1.60)	13.67 (1.07)
RSQ *	.4575 (.154)	.6275 (.123)

Note.

*
p < 0.01.

Table 5

Multiple Regression Analysis Output for Mesor, Amplitude, and Acrophase in Both Groups

Model 1	Dependent variable: Mesor	β	Std. Error	95% Confidence interval	
	Constant	228.91	19.11	190.15	267.66
	HF	-150.35 *	29.21	-209.58	-91.12
	Comorbidities	-1.65	5.54	-12.88	9.57
	Depression score	20.88	12.49	-4.45	46.21
Model 2	Dependent variable: Amplitude				
	Constant	194.10	17.15	159.32	228.89
	HF	-129.75 *	26.21	-182.91	-76.59
	Comorbidities	-1.10	4.97	-11.17	8.98
	Depression score	14.58	11.21	-8.16	37.32
Model 3	Dependent variable: Acrophase				
	Constant	13.79	0.37	13.03	14.55
	HF	0.49	0.57	-0.66	1.65
	Comorbidities	-0.06	0.11	-0.28	0.16
	Depression score	0.07	0.24	-0.42	0.57

Note.

*
p < 0.01.

Table 6

Sleep Characteristics of the HF and Reference Groups

	HF group (<i>n</i> = 20)	Reference group (<i>n</i> = 20)
Variable	mean (SD)	mean (SD)
TST (minutes) *	470.06 (67.08)	394.89 (75.05)
SL (minutes)	22.94 (23.82)	34.03 (20.91)
SE (%)	84.08 (5.68)	80.85 (11.43)
WASO (minutes)	52.90 (22.65)	45.89 (19.66)

Note.

*
 $p < 0.01$. TST = total sleep time; SE = sleep efficiency; SL = sleep latency; WASO = wake after sleep onset.