Published in final edited form as:

Chemistry. 2019 March 07; 25(14): 3662–3674. doi:10.1002/chem.201805919.

Evaluation of 1,2,3-Triazoles as Amide Bioisosteres In Cystic Fibrosis Transmembrane Conductance Regulator Modulators VX-770 and VX-809

Jake E. Doiron^[a], Christina A. Le^[b], Britton K. Ody^[a], Jonathon B. Brace^[a], Savannah J. Post^[a], Nathan L. Thacker^[a], Harrison M. Hill^[a], Prof. Gary W. Breton^[a], Matthew J. Mulder^[c], Sichen Chang^[c], Thomas M. Bridges^[c], Liping Tang^{[d],[e]}, Wei Wang^{[e],[f]}, Prof. Steven M. Rowe^{[d],[e],[f]}, Prof. Stephen G. Aller^[b], and Prof. Mark Turlington^[a]

^[a]Department of Chemistry and Biochemistry, Berry College, Mount Berry, Georgia 30165 (USA), mturlington@berry.edu

^[b]Department of Pharmacology and Toxicology, University of Alabama at Birmingham, Birmingham, Alabama 35205 (USA), sgaller@uab.edu

^[c]Vanderbilt Center for Neuroscience Drug Discovery, Nashville, Tennessee 37232 (USA)

^[d]Departments of Medicine and Pediatrics, University of Alabama at Birmingham, Birmingham, Alabama 35205 (USA)

^[e]Gregory Fleming James Cystic Fibrosis Research Center, University of Alabama at Birmingham, Birmingham, Alabama 35205 (USA)

^[f]Department of Cell, Development, and Integrative Biology, University of Alabama at Birmingham, Birmingham, Alabama 35205 (USA)

Abstract

The 1,2,3-triazole has been successfully utilized as an amide bioisostere in multiple therapeutic contexts. Based on this precedent, triazole analogs derived from VX-809 and VX-770, prominent amide-containing modulators of the cystic fibrosis transmembrane conductance regulator (CFTR), were synthesized and evaluated for CFTR modulation. Triazole **11**, derived from VX-809, displayed markedly reduced efficacy in F508del-CFTR correction in cellular TECC assays in comparison to VX-809. Surprisingly, triazole analogs derived from potentiator VX-770 displayed no potentiation of F508del, G551D, or WT-CFTR in cellular Ussing chamber assays. However, patch clamp analysis revealed that triazole **60** potentiates WT-CFTR similarly to VX-770. The efficacy of **60** in the cell-free patch clamp experiment suggests that the loss of activity in the cellular assay could be due to the inability of VX-770 triazole derivatives to reach the CFTR binding site. Moreover, in addition to the negative impact on biological activity, triazoles in both structural classes displayed decreased metabolic stability in human microsomes relative to the analogous amides. In contrast to the many studies that demonstrate the advantages of using the

Supporting information for this article is given via a link at the end of the document.

Conflict of interest

The authors declare no conflict of interest.

1,2,3-triazole, these findings highlight the negative impacts that can arise from replacement of the amide with the triazole and suggest that caution is warranted when considering use of the 1,2,3-triazole as an amide bioisostere.

Graphical Abstract

Trying out the triazole as an amide bioisostere in CFTR modulators VX-809 and VX-770 revealed that triazole derivatives of VX-809 and VX-770 displayed significantly decreased efficacy or inactivity in cellular assays. In addition, triazole derivatives displayed decreased metabolic stability in hepatic microsomal assays. These findings represent an important counterpoint to the many successful uses of the triazole as an amide bioisostere.

Keywords

triazole; bioisostere; CFTR; corrector; potentiator

Introduction

Since the development of efficient synthetic methodologies to prepare 1,2,3-triazoles in the early 2000's,^[1] the use of the 1,2,3-triazole as an amide bioisostere has proven to be a widely successful strategy in medicinal chemistry.^[2–5] As shown in Figure 1, the basis of the effective bioisosteric relationship is due to the similar topological and electronic characteristics of the 1,4-disubstituted 1,2,3-triazole and the *trans*-amide. Substituents located at the 1-and 4-positions of the triazole are separated by ~5.0 to 5.1 Å, which is close to the ~3.8 to 3.9 Å distance between the amide substituents.^[3] The triazole also exhibits similar H-bonding capabilities to the amide, with the triazole N3 serving as an H-bond acceptor in an analogous spatial location to the amide carbonyl.^[6] Furthermore, due to the substantial dipole moment of the triazole ring calculated to be ~5 Debye,^[5] the C5 hydrogen atom of the triazole has been demonstrated to be capable of functioning as an H-bond donor that occupies a comparable position to the amide hydrogen atom.^[7]

Impressively, numerous successful applications of the 1,2,3-triazole as an amide bioisostere have been reported as recently reviewed by Passarella and coauthors.^[2] These examples span diverse therapeutic contexts including antiviral agents,^[8] chemotherapeutics,^[9–11] antibiofilm agents,^[12] anti-nociceptives,^[13] and antipsychotics.^[14] Advantageously, use of the 1,2,3-triazole in place of the amide often affords increased biological activity, as demonstrated for viral infectivity factor (Vif) inhibitors **1** and **2** (Figure 2).^[8] In this example, the potency of amide **1** (IC₅₀ = 6 μM) was improved substantially in triazole **2** (IC₅₀ = 1.2 μM). Furthermore, the discovery that the triazole could be substituted for the amide enabled use of the Cu-catalyzed azide-alkyne cycloaddition (CuAAC) reaction to

rapidly discover highly potent triazole-containing inhibitor **3** ($IC_{50} = 10$ nM).^[8] In addition to CuAAC, the triazole linkage also opens the door to the use of target-guided in situ click chemistry (isCC)^[15] to probe protein binding sites and discover new chemical scaffolds. For example, the discovery that the amide in imatinib (**4**) could be substituted with the 1,2,3-triazole^[9] led to further studies demonstrating that the Abl protein could assemble triazole inhibitor **5** from click fragments **6** and **7** in situ, making possible the use of the isCC strategy for the discovery of new classes of Abl inhibitors.^[10] Finally, in addition to enabling rapid analog exploration using CuAAC or isCC, substitution with the 1,2,3-triazole has also been demonstrated to improve physiochemical properties such as aqueous solubility^[13] and metabolic stability.^[14]

Due to the widespread success of the 1,2,3-triazole as an amide bioisostere, we sought to evaluate the utility of this bioisosteric replacement within amide-containing small molecule modulators of the cystic fibrosis transmembrane conductance regulator (CFTR) protein. CFTR is a member of the ATP-binding cassette (ABC) transporter superfamily and is comprised of two membrane spanning domains (MSD1, MSD2), two nucleotide binding domains (NBD1, NBD2), and a regulatory domain that controls protein function via PKA-dependent phosphorylation.^[16] CFTR functions as a Cl^- and HCO_3^- ion channel and plays an essential role in fluid homeostasis across epithelial cells in the lungs, pancreas, intestines, reproductive tract, and sweat glands.^[17] Mutations to the CFTR ion channel are the causative factor for the devastating disease Cystic Fibrosis (CF).^[18,19]

While ~2,000 CFTR mutations have been identified,^[20] deletion of phenylalanine 508 (F508del) from NBD1 is the most prominent CFTR mutation, with ~90% of CF patients carrying at least one F508del-CFTR allele.^[21] The F508del mutation has especially severe consequences for protein stability and function. First, deletion of phenylalanine 508 destabilizes the NBD1 domain and disrupts key domain-domain interactions between NBD1 and intracellular loop 4 (ICL4) of MSD2.^[22] This combination of structural defects impairs F508del-CFTR folding, leading to significant ubiquitin-proteasomal degradation and low levels of F508del-CFTR expression in the apical membrane of epithelial cells.^[23] Exacerbating the problem, F508del-CFTR also exhibits a decreased half-life at the plasma membrane^[24] and a gating defect in which Cl^- transport is impaired due to a decreased open probability (P_o) of the channel.^[24–26] These combined deficiencies result in CFTR mediated Cl^- secretion in human bronchial epithelial (HBE) cells derived from patients homozygous for F508del-CFTR that is only ~3–4% of that observed in non-CF epithelia.^[24,27] Diminished Cl^- transport in CF patients results in dehydration of the airway surfaces, which concurrently leads to decreases in ciliary beat frequency and mucociliary clearance causing the buildup of thick mucus within the lungs.^[28] This mucus harbors harmful bacterial pathogens, leading to chronic lung infection and hyperinflammation that severely damage the lungs and ultimately causes lung failure, the primary cause of death in CF patients.

Groundbreaking work by Vertex Pharmaceuticals and others has identified small molecule modulation of mutant forms of CFTR as a revolutionary strategy to combat CF.^[16,18,19,29] CFTR modulators are grouped into two categories—potentiators and correctors. Potentiators increase Cl^- ion transport by counteracting the gating defect in the mutant protein while correctors counteract folding deficiencies to increase expression of the protein. Modulators

in both classes have gained FDA approval.^[19, 29] Potentiator VX-770 (Ivacaftor, Figure 3) has been approved for a number of gating mutations, and the combination of VX-770 with VX-809 (Lumacaftor, Figure 3) is approved to treat the F508del mutation that disrupts protein folding and channel gating. Recently, the combination of Ivacaftor with the second-generation corrector VX-661 (Tezacaftor, Figure 3), which exhibits an improved safety profile over VX-809, was also approved to treat patients with the F508del mutation.

Despite the recent breakthroughs in CF therapy, the search for improved CFTR modulators remains a priority.^[19] Given the presence of the amide linkage in both VX-809 and VX-770, the detailed pharmacological characterization of these modulators reported in the primary literature,^[24,27] and the promise of the triazole as an amide bioisostere, we investigated the suitability of the 1,2,3-triazole bioisosteric replacement within the VX-809 and VX-770 structural classes. Importantly, demonstration that the triazole substitution is tolerated in these molecular scaffolds would provide a starting point for future studies aimed at generating improved classes of modulators using CuAAC or isCC strategies. Herein, we report the synthesis and biological evaluation of 1,2,3-triazole analogs of CFTR modulators within the VX-809 and VX-770 structural classes.

Results and Discussion

Design of 1,2,3-Triazole Analogs in the VX-809 Structural Class

Among F508del-CFTR correctors, the benzo[d][1,3]dioxole cyclopropane carboxamide moiety has been maintained as an essential structural feature across otherwise diverse chemical scaffolds, suggesting that this moiety is a key pharmacophore for corrector efficacy. As seen in the comparison of VX-809, VX-661, corrector **8** reported by AbbVie Pharmaceuticals (Figure 4A),^[30] and the widely employed tool compound C18 (VRT-534, Figure 4A),^[31,32] considerable structural variability is tolerated within the amide nitrogen substituent provided that the benzo[d][1,3]dioxole cyclopropane carboxamide moiety is present. On the basis of this observation, we hypothesized that CuAAC or isCC methods using 2,2-difluorobenzo[d][1,3]dioxole cyclopropane alkyne **9** (Figure 4B) have strong potential to develop a new class of correctors, if the 2,2-difluorobenzo[d][1,3]dioxole cyclopropane 1,2,3-triazole can serve as a suitable pharmacophore for F508del-CFTR correction. To investigate this question, we targeted the synthesis of 1,2,3-triazole containing corrector **11** shown in Figure 4B, which is a close structural analog of amide-containing VX-809.

In addition to the 1,2,3-triazole substitution, **11** differs from VX-809 in that it contains a central phenyl ring in contrast to the pyridine found in VX-809. This modification was made to prevent challenges associated with the use of 2-azidopyridines in azide-alkyne cycloaddition reactions due to the unfavorable equilibrium between 2-azidopyridines and the corresponding tetrazole tautomer (Figure 4C).^[33] In addition to alkyne **9** we also targeted the synthesis of alkyne **12** containing the 4-methoxyphenyl group in order to gain insight into the importance of the 2,2-difluorobenzo[d][1,3]dioxole moiety for corrector efficacy. The 4-methoxyphenyl moiety was selected from the VX-809 patent literature as it appeared to display efficacy in F508del-CFTR correction^[34] and was readily accessible synthetically.

We hypothesized that **12** might also be a useful building block for CuAAC or isCC medicinal chemistry efforts.

Synthesis of 1,2,3-Triazole Analogs in the VX-809 Structural Class

Alkynes **9** and **12** were prepared from the corresponding nitriles as shown in Scheme 1. Treatment of nitrile **13** with DIBALH at -78°C followed by quenching with 3 M HCl produced aldehyde **14** in 89% yield. Corey-Fuchs alkynylation using CBr_4 and Ph_3P , followed by treatment of the dibromoolefin intermediate with $n\text{BuLi}$ produced targeted alkyne **9** in 60% yield over two steps. Preparation of alkyne **12** from nitrile **15** was accomplished in the same manner, with the Corey-Fuchs reaction proving more efficient for this substrate as **12** was produced in 81% yield over two steps.

Synthesis of the azide **10** was also accomplished in a straightforward manner as shown in Scheme 2. Suzuki coupling of 3-bromo-4-methylaniline with boronic acid **18** produced biaryl **19** in 98% yield. TFA-mediated deprotection of *tert*-butyl ester **19**, followed by diazotization with sodium nitrite and substitution of the aryl diazonium intermediate with sodium azide, produced azide **10** in 92% yield over the two-step sequence. With the targeted alkyne and azide intermediates in hand we then performed the Cu-mediated azide-alkyne cycloaddition using standard reaction conditions to produce triazoles **11** and **20** in quantitative yield.

To enable the direct comparison of triazoles **11** and **20** to their analogous amides, amides **23** and **26** were prepared as shown in Scheme 3. HATU-mediated coupling of carboxylic acids **21** and **24** with aniline **19** produced the corresponding amides in excellent yield. Removal of the *tert*-butyl ester with TFA quantitatively produced amide analogs **23** and **26**.

Finally, in order to fully understand the effect of the replacement of the pyridyl ring in the VX-809 scaffold with the phenyl ring in amides **23/26** and 1,2,3-triazoles **11/20**, we prepared pyridyl-containing corrector **33** as shown in Scheme 4. Following the synthetic procedure reported for this class of correctors in the patent literature,^[35] Suzuki coupling of 2-bromo-3-methylpyridine with boronic acid **18** first produced **28** in high yield which was then quantitatively converted to N-oxide **29** via treatment with mCPBA. Amination via reaction with Ms_2O and pyridine in MeCN at 70°C for 1.5 h, followed by treatment with ethanolamine yielded 2-aminopyridine **30** in 53% yield. Preparation of acid chloride **31** from carboxylic acid **24** using SOCl_2 , and reaction with amine **30** produced amide **32** which contained the pyridyl core. Finally, hydrolysis of the *tert*-butyl ester using TFA afforded **33**, the 4-methoxyphenyl analog of VX-809, in 77% yield.

In vitro Evaluation of 1,2,3-Triazole Analogs in the VX-809 Structural Class

1,2,3-triazole-containing correctors **11** and **20**, amide-containing correctors **23** and **26** containing the central phenyl ring, and the amide-containing corrector **33** bearing the central pyridine were evaluated in transepithelial chloride conductance (TECC) assays using F508del CFBE41o⁻ monolayers. The F508del CFBE41o⁻ cell line is an immortalized cell line derived from bronchial epithelial cells of a CF patient homozygous for the F508del CFTR mutation with complementary stable F508del-CFTR transgene expression,^[36] and

has been shown to be clinically relevant in characterizing small molecule CFTR modulators. [37,38] Prior to TECC experiments, F508del CFBE41o⁻ monolayers were incubated (24 h) with varying concentrations of corrector compound to increase functional F508del-CFTR present at the apical membrane as a result of corrector-assisted protein folding. Increased CFTR-mediated Cl⁻ ion passage (due to increased F508del-CFTR protein) was characterized via measurement of changes in equivalent current (I_{eq}). Since CFTR-mediated Cl⁻ ion passage and therefore I_{eq} is slightly variable between cell cultures, a 3 μ M dose of VX-809 was used as a positive control in all experiments. Data are presented as normalized area under the curve values following stimulation with the cAMP agonist forskolin (10 μ M) and VX-770 (1 μ M), followed by inhibition with the CFTR inhibitor CFTR_{Inh}-172 (20 μ M).

As shown in Table 1 and Figure 5, several trends were observed regarding the importance of the 2,2-difluorobenzo[d][1,3]dioxole moiety for corrector efficacy, the impact of substituting the central pyridyl ring with the phenyl ring, and the suitability of the 1,2,3-triazole as an amide bioisostere in the VX-809 chemical series. First, the 4-methoxyphenyl moiety in amide **33** displayed greatly reduced corrector efficacy, highlighting the importance of the 2,2-difluorobenzo[d][1,3]dioxole functionality for F508del-CFTR correction. Secondly, substitution of the pyridyl ring in VX-809 with the phenyl ring in corrector **23** resulted in minimal loss of potency and efficacy, as **23** displayed similar dose response activity to VX-809. The minimal impact of the phenyl substitution on corrector efficacy is also seen in comparison of the 4-methoxyphenyl amide-containing analogs (pyridyl-containing **33** and phenyl-containing **26**).

Importantly, as shown in Table 1 and Figure 5, triazole-containing corrector **11** displayed statistically significant correction of F508del-CFTR above the DMSO response, with an EC₅₀ = 0.660 ± 0.287 μ M and a maximal efficacy of 48% compared to the 3 μ M VX-809 response. This demonstrates that the 2,2-difluorobenzo[d][1,3]dioxole cyclopropane 1,2,3-triazole can serve as a pharmacophore for F508del-CFTR correction, albeit with significantly reduced efficacy in comparison to the analogous amide **23**. However, it is possible that optimization of the triazole N1 substituent via the CuAAC reaction of alkyne **9** with structurally diverse azides could result in correctors with improved efficacy.

In contrast, corrector **20** containing the 4-methoxyphenyl cyclopropane 1,2,3-triazole pharmacophore was inactive. As the directly analogous 4-methoxyphenyl amide-containing corrector **26** was significantly less efficacious than **23**, the inactivity of triazole **20** is consistent with the loss in efficacy observed upon substitution of the amide with the triazole in this chemical series. In total, these findings demonstrate that while the triazole can be used to replace the amide in the VX-809 structural class when the 2,2-difluorobenzo[d][1,3]dioxole moiety is retained, use of the triazole results in lowered efficacy and is not a suitable replacement for less efficacious amide-containing correctors.

Design of 1,2,3-Triazole Analogs in the VX-770 Structural Class

We next investigated use of the 1,2,3-triazole as an amide bioisostere in the VX-770 potentiator series. As shown in Figure 6, tautomer **34** of VX-770, in which the amide

carbonyl oxygen forms an intramolecular H-bond with the quinolinol hydroxyl group, has been proposed to be the bioactive structural isomer.^[39] Supporting this hypothesis, attempts to substitute the amide in the VX-770 series with ester, sulfonamide, or reverse amide linkers essentially abolished potentiator efficacy.^[39] We hypothesized that triazole containing tautomer **35** could form a similar intramolecular H-bonding interaction to stabilize the bioactive tautomer and would be a suitable linker replacement. Additionally, on the basis of the considerable structure activity relationship (SAR) data published for the amide nitrogen substituent in the VX-770 series, we also targeted triazole analogs of potentiators **36** and **37** reported by Hadida and coworkers during the discovery of VX-770.^[39] Amide **36** was selected as it was reported to be the most potent analog characterized in F508del-CFTR HBE cells, and **37** was chosen to explore the indole moiety that was also determined to be highly effective for F508del-CFTR potentiation.^[39]

Synthesis of 1,2,3-Triazole Analogs in the VX-770 Structural Class

In contrast to the straightforward synthesis of 1,2,3-triazole analogs within the VX-809 structural class, the synthesis of triazole derivatives in the VX-770 chemical series proved more challenging. We first focused on the synthesis of 3-ethynylquinolin-4(1*H*)-one **43** that would serve as a key CuAAC precursor as shown in Scheme 5. Iodination of quinolin-4(1*H*)-one with I₂ and Na₂CO₃ allowed facile synthesis of 3-iodoquinolin-4(1*H*)-one **39**^[40] which was subjected to Sonogoshira reaction conditions as reported by Corelli and coworkers.^[41] Disappointingly while Corelli and coworkers were able to use microwave irradiation for similar substrates to achieve Sonogoshira coupling without conversion to furo[3,2-*c*]quinoline derivatives such as **41**,^[42] reaction of **39** with thermal heating (reflux) under Corelli's reaction conditions afforded undesired furo[3,2-*c*]quinoline **41**, which was unambiguously identified by conversion to known compound **42**.^[43] The cyclization likely proceeds through the mechanism shown in Scheme 5 involving coordination of the acetylene by copper and deprotonation of the acidic proton on the quinolinone nitrogen atom.^[42] This suggests that use of **43** would lead to a similar cyclization reaction under the copper-catalyzed azide-alkyne cycloaddition reaction conditions (see Scheme 5), and this was found to be the case (see Supporting Information).

As a methyl substituent on the quinolin-4(1*H*)-one nitrogen prevents the cyclization reaction during the Sonogoshira coupling,^[42] we began to investigate suitable nitrogen protecting groups. As shown in Supplementary Schemes 1 and 2, use of the methyl and *tert*-butyl carbamate protecting groups proved ineffective (see Supporting Information). In contrast, use of the benzyl protecting group, which has been successfully used in the synthesis of VX-770,^[44] was effective. Benzyl protection of **39** via deprotonation with NaH and reaction with benzyl bromide produced **44**.^[45] Sonogoshira coupling of **44** with trimethylsilyl acetylene, and TMS deprotection using K₂CO₃ in MeOH produced terminal alkyne **45** in 57% total yield over the three step sequence as shown in Scheme 6. Pleasingly, use of **45** in the CuAAC reaction with test substrate 4-*tert*-butylphenyl azide produced the desired click product **46** in moderate yield (66%), and the benzyl group could be removed by refluxing **46** with ammonium formate and Pearlman's catalyst in DMF. Following these test reactions, the synthesis of **45** was readily scaled up to produce gram quantities of **45**.

In parallel with the synthesis of alkyne **45**, we also pursued the synthesis of azides **48**, **50**, and **53** as shown in Scheme 7. Precursor amines **47**, **49**, and **51** were prepared according to the synthetic procedures reported by Hadida and coworkers (see Supporting Information, Supplementary Scheme 3).^[39] The synthesis of azides **48** and **50** was readily accomplished in good yield by treatment of the precursor amine with NaNO₂ followed by NaN₃. Surprisingly, use of these reaction conditions with aniline **51** could not produce the desired azide, as vigorous gas evolution was observed after the addition of NaNO₂ to **51**. Fortunately, use of Zhang and Moses' method for azide formation using ^tBuONO and TMSN₃ in MeCN^[46] proved effective, although significant formation of trimethylsilyl protected phenol **52** was produced (73%) in addition to isolation of the desired azide **53** (20%). This did not prove to be problematic as the TMS group could be readily removed by treatment with TBAF to produce **53** in quantitative yield. Notably aniline **51** is not a bench stable solid, and must be transformed to the azide immediately after preparation to prevent decomposition to quinone **54**.^[47]

With azides **48**, **50**, and **53** and alkyne **45** in hand we completed the synthesis of the 1,2,3-triazole containing potentiators as shown in Scheme 8. While CuAAC could be achieved with azides **48** and **50** at rt, no product formation was observed for reaction of alkyne **45** with sterically hindered azide **53** at rt. Fortunately, heating to 60 °C for 24 h produced the desired product **60** in modest yield (45%), but attempts to improve yield by extending the reaction time (up to 3 days) were unsuccessful. Finally, removal of the benzyl protection group using ammonium formate and Pearlman's catalyst in refluxing DMF afforded the desired 1,2,3-triazole analogs in good yield. Additionally, in order to directly compare the potentiator ability of the triazole derivatives to their parent amides **36** and **37**, these compounds were also prepared. As shown in Scheme 9, HATU-mediated amide bond formation afforded the desired compounds in improved yields over the published reaction procedures.^[39]

In vitro Evaluation of 1,2,3-Triazole Analogs in the VX-770 Structural Class

1,2,3-triazole containing potentiators **56**, **58**, and **60** were evaluated for potentiator activity and compared to amide potentiators **36**, **37**, and VX-770 in Ussing Chamber experiments. VX-770 is known to potentiate WT-CFTR (that does not exhibit a gating defect) and F508del-CFTR and G551D-CFTR (that do display the gating defect) with varying levels of potency and efficacy.^[27,48] Accordingly, we evaluated our amide and triazole-containing potentiators in CFBE41o⁻ monolayers with complementary WT CFTR expression (WT CFBE41o⁻), F508del CFBE41o⁻ monolayers after 24 h treatment with VX-809 (3 μM) to rescue mutant F508del-CFTR to the cell surface, and CFBE41o⁻ monolayers with complementary G551D-CFTR expression (G551D CFBE41o⁻).^[49] In addition, we also utilized Fischer Rat Thyroid (FRT) cells expressing G551D as this cell line is commonly used to characterize potentiator compounds and exhibits elevated Cl⁻ ion conductance compared to CFBE cells.^[50] We reasoned the higher Cl⁻ ion conductance might allow the detection of weak potentiators and differentiate the response between potentiators to a greater extent than the CFBE41o⁻ cell line.

The effect of the potentiator compounds was quantified via measurement of changes in short circuit current (I_{sc}) due to increased CFTR-mediated Cl^- ion secretion. The response was measured as the cumulative I_{sc} observed with serial addition of each test agent after stimulation with forskolin, either 100 nM (the EC_{50}) in WT monolayers or 10 μ M in mutant (F508del, G551D) monolayers. As Cl^- ion transport and increases in I_{sc} can be variable between cell cultures, VX-770 was used as a positive control in all experiments and relative potentiator efficacy was determined by calculating the percentage of each analog's response against the VX-770 response as shown in Table 2 and Figure 7. Potency was quantified by determining EC_{50} values and is shown in Table 2.

Disappointingly, as shown in Table 2 and Figure 7, none of the 1,2,3-triazoles (**56**, **58**, **60**) achieved statistically significant increases in I_{sc} above the DMSO response in any of the CFTR variants or cell types tested. These findings demonstrate that use of the 1,2,3-triazole moiety significantly alters the VX-770 chemical scaffold such that potentiation is abolished in the in vitro cellular assay. Among the active amide-containing potentiators, VX-770 consistently displayed the highest efficacy by 1.5 to 3-fold (Table 2), a finding that to our knowledge has not previously been reported in the primary literature as Hadida and coworkers did not report a measurement of efficacy for **36** and **37**.^[39] Interestingly, **36** (which varies structurally from VX-770 only in the deletion of one *tert*-butyl group) did not produce statistically significant potentiation above the DMSO response except in the G551D CFBE41o⁻ cell line. In contrast, indole **37** displayed more robust efficacy across all cell lines, although it was less efficacious than VX-770. However, **37** did display improved potency over VX-770 in our experiments. In total, these results demonstrate the essential nature of the amide linker in the VX-770 series as substitution with the triazole resulted in inactivity in the cellular assay. These studies also highlight the important impact of the amide substituent on in vitro efficacy.

X-Ray Crystal Structures and Natural Bond Orbital Calculations for VX-770 and 1,2,3-Triazole Analog **60**

Following our disappointing findings that 1,2,3-triazole containing potentiators in the VX-770 structural class were inactive in the cellular Ussing Chamber assays, we verified that we correctly synthesized the targeted 1,4-substituted 1,2,3-triazoles by obtaining the crystal structure of triazole **60** as shown in Figure 8A. With the X-ray structure providing conclusive proof that we synthesized the targeted triazole analog, we also obtained the crystal structure of VX-770 as shown in Figure 8B. We then compared the two structures in order to gain insight into the structural differences in the solid-state (an overlay of the structures is shown in Figure 8C). In line with previous studies comparing the R1-R2 distance in amide and 1,2,3-triazole analogs,^[3] the C8-C12 distance across the triazole in **60** was determined to be 4.99 Å. In comparison the C7-C11 distance (i.e. the analogous R1-R2 distance) across the amide in VX-770 was 3.76 Å, representing a 1.23 Å lengthening of the linker between the quinolin-4(1*H*)-one core and the phenol substituent upon replacement of the amide with the triazole. Additionally, the orientation angle between the quinolin-4(1*H*)-one and phenol substituents as defined in Figure 8D was increased upon substitution of the amide (143.4°) with the triazole (161.5°). We wondered if the increased linker length and/or

the slight change in orientation angle could be responsible for an unfavorable steric interaction between potentiator **60** and the binding pocket.

More relevant than comparison of the solid-state structures was a structural comparison of the proposed bioactive tautomeric isomer and quantification of whether the triazole nitrogen atom could participate in H-bonding with the quinolinol hydroxyl to stabilize the tautomer as we had hypothesized (see Figure 6). To this end, we first performed a computational geometry optimization on the solid-state crystal structures of VX-770 and **60** at the M06–2X/6–31 + G(d) level of theory using the experimentally-derived X-ray structures as the input geometries. We then modeled the quinolinol tautomer for both molecules by tautomerizing the CO bond and rotating the amide or triazole moiety so that it could participate in an intramolecular H-bonding interaction (as shown in Figure 9). These geometries were then optimized computationally. Additionally, Natural Bond Orbital (NBO) analysis was performed on both of the resulting optimized geometries to quantify the degree of intramolecular H-bonding that stabilizes the tautomeric form. Stabilization afforded to the molecule by hydrogen bonding of the quinolinol O–H bond with the triazole nitrogen atom's lone pair was estimated to be 27.02 kcal/mol (as determined from 2nd order perturbation theory within the NBO analysis). This confirmed that a substantial intramolecular H-bonding interaction can exist with the triazole ring as a donor as shown in Figure 9B which displays the relevant orbital interactions of the H-bond as provided by NBO analysis. In comparison, stabilization afforded by hydrogen bonding of the quinolinol O–H bond with the amide group's carbonyl oxygen lone pairs was estimated to be 38.30 kcal/mol (see Figure 9A and Supporting Information). This analysis confirmed that the 1,2,3-triazole ring in **60** can participate in a H-bonding interaction with the quinolinol hydroxyl group to stabilize the bioactive tautomer similarly to the amide carbonyl within VX-770.

We also quantified the difference in energies (including zero-point energy corrections) between the optimized non-tautomeric and tautomeric conformations. This calculation allows comparison of the energetic cost of assuming the bioactive conformation. Interestingly, assuming bioactive conformation **34** from the optimized geometry of VX-770 requires 8.4 kcal/mol in energy, while the energetic cost to access **35** from **60** was significantly less at 3.8 kcal/mol. This analysis suggests that **60** is not energetically hindered from achieving the bioactive tautomeric form, as evidenced by the lower energetic cost required to assume the quinolinol tautomeric form in comparison to VX-770.

The lower energetic barrier in the conversion of **60** to **35** is likely due to the fact that an intramolecular H-bond already exists between the amide and quinolinone carbonyl in VX-770 (red dotted line in Figure 8B). Converting to tautomeric conformation **34**, therefore, involves trading one hydrogen bond in VX-770 (20.46 kcal/mol) for another, stronger hydrogen bond in the tautomeric form (38.30 kcal/mol). The net stabilization of 17.84 kcal/mol from the new hydrogen bonding interaction partially offsets the energetic cost of assuming the tautomeric form. In contrast, no H-bonding interaction is present in the non-tautomeric form of triazole analog **60** (Figure 8A) and so the energy required to populate tautomeric form **35** from **60** is offset to a greater degree by the formation of the new H-bond interaction in **35** for a net stabilization of 27.02 kcal/mol.

Finally, we quantified the R1-R2 distance between the amide and triazole substituents and angle of orientation between the R1 and R2 substituents in the energy optimized tautomeric forms as we did for the solid state crystal structures. Interestingly, the C8-C12 distance across the triazole in **35** was determined to be 4.99 Å and the C7-C11 distance across the amide in **34** was 3.78 Å, both of which were very similar to the solid state measurements. The orientation angles were also comparable between the tautomeric forms and the solid state as the orientation angle (as defined in Figure 8D) of the amide was found to be 145.6° and the orientation angle of the triazole was determined to be 163.7°.

In summary, the NBO calculations suggest that triazole **60** should be able to assume the bioactive conformation **35**. Comparison of the energy optimized structures and X-ray crystal structures highlight the lengthened R1-R2 distance and increased angle of orientation that would position the triazole substituents in a slightly different position in the CFTR binding pocket relative to the amide substituents.

Patch Clamp Analysis of VX-770 and 1,2,3-Triazole Analog **60**

Given the computational evidence that the 1,2,3-triazole isomers should be able to assume the bioactive tautomeric conformation, we investigated the potentiator ability of **60** using patch clamp analysis in order to study the interaction of our triazole potentiators directly with the CFTR protein. We hypothesized that this experiment would eliminate potential cellular factors that may be responsible for the inactivity of **60** in vitro and provide insight into whether structural changes arising from incorporation of the 1,2,3-triazole prevent interaction of the potentiator with the binding site of CFTR. Macroscopic currents for WT-CFTR were recorded in the excised inside-out patch as shown in Figure 10. Interestingly, we observed that triazole-containing potentiator **60** robustly stimulated WT-CFTR currents in a dose dependent manner. The potency and efficacy of triazole **60** was similar to VX-770, as **60** exhibited an EC₅₀ of 5.5 ± 0.9 nM in comparison to an EC₅₀ of 1.3 ± 0.3 nM for VX-770, and **60** increased the percent of the PKI control to a slightly greater extent than VX-770. These findings demonstrate that under the cell-free condition, the 1,2,3-triazole substitution within the VX-770 chemical series is tolerated and that triazole-containing analogs can directly interact with CFTR and potentiate the channel activity. Moreover, they raise the question as to the cause of inactivity in the cellular assay.

Comparison of the Metabolic Stability of 1,2,3-Triazole and Amide-Containing CFTR Modulators

Having uncovered some drawbacks in the use of the 1,2,3-triazole as an amide bioisostere in the VX-809 and VX-770 structural classes, we investigated whether the incorporation of the triazole would afford beneficial pharmacokinetic properties as has been demonstrated in some instances^[14] and is widely perceived to be the case.^[2–5] To this end, the metabolic stability of five triazoles and the analogous amides was evaluated in human hepatic microsomes.

Intrinsic clearance (CL_{INT}) and calculated human hepatic clearance (CL_{HEP}) are shown in Table 3 (triazole and amide pairs are listed side-by-side). Surprisingly, in all but one instance (**20** vs. **26**) the triazole derivatives showed reduced microsomal stability (i.e., increased

CL_{INT}) in comparison to the analogous amide. In particular, triazole corrector **11** displayed substantially increased CL_{INT} in comparison to amide **23**, and all triazole potentiators (**56**, **58**, **60**) showed a modest increase in microsomal clearance. In contrast to the common perception that the triazole affords increased metabolic stability (due to elimination of amide hydrolysis), the decreased stability of the triazole analogs studied here demonstrates that the triazole may also have a negative effect on hepatic clearance in certain applications. This impact on hepatic clearance must also be taken into account when considering the overall impact of the triazole substitution on metabolic stability.

Conclusions

In conclusion, despite the widespread success of the 1,2,3-triazole as an amide bioisostere, our results demonstrate that use of the triazole is not effective in all medicinal chemistry contexts and highlight several negative impacts that can arise from this substitution. First, in the context of the VX-809 chemical series, while the 2,2-difluorobenzo[d][1,3]-dioxole cyclopropane 1,2,3-triazole was found to be a suitable pharmacophore for F508del-CFTR correction, the triazole derivative displayed significantly reduced corrector efficacy as **11** increased Cl⁻ ion transport in F508del CFBE41o⁻ cells to only 48% of the 3 μM VX-809 response. Additionally, triazole corrector **20**, containing the 4-methoxyphenyl group, was inactive in contrast to the analogous amide **26** which displayed modest F508del-CFTR correction (54% of the 3 μM VX-809 response). These findings highlight that the triazole, while tolerated in the VX-809 chemical series when the 2,2-difluorobenzo[d][1,3]-dioxole cyclopropane moiety is maintained, is not widely applicable across less efficacious correctors in this series.

Secondly, in the VX-770 chemical series, the use of the triazole substitution completely abolished potentiator activity in the cellular assays. These findings point to a significant structural change to the VX-770 scaffold resulting from the triazole substitution. On the basis of Natural Bond Orbital calculations performed on triazole **60**, this structural change is not likely to be disruption of the proposed bioactive tautomeric form as the triazole was found to be able to participate in the necessary intramolecular H-bonding interaction that stabilizes the tautomer. Furthermore, quantification of the difference in energy between the optimized non-tautomeric and tautomeric conformations provided evidence that **60** is not energetically hindered from achieving the bioactive conformation. The conclusions drawn from these calculations were supported by the similar potency and efficacy of **60** in comparison to VX-770 in the cell-free patch clamp experiment. The patch clamp results suggest that the inactivity of triazole potentiators **56**, **58**, and **60** in the cellular assay could be due to the inability of these compounds to access the CFTR binding site under cellular conditions, either due to poor membrane permeability or active transport out of the cell. These findings highlight the considerable impact the triazole substitution can have on crucial physiochemical properties that impact target engagement.

Thirdly, comparison of the solid-state structures of **60** and VX-770 and comparison of the energy optimized tautomeric forms **34** and **35** highlight the structural differences resulting from the triazole substitution. The distance across the amide substituents increases by ~1.2 Å in the triazole derivative, and the angle of orientation between the amide substituents is

increased by $\sim 18.1^\circ$ in the triazole. While these structural changes can be tolerated in many instances without negative consequences on biological activity and/or physiochemical properties,^{2–5} this did not prove to be the case in the VX-809 and VX-770 chemical series.

Finally, despite the widespread perception that use of the 1,2,3-triazole can provide increased metabolic stability, in two distinct chemical series 1,2,3-triazole derivatives largely displayed similar or increased metabolic clearance in human hepatic microsomal assays. In the case of triazole **11** ($CL_{INT} = 11.8$) the increase in intrinsic clearance was substantial ($\sim 5x$) in comparison to the analogous amide **23** ($CL_{INT} = 2.49$). In total, these findings represent an important contrast to the many successful uses of the 1,2,3-triazole as an amide bioisostere in medicinal chemistry.

Experimental Section

Complete experimental procedures and Supplementary Schemes and Figures are provided in the Supporting Information. Chemistry: Supplementary Schemes and Figures, synthetic experimental procedures, compound characterization data, copies of 1H and ^{13}C NMR spectra, X-ray crystallography data, and details on computational studies are provided. Biology: Experimental procedures for cell culture, TECC assay, Ussing Chamber recordings, patch clamp analysis, and intrinsic clearance assays are provided.

CCDC data 1879006 (VX-770) and 1879007 (**60**) contain the supplementary crystallographic data for this paper. These data are provided free of charge by The Cambridge Crystallographic Data Centre.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

The authors gratefully acknowledge the following sources of funding: J.E.D. thanks the Richards Scholar Program for financial support. S.G.A. acknowledges funding by the Cystic Fibrosis Foundation (ALLER16P0 and ALLER16G0). S.M.R. acknowledges funding from the NIH (P30 DK072482 and R35 HL135816) and the Cystic Fibrosis Foundation (R464-CR11). The authors gratefully thank Prof. Craig Lindsley of Vanderbilt University for access to microsomal stability analysis, compound purity analysis, and HRMS analysis. The authors thank Dr. John Bacsa at the X-ray Crystallography Center at Emory University for X-ray data collection.

References

- [1]. Hein JE, Fokin VV, Chem. Soc. Rev 2010, 39, 1302–1315. [PubMed: 20309487]
- [2]. Bonandi E, Christodoulou MS, Fumagalli G, Perdicchia D, Rastelli G, Passarella D, Drug Discov. Today 2017, 22, 1572–1581. [PubMed: 28676407]
- [3]. Hou J, Liu X, Shen J, Zhao G, Wang PG, Expert Opin. Drug Discov 2012, 7, 489–501. [PubMed: 22607210]
- [4]. Tron GC, Pirali T, Billington RA, Canonico PL, Sorba G, Genazzani AA, Med. Res. Rev 2008, 28, 278–308. [PubMed: 17763363]
- [5]. Kolb HC, Sharpless KB, Drug Discov. Today 2003, 8, 1128–1137. [PubMed: 14678739]
- [6]. Chrysina ED, Bokor É, Alexacou K-M, Charavgi M-D, Oikonomakos GN, Zographos SE, Leonidas DD, Oikonomakos NG, Somsák L, Tetrahedron Asymmetry 2009, 20, 733–740.

- [7]. Horne WS, Yadav MK, Stout CD, Ghadiri MR, J. Am. Chem. Soc 2004, 126, 15366–15367. [PubMed: 15563148]
- [8]. Mohammed I, Kummetha IR, Singh G, Sharova N, Lichinchi G, Dang J, Stevenson M, Rana TM, J. Med. Chem 2016, 59, 7677–7682. [PubMed: 27509004]
- [9]. Arioli F, Borrelli S, Colombo F, Falchi F, Filippi I, Crespan E, Naldini A, Scalia G, Silvani A, Maga G, Carraro F, Botta M, Passarella D, ChemMedChem 2011, 6, 2009–2018. [PubMed: 21990039]
- [10]. Peruzzotti C, Borrelli S, Ventura M, Pantano R, Fumagalli G, Christodoulou MS, Monticelli D, Luzzani M, Fallacara AL, Tintori C, Botta M, Passarella D, ACS Med. Chem. Lett 2013, 4, 274–277. [PubMed: 24900659]
- [11]. Christodoulou MS, Mori M, Pantano R, Alfonsi R, Infante P, Botta M, Damia G, Ricci F, Sotiropoulou PA, Liekens S, Botta B, Passarella D, ChemPlusChem 2015, 80, 938–943.
- [12]. Ballard TE, Richards JJ, Wolfe AL, Melander C, Chem. – Eur. J 2008, 14, 10745–10761. [PubMed: 18942682]
- [13]. Mugnaini C, Nocerino S, Pedani V, Pasquini S, Tafi A, De Chiaro M, Bellucci L, Valoti M, Guida F, Luongo L, Dragoni S, Ligresti A, Rosenberg A, Bolognini D, Cascio MG, Pertwee RG, Moaddel R, Maione S, Di Marzo V, Corelli F, ChemMedChem 2012, 7, 920–934. [PubMed: 22383251]
- [14]. Keck TM, Banala AK, Slack RD, Burzynski C, Bonifazi A, Okunola-Bakare OM, Moore M, Deschamps JR, Rais R, Slusher BS, Newman AH, Bioorg. Med. Chem 2015, 23, 4000–4012. [PubMed: 25650314]
- [15]. Mamidyalala SK, Finn MG, Chem. Soc. Rev 2010, 39, 1252. [PubMed: 20309485]
- [16]. Hwang T-C, Yeh J-T, Zhang J, Yu Y-C, Yeh H-I, Destefano S, J. Gen. Physiol 2018, jgp. 201711946.
- [17]. Sheppard DN, Welsh MJ, Physiol. Rev 1999, 79, S23–S45. [PubMed: 9922375]
- [18]. Sloane PA, Rowe SM, Curr. Opin. Pulm. Med 2010, 16, 591–597. [PubMed: 20829696]
- [19]. Jennings MT, Flume PA, Ann. Am. Thorac. Soc 2018, 15, 897–902. [PubMed: 29812963]
- [20]. Sosnay PR, Raraigh KS, Gibson RL, Pediatr. Clin. North Am 2016, 63, 585–598. [PubMed: 27469177]
- [21]. Bobadilla JL, Macek M Jr., Fine JP, Farrell PM, Hum. Mutat 2002, 19, 575–606. [PubMed: 12007216]
- [22]. Serohijos AWR, Hegedus T, Aleksandrov AA, He L, Cui L, Dokholyan NV, Riordan JR, Proc. Natl. Acad. Sci 2008, 105, 3256–3261. [PubMed: 18305154]
- [23]. Penque D, Mendes F, Beck S, Farinha C, Pacheco P, Nogueira P, Lavinha J, Malhó R, Amaral MD, Lab. Invest 2000, 80, 857–868. [PubMed: 10879737]
- [24]. Van Goor F, Hadida S, Grootenhuis PDJ, Burton B, Stack JH, Straley KS, Decker CJ, Miller M, McCartney J, Olson ER, Wine JJ, Frizzell RA, Ashlock M, Negulescu PA, Proc. Natl. Acad. Sci 2011, 108, 18843–18848. [PubMed: 21976485]
- [25]. Wang W, Li G, Clancy JP, Kirk KL, J. Biol. Chem 2005, 280, 23622–23630. [PubMed: 15857825]
- [26]. Miki H, Zhou Z, Li M, Hwang T-C, Bompadre SG, J. Biol. Chem 2010, 285, 19967–19975. [PubMed: 20406820]
- [27]. Goor FV, Hadida S, Grootenhuis PDJ, Burton B, Cao D, Neuberger T, Turnbull A, Singh A, Joubran J, Hazlewood A, Zhou J, McCartney J, Arumugam V, Decker C, Yang J, Young C, Olson ER, Wine JJ, Frizzell RA, Ashlock M, Negulescu P, Proc. Natl. Acad. Sci 2009, 106, 18825–18830. [PubMed: 19846789]
- [28]. Matsui H, Grubb BR, Tarran R, Randell SH, Gatzky JT, Davis CW, Boucher RC, Cell 1998, 95, 1005–1015. [PubMed: 9875854]
- [29]. Kym PR, Wang X, Pizzonero M, Van der Plas SE, in Prog. Med. Chem (Eds.: Witty DR, Cox B), Elsevier, 2018, pp. 235–276.
- [30]. Wang X, Liu B, Searle X, Yeung C, Bogdan A, Greszler S, Singh A, Fan Y, Swensen AM, Vortherms T, Balut C, Jia Y, Desino K, Gao W, Yong H, Tse C, Kym P, J. Med. Chem 2018, 61, 1436–1449. [PubMed: 29251932]

- [31]. Okiyoneda T, Veit G, Dekkers JF, Bagdany M, Soya N, Xu H, Roldan A, Verkman AS, Kurth M, Simon A, Hegedus T, Beekman JM, Lukacs GL, Nat. Chem. Biol 2013, 9, 444–454. [PubMed: 23666117]
- [32]. Eckford PD, Ramjeesingh M, Molinski S, Pasyk S, Dekkers JF, Li C, Ahmadi S, Ip W, Chung TE, Du K, Yeger H, Beekman J, Gonska T, Bear CE, Chem. Biol 2014, 21, 666–678. [PubMed: 24726831]
- [33]. Bolje A, Urankar D, Košmrlj J, Eur. J. Org. Chem 2014, 8167–8181.
- [34]. Hadida R, Grootenhuis PDJ, Zhou J, Bear B, Miller M, McCartney J, Pyridine Compounds as Modulators of CFTR and Their Preparation and Use in the Treatment of Diseases WO 2008/141119, 11 20, 2008.
- [35]. Siesel D, Processes for Producing Cycloalkylcarboxamido-Pyridine Benzoic Acids US 8,124,781 B2, 2 28, 2012.
- [36]. Goncz KK, Kunzelmann K, Xu Z, Gruenert DC, Hum. Mol. Genet 1998, 7, 1913–1919. [PubMed: 9811935]
- [37]. Ehrhardt C, Collnot E-M, Baldes C, Becker U, Laue M, Kim K-J, Lehr C-M, Cell Tissue Res 2006, 323, 405–415. [PubMed: 16249874]
- [38]. Phuan P-W, Veit G, Tan J, Roldan A, Finkbeiner WE, Lukacs GL, Verkman AS, Mol. Pharmacol 2014, 86, 42–51. [PubMed: 24737137]
- [39]. Hadida S, Van Goor F, Zhou J, Arumugam V, McCartney J, Hazlewood A, Decker C, Negulescu P, Grootenhuis PDJ, J. Med. Chem 2014, 57, 9776–9795. [PubMed: 25441013]
- [40]. Almeida A, Silva A, Cavaleiro J, Synlett 2010, 2010, 462–466.
- [41]. Mugnaini C, Falciani C, De Rosa M, Brizzi A, Pasquini S, Corelli F, Tetrahedron 2011, 67, 5776–5783.
- [42]. Venkataraman S, Barange DK, Pal M, Tetrahedron Lett 2006, 47, 7317–7322.
- [43]. Beydoun K, Doucet H, Eur. J. Org. Chem 2012, 2012, 6745–6751.
- [44]. Thatipally S, Reddy VK, Dammalapati VLNR, Chava S, Processes for the Preparation of Ivacaftor WO 2017/037672 A1, 3 9, 2017.
- [45]. Audisio D, Messaoudi S, Peyrat J-F, Brion J-D, Alami M, J. Org. Chem 2011, 76, 4995–5005. [PubMed: 21524121]
- [46]. Zhang F, Moses JE, Org. Lett 2009, 11, 1587–1590. [PubMed: 19267454]
- [47]. Thatipally S, Reddy VK, Dammalapati VLNR, Gorantla SRA, Chava S, An Improved Process for the Preparation of 5-Amino-2,4,-Di-Tert-Butylphenol or an Acid Additional Salt Thereof WO 2016/075703 A2, 5 19, 2016.
- [48]. Jih K-Y, Hwang T-C, Proc. Natl. Acad. Sci 2013, 110, 4404–4409. [PubMed: 23440202]
- [49]. Gottschalk LB, Vecchio-Pagan B, Sharma N, Han ST, Franca A, Wohler ES, Batista DAS, Goff LA, Cutting GR, J. Cyst. Fibros 2016, 15, 285–294. [PubMed: 26694805]
- [50]. Yu H, Burton B, Huang C-J, Worley J, Cao D, Johnson JP, Urrutia A, Joubran J, Seepersaud S, Sussky K, Hoffman BJ, Van Goor F, J. Cyst. Fibros 2012, 11, 237–245. [PubMed: 22293084]
- [51]. Dolomanov OV, Bourhis LJ, Gildea RJ, Howard JAK, Puschmann H, J. Appl. Crystallogr 2009, 42, 339–341.

Figure 1. Bioisosteric relationship of 1,4-disubstituted 1,2,3-triazoles and *trans*-amides. A) Similar R1-R2 distance and arrangement of H-bond donor and acceptor groups in the *trans*-amide and 1,2,3-triazole. Adapted from references [3] and [4]. B) Overlay of *trans*-N-methylacetamide and 1,4-dimethyl-1H-1,2,3-triazole in which the amide carbonyl carbon and nitrogen atom are aligned with C4 and C5 of the 1,2,3-triazole.

HIV viral infectivity factor (Vif) antagonist:⁸ **Breakpoint cluster region-Abelson kinase (Bcr-Abl) inhibitors:^{9,10}** **Figure 2.**

Use of the 1,2,3-triazole as an amide bioisostere allows use of CuAAC or isCC to optimize and/or discover new chemical scaffolds.

Figure 3.
Structures of potentiator VX-770 and correctors VX-809 and VX-661.

Figure 4.

A) Structure of corrector **8**^[30] and C18 (VRT-534)^[31,32] bearing the benzo[d][1,3]dioxole cyclopropane carboxamide pharmacophore. B) Targeted alkyne **9** and azide **10** to explore the viability of the 2,2-difluorobenzo[d][1,3]dioxole cyclopropane 1,2,3-triazole pharmacophore. C) 2-azidopyridine and tetrazole tautomers. D) Targeted alkyne **12** to investigate the importance of the 2,2-difluorobenzo[d][1,3]dioxole moiety for corrector efficacy.

Figure 5.

Dose response effects of triazole and amide-containing correctors on Cl^- conductance in F508del CFBE41o $^-$ monolayers using the TECC assay. Current was measured by calculating area under the curve (AUC) after stimulation with forskolin (10 μM) + VX-770 (1 μM), followed by inhibition with CFTR $_{\text{Inh}}$ -172 (20 μM). A 3 μM dose of VX-809 was used as a positive control in all experiments. Relative corrector efficacy was determined by calculating the percentage of the analog's response against the 3 μM VX-809 response. Asterisks indicate significant ($p < 0.05$, Two-way ANOVA, Tukey's post-hoc test) increases in AUC against the vehicle (DMSO) response, $n = 3 - 5$.

Figure 6. A) Proposed bioactive tautomer of VX-770^[39] and triazole containing analog. B) Structures of highly potent compounds within the VX-770 structural class.^[39]

Figure 7.

Dose response effects of triazole and amide-containing potentiators on Cl^- conductance in WT CFBE41o⁻, F508del CFBE41o⁻ following 24 h exposure to VX-809 (3 μM), G551D CFBE41o⁻, and G551D FRT cell lines. Current was measured after stimulation with forskolin (100 nM in WT and 10 μM in mutants F508del and G551D). Inhibition with CFTR_{Inh}-172 (10 μM) demonstrated that increased current was due to a CFTR-mediated mechanism. Relative potentiator efficacy was determined by calculating the percentage of the analog's response against the VX-770 response. Data are normalized according to vehicle (DMSO) response. Asterisks indicate significant ($p < 0.05$, Two-way ANOVA, Tukey's post-hoc test) increases in I_{sc} against the vehicle (DMSO) response, $n = 3 - 5$.

Figure 8.

A) Crystal structure of **60** showing representation of the thermal ellipsoids. B) Crystal structure of VX-770 showing representation of the thermal ellipsoids. H-bonding interaction is shown in red. C) Overlay of crystal structures of **60** and VX-770. D) Calculation of orientation angle between quinolin-4(1*H*)-one and phenol substituents using Olex2.^[51] For VX-770 lines were generated from C7-C10, C10-C11, and C7-C11 to define a triangle. The C7-C10-C11 angle was measured to be 143.4°. For **60** lines were generated from C8-C10, C10-C12, and C8-C12 and the C8-C10-C12 angle was measured to be 161.5°.

Figure 9.

Optimized geometries for tautomeric isomers **34** and **35** at the M06-2X/6-31 + G(d) level of theory. The resulting hydrogen bonds between the tautomeric OH bond and the lone-pair donor atom in each molecule are illustrated by the relevant orbital interactions as provided by NBO analysis. A second, weaker, lone pair donation from the CO bond of **34** to the OH bond (see Supporting Information) is not shown for the sake of clarity.

Figure 10.

Dose response analysis for VX-770 and **60** by patch clamp analysis. A/B) VX-770 and **60** titrations for representative patches. Protein kinase A inhibitor (PKI) was applied to prevent further phosphorylation. C) Mean titration data fit to $Y = B_{max} \cdot X / (K_1 + X)$, $EC_{50} = 1.3 \pm 0.3$ (n=5) for VX-770 and 5.5 ± 0.9 (n=6) for **60**. Data were normalized to the peak current at 100–200 nM. The results are reported as means $\pm$ S.E. The lowest dose tested was 0.2 nM, which increased the current by $\sim 12\%$ for VX-770 and $\sim 7\%$ for **60**. D) Mean data % of maximal stimulation (efficacy) by VX-770 and **60**.

Scheme 1.

Synthesis of Alkynes **9** and **12**. Reagents and conditions: a) (i) DIBALH, toluene, -78°C , 3 h; (ii) HCl, 0°C , 15 min, **14**: 89%, **16**: 88%; b) CBr_4 , PPh_3 , CH_2Cl_2 , 0°C , 1 h; c) nBuLi (2.2 equiv) THF, -78°C , 1.5 h, **9**: 60% over 2 steps, **12**: 81% over 2 steps.

Scheme 2.

Synthesis of Azide **10** and 1,2,3-Triazole-Containing Correctors **11** and **20**. Reagents and conditions: (a) $\text{Pd(dppf)Cl}_2 \cdot \text{CH}_2\text{Cl}_2$, K_2CO_3 , toluene/ H_2O (3:1), reflux, 20 h, 98%; (b) TFA, CH_2Cl_2 , 0 °C to rt, 2 h; (c) (i) EtOAc, HCl, 0 °C, 5 min; (ii) NaNO_2 , H_2O , 0 °C, 1 h; (iii) NaN_3 , H_2O , 0 °C to rt, 3 h; 92% over 2 steps; (d) **9** or **12**, $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, sodium ascorbate, CH_2Cl_2 / $\text{tBuOH}/\text{H}_2\text{O}$ (1:2:2), rt, 24 h, **11**: 100%, **20**: 100%.

Scheme 3.

Synthesis of Amide-Containing Correctors **23** and **26**. Reagents and conditions: (a) **19**, HATU, Et₃N, DMF, rt, 24 h, **22**: 99%, **25**: 96%; (b) TFA, CH₂Cl₂, 0 °C to rt, 2 h, **23**: 100%, **26**: 99%.

Scheme 4.

Synthesis of Corrector **33**. Reagents and conditions: (a) **18**, Pd(dppf)Cl₂•CH₂Cl₂, K₂CO₃, toluene/H₂O, reflux, 18 h, 92%; (b) mCPBA, CH₂Cl₂, rt, 24 h, 99%; (c) (i) Ms₂O, pyridine, MeCN, 70 °C, 1.5 h; (ii) ethanolamine, rt, 2 h, 53% (d) SOCl₂, toluene, 60 °C, 2 h. (e) *i*-Pr₂EtN, DMAP, CH₂Cl₂, rt, 18 h, 56%; (f) TFA, CH₂Cl₂, 0 °C to rt, 2 h, 77%.

Scheme 5.

Unsuccessful Synthesis of **43** and Incompatibility of **43** with CuAAC Due to Cyclization of 3-Ethynylquinolin-4(1H)-ones to Furo[3,2-c]quinolines. Reagents and conditions: (a) I_2 , Na_2CO_3 , THF, rt, 24 h, 82%; (b) trimethylsilylacetylene, $Pd(PPh_3)_2Cl_2$, CuI, DIPEA, THF, rt, 18 h, 64%; (c) K_2CO_3 , MeOH, rt, 24 h, 67%.

Scheme 6.

Synthesis of Benzyl Protected 3-ethynylquinolin-4(1*H*)-one **45** and Compatibility with CuAAC. Reagents and conditions: (a) (i) NaH, THF, 0 °C, 30 min; (ii) benzyl bromide, 0 °C to rt, 19 h, 77%; (b) trimethylsilylacetylene, Pd(PPh₃)₂Cl₂, CuI, DIPEA, THF, rt, 18 h, 86%; (c) K₂CO₃, MeOH, rt, 24 h, 86%. (d) 4-*tert*-butylphenyl azide, CuSO₄•5H₂O, sodium ascorbate, CH₂Cl₂/tBuOH/H₂O (1:2:2), rt, 24 h, 66%.

Scheme 7.

Synthesis of Azides **48**, **50**, and **53**. Reagents and conditions: (a) (i) EtOAc, HCl, 0 °C, 5 min; (ii) NaNO₂, H₂O, 0 °C, 30 min; (iii) NaN₃, H₂O, 0 °C to rt, 2 h, **48**: 81%; **50**: 75%; **53**: 0%; (b) TMSN₃, ^tBuONO, MeCN, rt, 18 h, **52**: 73%, **53**: 20%; (c) TBAF, THF, 0 °C to rt, 2 h, 100%.

Scheme 8.

Synthesis of 1,2,3-Triazole Containing Potentiators **56**, **58**, and **60**. Reagents and conditions:

(a) $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, sodium ascorbate, $\text{CH}_2\text{Cl}_2/\text{BuOH}/\text{H}_2\text{O}$ (1:2:2), rt, 24 h, **55**: 69%, **57**: 71%, **59**: 0% (b) $\text{Pd}(\text{OH})_2$, ammonium formate, DMF, 80 °C, **56**: 10 h, 74%, **58**: 8 h, 88%, **60**: 8.5 h, 87%; (c) $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, sodium ascorbate, $\text{CH}_2\text{Cl}_2/\text{BuOH}/\text{H}_2\text{O}$ (1:2:2), 60 °C, 24 h, 45%.

Scheme 9.

Synthesis of Amide Containing Potentiators **36** and **37**. Reagents and conditions: (a) **47** or **49**, HATU, Et₃N, CH₂Cl₂, 70 °C, 16 h, **36**: 87%; **37**: 64%.

Table 1.

In Vitro Potency and Efficacy Data of Amide and Triazole-Containing Correctors in the VX-809 Chemical Series in F508del CFBE41o-Cells.

compd	EC ₅₀ (μM) ^[a]	Maximal Response ^[b]
11	0.660 ± 0.287	48.1%
20	inactive	--
23	1.960 ± 0.754	120.1%
26	2.971 ± 0.900	54.1%
33	0.427 ± 0.514	49.9%

^[a]Mean and standard deviation values from at least three determinations are reported.

^[b]Maximal response is reported as % activity of the 3 μM VX-809 response.

In Vitro Potency and Efficacy Data of Amide and Triazole-Containing Potentiators in the VX-770 Chemical Series in CFBE41o⁻ and FRT Cells.^[a]

Table 2.

compd	EC ₅₀ (nM) ^[b]			Maximal Response ^[c]				
	WT CFBE	F508del CFBE	G551D CFBE	G551D FRT	WT CFBE	F508del CFBE	G551D CFBE	G551D FRT
VX-770	36.5 ± 21.3	1889 ± 1388	77.7 ± 27.2	428.5 ± 167.3	100.0%	100.0%	100.0%	100.0%
36	--	--	26.7 ± 79.4	--	Inactive ^d	Inactive ^d	63.5%	Inactive ^d
37	21.8 ± 41.2	37.1 ± 48.2	18.5 ± 23.4	81.45 ± 102.8	47.5%	36.3%	72.0%	36.7%
56	Inactive across all cell and CFTR types ^d							
58	Inactive across all cell and CFTR types ^d							
60	Inactive across all cell and CFTR types ^d							

^aCFBE41o⁻ abbreviated as CFBE.

^bMean and standard deviation values from at least three determinations are reported.

^cMaximal response is reported as % activity of the VX-770 response.

^dNo statistically significant response above DMSO vehicle.

Table 3.

Intrinsic Clearance (CL_{INT}) in Human Microsomes and Predicted Hepatic Clearance (CL_{HEP}) of Analogous Triazole and Amide-Containing CFTR Modulators.

1,2,3-triazole			amide		
cmpd	$CL_{INT}^{[a]}$ (mL min ⁻¹ kg ⁻¹)	$CL_{HEP}^{[b]}$ (mL min ⁻¹ kg ⁻¹)	cmpd	$CL_{INT}^{[a]}$ (mL min ⁻¹ kg ⁻¹)	$CL_{HEP}^{[b]}$ (mL min ⁻¹ kg ⁻¹)
11	11.8	7.55	23	2.49	2.23
20	23.2	11.0	26	36.1	13.3
56	16.3	9.19	36	12.5	7.84
58	19.3	10.1	37	9.35	6.47
60	56.2	15.3	VX-770	45.6	14.4

^[a] Intrinsic clearance in human hepatic microsomes.

^[b] Predicted hepatic clearance, human.