Published in final edited form as:

ACS Chem Biol. 2018 December 21; 13(12): 3269–3278. doi:10.1021/acscchembio.8b00584.

Identification and Molecular Characterization of PPAR δ as a Novel Target for Covalent Modification by 15-deoxy-^{12, 14}-prostaglandin J₂

Aravind T. Reddy^{1,2}, Sowmya P. Lakshmi^{1,2}, Asoka Banno¹, and Raju C. Reddy^{1,2,*}

¹Department of Medicine, Division of Pulmonary, Allergy and Critical Care Medicine, University of Pittsburgh School of Medicine, Pittsburgh, PA 15213

²Veterans Affairs Pittsburgh Healthcare System, Pittsburgh, PA 15240

Abstract

PPAR δ belongs to the peroxisome proliferator-activated receptor (PPAR) family of nuclear receptors. Upon activation by an agonist, PPAR δ controls a variety of physiological processes via regulation of its target genes. 15-deoxy-^{12,14}-PGJ₂ (15d-PGJ₂) is a cyclopentenone prostaglandin that features an electrophilic, α , β -unsaturated ketone (an enone) in the cyclopentenone ring. Many of 15d-PGJ₂'s biological effects result from covalent interaction between C₉ and the thiol group of a catalytic cysteine (Cys) in target proteins. In this study, we investigated whether 15d-PGJ₂ activates PPAR δ by forming a covalent adduct. Our data show that 15d-PGJ₂ activates PPAR δ 's transcriptional activity through formation of a covalent adduct between its endocyclic enone at C₉ and Cys249 in the receptor's ligand-binding domain. As expected, no adduct formation was seen following a Cys-to-Ser mutation at residue 249 (C249S) of PPAR δ or with a PGD₂/PGJ₂ analog that lacks the electrophilic C₉. Furthermore, the PPAR δ C249S mutation reduced induction of the receptor's DNA binding activity by 15d-PGJ₂, which highlights the biological significance of our findings. Calculated chemical properties as well as data from molecular orbital calculations, reactive molecular dynamics simulations, and intrinsic reaction coordinate modeling also supported the selectivity of 15d-PGJ₂'s C₉ towards PPAR δ 's Cys thiol. In summary, our results provide the molecular, chemical, and structural basis of 15d-PGJ₂-mediated PPAR δ activation, designating 15d-PGJ₂ as the first covalent PPAR δ ligand to be identified.

Graphical Abstract

* Address correspondence to Raju C. Reddy, M.D.; reddyrc@upmc.edu.

Supporting Information Available: This material is available free of charge via the internet at <http://pubs.acs.org>.

The Supporting Information file for the current manuscript contains Supplemental Tables 1 and 2 with their captions; Supplemental Figures S1–S5 with their captions; and Supplemental Videos MovieS1–MovieS4 with their captions.

CONFLICT OF INTEREST

The authors declare that they have no conflicts of interest with the contents of this article.

FOOTNOTES

The contents of this article are solely the responsibility of the authors and do not necessarily represent the official views of the National Institutes of Health, the U.S. Department of Veterans Affairs, or the United States Government.

Keywords

nuclear receptor; transcription factor; fatty acid; covalent bond; ligand binding; 1,4 addition; Cysalkylation; cyclopentenone ring; molecular dynamics; molecular modeling

The peroxisome proliferator-activated receptor (PPAR) family of nuclear receptors, comprising PPAR α , PPAR δ , and PPAR γ ,^{1,2} is found in many cells and tissues of the body.^{3–5} Upon agonist-induced activation, PPARs control physiological processes such as lipid and glucose metabolism and immune responses^{6,7} by modulating gene expression through binding to specific DNA sequences, termed PPAR response elements (PPREs), in the enhancer regions of their targets. PPARs' transcriptional activity also requires formation of an activator complex with retinoid X receptors and additional co-activator proteins.¹

The three PPARs share more than 60% sequence homology in their ligand-binding domains (LBDs).⁸ X-ray crystallography of PPAR δ shows a large Y-shaped ligand-binding cavity of approximately 1,300 Å³ that is composed of three sub-arms (arms I, II and III) and enclosed by six α -helices (H3, H4, H6, H8, H11, and H12) and two β -strands (S2 and S3).^{9,10} Known

physiological PPAR ligands include saturated and unsaturated fatty acids as well as eicosanoids. Among the well-characterized PPAR ligands is a group of physiologically active lipids known as prostaglandins (PGs).^{1,9,11,12}

PG biosynthesis occurs in most tissues; it starts with an arachidonic acid intermediate generated by phospholipase-A₂ from phospholipids and proceeds through the cyclooxygenase pathway, finally producing thromboxane, prostacyclin, and PGD₂, PGE₂, or PGF₂ depending on the specific PG synthase involved. PGD₂ then undergoes a chain of dehydration, yielding members of the cyclopentenone PG (cyPG) family, such as PGJ₂,¹² PGJ₂, and 15-deoxy-^{12,14}PGJ₂ (15d-PGJ₂), that feature an electrophilic α , β -unsaturated ketone group (an enone) in the cyclopentenone ring.^{13–15} The electrophilic β -carbon covalently interacts via Michael addition with a nucleophile, primarily the sulfur atom of thiols such as glutathione and the cysteine residues (Cys) of cellular proteins.^{14,16–18} Such conjugation is stereoselective, creating cis or trans products depending on the type of ring involved.^{16,18} 15d-PGJ₂'s covalent adduction to the thiol group of a catalytic Cys in various proteins has been shown to mediate its biological effects; some key studies are summarized in Table S1.

Forman *et al.* identified 15d-PGJ₂ as a natural PPAR γ ligand.¹⁹ Subsequently, Soares *et al.* reported that 15d-PGJ₂ formed a covalent adduct with Cys285 within the PPAR γ LBD.²⁰ Indeed, covalent interactions with its target proteins account for many of 15d-PGJ₂'s biological effects. Although 15d-PGJ₂ was found to also activate PPAR δ complexed with retinoid X receptor¹⁹, the molecular and chemical nature of such interaction has so far been elusive. Our data present the molecular, chemical, and structural characteristics of the 15d-PGJ₂-PPAR δ complex and reveal 15d-PGJ₂ as the first covalent PPAR δ agonist to be identified so far. The chemical and biological modeling systems used and the information gained in this study can also serve as a foundation of future studies investigating 15d-PGJ₂'s interactions with other target proteins.

RESULTS AND DISCUSSION

15d-PGJ₂ activates PPAR δ .

15d-PGJ₂ is one of the cyPGs whose functions in inflammation, cell proliferation, survival, and apoptosis have been documented.^{21,22} Its ability to react as an electrophile with biological nucleophiles, predominantly Cys thiols, underlies many of its biological effects.²³ To determine whether 15d-PGJ₂ activates PPAR δ by forming a covalent adduct, we first examined 15d-PGJ₂'s ability to activate PPAR δ using a PPAR δ -specific cell reporter assay. As shown in Figure 1A, 15d-PGJ₂ activated PPAR δ in a dose-dependent manner. GW0742, a well-characterized PPAR δ synthetic agonist used as a positive control, also induced PPAR δ activation as expected (Figure 1A). We then performed an electrophoretic mobility shift assay (EMSA) to investigate whether 15d-PGJ₂ induces PPAR δ binding to the PPRE. 15d-PGJ₂ enhanced recombinant PPAR δ 's binding to the consensus PPRE but not to non-specific oligonucleotides (Figure 1B). Likewise, the binding of PPAR δ to the PPRE in normal human bronchial epithelial (NHBE) cell nuclear extracts was increased by 15d-PGJ₂ (Figure 1B). Supershift of the PPRE-PPAR δ complex in the presence of a PPAR δ -specific antibody further confirmed the specificity of this interaction (Figure 1B). Consistently, a

chromatin immunoprecipitation (ChIP) assay further proved that 15d-PGJ₂ stimulated PPAR δ binding to PPRE in NHBE cells (Figure 1C). Together, these data indicate that 15d-PGJ₂ activates PPAR δ 's transcriptional activity and provide the first evidence that PPAR δ functions as a mediator of 15d-PGJ₂'s biological effects.

cyPGs can be present in their free forms or bound to Cys, glutathione, or albumin via their cyclopentenone rings.^{16,24,25} Although pathological stimuli such as inflammation can elevate PG concentrations,^{22,26,27} 15d-PGJ₂'s basal level has been proposed to be in the picomolar-to-nanomolar range.^{21,22,28} This concentration is much lower than the micromolar concentrations required for exogenous 15d-PGJ₂ to elicit biological effects. The discrepancy may be explained by cyPGs' high metabolic rate, reactivity, and preference to be protein-bound as well as the nature of commonly used 15d-PGJ₂ assays to detect primarily free molecules.^{11,21} These factors are expected to interfere with accurate measurements and underestimate biologically available PGs. It should also be noted that most of these studies focus on extracellular/circulating PG levels. However, PGs have been shown to accumulate in the nucleus,^{29,30} either via nuclear production or indirectly via diffusion, and thus may be present at a higher concentration there than in the extracellular circulation. Since 15d-PGJ₂'s major targets are nuclear receptors, nuclear 15d-PGJ₂ levels may in fact be more appropriate. Indeed, not only concentrations but also intra/subcellular locations are significant factors to consider when studying ligand-induced activation.³¹ Kliewer et al. showed that the EC₅₀ for 15d-PGJ₂ required to activate PPAR γ is 2 μ M and noted that high nanomolar-to-micromolar concentrations of PGD₂ and PGJ₂ activate PPARs. These concentrations have been detected in human samples and thereby are potentially physiologically relevant.¹¹ Together, our findings that PPAR δ is activated by 15d-PGJ₂ are not only in agreement with the study by Forman *et al.*¹⁹ but also supported by our current knowledge about 15d-PGJ₂ concentrations and localization.

15d-PGJ₂ interacts covalently with PPAR δ .

Next, we investigated potential covalent interaction of 15d-PGJ₂ with PPAR δ by incubating PPAR δ with increasing concentrations of biotinylated 15d-PGJ₂ (15d-PGJ₂-Biotin), in the presence or absence of unlabeled 15d-PGJ₂, 9,10-dihydro-15-deoxy-^{12,14}-PGJ₂ (CAY10410), or dithiothreitol (DTT), followed by Western blotting of the resulting complexes. The interaction with PPAR δ increased with increasing concentrations of 15d-PGJ₂-Biotin (Figure 2). DTT, a competitive nucleophile, completely abolished this interaction via β -elimination followed by nucleophilic addition, thus confirming the covalent nature of the complex (Figure 2). Moreover, unlabeled 15d-PGJ₂ dose-dependently competed with formation of the 15d-PGJ₂-Biotin-PPAR δ complex and thus corroborated the specificity of the interaction (Figure 2). CAY10410 is a PGD₂/PGJ₂ analog that lacks the endocyclic β -carbon electrophile (C₉ enone) and the lowest energy unoccupied molecular orbital (LUMO) in the cyclopentenone ring (Figure S1). CAY10410's failure to compete for such interaction (Figure 2) indicates that the 15d-PGJ₂-PPAR δ covalent interaction relies on the electrophilic nature of the cyclopentenone ring as well as the higher LUMO coefficient of the C₉ (Figure S1). These results demonstrate that 15d-PGJ₂ interacts covalently with PPAR δ using the enone present in its cyclopentenone ring.

15d-PGJ₂ interacts with a Cys within PPAR δ 's LBD.

The covalent interactions of 15d-PGJ₂ with Cys in different target proteins are summarized in Table S1. Soares *et al.* showed that 15d-PGJ₂ covalently interacts with PPAR γ 's Cys285.²⁰ To explore the possibility of PPAR δ Cys-alkylation by 15d-PGJ₂, we performed an *in vitro* electrophilic addition reaction assay using the infrared dye (IRDye) 800CW maleimide with either NHBE cell extracts, the recombinant PPAR δ protein, or the recombinant PPAR δ LBD. Maleimide binds to molecules containing Cys with free sulfhydryl groups (-SH) and is a useful tool for investigating Cys modification. With each protein sample, 15d-PGJ₂-Biotin reduced the amount of signal representing the maleimide-bound Cys in the proteins (Figure 3), indicating that 15d-PGJ₂ binds to a Cys within PPAR δ 's LBD.

Molecular interactions of the 15d-PGJ₂-PPAR δ complex formation.

Analysis of the seven-crystallographic 15d-PGJ₂-PPAR δ complexes and their binding modes predicted that the ligand occupied the receptor's Y-shaped LBD by interacting with amino acids of arms I to III (Figure 4, Figure S2, and Table S2). To determine the covalent/non-covalent nature of the 15d-PGJ₂-PPAR δ interaction as well as the role of Cys249, we modeled and docked different 15d-PGJ₂-PPAR δ complexes. Figure 4 shows the lowest energy pose of each complex (based on the free energy perturbation), in all of which 15d-PGJ₂ occupied the Y-shaped LBD and adopted an orientation predicted by the pharmacophoric model of PPAR ligands.^{9,10} Conformations obtained from the covalent (Figure 4B) and the non-covalent (Figure 4C) docking present H12-stabilizing hydrogen bonds between 15d-PGJ₂'s acidic, hydrophilic head group (carboxylic acid group) and PPAR δ 's hydrophilic amino acids Thr253 (H3), His287 (H5), His413 (H11), and Tyr437 (H12). In all the docking models, the aliphatic tail group positioned mostly in arms II and III; it interacted with the hydrophobic amino acids Val245 (H3), Leu303 (S2), Leu317 (H6), and Ile328 (H7) of arm II and with Leu294 (H5) of arm III, which creates the entrance. The core cyclopentenone ring connecting the head and the tail groups lies close to H3 in the covalent docking (Figure 4B) and away from H3 in the non-covalent docking (Figure 4C).

In the covalent conformation, the electrophilic endocyclic β -carbon (C₉) in 15d-PGJ₂'s cyclopentenone ring was separated from the sulfur atom (S γ) of Cys249 by ~2–3 Å, a distance compatible with a 1,4 addition reaction and thus consistent with a covalent bond between them (Figure 4B). However, no such covalent bond was seen in the PPAR δ C249S mutant (the cysteine at the residue 249 was substituted with a serine) because the oxygen atom (O γ) in Ser249 was oriented away from the electrophilic C₉ (Figure 4D). These data support covalent interactions between 15d-PGJ₂'s C₉ and Cys249 in PPAR δ 's LBD, which was further confirmed by pharmacophore mapping (Figure S2). The hydrogen-bonding regions of the ligand and the receptor interact as expected from molecular modeling, as do the hydrophobic regions. The interatom distances (Figure S2 inset) are similarly compatible with a covalent bond between the 15d-PGJ₂ enone and Cys249 of PPAR δ .

A recent study described a covalent antagonist for PPAR δ ,³² but no covalently interacting molecule has so far been reported to activate PPAR δ . Therefore, our data revealing 15d-

PGJ₂ as a covalent PPAR δ agonist may provide new insights into other signaling pathways mediated by covalently activated PPAR δ .

15d-PGJ₂ covalently binds to Cys249 via Cys-alkylation.

To determine whether PPAR δ Cys249 was the specific Cys-alkylation target of 15d-PGJ₂, we examined the effect of the PPAR δ C249S mutation on Cys-alkylation by 15d-PGJ₂. After incubation with the PPAR δ wildtype (WT) or the PPAR δ C249S mutant, binding of 15d-PGJ₂Biotin was assessed by Western blotting. 15d-PGJ₂ bound to PPAR δ WT via Cys-alkylation (Figure 5A). This binding was completely abolished by DTT treatment (Figure 5A), supporting the covalent nature of the interaction. As predicted from the molecular modeling data (Figure 4D), 15d-PGJ₂ failed to interact with the PPAR δ C249S mutant (Figure 5A). These results indicate that the Cys-alkylation of PPAR δ by 15d-PGJ₂ occurs specifically at Cys249.

To investigate the functional consequences of the C249S mutation on 15d-PGJ₂-mediated PPAR δ activation, we studied PPAR δ 's DNA binding activity by an ELISA-based PPAR δ activity assay. No significant change was observed in the basal activity of PPAR δ WT and C249S mutant (Figure 5B). In the presence of 15d-PGJ₂, however, unlike PPAR δ WT whose DNA binding activity increased with increasing concentrations of the ligand, C249S mutant failed to show a substantial change in its DNA binding at any concentration, reflecting its inability to interact with the ligand (Figure 5B). Thus, the role of Cys249 as a target for Cys-alkylation by 15d-PGJ₂ renders it critical for covalent ligand-receptor interaction and the consequent activation of DNA binding.

Calculated chemical properties of 15d-PGJ₂ and CAY10410 and of WT and mutant PPAR δ .

The calculated chemical properties of 15d-PGJ₂ and CAY10410 as potential electrophiles and of Cys and Ser as potential nucleophiles are presented in Tables 1 and 2: hardness (η), softness (σ), chemical potential (μ), and electrophilicity (ω) (Table 1); Fukui functions, dual descriptors (DDs), and atomic charges (Table 2). Based on their molecular characteristics (Table 1), 15d-PGJ₂ is predicted to be soft and more electrophilic than CAY10410. The electron density distribution in the isosurface of Fukui functions and DDs also revealed that 15d-PGJ₂'s endocyclic β -carbon (C₉) in the cyclopentenone ring is more susceptible to nucleophilic attack than the exocyclic β -carbon electrophile (C₁₃) or the C₉ carbon of CAY10410 and is thus the preferred nucleophilic attack site (Table 2). The region/atom with a larger electrophilic Fukui function value is considered the better electrophilic attack site. CAY10410's small electrophilic Fukui function value at the endocyclic β -carbon (C₉) further explains its inferiority as an electrophile (Table 2). 15d-PGJ₂ has been proposed to use this carbon to interact with several other proteins also;^{33,34} our findings provide a convincing chemical and structural basis for the 15d-PGJ₂-PPAR δ complex that had only been implied by these earlier studies.

Compared to the serine residue (Ser)-O γ , the Cys-S γ is larger, less electronegative, and less able to attract the outermost valence electrons. These features, along with a higher relative reactivity ($>10^5$ vs. 10^4 of the Ser-O γ), make the Cys-S γ a highly polarizable/softer, more nucleophilic target. The lower pK_a value of the Cys-S γ than that of the Ser-O γ (pK_a 9 vs.

pKa 13, respectively) also agrees with its greater reactivity (Figure S3). Moreover, Kortemme *et al.* previously showed that α -helices are more positively charged at the N-terminal end, thereby lowering the pKa of Cys located in those regions.³⁵ Cys249's localization in the N-terminal region of the H3 thus contributes to its greater nucleophilicity.

The PPAR δ LBD contains two Cys residues in the H3, Cys249 (conserved as Cys276 in PPAR α and Cys285 in PPAR γ) and Cys251 (conserved only in PPAR α as Cys278) (Figure S4). Further structural analysis revealed that Cys251 is positioned away from the Y-shaped LBD towards the outer surface of the molecule and is thus unavailable for ligand interaction (Figure S5). Finally, the nucleophilic Fukui function predicted Cys-S γ as the preferred site for electrophilic attack (Table 2). Together, these data provide the chemical basis for 15d-PGJ₂'s covalent interactions with PPAR δ Cys249.

Frontier molecular orbital (MO) interaction.

In an electrophile-nucleophile addition reaction, a new bond is formed when the highest energy occupied molecular orbital (HOMO) of the nucleophile overlaps with the LUMO. The quantum chemical calculations and orbital density distributions revealed that the centered HOMO of the Cys249's S γ (Figure 6A) and the LUMO of 15d-PGJ₂ C β (C₉) (Figure 6B) are in phase and thereby overlap to form a covalent bond (Figure 6C). The HOMOs on the nucleophilic sulfur anion (the Cys-S γ 's lone pair) (Figure 6A) are perpendicular to each other and have almost identical energies. In the LUMOs, the largest orbital coefficient is on the endocyclic β -carbon electrophile (C₉) in 15d-PGJ₂'s cyclopentenone ring (Figure 6B). These LUMOs are perpendicular to the plane of the cyclopentenone ring; this is the site of attack by PPAR δ 's nucleophilic Cys. The reaction starts when one of the two bonding HOMOs on the Cys-S γ atom comes in the plane of the target cyclopentenone ring and competes with covalent bonding. Acting as an electron-withdrawer in the reaction complex, the ketone group in the cyclopentenone ring's enone increases the rate of nucleophilic attack by PPAR δ Cys249 and stabilizes the resulting covalent adduct.

15d-PGJ₂ and PPAR δ engage in a 1,4 addition reaction.

Some effects of 15d-PGJ₂ on intracellular proteins are mediated by conjugate addition of a thiol nucleophile to the electrophilic carbon in the cyclopentenone ring (Table S1). Specifically, this reaction involves a classical conjugate 1,4 addition to the C=C double bond located next to the C=O in an enone; an electron-withdrawing group polarizes the double bond so that the β -carbon acquires a partial positive character and becomes electrophilic. Thus, we propose that the enone in 15d-PGJ₂'s cyclopentenone ring polarizes the double bond and stabilizes the resulting negative charge on its own oxygen atom. Quantum chemical calculation showed this enone is a soft electrophile (Table 1; σ 0.229 eV⁻¹) that is expected to react with the Cys thiol (Table 1; σ 0.212 eV⁻¹), the softest biological nucleophile.³⁶ The hard-soft acid-base (HSAB) theory further agrees with our hypothesis by stating that a soft nucleophile (soft base) prefers to react with a soft electrophile (soft acid). As shown in Figures 6 (Movie S1) and 7 (Movies S2 and S3), the lone pair electrons in the HOMO of the Cys249-S γ attack the endocyclic β -carbon (C β) of 15d-PGJ₂ that has a large LUMO orbital coefficient and electrophilic Fukui function. This reaction causes HOMO-

LUMO overlap and forms a new covalent adduct. Finally, the intrinsic reaction coordinate showing the reactants, the transition state, and the products illustrates the predicted mechanism/path of the chemical reaction between 15d-PGJ₂ and a Cys (Figure 8 and Movie S4). Together, our data support that PPAR δ and 15d-PGJ₂ covalently interact by engaging in a 1,4 addition reaction. Importantly, our study is the first to our knowledge to provide the exact electronic structure, quantum chemical, and quantum mechanical basis of a 1,4 addition reaction between a protein thiol and a lipid electrophile. Furthermore, DFT calculations, reactive molecular dynamics simulations, and intrinsic reaction coordinate modeling prove the selectivity of PPAR δ Cys thiol towards the endocyclic enone in 15d-PGJ₂ and the mechanism underlying the overlap of HOMO-LUMO orbital forming a covalent bond.

15d-PGJ₂ as a covalent ligand may elicit PPAR δ -mediated effects that are distinct from those induced by non-covalent synthetic counterparts such as GW0742 and could conceivably offer additional clinical benefits.²⁰ Under physiological conditions, covalent interactions are essentially irreversible.^{20,30} Because ligand binding leads to a structural change in PPARs that allows coactivator recruitment, 15d-PGJ₂'s covalent interaction implies prolonged receptor activation.²⁰ This biochemical feature may translate to enhanced efficacy and improved pharmacological properties, which could lead to a reduction in drug dosage and dosing frequency as well as patients' systemic exposure to drugs.^{37–39}

PPAR δ is expressed ubiquitously throughout the body and is involved in a variety of cellular processes in a variety of pathophysiological contexts.^{40–42} Its clinical potentials have been proposed in many diseases (e.g. inflammatory diseases⁴⁰, metabolic diseases,^{43–45} a neurodegenerative disorder,⁴⁶ cardiovascular diseases,^{45,47} and cancer⁴¹). While there are currently no FDA-approved drugs that target PPAR δ ,^{43,44} some PPAR δ agonists are being investigated in clinical trials.⁴⁴ Taken together, our study strongly supports molecules with the chemical features of 15d-PGJ₂ as potential pharmacological agents that may offer unique benefits for patients with a variety of diseases.

METHODS

Cells.

NHBE cells, obtained from Lonza, were grown and maintained in BEGM (Lonza) supplemented with growth factors and hormones, as described in the manufacturer's instructions, at 37°C in a humidified atmosphere of 5% CO₂–95% air. After cells were treated with 15d-PGJ₂ or vehicle in basal medium for 24 h, nuclear extracts were prepared. The BCA Protein Assay kit (Pierce,) was used to determine protein concentrations.

PPAR δ Reporter Assay.

To determine the activation of PPAR δ by 15d-PGJ₂, we performed a PPAR δ -specific luciferase reporter assay, using the PPAR δ Reporter Assay System (IB00121; Indigo Biosciences) according to the manufacturer's instructions. Briefly, PPAR δ reporter cells were cultured in Cell Recovery Medium and treated with various concentrations of 15d-PGJ₂ (0.005–10 μ M) or GW0742 (0.005–10 μ M; positive control) for 24 h. At the end of the

treatment, the culture medium was discarded, and Luciferase Detection Reagent was added to each well of the assay plate. The plate was then incubated for 5 min at room temperature, and luminescence was quantified with a luminometer.

EMSA.

Recombinant PPAR δ protein (1 μ g) or nuclear proteins from NHBE cells (30 μ g) were incubated with 50 nM of double-stranded nonspecific or consensus PPRE oligonucleotides end-labeled with IRDye 700 in binding buffer (100 mM Tris, 500 mM KCl, 10 mM DTT [pH 7.5]), poly(deoxyinosinic-deoxycytidylic) (1 μ g/ μ l in 10 mM Tris, 1 mM EDTA), 25 mM DTT, and 2.5% (v/v) Tween 20. Samples were then separated on 5% non-denaturing polyacrylamide gels in 1 $\times$ Tris-Borate EDTA buffer (130 mM Tris [pH 8.3], 45 mM boric acid, 2.5 mM EDTA). In the supershift reaction, the reaction mixture was incubated with antibody against PPAR δ (101720; Cayman Chemical). The infrared signal was detected with an Odyssey Infrared Imager (LI-COR Biosciences).

ChIP Assay.

SimpleChIP Enzymatic Chromatin Immunoprecipitation Kit with magnetic beads (Cell Signaling Technology) was used, as described previously,⁴⁸ to assess DNA binding activity of PPAR δ in cells. An equal amount of chromatin was mixed with PPAR δ antibodies or IgG control. After overnight incubation at 4°C, protein G magnetic beads were added, and the chromatin-antibody-bead mixture was incubated for another 2 h at 4°C with rotation. An aliquot of chromatin that was incubated without an antibody was used as the input control. Antibody-bound protein-DNA complexes were eluted and subjected to real-time PCR⁴⁸ with specific primers that amplify the PPRE and β -Actin.

Western Blotting.

Western blotting was performed as described previously.⁴⁹ Primary antibody against PPAR δ was purchased from Cayman Chemical. The secondary antibodies, IRDye 800CW goat anti-rabbit IgG (H+L) (925–32211), IRDye 680RD streptavidin (925–68079), and IRDye 800CW maleimide (929–80020), were obtained from LI-COR Biosciences. The infrared signal was detected with an Odyssey Infrared Imager.

Recombinant PPAR δ Protein and PPAR δ Cys-alkylation.

Human recombinant PPAR δ protein was expressed by transforming pET28a plasmid (custom cloning core facility at Emory University) into One Shot BL21 Star (DE3) cells (C601003, Thermo Fisher Scientific). Cells were lysed, and the recombinant protein was purified and concentrated according to standard protocols, as described previously.⁵⁰ We excluded reducing agents from all buffers.

The resulting recombinant PPAR δ was incubated with 15d-PGJ₂-Biotin (10141; Cayman Chemical) for 1 h, which was followed by 30-min incubation with or without DTT (10 or 100 mM). Cys-alkylation of the samples was then analyzed on an SDS-PAGE gel under non-reducing conditions, followed by Western blotting to detect 15d-PGJ₂-Biotin (IRDye 680 streptavidin) and PPAR δ (IRDye 800). The infrared signal was detected with an Odyssey Infrared Imager.

***In Vitro* Electrophilic Addition Reaction.**

An *in vitro* electrophilic addition reaction was performed to assess 15d-PGJ₂'s ability to form covalent electrophile-protein (Cys) adducts. NHBE cell extracts, recombinant PPAR δ , or isolated PPAR δ LBD proteins were incubated with indicated concentrations of 15d-PGJ₂-Biotin or 1 μ M IRDye 800CW maleimide in PBS at room temperature for 30 min. In the competitive reactions including both electrophiles, IRDye 800CW maleimide was introduced 30 min after the addition of 15d-PGJ₂-Biotin. When the reactions were complete, the protein adducts formed were analyzed on an SDS-PAGE gel under non-reducing conditions, followed by Western blotting to detect 15d-PGJ₂ (IRDye 680 streptavidin) and maleimide (IRDye 800). The infrared signal was detected with an Odyssey Infrared Imager.

Molecular Modeling, Molecular Dynamics Simulations, and Docking.

Molecular modeling, molecular dynamics simulations, and docking analysis were carried out as described previously⁵⁰ with Discovery Studio software (BIOVIA) and Chem3D 17.0. Briefly, using seven X-ray structures of human PPAR δ (PDB IDs: 3tkm, 3sp9, 3et2, 2xyx, 2xyw, 2xyj and 2j14) obtained from the RCSB Protein Data Bank, we analyzed 15d-PGJ₂-PPAR δ interactions and generated a pharmacophore. The X-ray structure 3tkm⁵¹ with 1.95 Å resolution (the structure with the highest resolution among those selected) was chosen to build the PPAR δ C249S mutant with DS MODELER. All structures and complexes were energy minimized with CHARMM. 15d-PGJ₂ was used as a ligand to perform covalent and non-covalent docking.

Recombinant PPAR δ C249S Mutant Protein and PPAR δ Cys-alkylation.

Similarly to the recombinant PPAR δ (WT) protein described above, human recombinant PPAR δ C249S mutant protein was expressed by inserting the transforming pET28a plasmid into One Shot BL21 Star (DE3) cells. Cells were lysed, and recombinant mutant protein was purified and concentrated according to standard protocols. We excluded reducing agents from all buffers.

To assess Cys-alkylation by 15d-PGJ₂, we incubated PPAR δ WT and PPAR δ C249S proteins with or without 15d-PGJ₂-Biotin (10 μ M) for 1 h and then with or without DTT (10 mM) for 30 min. Cys-alkylation was analyzed on an SDS-PAGE gel under non-reducing conditions, followed by Western blotting to detect 15d-PGJ₂-Biotin (IRDye 680 streptavidin) and PPAR δ (IRDye 800).

PPAR δ Activity Assay.

PPAR δ Transcription Factor Assay Kit (10006914, Cayman Chemical) was used according to the manufacturer's instructions, to assess PPAR δ 's DNA binding activity.

Quantum Chemical Calculations.

The density functional theory (DFT) and the Quantum Theory of Atoms in Molecules (QTAIM) calculations were performed with the Amsterdam Density Functional (ADF) Modeling Suite,⁵² as implemented with default settings. Briefly, after the molecule has been inputted in ADFinput, the molecular structure was fully optimized with Generalized

Gradient Approximation (GGA) to the Becke and Perdew (BP) exchange-correlation energy functional. The Bader theory was selected for the calculation of critical points, bond paths, atomic properties, atomic energies, and reactivity indices. The electrophilic and nucleophilic Fukui functions were calculated with a charge change parameter set to one. Structures, orbitals, and molecular graphs were drawn and visualized with the graphical user interface of ADF (ADF-GUI). For each molecule, chemical hardness (η), chemical softness (σ), chemical potential (μ), and electrophilicity index (ω) were calculated by determining E_{HOMO} and E_{LUMO} and with the standard equations.

ReaxFF Molecular Dynamics Simulations and Reaction Coordinate.

DFT and ReaxFF features in ADF⁵² were used, as implemented by the program, for all the reactive dynamics simulations and transition state search. The geometry-optimized structures, generated based on DFT and non-reactive molecular dynamics, were used as starting structures. The 1,4 addition reaction was studied with a Cys molecule as the nucleophile and 15d-PGJ₂ (cyclopentenone in ReaxFF) as the electrophile. Reactive simulations were performed in the NVE ensemble using the CHONSMgPNaCuCl_v2 force field. To find the minimum energy path for the 1,4 addition reaction, we performed the transition state search using Atomic Simulation Environment (ASE) with the Nudged Elastic Band (ASE-NEB) method in ADF.

Quantum mechanics were performed to investigate transition state structures, reaction path structures, and intrinsic reaction coordinate. Specifically, we used the semiempirical parametric method (PM6) and COSMO to simulate water solvation effects, as implemented by SCiGRESS (v 2.8.1, Fujitsu Ltd.). Potential energy mapping, reaction saddle point, vibrational transitions, and the reaction path from the reactants through a transition state to the products were calculated with the same parameter set and solvent effects.

Statistical Analysis.

Data are presented as the mean $\pm$ SD. We determined the differences between experimental groups using analysis of variance followed by a Bonferroni multiple comparison correction. GraphPad Prism 7 (GraphPad Software) was used to perform these statistical analyses. $p < 0.05$ was considered significant.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

FUNDING SOURCES

This work was supported by a merit review award from the U.S. Department of Veterans Affairs, Flight Attendant Medical Research Institute Clinical Innovator Award 16006, and National Institutes of Health Grants HL137842 and AI125338 (R.C.R.).

The abbreviations used are:

PPAR peroxisome proliferator-activated receptor

PPRE	PPAR response element
LBD	ligand-binding domain
H	α -helix
S	β -strand
PG	prostaglandin
cyPG	cyclopentenone PG
15d-PGJ₂	15-deoxy- ^{12,14} -PGJ ₂
Cys	cysteine residue(s)
EMSA	electrophoretic mobility shift assay
NHBE	normal human bronchial epithelial
ChIP	chromatin immunoprecipitation
15d-PGJ₂-Biotin	biotinylated 15d-PGJ ₂
CAY10410	9,10-dihydro-15-deoxy- ^{12,14} -PGJ ₂
LUMO	lowest energy unoccupied molecular orbital
C	carbon atom
S	sulfur atom
O	oxygen atom
WT	wildtype
DD	dual descriptor
Ser	serine residue
MO	molecular orbital
HOMO	highest energy occupied molecular orbital
HSAB	hard-soft acid base
IRDye	infrared dye
DFT	density functional theory
QTAIM	Quantum Theory of Atoms in Molecules
ADF	Amsterdam Density Functional
GGA	Generalized Gradient Approximation
BP	Becke and Perdew

GUI	graphical user interface
ASE	Atomic Simulation Environment
NEB	Nudged Elastic Band
S/B	signal-to-background
HBD	hydrogen bond donor
HBA	hydrogen bond acceptor
RA	ring aromatic
HY	hydrophobic feature and location

REFERENCES

- Berger J, and Moller DE (2002) The mechanisms of action of PPARs. *Annu Rev Med* 53, 409–435 [PubMed: 11818483]
- Issemann I, and Green S (1990) Activation of a member of the steroid hormone receptor superfamily by peroxisome proliferators. *Nature* 347, 645–650 [PubMed: 2129546]
- Becker J, Delayre-Orthez C, Frossard N, and Pons F (2006) Regulation of inflammation by PPARs: a future approach to treat lung inflammatory diseases? *Fundam Clin Pharmacol* 20, 429–447 [PubMed: 16968414]
- Wahli W, and Michalik L (2012) PPARs at the crossroads of lipid signaling and inflammation. *Trends Endocrinol Metab* 23, 351–363 [PubMed: 22704720]
- Tyagi S, Gupta P, Saini AS, Kaushal C, and Sharma S (2011) The peroxisome proliferator-activated receptor: A family of nuclear receptors role in various diseases. *J Adv Pharm Technol Res* 2, 236–240 [PubMed: 22247890]
- Belvisi MG, and Mitchell JA (2009) Targeting PPAR receptors in the airway for the treatment of inflammatory lung disease. *Br J Pharmacol* 158, 994–1003 [PubMed: 19703165]
- Daynes RA, and Jones DC (2002) Emerging roles of PPARs in inflammation and immunity. *Nat Rev Immunol* 2, 748–759 [PubMed: 12360213]
- Xu HE, Lambert MH, Montana VG, Plunket KD, Moore LB, Collins JL, Oplinger JA, Kliewer SA, Gampe RT, and McKee DD (2001) Structural determinants of ligand binding selectivity between the peroxisome proliferator-activated receptors. *Proceedings of the National Academy of Sciences* 98, 13919–13924
- Xu HE, Lambert MH, Montana VG, Parks DJ, Blanchard SG, Brown PJ, Sternbach DD, Lehmann JM, Wisely GB, Willson TM, Kliewer SA, and Milburn MV (1999) Molecular recognition of fatty acids by peroxisome proliferator-activated receptors. *Mol Cell* 3, 397–403 [PubMed: 10198642]
- Fyffe SA, Alphey MS, Buetow L, Smith TK, Ferguson MA, Sorensen MD, Bjorkling F, and Hunter WN (2006) Recombinant human PPAR-beta/delta ligand-binding domain is locked in an activated conformation by endogenous fatty acids. *J Mol Biol* 356, 10051013
- Kliewer SA, Lenhard JM, Willson TM, Patel I, Morris DC, and Lehmann JM (1995) A prostaglandin J2 metabolite binds peroxisome proliferator-activated receptor gamma and promotes adipocyte differentiation. *Cell* 83, 813–819 [PubMed: 8521498]
- Forman BM, Chen J, and Evans RM (1997) Hypolipidemic drugs, polyunsaturated fatty acids, and eicosanoids are ligands for peroxisome proliferator-activated receptors alpha and delta. *Proc Natl Acad Sci U S A* 94, 4312–4317 [PubMed: 9113986]
- Giles H, and Leff P (1988) The biology and pharmacology of PGD2. *Prostaglandins* 35, 277–300 [PubMed: 3283848]
- Fukushima M (1992) Biological activities and mechanisms of action of PGJ2 and related compounds: an update. *Prostaglandins Leukot Essent Fatty Acids* 47, 1–12 [PubMed: 1438462]

15. Shibata T, Kondo M, Osawa T, Shibata N, Kobayashi M, and Uchida K (2002) 15-deoxy-delta 12,14-prostaglandin J2. A prostaglandin D2 metabolite generated during inflammatory processes. *J Biol Chem* 277, 10459–10466 [PubMed: 11786541]
16. Noyori R, and Suzuki M (1993) Organic synthesis of prostaglandins: advancing biology. *Science* 259, 44–45 [PubMed: 8418493]
17. Atsmon J, Freeman ML, Meredith MJ, Sweetman BJ, and Roberts LJ (1990) Conjugation of 9-deoxy- 9, 12 (E)-prostaglandin D2 with intracellular glutathione and enhancement of its antiproliferative activity by glutathione depletion. *Cancer research* 50, 18791885
18. Bogaards JJ, Venekamp JC, and van Bladeren PJ (1997) Stereoselective conjugation of prostaglandin A2 and prostaglandin J2 with glutathione, catalyzed by the human glutathione S-transferases A1–1, A2–2, M1a-1a, and P1–1. *Chem Res Toxicol* 10, 310–317 [PubMed: 9084911]
19. Forman BM, Tontonoz P, Chen J, Brun RP, Spiegelman BM, and Evans RM (1995) 15-Deoxy-delta 12, 14-prostaglandin J2 is a ligand for the adipocyte determination factor PPAR gamma. *Cell* 83, 803–812 [PubMed: 8521497]
20. Soares AF, Nosjean O, Cozzone D, D’Orazio D, Becchi M, Guichardant M, Ferry G, Boutin JA, Lagarde M, and Geloën A (2005) Covalent binding of 15-deoxy-delta12,14prostaglandin J2 to PPARgamma. *Biochem Biophys Res Commun* 337, 521–525 [PubMed: 16198309]
21. Marcone S, and Fitzgerald DJ (2013) Proteomic identification of the candidate target proteins of 15-deoxy-delta12,14-prostaglandin J2. *Proteomics* 13, 2135–2139 [PubMed: 23606334]
22. Morgenstern J, Fleming T, Kadiyska I, Brings S, Groener JB, Nawroth P, Hecker M, and Brune M (2018) Sensitive mass spectrometric assay for determination of 15-deoxyDelta(12,14)-prostaglandin J2 and its application in human plasma samples of patients with diabetes. *Anal Bioanal Chem* 410, 521–528 [PubMed: 29143878]
23. Shibata T (2015) 15-Deoxy-Delta(1)(2),(1)(4)-prostaglandin J(2) as an electrophilic mediator. *Biosci Biotechnol Biochem* 79, 1044–1049 [PubMed: 26011133]
24. Atsmon J, Sweetman BJ, Baertschi SW, Harris TM, and Roberts LJ 2nd. (1990) Formation of thiol conjugates of 9-deoxy-delta 9,delta 12(E)-prostaglandin D2 and delta 12(E)prostaglandin D2. *Biochemistry* 29, 3760–3765 [PubMed: 2340271]
25. Person EC, Waite LL, Taylor RN, and Scanlan TS (2001) Albumin regulates induction of peroxisome proliferator-activated receptor-gamma (PPARgamma) by 15-deoxy-delta(12–14)prostaglandin J(2) in vitro and may be an important regulator of PPARgamma function in vivo. *Endocrinology* 142, 551–556 [PubMed: 11159824]
26. Gilroy DW, Colville-Nash PR, Willis D, Chivers J, Paul-Clark MJ, and Willoughby DA (1999) Inducible cyclooxygenase may have anti-inflammatory properties. *Nat Med* 5, 698701
27. McAdam BF, Mardini IA, Habib A, Burke A, Lawson JA, Kapoor S, and FitzGerald GA (2000) Effect of regulated expression of human cyclooxygenase isoforms on eicosanoid and isoeicosanoid production in inflammation. *J Clin Invest* 105, 1473–1482 [PubMed: 10811855]
28. Bell-Parikh LC, Ide T, Lawson JA, McNamara P, Reilly M, and FitzGerald GA (2003) Biosynthesis of 15-deoxy-delta12,14-PGJ2 and the ligation of PPARgamma. *J Clin Invest* 112, 945–955 [PubMed: 12975479]
29. Narumiya S, Ohno K, Fukushima M, and Fujiwara M (1987) Site and mechanism of growth inhibition by prostaglandins. III. Distribution and binding of prostaglandin A2 and delta 12-prostaglandin J2 in nuclei. *J Pharmacol Exp Ther* 242, 306–311 [PubMed: 3302206]
30. Suzuki M, Mori M, Niwa T, Hirata R, Furuta K, Ishikawa T, and Noyori R (1997) Chemical Implications for Antitumor and Antiviral Prostaglandins: Reaction of 7-Prostaglandin A1 and Prostaglandin A1 Methyl Esters with Thiols. *Journal of the American Chemical Society* 119, 2376–2385
31. Narala VR, Adapala RK, Suresh MV, Brock TG, Peters-Golden M, and Reddy RC (2010) Leukotriene B4 is a physiologically relevant endogenous peroxisome proliferator-activated receptor-alpha agonist. *J Biol Chem* 285, 22067–22074 [PubMed: 20400503]
32. Kaupang A, Paulsen SM, Steindal CC, Ravna AW, Sylte I, Halvorsen TG, Thoresen GH, and Hansen TV (2015) Synthesis, biological evaluation and molecular modeling studies of the PPARbeta/delta antagonist CC618. *Eur J Med Chem* 94, 229–236 [PubMed: 25768705]

33. Sanchez-Gomez FJ, Cernuda-Morollon E, Stamatakis K, and Perez-Sala D (2004) Protein thiol modification by 15-deoxy-Delta12,14-prostaglandin J2 addition in mesangial cells: role in the inhibition of pro-inflammatory genes. *Mol Pharmacol* 66, 1349–1358 [PubMed: 15317873]
34. Straus DS, Pascual G, Li M, Welch JS, Ricote M, Hsiang CH, Sengchanthalangsy LL, Ghosh G, and Glass CK (2000) 15-deoxy-delta 12,14-prostaglandin J2 inhibits multiple steps in the NF-kappa B signaling pathway. *Proc Natl Acad Sci U S A* 97, 4844–4849 [PubMed: 10781090]
35. Kortemme T, and Creighton TE (1995) Ionisation of Cysteine Residues at the Termini of Model α -Helical Peptides. Relevance to Unusual Thiol pKa Values in Proteins of the Thioredoxin Family. *Journal of Molecular Biology* 253, 799–812 [PubMed: 7473753]
36. LoPachin RM, Gavin T, Petersen DR, and Barber DS (2009) Molecular mechanisms of 4-hydroxy-2-nonenal and acrolein toxicity: nucleophilic targets and adduct formation. *Chem Res Toxicol* 22, 1499–1508 [PubMed: 19610654]
37. Bauer RA (2015) Covalent inhibitors in drug discovery: from accidental discoveries to avoided liabilities and designed therapies. *Drug Discov Today* 20, 1061–1073 [PubMed: 26002380]
38. Singh J, Petter RC, Baillie TA, and Whitty A (2011) The resurgence of covalent drugs. *Nat Rev Drug Discov* 10, 307–317 [PubMed: 21455239]
39. Wang L, Zhao J, Yao Y, Wang C, Zhang J, Shu X, Sun X, Li Y, Liu K, Yuan H, and Ma X (2017) Covalent binding design strategy: A prospective method for discovery of potent targeted anticancer agents. *Eur J Med Chem* 142, 493–505 [PubMed: 28986130]
40. Banno A, Reddy AT, Lakshmi SP, and Reddy RC (2018) PPARs: Key Regulators of Airway Inflammation and Potential Therapeutic Targets in Asthma. *Nucl Receptor Res* 5
41. Lakshmi SP, Reddy AT, Banno A, and Reddy RC (2017) PPAR Agonists for the Prevention and Treatment of Lung Cancer. *PPAR Res* 2017, 8252796 [PubMed: 28316613]
42. Michalik L, Desvergne B, and Wahli W (2004) Peroxisome-proliferator-activated receptors and cancers: complex stories. *Nat Rev Cancer* 4, 61–70 [PubMed: 14708026]
43. Beyaz S, and Yilmaz OH (2016) Molecular Pathways: Dietary Regulation of Stemness and Tumor Initiation by the PPAR-delta Pathway. *Clin Cancer Res* 22, 5636–5641 [PubMed: 27702819]
44. Palomer X, Barroso E, Pizarro-Delgado J, Pena L, Botteri G, Zarei M, Aguilar D, Montori-Grau M, and Vazquez-Carrera M (2018) PPARbeta/delta: A Key Therapeutic Target in Metabolic Disorders. *Int J Mol Sci* 19
45. Vazquez-Carrera M (2016) Unraveling the Effects of PPARbeta/delta on Insulin Resistance and Cardiovascular Disease. *Trends Endocrinol Metab* 27, 319–334 [PubMed: 27005447]
46. Dickey AS, Pineda VV, Tsunemi T, Liu PP, Miranda HC, Gilmore-Hall SK, Lomas N, Sampat KR, Buttgerit A, Torres MJ, Flores AL, Arreola M, Arbez N, Akimov SS, Gaasterland T, Lazarowski ER, Ross CA, Yeo GW, Sopher BL, Magnuson GK, Pinkerton AB, Masliah E, and La Spada AR (2016) PPAR-delta is repressed in Huntington's disease, is required for normal neuronal function and can be targeted therapeutically. *Nat Med* 22, 37–45 [PubMed: 26642438]
47. Ehrenborg E, and Skogberg J (2013) Peroxisome proliferator-activated receptor delta and cardiovascular disease. *Atherosclerosis* 231, 95–106 [PubMed: 24125418]
48. Lakshmi SP, Reddy AT, and Reddy RC (2017) Transforming growth factor beta suppresses peroxisome proliferator-activated receptor gamma expression via both SMAD binding and novel TGF-beta inhibitory elements. *Biochem J* 474, 1531–1546 [PubMed: 28100650]
49. Reddy AT, Lakshmi SP, Zhang Y, and Reddy RC (2014) Nitrated fatty acids reverse pulmonary fibrosis by dedifferentiating myofibroblasts and promoting collagen uptake by alveolar macrophages. *FASEB J* 28, 5299–5310 [PubMed: 25252739]
50. Reddy AT, Lakshmi SP, Muchumarri RR, and Reddy RC (2016) Nitrated Fatty Acids Reverse Cigarette Smoke-Induced Alveolar Macrophage Activation and Inhibit Protease Activity via Electrophilic S-Alkylation. *PLoS One* 11, e0153336 [PubMed: 27119365]
51. Batista FA, Trivella DB, Bernardes A, Gratieri J, Oliveira PS, Figueira AC, Webb P, and Polikarpov I (2012) Structural insights into human peroxisome proliferator activated receptor delta (PPAR-delta) selective ligand binding. *PLoS One* 7, e33643 [PubMed: 22606221]
52. Baerends EJ, Ziegler T, Atkins AJ, Autschbach J, Bashford D, Baseggio O, Brces A, Bickelhaupt FM, Bo C, Boeritger PM, Cavallo L, Daul C, Chong DP, Chulhai DV, Deng L, Dickson RM, Dieterich JM, Ellis DE, van Faassen M, Ghysels A, Giammona A, van Gisbergen SJA, Goetz A,

Gtz AW, Gusarov S, Harris FE, van den Hoek P, Hu Z, Jacob CR, Jacobsen H, Jensen L, Joubert L, Kaminski JW, van Kessel G, Knig C, Kootstra F, Kovalenko A, Krykunov M, van Lenthe E, McCormack DA, Michalak A, Mitoraj M, Morton SM, Neugebauer J, Nicu VP, Noodleman L, Osinga VP, Patchkovskii S, Pavanello M, Peeples CA, Philipsen PHT, Post D, Pye CC, Ramanantoanina H, Ramos P, Ravenek W, Rodrguez JI, Ros P, Rger R, Schipper PRT, Schlus D, van Schoot H, Schreckenbach G, Seldenthuis JS, Seth M, Snijders JG, and Sol. ADF2017, SCM, Theoretical Chemistry, Vrije Universiteit, Amsterdam, The Netherlands, <https://www.scm.com/>.

Figure 1. Identification of 15d-PGJ₂ as a PPAR δ agonist.

(A) 15d-PGJ₂ and GW0742 (positive control) activated PPAR δ in a PPAR δ -specific reporter assay. PPAR δ activity is represented as the mean luminescence signal-to-background (S/B) $\pm$ SD with $n = 3$. The results were reproduced independently at least two times. (B) The PPRE-binding activity of recombinant PPAR δ as well as NHBE cell nuclear extracts was increased by 15d-PGJ₂. PPAR δ -specific antibody caused supershift of 15d-PGJ₂-induced PPAR δ -PPRE complex. The nonspecific oligonucleotides showed no shift with the nuclear extracts or PPAR δ with or without 15d-PGJ₂ (1 μM). The results were reproduced independently at least two times. A representative image is shown. (C) 15d-PGJ₂ (1 μM) increased binding of PPAR δ to the PPRE-containing DNA. The data are expressed as the mean $\pm$ SD with $n = 3$, and the results reproduced at least two times independently; ** $p < 0.01$.

Figure 2. Covalent interaction between 15d-PGJ₂ and PPAR δ .

Recombinant human PPAR δ was incubated with varying concentrations of 15d-PGJ₂-Biotin in the presence or absence of unlabeled 15d-PGJ₂, the noncovalent ligand CAY10410, or DTT. Binding was assessed by Western blotting. 15d-PGJ₂-Biotin bound to PPAR δ in a dose-dependent manner. Unlabeled 15d-PGJ₂ competed with the 15d-PGJ₂-Biotin-PPAR δ interaction whereas CAY10410 failed to do so. DTT completely abolished complex formation. The results were reproduced independently at least two times. A representative image is shown.

Protein Sample	NHBE Cell Lysate								PPAR δ		PPAR δ LBD			
15d-PGJ ₂ -Biotin (μ M)	–	1	5	10	–	1	5	10	10	–	10	10	–	10
Maleimide (μ M)	–	–	–	–	1	1	1	1	–	1	1	–	1	1

15d-PGJ₂-Biotin

Maleimide

Merge

Figure 3. 15d-PGJ₂ interacts with Cys of PPAR δ .

NHBE cell extracts, recombinant PPAR δ proteins, and recombinant, isolated PPAR δ LBD bound to 15d-PGJ₂-Biotin (top panel). Similarly, they interacted with IRDye 800CW maleimide via the free Cys (middle panel). When both electrophiles were present, 15d-PGJ₂-Biotin competed for the reactive Cys in the protein samples and thereby reduced the maleimide signal. The bottom panel shows a merged image of the top (red) and the middle (green) panels. The results were reproduced at least two times independently. A representative image is shown.

Figure 4. Molecular modeling and docking of the 15d-PGJ₂-PPAR δ complex.

(A) Molecular model of 15d-PGJ₂ (yellow) bound to PPAR δ 's active site. (B-D) Stereo views of the binding site containing 15d-PGJ₂ and the PPAR δ residues that interact and stabilize the ligand from the selected docking pose; the presence of a covalent interaction between 15d-PGJ₂ and Cys249 of PPAR δ WT is shown in covalent docking (B). Such interaction is absent in non-covalent docking (C) as well as with Ser249 of the PPAR δ C249S mutant in covalent docking (D). PPAR δ is shown as a colored ribbon with helices indicated and with a hydrogen bond donor-acceptor surface. 15d-PGJ₂ is shown as a red stick model with interacting amino acids represented as scaled-sticks. The covalent bond between 15d-PGJ₂ and Cys249 is indicated by a red arrow (B); its absence is indicated by

pink arrows (C, D). Hydrogen bonds are indicated by orange arrows. Figures were generated with Discovery Studio software.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 5. 15d-PGJ₂ alkylates PPAR δ specifically at Cys249 and enhances its activity.

(A) 15d-PGJ₂-Biotin bound covalently to PPAR δ WT but not the PPAR δ C249S mutant (top panel, 15d-PGJ₂-Biotin (10 μ M) in red; middle panel, PPAR δ in green). DTT abolished the 15d-PGJ₂-Biotin-PPAR δ interaction. The bottom panel shows a merged image of the top (red) and the middle (green) panels. The results were reproduced independently at least two times. A representative image is shown. (B) The C249S mutation reduced 15d-PGJ₂-stimulated PPAR δ binding to the double-stranded PPRE-containing DNA. The results were reproduced independently at least two times; ($p < 0.001$ mutant vs. WT). The data are expressed as the mean $\pm$ SD with $n = 3$, and the results reproduced at least two times independently.

Figure 6. Frontier MOs of the 15d-PGJ₂-PPAR δ complex.

MOs showing (A) the HOMO of PPAR δ Cys249-S γ , (B) the LUMO of 15d-PGJ₂, and (C) the HOMO–LUMO overlap of the 15d-PGJ₂-PPAR δ complex generated based on DFT and QIATM calculations, with an isosurface 0.03. Figures were generated with ADF software.

Figure 7. A proposed mechanism for the 1,4 addition between 15d-PGJ $_2$ and PPAR δ Cys249. The reaction starts with the nucleophilic attack of 15d-PGJ $_2$'s endocyclic β -carbon (C $_{\beta}$) by PPAR δ 's Cys249-S $_{\gamma}$. Proton transfer and tautomerization follow and lead to the covalent bonding between 15d-PGJ $_2$ and PPAR δ Cys249.

Figure 8. Reaction coordinate of the 1,4 addition between 15d-PGJ₂ and a Cys. Progression of the reaction, calculated with the semiempirical parametric method (PM6) and COSMO, is shown against the potential energy (red; the left y-axis) as well as the bond distance (green; the right y-axis) associated with the interaction. Structures and MOs of the reactant, the transition, and the product states are shown. The reaction starts with a distance of 3.296 Å between the reactants, proceeds through the transition state with a 2.064 Å distance, and finally forms a covalent bond of 1.888 Å (also shown in Movie S4).

Table 1.Calculated quantum chemical parameters of 15d-PGJ₂, CAY10410, and PPAR δ Cys, and Ser.

Molecule	η (eV)	σ (eV ⁻¹)	μ (eV)	ω (eV)
15d-PGJ ₂	4.375	0.229	-5.005	<u>2.863</u>
CAY10410	2.570	0.389	-2.700	1.418
Cysteine	4.710	0.212	-5.120	—
Serine	5.265	0.190	-4.705	—

Table 2.

Calculated selectivity parameters and atomic charges of 15d-PGJ₂, CAY10410, and PPAR δ Cys, and Ser.

Molecule	Atom	Fukui		DD	Mulliken Charge	Hirshfeld Charge
		Nucleophilic	Electrophilic			
15d-PGJ ₂	C ₉	0.119	<u>0.613</u>	-0.494	0.058	-0.039
	C ₁₃	0.188	0.007	0.181	-0.278	-0.031
CAY10410	C ₉	0.006	0.004	0.002	-0.573	-0.080
	C ₁₃	0.256	0.340	-0.084	-0.469	0.007
Cysteine	S _{γ}	<u>0.044</u>	0.012	0.033	-0.034	-0.036
Serine	O _{γ}	0.013	0.009	0.004	-0.686	-0.228