Published in final edited form as:

Cardiovasc Toxicol. 2019 April ; 19(2): 168–177. doi:10.1007/s12012-018-9487-6.

Airway exposure to modified multi-walled carbon nanotubes perturbs cardiovascular adenosinergic signaling in mice

Leslie C. Thompson¹, Nicole L. Sheehan¹, Dianne M. Walters¹, Robert M. Lust¹, Jared M. Brown², and Christopher J. Wingard³

¹Department of Physiology, Brody School of Medicine at East Carolina University, Greenville, NC, USA

²Department of Pharmaceutical Sciences, Skaggs School of Pharmacy and Pharmaceutical Science, University of Colorado, Aurora, CO, USA

³College of Health Professions, School of Movement and Rehabilitation Sciences, Bellarmine University, Louisville, KY, USA

Abstract

The broad list of commercial applications for multi-walled carbon nanotubes (MWCNT) can be further expanded with the addition of various surface chemistry modifications. For example, standard commercial grade MWCNT (C-grade) can be carboxylated (COOH) or nitrogen-doped (N-doped) to suite specific utilities. We previously reported dose-dependent expansions of cardiac ischemia/reperfusion (I/R) injury, 24 hours after intratracheal instillation of C-grade, COOH, or N-doped MWCNT in mice. Here we have tested the hypothesis that airway exposure to MWCNT perturbs cardiovascular adenosinergic signaling, which could contribute to exacerbation of cardiac I/R injury. One hundred microliters of Vehicle or identical suspension volumes containing 100 µg of C-grade, COOH, or N-doped MWCNT were instilled into the trachea of CD-1 ICR mice. One day later we measured cyclic adenosine monophosphate (cAMP) concentrations in cardiac tissue and evaluated arterial adenosinergic smooth muscle signaling mechanisms related to nitric oxide synthase (NOS) and cyclooxygenase (COX) in isolated aortic tissue. We also verified cardiac I/R injury expansion and examined both lung histology and bronchoalveolar lavage fluid cellularity in MWCNT exposed mice. Myocardial cAMP concentrations were reduced ($p < 0.05$) in the C-grade group by 17.4% and N-doped group by 13.7% compared to the Vehicle group. Curve fits to aortic ring 2-Cl-Adenosine concentration responses were significantly greater in the MWCNT groups vs. the Vehicle group. Aortic constrictor responses were more pronounced with NOS inhibition and were abolished with COX inhibition. These findings indicate that addition of functional chemical

Corresponding Author: Dr. Christopher J. Wingard, College of Health Professions, School of Movement and Rehabilitation Sciences, Bellarmine University, Louisville, KY, USA, 2001 Newburg Road, Louisville, KY 40205, USA, cwingard@bellarmine.edu.

Conflict of Interest:

The authors report no conflicts of interests related to the research reported in this manuscript. The conclusions reported in the manuscript are those of the authors and do not necessarily reflect those of the National Institute of Environment Health Sciences, East Carolina University, Bellarmine University, or the University of Colorado at Denver.

Statement on the Welfare of Animals:

All applicable international, national, and/or institutional guidelines for the care and use of animals were followed. All procedures performed in studies involving animals were approved and in accordance with the ethical standards of the Institutional Animal Care and Use Committee at East Carolina University (Greenville, NC, USA).

moieties on the surface of MWCNT may alter the biological responses to exposure by influencing cardiovascular adenosinergic signaling and promoting cardiac injury.

Keywords

Multi-walled carbon nanotubes; cardiac ischemia/reperfusion injury; adenosine; cAMP

INTRODUCTION

Multi-walled carbon nanotubes (MWCNT) have dynamic applicability across industries and present potential for occupational exposure by inhalation during production and handling [1]. The physicochemical properties of MWCNT have been shown to dictate biological responses to airway exposures [2], which raises questions regarding what MWCNT modifications can contribute to these observations. Standard commercial grade MWCNT (C-grade) have high chemical and thermal stability. In order to increase MWCNT electrothermal conduction, the surface of MWCNT can be doped with nitrogen moieties (N-doped) [3]. MWCNT can be functionalized with carboxylic acid groups (COOH) to enhance solubility [4]. We have previously reported that intratracheal instillation of C-grade, COOH, and N-doped MWCNT caused a dose-dependent and time-dependent expansion of cardiac ischemia/reperfusion (I/R) injury in C57BL/6 mice as early as 24 hours after exposure [5]. However, it remains unclear what physiological mechanisms contributed to the expansion of cardiac I/R.

Adenosinergic signaling regulates an array of cardiovascular functions [6]. Reduced concentrations of 3',5'-cyclic adenosine monophosphate (cAMP) have been implicated in cardiovascular pathologies [7, 8] and increased cAMP concentration provides cardioprotection against I/R injury [9, 10]. Generally, this protection is thought to occur because cAMP activates protein kinase A signaling cascades and influences mitochondrial permeability transition pore, ROS generation, and various channels and cell types in cardiovascular tissues [11, 12]. Cardiac cAMP may also be linked to adenosinergic dysfunction because cAMP is converted to adenosine via the cAMP-adenosine pathway [13]. Adenosine itself is an important autocrine-paracrine signaling agent in both heart [14] and vascular tissue [15], and is usually regarded as tissue protective in acute settings [16, 17]. The local action of adenosine allows for the titration of local blood flow during changes in autonomic tone. For example, adenosine has been documented to attenuate α -adrenergic vascular tone [18]. Adenosine regulates vascular tone via activation of both nitric oxide (NO) [19] and cyclooxygenase (COX) signaling [20], and is thought to be released during ischemia [21, 22]. In contrast, adenosine has also been reported to augment aortic constrictor responses mediated through selective adenosine receptors and endothelial cell signaling [23-25]. Thus, altered vascular responsiveness to adenosine associated with MWCNT exposure could negatively impact cardiac I/R injury.

The purpose of this study was to test the influence of modified MWCNT exposure on cardiovascular adenosinergic systems. We hypothesized that pulmonary exposure to MWCNT would decrease cAMP concentrations in the heart and alter adenosinergic vascular

responses through changes in NO and COX signaling. We selected the MWCNT dose based on similar levels of pulmonary injury between MWCNT types, one day after exposure in C57BL/6 mice in our previous study [5]. Here we have assessed pulmonary inflammation, verified cardiac I/R injury, measured heart and lung cAMP concentrations, and interrogated adenosinergic responsiveness in vascular tissue one day after intratracheal instillation of MWCNT in CD-1 ICR mice.

MATERIALS AND METHODS

Animals:

Male CD-1 ICR mice were purchased from Charles River (Raleigh NC, USA) at 4 to 6 weeks of age. After arrival at East Carolina University's Department of Comparative Medicine the mice were allowed to acclimate for at least one week prior to experimental procedures. Mice were housed in temperature controlled rooms ($23 \pm 1^\circ\text{C}$) with 12-hour light/dark cycles and provided access to standard laboratory chow and fresh water *ad libitum*. The use of mice in this study complied with protocols approved by the Institutional Animal Care and Use Committee at East Carolina University (Greenville, NC, USA).

MWCNT Characterization:

In this study, unmodified commercial grade (C-grade), acid-washed carboxylated (COOH), and nitrogen-doped (N-doped) MWCNT were examined. All MWCNT were derived by chemical vapor-deposition and kindly provided by NanoTecLabs, Yadkinville NC, USA. The MWCNT used in this study have been characterized previously in both dry form and in suspension and are free of endotoxin contamination [5].

MWCNT Instillation:

Experimental suspensions of C-grade, COOH, and N-doped MWCNT were prepared at 1 mg/mL in sterile 10% pulmonary surfactant/saline using Infasurf™ (ONY, Inc., Amherst NY, USA). Infasurf™ contains 35 mg of bovine phospholipids, including 16 mg disaturated phosphatidylcholine; 0.7 mg bovine proteins, including 0.44 mg hydrophobic surfactant-associated protein C and 0.26 mg hydrophobic surfactant-associated protein B (<http://www.infasurf.com/about-infasurf/#composition>). Twenty-four hours prior to cardiopulmonary assessments the suspensions were sonicated in a bath sonicator for twenty minutes to generate approximately 10,000 J. Immediately after sonicating the MWCNT, mice were anesthetized with Isoflurane and then intratracheally instilled with 100 μL suspensions containing 0 (Vehicle) or 100 μg of either C-grade, COOH, or N-doped MWCNT.

Heart, lung and aorta collection:

In the cohort of mice not used for ischemia/reperfusion experiments, mice were anesthetized with an intraperitoneal injection of ketamine/xylazine (90/10 mg/kg, respectively) and exsanguinated by cutting the inferior vena cava. Hearts and thoracic aortas were resected and placed on ice. Aortas were placed in cold physiological saline solution (PSS) containing (in mM) 140 NaCl, 5.0 KCl, 1.6 CaCl_2 , 1.2 MgSO_4 , 1.2 3-[N-morpholino]-propane sulfonic acid (MOPS), 5.6 D-glucose, and 0.02 EDTA for myographic assessments. Hearts were snap

frozen in liquid nitrogen and stored at -80°C until analyzed. The intact lung tissue was either used for bronchoalveolar lavage studies, histological fixation, or was collected for tissue analysis, in which case it was snap frozen in liquid nitrogen and stored at -80°C .

Bronchoalveolar Lavage (BAL) and Lung Histology:

Right lungs were lavaged *in situ* four times with 26.25 mL/kg of cold Hanks balanced salt solution (HBSS) as previously described [26]. Each of the four individual lavage samples were collected separately and centrifuged at 4°C for 10 minutes at $1000 \times g$. The supernatant from the first lavage sample was used for protein quantification. Each cell pellet was combined and re-suspended in 1 mL of HBSS to derive total cell counts using a Cellometer® Auto X4 (Nexcelcom Biosciences, LLC., Lawrence, MA, USA). A volume of cell suspension containing approximately 20,000 cells was centrifuged onto glass slides using a Cytospin IV (Shandon Scientific Ltd., Cheshire, UK) and stained with three-step hematology stain (Richard Allan Scientific, Kalamazoo, MI, USA). Differential cell counts were derived by microscopically evaluating 300 cells per slide based on hematological staining and morphology. For each cell type per slide, the percentage of 300 cells was calculated and multiplied by the total cell count for data reporting. Unlavaged left lungs were inflated with 10 % neutral buffered formalin at 25 cm of H_2O and incubated at room temperature for 24 - 72 hrs. Fixed lung tissue was cut, processed, embedded in paraffin, sectioned 5 μm thick, mounted on slides, and stained with hematoxylin and eosin (H&E).

Ischemia/Reperfusion (I/R) Injury:

Twenty-four hours after instillation, one cohort of mice was subjected to an experimental cardiac I/R protocol as previously described [5]. In short, mice were anesthetized with an intraperitoneal injection of ketamine/xylazine (90/10 mg/kg, respectively), then intubated, and mechanically ventilated. A midline parasternal thoracotomy was conducted to access the left anterior descending coronary artery (LAD), which was then ligated with a reversible tourniquet to induce ischemia in the zone of tissue distal to the occlusion (*i.e.* zone at risk; ZAR). After 20 minutes the ligature was reversed and the ZAR was allowed to reperfuse for 2 hours, followed by exsanguination of the inferior vena cava. In order to demarcate the ZAR, the LAD was religated, the aorta was cannulated, and Evans blue dye was perfused in a retrograde fashion to stain myocardial tissue not in the ZAR. The left ventricle was then excised, serially sectioned, and incubated in a 1 % triphenyltetrazolium chloride solution to demarcate the infarcted tissue within the ZAR. Triphenyltetrazolium is enzymatically converted into a brick red substance in viable tissue. Infarcted tissue remains pale. Differential staining was analyzed by computer planimetry and infarction is reported as percent of the ZAR.

Adenosine 3', 5'-Cyclic Monophosphate (cAMP) ELISA:

An enzyme-linked immunosorbent assay (ELISA) for cAMP (cat# ADI-900-163, Enzo Life Sciences, Farmingdale, NY, USA) was used to determine the cAMP concentrations present in heart and lung tissues. Portions of heart and lung were individually weighed and homogenized in 0.1 N HCl (HCl to tissue mass ratio) using a Mini bead beater (Cole Parmer, Vernon Hills, IL, USA). Protein concentrations were analyzed using a Bradford assay, using a Synergy HT plate reader and Gen5 software (BioTek Instruments, Winooski,

VT, USA). Heart homogenates were further diluted 1:50 and lung homogenates were further diluted 1:100 using 0.1 N HCl. Using an option provided with the assay, we performed an acetylation step to increase the sensitivity of the ELISA by 10-fold. Then the assay was prepared per the manufacturer's instructions and the experimental samples were plated in duplicate. Optical densities were read at 405 nm using a Synergy HT plate reader and Gen5 software (BioTek Instruments). In assays of heart homogenates optical densities below 0.3 were taken as 0 pmol/mL. In assays of lung homogenates optical densities below 0.27 were taken as 0 pmol/mL. Concentration values for cAMP were then normalized to mg of total protein for data reporting.

Adenosinergic modulation of vascular tone in isolated aortas:

Thoracic aortas were cleaned of surrounding perivascular tissue, cut into 2 mm length rings, and mounted into a DMT 610M multi-channel myograph system (Ann Arbor, MI, USA). Aortic rings were maintained in 37°C PSS (see above description), bubbled with medical grade breathing air, stretched to a tension yielding 20 mN of force on the transducer, and allowed to relax for 15 minutes. Then each ring was washed with fresh PSS and re-stretched to a 20 mN force readout. This process was repeated until rings maintained a resting tension at 20 mN of force. Tension was then quickly released to a passive tension yielding 10 mN of force and rings were allowed to equilibrate for 10 minutes. Aortic rings were assessed for viability via K⁺-depolarization using 109 mM K⁺PSS (PSS with equal molar substitution of K⁺ for Na⁺) and basic endothelial function using 1 μM phenylephrine (PE), an α-adrenergic agonist (cat# P6126, Sigma-Aldrich, St. Louis, MO, USA), to contract aortic smooth muscle and 3.0 μM acetylcholine (cat# A6625, Sigma-Aldrich) to stimulate endothelial-dependent relaxation. After an additional 3 × 10-minute washout periods, aortic rings were incubated for 20 minutes with no inhibitor, 10 μM indomethacin – a general COX inhibitor (cat# I7378, Sigma-Aldrich), or 1 mM Nω-Nitro-L-Arginine Methyl Ester Hydrochloride (L-NAME) – a general NO synthase inhibitor (Cat# N5751, Sigma-Aldrich). Aortic rings were then stimulated with 1 μM PE for 5 minutes to allow active stress development to stabilize, followed by cumulative additions of 2-Cl-Adenosine – stable purinergic adenosine analog (Cat# 3136, Tocris Bioscience/R&D Systems, Inc., Minneapolis, MN, USA) every 5 minutes from 0.1 nM - 30.0 μM.

Statistics:

Data are expressed as mean ± SEM. Differences were considered significantly different when $p < 0.05$. BAL fluid protein, BAL cell differentials, cAMP concentrations, and maximal aortic responses were analyzed by ANOVA with Tukey's post-test. In Figure 4, mean data points of concentration-response curves for 2-Cl-adenosine were analyzed by Repeated Measures ANOVA (marked with *for $p < 0.05$ vs. Vehicle). Additionally, a non-linear curve fit was applied to the response profile and analyzed for statistical difference of the resultant fitted lines. This approach allows for the interrogating of complex responses often associated with second messenger systems [27].

RESULTS

BAL fluid analysis:

The results from fluid protein and cell differential analyses from BAL samples collected 24 hours following MWCNT instillation are provided in Table 1. The following results were significantly different than the Vehicle group: (i) mice instilled with C-grade were found to have significantly elevated protein concentration in BALF, increased cellularity, and neutrophilia; (ii) mice instilled with COOH were found to have eosinophilia; (iii) mice instilled with N-doped were found to have significantly elevated cellularity, neutrophilia, and eosinophilia. The extent of eosinophilia found in mice instilled with COOH was significantly different from that of the C-grade group. The extent of neutrophilia found in mice instilled with N-doped was also significantly different from the COOH group.

Lung histology:

Representative histological images of the left lung, 24 hours after MWCNT instillation are shown in Figure 1. Histological analysis had consistent results with BAL analysis. When taking histological lung samples from the Vehicle group (Figure 1A) as a baseline comparison, lung samples from the C-grade group showed some increased cellularity and wall thickening (Figure 1B). Histological lung samples from the COOH group appeared to fall between the Vehicle and C-grade groups in terms of cellularity and wall thickening (Figure 1C). When compared to Vehicle, C-grade, and COOH, the N-doped group appeared to have the highest degree of pulmonary cellularity and wall thickening in response to MWCNT instillation (Figure 1D).

Cardiac I/R injury:

We verified that the CD-1 ICR mouse model would respond to MWCNT instillation with expansion of cardiac I/R injury, as we have previously reported for C57BL/6 mice [5]. The sizes of myocardial infarctions generated by cardiac I/R, 24 hours following MWCNT exposure are presented in Figure 2 as a percent of the zone at risk for infarction. Myocardial infarct size was significantly larger following I/R in the C-grade, COOH, and N-doped groups when compared to the Vehicle group by 33%, 60%, and 87%, respectively. Myocardial infarct size in the COOH and N-doped groups was also significantly larger than the infarction in the C-grade group by 21% and 41%, respectively. Myocardial infarct size in the N-doped group was also significantly larger by 16% than the infarct size in the COOH group.

Heart and lung cAMP concentrations:

Data from cAMP measurements in heart and right lung tissues collected 24 hours after MWCNT instillation are reported in Figure 3. Analysis of heart tissue revealed that cAMP concentrations were significantly decreased in the C-grade and N-doped groups (17.4% and 13.7% respectively) when compared to the Vehicle group (Figure 3A). Heart cAMP levels were not different between the COOH and Vehicle groups. We also examined lung tissue for changes in cAMP following MWCNT instillation but found no significant differences between groups (Figure 3B).

Adenosinergic modulation of aortic tone:

Aortic adenosinergic modulation of α -adrenergic (phenylephrine - PE) force generation is presented in Figure 4. We conducted these experiments either in the presence or absence of the general COX inhibitor indomethacin (10 μ M) or the NO synthase inhibitor L-NAME (1 mM). Use of a best-fit comparison of non-linear fits of the cumulative 2-Cl-adenosine response curves revealed significant differences between the MWCNT groups and the Vehicle group when no inhibitors were present (Figure 4A). The same analysis of best fit comparison of the cumulative response curves revealed a significant difference between the N-doped and C-grade groups with no inhibition. When aortic 2-Cl-adenosine best fit responses lines were assessed in the presence of 10 μ M indomethacin no significant differences were detected between any groups (Figure 4B). When aortic 2-Cl-adenosine best fit responses lines were compared in the presence of 1 mM L-NAME, the fits of the COOH and C-grade groups were significantly different from the Vehicle group (Figure 4C). 2-Cl-adenosine responses in the COOH group were also significantly different from the Vehicle group when compared by Repeated Measures ANOVA. Aortic 2-Cl-adenosine responses of the N-doped group were not significantly different from Vehicle group.

Peak aortic responses:

The peak aortic constrictor responses to PE and relaxation response to 2-Cl-adenosine are reported in Figure 5. Despite some separation in the mean constrictor stress responses generated by PE stimulation, there were no significant differences between MWCNT treatment groups within any of the inhibitor groups (Figure 5A). We did find eNOS inhibition with L-NAME enhanced the PE stress responses in the Vehicle and COOH groups when compared to PE stress when no inhibitor was present. Maximum aortic relaxation achieved with 30 μ M 2-Cl-adenosine stimulation was not different between groups when no inhibitors were present or in the presence of the COX inhibitor indomethacin (Figure 5B). However, maximum 2-Cl-adenosine relaxation was blunted by 80% in the N-doped group compared to the Vehicle group when aortas were incubated with 1 mM L-NAME a NOS inhibitor.

DISCUSSION

The key finding in this study is that pulmonary exposure to modified MWCNT influences cardiovascular adenosinergic responses in the CD-1 ICR mouse model promoting constriction in the face of adrenergic stimulation. In particular, intratracheal instillation of modified MWCNT diminishes cAMP in myocardial tissue and adapts adenosine modulation of aortic smooth muscle tone via imbalanced NO and COX mechanisms. We found that these endpoints were uniquely impacted depending on exposure to either C-grade, COOH, and N-doped MWCNT. Moreover, responses in the lung, as well as cardiac I/R injury are also dependent on the type of MWCNT encountered during exposure. The CD-1 mouse strain has been used for receptor subtype knockout studies of adenosine receptor function [11, 28] and allows the comparison of cardiac I/R injury after MWCNT instillation between this study and those we previously reported in the C57BL/6 model [5].

Accumulating evidence as recently reviewed by Abukabda et al. and Stone et al. [29 and 30] suggests that engineered nanomaterials detrimentally impacts cardiovascular system health by a diversity of postulated modes of action. Many of the modes are borne out of the extensive work reported on ambient particulate matter exposures [31] and include: central nervous system effects, pulmonary and systemic inflammation responses and possible direct particle effects. A common target for the all of these modes include the endothelium. This is a tissue critical in the regulation of vascular function to all organs by autocrine, paracrine and hormonal signaling cascades. The endothelium may be sensitive to the physiochemical distinction of the MWCNT impacting a NO/reactive oxygen species imbalance and create a tissue environment that promotes injury through thrombosis, vascular dysfunction and autonomic dysreflexia [reviewed in 29 - 31].

Exposure to MWCNT produces unique airway responses and I/R-induced myocardial infarct size depending on the modification, which is consistent with our previous report [5]. In both these studies, infarct sizes graded in size with smallest to largest in the order of: Vehicle < C-grade < COOH, with N-doped. In the present study, mice instilled with C-grade had significant increases in BALF protein, cellularity, and neutrophilia compared to the Vehicle groups. Both studies showed that N-doped exposure increased cellularity and neutrophilia compared to the Vehicle instillation. This type of cellularity/neutrophilia has also been demonstrated with MWCNT exposure in mice [32] and rats [33]. One contrast between our two studies were that CD-1 mice did not present with increased BALF protein concentration after N-doped exposure, as was seen with the C57BL/6 mice [5]. We did find eosinophilia in the COOH and N-doped groups, which we have reported in rats 24 hours after instillation of C-grade MWCNT [34] and 60-carbon fullerenes [35].

One of the key findings here is that exposure to modified MWCNT decreases cardiac cAMP concentrations. More specifically, we found that cardiac cAMP was significantly lower in the C-grade and N-doped groups when compared to the Vehicle group, with the mean value of cAMP in the COOH group also being lower (but not statistically different) than that of the Vehicle group. Decreased tissue cAMP concentration can result from excess hydrolysis of cAMP by phosphodiesterases, decreased synthesis by adenylate cyclase, or by extrusion from cardiomyocytes by ATP-binding cassette transporters [36]. One important function of cAMP is activation of protein kinase A, which has reported involvement in I/R cardioprotection [37, 38]. The myocyte cyclic nucleotide system is also important for appropriate heart responses to functional regulation by the autonomic nervous system, endocrine system, and autocrine/paracrine systems via G-protein coupled receptor activation [39]. Thus, reductions in cAMP concentrations after MWCNT exposure may contribute to cardiac injuries and blunting of cAMP-dependent heart function.

We also found adaptations in aortic responsiveness to adenosine following exposure to all three types of MWCNT but the responses were unique depending on the surface modifications. Best-fit analysis of the fitted curves revealed that that N-doped and COOH exposure augmented cumulative response profile to 2-Cl-adenosine. These alterations were prevented by treating aortic rings with indomethacin, a general COX inhibitor. However, NOS inhibition produced enhanced aortic contractile and relaxation responses to 2-Cl-adenosine in the COOH group but blunted the relaxation responses in the N-doped group.

Extracellular adenosine is an important arm of the cardiovascular adenosinergic system, which includes the regulation of heart rate, myocardial contractility, coronary flow, and vascular tone [6], so one may suggest alterations in adenosinergic signaling after MWCNT exposure may impact functions beyond the vascular system. Similar alterations in vascular NO and COX signaling following MWCNT instillation have been reported, including blunted endothelium-dependent dilation [40-42] and enhanced vasoconstriction via COX [28]. While other mouse models have reported an augmented aortic constriction to adenosine in the face of modulated NO/COX signaling of the endothelium [23-25]. Given that myocardial ischemia can cause the release of adenosine [22, 43], altered vascular responses to extracellular adenosine may have played a role in the expansion of cardiac I/R injury after exposure to the COOH and N-doped forms of MWCNT. When considered together, these findings indicate that MWCNT exposure can alter adenosinergic signaling and that nitrogen-doping and carboxylation modifies these responses, which may make predicting health outcomes of exposure more difficult.

Before drawing final conclusions from our findings some limitations must be acknowledged. First, we have used a high dose of MWCNT for instillation in this study. We have previously reported dose-response and time-course findings associated with the MWCNT used in this study [5]. The purpose of our present study was to investigate the impact of MWCNT exposure on adenosinergic cardiovascular endpoints and further determine if the responses were unique based on form of modified MWCNT. We selected the high dose of MWCNT used in our previous study to interrogate a mechanism of dysfunction, future efforts need to be undertaken to determine any dose- and time-dependent aspects of altered adenosinergic signaling contributing to the apparent cardiovascular detriment. We have also examined vascular responses in aortic tissue as general assessments of arterial endothelial and smooth muscle function as this tissue is routinely used to support studies examining function in other arteries. However, it will be informative to examine adenosine responsiveness in small resistive arteries. This question will be particularly relevant in assessing the role of MWCNT exposure on the adenosinergic signaling in coronary vasculature due to its prominent role in protecting the myocardium from ischemic damage.

In conclusion, modified versions of MWCNT can uniquely alter cAMP concentrations in the heart and influence aortic adenosinergic responsiveness via dependence on NO and COX signaling. The impact of MWCNT exposure on cAMP concentrations in the heart could possibly disrupt other signaling mechanisms that rely on cyclic nucleotides. The alterations in vascular adenosinergic signaling following exposure to surface-modified MWCNT could cause circulatory supply and demand mismatching. The combination of impacted adenosinergic systems after exposure to C-grade and modified MWCNT could narrow cardiovascular homeostasis to a degree that episodes like cardiac I/R may contribute to the observed exacerbated injury. Given that the utility of MWCNT is expanding across industries, a more thorough understanding of physiological consequences occurring because of exposure is relevant.

ACKNOWLEDGEMENTS

We would like to thank Cathy Stang and Alvin Tsang for help with tissue collection and vascular data acquisition. We thank Dr. Walter Klein of ONY, Inc. for donating the Infasurf™, Dr. Benjamin Harrison and Dr. Richard Czerw of NanoTechLabs, Inc. for providing the MWCNT used in this study.

Funding:

This research was supported by East Carolina University and funded through the National Institutes of Environmental Health Sciences: NIH R01 ES016246 (CJW) and U19 ES019525 (JMB and CJW).

REFERENCES

1. Kuijpers E, Bekker C, Fransman W, Brouwer D, Tromp P, Vlaanderen J, Godderis L, Hoet P, Lan Q, Silverman D, Vermeulen R and Pronk A (2016). Occupational Exposure to Multi-Walled Carbon Nanotubes During Commercial Production Synthesis and Handling. *Ann Occup Hyg* 60: 305–317. [PubMed: 26613611]
2. Poulsen SS, Knudsen KB, Jackson P, Weydahl IE, Saber AT, Wallin H and Vogel U (2017). Multi-walled carbon nanotube-physicochemical properties predict the systemic acute phase response following pulmonary exposure in mice. *PloS One* 12: e0174167. [PubMed: 28380028]
3. Ayala P, Arenal R, Rummeli M, Rubio A and Pichler T (2010). The doping of carbon nanotubes with nitrogen and their potential applications. *Carbon* 48: 575–586.
4. Min C, Zhang Q, Shen C, Liu D, Shen X, Song H, Li S, Xu D, Lin X and Zhang K (2017). Graphene oxide/carboxyl-functionalized multi-walled carbon nanotube hybrids: powerful additives for water-based lubrication. *RSC Advances* 7: 32574–32580.
5. Urankar RN, Lust RM, Mann E, Katwa P, Wang X, Podila R, Hilderbrand SC, Harrison BS, Chen P, Ke PC, Rao AM, Brown JM and Wingard CJ (2012). Expansion of cardiac ischemia/reperfusion injury after instillation of three forms of multi-walled carbon nanotubes. *Part Fibre Toxicol* 9: 38. [PubMed: 23072542]
6. Sousa JB and Diniz C (2017). The Adenosinergic System as a Therapeutic Target in the Vasculature: New Ligands and Challenges. *Molecules* (Basel, Switzerland) 22 DOI: 10.3390/molecules22050752
7. Sprenger JU, Perera RK, Steinbrecher JH, Lehnart SE, Maier LS, Hasenfuss G and Nikolaev VO (2015). In vivo model with targeted cAMP biosensor reveals changes in receptor-microdomain communication in cardiac disease. *Nat Commun* 6: 6965. [PubMed: 25917898]
8. Weber S, Zeller M, Guan K, Wunder F, Wagner M and El-Armouche A (2017). PDE2 at the crossway between cAMP and cGMP signalling in the heart. *Cell Signal* 38: 76–84. [PubMed: 28668721]
9. Khaliulin I, Bond M, James AF, Dyar Z, Amini R, Johnson JL and Suleiman MS (2017). Functional and cardioprotective effects of simultaneous and individual activation of protein kinase A and Epac. *Br J Pharmacol* 174: 438–453. [PubMed: 28071786]
10. Lochner A, Genade S, Tromp E, Podzuweit T and Moolman JA (1999). Ischemic preconditioning and the beta-adrenergic signal transduction pathway. *Circulation* 100: 958–966. [PubMed: 10468527]
11. Headrick JP, Ashton KJ, Rose'meyer RB and Peart JN (2013). Cardiovascular adenosine receptors: expression, actions and interactions. *Pharmacol Ther* 140: 92–111. [PubMed: 23764371]
12. Ladilov Y and Appukuttan A (2014). Role of soluble adenylyl cyclase in cell death and growth. *Biochim Biophys Acta* 1842: 2646–2655. [PubMed: 25010002]
13. Jackson EK, Mi Z and Dubey RK (2007). The extracellular cAMP-adenosine pathway significantly contributes to the in vivo production of adenosine. *J Pharmacol Exp Ther* 320: 117–123. [PubMed: 17028245]
14. Vecchio EA, White PJ and May LT (2017). Targeting Adenosine Receptors for the Treatment of Cardiac Fibrosis. *Front Pharmacol* 8: 243. [PubMed: 28529484]
15. Ho MF and Rose'Meyer RB (2013). Vascular adenosine receptors; potential clinical applications. *Curr Vasc Pharmacol* 11: 327–337. [PubMed: 22724462]

16. Antonioli L, Fornai M, Blandizzi C, Pacher P and Hasko G (2018). Adenosine signaling and the immune system: When a lot could be too much. *Immunol Lett* DOI: 10.1016/j.imlet.2018.04.006
17. Bowser JL, Lee JW, Yuan X and Eltzschig HK (2017). The hypoxia-adenosine link during inflammation. *J Appl Physiol* (Bethesda, Md. : 1985) 123: 1303–1320. [PubMed: 28798196]
18. Smits P, Lenders JW, Willemsen JJ and Thien T (1991). Adenosine attenuates the response to sympathetic stimuli in humans. *Hypertension* (Dallas, Tex : 1979) 18: 216–223. [PubMed: 1885230]
19. Lynch FM, Austin C, Heagerty AM and Izzard AS (2006). Adenosine and hypoxic dilation of rat coronary small arteries: roles of the ATP-sensitive potassium channel, endothelium, and nitric oxide. *Am J Physiol Heart Circ Physiol* 290: H1145–1150. [PubMed: 16243919]
20. Ansari HR, Nadeem A, Tilley SL and Mustafa SJ (2007). Involvement of COX-1 in A3 adenosine receptor-mediated contraction through endothelium in mice aorta. *Am J Physiol Heart Circ Physiol* 293: H3448–3455. [PubMed: 17921329]
21. Burnstock G and Pelleg A (2015). Cardiac purinergic signalling in health and disease. *Purinergic Signal* 11: 1–46. [PubMed: 25527177]
22. Fox AC, Reed GE, Glassman E, Kaltman AJ and Silk BB (1974). Release of adenosine from human hearts during angina induced by rapid atrial pacing. *J Clin Invest* 53: 1447–1457. [PubMed: 4825235]
23. Zhou Z, Sun C, Tilley SL, Mustafa SJ (2015). Mechanisms underlying uridine adenosine tetraphosphate-induced vascular contraction in mouse aorta: Role of thromboxane and purinergic receptors. *Vascul Pharmacol* 73:78–85. [PubMed: 25921923]
24. Ponnoth DS, Nadeem A, Tilley S, Mustafa SJ. (2010). Involvement of A1 adenosine receptors in altered vascular responses and inflammation in an allergic mouse model of asthma. *Am J Physiol Heart Circ Physiol* 299: H81–87. [PubMed: 20400685]
25. Hansen PB, Hristovska A, Wolff H, Vanhoutte P, Jensen BL, Bie P (2010). Uridine adenosine tetraphosphate affects contractility of mouse aorta and decreases blood pressure in conscious rats and mice. *Acta Physiol (Oxf)*. 200: 171–179. [PubMed: 20384597]
26. Wang X, Katwa P, Podila R, Chen P, Ke PC, Rao AM, Walters DM, Wingard CJ and Brown JM (2011). Multi-walled carbon nanotube instillation impairs pulmonary function in C57BL/6 mice. *Part Fibre Toxicol* 8: 24. [PubMed: 21851604]
27. Ludbrook J (1994). Repeated measurements and multiple comparisons in cardiovascular research. *Cardiovasc Res* 28: 303–311. [PubMed: 8174149]
28. Yaar R, Jones MR, Chen JF and Ravid K (2005). Animal models for the study of adenosine receptor function. *J Cell Physiol* 202: 9–20. [PubMed: 15389588]
29. Abukabda AB, Stapleton PA, Nurkiewicz TR (2016). Metal Nanomaterial Toxicity Variations Within the Vascular System. *Curr Environ Health Rep* 3: 379–391. [PubMed: 27686080]
30. Stone V, Miller MR, Clift MJD, Elder A, Mills NL, Møller P, Schins RPF, Vogel U, Kreyling WG, Alstrup Jensen K, Kuhlbusch TAJ, Schwarze PE, Hoet P, Pietroiusti A, De Vizcaya-Ruiz A, Baeza-Squiban A, Teixeira JP, Tran CL, Cassee FR (2017). Nanomaterials Versus Ambient Ultrafine Particles: An Opportunity to Exchange Toxicology Knowledge. *Environ Health Perspect* 125(10):106002. [PubMed: 29017987]
31. Brook RD, Rajagopalan S, Pope CA 3rd, Brook JR, Bhatnagar A, Diez-Roux AV, Holguin F, Hong Y, Luepker RV, Mittleman MA, Peters A, Siscovick D, Smith SC Jr, Whitsel L, Kaufman JD (2010). Particulate matter air pollution and cardiovascular disease: An update to the scientific statement from the American Heart Association. *Circulation* 121:2331–2378. [PubMed: 20458016]
32. Cartwright MM, Schmuck SC, Corredor C, Wang B, Scoville DK, Chisholm CR, Wilkerson HW, Afsharinejad Z, Bammler TK, Posner JD, Shutthanandan V, Baer DR, Mitra S, Altemeier WA and Kavanagh TJ (2016). The pulmonary inflammatory response to multiwalled carbon nanotubes is influenced by gender and glutathione synthesis. *Redox Biol* 9: 264–275. [PubMed: 27596734]
33. Bonner JC, Silva RM, Taylor AJ, Brown JM, Hilderbrand SC, Castranova V, Porter D, Elder A, Oberdorster G, Harkema JR, Bramble LA, Kavanagh TJ, Botta D, Nel A and Pinkerton KE (2013). Interlaboratory evaluation of rodent pulmonary responses to engineered nanomaterials: the NIEHS Nano GO Consortium. *Environ Health Perspect* 121: 676–682. [PubMed: 23649427]

34. Thompson LC, Frasier CR, Sloan RC, Mann EE, Harrison BS, Brown JM, Brown DA and Wingard CJ (2014). Pulmonary instillation of multi-walled carbon nanotubes promotes coronary vasoconstriction and exacerbates injury in isolated hearts. *Nanotoxicology* 8:38–49. [PubMed: 23102262]
35. Thompson LC, Urankar RN, Holland NA, Vidanapathirana AK, Pitzer JE, Han L, Sumner SJ, Lewin AH, Fennell TR, Lust RM, Brown JM and Wingard CJ (2014). C(6)(0) exposure augments cardiac ischemia/reperfusion injury and coronary artery contraction in Sprague Dawley rats. *Toxicol Sci* 138: 365–378. [PubMed: 24431213]
36. Boularan C and Gales C (2015). Cardiac cAMP: production, hydrolysis, modulation and detection. *Front Pharmacol* 6: 203. [PubMed: 26483685]
37. Chung YW, Lagranha C, Chen Y, Sun J, Tong G, Hockman SC, Ahmad F, Esfahani SG, Bae DH, Polidovitch N, Wu J, Rhee DK, Lee BS, Gucek M, Daniels MP, Brantner CA, Backx PH, Murphy E and Manganiello VC (2015). Targeted disruption of PDE3B, but not PDE3A, protects murine heart from ischemia/reperfusion injury. *Proc Natl Acad Sci* 112: E2253–2262. [PubMed: 25877153]
38. Huang MH, Poh KK, Tan HC, Welt FG and Lui CY (2016). Therapeutic synergy and complementarity for ischemia/reperfusion injury: beta1-adrenergic blockade and phosphodiesterase-3 inhibition. *Int J Cardiol* 214: 374–380. [PubMed: 27085132]
39. Baldwin TA and Dessauer CW (2018). Function of Adenylyl Cyclase in Heart: the AKAP Connection. *J Cardiovasc Dev Dis* 5 DOI: 10.3390/jcdd5010002
40. Aragon M, Erdely A, Bishop L, Salmen R, Weaver J, Liu J, Hall P, Eye T, Kodali V, Zeidler-Erdely P, Stafflinger JE, Ottens AK and Campen MJ (2016). MMP-9-Dependent Serum-Borne Bioactivity Caused by Multiwalled Carbon Nanotube Exposure Induces Vascular Dysfunction via the CD36 Scavenger Receptor. *Toxicol Sci* 150: 488–498. [PubMed: 26801584]
41. Mandler WK, Nurkiewicz TR, Porter DW and Olfert IM (2017). Thrombospondin-1 mediates multi-walled carbon nanotube induced impairment of arteriolar dilation. *Nanotoxicology* 11: 112–122. [PubMed: 28024456]
42. Stapleton PA, Minarchick VC, Cumpston AM, McKinney W, Chen BT, Sager TM, Frazer DG, Mercer RR, Scabilloni J, Andrew ME, Castranova V and Nurkiewicz TR (2012). Impairment of coronary arteriolar endothelium-dependent dilation after multi-walled carbon nanotube inhalation: a time-course study. *Int J Mol Sci* 13: 13781–13803. [PubMed: 23203034]
43. Achterberg PW, Harmsen E and de Jong JW (1985). Adenosine deaminase inhibition and myocardial purine release during normoxia and ischaemia. *Cardiovasc Res* 19:593–598. [PubMed: 4053134]

Figure 1. Lung histology.

Left lungs were collected 24 hours following Vehicle or MWCNT instillation and examined histologically by H&E staining in order to visually assess pulmonary responses to MWCNT. Images were collected using the 10× objective and inlays were collected using the 40× objective. (A) Representative images of a left lung tissue collected from mice instilled with Vehicle. (B) Representative images of lung tissue collected from mice instilled with C-grade. (C) Representative images of left lung tissue collected from mice instilled with COOH. (D) Representative images of left lung tissue collected from mice instilled with N-doped.

Figure 2. Cardiac I/R injury.

We tested the hypothesis that 24 hours after MWCNT instillation cardiac I/R injury would result in expansion of myocardial infarction. Myocardial infarct sizes from CD-1 mice instilled with Vehicle, 100 μ g C-grade, COOH, or N-doped are presented as a percent of the zone at risk. *for $P < 0.05$ vs. Vehicle; † $P < 0.05$ vs. C-grade; ‡ $P < 0.05$ vs. COOH as determined by ANOVA. $N = 5$ for all groups.

Figure 3. Heart and Lung cAMP concentrations.

Heart and right lung tissues were collected from mice 24 hours after instillation with Vehicle or MWCNT and assessed for cAMP concentrations via ELISA. (A) cAMP concentrations in heart tissue collected from mice instilled with Vehicle (n = 6), C-grade (n = 5), COOH (n = 5), or N-doped (n = 6) normalized to total protein. (B) cAMP concentrations in lung tissue collected from mice instilled with Vehicle (n = 8), 100 μ g C-grade (n = 4), COOH (n = 4), and N-doped (n = 8) normalized to total protein. *P < 0.05 vs. Vehicle.

Adenosinergic Modulation of Aortic Tone

Figure 4. Adenosinergic modulation of aortic tone.

Isolated rings of thoracic aorta were stimulated with 1 μ M phenylephrine (PE) for 5 minutes followed by additions of 2-Cl-adenosine every 5 minutes to generate cumulative concentration-responses from 0.1 nM to 30 μ M. (A) Concentration-response profiles with NO Inhibitor present (n = 8-10). (B) Concentration-response profiles with 10 μ M general cyclooxygenase (COX) inhibitor indomethacin present (n = 3-5). (C) Concentration-response profiles with 1 mM nitric oxide synthase (NOS) inhibitor L-NAME present (n = 3-5). $P < 0.05$ for comparisons of the nonlinear fits of the treatment response curve was determined by best fit regression analysis [24]. *indicates $P < 0.05$ vs. Vehicle for mean data reported as done by Repeated Measures ANOVA with Bonferroni post-test. (n = 3-10).

Peak Vascular Responses

Figure 5. Aortic contraction with Phenylephrine (PE) and relaxation with 2-Cl-adenosine.

(A) Maximal aortic stress generated 5 minutes following stimulation with 1 μ M PE, just

prior to the start of 2-Cl-adenosine protocols ($n = 3-10$). (B) The maximal aortic

vasorelaxation generated by stimulation with 30 μ M 2-Cl-adenosine taken as a function of

peak stress generated during experimental protocols with PE + 2-Cl-adenosine ($n = 3-10$).

*for $P < 0.05$ by Two-way ANOVA with Bonferroni post-test.

Table 1

Bronchoalveolar Lavage Fluid Analysis

Bronchoalveolar Lavage Fluid Analysis						
	[Protein]	Total Cells	Macrophages	Neutrophils	Eosinophils	Lymphocytes
	µg/mL	× 10 ⁴	× 10 ⁴	× 10 ³	× 10 ³	× 10 ³
Vehicle	167 ± 27	15.9 ± 1.1	14.4 ± 1.2	8.1 ± 4.1	1.1 ± 0.4	2.7 ± 0.6
C-grade	409 ± 40 *	39.7 ± 3.7 *	21.4 ± 2.0	148 ± 25 *	11.2 ± 1.1	4.3 ± 1.2
COOH	255 ± 40	26.5 ± 2.3	14.5 ± 1.2	41.4 ± 2.0	69.1 ± 9.0 * [†]	5.2 ± 1.0
N-doped	324 ± 60	36.2 ± 6.4 *	13.7 ± 2.1 [†]	163 ± 50 * [‡]	42.2 ± 17 *	6.1 ± 1.0
Mean ± SEM (n = 6)						

* P < 0.05 vs. Vehicle;

[†]P < 0.05 vs. C-grade;

[‡]P < 0.05 vs. COOH