Published in final edited form as:

J Am Chem Soc. 2018 December 19; 140(50): 17349–17355. doi:10.1021/jacs.8b09677.

Nickel-Catalyzed Reductive [2+2] Cycloaddition of Alkynes

Santiago Cañellas^{†,§}, John Montgomery[#], and Miquel À. Pericàs^{†,‡,*}

[†]Institute of Chemical Research of Catalonia (ICIQ), The Barcelona Institute of Science and Technology, Avda. Països Catalans 16, E-43007, Tarragona, Spain

[‡]Departament de Química Inorgànica i Orgànica, Universitat de Barcelona, 08028, Barcelona, Spain

[§]Universitat Rovira i Virgili, Departament de Química Analítica i Química Orgànica, Marcel·lí Domingo, 1, 43007 Tarragona, Spain

[#]Department of Chemistry, University of Michigan, 930 North University Avenue, Ann Arbor, MI 48019-1055, United States

Abstract

The nickel-catalyzed synthesis of tetrasubstituted cyclobutenes from alkynes is reported. This transformation is uniquely promoted by the use of a primary aminophosphine, an unusual ligand in nickel catalysis. Mechanistic insights for this new transformation are provided, and post-reaction modifications of the cyclobutene products to stereodefined cyclic and acyclic compounds are reported, including the synthesis of *epi*-truxillic acid.

Graphical Abstract

*Corresponding Author: To whom correspondence should be addressed: mapericas@iciq.es.

The Supporting Information is available free of charge on the ACS Publications website.

Experimental procedures and spectral data (PDF)

Crystallographic data for **4a** (CIF)

Crystallographic data for **4b** (CIF)

Crystallographic data for **10** (CIF)

Notes

The authors declare no competing financial interest.

Since the pioneering work by Reppe and Wilke on the cyclotetramerization of acetylene and ensuing ligand-influenced outcomes in catalytic cycloadditions,¹ nickel catalysis has contributed to the development of many C–C and C–heteroatom bond-forming processes.² Building on these early studies, nickel-catalyzed methods have led to many advances in cycloadditions, allowing the assembly of various ring sizes from simple π -components depending on the ligand system and precursors employed.^{2a,2d} While the assembly of six- and eight-membered rings have been well developed, accessing cyclobutane and cyclobutene products is more limited and typically requires conjugated and activated substrates such as allenes, 1,3-enynes, or strained rings such as norbornene derivatives. The assembly of cyclobutanes and cyclobutenes by the catalytic cycloaddition of simple, non-conjugated π -components has thus proven elusive by using existing methods.

Cyclobutenes in particular are highly versatile synthetic intermediates because of their ring strain and high reactivity.³ In addition, they are also found in many bioactive metabolites, natural products, drugs, and organic dyes.⁴ In recent years, much progress has been made in strategies for the direct formation of the cyclobutene skeleton. Most commonly, the catalytic synthesis of cyclobutenes involves the use of an alkene-alkyne [2+2]-cycloaddition reaction typically involving conjugated and activated substrates (Figure 1a). Using this approach, many useful methodologies have been reported using transition metal^{5–6} and Lewis acid⁷ catalysis. An alternate approach involves the reductive cyclodimerization of two alkynes. This latter approach has been developed using a zirconium-mediated pyridine-directed strategy (Figure 1b),⁸ but variations of this process that involve sub-stoichiometric catalyst loadings or simple alkynes that lack directing groups have not been previously described. Despite the advances realized, the development of an intermolecular catalytic route with two different alkyne coupling partners is currently not available. We recognized that such a process would provide new opportunities for rapidly building up useful synthetic intermediates from simple starting materials.

In the course of exploring new types of catalytic additions to alkynes, we unexpectedly found that the reductive dimerization of alkynes to produce cyclobutenes is the major pathway when primary aminophosphine ligands are employed. (Figure 1c). Primary β -aminophosphines have rarely been used as ligands in catalysis,⁹ and are not typically employed in ligand screens of nickel-catalyzed processes. The reactivity is notable given that many nickel-catalyzed reductive transformations have been described with alkynes, whereas the alkyne reductive cyclodimerization has not previously been described. Other cycloadditions and reductive couplings involving nickel catalysis typically involve monodentate phosphines or NHC ligands,^{2d} suggesting that the unique behavior of primary aminophosphines will serve as an important counterpart to these well-studied catalytic systems. Given the novelty and potential utility of the reductive cyclodimerization of alkynes in the assembly of stereo- and regiochemically-defined cyclobutenes, as well as the lack of information about the unique reactivity of nickel complexes of primary aminophosphines, we report here the development, scope, and mechanistic insights of this newly discovered process.

We began our investigations by studying the reaction of **1a** in a High Throughput Experimentation (HTE) platform with Ni(COD)₂. We screened a wide variety of ligands

(**L1-L23**) such as aminophosphines, phosphoramidites, phosphines, amines, heterocyclic amines, NHCs and diols (Table 1, entries 4–5), several Brønsted acids such as methanol, isopropanol and benzoic acid (Table 1, entries 4, 11–12), and reducing agents such as triethylborane and dimethylphenylsilane (Table 1, entries 4 and 14).¹⁰ Interestingly, we only found significant amounts (>10% yield) of the desired cyclobutene **2a** with a combination of the primary aminophosphine ligand **L1**, methanol or isopropanol, and triethylborane. Notably, traces of product were found with a binary ligand system consisting of a primary amine (BuNH₂) and a tertiary phosphine (PPh₃), thus reinforcing the need for a tethered aminophosphine backbone. No significant amount of cyclobutene product was observed when using NiCl₂ reduced *in situ* by Mn or Zn (Table 1, entry 15) or the corresponding control experiments (Table 1, entries 2–3).

Several ligands of this class (**L1**, **L24-L28**) were screened on a preparative scale (Table 1, entries 6–11), after which **L1** was revealed as the ligand of choice. After further optimization of reaction conditions,¹⁰ the head-to-tail *trans*-cyclobutene **2a** was obtained in 73% isolated yield. Excellent regio- and stereoselectivities were observed, and all reagents and catalysts are commercially available, thus making the procedure simple and selective for the desired cyclobutenes.

We then turned our attention to the generality of this transformation. First, a robustness screen was performed, showing a high functional group tolerance.¹¹ Then, a series of examples of nickel-catalyzed reductive [2+2] cycloadditions of alkynes to produce *trans*-cyclobutenes were illustrated under optimized conditions (Scheme 1). The reaction was efficient with alkynes bearing alkyl groups with different chain lengths (**2a-d**). Electronic properties of the alkyne were evaluated by introducing electron-withdrawing and electron-donating groups on the aryl moiety. Although electron rich alkynes (**2f** and **2h**) underwent coupling faster than electron-poor ones (**2g**), the desired cyclobutene was successfully obtained in all cases. The reaction was effective in the presence of unprotected polar functional groups such as an alcohol (**2e**), aldehyde (**2k**), amide (**2j**), aryl ether (**2f** and **2i**) or amine (**2h**). Alkynes bearing heterocyclic moieties such as 2-pyridyl (**2l-m**), 2-pyrimidinyl (**2n**) and 2-thiazolyl (**2o**) also gave rise to the desired cyclobutenes. TMS-alkynes showed a silyl group migration on the cyclobutene products (**2p** and **2q**).¹² Surprisingly, aldehyde/alkyne reductive coupling¹³ or demethoxylation¹⁴ pathways were not observed. The scalability of this protocol was tested through synthesis of the cyclobutene **2e** on a 2-gram scale. In this case, the catalyst loading was successfully lowered to 2 mol% and the desired cyclobutene was obtained in 62% isolated yield.

The suppression of alkyne homocoupling presents a clear challenge in the development of synthetically desirable reductive heterocouplings of two different alkynes.¹⁶ In order to explore the feasibility of this approach, alkynes of different electronic and steric properties were examined to achieve the desired cross-reductive [2+2] alkyne cycloaddition. First, we attempted the reaction using two electronically opposed alkynes, **1g** and **1f**. To our delight, the corresponding heterocoupling product **2r** was obtained in 62% yield, accompanied by minor amounts of the homocoupling products (7% of **2g** and 21% of **2f**). Nonetheless, the two regioisomers arising from the position of the double bond in the cyclobutene ring were formed in 1:1 ratio. Then, we subjected the steroid phenyl-17 α -ethynylestradiol **1s** to the

reductive [2+2] cycloaddition reaction with **1a**. Due to the bulkiness of the steroid, it was unreactive towards the reductive homo-cyclodimerization. This allowed us to perform the heterocoupling in a very clean manner by slowly adding the alkyne **1a** *via* syringe pump. The corresponding hetero-coupling product **2s** was obtained in 59% yield (10:1 d.r.), with 40% of the starting material **1s** being recovered. Furthermore, diphenylacetylene **3a** gave rise to the *cis,cis*-diene **4a** in good yield (Scheme 2a).^{17a} We also attempted the cyclization of diynes to form bicyclic cyclobutenes (Scheme 2b). In both cases, the dienes **4b** and **4c** were obtained as the major products in good yields.¹⁸ The isomer obtained in products **4a-c** suggests the formation of the *trans*-cyclobutene followed by a thermal conrotatory electrocyclic ring-opening step.¹⁷

The potential synthetic utility of this protocol was then demonstrated through the diversification of the cyclobutene products (Scheme 3). First, ozonolysis of cyclobutene **2a** afforded the stereodefined acyclic 1,4-diketone **5** as a single isomer.¹⁹ On the other hand, epoxidation of **2a** gave rise to the cyclobutane epoxide **6** in good yield and 4:1 dr.²⁰ This product was then subjected to acidic conditions to perform a ring-contraction, thus affording the cyclopropyl ketone **7** in good yield and excellent diastereomeric ratio.²¹ Finally, we targeted the synthesis of *epi*-truxillic acid, a naturally occurring cyclobutanedicarboxylic acid with interesting biological properties.^{22,23} To our delight, the hydrogenation of cyclobutene **2e** to cyclo-butane **8** took place with exquisite diastereoselectivity, with **8** being isolated as a single isomer in good yield.²⁴ Finally, Jones oxidation afforded the corresponding *epi*-truxillic acid **9** in quantitative yield.^{25,26}

A number of experiments were conducted to shed light on the mechanistic features of this reductive cyclodimerization process. Deuterium-labelling studies displayed 90–92% deuterium incorporation using CD₃OD and CH₃OD in a regio- and stereoselective manner. On the other hand, no deuterium incorporation was observed using CD₃OH (Scheme 4a). With these results in hand, we performed the kinetic analysis of two parallel reactions run using CD₃OD and CH₃OH respectively. A primary kinetic isotope effect (KIE) of 3.0 was observed, indicating that a proton transfer is involved in the rate-determining step.²⁷ Motivated by the different reaction rate of electron-poor and electron-rich alkynes, we studied linear free-energy relationships to unveil potential cationic or anionic intermediates. The reactions using substrates **1a**, **1f**, **1g** and **1h** were monitored by ¹H-NMR to determine the corresponding kinetic behaviors (Scheme 4b). The observed Hammett plots, characterized by $\rho < 0$, suggest that a positive charge accumulates in the rate-determining step.²⁸

With this mechanistic information in hand, a plausible catalytic cycle is proposed for this reaction (Scheme 5). After rapid ligand exchange with **L1** (**I**) and coordination of two alkynes to the nickel center (**II**),²⁹ Ni(0) may undergo an oxidative cyclization to form the first C–C bond, generating the nickelacycle (**III**).^{1,2a,30} Then, the methanol-triethylborane adduct could protonate³¹ the π -system generating a stabilized cationic nickel carbene **IV**.^{32,33} Next, a 1,2-alkyl migration would generate the 4-membered ring and the second C–C bond (**V**).^{12,34} Finally, an ethyl transfer to nickel (**VI**), β -hydride elimination³⁵ (**VII**) and reductive elimination³⁶ gives rise to the desired cyclobutene, regenerating the nickel(0) catalyst. *In situ* HRMS (ESI+) analysis of the reaction mixture using alkyne **1e** revealed a

very intense m/z signal at 552.1609 with the characteristic isotopic distribution of a nickel species. This data suggests the presence of a nickel species that incorporates the aminophosphine and the two units of the corresponding alkyne (i.e. structures **II–V**). In combination with the Hammett studies and KIE experiments, the data overall is consistent with **III** being the catalyst resting state.

This overall mechanism bears analogy to [3+2] reductive cycloadditions of enals and alkynes that proceed through the protonation of enolate motifs embedded with a nickel metallacycle.^{37–38} However, the only previous examples where metallacyclopentadienes are converted to cyclobutenes are the stoichiometric zirconium-mediated processes described above.⁸ Notably, protonation of a Ni(II) metallacyclopentene species by a water/borate adduct^{31a} and the protonation of Ni(0) species by a similar methanol/borane mixture^{31b–c} have recently been proposed in the hydroalkylation and hydroarylation of allenes, styrenes and dienes under similar conditions to our study.³¹ Although Ni(0) protonation would be expected to be faster than protonation of **III**, products resulting from hydroalkylation or reduction of the alkyne are only observed as minor by-products. Furthermore, exogenous alkenes such as styrene and *E/Z*- β -methylstyrene are not incorporated in the cyclobutene products,³⁹ suggesting that nickel hydride mediated reduction of the alkyne and subsequent alkene-alkyne [2+2] cycloaddition reaction is unlikely for the production of cyclobutene products. The basis for the unique promotion exhibited by the aminophosphine **L1** is currently unclear. The role of this ligand architecture may involve features other than simple bidentate coordination, such as proton shuttling or generation of a Lewis adduct with the borane component, and further studies to understand the unique behavior of this ligand class are in progress.

In summary, a nickel-catalyzed reductive [2+2] cycloaddition reaction of alkynes towards the synthesis of *trans*-cyclobutenes has been developed. The use of an unusual primary aminophosphine ligand was key to the discovery of this new reaction. Post-reaction modifications highlighted the synthetic versatility of these products, including the synthesis of the natural product *epi*-truxillic acid.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENT

Financial support from the MINECO (grant CTQ2015–69136-R (MINECO/FEDER)) and CERCA Program/ Generalitat de Catalunya is gratefully acknowledged. We also thank the MINECO for support through the Severo Ochoa Excellence Accreditation 2014–2018 (SEV-2013–0319). S.C. thanks the MINECO for a Severo Ochoa FPI fellowship (BES-2015–072152). J.M. thanks the National Institutes of Health (R35-GM118133) for support. We are grateful to the ICIQ X-ray diffraction unit for the X-ray single crystal analyses of **4a**, **4b** and **10**. Prof. Ruben Martin (ICIQ) and Dr. Alex J. Nett (University of Michigan) are acknowledged for very useful discussions. We thank Mr. Sergio Fernández (ICIQ) for the assistance on the head-space analysis.

REFERENCES

- (1). (a) Reppe W; Schlichting O; Klager K; Toepel T Cyclisierende Polymerisation von Acetylen I Über Cyclooctatetraen. *Justus Liebig Ann. Chem* 1948, 560, 1–92. (b) Wilke G Contributions to Organo-Nickel Chemistry. *Angew. Chem. Int. Ed. Engl* 1988, 27, 185–206.
- (2). (a) Montgomery J Organonickel Chemistry In *Organometallics in Synthesis: Fourth Manual*; Lipshutz BH, Ed.; Wiley: Hoboken, NJ, 2013; pp 319–428. (b) Tasker SZ; Standley EA; Jamison TF Recent advances in homogeneous nickel catalysis. *Nature*, 2014, 509, 299–309. [PubMed: 24828188] (c) Standley EA; Tasker SZ; Jensen KL; Jamison TF Nickel Catalysis: Synergy between Method Development and Total Synthesis. *Acc. Chem. Res* 2015, 48, 1503–1514. [PubMed: 25905431] (d) Thakur A; Louie J Advances in Nickel-Catalyzed Cycloaddition Reactions To Construct Carbocycles and Heterocycles. *Acc. Chem. Res* 2015, 48, 2354–2365. [PubMed: 26200651] (e) Moslin RM; Miller-Moslin K; Jamison TF Regioselectivity and enantioselectivity in nickel-catalysed reductive coupling reactions of alkynes. *Chem. Commun* 2007, 4441–4449.
- (3). a) Namyslo JC; Kaufmann DE The Application of Cyclobutane Derivatives in Organic Synthesis. *Chem. Rev* 2003, 103, 1485–1537. [PubMed: 12683789] b) Misale A; Niyomchon S; Maulide N Cyclobutenes: At a Crossroad between Diastereoselective Syntheses of Dienes and Unique Palladium-Catalyzed Asymmetric Allylic Substitutions. *Acc. Chem. Res* 2016, 49, 2444–2458. [PubMed: 27797480]
- (4). (a) Sergeiko A; Poroikov VV; Hanus LO; Dembitsky VM Cyclobutane-containing alkaloids: origin, synthesis, and biological activities. *Open Med. Chem. J* 2008, 2, 26–37. [PubMed: 19696873] (b) Dembitsky VM Naturally occurring bioactive Cyclobutane-containing (CBC) alkaloids in fungi, fungal endophytes, and plants. *Phytomedicine* 2014, 21, 1559–1581. [PubMed: 25442265] (c) Chen G; Sasabe H; Igarashi T; Hong Z; Kido J Squaraine dyes for organic photovoltaic cells. *J. Mater. Chem. A* 2015, 3, 14517–14534.
- (5). For examples of nickel-catalyzed activated alkene-alkyne [2+2] cycloaddition reactions, see: Huang D-J; Cheng C-H [2+2] Dimerization of norbornadiene and its derivatives in the presence of nickel complexes and zinc metal. *J. Organomet. Chem* 1995, 490, C1–C7. Huang D-J; Rayabarapu DK; Li L-P; Sambaiiah T; Cheng C-H Nickel-Catalyzed [2+2] Cycloaddition of Alkynes with Activated Cyclic Alkenes: Synthesis and Novel Ring Expansion Studies of Cyclobutene Products. *Chem. Eur. J* 2000, 6, 3706–3713. [PubMed: 11073240] Saito S; Hirayama K; Kabuto C; Yamamoto Y Nickel(0)-Catalyzed [2+2] Annulation of Electron-Deficient Allenes. Highly Regioselective Synthesis of Cyclobutanes. *J. Am. Chem. Soc* 2000, 122, 10776–10780. Nishimura A; Ohashi M; Ogoshi S Nickel-Catalyzed Intermolecular [2+2] Cycloaddition of Conjugated Enynes with Alkenes. *J. Am. Chem. Soc* 2012, 134, 15692–15695. [PubMed: 22966854] Abulimiti A; Nishimura A; Ohashi M; Ogoshi S Nickel-catalyzed [2+2] Cycloaddition Reaction of Bulky Enones with Simple Alkynes. The Effect of Bulkiness of Substituent Attached at β -Carbon. *Chem. Lett* 2013, 42, 904–905. Hori H; Arai S; Nishida AA 2-Benzothiazolylphenyl Group Accelerates the Intramolecular [2+2] Cycloaddition of Allene Ynes. *Asian J. Org. Chem* 2014, 3, 41–43. Noucti NN; Alexanian EJ Stereoselective Nickel-Catalyzed [2+2] Cycloadditions of Ene Allenes. *Angew. Chem. Int. Ed* 2015, 54, 5447–5450. Kumar R; Tamai E; Ohnishi A; Nishimura A; Hoshimoto Y; Ohashi M; Ogoshi S Nickel-Catalyzed Enantioselective Synthesis of Cyclobutenes via [2+2] Cycloaddition of α,β -Unsaturated Carbonyls with 1,3-Enynes. *Synthesis* 2016, 48, 2789–2794. Qin H; Chen J; Li K; He Z; Zhou Y; Fan B Nickel Catalyzed Asymmetric [2+2] Cycloaddition Reaction of Hetero Bicyclic Alkenes with Internal Alkynes. *Chem. Asian J* 2018, DOI: 10.1002/asia.201800492.
- (6). For selected examples of [2+2] cycloaddition reactions catalyzed by other metals different than nickel, see: Treutwein J; Hilt G Cobalt-Catalyzed [2+2] Cycloaddition. *Angew. Chem. Int. Ed* 2008, 47, 6811–6813. López-Carrillo V; Echavarren AM Gold(I)-Catalyzed Intermolecular [2+2] Cycloaddition of Alkynes with Alkenes. *J. Am. Chem. Soc* 2010, 132, 9292–9294. [PubMed: 20568750] Schotes C; Mezzetti A Enantioselective Ficini Reaction: Ruthenium/PNNP-Catalyzed [2+2] Cycloaddition of Ynamides with Cyclic Enones. *Angew. Chem. Int. Ed* 2011, 50, 3072–3074. Nishimura A; Ohashi M; Ogoshi S Nickel-Catalyzed Inter-molecular [2+2] Cycloaddition of Conjugated Enynes with Alkenes. *J. Am. Chem. Soc* 2012, 134, 15692–15695. [PubMed: 22966854] Nishimura A; Tamai E; Ohashi M; Ogoshi S Synthesis of Cyclobutenes and Allenes

by Cobalt-Catalyzed Cross-Dimerization of Simple Alkenes with 1,3-Enynes. *Chem. Eur. J* 2014, 20, 6613–6617. [PubMed: 24782325] de Orbe ME; Echavarren AM Broadening the Scope of the Gold-Catalyzed [2+2] Cycloaddition Reaction: Synthesis of Vinylcyclobutenes and Further Transformations. *Eur. J. Org. Chem* 2018, 2740–2752. Bai Y-B; Luo Z; Wang Y; Gao J-M; Zhang L Au-Catalyzed Intermolecular [2+2] Cycloadditions between Chloroalkynes and Unactivated Alkenes. *J. Am. Chem. Soc.* 2018, 140, 5860–5865. [PubMed: 29618202]

- (7). For selected examples of Lewis Acid catalyzed [2+2] cycloaddition reactions, see: Snider BB; Rodini DJ; Conn RSE; Sealfon S Lewis acid catalyzed reactions of methyl propiolate with unactivated alkenes. *J. Am. Chem. Soc* 1979, 101, 5283–5293. Ito H; Hasegawa M; Takenaka Y; Kobayashi T; Iguchi K Enantioselective Total Synthesis of (+)-Tricycloclavulone. *J. Am. Chem. Soc* 2004, 126, 4520–4521. [PubMed: 15070361] Sweis RF; Schramm MP; Kozmin SA Silver-Catalyzed [2 + 2] Cycloadditions of Siloxy Alkynes. *J. Am. Chem. Soc* 2004, 126, 7442–7443. [PubMed: 15198582] Kang T; Ge S; Lin L; Lu Y; Liu X; Feng X A Chiral *N,N'*-Dioxide-Zn^{II} Complex Catalyzes the Enantioselective [2+2] Cycloaddition of Alkynones with Cyclic Enol Silyl Ethers. *Angew. Chem. Int. Ed* 2016, 55, 5541–5544. Shen L; Zhao K; Doitomi K; Ganguly R; Li Y-X; Shen Z-L; Hirao H; Loh T-P Lewis Acid-Catalyzed Selective [2+2]-Cycloaddition and Dearomatizing Cascade Reaction of Aryl Alkynes with Acrylates. *J. Am. Chem. Soc* 2017, 139, 13570–13578. [PubMed: 28880536]
- (8). Liu Y; Liu M; Song Z Highly Regio- and Stereoselective Synthesis of Tetrasubstituted Cyclobutenes via Cyclodimerization of Alkynes Mediated by Zirconium. *J. Am. Chem. Soc* 2005, 127, 3662–3663. [PubMed: 15771477]
- (9). For examples using primary β -aminophosphine ligands, see: Abdur-Rashid K; Guo R; Lough AJ; Morris RH; Song D Synthesis of Ruthenium Hydride Complexes Containing beta Aminophosphine Ligands Derived from Amino Acids and their use in the H₂-Hydrogenation of Ketones and Imines. *Adv. Synth. Catal* 2005, 347, 571–579. Guo M; Jian F; He R The air-stable and highly efficient P,N-chelated palladium(II) complexes as catalysts for the Suzuki cross-coupling reaction at room temperature *Tetrahedron Lett.* 2006, 45, 2033–2036. John JM; Bergens SH A Highly Active Catalyst for the Hydrogenation of Amides to Alcohols and Amines. *Angew. Chem. Int. Ed* 2011, 50, 10377–10380. Higuchi T; Tagawa R; Iimuro A; Akiyama S; Nagae H; Mashima K Tunable Ligand Effects on Ruthenium Catalyst Activity for Selectively Preparing Imines or Amides by Dehydrogenative Coupling Reactions of Alcohols and Amines. *Chem. Eur. J* 2017, 23, 12795–12804. [PubMed: 28557018] Van Putten R; Uslamin EA; Garbe M; Liu C; Gonzalez-de-Castro A; Lutz M; Junge K; Hensen EJM; Beller M; Lefort L; Pidko EA Non-Pincer-Type Manganese Complexes as Efficient Catalysts for the Hydrogenation of Esters. *Angew. Chem. Int. Ed* 2017, 56, 7531–7534. For a review on the use of tertiary β -aminophosphine ligands, see: Li W; Zhang J Recent developments in the synthesis and utilization of chiral β -aminophosphine derivatives as catalysts or ligands. *Chem. Soc. Rev* 2016, 45, 1657–1677. [PubMed: 26776280]
- (10). See Supporting Information for more details.
- (11). Functional groups such as nitrile, aldehyde, amide, ester, ketone, bromide, phenol, alcohols, pyridine, alkenes and amine were tolerated, while imidazole shut down the desired reaction. See Supporting Information for more details. Collins KD; Glorius F A robustness screen for the rapid assessment of chemical reactions. *Nat. Chem* 2013, 5, 597–601. [PubMed: 23787750] Richardson J; Ruble JC; Love EA; Berritt S A Method for Identifying and Developing Functional Group Tolerant Catalytic Reactions: Application to the Buchwald–Hartwig Amination. *J. Org. Chem* 2017, 82, 3741–3750. [PubMed: 28245358]
- (12). This migration observed gives support to the cationic species proposed to be generated at β -position of the silyl group (species V on Scheme 5). For selected examples of silyl group migrations to β -cationic positions, see: Groaning MD; Brengel GP; Meyers A I. Allyldimethyltritylsilane. Synthesis of Cyclopentanols, Oxetanes, and Tetrahydrofurans by Reaction with Electron Deficient Olefins. *J. Org. Chem* 1998, 63, 5517–5522. Knölker H-J; Baum E; Graf R; Jones PG; Spieb O An Unprecedented Domino Double Allylsilane [3+2] Cycloaddition/Wagner-Meerwein Rearrangement/Friedel-Crafts Alkylation/Elimination Reaction Sequence Leading to a Novel Pentacyclic Ring System. *Angew. Chem. Int. Ed* 1999, 38, 2583–2585. Ball-Jones NR; Badillo JJ; Tran NT; Franz AK Catalytic Enantioselective Carboannulation with Allylsilanes. *Angew. Chem. Int. Ed* 2014, 53, 9462–9465.

- (13). a)Oblinger E; Montgomery J A New Stereoselective Method for the Preparation of Allylic Alcohols. *J. Am. Chem. Soc* 1997, 119, 9065–9066.b)Montgomery J Nickel Catalyzed Reductive Cyclizations and Couplings. *Angew. Chem. Int. Ed* 2004, 43, 3890–3908.c)Malik HA; Sormunen GJ; Montgomery J A General Strategy for Regiocontrol in Nickel-Catalyzed Reductive Couplings of Aldehydes and Alkynes. *J. Am. Chem. Soc* 2010, 132, 6304–6305. [PubMed: 20394367] d)Todd DP; Thompson BB; Nett AJ; Montgomery J Deoxygenative C–C Bond-Forming Processes via a Net Four-Electron Reductive Coupling. *J. Am. Chem. Soc* 2015, 137, 12788–12791. [PubMed: 26436636] e)Wang H; Lu G; Sormunen GJ; Malik HA; Liu P; Montgomery J NHC Ligands Tailored for Simultaneous Regio- and Enantiocontrol in Nickel-Catalyzed Reductive Couplings. *J. Am. Chem. Soc* 2017, 139, 9317–9324. [PubMed: 28621131] f)Nett AJ; Cañellas S; Higuchi Y; Robo MT; Kochkodan JM; Haynes MT II; Kampf JW; Montgomery J Stable, Well-Defined Nickel(0) Catalysts for Catalytic C–C and C–N Bond Formation. *ACS Catal.* 2018, 8, 6606–6611. [PubMed: 30079273]
- (14). Igarashi T; Haito A; Chatani N; Tobisu M Nickel-Catalyzed Reductive Cleavage of Carbon–Oxygen Bonds in Anisole Derivatives Using Diisopropylaminoborane. *ACS Catal.* 2018, 8, 7475–7483.
- (15). The major diastereomer is a trans-cyclobutene. However, attempts to obtain crystals of 2s, or derivatives thereof, suitable for X-ray crystal analysis have been unsuccessful.
- (16). Reichard HA; McLaughlin M; Chen MZ; Micalizio GC Regioselective Reductive Cross-Coupling Reactions of Unsymmetrical Alkynes. *Eur. J. Org. Chem* 2010, 391–409.
- (17). The hypothetical trans-1,2,3,4-tetraphenylcyclobutene is known to undergo rapid ring-opening to afford the corresponding cis, cis-1,2,3,4-tetraphenyl-1,3-butadiene 4a. See:Freedman HH; Doorakian GA; Sandel VR The Valence Isomerization of 1,2,3,4-Tetraphenylcyclobutene and Its Anion. *J. Am. Chem. Soc* 1965, 87, 3019–3020.Lee PS; Sakai S; Hörstermann P; Roth WR; Kallel EA; Houk KN Altering the Allowed/Forbidden Gap in Cyclobutene Electrocyclic Reactions: Experimental and Theoretical Evaluations of the Effect of Planarity Constraints. *J. Am. Chem. Soc* 2003, 125, 5839–5848. [PubMed: 12733925]
- (18). The diene structures of 4a and 4b were unambiguously confirmed by single X-ray crystal analyses. See the Supporting Information file.
- (19). For a selected example, see:Arnó M; Betancur-Galvis L; González MA; Sierra J; Zaragoza RJ Synthesis and cytotoxic activity of novel C7-Functionalized spongiane diterpenes. *Bioorganic Med. Chem* 2003, 11, 3171–3177.
- (20). For a selected example, see:Paquette LA; Wang T-Z; Cottrell CE Flattening of the cyclooctatetraene ring by annulation. *J. Am. Chem. Soc* 1987, 109, 3730–3734.
- (21). Baumann AN; Schüppel F; Eisold M; Kreppel A; de Vivie-Riedle R; Didier D Oxidative Ring Contraction of Cyclobutenes: General Approach to Cyclopropylketones including Mechanistic Insights. *J. Org. Chem* 2018, 83, 4905–4921. [PubMed: 29641195]
- (22). a)Liebermann C Ueber Cinnamylcocaïn. *Ber. Dtsch. Chem. Ges* 1888, 21, 3372–3376.b)Pictet A in *The Vegetable Alkaloids, With Particular Reference to Their Chemical Constitution*, Wiley, New York, 1913, p. 236.
- (23). Antinociceptive activity:Chi Y-M; Nakamura M; Zhao X-Y; Yoshizawa T; Yan W-M; Hashimoto F; Kinjo J; Nohara T; Sakurada S Antinociceptive Activities of α -Truxillic Acid and β -Truxinic Acid Derivatives. *Biol. Pharm. Bull* 2006, 29, 580–584. [PubMed: 16508173] Kaczocha M; Glaser ST; Deutsch DG Identification of intracellular carriers for the endocannabinoid anandamide. *Proc. Natl. Acad. Sci. USA* 2009, 106, 6375. [PubMed: 19307565] Liu SX; Jin HZ; Shan L; Zeng HW; Chen BY; Sun QY; Zhang WD Inhibitory effect of 4,4'-dihydroxy- α -truxillic acid derivatives on NO production in lipopolysaccharide-induced RAW 264.7 macrophages and exploration of structure–activity relationships. *Bioorg. Med. Chem. Lett* 2013, 23, 2207–2211. [PubMed: 23454014] Antimuscarinic activity:Lysíková M; Fuksová K; Elbert T; Jakubík J; Tuček S Subtype-selective inhibition of [methyl-3H]-N-methylscopolamine binding to muscarinic receptors by α -truxillic acid esters. *Br. J. Pharmacol* 1999, 127, 1240–1246. [PubMed: 10455271]
- (24). Please note that products 8 and 9 are meso compounds, so they lack chirality.

- (25). This isomeric form possessing a characteristic S1 axis is easily distinguished by NMR from the other isomers of truxillic acid, 20 and the structure of 9 was unambiguously assigned by single X-ray crystal of its methyl ester 10. See the Supporting Information file.
- (26). For a selected examples of Jones oxidation reactions, see: Bowden K; Heilbron IM; Jones ERH; Weedon BCL 13. Researches on acetylenic compounds. Part I. The preparation of acetylenic ketones by oxidation of acetylenic carbinols and glycols. *J. Chem. Soc* 1946, 0, 39–45. Heller L; Schwarz S; Perl V; Köwitsch A; Siewert B; Csuk R Incorporation of a Michael acceptor enhances the antitumor activity of triterpenic acids. *Eur. J. Med. Chem* 2015, 101, 391–399. [PubMed: 26177446]
- (27). Gómez-Gallego M; Sierra MA Kinetic Isotope Effects in the Study of Organometallic Reaction Mechanisms. *Chem. Rev* 2011, 111, 4857–4963. [PubMed: 21545118]
- (28). At present, we cannot rule out other mechanistic interpretation of these results. For further insight on linear free energy relationships, see: Wells PR Linear Free Energy Relationships. *Chem. Rev* 1963, 63, 171–219.
- (29). Kinetic analyses of the model reaction with different L1:Ni ratios (1:1, 2:1 and 4:1) suggest that the active species consist of equimolar quantities of ligand and nickel. However, the use of an excess of ligand prevents the generation of other byproducts, achieving higher yield of the desired cyclobutene. See Supporting Information file for more details.
- (30). a) Ogoshi S; Oka M, Kurosawa H Direct Observation of Oxidative Cyclization of η^2 -Alkene and η^2 -Aldehyde on Ni(0) Center. Significant Acceleration by Addition of Me₃SiOTf. *J. Am. Chem. Soc* 2004, 126, 11802–11803. [PubMed: 15382909] b) Hong X; Holte D; Götz DCG; Baran PS; Houk KN Mechanism, Reactivity, and Selectivity of Nickel-Catalyzed [4+4+2] Cycloadditions of Dienes and Alkynes. *J. Org. Chem* 2014, 79, 12177–12184. [PubMed: 25325891] c) Ma W; Yu C; Chen T; Xu L; Zhang W-X; Xi Z Metallacyclopentadienes: synthesis, structure and reactivity. *Chem. Soc. Rev* 2017, 46, 1160–1192. [PubMed: 28119972]
- (31). (a) Takahashi G; Shirakawa E; Tsuchimoto T; Kawakami Y Nickel-Catalyzed Addition of Organoboronates to 1,2-Dienes and the Corresponding Three-Component Reaction with an Alkyne. *Adv. Synth. Catal* 2006, 348, 837–840. (b) Xiao L-J; Cheng L; Feng W-M; Li M-L; Xie J-H; Zhou Q-L Nickel(0)-Catalyzed Hydroarylation of Styrenes and 1,3-Dienes with Organoboron Compounds. *Angew. Chem. Int. Ed* 2018, 57, 461–464. (c) Han X-W; Zhang T; Zheng Y-L; Yao W-W; Li J-F; Pu Y-G; Ye M; Zhou Q-L Brønsted Acid Enabled Nickel-Catalyzed Hydroalkenylation of Aldehydes with Styrene and its Derivatives. *Angew. Chem. Int. Ed* 2018, 57, 5068–5071.
- (32). This transformation is possibly the rate-determining step, as suggested by the mechanistic experiments, since it involves both the formation of a cation and a proton transfer of the former methanol proton.
- (33). For examples of nickel carbene intermediates, see: Barluenga J; Barrio P; López LA; Tomás M; García-Granda S; Alvarez-Rúa C Nickel(0)-Mediated [3+2+2] and [2+2+2+1] Cyclization Reactions of Chromium Fischer Carbene Complexes and Alkynes. *Angew. Chem. Int. Ed* 2003, 42, 3008–3011. Ni Y; Montgomery, J. Synthetic Studies and Mechanistic Insight in Nickel-Catalyzed [4+2+1] Cycloadditions. *J. Am. Chem. Soc* 2006, 128, 2609–2614. [PubMed: 16492045]
- (34). a) Kociński P; Barber C Synthetic applications of metallate rearrangements. *Pure & Appl. Chem* 1990, 62, 1933–1940. b) Clement ND; Cavell KJ Transition-Metal-Catalyzed Reactions Involving Imidazolium Salt/N Heterocyclic Carbene Couples as Substrates. *Angew. Chem. Int. Ed* 2004, 43, 3845–3847. c) Steinke T; Shaw BK; Jong H; Patrick BO; Fryzuk MD; Green JC Noninnocent Behavior of Ancillary Ligands: Apparent Trans Coupling of a Saturated N-Heterocyclic Carbene Unit with an Ethyl Ligand Mediated by Nickel. *J. Am. Chem. Soc* 2009, 131, 10461–10466. [PubMed: 19585991]
- (35). Head-space analysis reveals the generation of ethylene in the reaction. See Supporting Information file for more details.
- (36). a) Patel SJ; Jamison TF Catalytic Three-Component Coupling of Alkynes, Imines, and Organoboron Reagents. *Angew. Chem. Int. Ed* 2003, 42, 1364–1367. b) Molinaro C; Jamison TF Nickel-Catalyzed Reductive Coupling of Alkynes and Epoxides. *J. Am. Chem. Soc* 2003, 125, 8076–8077. [PubMed: 12837057] (c) McCarren PR; Liu P; Cheong PH-Y; Jamison TF; Houk KN

- Mechanism and Transition-State Structures for Nickel-Catalyzed Reductive Alkyne–Aldehyde Coupling Reactions. *J. Am. Chem. Soc.* 2009, 131, 6654–6655. [PubMed: 19397371]
- (37). (a) Herath A; Montgomery J Catalytic Intermolecular Enal–Alkyne [3+2] Reductive Cycloadditions. *J. Am. Chem. Soc.* 2006, 128, 14030–14031. [PubMed: 17061877] (b) Jenkins AD; Herath A; Song M; Montgomery J Synthesis of Cyclopentenols and Cyclopentenones via Nickel-Catalyzed Reductive Cycloaddition. *J. Am. Chem. Soc.* 2011, 133, 14460–14466. [PubMed: 21819132] (c) Ohashi M; Taniguchi T; Ogoshi S Nickel-Catalyzed Formation of Cyclopentenone Derivatives via the Unique Cycloaddition of α,β -Unsaturated Phenyl Esters with Alkynes. *J. Am. Chem. Soc.* 2011, 133, 14900–14903. [PubMed: 21877707]
- (38). For other examples of reductive cycloaddition: Enholm EJ; Kinter KS Free radical cyclizations promoted by allylic O-stannyl ketyls: the intramolecular coupling of the beta-carbons of activated alkenes. *J. Am. Chem. Soc.* 1991, 113, 7784–7785. Hays DS; Fu GC Organotin Hydride Catalyzed Carbon–Carbon Bond Formation: Radical-Mediated Reductive Cyclization of Enals and Enones. *J. Org. Chem.* 1996, 61, 4–5. Savchenko AV; Montgomery J Organozinc/Nickel(0)-Promoted Cyclizations of Bis-Enones. *J. Org. Chem.* 1996, 61, 1562–1563. [PubMed: 11667020] Lee J; Kim H; Cha JK A New Variant of the Kulinkovich Hydroxycyclopropanation. Reductive Coupling of Carboxylic Esters with Terminal Olefins. *J. Am. Chem. Soc.* 1996, 118, 4198–4199. Kulinkovich OG The Chemistry of Cyclopropanols. *Chem. Rev.* 2003, 103, 2597–2632. [PubMed: 12848581] Zhou YY; Uyeda C Reductive Cyclopropanations Catalyzed by Dinuclear Nickel Complexes. *Angew. Chem. Int. Ed.* 2016, 55, 3171–3175.
- (39). Kinetic profile of the reaction for the formation of 2c in the presence of 1 equivalent of E- and Z- β -methylstyrene overlap with the one where no additive is added. See kinetic analyses in the Supporting Information file.

a) Alkene-alkyne [2+2]-cycloaddition

b) Zirconium-mediated reductive cyclodimerization of ortho-alkynylpyridines

c) This work

Figure 1.
Approaches towards the synthesis of cyclobutenes

Scheme 1.

Substrate scope. Isomeric ratios (d.r. and r.r.) calculated by GC-FID. ^a Ni(COD)₂ (2 mol %) and **L1** (4 mol %) were used. ^b See reference 15.

Scheme 2.

a) Reductive homocoupling of diphenylacetylene. b) Reductive cyclization of diynes towards dienes.

Scheme 3.

Synthetic diversification of cyclobutene **2a** and **2f**. a) O₃, CHCl₃, -78 °C. b) *m*-CPBA, NaHCO₃, CHCl₃, 0 °C. c) aq HCl (0.1 M), rt. d) H-Cube, 1 mL/min, H₂ (60 bar), Pd/C (10%), 60 °C, MeOH (0.25 M). e) CrO₃, H₂SO₄, H₂O, acetone.

Scheme 4.
Mechanistic experiments.

Scheme 5.
Proposed catalytic cycle.

Table 1.

Screening and optimization of the reaction conditions.^a

Entry	Variation from standard conditions	Yield (%)
1	None	73
2 ^b	w/o Ni(COD) ₂ , L1 or BEt ₃	0
3 ^b	w/o MeOH	7
4 ^b	THF:MeOH (4:1)	31
5 ^b	Other class of ligands instead of L1	<5–9
6 ^c	L24 instead of L1	30
7 ^c	L25 instead of L1	19
8 ^c	L26 instead of L1	11
9 ^c	L27 instead of L1	47
10 ^c	L28 instead of L1	37
11 ^c	L1 , 4 hours	61
12 ^b	iPrOH instead of MeOH	12
13 ^b	Benzoic acid instead of MeOH	0
14 ^b	PhMe ₂ SiH instead of BEt ₃	<5
15 ^c	NiCl ₂ + Mn or Zn instead of Ni(COD) ₂	<5

^a Optimized reaction conditions: **1a** (0.45 mmol), Ni(COD)₂ (0.045 mmol), **L1** (0.09 mmol), BEt₃ (0.9 mmol), THF:MeOH (1:4 ratio, 0.08 M), 50 °C, 2.5 h.

^b 4:1 ratio THF:MeOH was used in the HTE (10 μmol **1a**).

^c 4 hours.