Published in final edited form as:

Curr Opin Immunol. 2019 April ; 57: 15–22. doi:10.1016/j.coi.2018.09.020.

Epigenetic dynamics in normal and malignant B cells: Die a hero or live to become a villain

Jared M. Andrews¹ and Jacqueline E. Payton^{1,*}

¹Department of Pathology and Immunology, Washington University in St. Louis. St. Louis, MO 63110, USA.

Abstract

Normal B cell development, activation, and terminal differentiation depend on the intricate dynamics of cooperating epigenetic and non-coding components to control the level and timing of expression of thousands of genes. Recent genome-wide studies have integratively mapped changes in the chromatin landscape, DNA methylome, 3-dimensional interactome, and coding and non-coding transcriptomes of normal and malignant B cells. Genetic ablation in human cells and mouse models has begun to elucidate the coordinated roles of essential epigenetic modifiers, key transcription factors, and long non-coding RNAs in B cell biology. Perturbation of these stewards of the epigenome drive B cell oncogenesis, but may be exploited to develop new avenues of therapy.

Introduction

Long regarded as “junk DNA”, the non-protein-coding portion of the genome has been liberated from this denigrating moniker by high-throughput sequencing-enabled discoveries demonstrating its essential role in the regulation of genome integrity, structure, replication, and gene expression. Sometimes referred to as the “regulome”, these epigenetic mechanisms are exquisitely controlled and essential for development and homeostasis, and when disturbed, contribute to a range of diseases, from auto-immunity to cancer.

The following review discusses the most recent discoveries regarding the role of epigenetic regulation in B lymphocytes, focusing on latter stages of maturation and activation, as well as how mature B cell cancers subvert these mechanisms for their own survival and progression. Figure 1 provides a basic overview of each epigenetic mechanism and its role in B cell activation and Figures 2 & 3 illustrate how lymphomas hijack these mechanisms. We summarize and highlight three epigenetic modes of regulation and discuss each separately: first DNA methylation; second, histone modifications and chromatin architecture; and last, long non-coding RNAs.

*To whom correspondence should be addressed: jpayton@wustl.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Declarations of interest: none

Dynamic flux of the B cell epigenome throughout development and in B cell cancers

Genome-wide studies revealed the extensive plasticity of the DNA methylome during B cell development (Figure 1A), with dynamic methylation at nearly 5 million CpGs (30% of sites surveyed) [1,2]. These changes are temporally enriched in specific regulatory regions, with enhancers and heterochromatin preferentially demethylated in precursors and germinal center (GC) cells, respectively, and methylation gains in polycomb-repressed regions in the most differentiated B cells [3]. Dynamic methylome changes exhibited no direct correlation with gene expression, though demethylation of key B cell transcription factor (TF) binding sites correlated with expression of those TFs and their transcriptional programs (Figure 1B) [1,2,4,5]. The importance of enhancer demethylation was directly demonstrated by knockouts of TET family demethylase genes in early stage B cells, which prevented lineage-specific programmed demethylation of enhancers of B-cell lineage genes, causing decreased expression of these key genes, defects in development at the pro- to pre-B cell stages, and impaired IgK rearrangement [6,7]. The importance of de novo methylation was shown by double knock-out of DNMT3A and DNMT3B in pre-B cells, which did not affect B cell development and maturation, but, upon immunization, increased cell activation and expansion of GC B cell and plasma cell (PC) populations by 2–3-fold [8*].

Given the essential nature of epigenetic control in B cell differentiation, it is not surprising that aberrant DNA methylomes are hallmarks of B cell cancers and the degree of methylation heterogeneity correlates with more aggressive disease [9,10*]. While earlier studies reported “global hypomethylation” in tumors, more recent comparisons to normal B cell differentiation stages revealed that these widespread changes parallel those of each tumor type’s “cell of origin” (Figure 2A) [11,12]. Focusing instead on differences between B cell tumors and matched-stage normal cells demonstrated that altered DNA methylation is concentrated at enhancers, and particularly enriched in the binding sites of stage- and tumor-specific TFs that are crucial for B cell maturation and activation (Figure 2B) [2,12,13]. Importantly, methylation changes in these regulatory regions correspond to alterations in the transcriptional programs of the associated TFs [11,14].

The underlying cause(s) of DNA methylation changes in B cell cancers is less clear. In GC B cells and GC-derived lymphomas, activation-induced cytosine deaminase (AID), which drives somatic hypermutation, an essential step in affinity maturation, also mediates DNA hypomethylation and increased methylome heterogeneity in intergenic and distal regions associated with essential B lineage genes (Figure 1B) [3,15,16]. Over-expression of AID accelerates a mouse model of BCL2-driven lymphoma without increasing off-target mutations [17,18*]. However, most B cell cancers do not overexpress AID; mutations in DNA methylation and demethylation factors are not common – though chromatin modifiers are frequently mutated; and there is little association between genetic mutation and DNA methylation profiles (reviewed in [11]). Instead, the coincidence of DNA methylome changes with B cell TF binding sites and B lineage enhancers in all B cell cancers suggest a common mechanism, and one that is well-established in cancer: perturbations in the activities of key TFs and chromatin modifiers involved in B cell activation and maturation are major contributors to aberrant methylation patterns.

The 3-dimensional (3D) interactome and chromatin modifiers mediate transcriptional programs integral to B cell activation

Globally increased transcription and histone acetylation were linked to B cell activation more than 50 years ago [19]. More recently, integrative analysis of activating and repressive histone marks, chromatin accessibility, and gene expression have defined regulatory landscape transitions in normal development from human or murine naïve B through plasma cells, and have revealed key regulatory elements that are reprogrammed in human B cell cancers and may be prognostic [20**,21*–23]. The mechanisms driving these epigenetic changes have been elucidated through B cell specific knockout of histone modifier genes, many of which are recurrently mutated in B cell cancers (Figure 2C) [24–27]. Conditional deletion of the histone acetyltransferase CREBBP perturbed B cell development and accelerated the development of lymphoma in BCL2- and MYC-driven mouse models [28,29,30*,31]. These effects are likely related to CREBBP's role in programs crucial to the GC reaction, including counteracting the repressive effects of BCL6 on transcription via direct acetylation of the BCL6 protein itself, and by H3K27 acetylation at promoters and enhancers of BCL6 target genes [25,28,29]. Several histone methyltransferases are also commonly mutated in NHL and crucial for B cell development and GC formation [24,25]. EZH2, a component of the polycomb repressive complex 2 (PRC2) that mediates trimethylation of H3K27 (H3K27me3), interacts with a distinct set of TFs to direct stage-specific gene repression programs, such as BCL6 in GC B cells and Blimp-1/PRDM1 in plasma cells (PC) [32,33]. Conditional ablation of EZH2 in B cells led to inappropriate activation of immune response pathways, B cell transcription factors, and regulators of cell division and metabolism normally repressed during terminal differentiation of PC, resulting in decreased proliferation and poor antibody production [22,34]. Thus, several chromatin modifiers coordinate with key B cell TFs to temporally regulate developmental transcriptional programs, and, when mutated, are lymphomagenic.

In concert with dynamic epigenetic landscapes, higher order genome organization and nuclear sub-compartmentalization orchestrate transcriptional activity [35]. Chromosome conformation capture and super-resolution microscopy methods have enabled mapping of the dramatic reorganization of 3D genome architecture during normal B cell development and activation [36]. These changes include chromatin decondensation via amplification of histone acetylation and nucleosome spreading, as well as chromatin decompaction mediated by new short-range CTCF loop interactions (Figure 1C) [20]. The latter changes facilitate increased promoter-enhancer contacts between, and augmented expression of, B cell activation genes and require the proto-oncogene Myc. Additional studies corroborated these findings and implicated lineage-specific transcription and non-coding regulatory regions as contributors to these changes in chromatin architecture [36,37**]. In GC B cells, a large (>100kb) enhancer cluster upstream of BCL6 comprises a locus control region (LCR) that forms the crux of intra-chromosomal interactions with promoters and enhancers of hundreds of upregulated GC B cell-specific genes [37**]. Strikingly, this BCL6 LCR was necessary for GC formation, but was dispensable for BCL6 expression in non-B cells [37**]. Taken together, these epigenomic findings underscore the complex choreography of chromatin landscape and 3D-genome organization with TF identity, level and timing that is crucial for normal B cell differentiation.

The roles of long non-coding RNAs in B cell activation and lymphomagenesis

LncRNAs are a heterogeneous group of ncRNAs that exhibit remarkable cell type specificity and disease association. They are differentiated from other ncRNAs (miRNA, tRNA, etc.) based on their length (>200 nucleotides), and are sub-classified by genomic location (e.g., intergenic, antisense, and enhancer) and structure (e.g., un-/spliced, uni-/bi-directionally transcribed, circRNAs) [38*]. LncRNA's functional versatility includes epigenetic modification, nuclear domain organization, transcriptional control, regulation of RNA splicing and translation, and modulating protein activity (reviewed in [39]).

Developmentally-controlled and tissue-specific production of lncRNA “germline” transcripts from unrearranged VH and VK gene segments in B lineage cells was identified nearly four decades ago [40] and was subsequently linked to enhanced chromatin accessibility and recombination of VDJ loci (reviewed in [41]). More recently, complementary roles in class-switch recombination (CSR) and somatic hypermutation (SHM) have been shown for intronic and anti-sense lncRNAs derived from switch regions and divergent transcription at promoters and enhancers. These lncRNAs create triplex RNA/DNA hybrid R-loops and G-quadruplexes, recruiting AID, the RNA exosome complex, and the RNA helicase DDX1, thereby facilitating CSR (Figure 1D) [42*–44].

Some super-enhancer loci, such as *MYC* and *BCL6*, also exhibit high levels of divergent and convergent lncRNA transcription, which may underlie off-target AID-mediated hypermutation and translocations that are oncogenic [16,45,46].

Several profiling studies have catalogued the patterns of lncRNA expression across B cell development and activation in human and mouse. One surprising finding is the loss of the repressive lncRNA Xist from the inactive X (Xi) chromosome at the pro-B cell stage, leading to loss of heterochromatin and re-expression of immune genes from the Xi. Activation of mature B cells restored Xist RNA and heterochromatin to the Xi. These findings suggest that chromatin changes on the Xi during B cell development and the dynamic nature of Xi maintenance in mature B cells predisposes X-linked immunity genes to reactivation [47,48]. Across B cell differentiation, hundreds of lncRNAs are expressed in a cell-type/stage-specific manner, which is consistent with studies in other tissues and suggests a *cis*-regulatory function for many of them [38,49*–55]. Compared to coding genes, lncRNAs exhibit significantly less sequence conservation, but alternative comparative methods, such as structure, profile, or positional conservation can detect homologues without primary sequence conservation [56]. In support of this approach, only 185 of 4516 lncRNAs expressed during murine B cell development had a human syntenic ortholog based on pairwise genomic alignments, but positional comparison of murine B cell eRNAs near orthologous coding genes revealed an additional 228 potentially orthologous lncRNAs [47]. One such syntenic lncRNA, RP11–132N15.3, lies in an intergenic enhancer region ~200kb upstream of *BCL6* within a cluster of lncRNAs that is highly and specifically expressed in both mouse and human GC B cells and strongly correlates with expression of the GC transcription factor *BCL6* [23,37**,47,52]. The robust expression and conservation of this lncRNA suggest that it is involved in the *cis*-regulation of *BCL6* expression. To answer this question, future experiments are needed that compare suppression and truncation of the

lncRNA transcript to epigenetic repression of the enhancer locus, to determine the impact on BCL6 expression.

Profiling and clinical retrospective studies in many lymphoma and leukemia subtypes identified altered expression of lncRNAs, suggesting a potential oncogenic role. MINCR (MYC-induced long noncoding RNA) is upregulated in MYC -rearranged or amplified primary lymphomas and cell lines. Knock-down of MINCR reduced MYC binding to the promoters of cell cycle genes and decreased their expression, impairing cell cycle progression [57]. Another lncRNA gene, PVT1, plays essential roles in the regulation of MYC-mediated growth and oncogenesis via epigenetic mechanisms (promoter competition) and direct stabilization of MYC protein levels (Figure 3) [58**,59]. Other lncRNAs, such as MALAT1, lncRNA-p21 (PANDAR), and NEAT1, have significant associations with clinicopathologic metrics of increased disease severity and decreased survival in lymphomas and MM and may be promising therapeutic targets [50,60,61]. NEAT1 is induced by p53 in response to DNA damage in CLL and lymphoma; primary CLL cells with low NEAT1 expression exhibited reduced cell death in response to DNA damage, even if TP53 was unmutated [60]. Depletion of NEAT1 in non-lymphoid cancer cells induced synthetic lethality with genotoxic chemotherapies as well as with activation of p53, suggesting that NEAT1 may be a promising target in B cell cancers as well [62]. The novel mechanisms of action exhibited by the few lncRNAs studied to date strongly suggest that these molecules hold vast potential for greater understanding of B cell biology, and new avenues for therapeutic development.

Conclusion

Significant progress has been made in charting the dynamic, non-coding portions of the epigenome and transcriptome in normal B lymphocytes and B cell cancers. However, outstanding questions remain. Though performed on sorted cells, the profiling studies to date are bulk analyses, which can only detect the average epigenomes and transcriptomes of component cells. Given the heterogeneity of differentiating B cells, single cell transcriptional and epigenetic profiling at many time points is necessary to definitively determine the causative links between temporally regulated epigenetic modifications, binding of regulatory factors, and gene expression levels. In addition, to more accurately elucidate their roles in lymphomagenesis, it is necessary to perturb multiple of epigenetic modifiers and transcription factors in a single model in the sequence that they are known to occur in human lymphomas. Finally, arguably the largest outstanding question is defining the functions of thousands of long non-coding RNAs with dynamic transcriptional patterns during B cell differentiation and/or in B cell cancers.

Acknowledgments

Funding

This work was supported by the National Institutes of Health [F31CA221012 to J. .; R01CA188286 to J.P.].

References

1. Lee S-T, Xiao Y, Muench MO, Xiao J, Fomin ME, Wiencke JK, Zheng S, Dou X, de Smith A, Chokkalingam A, et al.: A global DNA methylation and gene expression analysis of early human B-cell development reveals a demethylation signature and transcription factor network. *Nucleic Acids Res* 2012, 40:11339–11351. [PubMed: 23074194]
2. Kulis M, Merkel A, Heath S, Queirós AC, Schuyler RP, Castellano G, Beekman R, Raineri E, Esteve A, Clot G, et al.: Whole-genome fingerprint of the DNA methylome during human B cell differentiation. *Nat Genet* 2015, 47:746–756. [PubMed: 26053498]
3. Martin-Subero JI, Oakes CC: Charting the dynamic epigenome during B-cell development. *Semin Cancer Biol* 2017, doi:10.1016/j.semcancer.2017.08.008.
4. Almamun M, Levinson BT, Gater ST, Schnabel RD, Arthur GL, Davis JW, Taylor KH: Genome-wide DNA methylation analysis in precursor B-cells. *Epigenetics* 2014, 9:1588–1595. [PubMed: 25484143]
5. Barwick BG, Scharer CD, Bally APR, Boss JM: Plasma cell differentiation is coupled to division-dependent DNA hypomethylation and gene regulation. *Nat Immunol* 2016, 17:1216–1225. [PubMed: 27500631]
6. Lio C-W, Zhang J, González-Avalos E, Hogan PG, Chang X, Rao A: Tet2 and Tet3 cooperate with B-lineage transcription factors to regulate DNA modification and chromatin accessibility. *eLife* 2016, 5.
7. Orlanski S, Labi V, Reizel Y, Spiro A, Lichtenstein M, Levin-Klein R, Koralov SB, Skversky Y, Rajewsky K, Cedar H, et al.: Tissue-specific DNA demethylation is required for proper B-cell differentiation and function. *Proc Natl Acad Sci* 2016, 113:5018–5023. [PubMed: 27091986]
- 8*. Barwick BG, Scharer CD, Martinez RJ, Price MJ, Wein AN, Haines RR, Bally APR, Kohlmeier JE, Boss JM: B cell activation and plasma cell differentiation are inhibited by de novo DNA methylation. *Nat Commun* 2018, 9:1900. [PubMed: 29765016] Using a conditional Dnmt3a/b knockout mouse, the authors mapped de novo DNA methylation and its impact on B cell differentiation and plasma cell function.
9. Dominguez PM, Teater M, Shaknovich R: The new frontier of epigenetic heterogeneity in B-cell neoplasms. *Curr Opin Hematol* 2017, 24:402. [PubMed: 28375986]
- 10*. Flavahan WA, Gaskell E, Bernstein BE: Epigenetic plasticity and the hallmarks of cancer. *Science* 2017, 357:eaal2380. [PubMed: 28729483] This review provides a comprehensive look at the deregulation of the epigenome in cancer and argues that epigenetic aberrations contribute to all of the classic hallmarks of cancer.
11. Oakes CC, Martin-Subero JI: Insight into origins, mechanisms & utility of DNA methylation in B cell malignancies. *Blood* 2018, doi:10.1182/Blood-2018-02-692970.
12. Oakes CC, Seifert M, Assenov Y, Gu L, Przekopowicz M, Ruppert AS, Wang Q, Imbusch CD, Serva A, Koser SD, et al.: DNA methylation dynamics during B cell maturation underlie a continuum of disease phenotypes in chronic lymphocytic leukemia. *Nat Genet* 2016, 48:253–264. [PubMed: 26780610]
13. Queirós AC, Beekman R, Vilarrasa-Blasi R, Duran-Ferrer M, Clot G, Merkel A, Raineri E, Russiñol N, Castellano G, Beà S, et al.: Decoding the DNA Methylome of Mantle Cell Lymphoma in the Light of the Entire B Cell Lineage. *Cancer Cell* 2016, 30:806–821. [PubMed: 27846393]
14. Almamun M, Levinson BT, Swaay AC van, Johnson NT, McKay SD, Arthur GL, Davis JW, Taylor KH: Integrated methylome and transcriptome analysis reveals novel regulatory elements in pediatric acute lymphoblastic leukemia. *Epigenetics* 2015, 10:882–890. [PubMed: 26308964]
15. Dominguez PM, Teater M, Chambwe N, Kormaksson M, Redmond D, Ishii J, Vuong B, Chaudhuri J, Melnick A, Vasanthakumar A, et al.: DNA Methylation Dynamics of Germinal Center B Cells Are Mediated by AID. *Cell Rep* 2015, 12:2086–2098. [PubMed: 26365193]
16. Casellas R, Basu U, Yewdell WT, Chaudhuri J, Robbiani DF, Di Noia JM: Mutations, kataegis, and translocations in B lymphocytes: towards a mechanistic understanding of AID promiscuous activity. *Nat Rev Immunol* 2016, 16:164–176. [PubMed: 26898111]

17. Dominguez MDP, Teater M, Redmond D, Cheng Z, Chaudhuri J, Elemento O, Melnick A, Shklovich R: AICDA Introduces Epigenetic Plasticity in Germinal Center-Derived Lymphomas and Accelerates Lymphomagenesis. *Blood* 2016, 128:1045–1045.
- 18*. Teater M, Dominguez PM, Redmond D, Chen Z, Ennishi D, Cott DW, Cimmino L, Ghione P, Chaudhuri J, Gascoyne RD, et al.: AICDA drives epigenetic heterogeneity and accelerates germinal center-derived lymphomagenesis. *Nat Commun* 2018, 9:222. [PubMed: 29335468] Using a BCL2-driven lymphoma mouse model, this work showed that AID contributes to increased disease severity and drives DNA methylome heterogeneity.
19. Pogo BG, Allfrey VG, Mirsky AE: RNA synthesis and histone acetylation during the course of gene activation in lymphocytes. *Proc Natl Acad Sci* 1966, 55:805–812. [PubMed: 5219687]
- 20*. Kieffer-Kwon K-R, Nimura K, Rao SSP, Xu J, Jung S, Pekowska A, Dose M, Stevens E, Mathe E, Dong P, et al.: Myc Regulates Chromatin Decompaction and Nuclear Architecture during B Cell Activation. *Mol Cell* 2017, 67:566–578.e10. [PubMed: 28803781] This work revealed the importance of chromatin remodeling as a general mechanism of B cell activation. This mechanism is regulated by Myc and leads to dramatic chromatin decompaction and increased short-range chromatin interactions.
- 21*. Beekman R, Chapaprieta V, Russiñol N, Vilarasa-Blasi R, Verdaguer-Dot N, Martens JHA, Duran-Ferrer M, Kulis M, Serra F, Javierre BM, et al.: The reference epigenome and regulatory chromatin landscape of chronic lymphocytic leukemia. *Nat Med* 2018, 24:868–880. [PubMed: 29785028] This resource provides one of the most complete integrative datasets of both normal B cells and chronic lymphocytic leukemia to date.
22. Scharer CD, Barwick BG, Guo M, Bally APR, Boss JM: Plasma cell differentiation is controlled by multiple cell division-coupled epigenetic programs. *Nat Commun* 2018, 9:1698. [PubMed: 29703886]
23. Koues OI, Kowalewski RA, Chang L-W, Pyfrom SC, Schmidt JA, Luo H, Sandoval LE, Hughes TB, Bednarski JJ, Cashen AF, et al.: Enhancer sequence variants and transcription-factor deregulation synergize to construct pathogenic regulatory circuits in B-cell lymphoma. *Immunity* 2015, 42:186–198. [PubMed: 25607463]
24. Green MR: Chromatin modifying gene mutations in follicular lymphoma. *Blood* 2018, 131:595–604. [PubMed: 29158360]
25. Pasqualucci L, Dalla-Favera R: Genetics of diffuse large B-cell lymphoma. *Blood* 2018, 131:2307–2319. [PubMed: 29666115]
26. Puente XS, Jares P, Campo E: Chronic lymphocytic leukemia and mantle cell lymphoma: Crossroads of genetic and microenvironment interactions. *Blood* 2018, doi:10.1182/blood-2017-10-764373.
27. Gascoyne RD, Nadel B, Pasqualucci L, Fitzgibbon J, Payton JE, Melnick A, Weigert O, Tarte K, Gribben JG, Friedberg JW, et al.: Follicular lymphoma: State-of-the-art ICML workshop in Lugano 2015. *Hematol Oncol* 2017, 35:397–407. [PubMed: 28378425]
28. Zhang J, Vlasevska S, Wells VA, Nataraj S, Holmes AB, Duval R, Meyer SN, Mo T, Basso K, Brindle PK, et al.: The CREBBP Acetyltransferase Is a Haploinsufficient Tumor Suppressor in B-cell Lymphoma. *Cancer Discov* 2017, doi:10.1158/2159-8290.CD-16-1417.
29. Jiang Y, Ortega-Molina A, Geng H, Ying H-Y, Hatzi K, Parsa S, McNally D, Wang L, Doane AS, Agirre X, et al.: CREBBP Inactivation Promotes the Development of HDAC3-Dependent Lymphomas. *Cancer Discov* 2017, 7:38–53. [PubMed: 27733359]
- 30*. Hashwah H, Schmid CA, Kasser S, Bertram K, Stelling A, Manz MG, Müller A: Inactivation of CREBBP expands the germinal center B cell compartment, down-regulates MHCII expression and promotes DLBCL growth. *Proc Natl Acad Sci* 2017, 114:9701–9706. [PubMed: 28831000] This paper, along with [29–31], highlight the critical nature of master epigenetic modifiers for proper B cell development and how their deregulation can promote B cell oncogenesis.
31. García-Ramírez I, Tadros S, González-Herrero I, Martín-Lorenzo A, Rodríguez-Hernández G, Moore D, Ruiz-Roca L, Blanco O, Alonso-López D, Rivas JDL, et al.: Crebbp loss cooperates with Bcl2 over-expression to promote lymphoma in mice. *Blood* 2017, doi:10.1182/blood-2016-08-733469.

32. Minnich M, Tagoh H, Bönelt P, Axelsson E, Fischer M, Cebolla B, Tarakhovsky A, Nutt SL, Jaritz M, Busslinger M: Multifunctional role of the transcription factor Blimp-1 in coordinating plasma cell differentiation. *Nat Immunol* 2016, 17:331–343. [PubMed: 26779602]
33. Béguelin W, Teater M, Gearhart MD, Calvo Fernández MT, Goldstein RL, Cárdenas MG, Hatzi K, Rosen M, Shen H, Corcoran CM, et al.: EZH2 and BCL6 Cooperate to Assemble CBX8-BCOR Complex to Repress Bivalent Promoters, Mediate Germinal Center Formation and Lymphomagenesis. *Cancer Cell* 2016, 30:197–213. [PubMed: 27505670]
34. Guo M, Price MJ, Patterson DG, Barwick BG, Haines RR, Kania AK, Bradley JE, Randall TD, Boss JM, Scharer CD: EZH2 Represses the B Cell Transcriptional Program and Regulates Antibody-Secreting Cell Metabolism and Antibody Production. *J Immunol* 2018, 200:1039–1052. [PubMed: 29288200]
35. Cuartero S, Merkenschlager M: Three-dimensional genome organization in normal and malignant haematopoiesis. *Curr Opin Hematol* 2018, 25:323. [PubMed: 29702522]
36. Boya R, Yadavalli AD, Nikhat S, Kurukuti S, Palakodeti D, Pongubala JMR: Developmentally regulated higher-order chromatin interactions orchestrate B cell fate commitment. *Nucleic Acids Res* 2017, 45:11070–11087. [PubMed: 28977418]
- 37*. Bunting KL, Soong TD, Singh R, Jiang Y, Béguelin W, Poloway DW, Swed BL, Hatzi K, Reisacher W, Teater M, et al.: Multi-tiered Reorganization of the Genome during B Cell Affinity Maturation Anchored by a Germinal Center-Specific Locus Control Region. *Immunity* 2016, 45:497–512. [PubMed: 27637145] This study identified a locus control region upstream of BCL6 and revealed its role in the dramatic remodeling of high-order chromatin interactions necessary for B cell activation and proper germinal center formation.
- 38*. Chen YG, Satpathy AT, Chang HY: Gene regulation in the immune system by long noncoding RNAs. *Nat Immunol* 2017, 18:962–972. [PubMed: 28829444] This excellent review provides a detailed survey of mechanistic studies of lncRNA as transcriptional regulators in immune cells.
39. Marchese FP, Raimondi I, Huarte M: The multidimensional mechanisms of long noncoding RNA function. *Genome Biol* 2017, 18:206. [PubMed: 29084573]
40. Yancopoulos GD, Alt FW: Developmentally controlled and tissue-specific expression of unrearranged VH gene segments. *Cell* 1985, 40:271–81. [PubMed: 2578321]
41. Cobb RM, Oestreich KJ, Osipovich OA, Oltz EM: Accessibility control of V(D)J recombination. *Adv Immunol* 2006, 91:45–109 [PubMed: 16938538]
- 42*. Ribeiro de Almeida C, Dhir S, Dhir A, Moghaddam AE, Sattentau Q, Meinhart A, Proudfoot NJ: RNA Helicase DDX1 Converts RNA G-Quadruplex Structures into R-Loops to Promote IgH Class Switch Recombination. *Mol Cell* 2018, 70:650–662.e8. [PubMed: 29731414] In combination with [43], this work showed the importance of lncRNA transcription at switch regions for proper class switch recombination through the formation of R-loops targeted by AID.
43. Abdouni HS, King JJ, Ghorbani A, Fifield H, Berghuis L, Larijani M: DNA/RNA hybrid substrates modulate the catalytic activity of purified AID. *Mol Immunol* 2018, 93:94–106. [PubMed: 29161581]
44. Yewdell WT, Chaudhuri J: A transcriptional serenAID: the role of noncoding RNAs in class switch recombination. *Int Immunol* 2017, 29:183–196. [PubMed: 28535205]
45. Zan H, Casali P: Epigenetics of Peripheral B-Cell Differentiation and the Antibody Response. *B Cell Biol* 2015, doi:10.3389/fimmu.2015.00631.
46. Boller S, Li R, Grosschedl R: Defining B Cell Chromatin: Lessons from EBF1. *Trends Genet* 2018, 34:257–269. [PubMed: 29336845]
47. Syrett CM, Sindhava V, Hodawadekar S, Myles A, Liang G, Zhang Y, Nandi S, Cancro M, Atchison M, Anguera MC: Loss of Xist RNA from the inactive X during B cell development is restored in a dynamic YY1-dependent two-step process in activated B cells. *PLOS Genet* 2017, 13:e1007050. [PubMed: 28991910]
48. Wang J, Syrett CM, Kramer MC, Basu A, Atchison ML, Anguera MC: Unusual maintenance of X chromosome inactivation predisposes female lymphocytes for increased expression from the inactive X. *Proc Natl Acad Sci* 2016, doi:10.1073/pnas.1520113113.
- 49*. Brazão TF, Johnson JS, Müller J, Heger A, Ponting CP, Tybulewicz VLJ: Long noncoding RNAs in B-cell development and activation. *Blood* 2016, 128:e10–e19. [PubMed: 27381906] Through

expression profiling via RNA-seq of B cells at several stages of development, this work developed a map of temporally regulated lncRNAs during B cell differentiation that may serve as a springboard for further mechanistic studies.

50. Winkle M, Kluiver JL, Diepstra A, van den Berg A: Emerging roles for long noncoding RNAs in B-cell development and malignancy. *Crit Rev Oncol Hematol* 2017, 120:77–85. [PubMed: 29198340]
51. Casero D, Sandoval S, Seet CS, Scholes J, Zhu Y, Ha VL, Luong A, Parekh C, Crooks GM: Long non-coding RNA profiling of human lymphoid progenitor cells reveals transcriptional divergence of B cell and T cell lineages. *Nat Immunol* 2015, 16:1282–1291 [PubMed: 26502406]
52. Tayari MM, Winkle M, Kortman G, Sietzema J, de Jong D, Terpstra M, Mestdagh P, Kroese FGM, Visser L, Diepstra A, et al.: Long Noncoding RNA Expression Profiling in Normal B-Cell Subsets and Hodgkin Lymphoma Reveals Hodgkin and ReedSternberg Cell-Specific Long Noncoding RNAs. *Am J Pathol* 2016, 186:2462–2472. [PubMed: 27423697]
53. Ranzani V, Rossetti G, Panzeri I, Arrigoni A, Bonnal RJP, Curti S, Gruarin P, Provati E, Sugliano E, Marconi M, et al.: The long intergenic noncoding RNA landscape of human lymphocytes highlights the regulation of T cell differentiation by linc-MAF-4. *Nat Immunol* 2015, 16:318–325 [PubMed: 25621826]
54. Petri A, Dybkær K, Bøgsted M, Thue CA, Hagedorn PH, Schmitz A, Bødker JS, Johnsen HE, Kauppinen S: Long Noncoding RNA Expression during Human B-Cell Development. *PLoS ONE* 2015, 10:e0138236. [PubMed: 26394393]
55. Liu SJ, Horlbeck MA, Cho SW, Birk HS, Malatesta M, He D, Attenello FJ, Villalta JE, Cho MY, Chen Y, et al.: CRISPRi-based genome-scale identification of functional long noncoding RNA loci in human cells. *Science* 2017, 355:eaah7111.
56. Ulitsky I: Evolution to the rescue: using comparative genomics to understand long non-coding RNAs. *Nat Rev Genet* 2016, 17:601–614. [PubMed: 27573374]
57. Dooze G, Haake A, Bernhart SH, López C, Duggimpudi S, Wojciech F, Bergmann AK, Borkhardt A, Burkhardt B, Claviez A, et al.: MINCR is a MYC-induced lncRNA able to modulate MYC's transcriptional network in Burkitt lymphoma cells. *Proc Natl Acad Sci* 2015, 112:E5261–E5270. [PubMed: 26351698]
- 58*. Cho SW, Xu J, Sun R, Mumbach MR, Carter AC, Chen YG, Yost KE, Kim J, He J, Nevins SA, et al.: Promoter of lncRNA Gene PVT1 Is a Tumor-Suppressor DNA Boundary Element. *Cell* 2018, 173:1398–1412.e22. [PubMed: 29731168] This study provides an exquisite example of the intricate interplay between genetic perturbations, regulatory sequences, and 3D genome structure in gene regulation. The authors utilized CRISPRi-methods to reveal tumor suppressor function of the promoter of the oncogenic lncRNA PVT1.
59. Tseng Y-Y, Moriarity BS, Gong W, Akiyama R, Tiwari A, Kawakami H, Ronning P, Reuland B, Guenther K, Beadnell TC, et al.: PVT1 dependence in cancer with MYC copy-number increase. *Nature* 2014, 512:82–86. [PubMed: 25043044]
60. Blume CJ, Hotz-Wagenblatt A, Hüllein J, Sellner L, Jethwa A, Stolz T, Slabicki M, Lee K, Sharathchandra A, Benner A, et al.: p53-dependent non-coding RNA networks in chronic lymphocytic leukemia. *Leukemia* 2015, 29:2015–2023. [PubMed: 25971364]
61. Amodio N, Stamato MA, Juli G, Morelli E, Fulciniti M, Manzoni M, Taiana E, Agnelli L, Cantafio MEG, Romeo E, et al.: Drugging the lncRNA MALAT1 via LNA gapmeR ASO inhibits gene expression of proteasome subunits and triggers anti-multiple myeloma activity. *Leukemia* 2018, doi:10.1038/s41375-018-0067-3.
62. Adriaens C, Standaert L, Barra J, Latil M, Verfaillie A, Kalev P, Boeckx B, Wijnhoven PWG, Radaelli E, Vermi W, et al.: p53 induces formation of *NEAT1* lncRNA-containing paraspeckles that modulate replication stress response and chemosensitivity. *Nat Med* 2016, 22:861–868. [PubMed: 27376578]

Highlights

- B cell development requires coordinated, sequential changes in epigenome landscapes.
- 3-D nuclear re-organization is essential for B cell differentiation.
- LncRNAs are cis-regulators of essential B cell factors.
- Epigenetic deregulation is a cornerstone of B cell oncogenesis.

Figure 1. Epigenetic mechanisms regulating B cell activation and the GC reaction.

A) Simplified diagram shows the stages of peripheral B cell activation and maturation in the GC reaction. B) DNA hypomethylation leads to methylation heterogeneity during B cell maturation. As naïve B cells are activated and form GCs, i) increased histone acetylation and loss of repressive methylation drastically alter the chromatin landscape, which ii) decompacts chromatin and generates chromatin loops, iii) facilitating increased promoter-enhancer contacts and a massive increase in gene expression. D) lncRNA transcription at switch regions contributes to SHM and CSR through two mechanisms; i) R-loops -- DNA-RNA hybrid and ssDNA -- are formed during transcription and the ssDNA can be targeted by AID or ii) switch transcripts form G-quadruplexes that are bound by AICDA and unwound by DDX1 to facilitate R-loop formation at switch regions. SHM – somatic hypermutation; CSR – class-switch recombination; BCR – B cell receptor.

Figure 2. Subversion of epigenetic mechanisms in B cell cancers.

A) Overview of the cell of origin for many B cell cancers. B) B cell lymphomas exhibit increased intra- and inter-sample DNA methylation heterogeneity due to i) aberrant hypomethylation at major transcription factor and AICDA binding sites, and ii) aberrant histone and DNA hypermethylation at gene promoters regulatory elements that may result in transcriptional repression. C) Frequencies of select mutations in epigenetic modifiers, chromatin remodelers, and histone proteins in several B cell cancers. ABC-DLBCL – Activated B-cell-like subtype of diffuse large B cell lymphoma; GCB-DLBCL – Germinal

center B-cell-like subtype of diffuse large B cell lymphoma; U-CLL – unmutated chronic lymphocytic leukemia; M-CLL – mutated chronic lymphocytic leukemia; FL – Follicular lymphoma; MM – Multiple myeloma; BL - Burkitt lymphoma; MCL – Mantle cell lymphoma.

Figure 3. The complex, synergistic interplay between genetic and epigenetic aberrations of the *PVT1/MYC* locus.

i) The *MYC* and *PVT1* promoters exhibit competitive binding to a group of enhancers (orange bars) located within the *PVT1* gene body. ii) Mutations to the *PVT1* promoter eliminate its interaction with this regulatory cluster, allowing the *MYC* promoter to freely interact with the enhancers and increase *MYC* expression. iii) Independently, IgH promoter/enhancer translocations or amplifications of the locus can result in *PVT1* overexpression, which binds to the *MYC* protein and prevents its degradation.