Published in final edited form as:

Curr Opin Physiol. 2018 October ; 5: 75–79. doi:10.1016/j.cophys.2018.08.002.

The circadian clock: A tale of genetic-electrical interplay and synaptic integration

Mino D C Belle¹ and Charles N Allen²

¹Institute of Biomedical and Clinical Sciences, University of Exeter Medical School, University of Exeter, Exeter, EX4 4PS, UK. m.d.c.belle@exeter.ac.uk

²Oregon Institute of Occupational Health Sciences and Department of Behavioral Neuroscience, Oregon Health & Science University, Portland, Oregon 97239, USA. allenc@ohsu.edu

Abstract

Pioneering work in *Drosophila* uncovered the building blocks of the molecular clock, consisting of transcription-translation feedback loops (TTFLs). Subsequent experimental work demonstrated that the mammalian TTFL is localized in cells and tissues throughout the brain and body. Further research established that neuronal activity forms an essential aspect of clock function. However, how the membrane electrical activity of clock neurons of the suprachiasmatic nucleus collaborate with the TTFL to drive circadian behaviors remains mostly unknown. Intercellular communication synchronizes the individual circadian oscillators to produce a precise and coherent circadian output. Here, we briefly review significant research that is increasing our understanding of the critical interactions between the TTFL and neuronal and glial activity in the generation of circadian timing signals.

Keywords

Suprachiasmatic nucleus; circadian rhythms; clock genes; electrical activity; ion channels

TTFL and membrane signaling

Individual neurons in the suprachiasmatic nucleus (SCN) contain molecular circadian clocks, consisting of the *Period1/2* (*Per1/2*), *Cryptochrome1/2* (*Cry1/2*), *Clock* and *Bmal1* genes. The molecular circadian signal is translated into a circadian pattern of action potential firing **and, in some SCN neurons, electrical silencing by hyperand hypo-excitation** [1], which is required for the generation of circadian behaviors. The interaction between SCN

Correspondence: Belle, Mino (m.d.c.belle@exeter.ac.uk).

The authors declare no conflicts of interest.

Nothing declared.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

neuron electrical activity and the molecular clock, however, is more **complicated** than just a driven rhythm.

Inhibiting action potential firing in the SCN with tetrodotoxin (TTX) abolishes behavioral circadian rhythmicity but does not affect the timing of the circadian clock [2]. Therefore, SCN action potential firing is an output of the circadian clock required to drive circadian behaviors, but pacemaker timing persists in the absence of action potential activity at the single cell level. However, TTX application to SCN brain slices for an extended period of time produces a rundown of the molecular clock amplitude [3]. This suggests that for proper clock functioning the membrane electrical excitability feeds back onto the molecular clock machinery to support and maintain circadian oscillations [3]. Two **essential** and unresolved experimental questions in circadian biology are how the molecular clock communicates with the membrane ion channels that regulate membrane excitability, and how the membrane electrical activity signals to the molecular circadian clock. Knowledge of these mechanisms will provide insights into how the molecular clock appropriately synchronizes its activity across the SCN network and outputs its collective phase to downstream targets. In addition, this information will provide a mechanistic understanding of how environmental and internal physiological signals influence the timing precision in SCN clock cells (Figure 1). Indeed, an ingenious study in flies has unequivocally shown that the membrane activity feeds back to impose time-of-day stamps onto the molecular clock programs [4]. **In mammals, the *Fbxl3*^{Afh} mutation delays the degradation of CRY1/2 and increases the circadian period [5]. The disrupted molecular clock alters the membrane excitability and GABA neurotransmission of the SCN neurons.**

Recent work has provided new insights into possible signaling pathways regulating the circadian rhythm in SCN neuronal activity. It had been postulated that the **excitable states** represent a balance between the depolarizing activity of voltagegated sodium channels and hyperpolarizing potassium channels [6–8]. Recently, a voltage-independent sodium conductance, mediated by the NA/NALCN ion channel, has been shown to depolarize SCN neurons [9]. This current is driven by the rhythmic expression of **NCA Localization Factor-1**, providing an example of signaling pathway linking the molecular clock to ion channel function [9]. This channel pathway is a new addition to the families of cation channels that provide depolarizing forces to SCN neurons during the day, elevating their resting membrane potential and increasing firing rate [10–12]. The L-type calcium channel is another key cation channel involved in sustaining excitation in SCN neurons [1,8]. Recently, L-type calcium channel activity has also been shown to be under the direct control of the TTFL component REV-ERBa [13], a negative feedback loop in the molecular circadian clock. This provides yet another example of a pathway where **the TTFL** can influence ion channel activity. Some potassium channels may reduce their conductivity to support such depolarization during the day [10–12]. For example, a reduction in the activity of the small-conductance calcium-activated potassium channels transits a proportion of daytime SCN neurons into hyperexcitation and depolarization blockade states, where they ceased firing [1,14].

In the evening, the activity of a number of potassium currents peaks to hyperpolarize and silence clock neurons [10–12]. Reduction of *Per1* activity by antisense oligonucleotides

suppressed firing of SCN neurons by reducing intracellular calcium levels and the conductance of the large calcium-activated potassium currents (BK; [15]). Indeed, the pharmacological blockade of BK channels mimicked the effects of the antisense on SCN firing rate [15]. Circadian clock regulation of BK channels and their biophysical properties, particularly the rate of inactivation, alters the subthreshold membrane properties contributing to the day-night firing rhythm [16,17]. This permits SCN neurons to readily fire action potentials during the day and make them less likely to spike at night. Additional evidence for this inactivation has also been provided for the fast-delayed rectifier potassium channels [18].

One of the **significant** challenges facing circadian biologists is that SCN neurons are exceedingly heterogeneous. Indeed, with this diversity comes complexity and the daunting task of categorizing and cataloguing neurons based on their intrinsic excitability [19]. That is, grouping neurons that show similar firing patterns when manually presented with depolarizing stimuli in the absence of synaptic communication. **This measure will naturally be aided by targeted recordings in identified SCN neuronal populations, using animals expressing appropriate genetic fluorescence reporters. For example, previous work in the SCN indicated that when presented with a depolarizing pulse, *Per1*-EGFP positive neurons (presumed clock neurons) show firing characteristics that are broadly different from neurons in which EGFP could not be detected (presumed nonclock cells) [1]. This form of neuronal targeting can be used in new models, such as the *Per1*-Venus mouse [20], and animals in which the SCN's peptidergic populations can also be labelled. It is noteworthy, however, that recent work suggests that *Per1*- and *Per2*-containing neurons in the SCN form different but overlapping neuronal population, adding complexity to analysis [21]. Nevertheless, we now also understand that similar classes of neurons can generate comparable action potential firing patterns using different complements of ion channels [22]. Ion channel expression is regulated by homeostatic mechanisms that couple channel expression to specific neuronal activity patterns [22]. Therefore, even similar types of SCN neurons may recruit subtly different ionic mechanisms in order to regulate the circadian activity, and these ion channel activity patterns may show day-night expression patterns.**

Cell to cell signaling

Indeed, the governing principle in SCN circadian rhythm generation is the maintenance of synchrony and appropriate phase among its neurons. GABAergic neurotransmission is a fundamental component of the SCN neural network, and virtually all SCN neurons communicate using GABA [23,24]. GABA regulates many functions in the SCN, including light-induced phase shifts, synchronization of the dorsal and ventral SCN, and the sensitivity of the circadian clock to light-entraining signals. The specific role of GABA neurotransmission in maintaining cellular synchrony remains controversial (see recent reviews by [23–25]). GABA acts on synaptic GABA_A receptors to mediate fast “phasic” signaling between SCN neurons, and extrasynaptic GABA_A receptor activation provides SCN cells with a “tonic” GABA_A current [26]. As a network, activity signaling by the tonic GABA-GABA_A receptor current is a strong candidate to regulate the coupling strength between individual SCN neuronal oscillators [27,28]. Modulation of the intracellular

chloride concentration ($[Cl^-]_i$) regulates the strength of the GABA currents by altering the magnitude of the current flowing through open GABA-activated channels at a given membrane potential [29]. In the adult SCN, GABA acts both as an inhibitory or excitatory neurotransmitter depending on the postsynaptic $[Cl^-]_i$ concentration and the time-of-day [29–33]. Excitatory GABA neurotransmission is more prevalent in the dorsal SCN compared to the ventral SCN, and the overall consensus is that there is more excitatory GABA transmission during the night than during the day [29–33]. The regional differences in GABA activity may reflect different intracellular Cl^- regulation in arginine vasopressin (AVP)- and vasoactive intestinal polypeptide (VIP)-expressing SCN neurons [34]. Computer simulations and physiological recordings suggest that this inhibitory/excitatory switch in GABA action is an essential component of the SCN network activity, regulating period length and photoperiod encoding [27,28,35].

Glia

Astrocytes in the SCN also express a molecular TTFL circadian clock and play an essential role in regulating activity and entrainment of the SCN clock [36–38]. There is one astrocyte for every three SCN neurons, and these glial cells have a soma and a large number of fine processes that encase the neurons and synapses [39,40]. These fine processes allow a single astrocyte to influence the activity of a significant number of synapses (Figure 1). Modulation of astrocyte activity can alter the timing of the circadian clock. In *Drosophila*, inhibition of intracellular calcium signaling in astrocytes disrupts functional circadian rhythms [41,42]. Genetic deletion of BMAL1 in astrocytes lengthens the circadian period and represses the rhythmic expression of VIP and several clock genes [37,38,43].

Astrocytes modify neuronal activity and neurotransmission through a number of mechanisms including the release of transmitter substances (gliotransmitters), such as adenosine triphosphate (ATP) and glutamate [44]. ATP is released by astrocytes in a circadian pattern with the maximum concentration observed in the middle of the night [45–47]. ATP can activate ionotropic P2X receptors to potentiate GABA release from SCN synapses and metabotropic P2Y receptors to inhibit GABA release [48,49]. Enzymatic conversion of ATP to adenosine leads to activation of presynaptic adenosine A1 receptors located on terminals of the retinohypothalamic tract (RHT). This reduces light-induced phase changes by decreasing glutamate release by the RHT [50,51]. These data indicate that the role of glial-released ATP depends on the location and timing of its release. SCN astrocytes also release glutamate in a circadian manner, with higher concentrations reported during the subjective night [37]. The glutamate may act on presynaptic NMDA receptors containing NR2C subunits to facilitate GABA release [37].

Conclusion and perspectives

Emerging work in the circadian field is revealing a collaborative, but intricate and complex relationship between the molecular clockwork and the electrical activity in SCN neurons. Progress in understanding the mechanistic nature of this relationship is slow because despite our vast understanding of the cell-autonomous processes causing daily oscillations in clock gene expression, our knowledge of how the molecular clockwork interacts with the

membrane to regulate **the** excitability of SCN neurons is severely lacking. Feedback cues from the environment and internal physiology signal to SCN neurons. **However**, here too, the mechanisms involved in this electrical-genetic interaction remain elusive. Despite these knowledge gaps, **the** intracellular calcium level and the activity of its associated signaling pathways are known to play important roles. Here, a **critical** observation is that both the intracellular and extracellular sources of calcium are important for circadian rhythm generation and communication in the SCN [52–55].

Indeed, evidence in mammals and *Drosophila* supports the concept that the plasma membrane is not merely the proximal target of the molecular clock, but its excitability is integral to clock function. Across this partnership also lies the glial circadian clock. As in neurons, the relationship between the molecular clock and glio-physiology remains poorly understood. Uncovering the signaling pathways and mechanisms involved are daunting challenges, but a necessary task if we are to understand how circadian rhythms are generated and communicated in the SCN and across the brain.

Acknowledgements

M.D.C.B is supported by the University of Exeter Medical School (UEMS) and the Royal Society (Grant: RSG \R1\180131). C. N. Allen is supported by the National Institute of Neurological Disorders and Stroke of the National Institutes of Health under award number R56NS036607.

We would like to acknowledge the excellent experimental work of many research groups that we are not able to reference due to space constraints.

References

1. Belle MD, Diekmann CO, Forger DB, Piggins HD: Daily electrical silencing in the mammalian circadian clock. *Science* 2009, 326:281–284. [PubMed: 19815775]
 2. Schwartz WJ, Gross RA, Morton MT: The suprachiasmatic nuclei contain a tetrodotoxin-resistant circadian pacemaker. *Proc Natl Acad Sci U S A* 1987, 84:1694–1698. [PubMed: 3470750]
 3. Yamaguchi S, Isejima H, Matsuo T, Okura R, Yagita K, Kobayashi M, Okamura H: Synchronization of cellular clocks in the suprachiasmatic nucleus. *Science* 2003, 302:1408–1412. [PubMed: 14631044]
 4. Mizrak D, Ruben M, Myers GN, Rhrissorakrai K, Gunsalus KC, Blau J: Electrical activity can impose time of day on the circadian transcriptome of pacemaker neurons. *Curr. Biol* 2012, 22:1871–1880. [PubMed: 22940468]
 - 5**. Wegner S, Belle MDC, Hughes ATL, Diekmann CO, Piggins HD: Delayed Cryptochrome Degradation Asymmetrically Alters the Daily Rhythm in Suprachiasmatic Clock Neuron Excitability. *J Neurosci* 2017, 37:7824–7836.
- The Fxlb^{Afh}, which changes the rate of CRY1/2 degradation produces changes in the regulation of neuronal excitability of SCN neurons by changing ion conductances and the strength of GABAergic neurotransmission.
- [PubMed: 28698388]
6. Kononenko NI, Honma S, Dudek FE, Honma K: On the role of calcium and potassium currents in circadian modulation of firing rate in rat suprachiasmatic nucleus neurons: multielectrode dish analysis. *Neurosci Res* 2008, 62:51–57. [PubMed: 18602427]
 7. Jiang ZG, Yang Y, Liu ZP, Allen CN: Membrane properties and synaptic inputs of suprachiasmatic nucleus neurons in rat brain slices. *J Physiol* 1997, 499 (Pt 1):141–159. [PubMed: 9061646]
 8. Jackson AC, Yao GL, Bean BP: Mechanism of spontaneous firing in dorsomedial suprachiasmatic nucleus neurons. *J Neurosci* 2004, 24:7985–7998. [PubMed: 15371499]

9**. Flourakis M, Kula-Eversole E, Hutchison AL, Han TH, Aranda K, Moose DL, White KP, Dinner AR, Lear BC, Ren D, et al.: A Conserved Bicycle Model for Circadian Clock Control of Membrane Excitability. *Cell* 2015, 162:836–848.

A NA/NALCN ion channel was identified that regulated circadian activity in *Drosophila* clock neurons. The authors then demonstrated in mice that a similar current is under circadian clock control and is responsible for the circadian activity of SCN neurons.

[PubMed: 26276633]

10. Allen CN, Nitabach MN, Colwell CS: Membrane Currents, Gene Expression, and Circadian Clocks. *Cold Spring Harb Perspect Biol* 2017, 9.
 11. Colwell CS: Linking neural activity and molecular oscillations in the SCN. *Nat. Rev. Neurosci* 2011, 12:553–569. [PubMed: 21886186]
 12. Belle MDC, Diekmann CO: Neuronal oscillations on an ultra-slow timescale: daily rhythms in electrical activity and gene expression in the mammalian master circadian clockwork. *Eur J Neurosci* 2018.
 13. Schmutz I, Chavan R, Ripperger JA, Maywood ES, Langwieser N, Jurik A, Stauffer A, Delorme JE, Moosmang S, Hastings MH, et al.: A specific role for the REV-ERB α -controlled L-Type Voltage-Gated Calcium Channel CaV1.2 in resetting the circadian clock in the late night. *J Biol Rhythms* 2014, 29:288–298. [PubMed: 25238857]
 14. Paul JR, DeWoskin D, McMeekin LJ, Cowell RM, Forger DB, Gamble KL: Regulation of persistent sodium currents by glycogen synthase kinase 3 encodes daily rhythms of neuronal excitability. *Nat Commun* 2016, 7:13470. [PubMed: 27841351]
 15. Kudo T, Block GD, Colwell CS: The Circadian Clock Gene *Period1* Connects the Molecular Clock to Neural Activity in the Suprachiasmatic Nucleus. *ASN Neuro* 2015, 7.
 16. Whitt JP, Montgomery JR, Meredith AL: BK channel inactivation gates daytime excitability in the circadian clock. *Nat Commun* 2016, 7:10837 A comprehensive set of experiments demonstrating that the circadian clock modifies the activity of the large conductance calcium-activated potassium channel inactivation gate as a biophysical switch to regulate the diurnal variation in SCN neuronal activity. [PubMed: 26940770]
 17. Montgomery JR, Whitt JP, Wright BN, Lai MH, Meredith AL: Mis-expression of the BK K(+) channel disrupts suprachiasmatic nucleus circuit rhythmicity and alters clock-controlled behavior. *Am J Physiol Cell Physiol* 2013, 304:C299–311. [PubMed: 23174562]
 18. Kudo T, Loh DH, Kuljis D, Constance C, Colwell CS: Fast delayed rectifier potassium current: critical for input and output of the circadian system. *J Neurosci* 2011, 31:2746–2755. [PubMed: 21414897]
 19. Pennartz CM, De Jeu MT, Geurtsen AM, Sluiter AA, Hermes ML: Electrophysiological and morphological heterogeneity of neurons in slices of rat suprachiasmatic nucleus. *J Physiol* 1998, 506 (Pt 3):775–793. [PubMed: 9503337]
 20. Cheng HY, Alvarez-Saavedra M, Dziema H, Choi YS, Li A, Obrietan K: Segregation of expression of *mPeriod* gene homologs in neurons and glia: possible divergent roles of *mPeriod1* and *mPeriod2* in the brain. *Hum Mol Genet* 2009, 18:3110–3124. [PubMed: 19477955]
 21. Riddle M, Mezas E, Foley D, LeSauter J, Silver R: Differential localization of *PER1* and *PER2* in the brain master circadian clock. *Eur J Neurosci* 2017, 45:1357–1367. [PubMed: 27740710]
 22. O'Leary T, Williams AH, Caplan JS, Marder E: Correlations in ion channel expression emerge from homeostatic tuning rules. *Proc Natl Acad Sci U S A* 2013, 110:E2645–2654. [PubMed: 23798391]
 - 23*. Albers HE, Walton JC, Gamble KL, McNeill JKt, Hummer DL: The dynamics of GABA signaling: Revelations from the circadian pacemaker in the suprachiasmatic nucleus. *Front Neuroendocrinol* 2017, 44:35–82.
- A comprehensive review covering decades of work exploring the importance of GABA signaling to SCN function.

[PubMed: 27894927]

- 24*. Ono D, Honma KI, Yanagawa Y, Yamanaka A, Honma S: Role of GABA in the regulation of the central circadian clock of the suprachiasmatic nucleus. *J Physiol Sci* 2018
- This recent review focusses on the controversies to be resolved and technical challenges to be overcome to better understand the roles that GABA signaling plays in the SCN.
25. Evans JA, Gorman MR: In synch but not in step: Circadian clock circuits regulating plasticity in daily rhythms. *Neuroscience* 2016, 320:259–280 [PubMed: 26861419]
26. Moldavan MG, Cravetchi O, Allen CN: GABA transporters regulate tonic and synaptic GABA_A receptor-mediated currents in the Suprachiasmatic Nucleus Neurons. *J Neurophysiol* 2017:jn00194 02017.
27. DeWoskin D, Myung J, Belle MD, Piggins HD, Takumi T, Forger DB: Distinct roles for GABA across multiple timescales in mammalian circadian timekeeping. *Proc Natl Acad Sci U S A* 2015, 112:E3911–3919. [PubMed: 26130805]
28. Myung J, Hong S, DeWoskin D, De Schutter E, Forger DB, Takumi T: GABA-mediated repulsive coupling between circadian clock neurons in the SCN encodes seasonal time. *Proc Natl Acad Sci U S A* 2015, 112:E3920–3929. [PubMed: 26130804]
29. Wagner S, Sagiv N, Yarom Y: GABA-induced current and circadian regulation of chloride in neurones of the rat suprachiasmatic nucleus. *J Physiol* 2001, 537:853–869. [PubMed: 11744760]
30. Irwin RP, Allen CN: GABAergic signaling induces divergent neuronal Ca²⁺ responses in the suprachiasmatic nucleus network. *Eur J Neurosci* 2009, 30:1462–1475. [PubMed: 19821838]
31. Wagner S, Castel M, Gainer H, Yarom Y: GABA in the mammalian suprachiasmatic nucleus and its role in diurnal rhythmicity. *Nature* 1997, 387:598–603. [PubMed: 9177347]
32. De Jeu M, Pennartz CMA: Circadian modulation of GABA function in the rat suprachiasmatic nucleus: excitatory effects during the night phase. *J Neurophysiol* 2002, 87:834–844. [PubMed: 11826050]
33. Choi HJ, Lee CJ, Schroeder A, Kim YS, Jung SH, Kim JS, Kim do Y, Son EJ, Han HC, Hong SK, et al.: Excitatory Actions of GABA in the Suprachiasmatic Nucleus. *J Neurosci* 2008, 28:5450–5459. [PubMed: 18495878]
34. Klett NJ, Allen CN: Intracellular Chloride Regulation in AVP+ and VIP+ Neurons of the Suprachiasmatic Nucleus. *Sci Rep* 2017, 7:10226. [PubMed: 28860458]
35. Farajnia S, van Westering TL, Meijer JH, Michel S: Seasonal induction of GABAergic excitation in the central mammalian clock. *Proc Natl Acad Sci U S A* 2014, 111:9627–9632. [PubMed: 24979761]
36. Prolo LM, Takahashi JS, Herzog ED: Circadian rhythm generation and entrainment in astrocytes. *J Neurosci* 2005, 25:404–408. [PubMed: 15647483]
- 37*. Brancaccio M, Patton AP, Chesham JE, Maywood ES, Hastings MH: Astrocytes Control Circadian Timekeeping in the Suprachiasmatic Nucleus via Glutamatergic Signaling. *Neuron* 2017, 93:1420–1435 e1425.
- SCN astrocytes express a circadian clock that drives an intracellular calcium rhythm that is antiphase to the calcium rhythm observed in SCN neurons. The astrocytes release glutamate, which is proposed to act on presynaptic NR2-expressing NMDA receptors to modulate GABA neurotransmission and regulate circadian timing.
- [PubMed: 28285822]
- 38*. Tso CF, Simon T, Greenlaw AC, Puri T, Mieda M, Herzog ED: Astrocytes Regulate Daily Rhythms in the Suprachiasmatic Nucleus and Behavior. *Curr Biol* 2017
- This study demonstrates that SCN astrocyte activity contributes to setting the SCN and behavioral circadian period. *Bmal1* deletion, or lengthening the period of the astrocyte circadian clock, lengthens the behavioral circadian period.
39. Moldavan M, Cravetchi O, Williams M, Irwin RP, Aicher SA, Allen CN: Localization and expression of GABA transporters in the suprachiasmatic nucleus. *Eur J Neurosci* 2015, 42:3018–3032. [PubMed: 26390912]

40. Guldner FH: Numbers of neurons and astroglial cells in the suprachiasmatic nucleus of male and female rats. *Exp Brain Res* 1983, 50:373–376. [PubMed: 6641871]
41. Jackson FR: Glial cell modulation of circadian rhythms. *Glia* 2011, 59:1341–1350. [PubMed: 21732426]
42. Ng FS, Tangredi MM, Jackson FR: Glial cells physiologically modulate clock neurons and circadian behavior in a calcium-dependent manner. *Curr Biol* 2011, 21:625–634. [PubMed: 21497088]
- 43**. Barca-Mayo O, Pons-Espinal M, Follert P, Armirotti A, Berdondini L, De PietriTonelli D: Astrocyte deletion of Bmal1 alters daily locomotor activity and cognitive functions via GABA signalling. *Nat Commun* 2017, 8:14336

This study provides a mechanism by which SCN astrocytes contribute to the generation of circadian timing signals. Disruption of the circadian clock in astrocytes reduces GABA uptake and alters behavioral circadian rhythms, possibly by changing GABAergic neurotransmission.

[PubMed: 28186121]

44. Halassa MM, Haydon PG: Integrated brain circuits: astrocytic networks modulate neuronal activity and behavior. *Annu Rev Physiol* 2010, 72:335–355.
45. Womac AD, Burkeen JF, Neuendorff N, Earnest DJ, Zoran MJ: Circadian rhythms of extracellular ATP accumulation in suprachiasmatic nucleus cells and cultured astrocytes. *Eur J Neurosci* 2009, 30:869–876. [PubMed: 19712092]
46. Marpegan L, Swannstrom AE, Chung K, Simon T, Haydon PG, Khan SK, Liu AC, Herzog ED, Beaulieu C: Circadian regulation of ATP release in astrocytes. *J Neurosci* 2011, 31:8342–8350. [PubMed: 21653839]
47. Burkeen JF, Womac AD, Earnest DJ, Zoran MJ: Mitochondrial calcium signaling mediates rhythmic extracellular ATP accumulation in suprachiasmatic nucleus astrocytes. *J Neurosci* 2011, 31:8432–8440 [PubMed: 21653847]
48. Bhattacharya A, Vavra V, Svobodova I, Bendova Z, Vereb G, Zemkova H: Potentiation of inhibitory synaptic transmission by extracellular ATP in rat suprachiasmatic nuclei. *J Neurosci* 2013, 33:8035–8044. [PubMed: 23637193]
49. Chen G, van den Pol AN: Adenosine modulation of calcium currents and presynaptic inhibition of GABA release in suprachiasmatic and arcuate nucleus neurons. *J Neurophysiol* 1997, 77:3035–3047. [PubMed: 9212255]
50. Hallworth R, Cato M, Colbert C, Rea MA: Presynaptic adenosine A1 receptors regulate retinohypothalamic neurotransmission in the hamster suprachiasmatic nucleus. *J Neurobiol* 2002, 52:230–240. [PubMed: 12210106]
51. Sigworth LA, Rea MA: Adenosine A1 receptors regulate the response of the mouse circadian clock to light. *Brain Res* 2003, 960:246–251. [PubMed: 12505678]
52. Ikeda M, Sugiyama T, Wallace CS, Gompf HS, Yoshioka T, Miyawaki A, Allen CN: Circadian dynamics of cytosolic and nuclear Ca²⁺ in single suprachiasmatic nucleus neurons. *Neuron* 2003, 38:253–263. [PubMed: 12718859]
53. Mercado C, Diaz-Munoz M, Alamilla J, Valderrama K, Morales-Tlalpan V, Aguilar-Roblero R: Ryanodine-sensitive intracellular Ca²⁺ channels in rat suprachiasmatic nuclei are required for circadian clock control of behavior. *J Biol Rhythms* 2009, 24:203–210. [PubMed: 19465697]
54. Lundkvist GB, Kwak Y, Davis EK, Tei H, Block GD: A calcium flux is required for circadian rhythm generation in mammalian pacemaker neurons. *J. Neurosci* 2005, 25:7682–7686. [PubMed: 16107654]
- 55**. Enoki R, Ono D, Kuroda S, Honma S, Honma KI: Dual origins of the intracellular circadian calcium rhythm in the suprachiasmatic nucleus. *Sci Rep* 2017, 7:41733

This study uses ex vivo methods to simultaneously monitor intracellular calcium levels, clock gene activity and spontaneous spiking in SCN slices. The overall conclusions support previous work, demonstrating that in SCN neurons, intracellular calcium rhythms are dependent on both intracellular release and extracellular entry of calcium.

[PubMed: 28155916]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Highlights

- In this review we briefly highlight significant research that is increasing our understanding of the critical interactions between the circadian molecular clock and neuronal and glial activity in the generation and communication of circadian timing signals.

Figure 1.

A representation of how the circadian signal is generated and processed by suprachiasmatic nucleus (SCN) neurons and glia. Here, signals travelling to and from the transcription–translation feedback loop (TTFL) in SCN cells must first flow through the membrane by poorly understood mechanisms. This communication pathway is central to how SCN cells maintain phase synchrony. Through similar unknown processes, environmental light and homeostatic cues from physiology and behaviour feedback to appropriately adjust TTFL timing in SCN cells. This ensures that circadian rhythms generated are aligned with the animal’s internal physiology and the prevailing light–dark cycle. Solid and broken axon and glial lines show established and possible contacts, respectively.