Published in final edited form as:

Curr Opin Virol. 2019 February ; 34: 130–139. doi:10.1016/j.coviro.2019.02.005.

Mutually assured destruction: the cold war between viruses and natural killer cells

Ayad Ali^{1,2,3}, Ivayla E. Gyurova^{1,4}, and Stephen N. Waggoner^{1,2,3,4,5,*}

¹Center for Autoimmune Genomics and Etiology, Cincinnati Children's Hospital Medical Center

²Medical Scientist Training Program, University of Cincinnati College of Medicine

³Immunology Graduate Training Program, University of Cincinnati College of Medicine

⁴Pathobiology and Molecular Medicine Graduate Program, University of Cincinnati College of Medicine

⁵Department of Pediatrics, University of Cincinnati College of Medicine

Abstract

Natural killer (NK) cells play a multitude of antiviral roles that are significant enough to provoke viral counter efforts to subvert their activity. As innate lymphocytes, NK cells provide a rapid source of pro-inflammatory antiviral cytokines and bring to bear cytolytic activities that are collectively meant to constrain viral replication and dissemination. Additionally, NK cells participate in adaptive immunity both by shaping virus-specific T-cell responses and by developing adaptive features themselves, including enhanced antibody-dependent effector functions. The relative importance of different functional activities of NK cells are poorly understood, thereby obfuscating clinical use of these cells. Here we focus on opposing efforts of NK cells and viruses to gain tactical superiority during infection.

Graphical abstract

Introduction

There is strong evidence that natural killer (NK) cells make crucial contributions to host defense during virus infection. In humans, genetic lesions that result in functional or cellular deficiencies of NK cells are strongly linked to heightened susceptibility to herpesvirus and

*Address correspondence to: Stephen.Waggoner@cchmc.org.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

papillomavirus infections [1–3]. Absence or dysfunction of NK cells likewise contributes to loss of viral control and altered pathogenesis of virus infection in mice [4], most notably in the context of murine cytomegalovirus (MCMV) infection. NK cells also play a role in pathogenesis of simian immunodeficiency virus (SIV) infection in non-human primates [••5]. The antiviral activities of NK cells span production of pro-inflammatory cytokines, like interferon gamma (IFN- γ) [6], and lysis of infected cells [7]. NK cells also shape adaptive antiviral responses by editing the available pool of antigen-presenting cells [8] and directly inhibiting T-cell responses [9]. Remarkably, NK cells can also develop memory-like features of antiviral T cells [10,11] and are capable of cooperating with B cells to suppress virus replication via antibody-dependent cellular cytotoxicity (ADCC) [12]. While the relative importance of these diverse functions of NK cells in virus infection remains undefined, the evolutionary trade-offs exhibited by viruses to facilitate suppression or evasion of NK cell effector function [13,14] highlights the strong antiviral potential of these cells. In this review, we highlight contributions made by NK cells to pathogenesis of virus infection and describe mechanisms used by viruses to fight back.

Innate functions of NK cells during virus infection

Proinflammatory cytokine release

NK cells provide a crucial, early source of IFN- γ that is necessary for host defense against multiple viruses, including MCMV [6], vaccinia virus [15], herpes simplex virus-2 (HSV-2) [•16], ectromelia virus (ECTV) [•17], and influenza virus [••18]. NK-cell derived IFN- γ promotes non-cytolytic control of virus replication [19] and enhances antiviral T-cell responses [20] (Figure 1). Notably, activated NK cells make additional cytokines (e.g. tumor necrosis factor alpha, TNF- α), growth factors (e.g. Granulocyte-macrophage colony-stimulating factor, GM-CSF), and inflammatory chemokines (e.g. RANTES) [21,••22], with the potential to influence antiviral immunity.

The closely related but distinct lineage of type 1 innate lymphoid cells (ILC1) [23] are an additional source of early IFN- γ after virus infection. In fact, tissue resident ILC1 in the liver and at other sites of initial virus replication confer IFN- γ -dependent control of MCMV infection [24]. The context-dependent requirements and potential overlap of NK or ILC1 antiviral IFN- γ production, particularly where both cells are implicated in viral control at initial sites of virus entry [6,••18,24,25], remains to be determined.

Antiviral IFN- γ production by NK cells is driven by the pro-inflammatory cytokine milieu elicited by virus infection. This frequently includes interleukin-12 (IL-12) and IL-18 induction of IFN- γ [26,27], although type I IFN and coordination among innate cells is important as well [28] (Figure 1). Mechanistically, type I IFN critically primes IL-18 production by inflammatory monocytes, which in turn elicits antiviral IFN- γ production by NK cells during HSV-2 infection [•16]. In addition, NK cells can promote coordination among innate leukocytes to enhance their own antiviral function. Infected dendritic cells trigger initial release of IFN- γ by NK cells in draining lymph node after ECTV infection, thereby simulating chemokine (C-X-C motif) ligand 9 (CXCL9) expression by inflammatory monocytes and subsequent recruitment of CXCR3-expressing NK cells to constrain viral dissemination [•17]. Recent work in cancer immunology reveals the capacity of NK cells to

stimulate inflammatory dendritic cell (DC) recruitment into tumors [29,30], highlighting the intriguing possibility that similar mechanisms exist at sites of virus infection.

Direct lysis of virus-infected cells

While inflammatory cytokines can enhance IFN- γ -producing and cytolytic functions of NK cells, variegated expression of an array of germline encoded receptors that recognize viral, self, or altered self ligands ultimately controls killing of virus-infected cells by NK cells (reviewed in [7]) (Figure 2). Natural Killer Group 2D (NKG2D) provides an activating signal upon recognition of stress-induced ligands on infected cells [•17,31–33]. NKP46 stimulates NK-cell activation in response to viral ligands encoded by parainfluenza [34], metapneumovirus [35], reovirus [36], and influenza A [37,38] viruses. In C57BL/6 mice, the Ly49H receptor directly recognizes the m157 gene product of MCMV [39–41], whereas human NK cell responses against CMV are driven largely by NKG2C [••42] and activating killer cell immunoglobulin-like receptors (KIRs) [43,44]. Notably, the recent discovery that NKG2C recognized CMV UL40-derived peptides [••42] underscores the phenomenon of receptor driven expansion of specific subsets of NK cells expressing virus-specific recognition receptors in mice [45–47], humans [48], and non-human primates [49].

Variable engagement of NK cell receptors by viral peptide-containing class I human leukocyte antigen (HLA) and major histocompatibility complex (MHC) molecules is another mechanism controlling NK cell effector function in responses to virus infection (Figure 2). This concept is reinforced by the strong correlations between various KIR/HLA combinations and outcomes of virus infection, including those with hepatitis C virus (HCV) and HIV-1 [50,51]. Preservation of surface expression of HLA-C can limit viral elimination by NK cells via engagement of inhibitory KIR [52], yet KIR2DL2/S2 is implicated in protection against HIV in the context of specific HLA-C alleles. Expression of KIR2DL2/S2 in combination with HLA-C*14:03 or HLA-C*12:02 correlated with reduced viral load and increased NK-cell mediated elimination of infected CD4 T cells [•53]. Mechanistically, low HIV peptide affinity with these HLA subtypes resulted in reduced inhibitory signaling via KIR2DL2. In a similar vein, downmodulation of HLA-C does not always equate to enhanced NK cell antiviral function. Korner and colleagues witnessed impaired antiviral function of KIR2DL+ NK cells against HIV-1-infected T cells that had downregulated HLA-C, revealing that NK cells are sensitive to virus-induced changes in HLA expression, but that the response to reduced HLA depends on genetics of HLA-KIR interaction [•54].

Activating KIR are also associated with pathogenesis of virus infection [55] (Figure 2). KIR2DS2 stimulates NK cell function upon recognition of a well-conserved flavivirus helicase peptide bound to HLA-C*0102, thereby implicating KIR2DS2 in immunity to HCV [•56]. In a similar fashion, KIR3DS1 is associated with delayed progression to AIDS in HIV-1 infection [50]. Recently, open-conformers of HLA-F were revealed as high-affinity ligands for KIR3DS1 that trigger NK cell cytokine production and degranulation [••57]. Moreover, activated and HIV-infected CD4 T cells exhibit increased expression of HLA-F. However, peptide-bound HLA-F preferentially binds leukocyte immunoglobulin-like receptor 1 (LIR-1), potentially providing an inhibitory signal to LIR-1 expressing NK cells [•58]

The inhibitory receptor NKG2A that interacts with the non-classical MHC receptor HLA-E (Figure 2), has emerged as a potent regulator of antiviral NK cell function [59,60]. Barker and colleagues [61] discovered that a conserved HIV-1 peptide bound to HLA-E renders infected T cells highly susceptible to NK cell killing by limiting NKG2A-derived inhibitory signals. In addition, elegant genetic work reveals that expression of the NKG2A ligand HLA-E is driven by a combination of genetic alleles that dictate HLA-A and HLA-B expression and ultimately determine HIV-1 control [62]. High expression levels of HLA-A promotes HLA-E expression by providing a source of HLA-E stabilizing peptides, thereby provoking a strong NKG2A inhibitory signal and weak NK cell antiviral responses. Moreover, HLA-B haplotypes that characterize strong education of NK cells via NKG2A [63] exacerbate the effect of HLA-A expression on HLA-E/NKG2A inhibitory signaling [62].

NK cell migration to sites of virus replication

Migration of NK cells to sites of virus replication and specific regions of lymphoid tissues is implicated in both viral control and determination of disease pathology [64,65]. Recent studies revealed that NK cell migration into the B-cell zone of lymph nodes during SIV infection is associated with better viral control and non-pathogenic infection in the natural SIV host species, the African Green Monkeys [5]. Depletion of NK cells in these animals significantly increased burden of SIV in lymph nodes, highlighting the potential antiviral role of NK cells in natural SIV hosts. In contrast, SIV infection of macaques is associated with reduced NK cell migration to B-cell follicles, poor viral control, and development of AIDS [5]. Of note, NK cell localization within the follicle is a shared feature of MCMV and lymphocytic choriomeningitis virus (LCMV) infections of mice [66,67], reflecting the general importance of NK migration to sites of infection to mediate viral elimination or control.

NK cell participation in antiviral adaptive immunity

NK-cell regulation of adaptive immunity

Perforin- and IL-10-dependent immunoregulatory mechanisms of NK cells are implicated in restraint of antiviral responses by T and B cells [9,68,69] (Figure 1). This immunoregulatory activity of NK cells crucially determines immune exhaustion, viral persistence, and immunopathology during chronic infection with LCMV [70–72]. Moreover, NK cell suppression affects development of T and B cell memory responses after acute virus infection [73]. However, neither the generation of memory T cells after adenovirus immunization nor the recall of these memory T cells after LCMV challenge was measurably impacted by NK cells [74]. Collective results from several labs indicate that induction of NK cell regulatory function is a shared feature of many virus infections [72,75–79], but that differential gene expression, including that of NKG2A, may tune the regulatory capacity of NK cells and the susceptibility of T cells to this effect [74,80,81]. In fact, NKG2A has emerged as a crucial negative regulator of the immunosuppressive capacity of NK cells [80–82], which highlights an alternative explanation for links between NKG2A and virus infection outcome [62]. Distinct subsets of NK cells may differ in their regulatory capacity, in part due to how they are educated during development [22]. A critical new finding

concerning the immunoregulatory role of NK cells in virus infection recently emerged from investigation of HIV-1-infected individuals who do or do not develop broadly neutralizing antibodies (bnAbs) against HIV-1. Induction of bnAbs was associated with high expression of *RAB11FIP5* and dysfunction of NK cells [••83]. These data extend past mouse studies into humans and reveal the potential utility of targeting NK cells to enhance vaccine-elicited antibody responses [73,•84].

Adaptive features of NK cells

The recognition that subsets of NK cells can clonally expand and form pools of memory-like cells in response to virus infection or immunization has revolutionized NK cell research (reviewed in [85]) (Figure 1). NK cells with adaptive features are observed in MCMV [46], HCMV [••42,86,87], vaccinia virus [88], and SIV [49,89] infection, as well as after cytokine, hapten, or vaccine administration [89–92]. Although a wealth of studies reveal precise epigenetic [•93,94], metabolic [95], and transcription factors [96–98] involved in generation of antiviral NK cells with adaptive features, the mechanisms governing antigen-specific induction of these memory NK cells remain unclear. Recent evidence points to HCMV UL40-derived peptides as drivers of NKG2C+ NK cell expansion [••42]. In the case of influenza A virus, NK cell expression of IFN- γ varies in a viral strain-dependent manner linked to differential type I IFN, CD54, and CD112 expression [••18]. Likewise, peptides derived from HCV and HIV alter KIR interaction with HLA, resulting in enhanced activation of NK cell subsets expressing particular KIR [56,99–102]. The importance of viral peptide-HLA interaction with KIR is reinforced by observations of HIV adaptation to KIR [103,104]. Alteration of HLA interactions with NK cell receptors by viral peptides may also explain the observation of NK cell specificity for SIV-derived antigens in SIV- or SHIV-infected, as well as Ad26 based SIV vaccine immunized resus macaques [89]. Notably, another type of adaptive NK cell demarcated by reduced expression of Fc ϵ R γ , SYK, and/or EAT-2 expands in CMV infected individuals, but appears to do so in response to antibody-dependent signals [86,87,105]. Remarkably, this NK cell subset displays enhanced antibody-dependent effector functions that reflect cooperation between innate (NK) and adaptive (B-cell) memory cells (Figure 1).

Antibody-dependent cellular cytotoxicity (ADCC)

ADCC involves CD16-dependent recognition and killing of antibody-opsonized targets by NK cells [12] (Figure 1). This mechanism couples NK cells to antibody production by B cells, and represents a possible correlate of protection against infections, including those with HIV-1 [106–108]. Notably, the education of NK cells by self-specific HLA-binding receptors [109,110] contributes to capacity of mature NK cells to mediate ADCC [63,•111]. Some pathogens, including HSV-1, express viral antigens (gE) with the capacity to bind Fc domains of immunoglobulin molecules (IgG-Fc) to block ADCC. However, a recent study revealed that HSV-1 gE binding to IgG-Fc forms a molecular bridge between NK cells and infected cells that promotes NK cell-mediated lysis and clearance of HSV-1-infected cells [••112]. Bacteria-derived proteins are capable of similar cross-linking, potentially highlighting a common mechanism of immune defense.

Viral subversion of NK cell functions

Virus infection of NK cells

Some viruses may evade innate immunity by directly infecting NK cells. HSV [113,114], Epstein-Barr virus [115], and human herpesvirus 6 [116] were all previously shown to productively infect NK cells, while influenza and vaccinia viruses can establish non-productive infections in these cells [117,118]. HIV also infects CD4+ NK cells [119], although more recent work suggests this may reflect infection of CD4+ ILC1, which are subsequently depleted from the gut [120]. More recently, another herpesvirus, varicella zoster virus (VZV), was shown to preferentially infect CD56dim NK cells, thereby driving CD57 expression, loss of cell-surface CD16, and impairment of NK-cell function [121].

Viral evasion of NK cells

Many viruses work to actively suppress or avoid NK cell activation (reviewed in [122]). During HIV-1 infection, viral gene products actively subvert NK-cell responses [123,124] while the virus itself appears to evolve in response to NK cell mediated pressure [103,104]. Most, if not all, members of the herpesvirus family suppress NK cells through a myriad of methods (reviewed in [14]). These mechanisms often reflect complicated evolutionary relationships between virus and host. For example, MCMV m157 binds to Ly49H [39] promoting viral resistance in a singular strain of inbred mice (C57BL/6), yet many MCMV isolates have mutations in m157 that evade Ly49H recognition [125] or promote interaction with inhibitory Ly49C receptors [126]. Another MCMV protein, m12, acts as a decoy immunoevasin by engaging with both inhibitory NKR-P1B and activating NKR-P1A/C receptors to modulate NK cell cytotoxic function [127]. Remarkably, gain- and loss-of-function mutations in m12 reveal evolution of host-virus interaction.

NK cell exhaustion

Persistent virus infections are known to promote dysfunction of virus-specific T cells [128] (Figure 1). This phenomenon led to the discovery that blockade of programmed death-1 (PD-1) interactions with its ligand, PD-L1, can rescue exhausted T cells in the context of chronic virus infection [129] and cancer, with the latter finding revolutionizing the cancer clinic [130]. NK cells also undergo functional exhaustion during chronic inflammation [131] and may contribute to clinical responses following PD-1 blockade [132,133]. However, TIGIT [134], IL1R8 [135], and TIM3 [136] also appear to be highly relevant checkpoint receptors in NK cell functional exhaustion. Furthermore, a wealth of new studies indicate the metabolism stringently regulates the function of NK cells [137–143], which can limit their antiviral capacity in the context of CMV or HCV infection [144,145].

Conclusions

NK cells play a variety of roles in response to virus infection. These activities include both independent and cooperative, multi-lineage functions that span innate and adaptive immunity. The clinical importance of NK cell function is clearly seen both from disease in patients who lack NK cells as well as from the evolutionary sacrifices made by many viruses to manipulate NK-cell functional activity. An improved understanding of context-specific

contributions of NK cells in virus infection will be necessary to develop means to therapeutically modulate NK cells during infection.

Acknowledgements

The authors are supported by National Institutes of Health grants DA038017 and TR001425 (subaward), the Cincinnati Children's Research Foundation, The Lawrence Ellison Foundation, and the Ralph and Marion Falk Medical Research Trust.

References

1. Voss M, Bryceson YT: Natural killer cell biology illuminated by primary immunodeficiency syndromes in humans. *Clin Immunol* 2017, 177:29–42. [PubMed: 26592356]
2. Mace EM, Orange JS: Genetic Causes of Human NK Cell Deficiency and Their Effect on NK Cell Subsets. *Front Immunol* 2016, 7:545. [PubMed: 27994588]
3. Mace EM, Bigley V, Gunesch JT, Chinn IK, Angelo LS, Care MA, Maisuria S, Keller MD, Togi S, Watkin LB, et al.: Biallelic mutations in IRF8 impair human NK cell maturation and function. *J Clin Invest* 2017, 127:306–320. [PubMed: 27893462]
4. Waggoner SN, Reighard SD, Gyurova IE, Cranert SA, Mahl SE, Karnele EP, McNally JP, Moran MT, Brooks TR, Yaqoob F, et al.: Roles of natural killer cells in antiviral immunity. *Curr Opin Virol* 2016, 16:15–23. [PubMed: 26590692]
5. Huot N, Jacquelin B, Garcia-Tellez T, Rasclé P, Ploquin MJ, Madec Y, Reeves RK, Derreudre-Bosquet N, Muller-Trutwin M: Natural killer cells migrate into and control simian immunodeficiency virus replication in lymph node follicles in African green monkeys. *Nat Med* 2017, 23:1277–1286. [PubMed: 29035370] Demonstrates that migration of NK cells into B-cell follicles is associated with viral control and non-pathogenic course of SIV infection in natural host species.
6. Orange JS, Wang B, Terhorst C, Biron CA: Requirement for natural killer cell-produced interferon gamma in defense against murine cytomegalovirus infection and enhancement of this defense pathway by interleukin 12 administration. *J Exp Med* 1995, 182:1045–1056. [PubMed: 7561678]
7. Hammer Q, Ruckert T, Romagnani C: Natural killer cell specificity for viral infections. *Nat Immunol* 2018, 19:800–808. [PubMed: 30026479]
8. Moretta A: Natural killer cells and dendritic cells: rendezvous in abused tissues. *Nat Rev Immunol* 2002, 2:957–964. [PubMed: 12461568]
9. Welsh RM, Waggoner SN: NK cells controlling virus-specific T cells: Rheostats for acute vs. persistent infections. *Virology* 2013, 435:37–45. [PubMed: 23217614]
10. Lau CM, Sun JC: The widening spectrum of immunological memory. *Curr Opin Immunol* 2018, 54:42–49. [PubMed: 29902670]
11. Paust S, Blish CA, Reeves RK: Redefining Memory: Building the Case for Adaptive NK Cells. *J Virol* 2017, 91.
12. Ochoa MC, Minute L, Rodriguez I, Garasa S, Perez-Ruiz E, Inoges S, Melero I, Berraondo P: Antibody-dependent cell cytotoxicity: immunotherapy strategies enhancing effector NK cells. *Immunol Cell Biol* 2017, 95:347–355. [PubMed: 28138156]
13. Lanier LL: Evolutionary struggles between NK cells and viruses. *Nat Rev Immunol* 2008, 8:259–268. [PubMed: 18340344]
14. De Pelsmaeker S, Romero N, Vitale M, Favoreel HW: Herpesvirus Evasion of Natural Killer Cells. *J Virol* 2018, 92.
15. Abboud G, Tahiliani V, Desai P, Varkoly K, Driver J, Hutchinson TE, Salek-Ardakani S: Natural Killer Cells and Innate Interferon Gamma Participate in the Host Defense against Respiratory Vaccinia Virus Infection. *J Virol* 2016, 90:129–141. [PubMed: 26468539]
16. Lee AJ, Chen B, Chew MV, Barra NG, Shenouda MM, Nham T, van Rooijen N, Jordana M, Mossman KL, Schreiber RD, et al.: Inflammatory monocytes require type I interferon receptor signaling to activate NK cells via IL-18 during a mucosal viral infection. *J Exp Med* 2017,

- 214:1153–1167. [PubMed: 28264883] In a mouse model of HSV-2 infection, the authors show that type I IFN signaling in inflammatory monocytes is critical for NK cell function.
- 17•. Wong E, Xu RH, Rubio D, Lev A, Stotesbury C, Fang M, Sigal LJ: Migratory Dendritic Cells, Group 1 Innate Lymphoid Cells, and Inflammatory Monocytes Collaborate to Recruit NK Cells to the Virus-Infected Lymph Node. *Cell Rep* 2018, 24:142–154. [PubMed: 29972776] The authors unveil a TLR9/MyD88-dependent mechanism of recruitment of NK cells to draining lymph nodes in order to establish NK cell viral control.
 - 18••. Kronstad LM, Seiler C, Vergara R, Holmes SP, Blish CA: Differential Induction of IFN-alpha and Modulation of CD112 and CD54 Expression Govern the Magnitude of NK Cell IFN-gamma Response to Influenza A Viruses. *J Immunol* 2018, 201:2117–2131. [PubMed: 30143589] This study unveils strain-specific functional responses of NK cells to influenza A virus that depend on differential induction of type I IFN and receptors for NK cell ligands.
 19. Guidotti LG, Chisari FV: Noncytolytic control of viral infections by the innate and adaptive immune response. *Annu Rev Immunol* 2001, 19:65–91. [PubMed: 11244031]
 20. Martin-Fontecha A, Thomsen LL, Brett S, Gerard C, Lipp M, Lanzavecchia A, Sallusto F: Induced recruitment of NK cells to lymph nodes provides IFN-gamma for T(H)1 priming. *Nat Immunol* 2004, 5:1260–1265. [PubMed: 15531883]
 21. Wagstaffe HR, Nielsen CM, Riley EM, Goodier MR: IL-15 Promotes Polyfunctional NK Cell Responses to Influenza by Boosting IL-12 Production. *J Immunol* 2018, 200:2738–2747. [PubMed: 29491009]
 - 22••. Zamora AE, Aguilar EG, Sungur CM, Khuat LT, Dunai C, Lochhead GR, Du J, Pomeroy C, Blazar BR, Longo DL, et al.: Licensing delineates helper and effector NK cell subsets during viral infection. *JCI Insight* 2017, 2. The authors reveal that distinct subsets of NK cells that are licensed via different inhibitory receptors have disparate functional capacities during virus infection.
 23. Vivier E, Artis D, Colonna M, Diefenbach A, Di Santo JP, Eberl G, Koyasu S, Locksley RM, McKenzie ANJ, Mebius RE, et al.: Innate Lymphoid Cells: 10 Years On. *Cell* 2018, 174:1054–1066. [PubMed: 30142344]
 24. Weizman OE, Adams NM, Schuster IS, Krishna C, Pritykin Y, Lau C, Degli-Esposti MA, Leslie CS, Sun JC, O'Sullivan TE: ILC1 Confer Early Host Protection at Initial Sites of Viral Infection. *Cell* 2017, 171:795–808 e712. [PubMed: 29056343]
 25. Vashist N, Trittel S, Ebensen T, Chambers BJ, Guzman CA, Riese P: Influenza-Activated ILC1s Contribute to Antiviral Immunity Partially Influenced by Differential GTR Expression. *Front Immunol* 2018, 9:505. [PubMed: 29623077]
 26. Chaix J, Tessmer MS, Hoebe K, Fuseri N, Ryffel B, Dalod M, Alexopoulou L, Beutler B, Brossay L, Vivier E, et al.: Cutting edge: Priming of NK cells by IL-18. *J Immunol* 2008, 181:1627–1631. [PubMed: 18641298]
 27. Chan SH, Perussia B, Gupta JW, Kobayashi M, Pospisil M, Young HA, Wolf SF, Young D, Clark SC, Trinchieri G: Induction of interferon gamma production by natural killer cell stimulatory factor: characterization of the responder cells and synergy with other inducers. *J Exp Med* 1991, 173:869–879. [PubMed: 1672545]
 28. Hunter CA, Gabriel KE, Radzanowski T, Neyer LE, Remington JS: Type I interferons enhance production of IFN-gamma by NK cells. *Immunol Lett* 1997, 59:1–5. [PubMed: 9334850]
 29. Barry KC, Hsu J, Broz ML, Cueto FJ, Binnewies M, Combes AJ, Nelson AE, Loo K, Kumar R, Rosenblum MD, et al.: A natural killer-dendritic cell axis defines checkpoint therapy-responsive tumor microenvironments. *Nat Med* 2018, 24:1178–1191. [PubMed: 29942093]
 30. Bottcher JP, Bonavita E, Chakravarty P, Blees H, Cabeza-Cabrero M, Sammiceli S, Rogers NC, Sahai E, Zelenay S, Reis e Sousa C: NK Cells Stimulate Recruitment of cDC1 into the Tumor Microenvironment Promoting Cancer Immune Control. *Cell* 2018, 172:1022–1037 e1014. [PubMed: 29429633]
 31. Fang M, Lanier LL, Sigal LJ: A role for NKG2D in NK cell-mediated resistance to poxvirus disease. *PLoS Pathog* 2008, 4:e30. [PubMed: 18266471]
 32. Draghi M, Pashine A, Sanjanwala B, Gendzekhadze K, Cantoni C, Cosman D, Moretta A, Valiante NM, Parham P: NKG2D recognition of infected dendritic cells is necessary for NK

- cell activation in the human response to influenza infection. *J Immunol* 2007, 178:2688–2698. [PubMed: 17312110]
33. Cosman D, Mullberg J, Sutherland CL, Chin W, Armitage R, Fanslow W, Kubin M, Chalupny NJ: ULBPs, novel MHC class I-related molecules, bind to CMV glycoprotein UL16 and stimulate NK cytotoxicity through the NKG2D receptor. *Immunity* 2001, 14:123–133. [PubMed: 11239445]
 34. McQuaid S, Loughran S, Power P, Maguire P, Walls D, Cusi MG, Orvell C, Johnson P: Haemagglutinin-neuraminidase from HPIV3 mediates human NK regulation of T cell proliferation via NKp44 and NKp46. *J Gen Virol* 2018, 99:763–767. [PubMed: 29683419]
 35. Diab M, Glasner A, Isaacson B, Bar-On Y, Drori Y, Yamin R, Duev-Cohen A, Danziger O, Zamostiano R, Mandelboim M, et al.: NK-cell receptors NKp46 and NCR1 control human metapneumovirus infection. *Eur J Immunol* 2017, 47:692–703. [PubMed: 28191644]
 36. Bar-On Y, Charpak-Amikam Y, Glasner A, Isaacson B, Duev-Cohen A, Tsukerman P, Varvak A, Mandelboim M, Mandelboim O: NKp46 Recognizes the Sigma1 Protein of Reovirus: Implications for Reovirus-Based Cancer Therapy. *J Virol* 2017, 91.
 37. Mendelson M, Tekoah Y, Zilka A, Gershoni-Yahalom O, Gazit R, Achdout H, Bovin NV, Meninger T, Mandelboim M, Mandelboim O, et al.: NKp46 O-glycan sequences that are involved in the interaction with hemagglutinin type 1 of influenza virus. *J Virol* 2010, 84:3789–3797. [PubMed: 20147410]
 38. Mandelboim O, Lieberman N, Lev M, Paul L, Arnon TI, Bushkin Y, Davis DM, Strominger JL, Yewdell JW, Porgador A: Recognition of haemagglutinins on virus-infected cells by NKp46 activates lysis by human NK cells. *Nature* 2001, 409:1055–1060. [PubMed: 11234016]
 39. Arase H, Mocarski ES, Campbell AE, Hill AB, Lanier LL: Direct recognition of cytomegalovirus by activating and inhibitory NK cell receptors. *Science* 2002, 296:1323–1326. [PubMed: 11950999]
 40. Smith HR, Heusel JW, Mehta IK, Kim S, Dorner BG, Naidenko OV, Iizuka K, Furukawa H, Beckman DL, Pingel JT, et al.: Recognition of a virus-encoded ligand by a natural killer cell activation receptor. *Proc Natl Acad Sci U S A* 2002, 99:8826–8831. [PubMed: 12060703]
 41. Lee SH, Girard S, Macina D, Busa M, Zafer A, Belouchi A, Gros P, Vidal SM: Susceptibility to mouse cytomegalovirus is associated with deletion of an activating natural killer cell receptor of the C-type lectin superfamily. *Nat Genet* 2001, 28:42–45. [PubMed: 11326273]
 42. Hammer Q, Ruckert T, Borst EM, Dunst J, Haubner A, Durek P, Heinrich F, Gasparoni G, Babic M, Tomic A, et al.: Peptide-specific recognition of human cytomegalovirus strains controls adaptive natural killer cells. *Nat Immunol* 2018, 19:453–463. [PubMed: 29632329] The authors discover that viral peptides are the ligands recognized by NKG2C that promote expansion and differentiation of adaptive NK cells.
 43. van der Ploeg K, Chang C, Ivarsson MA, Moffett A, Wills MR, Trowsdale J: Modulation of Human Leukocyte Antigen-C by Human Cytomegalovirus Stimulates KIR2DS1 Recognition by Natural Killer Cells. *Front Immunol* 2017, 8:298. [PubMed: 28424684]
 44. Beziat V, Liu LL, Malmberg JA, Ivarsson MA, Sohlberg E, Bjorklund AT, Retiere C, Sverremark-Ekstrom E, Traherne J, Ljungman P, et al.: NK cell responses to cytomegalovirus infection lead to stable imprints in the human KIR repertoire and involve activating KIRs. *Blood* 2013, 121:2678–2688. [PubMed: 23325834]
 45. Daniels KA, Devora G, Lai WC, O'Donnell CL, Bennett M, Welsh RM: Murine cytomegalovirus is regulated by a discrete subset of natural killer cells reactive with monoclonal antibody to Ly49H. *J Exp Med* 2001, 194:29–44. [PubMed: 11435470]
 46. Sun JC, Beilke JN, Lanier LL: Adaptive immune features of natural killer cells. *Nature* 2009, 457:557–561. [PubMed: 19136945]
 47. Dokun AO, Kim S, Smith HR, Kang HS, Chu DT, Yokoyama WM: Specific and nonspecific NK cell activation during virus infection. *Nat Immunol* 2001, 2:951–956. [PubMed: 11550009]
 48. Guma M, Angulo A, Vilches C, Gomez-Lozano N, Malats N, Lopez-Botet M: Imprint of human cytomegalovirus infection on the NK cell receptor repertoire. *Blood* 2004, 104:3664–3671. [PubMed: 15304389]

49. Ram DR, Manickam C, Hueber B, Itell HL, Permar SR, Varner V, Reeves RK: Tracking KLRC2 (NKG2C)+ memory-like NK cells in SIV+ and rhCMV+ rhesus macaques. *PLoS Pathog* 2018, 14:e1007104. [PubMed: 29851983]
50. Martin MP, Gao X, Lee JH, Nelson GW, Detels R, Goedert JJ, Buchbinder S, Hoots K, Vlahov D, Trowsdale J, et al.: Epistatic interaction between KIR3DS1 and HLA-B delays the progression to AIDS. *Nat Genet* 2002, 31:429–434. [PubMed: 12134147]
51. Khakoo SI, Thio CL, Martin MP, Brooks CR, Gao X, Astemborski J, Cheng J, Goedert JJ, Vlahov D, Hilgartner M, et al.: HLA and NK cell inhibitory receptor genes in resolving hepatitis C virus infection. *Science* 2004, 305:872–874. [PubMed: 15297676]
52. Cohen GB, Gandhi RT, Davis DM, Mandelboim O, Chen BK, Strominger JL, Baltimore D: The selective downregulation of class I major histocompatibility complex proteins by HIV-1 protects HIV-infected cells from NK cells. *Immunity* 1999, 10:661–671. [PubMed: 10403641]
- 53•. Lin Z, Kuroki K, Kuse N, Sun X, Akahoshi T, Qi Y, Chikata T, Naruto T, Koyanagi M, Murakoshi H, et al.: HIV-1 Control by NK Cells via Reduced Interaction between KIR2DL2 and HLA-C(*):12:02/C(*):14:03. *Cell Rep* 2016, 17:2210–2220. [PubMed: 27880898] The authors demonstrate that KIR2DL2/S2 expression along with KIR2DL2/HLA-C*12:02 or KIR2DL2/HLA-C*14:03 have protective effects on control of HIV-1.
- 54•. Korner C, Simoneau CR, Schommers P, Granoff M, Ziegler M, Holzemer A, Lunemann, Chukwukelu J, Corleis B, Naranbhai V, et al.: HIV-1-Mediated Downmodulation of HLA-C Impacts Target Cell Recognition and Antiviral Activity of NK Cells. *Cell Host Microbe* 2017, 22:111–119 e114. [PubMed: 28704647] This study reveals that NK cells are ultrasensitive to virus-induced changes in HLA expression, and that the responses of NK cells depends on education genetics.
55. Jost S, Altfeld M: Control of human viral infections by natural killer cells. *Annu Rev Immunol* 2013, 31:163–194. [PubMed: 23298212]
- 56•. Naiyer MM, Cassidy SA, Magri A, Cowton V, Chen K, Mansour S, Kranidioti H, Mbirindi B, Rettman P, Harris S, et al.: KIR2DS2 recognizes conserved peptides derived from viral helicases in the context of HLA-C. *Sci Immunol* 2017, 2:The authors reveal a viral peptide that promotes KIR mediated activation of NK cells.
- 57•. Garcia-Beltran WF, Holzemer A, Martrus G, Chung AW, Pacheco Y, Simoneau CR, Rucevic M, Lamothe-Molina PA, Pertel T, Kim TE, et al.: Open conformers of HLA-F are high-affinity ligands of the activating NK-cell receptor KIR3DS1. *Nat Immunol* 2016, 17:1067–1074. [PubMed: 27455421] The authors discover that HLA-F is the high affinity ligand for KIR3DS1 on NK cells that triggers NK cell cytokine production and degranulation.
- 58•. Dulberger CL, McMurtrey CP, Holzemer, Neu KE, Liu V, Steinbach AM, Garcia-Beltran WF, Sulak M, Jabri B, Lynch VJ, et al.: Human Leukocyte Antigen F Presents Peptides and Regulates Immunity through Interactions with NK Cell Receptors. *Immunity* 2017, 46:1018–1029 e1017. [PubMed: 28636952] This study reveals that peptide-bound HLA-F can inhibit NK cells LIR-1.
59. Lee N, Llano M, Carretero M, Ishitani A, Navarro F, Lopez-Botet M, Geraghty DE: HLA-E is a major ligand for the natural killer inhibitory receptor CD94/NKG2A. *Proc Natl Acad Sci U S A* 1998, 95:5199–5204. [PubMed: 9560253]
60. Braud VM, Allan DS, O'Callaghan CA, Soderstrom K, D'Andrea A, Ogg GS, Lazetic S, Young NT, Bell JI, Phillips JH, et al.: HLA-E binds to natural killer cell receptors CD94/NKG2A, B and C. *Nature* 1998, 391:795–799. [PubMed: 9486650]
- 61•. Davis ZB, Cogswell A, Scott H, Mertsching A, Boucau J, Wambua D, Le Gall S, Planelles V, Campbell KS, Barker E: A Conserved HIV-1-Derived Peptide Presented by HLA-E Renders Infected T-cells Highly Susceptible to Attack by NKG2A/CD94-Bearing Natural Killer Cells. *PLoS Pathog* 2016, 12:e1005421. [PubMed: 26828202] The authors show that a conserved HIV-1 peptide binds to HLA-E, thereby rendering infected T cells susceptible to NK cell killing by limiting NKG2A-derived inhibitory signals.
- 62•. Ramsuran V, Naranbhai V, Horowitz A, Qi Y, Martin MP, Yuki Y, Gao X, Walker-Sperling V, Del Prete GQ, Schneider DK, et al.: Elevated HLA-A expression impairs HIV control through inhibition of NKG2A-expressing cells. *Science* 2018, 359:86–90. [PubMed: 29302013] Using HIV-infected individuals from 21 distinct cohorts, the authors uncover HLA genetic control of

HLA-A expression and show that HLA-A:NKG2A inhibition of NK cells impairs viral control and clearance.

63. Horowitz A, Djaoud Z, Nemat-Gorgani N, Blokhuis J, Hilton HG, Beziat V, Malmberg KJ, Norman PJ, Guethlein LA, Parham P: Class I HLA haplotypes form two schools that educate NK cells in different ways. *Sci Immunol* 2016, 1:eaag1672. [PubMed: 27868107]
64. Bernardini G, Antonangeli F, Bonanni V, Santoni A: Dysregulation of Chemokine/Chemokine Receptor Axes and NK Cell Tissue Localization during Diseases. *Front Immunol* 2016, 7:402. [PubMed: 27766097]
65. Pesce S, Moretta L, Moretta A, Marcenaro E: Human NK Cell Subsets Redistribution in Pathological Conditions: A Role for CCR7 Receptor. *Front Immunol* 2016, 7:414. [PubMed: 27774094]
66. Salazar-Mather TP, Ishikawa R, Biron CA: NK cell trafficking and cytokine expression in splenic compartments after IFN induction and viral infection. *J Immunol* 1996, 157:3054–3064. [PubMed: 8816415]
67. Medaglia C, Giladi A, Stoler-Barak L, De Giovanni M, Salame TM, Biram A, David E, Li H, Iannacone M, Shulman Z, et al.: Spatial reconstruction of immune niches by combining photoactivatable reporters and scRNA-seq. *Science* 2017, 358:1622–1626. [PubMed: 29217582]
68. Crome SQ, Lang PA, Lang KS, Ohashi PS: Natural killer cells regulate diverse T cell responses. *Trends Immunol* 2013, 34:342–349. [PubMed: 23601842]
69. Crouse J, Xu HC, Lang PA, Oxenius A: NK cells regulating T cell responses: mechanisms and outcome. *Trends Immunol* 2015, 36:49–58. [PubMed: 25432489]
70. Waggoner SN, Daniels KA, Welsh RM: Therapeutic depletion of natural killer cells controls persistent infection. *J Virol* 2014, 88:1953–1960. [PubMed: 24284324]
71. Lang PA, Lang KS, Xu HC, Grusdat M, Parish IA, Recher M, Elford AR, Dhanji S, Shaabani N, Tran CW, et al.: Natural killer cell activation enhances immune pathology and promotes chronic infection by limiting CD8+ T-cell immunity. *Proc Natl Acad Sci U S A* 2012, 109:1210–1215. [PubMed: 22167808]
72. Waggoner SN, Cornberg M, Selin LK, Welsh RM: Natural killer cells act as rheostats modulating antiviral T cells. *Nature* 2012, 481:394–398.
73. Rydzynski C, Daniels KA, Karmele EP, Brooks TR, Mahl SE, Moran MT, Li C, Sutiwisesak R, Welsh RM, Waggoner SN: Generation of cellular immune memory and B-cell immunity is impaired by natural killer cells. *Nat Commun* 2015, 6:6375. [PubMed: 25721802]
74. Blass E, Aid M, Martinot AJ, Larocca RA, Kang ZH, Badamchi-Zadeh A, Penaloza-MacMaster P, Reeves RK, Barouch DH: Adenovirus Vector Vaccination Impacts NK Cell Rheostat Function following Lymphocytic Choriomeningitis Virus Infection. *J Virol* 2018, 92. The authors demonstrate that adenovirus elicited memory T cells are less amenable to the immunoregulatory functions of NK cells during virus challenge.
75. Schuster IS, Wikstrom ME, Brizard G, Coudert JD, Estcourt MJ, Manzur M, O'Reilly LA, Smyth MJ, Trapani JA, Hill GR, et al.: TRAIL+ NK cells control CD4+ T cell responses during chronic viral infection to limit autoimmunity. *Immunity* 2014, 41:646–656. [PubMed: 25367576]
76. Andrews DM, Estcourt MJ, Andoniou CE, Wikstrom ME, Khong A, Voigt V, Fleming P, Tabarias H, Hill GR, van der Most RG, et al.: Innate immunity defines the capacity of antiviral T cells to limit persistent infection. *J Exp Med* 2010, 207:1333–1343. [PubMed: 20513749]
77. Su HC, Nguyen KB, Salazar-Mather TP, Ruzek MC, Dalod MY, Biron CA: NK cell functions restrain T cell responses during viral infections. *Eur J Immunol* 2001, 31:3048–3055. [PubMed: 11592081]
78. Soderquest K, Walzer T, Zafirova B, Klavinskis LS, Polic B, Vivier E, Lord GM, Martin-Fontecha A: Cutting edge: CD8+ T cell priming in the absence of NK cells leads to enhanced memory responses. *J Immunol* 2011, 186:3304–3308. [PubMed: 21307295]
79. Peppas D, Gill US, Reynolds G, Easom NJ, Pallett LJ, Schurich A, Micco L, Nebbia G, Singh HD, Adams DH, et al.: Up-regulation of a death receptor renders antiviral T cells susceptible to NK cell-mediated deletion. *J Exp Med* 2013, 210:99–114. [PubMed: 23254287]

80. Hatfield SD, Daniels KA, O'Donnell CL, Waggoner SN, Welsh RM: Weak vaccinia virus-induced NK cell regulation of CD4 T cells is associated with reduced NK cell differentiation and cytolytic activity. *Virology* 2018, 519:131–144. [PubMed: 29715623]
81. Xu HC, Huang J, Pandya AA, Lang E, Zhuang Y, Thons C, Timm J, Haussinger D, Colonna M, Cantor H, et al.: Lymphocytes Negatively Regulate NK Cell Activity via Qa-1b following Viral Infection. *Cell Rep* 2017, 21:2528–2540. [PubMed: 29186689]
82. Lu L, Ikizawa K, Hu D, Werneck MB, Wucherpfennig KW, Cantor H: Regulation of activated CD4+ T cells by NK cells via the Qa-1-NKG2A inhibitory pathway. *Immunity* 2007, 26:593–604. [PubMed: 17509909]
- 83•. Bradley TPD, Pedroza-Pacheco I, Li D, Cain D, Henao R, Venkat V, Hora B, Chen Y, Vandergrift N, Overman G, Edwards R, Woods C, Tomaras G, Ferrari G, Ginsburg G, M. CCohen M, Moody M, Borrow P, Haynes B: RAB11FIP5 expression and altered natural killer cell function are associated with induction of HIV broadly neutralizing antibody responses. *Cell* 2018, In press. In cohorts of individuals infected with HIV-1, the authors discover that development of broadly neutralizing antibodies against the virus is correlated with accumulation of dysfunctional NK cells that express high levels of RAB11FIP5.
- 84•. Rydzynski CE, Cranert SA, Zhou JQ, Xu H, Kleinstein SH, Singh H, Waggoner SN: Affinity maturation is impaired by natural killer cell suppression of germinal centers. *Cell Reports* 2018, In press. This study reveals that NK cell immunoregulatory functions limit somatic mutation and affinity maturation of vaccine-elicited immunoglobulin responses in the germinal center.
85. Cerwenka A, Lanier LL: Natural killer cell memory in infection, inflammation and cancer. *Nat Rev Immunol* 2016, 16:112–123. [PubMed: 26806484]
86. Schlums H, Cichocki F, Tesi B, Theorell J, Beziat V, Holmes TD, Han H, Chiang SC, Foley B, Mattsson K, et al.: Cytomegalovirus infection drives adaptive epigenetic diversification of NK cells with altered signaling and effector function. *Immunity* 2015, 42:443–456. [PubMed: 25786176]
87. Lee J, Zhang T, Hwang I, Kim A, Nitschke L, Kim M, Scott JM, Kamimura Y, Lanier LL, Kim S: Epigenetic modification and antibody-dependent expansion of memory-like NK cells in human cytomegalovirus-infected individuals. *Immunity* 2015, 42:431–442. [PubMed: 25786175]
88. Gillard GO, Bivas-Benita M, Hovav AH, Grandpre LE, Panas MW, Seaman MS, Haynes BF, Letvin NL: Thy1+ NK [corrected] cells from vaccinia virus-primed mice confer protection against vaccinia virus challenge in the absence of adaptive lymphocytes. *PLoS Pathog* 2011, 7:e1002141. [PubMed: 21829360]
89. Reeves RK, Li H, Jost S, Blass E, Li H, Schafer JL, Varner V, Manickam C, Eslamizar L, Altfeld M, et al.: Antigen-specific NK cell memory in rhesus macaques. *Nat Immunol* 2015, 16:927–932. [PubMed: 26193080]
90. Paust S, Gill HS, Wang BZ, Flynn MP, Moseman EA, Senman B, Szczepanik M, Telenti A, Askenase PW, Compans RW, et al.: Critical role for the chemokine receptor CXCR6 in NK cell-mediated antigen-specific memory of haptens and viruses. *Nat Immunol* 2010, 11:1127–1135. [PubMed: 20972432]
91. O'Leary JG, Goodarzi M, Drayton DL, von Andrian UH: T cell- and B cell-independent adaptive immunity mediated by natural killer cells. *Nat Immunol* 2006, 7:507–516. [PubMed: 16617337]
92. Cooper MA, Elliott JM, Keyel PA, Yang L, Carrero JA, Yokoyama WM: Cytokine-induced memory-like natural killer cells. *Proc Natl Acad Sci U S A* 2009, 106:1915–1919. [PubMed: 19181844]
- 93•. Lau CM, Adams NM, Geary CD, Weizman OE, Rapp M, Pritykin Y, Leslie CS, Sun JC: Epigenetic control of innate and adaptive immune memory. *Nat Immunol* 2018, 19:963–972. [PubMed: 30082830] This study reveals dynamic changes in epigenetic modifications during evolution of memory NK cell responses to virus infection.
94. Tesi B, Schlums H, Cichocki F, Bryceson YT: Epigenetic Regulation of Adaptive NK Cell Diversification. *Trends Immunol* 2016, 37:451–461. [PubMed: 27160662]
95. Cichocki F, Wu CY, Zhang B, Felices M, Tesi B, Tuininga K, Dougherty P, Taras E, Hinderlie P, Blazar BR, et al.: ARID5B regulates metabolic programming in human adaptive NK cells. *J Exp Med* 2018, 215:2379–2395. [PubMed: 30061358]

96. Rapp M, Lau CM, Adams NM, Weizman OE, O'Sullivan TE, Geary CD, Sun JC: Core-binding factor beta and Runx transcription factors promote adaptive natural killer cell responses. *Sci Immunol* 2017, 2.
97. Adams NM, Lau CM, Fan X, Rapp M, Geary CD, Weizman OE, Diaz-Salazar C, Sun JC: Transcription Factor IRF8 Orchestrates the Adaptive Natural Killer Cell Response. *Immunity* 2018, 48:1172–1182 e1176. [PubMed: 29858012]
98. Geary CD, Krishna C, Lau CM, Adams NM, Gearty SV, Pritykin Y, Thomsen AR, Leslie CS, Sun JC: Non-redundant ISGF3 Components Promote NK Cell Survival in an Auto-regulatory Manner during Viral Infection. *Cell Rep* 2018, 24:1949–1957 e1946. [PubMed: 30134157]
99. Lunemann S, Martrus G, Holzemer A, Chapel A, Ziegler M, Korner C, Garcia Beltran W, Carrington M, Wedemeyer H, Altfeld M: Sequence variations in HCV core-derived epitopes alter binding of KIR2DL3 to HLA-C *03:04 and modulate NK cell function. *J Hepatol* 2016, 65:252–258. [PubMed: 27057987]
100. Holzemer A, Thobakgale CF, Jimenez Cruz CA, Garcia-Beltran WF, Carlson JM, van Teijlingen NH, Mann JK, Jaggernath M, Kang SG, Korner C, et al.: Selection of an HLA-C*03:04-Restricted HIV-1 p24 Gag Sequence Variant Is Associated with Viral Escape from KIR2DL3+ Natural Killer Cells: Data from an Observational Cohort in South Africa. *PLoS Med* 2015, 12:e1001900; discussion e1001900. [PubMed: 26575988]
101. van Teijlingen NH, Holzemer A, Korner C, Garcia-Beltran WF, Schafer JL, Fadda L, Suscovich TJ, Brander C, Carrington M, Evans DT, et al.: Sequence variations in HIV-1 p24 Gag-derived epitopes can alter binding of KIR2DL2 to HLA-C*03:04 and modulate primary natural killer cell function. *AIDS* 2014, 28:1399–1408. [PubMed: 24785948]
102. O'Connor GM, Vivian JP, Gostick E, Pymm P, Lafont BA, Price DA, Rossjohn J, Brooks AG, McVicar DW: Peptide-Dependent Recognition of HLA-B*57:01 by KIR3DS1. *J Virol* 2015, 89:5213–5221. [PubMed: 25740999]
103. Alter G, Heckerman D, Schneidewind A, Fadda L, Kadie CM, Carlson JM, Oniangue-Ndza C, Martin M, Li B, Khakoo SI, et al.: HIV-1 adaptation to NK-cell-mediated immune pressure. *Nature* 2011, 476:96–100. [PubMed: 21814282]
104. Elemans M, Boelen L, Rasmussen M, Buus S, Asquith B: HIV-1 adaptation to NK cell-mediated immune pressure. *PLoS Pathog* 2017, 13:e1006361. [PubMed: 28582449]
105. Zhang T, Scott JM, Hwang I, Kim S: Cutting edge: antibody-dependent memory-like NK cells distinguished by FcRgamma deficiency. *J Immunol* 2013, 190:1402–1406. [PubMed: 23345329]
106. Haynes BF, Gilbert PB, McElrath MJ, Zolla-Pazner S, Tomaras GD, Alam SM, Evans DT, Montefiori DC, Karnasuta C, Sutthent R, et al.: Immune-correlates analysis of an HIV-1 vaccine efficacy trial. *N Engl J Med* 2012, 366:1275–1286. [PubMed: 22475592]
107. Bernard NF, Kiani Z, Tremblay-McLean A, Kant SA, Leeks CE, Dupuy FP: Natural Killer (NK) Cell Education Differentially Influences HIV Antibody-Dependent NK Cell Activation and Antibody-Dependent Cellular Cytotoxicity. *Front Immunol* 2017, 8:1033. [PubMed: 28883824]
108. Lambotte O, Ferrari G, Moog C, Yates NL, Liao HX, Parks RJ, Hicks CB, Owzar K, Tomaras GD, Montefiori DC, et al.: Heterogeneous neutralizing antibody and antibody-dependent cell cytotoxicity responses in HIV-1 elite controllers. *AIDS* 2009, 23:897–906. [PubMed: 19414990]
109. Kim S, Sunwoo JB, Yang L, Choi T, Song YJ, French AR, Vlahiotis A, Piccirillo JF, Cella M, Colonna M, et al.: HLA alleles determine differences in human natural killer cell responsiveness and potency. *Proc Natl Acad Sci U S A* 2008, 105:3053–3058. [PubMed: 18287063]
110. Kim S, Poursine-Laurent J, Truscott SM, Lybarger L, Song YJ, Yang L, French AR, Sunwoo JB, Lemieux S, Hansen TH, et al.: Licensing of natural killer cells by host major histocompatibility complex class I molecules. *Nature* 2005, 436:709–713. [PubMed: 16079848]
111. Kristensen AB, Kent SJ, Parsons MS: Contribution of NK Cell Education to both Direct and Anti-HIV-1 Antibody-Dependent NK Cell Functions. *J Virol* 2018, 92. The authors demonstrate that distinct pathways to NK cell education impact the functional capacity of mature NK cells to mediate ADCC.
112. Dai HS, Griffin N, Bolyard C, Mao HC, Zhang J, Cripe TP, Suenaga T, Arase H, Nakano, Chiocca EA, et al.: The Fc Domain of Immunoglobulin Is Sufficient to Bridge NK Cells with Virally Infected Cells. *Immunity* 2017, 47:159–170 e110. [PubMed: 28723548] The authors are

the first to demonstrate that viral IgG-Fc binding proteins can form a bridge between NK cells and virus infected cells that enhances viral control.

113. York IA, Johnson DC: Direct contact with herpes simplex virus-infected cells results in inhibition of lymphokine-activated killer cells because of cell-to-cell spread of virus. *J Infect Dis* 1993, 168:1127–1132. [PubMed: 8228345]
114. Strickland P, Dunston G, Roane P, Vanderpool E: Multiplication of herpes simplex virus in large granular lymphocytes that co-fractionate with human natural killer cell activity. *Immunol Invest* 1989, 18:785–795. [PubMed: 2548958]
115. Isobe Y, Sugimoto K, Yang L, Tamayose K, Egashira M, Kaneko T, Takada K, Oshimi K: Epstein-Barr virus infection of human natural killer cell lines and peripheral blood natural killer cells. *Cancer Res* 2004, 64:2167–2174. [PubMed: 15026359]
116. Lusso P, Malnati MS, Garzino-Demo A, Crowley RW, Long EO, Gallo RC: Infection of natural killer cells by human herpesvirus 6. *Nature* 1993, 362:458–462. [PubMed: 7681936]
117. Kirwan S, Merriam D, Barsby N, McKinnon A, Burshtyn DN: Vaccinia virus modulation of natural killer cell function by direct infection. *Virology* 2006, 347:75–87. [PubMed: 16387342]
118. Mao H, Tu W, Qin G, Law HK, Sia SF, Chan PL, Liu Y, Lam KT, Zheng J, Peiris M, et al.: Influenza virus directly infects human natural killer cells and induces cell apoptosis. *J Virol* 2009, 83:9215–9222. [PubMed: 19587043]
119. Bernstein HB, Wang G, Plasterer MC, Zack JA, Ramasastry P, Mumenthaler SM, Kitchen CM: CD4+ NK cells can be productively infected with HIV, leading to downregulation of CD4 expression and changes in function. *Virology* 2009, 387:59–66. [PubMed: 19251297]
120. Zhao J, Cheng L, Wang H, Yu H, Tu B, Fu Q, Li G, Wang Q, Sun Y, Zhang X, et al.: Infection and depletion of CD4+ group-1 innate lymphoid cells by HIV-1 via type-I interferon pathway. *PLoS Pathog* 2018, 14:e1006819. [PubMed: 29304123]
121. Campbell TM, McSharry BP, Steain M, Ashhurst TM, Slobedman B, Abendroth A: Varicella zoster virus productively infects human natural killer cells and manipulates phenotype. *PLoS Pathog* 2018, 14:e1006999. [PubMed: 29709039] This paper reveals how a herpesvirus can infect and manipulate NK cells to promote viral persistence and pathogenesis.
122. Orange JS, Fassett MS, Koopman LA, Boyson JE, Strominger JL: Viral evasion of natural killer cells. *Nat Immunol* 2002, 3:1006–1012. [PubMed: 12407408]
123. Matusali G, Potesta M, Santoni A, Cerboni C, Doria M: The human immunodeficiency virus type 1 Nef and Vpu proteins downregulate the natural killer cell-activating ligand PVR. *J Virol* 2012, 86:4496–4504. [PubMed: 22301152]
124. Shah AH, Sowrirajan B, Davis ZB, Ward JP, Campbell EM, Planelles V, Barker E: Degranulation of natural killer cells following interaction with HIV-1-infected cells is hindered by downmodulation of NTB-A by Vpu. *Cell Host Microbe* 2010, 8:397–409. [PubMed: 21075351]
125. Voigt V, Forbes CA, Tonkin JN, Degli-Esposti MA, Smith HR, Yokoyama WM, Scalzo AA: Murine cytomegalovirus m157 mutation and variation leads to immune evasion of natural killer cells. *Proc Natl Acad Sci U S A* 2003, 100:13483–13488. [PubMed: 14597723]
126. Pyzik M, Dumaine A, Charbonneau B, Fodil-Cornu N, Jonjic S, Vidal SM: Viral MHC class I-like molecule allows evasion of NK cell effector responses in vivo. *J Immunol* 2014, 193:6061–6069. [PubMed: 25392524]
127. Aguilar OA, Berry R, Rahim MMA, Reichel JJ, Popovic B, Tanaka M, Fu Z, Balaji GR, Lau TNH, Tu MM, et al.: A Viral Immuno-evasin Controls Innate Immunity by Targeting the Prototypical Natural Killer Cell Receptor Family. *Cell* 2017, 169:58–71 e14. [PubMed: 28340350] The authors dissect a long-standing conundrum by discovering complex interplay between the m12 immuno-evasin of MCMV and the NKR-P1 receptors in control of NK cell antiviral functions.
128. McKinney EF, Smith KG: T cell exhaustion and immune-mediated disease-the potential for therapeutic exhaustion. *Curr Opin Immunol* 2016, 43:74–80. [PubMed: 27744240]
129. Barber DL, Wherry EJ, Masopust D, Zhu B, Allison JP, Sharpe AH, Freeman GJ, Ahmed R: Restoring function in exhausted CD8 T cells during chronic viral infection. *Nature* 2006, 439:682–687. [PubMed: 16382236]

130. Zou W, Wolchok JD, Chen L: PD-L1 (B7-H1) and PD-1 pathway blockade for cancer therapy: Mechanisms, response biomarkers, and combinations. *Sci Transl Med* 2016, 8:328rv324.
131. Gill S, Vasey AE, De Souza A, Baker J, Smith AT, Kohrt HE, Florek M, Gibbs KD Jr., Tate K, Ritchie DS, et al.: Rapid development of exhaustion and down-regulation of eomesodermin limit the antitumor activity of adoptively transferred murine natural killer cells. *Blood* 2012, 119:5758–5768. [PubMed: 22544698]
132. Beldi-Ferchiou A, Lambert M, Dogniaux S, Vely F, Vivier E, Olive D, Dupuy S, Levasseur F, Zucman D, Lebbe C, et al.: PD-1 mediates functional exhaustion of activated NK cells in patients with Kaposi sarcoma. *Oncotarget* 2016, 7:72961–72977. [PubMed: 27662664]
133. Hsu J, Hodgins JJ, Marathe M, Nicolai CJ, Bourgeois-Daigneault MC, Trevino TN, Azimi CS, Scheer AK, Randolph HE, Thompson TW, et al.: Contribution of NK cells to immunotherapy mediated by PD-1/PD-L1 blockade. *J Clin Invest* 2018. This study reveals that blockade of PD-1/PD-L1 interactions unleashes NK cell antitumor functions, suggesting that NK cells may contribute to clinical efficacy of checkpoint inhibitors.
134. Zhang Q, Bi J, Zheng X, Chen Y, Wang H, Wu W, Wang Z, Wu Q, Peng H, Wei H, et al.: Blockade of the checkpoint receptor TIGIT prevents NK cell exhaustion and elicits potent antitumor immunity. *Nat Immunol* 2018. The authors discover that TIGIT is a potent checkpoint inhibitor of NK cell function.
135. Molgora M, Bonavita E, Ponzetta A, Riva F, Barbagallo M, Jaillon S, Popovic B, Bernardini G, Magrini E, Gianni F, et al.: IL-1R8 is a checkpoint in NK cells regulating anti-tumour and antiviral activity. *Nature* 2017, 551:110–114. [PubMed: 29072292] This study highlights a role for a member of the IL-1 receptor family in restraint of NK cell function in the context of both cancer and virus infection.
136. da Silva IP, Gallois A, Jimenez-Baranda S, Khan S, Anderson AC, Kuchroo VK, Osman I, Bhardwaj N: Reversal of NK-cell exhaustion in advanced melanoma by Tim-3 blockade. *Cancer Immunol Res* 2014, 2:410–422. [PubMed: 24795354]
137. Castro W, Chelbi ST, Niogret C, Ramon-Barros C, Welten SPM, Osterheld K, Wang H, Rota G, Morgado L, Vivier E, et al.: The transcription factor Rfx7 limits metabolism of NK cells and promotes their maintenance and immunity. *Nat Immunol* 2018, 19:809–820. [PubMed: 29967452]
138. Loftus RM, Assmann N, Kedia-Mehta N, O'Brien KL, Garcia A, Gillespie C, Hukelmann JL, Oefner PJ, Lamond AI, Gardiner CM, et al.: Amino acid-dependent cMyc expression is essential for NK cell metabolic and functional responses in mice. *Nat Commun* 2018, 9:2341. [PubMed: 29904050]
139. Cong J, Wang X, Zheng X, Wang D, Fu B, Sun R, Tian Z, Wei H: Dysfunction of Natural Killer Cells by FBP1-Induced Inhibition of Glycolysis during Lung Cancer Progression. *Cell Metab* 2018, 28:243–255 e245. [PubMed: 30033198]
140. Assmann N, O'Brien KL, Donnelly RP, Dyck L, Zaiatz-Bittencourt V, Loftus RM, Heinrich P, Oefner PJ, Lynch L, Gardiner CM, et al.: Srebp-controlled glucose metabolism is essential for NK cell functional responses. *Nat Immunol* 2017, 18:1197–1206. [PubMed: 28920951]
141. Mao Y, van Hoef V, Zhang X, Wennerberg E, Lorent J, Witt K, Masvidal L, Liang S, Murray S, Larsson O, et al.: IL-15 activates mTOR and primes stress-activated gene expression leading to prolonged antitumor capacity of NK cells. *Blood* 2016, 128:1475–1489. [PubMed: 27465917]
142. Keating SE, Zaiatz-Bittencourt V, Loftus RM, Keane C, Brennan K, Finlay DK, Gardiner CM: Metabolic Reprogramming Supports IFN-gamma Production by CD56bright NK Cells. *J Immunol* 2016, 196:2552–2560. [PubMed: 26873994]
143. Schafer JR, Salzillo TC, Chakravarti N, Kararoudi MN, Trikha P, Foltz JA, Wang R, Li S, Lee DA: Education-dependent activation of glycolysis promotes the cytolytic potency of licensed human natural killer cells. *J Allergy Clin Immunol* 2018.
144. Goh CC, Roggerson KM, Lee HC, Golden-Mason L, Rosen HR, Hahn YS: Hepatitis C Virus-Induced Myeloid-Derived Suppressor Cells Suppress NK Cell IFN-gamma Production by Altering Cellular Metabolism via Arginase-1. *J Immunol* 2016, 196:2283–2292. [PubMed: 26826241]
145. Mah AY, Rashidi A, Keppel MP, Saucier N, Moore EK, Alinger JB, Tripathy SK, Agarwal SK, Jeng EK, Wong HC, et al.: Glycolytic requirement for NK cell cytotoxicity and cytomegalovirus

control. JCI Insight 2017, 2. The authors demonstrate that NK cell control of CMV infection requires glucose metabolism, and that IL-15 therapy can overcome this metabolic demand.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Highlights

- NK cells protect by killing virus-infected cells and making IFN- γ
- Viruses evade immunity by modulating NK-cell function
- NK cells can participate in and regulate long-lived adaptive immunity

Figure 1. Contributions of NK cells to acute, chronic, and secondary infections with viruses. During acute challenge with new virus, NK cells contribute to host immunity by secreting proinflammatory cytokines (e.g. IFN- γ), lysing virus-infected cells, and exerting immunosuppressive effects indirectly via antigen-presenting cells (APC) or directly on T and B cells. Establishment of chronic infection is associated with functional and phenotypic exhaustion of both T (Tex) and NK cells (exNK), where some NK cells contribute to maintenance of T cell exhaustion and viral persistence. Viral exposure is associated with development of memory NK cells (mNK), T cells (Tm), and B cells (Bm). Following resolution of acute infection, re-exposure to the original virus prompts more potent antiviral and ADCC responses of mNK cells, altered capacity of naïve NK or mNK cells to suppress Tm, and amplification of NK cell responses by Tm-derived cytokines (e.g. IL-2).

Figure 2. NK cell sensitivity to viral modulation of HLA.

A) Wild type viral peptides bind to HLA-C and suppress NK-cell cytotoxicity upon interaction with KIR2DL1/2 on NK cells. Certain viral variants decrease HLA-C expression or produce mutant peptides with reduced binding affinity to HLA-C, resulting in reduced KIR2DL1/2-mediated inhibition of NK cells. B) Expression of HLA-E bound self-peptides on uninfected cells protects from NK-cell killing via NKG2A inhibitory signaling. Certain viral peptides displayed on HLA-E of virus-infected cells obstruct HLA-E interaction with NKG2A, rendering virus-infected cells susceptible to NK-cell cytotoxicity. C) Open conformers of HLA-F preferentially interact with KIR3DS1 and activate NK cells. HLA-F occupation by viral peptides leads to preferential interaction with LIR-1, thereby suppressing NK cell activity. D) Host alleles promoting low HLA-A expression coupled

with specific HLA-B alleles provoke low expression of HLA-E and weaker NKG2A control of NK cells. Alternatively, alleles associated with high HLA-A expression and HLA-B alleles associated with education via NKG2A render high HLA-E expression and strong NKG2A modulation of NK cell function.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript