Published in final edited form as:

Curr Opin Virol. 2019 February ; 34: 10–17. doi:10.1016/j.coviro.2018.11.006.

Hepatitis E virus: Reasons for emergence in humans

Harini Sooryanarain and Xiang-Jin Meng*

Department of Biomedical Sciences and Pathobiology, Virginia-Maryland College of Veterinary Medicine, Virginia Polytechnic Institute and State University, Blacksburg, VA

Abstract

Hepatitis E virus (HEV) infects both humans and other animal species. Recently, we have seen a steady increase in autochthonous cases of human HEV infection in certain areas especially in Europe, and large outbreaks in several African countries among the displaced population. This mini-review critically analyzes potential host, environmental, and viral factors that may be associated with the emergence of hepatitis E in humans. The existence of numerous HEV reservoir animals such as pig, deer and rabbit results in human exposure to infected animals via direct contact or through animal meat consumption. Contamination of drinking, irrigation and coastal water by animal and human wastes lead to emergence of endemic cases in industrialized countries and outbreaks in displaced communities especially in war-torn countries.

Introduction

Hepatitis E virus (HEV) is a single-stranded positive-sense RNA (~7.2 kb size) virus in the family *Hepeviridae*, which comprised of two genera *Orthohepevirus* (all mammalian and avian HEV strains) and *Piscihepevirus* (Cutthroat trout virus) [1,2]. The genus *Orthohepevirus* comprises of four species *Orthohepevirus* A-D. Within the species *Orthohepevirus* A, at least 8 distinct genotypes have been identified [2]: genotypes 1 and 2 HEVs infect only humans, genotypes 3 and 4 HEVs infect humans and several other animal species, genotypes 5 and 6 HEVs infect wild boars, genotype 7 HEV infects dromedary camels and possibly humans [3–5], and genotype 8 HEV infects Bactrian camel [6].

In humans, HEV infection usually results in an acute self-limiting disease, and the majority of infections are asymptomatic. The patterns of clinical disease vary among different HEV genotypes [7]. Genotypes 1 and 2 HEVs typically cause self-limiting acute infections, and are transmitted through contaminated water. Infected pregnant women, especially with genotype 1 HEV, have higher risk of developing fulminant hepatitis with a higher mortality rate of up to 30% [8]. In contrast, genotypes 3 and 4 HEVs are zoonotic, transmitted via consumption of animal meats or direct contact with infected animals, and causes mainly sporadic and cluster cases of hepatitis E [3]. Genotype 3, and to lesser extent, genotype 4

*Address for correspondence: X.J. Meng, Department of Biomedical Sciences and Pathobiology, Virginia Tech, 1981 Kraft Drive, Blacksburg, VA 24061-0913, xjmeng@vt.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

HEVs can establish chronic infection in immunocompromised individuals such as organ transplant recipients [8].

In this mini-review, we critically analyze the potential host, viral, and environmental factors that are associated with the emergence of HEV infection in humans.

Epidemiology

The seroprevalence of HEV infection varies in populations and the serological assays used worldwide. For example, more than 30% in German general population [9], 60.5% blood donor population in India [10], and 52.5% in Southwestern France were tested seropositive for HEV antibodies [11]. Cases of hepatitis E in humans are associated with genotypes 1–4 HEV, and in one rare instance, genotype 7 HEV [7] (Fig 1). The genotypes 1 and 2 HEV strains are endemic to countries in Asia and Africa, and also in Mexico [7,12,13]. In addition to historical large explosive outbreaks in Asia [14–16], in recent years, genotype 1 HEV infections in humans were also reported from individuals in Israel [17], Italy [18], and Germany [19] who are usually associated with prior travel to endemic countries. Two recent large outbreaks associated with genotype 1 HEV infection have been reported from Niger [20], and Uganda [21,22]. In the Niger outbreak, a total of 1,917 suspected cases were reported between January and September of 2017. During the 2013–2014 outbreak in Uganda, approximately 1,359 cases were suspected with an overall mortality rate of 2.2%, while a significantly higher mortality rate of ~65% was observed in pregnant women [22]. Smaller outbreaks have also been reported from Bangladesh [23], and India [24] between 2012 and 2014. A higher HEV seroprevalence is also reported amongst refugee populations in war-torn countries such as South Sudan and Ethiopia [25,26]. Poor sanitation condition or inadequate access to clean drinking water in overcrowded setting enhances genotype 1 HEV transmission rates in humans from developing countries, while genotype 1 HEV cases in industrialized countries are primarily associated with a history of travel to HEV endemic regions.

Autochthonous cases associated with genotypes 3 and 4 HEV infection are commonly seen in industrialized countries, but also in certain developing countries [7,27,28]. A rise in autochthonous cases of HEV infection in humans from 2007 to 2013 was reported in Japan [29]. Interestingly, genotype 4 HEV was more predominant during 2007 – 2009 in Japan, whereas genotype 3 HEV infections have increased from 2010 – 2013 [29]. Importantly, genotypes 3 and 4 HEV strains are also prevalent amongst various animal species, including domestic pigs, wild boars, deer, mongoose, and rabbit in many countries including the United States, China, Japan, and numerous countries in Europe [3,30]. Interestingly, during the last decade a decline in anti-HEV prevalence in humans has been observed in the United States [31] as well as from several European countries [32,33]. Meanwhile, in the last 25 years in China, the frequency of genotype 4 HEV infection has overtaken genotype 1 HEV-associated infections [34], therefore suggesting that a country's sanitation conditions and zoonotic HEV sources likely determine the epidemiological shift within circulating HEV genotypes in a given human population.

Factors associated with emergence of hepatitis E cases caused by genotypes 1 and 2 HEVs

The prevalence of hepatitis E among a given human population is influenced by multilevel interactions between various factors, including living conditions and/or behavioral habits, host immune status, environmental factors that affect HEV persistence in a given geographical area, and viral factors which contribute to genetic diversity and host range of the virus. Genotypes 1 and 2 HEVs are typically transmitted to humans through contaminated water, and pregnancy increases mortality rates.

(1). Poor sanitation and lack of clean drinking water: The first HEV outbreak was reported in 1955 from India [16,35]. During 1980s and 1990s, numerous explosive large outbreaks were reported from India, Pakistan, Bangladesh, and China [14,16]. However, recent outbreaks have mainly been reported from war torn countries or countries with major humanitarian crises, such as Niger [20], Uganda [22], South Sudan [26], Ethiopia [26], and Chad [36], largely due to poor sanitation conditions and lack of clean drinking water [25]. In endemic area, such as China, a spatio-temporal study predicted 4.03 infection per 100,000 person [37]. The average attack rate of HEV infection varies from 2 – 4%, however, in regions with poor sanitation conditions and without clean drinking water, as seen in overcrowded communities in slums, refugee camps, and/or displaced population, the HEV attack rate can go up to 25% [25]. Since genotype 1 HEV is transmitted through fecal-oral route, an increased attack rate amongst humans living in a densely-populated region with poor sanitation conditions points out to the existence of human-to-human transmission. Further studies are needed to understand this mode of HEV transmission in a large population in order to develop more practical public health measures that could be implemented within economically impoverished and overcrowded communities such as refugee camps.

(2). High mortality during pregnancy: The case fatality rate during genotype 1 HEV infection is usually between 1–3%, while pregnancy significantly increases the mortality rate. Amongst HEV-infected pregnant women from Asian and African countries, a maternal case fatality rate of 20.8%, and fetal case fatality rate of 34.2% have been reported [38]. This case fatality rate can increase up to a staggering 61.2% in mothers with fulminant hepatic failure [38]. In addition to preexisting liver condition, serum viral loads in mothers were a significant predictor of vertical transmission [39]. Sporadic cases of genotype 3 HEV-associated infection of pregnant women were also reported in some countries such as Germany [40] and France [41,42]. However, unlike the genotype 1 HEV infections, the genotype 3 HEV infection spontaneously resolved in mothers with no evidence of vertical transmission and no HEV viral RNA was detected in the newborn. The exact reason contributing to fulminant hepatitis and high mortality in genotype 1 HEV-infected pregnant women remains unknown, however changes in hormonal and immune status and/or response during pregnancy may contribute to this phenomena. However, it should be pointed out that severe hepatitis was not reproduced in pregnant rhesus monkeys experimentally infected with genotype 1 HEV [43], or in pregnant sows experimentally-infected with genotype 3 HEV [44]. Therefore, future studies are warranted to delineate the mechanism of HEV-associated fulminant hepatitis and high mortality during pregnancy.

Factors associated with emergence of human hepatitis E caused by genotypes 3 and 4 HEVs

Genotypes 3 and 4 HEVs can establish chronic infections in immunocompromised patients [45,46], and are usually transmitted through consumption of contaminated animal meat, direct contact with infected animals, or less commonly through contaminated blood products. Since genotypes 3 and 4 HEVs are zoonotic in nature, with multiple animal reservoirs, cross-species HEV infections from reservoir animals to humans also influence the emergence of human hepatitis E within a geographical area.

Foodborne HEV infection via consumption of contaminated animal meats: It has been demonstrated that consumption of raw or undercooked meats or meat products from reservoir animals such as pig and deer represents an important risk factor for the emergence of hepatitis E in human population [8,28]. In liver transplant patients, consumption of undercooked game meat or pork products was associated with HEV infection and chronic liver disease [47]. Autochthonous HEV infection from Australia was reported in people who ate pork products in one restaurant during 2013–2014 [48]. A familial HEV outbreak was definitively linked to consumption of HEV-infected wild-boar meat in Spain [49]. Autochthonous cases of genotype 3 HEV from Germany and Italy have also been attributed to consumption of pork products. Pork liver sausages have been documented as sources for numerous cluster cases of foodborne hepatitis E. For example, figatelli, a traditional pig liver sausage widely consumed in France, was linked to cluster cases of HEV infection [50]. Similarly, in Japan, cluster cases of autochthonous hepatitis E are linked to consumption of raw meat [51,52]. Four patients who consumed raw deer meat were infected with genotype 3 HEV [52]. Autochthonous genotype 4 HEV infection due to consumption of raw deer meat have also been reported from South Korea [53]. It has been reported that a significant proportion of wild deer population is infected by HEV [54]. Recently, chronic/persistent HEV infections caused by genotype 4 HEV have been reported in a child with acute lymphoblastic leukemia in China [55], in a woman post-liver transplant from Taiwan [45], and in an individual from the United States with a fatally accelerated cirrhosis [56]. Clearly, foodborne transmission is partially responsible for the emergence of hepatitis E in humans, and the prevalence of diverse HEV genotypes in various farm and wild food animal populations is a risk factor to humans who consume raw or undercooked animal meat products.

Since HEV can contaminate animal meat and meat products, it is important to properly cook the meats prior to consumption. Incubation of HEV at 60°C or greater could inactivate up to 80% infectious virions [57], and can reduce HEV RNA levels [58,59]. Infectious HEV present in contaminated pig livers could be inactivated by cooking the meat product to an internal temperature of at least 71°C for 5 min [60]. Understanding the thermal stability of HEV will enable consumers to properly cook animal meats to reduce the chance for the emergence of foodborne hepatitis E.

Cross-species infection via direct contact with HEV-infected reservoir animals: Under experimental conditions, several animal strains of HEV can infect across species barriers. Swine HEV infects both rhesus macaques and chimpanzee [61,62], the

surrogates of man. Pigs may serve as a mixing vessel for different HEV strains due to their susceptibility to multiple HEV genotypes including genotypes 3, 4, 5 and 6 strains, and thus pose a public health risk [3]. Avian HEV infect chickens and turkeys but not rhesus monkeys [63,64], suggesting that avian HEV likely does not infect humans. The rabbit HEV can be experimentally transmitted to cynomolgus macaques [65], and infections of humans by rabbit HEV have been reported [66,67].

A high rate of anti-HEV seropositivity has been reported amongst veterinarians, swine farmers and hunters from many countries including the United States [68], Finland [69], China [70], Italy, and Poland [71,72]. For example, U.S. swine veterinarians were 1.51 times more likely to be IgG anti-HEV seropositive than blood donor controls. Importantly, subjects from traditionally major swine states such as Minnesota are more likely to be seropositive than those from traditionally non-swine states such as Alabama [68]. A significantly higher prevalence of IgG anti-HEV was also reported in individuals with occupational exposure to wild animals and environmental sources of animal wastes [73]. However, there was no association between anti-HEV seropositivity and exposure to pigs in the Netherlands [74]. Therefore, zoonotic HEV transmission in human population would be dependent on frequency and nature of contact exposure, the amount of viral loads in the infected animals, and the animal species.

Blood transfusion: Transfusion-transmitted cases of hepatitis E have been reported from many countries including Japan [75], U.K [76], France [77], Germany [78], Australia [79], and Canada [80]. HEV seroprevalence among blood donors varies from 3.4 % in Japan [81], 5.9% in Canada [82], 6.8% in Germany [78,83], to 22.4% in France [84]. A seroprevalence of 18.8% was reported in blood donors in the United States, however, during the follow-up study, there was no transfusion-transmitted HEV infection among the recipients [85]. A recent large-scale survey study from United States (N = 128,020) reported a 0.002% HEV RNA positivity amongst the plasma donors [86]. The risks associated with transfusion-transmitted HEV depend upon the frequency of viremia in blood donors, the viral loads in the donor blood, and the volume of plasma in the final transfused blood component [87].

Asymptomatic viremia has been reported in healthy blood donors from many countries, and the rate of HEV viremia in blood donors ranges from 1:600 to 1:42,000 [86,87]. The majority of viremic donors remained seronegative at the time of blood donation in Germany [88] and U.K [89], and had spontaneous virus clearance within a median time of 6 – 8 weeks. The lowest viral dose required for transmission was reportedly at 2×10^4 IUs, and 55% of the components containing this dose transmitted HEV infection [90].

Immunosuppressed patients with an extensive need for blood transfusion are at higher risk of developing transfusion-transmitted chronic HEV sequel [75,90]. Interestingly, a severe case of genotype 3 HEV infection was reported from France in a previously healthy male trauma patient who received massive transfusion of HEV-contaminated platelet pool [91]. Taken together, although blood transfusion is associated with the emergence of hepatitis E in humans especially in immunosuppressed patients requiring extensive transfusion, such cases are low. Therefore, this raises a question as to whether blood donors should be required to screen for HEV RNA. Although some European countries have already implemented

screening of blood donors for acute HEV infection marker such as HEV RNA [92], however, this practice is not yet implemented universally.

Environmental factors contributing to the persistence and emergence of HEV in humans

Studies on the environmental factors affecting persistence and emergence of HEV is very limited. HEV has been detected in environmental samples such as urban wastewater effluents, soil, and effluents from animal slaughterhouses. For example, approximately 25% of urban wastewater effluent was tested positive for genotype 3 HEV in France [93], and approximately 4.9% soil samples near river were tested positive for genotype 1 HEV in India [94]. Infectious swine HEV was detected in pig manure storage facilities in U.S. swine farms including lagoons and concrete pits, although evidence of drinking or surface water contamination on or near pig farms was not found [95]. Swine HEV has also been detected in effluents from pig farms [93,96], slaughter houses [93,96], and pig manure composite plants [97]. In regions with large operations of hog farms, swine manure or sewage water could potentially contaminate irrigation water or coastal waters with concomitant contamination of produce or shellfish. For example, genotype 3 HEV of swine origin was detected in mussels from Spain [98]. In Canada, genotype 3 HEV of swine origin was detected on farm-grown strawberries and irrigation water is suspected as the source of contamination [99]. Taken together, the available data clearly suggest a conducive fecal-contaminated wet environment that potentially aids in emergence of hepatitis E in humans.

Viral factors contributing to the host range and cross-species infection of HEV

Two major events, point mutations and recombination, attribute to the genetic variations and evolution of HEV genome. Accumulating evidence suggest that the ORF1 appears to be responsible for determining the host range and cross-species infection. It has been documented that point mutations affect HEV heterogeneity, and contribute to viral pathogenesis and susceptibility [100]. Recombination events can occur within different HEV genotypes, and between human and other animal HEV strains. An extensive heterogeneity in the hypervariable region (HVR) and macro domain of HEV genome is seen amongst the immunocompromised patients developing a chronic disease as compared to patients with resolving HEV infections [101]. Insertion and/or deletions in HVR of HEV ORF1 contributes to viral replication efficiency and host adaptations [102,103]. By utilizing intergenotypic chimeric viruses between human-exclusive genotype 1 HEV, and zoonotic genotypes 3 and 4 HEVs, it was found that the HEV ORF2 capsid gene does not determine host range [104,105]. Instead, the ORF1 is thought to be involved in host tropism of HEV. It has been shown that insertion of host RPS17 gene within the HEV HVR led to increased adaptability and infection of the virus in cells from different animal origins [106]. A clinical isolate of HEV from a chronically-infected patient with a 258 bp duplicated HVR fragment and a 24 bp RdRp-derived fragment was associated with increased viral replication [107]. HVR appears to be exchangeable between different HEV genotypes, however the chimeric viruses displayed a reduced virus replication kinetics [103]. Thus, it appears that HVR may contain genotype-specific sequences that are crucial for viral RNA replication and host adaptability, which may further influence the disease outcome.

Conclusion

Human behavioral and/or living conditions, as well as the presence of zoonotic HEV strains among animal species at a given geographic area determine changes in epidemiology of human HEV infection. More extensive studies are needed to understand the ecology and natural history of animal HEV strains as well as various modes of HEV transmission among host species, in order to formulate public health policies to prevent and control HEV infection in humans.

References

1. Smith DB, Simmonds P, International Committee on Taxonomy of Viruses Hepeviridae Study G, Jameel S, Emerson SU, Harrison TJ, Meng XJ, Okamoto H, Van der Poel WH, Purdy MA: Consensus proposals for classification of the family Hepeviridae. *J Gen Virol* 2014, 95:2223–2232. [PubMed: 24989172]
2. Purdy MA, Harrison TJ, Jameel S, Meng XJ, Okamoto H, Van der Poel WHM, Smith DB, Ictv Report C: ICTV Virus Taxonomy Profile: Hepeviridae. *J Gen Virol* 2017, 98:2645–2646. [PubMed: 29022866] * Recent ICTV report on taxonomy of the family Hepeviridae.
3. Meng XJ: Expanding Host Range and Cross-Species Infection of Hepatitis E Virus. *PLoS Pathog* 2016, 12:e1005695. [PubMed: 27490119] * This review gives an comprehensive overview on host ranges, cross-species infection, zoonotic risk, and food safety of HEV.
4. Spahr C, Knauf-Witzens T, Vahlenkamp T, Ulrich RG, John R: Hepatitis E virus and related viruses in wild, domestic and zoo animals: A review. *Zoonoses Public Health* 2018, 65:11–29. [PubMed: 28944602]
5. Lee GH, Tan BH, Teo EC, Lim SG, Dan YY, Wee A, Aw PP, Zhu Y, Hibberd ML, Tan CK, et al.: Chronic Infection With Camelid Hepatitis E Virus in a Liver Transplant Recipient Who Regularly Consumes Camel Meat and Milk. *Gastroenterology* 2016, 150:355–357 e353. [PubMed: 26551551]
6. Woo PC, Lau SK, Teng JL, Cao KY, Wernery U, Schountz T, Chiu TH, Tsang AK, Wong PC, Wong EY, et al.: New Hepatitis E Virus Genotype in Bactrian Camels, Xinjiang, China, 2013. *Emerg Infect Dis* 2016, 22:2219–2221. [PubMed: 27869607]
7. Sridhar S, Lau SK, Woo PC: Hepatitis E: A disease of reemerging importance. *J Formos Med Assoc* 2015, 114:681–690. [PubMed: 25773541]
8. Kmush BL, Nelson KE, Labrique AB: Risk factors for hepatitis E virus infection and disease. *Expert Rev Anti Infect Ther* 2015, 13:41–53. [PubMed: 25399510]
9. Wenzel JJ, Preiss J, Schemmerer M, Huber B, Jilg W: Test performance characteristics of Anti-HEV IgG assays strongly influence hepatitis E seroprevalence estimates. *J Infect Dis* 2013, 207:497–500. [PubMed: 23148290]
10. Katiyar H, Goel A, Sonker A, Yadav V, Sapun S, Chaudhary R, Aggarwal R: Prevalence of hepatitis E virus viremia and antibodies among healthy blood donors in India. *Indian J Gastroenterol* 2018, 37:342–346. [PubMed: 30159666]
11. Mansuy JM, Bendall R, Legrand-Abravanel F, Saune K, Miedouge M, Ellis V, Rech H, Destruel F, Kamar N, Dalton HR, et al.: Hepatitis E virus antibodies in blood donors, France. *Emerg Infect Dis* 2011, 17:2309–2312. [PubMed: 22172156]
12. Kim JH, Nelson KE, Panzner U, Kasture Y, Labrique AB, Wierzb TF: A systematic review of the epidemiology of hepatitis E virus in Africa. *BMC Infect Dis* 2014, 14:308. [PubMed: 24902967]
13. Echevarria JM, Gonzalez JE, Lewis-Ximenez LL, Dos Santos DR, Munne MS, Pinto MA, Pujol FH, Rodriguez-Lay LA: Hepatitis E virus infection in Latin America: a review. *J Med Virol* 2013, 85:1037–1045. [PubMed: 23588729]
14. Khuroo MS: Discovery of hepatitis E: the epidemic non-A, non-B hepatitis 30 years down the memory lane. *Virus Res* 2011, 161:3–14. [PubMed: 21320558]
15. Khuroo MS: Study of an epidemic of non-A, non-B hepatitis. Possibility of another human hepatitis virus distinct from post-transfusion non-A, non-B type. *Am J Med* 1980, 68:818–824. [PubMed: 6770682]

16. Balayan MS: Epidemiology of hepatitis E virus infection. *J Viral Hepat* 1997, 4:155–165. [PubMed: 9181524]
17. Erez-Granat O, Lachish T, Daudi N, Shouval D, Schwartz E: Hepatitis E in Israel: A nation-wide retrospective study. *World J Gastroenterol* 2016, 22:5568–5577. [PubMed: 27350735]
18. Bradanini L, Youkee D, Fabris P, Romano L, Brunetti E, Giordani MT: Acute hepatitis E virus infection in a migrant population in North East Italy: A retrospective analysis. *Travel Med Infect Dis* 2017, 20:37–42. [PubMed: 28411095]
19. Pischke S, Schulze-Zur-Wiesch J, Lutgehetmann M, Kreuels B, Lueth S, Kapaun P, Benten D, Schmiedel S, Sterneck M, Lohse AW, et al.: High clinical manifestation rate in an imported outbreak of hepatitis E genotype 1 infection in a German group of travellers returning from India. *Ann Hepatol* 2017, 16:57–62. [PubMed: 28051793]
20. Lagare A, Ibrahim A, Ousmane S, Issaka B, Zaneidou M, Kadade G, Testa J: Outbreak of Hepatitis E Virus Infection in Displaced Persons Camps in Diffa Region, Niger, 2017. *Am J Trop Med Hyg* 2018.
21. Teshale EH, Howard CM, Grytdal SP, Handzel TR, Barry V, Kamili S, Drobeniuc J, Okware S, Downing R, Tappero JW, et al.: Hepatitis E epidemic, Uganda. *Emerg Infect Dis* 2010, 16:126–129. [PubMed: 20031058]
22. Amany G, Kizito S, Nabukenya I, Kalyango J, Atuheire C, Nansumba H, Abwoye SA, Opio DN, Kibuuka E, Karamagi C: Risk factors, person, place and time characteristics associated with Hepatitis E Virus outbreak in Napak District, Uganda. *BMC Infect Dis* 2017, 17:451. [PubMed: 28651629]
23. Sazzad HM, Labrique AB, Teo CG, Luby SP, Gurley ES: Surveillance at Private Laboratories Identifies Small Outbreaks of Hepatitis E in Urban Bangladesh. *Am J Trop Med Hyg* 2017, 96:395–399. [PubMed: 27994104]
24. Singh MP, Majumdar M, Goyal K, Lakshmi PV, Bhatia D, Ratho RK: Investigation of suspected viral hepatitis outbreaks in North West India. *Diagn Microbiol Infect Dis* 2016, 84:309–314. [PubMed: 26853491]
25. Hakim MS, Wang W, Bramer WM, Geng J, Huang F, de Man RA, Peppelenbosch MP, Pan Q: The global burden of hepatitis E outbreaks: a systematic review. *Liver Int* 2017, 37:19–31. * This systematic review gives an detailed analysis of epidemiology of HEV outbreaks, outbreak settings and modes of transmission, control measures, and the distribution of HEV genotype responsible for the outbreaks.
26. Browne LB, Menkir Z, Kahi V, Maina G, Asnakew S, Tubman M, Elyas HZ, Nigatu A, Dak D, Maung UA, et al.: Notes from the field: hepatitis E outbreak among refugees from South Sudan - Gambella, Ethiopia, April 2014-January 2015. *MMWR Morb Mortal Wkly Rep* 2015, 64:537. [PubMed: 25996097]
27. Pisano MB, Martinez-Wassaf MG, Mirazo S, Fantilli A, Arbiza J, Debes JD, Re VE: Hepatitis E virus in South America: The current scenario. *Liver Int* 2018.
28. Clemente-Casares P, Ramos-Romero C, Ramirez-Gonzalez E, Mas A: Hepatitis E Virus in Industrialized Countries: The Silent Threat. *Biomed Res Int* 2016, 2016:9838041. [PubMed: 28070522] * A comprehensive review on HEV epidemiology in Europe and its associated risk factors.
29. Kanayama A, Arima Y, Yamagishi T, Kinoshita H, Sunagawa T, Yahata Y, Matsui T, Ishii K, Wakita T, Oishi K: Epidemiology of domestically acquired hepatitis E virus infection in Japan: assessment of the nationally reported surveillance data, 2007–2013. *J Med Microbiol* 2015, 64:752–758. [PubMed: 25976003]
30. Meng XJ: Zoonotic and foodborne transmission of hepatitis E virus. *Semin Liver Dis* 2013, 33:41–49. [PubMed: 23564388]
31. Teshale EH, Denniston MM, Drobeniuc J, Kamili S, Teo CG, Holmberg SD: Decline in hepatitis E virus antibody prevalence in the United States from 1988–1994 to 2009–2010. *J Infect Dis* 2015, 211:366–373. [PubMed: 25147277]
32. Niederhauser C, Widmer N, Hotz M, Tinguely C, Fontana S, Allemann G, Borri M, Infanti L, Sarraj A, Sigle J, et al.: Current hepatitis E virus seroprevalence in Swiss blood donors and apparent decline from 1997 to 2016. *Euro Surveill* 2018, 23.

33. Mahrt H, Schemmerer M, Behrens G, Leitzmann M, Jilg W, Wenzel JJ: Continuous decline of hepatitis E virus seroprevalence in southern Germany despite increasing notifications, 2003–2015. *Emerg Microbes Infect* 2018, 7:133. [PubMed: 30042475]
34. Liu P, Li L, Wang L, Bu Q, Fu H, Han J, Zhu Y, Lu F, Zhuang H: Phylogenetic analysis of 626 hepatitis E virus (HEV) isolates from humans and animals in China (1986–2011) showing genotype diversity and zoonotic transmission. *Infect Genet Evol* 2012, 12:428–434. [PubMed: 22306814]
35. Bhattacharjya B: The Delhi outbreak of infectious hepatitis: clinical features. *Indian J Med Res* 1957, 45:85–89.
36. Spina A, Lenglet A, Beversluis D, de Jong M, Vernier L, Spencer C, Andayi F, Kamau C, Vollmer S, Hogema B, et al.: A large outbreak of Hepatitis E virus genotype 1 infection in an urban setting in Chad likely linked to household level transmission factors, 2016–2017. *PLoS One* 2017, 12:e0188240. [PubMed: 29176816]
37. Liu K, Cai J, Wang S, Wu Z, Li L, Jiang T, Chen B, Cai G, Jiang Z, Chen Y, et al.: Identification of Distribution Characteristics and Epidemic Trends of Hepatitis E in Zhejiang Province, China from 2007 to 2012. *Sci Rep* 2016, 6:25407. [PubMed: 27146250]
38. Jin H, Zhao Y, Zhang X, Wang B, Liu P: Case-fatality risk of pregnant women with acute viral hepatitis type E: a systematic review and meta-analysis. *Epidemiol Infect* 2016, 144:2098–2106. [PubMed: 26939626]
39. Sharma S, Kumar A, Kar P, Agarwal S, Ramji S, Husain SA, Prasad S, Sharma S: Risk factors for vertical transmission of hepatitis E virus infection. *J Viral Hepat* 2017, 24:1067–1075. [PubMed: 28570034]
40. Tabatabai J, Wenzel JJ, Soboletski M, Flux C, Navid MH, Schnitzler P: First case report of an acute hepatitis E subgenotype 3c infection during pregnancy in Germany. *J Clin Virol* 2014, 61:170–172. [PubMed: 24996764]
41. Bouthry E, Benachi A, Vivanti AJ, Letamendia E, Vauloup-Fellous C, Roque-Afonso AM: Autochthonous Hepatitis E during Pregnancy, France. *Emerg Infect Dis* 2018, 24:1586–1587. [PubMed: 30016249]
42. Anty R, Ollier L, Peron JM, Nicand E, Cannavo I, Bongain A, Giordanengo V, Tran A: First case report of an acute genotype 3 hepatitis E infected pregnant woman living in South-Eastern France. *J Clin Virol* 2012, 54:76–78. [PubMed: 22336086]
43. Tsarev SA, Tsareva TS, Emerson SU, Rippey MK, Zack P, Shapiro M, Purcell RH: Experimental hepatitis E in pregnant rhesus monkeys: failure to transmit hepatitis E virus (HEV) to offspring and evidence of naturally acquired antibodies to HEV. *J Infect Dis* 1995, 172:31–37. [PubMed: 7797939]
44. Kasorndorkbua C, Thacker BJ, Halbur PG, Guenette DK, Buitenwerf RM, Royer RL, Meng XJ: Experimental infection of pregnant gilts with swine hepatitis E virus. *Can J Vet Res* 2003, 67:303–306. [PubMed: 14620868]
45. Wu CH, Ho CM, Tsai JH, Sun HY, Hu RH, Lee PH: First Case Genotype 4 Hepatitis E Infection After a Liver Transplant. *Exp Clin Transplant* 2017, 15:228–230. [PubMed: 26221721]
46. Kamar N, Izopet J, Dalton HR: Chronic hepatitis e virus infection and treatment. *J Clin Exp Hepatol* 2013, 3:134–140. [PubMed: 25755487]
47. Legrand-Abravanel F, Kamar N, Sandres-Saune K, Garrouste C, Dubois M, Mansuy JM, Muscari F, Sallusto F, Rostaing L, Izopet J: Characteristics of autochthonous hepatitis E virus infection in solid-organ transplant recipients in France. *J Infect Dis* 2010, 202:835–844. [PubMed: 20695798]
48. Yapa CM, Furlong C, Rosewell A, Ward KA, Adamson S, Shadbolt C, Kok J, Tracy SL, Bowden S, Smedley EJ, et al.: First reported outbreak of locally acquired hepatitis E virus infection in Australia. *Med J Aust* 2016, 204:274. [PubMed: 27078603]
49. Rivero-Juarez A, Frias M, Martinez-Peinado A, Rialde MA, Rodriguez-Cano D, Camacho A, Garcia-Bocanegra I, Cuenca-Lopez F, Gomez-Villamandos JC, Rivero A: Familial Hepatitis E Outbreak Linked to Wild Boar Meat Consumption. *Zoonoses Public Health* 2017, 64:561–565. [PubMed: 28067990]
50. Colson P, Borentain P, Queyriaux B, Kaba M, Moal V, Gallian P, Heyries L, Raoult D, Gerolami R: Pig liver sausage as a source of hepatitis E virus transmission to humans. *J Infect Dis* 2010,

202:825–834. [PubMed: 20695796] *This study reported cluster cases of hepatitis E virus infections due to consumption of contaminated pork products.

51. Tei S, Kitajima N, Ohara S, Inoue Y, Miki M, Yamatani T, Yamabe H, Mishiro S, Kinoshita Y: Consumption of uncooked deer meat as a risk factor for hepatitis E virus infection: an age- and sex-matched case-control study. *J Med Virol* 2004, 74:67–70. [PubMed: 15258970]
52. Takahashi K, Kitajima N, Abe N, Mishiro S: Complete or near-complete nucleotide sequences of hepatitis E virus genome recovered from a wild boar, a deer, and four patients who ate the deer. *Virology* 2004, 330:501–505. [PubMed: 15567444]
53. Choi JY, Lee JM, Jo YW, Min HJ, Kim HJ, Jung WT, Lee OJ, Yun H, Yoon YS: Genotype-4 hepatitis E in a human after ingesting roe deer meat in South Korea. *Clin Mol Hepatol* 2013, 19:309–314. [PubMed: 24133670]
54. Kukiella D, Rodriguez-Prieto V, Vicente J, Sanchez-Vizcaino JM: Constant Hepatitis E Virus (HEV) Circulation in Wild Boar and Red Deer in Spain: An Increasing Concern Source of HEV Zoonotic Transmission. *Transbound Emerg Dis* 2016, 63:e360–368. [PubMed: 25571944]
55. Geng Y, Zhang H, Huang W, T JH, Geng K, Li Z, Wang Y: Persistent hepatitis e virus genotype 4 infection in a child with acute lymphoblastic leukemia. *Hepat Mon* 2014, 14:e15618. [PubMed: 24596581]
56. Perumpail RB, Ahmed A, Higgins JP, So SK, Cochran JL, Drobeniuc J, Mixson-Hayden TR, Teo CG: Fatal Accelerated Cirrhosis after Imported HEV Genotype 4 Infection. *Emerg Infect Dis* 2015, 21:1679–1681. [PubMed: 26291424]
57. Emerson SU, Arankalle VA, Purcell RH: Thermal stability of hepatitis E virus. *J Infect Dis* 2005, 192:930–933. [PubMed: 16088844]
58. Barnaud E, Rogee S, Garry P, Rose N, Pavio N: Thermal inactivation of infectious hepatitis E virus in experimentally contaminated food. *Appl Environ Microbiol* 2012, 78:5153–5159. [PubMed: 22610436]
59. Jones TH, Muehlhauser V: Effect of handling and storage conditions and stabilizing agent on the recovery of viral RNA from oral fluid of pigs. *J Virol Methods* 2014, 198:26–31. [PubMed: 24384096]
60. Feagins AR, Opriessnig T, Guenette DK, Halbur PG, Meng XJ: Inactivation of infectious hepatitis E virus present in commercial pig livers sold in local grocery stores in the United States. *Int J Food Microbiol* 2008, 123:32–37. [PubMed: 18166239]
61. Meng XJ, Halbur PG, Shapiro MS, Govindarajan S, Bruna JD, Mushahwar IK, Purcell RH, Emerson SU: Genetic and experimental evidence for cross-species infection by swine hepatitis E virus. *J Virol* 1998, 72:9714–9721. [PubMed: 9811705] *This study reports for the first time experimental evidence of cross-species infection by genotype 3 hepatitis E virus.
62. Arankalle VA, Chobe LP, Chadha MS: Type-IV Indian swine HEV infects rhesus monkeys. *J Viral Hepat* 2006, 13:742–745. [PubMed: 17052273]
63. Sun ZF, Larsen CT, Huang FF, Billam P, Pierson FW, Toth TE, Meng XJ: Generation and infectivity titration of an infectious stock of avian hepatitis E virus (HEV) in chickens and cross-species infection of turkeys with avian HEV. *J Clin Microbiol* 2004, 42:2658–2662. [PubMed: 15184448]
64. Huang FF, Sun ZF, Emerson SU, Purcell RH, Shivaprasad HL, Pierson FW, Toth TE, Meng XJ: Determination and analysis of the complete genomic sequence of avian hepatitis E virus (avian HEV) and attempts to infect rhesus monkeys with avian HEV. *J Gen Virol* 2004, 85:1609–1618. [PubMed: 15166445]
65. Liu P, Bu QN, Wang L, Han J, Du RJ, Lei YX, Ouyang YQ, Li J, Zhu YH, Lu FM, et al.: Transmission of hepatitis E virus from rabbits to cynomolgus macaques. *Emerg Infect Dis* 2013, 19:559–565. [PubMed: 23628346]
66. Abravanel F, Lhomme S, El Costa H, Schvartz B, Peron JM, Kamar N, Izopet J: Rabbit Hepatitis E Virus Infections in Humans, France. *Emerg Infect Dis* 2017, 23:1191–1193. [PubMed: 28628452]
67. Kaiser M, Delaune D, Chazouilleres O, Blumel J, Roque-Afonso AM, Baylis SA: A World Health Organization Human Hepatitis E Virus Reference Strain Related to Similar Strains Isolated from Rabbits. *Genome Announc* 2018, 6.

68. Meng XJ, Wiseman B, Elvinger F, Guenette DK, Toth TE, Engle RE, Emerson SU, Purcell RH: Prevalence of antibodies to hepatitis E virus in veterinarians working with swine and in normal blood donors in the United States and other countries. *J Clin Microbiol* 2002, 40:117–122. [PubMed: 11773103]
69. Kantala T, Kinnunen PM, Oristo S, Jokelainen P, Vapalahti O, Maunula L: Hepatitis E Virus Antibodies in Finnish Veterinarians. *Zoonoses Public Health* 2017, 64:232–238. [PubMed: 27621202]
70. Kang YH, Cong W, Zhang XY, Wang CF, Shan XF, Qian AD: Hepatitis E virus seroprevalence among farmers, veterinarians and control subjects in Jilin province, Shandong province and Inner Mongolia Autonomous Region, China. *J Med Virol* 2017, 89:872–877. [PubMed: 27664799]
71. Baumann-Popczyk A, Popczyk B, Golab E, Rozej-Bielicka W, Sadkowska-Todys M: A cross-sectional study among Polish hunters: seroprevalence of hepatitis E and the analysis of factors contributing to HEV infections. *Med Microbiol Immunol* 2017, 206:367–378. [PubMed: 28776194]
72. Mughini-Gras L, Angeloni G, Salata C, Vonesch N, D'Amico W, Campagna G, Natale A, Zuliani F, Ceglie L, Monne I, et al.: Hepatitis E virus infection in North Italy: high seroprevalence in swine herds and increased risk for swine workers. *Epidemiol Infect* 2017, 145:3375–3384. [PubMed: 29145911]
73. Karetnyi YV, Gilchrist MJ, Naides SJ: Hepatitis E virus infection prevalence among selected populations in Iowa. *J Clin Virol* 1999, 14:51–55. [PubMed: 10548130]
74. van Gageldonk-Lafeber AB, van der Hoek W, Borlee F, Heederik DJ, Mooi SH, Maassen CB, Yzermans CJ, Rockx B, Smit LA, Reimerink JH: Hepatitis E virus seroprevalence among the general population in a livestock-dense area in the Netherlands: a cross-sectional population-based serological survey. *BMC Infect Dis* 2017, 17:21. [PubMed: 28056844]
75. Satake M, Matsubayashi K, Hoshi Y, Taira R, Furui Y, Kokudo N, Akamatsu N, Yoshizumi T, Ohkohchi N, Okamoto H, et al.: Unique clinical courses of transfusion-transmitted hepatitis E in patients with immunosuppression. *Transfusion* 2017, 57:280–288. [PubMed: 28144952]
76. Hewitt PE, Ijaz S, Brailsford SR, Brett R, Dicks S, Haywood B, Kennedy IT, Kitchen A, Patel P, Poh J, et al.: Hepatitis E virus in blood components: a prevalence and transmission study in southeast England. *Lancet* 2014, 384:1766–1773. [PubMed: 25078306]
77. Colson P, Coze C, Gallian P, Henry M, De Micco P, Tamalet C: Transfusion-associated hepatitis E, France. *Emerg Infect Dis* 2007, 13:648–649. [PubMed: 17561564]
78. Huzly D, Umhau M, Bettinger D, Cathomen T, Emmerich F, Hasselblatt P, Hengel H, Herzog R, Kappert O, Maassen S, et al.: Transfusion-transmitted hepatitis E in Germany, 2013. *Euro Surveill* 2014, 19.
79. Hoad VC, Gibbs T, Ravikumara M, Nash M, Levy A, Tracy SL, Mews C, Perkowska-Guse Z, Faddy HM, Bowden S: First confirmed case of transfusion-transmitted hepatitis E in Australia. *Med J Aust* 2017, 206:289–290. [PubMed: 28403756]
80. Andonov A, Rock G, Lin L, Borlang J, Hooper J, Grudeski E, Wu J, Members of the Canadian Apheresis G: **Serological and molecular evidence of a plausible transmission of hepatitis E virus through pooled plasma.** *Vox Sang* 2014, 107:213–219. [PubMed: 24830322]
81. Takeda H, Matsubayashi K, Sakata H, Sato S, Kato T, Hino S, Tadokoro K, Ikeda H: A nationwide survey for prevalence of hepatitis E virus antibody in qualified blood donors in Japan. *Vox Sang* 2010, 99:307–313. [PubMed: 20576022]
82. Fearon MA, O'Brien SF, Delage G, Scalia V, Bernier F, Bigham M, Weger S, Prabhu S, Andonov A: Hepatitis E in Canadian blood donors. *Transfusion* 2017, 57:1420–1425. [PubMed: 28394029]
83. Juhl D, Baylis SA, Blumel J, Gorg S, Hennig H: Seroprevalence and incidence of hepatitis E virus infection in German blood donors. *Transfusion* 2014, 54:49–56. [PubMed: 23441647]
84. Mansuy JM, Gallian P, Dimeglio C, Saune K, Arnaud C, Pelletier B, Morel P, Legrand D, Tiberghien P, Izopet J: A nationwide survey of hepatitis E viral infection in French blood donors. *Hepatology* 2016, 63:1145–1154. [PubMed: 27008201]
85. Xu C, Wang RY, Schechterly CA, Ge S, Shih JW, Xia NS, Luban NL, Alter HJ: An assessment of hepatitis E virus (HEV) in US blood donors and recipients: no detectable HEV RNA in 1939

- donors tested and no evidence for HEV transmission to 362 prospectively followed recipients. *Transfusion* 2013, 53:2505–2511. [PubMed: 23829163]
86. Roth NJ, Schafer W, Alexander R, Elliott K, Elliott-Browne W, Knowles J, Wenzel JJ, Simon TL: Low hepatitis E virus RNA prevalence in a large-scale survey of United States source plasma donors. *Transfusion* 2017, 57:2958–2964. [PubMed: 28833188]
 87. Izopet J, Lhomme S, Chapuy-Regaud S, Mansuy JM, Kamar N, Abravanel F: HEV and transfusion-recipient risk. *Transfus Clin Biol* 2017, 24:176–181. [PubMed: 28690036]
 88. Kraef C, Schlein C, Hiller J, Westholter D, Denzer U, Horvatits T, Peine S, Lohse AW, Lutgehetmann M, Polywka S, et al.: Course of HEV viremia and anti-HEV IgM/IgG response in asymptomatic blood donors. *J Clin Virol* 2018, 105:26–30. [PubMed: 29860111] * This study reports the course of HEV viremia and epitope specific anti-HEV IgM/IgG response in asymptomatic blood donors in Germany.
 89. Tedder RS, Tettmar KI, Brailsford SR, Said B, Ushiro-Lumb I, Kitchen A, Morgan D, Lattimore S, Tossell J, Ijaz S, et al.: Virology, serology, and demography of hepatitis E viremic blood donors in South East England. *Transfusion* 2016, 56:1529–1536. [PubMed: 26841005]
 90. Tedder RS, Ijaz S, Kitchen A, Ushiro-Lumb I, Tettmar KI, Hewitt P, Andrews N: Hepatitis E risks: pigs or blood—that is the question. *Transfusion* 2017, 57:267–272. [PubMed: 28194857] * This study determined the minimum infective dose for HEV transmission by transfusion, and estimate relative risk between transfusion and food-borne transmission of HEV.
 91. Loyrion E, Trouve-Buisson T, Pouzol P, Larrat S, Decaens T, Payen JF: Hepatitis E Virus Infection after Platelet Transfusion in an Immunocompetent Trauma Patient. *Emerg Infect Dis* 2017, 23:146–147. [PubMed: 27983485]
 92. Domanovic D, Tedder R, Blumel J, Zaaier H, Gallian P, Niederhauser C, Saulea Oliveras S, O’Riordan J, Boland F, Harritshoj L, et al.: Hepatitis E and blood donation safety in selected European countries: a shift to screening? *Euro Surveill* 2017, 22.* This review summarizes epidemiology of HEV among the European blood donors and proposes strategies to prevent transfusion-transmitted HEV.
 93. Fenaux H, Chassaing M, Berger S, Jeulin H, Gentilhomme A, Bensenane M, Bronowicki JP, Gantzer C, Bertrand I, Schvoerer E: Molecular features of Hepatitis E Virus circulation in environmental and human samples. *J Clin Virol* 2018, 103:63–70. [PubMed: 29656087]
 94. Parashar D, Khalkar P, Arankalle VA: Survival of hepatitis A and E viruses in soil samples. *Clin Microbiol Infect* 2011, 17:E1–4.
 95. Kasorndorkbua C, Opriessnig T, Huang FF, Guenette DK, Thomas PJ, Meng XJ, Halbur PG: Infectious swine hepatitis E virus is present in pig manure storage facilities on United States farms, but evidence of water contamination is lacking. *Appl Environ Microbiol* 2005, 71:7831–7837. [PubMed: 16332757]
 96. dos Santos DR, de Paula VS, de Oliveira JM, Marchevsky RS, Pinto MA: Hepatitis E virus in swine and effluent samples from slaughterhouses in Brazil. *Vet Microbiol* 2011, 149:236–241. [PubMed: 21129863]
 97. Garcia M, Fernandez-Barredo S, Perez-Gracia MT: Detection of hepatitis E virus (HEV) through the different stages of pig manure composting plants. *Microb Biotechnol* 2014, 7:26–31. [PubMed: 24206540]
 98. Mesquita JR, Oliveira D, Rivadulla E, Abreu-Silva J, Varela MF, Romalde JL, Nascimento MS: Hepatitis E virus genotype 3 in mussels (*Mytilus galloprovincialis*), Spain. *Food Microbiol* 2016, 58:13–15. [PubMed: 27217353]
 99. Brassard J, Gagne MJ, Genereux M, Cote C: Detection of human food-borne and zoonotic viruses on irrigated, field-grown strawberries. *Appl Environ Microbiol* 2012, 78:3763–3766. [PubMed: 22427499]
 100. van Tong H, Hoan NX, Wang B, Wedemeyer H, Bock CT, Velavan TP: Hepatitis E Virus Mutations: Functional and Clinical Relevance. *EBioMedicine* 2016, 11:31–42. [PubMed: 27528267] * A detailed review on recent advances on the role of mutations in HEV heterogeneity and its variants on viral morphogenesis, pathogenesis, clinical outcome and antiviral resistance.
 101. Lhomme S, Garrouste C, Kamar N, Saune K, Abravanel F, Mansuy JM, Dubois M, Rostaing L, Izopet J: Influence of polyproline region and macro domain genetic heterogeneity on HEV

- persistence in immunocompromised patients. *J Infect Dis* 2014, 209:300–303. [PubMed: 23964111]
102. Pudupakam RS, Huang YW, Opriessnig T, Halbur PG, Pierson FW, Meng XJ: Deletions of the hypervariable region (HVR) in open reading frame 1 of hepatitis E virus do not abolish virus infectivity: evidence for attenuation of HVR deletion mutants in vivo. *J Virol* 2009, 83:384–395. [PubMed: 18945785]
103. Pudupakam RS, Kenney SP, Cordoba L, Huang YW, Dryman BA, Leroith T, Pierson FW, Meng XJ: Mutational analysis of the hypervariable region of hepatitis e virus reveals its involvement in the efficiency of viral RNA replication. *J Virol* 2011, 85:10031–10040. [PubMed: 21775444]
104. Cordoba L, Feagins AR, Opriessnig T, Cossaboom CM, Dryman BA, Huang YW, Meng XJ: Rescue of a genotype 4 human hepatitis E virus from cloned cDNA and characterization of intergenotypic chimeric viruses in cultured human liver cells and in pigs. *J Gen Virol* 2012, 93:2183–2194. [PubMed: 22837416]
105. Feagins AR, Cordoba L, Sanford BJ, Dryman BA, Huang YW, LeRoith T, Emerson SU, Meng XJ: Intergenotypic chimeric hepatitis E viruses (HEVs) with the genotype 4 human HEV capsid gene in the backbone of genotype 3 swine HEV are infectious in pigs. *Virus Res* 2011, 156:141–146. [PubMed: 21195119]
106. Shukla P, Nguyen HT, Faulk K, Mather K, Torian U, Engle RE, Emerson SU: Adaptation of a genotype 3 hepatitis E virus to efficient growth in cell culture depends on an inserted human gene segment acquired by recombination. *J Virol* 2012, 86:5697–5707. [PubMed: 22398290] *This study reports that the acquisition of a human ribosomal protein sequence S17 in the hypervariable region of HEV genome confers adaptation of HEV growth in cell cultures.
107. Debing Y, Ramiere C, Dallmeier K, Piorkowski G, Traub MA, Lebosse F, Scholtes C, Roche M, Legras-Lachuer C, de Lamballerie X, et al.: Hepatitis E virus mutations associated with ribavirin treatment failure result in altered viral fitness and ribavirin sensitivity. *J Hepatol* 2016, 65:499–508. [PubMed: 27174035]

Highlights

- Emergence of novel animal strains of HEV transmissible to human.
- Genotype 1 HEV outbreaks among displaced population with poor sanitation conditions.
- Emergence of autochthonous cases of hepatitis E caused by zoonotic genotypes 3 and 4 HEVs.
- Point mutations and recombination attribute to HEV heterogeneity.

Figure 1. Worldwide prevalence of human hepatitis E cases associated with genotypes 1, 2, 3, 4, and 7 HEV infections.

Genotypes (Gt) 1 and 2 HEV mainly infect humans and are transmitted via fecal-oral route through contaminated water or water supplies. Gt3 and Gt4 HEVs are zoonotic in nature, and are transmitted to human through direct contact with infected animals or through eating undercooked or raw animal meat products. Pigs, wild boar and deer meats are most commonly associated with Gt3 and Gt4 transmission to humans. Gt3 rabbit HEV infection of human is reported from France, and Gt7 camel HEV infection of human is reported from Middle East.