Cochrane
Library

Cochrane Database of Systematic Reviews

Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery (Review)

Grocott MPW, Dushianthan A, Hamilton MA, Mythen MG, Harrison D, Rowan K, Optimisation Systematic Review Steering Group

Grocott MPW, Dushianthan A, Hamilton MA, Mythen MG, Harrison D, Rowan K, Optimisation Systematic Review Steering Group.

Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery.

Cochrane Database of Systematic Reviews 2012, Issue 11. Art. No.: CD004082.

DOI: 10.1002/14651858.CD004082.pub5.

www.cochranelibrary.com

TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
SUMMARY OF FINDINGS FOR THE MAIN COMPARISON	3
BACKGROUND	5
OBJECTIVES	5
METHODS	5
RESULTS	8
Figure 1.	9
Figure 2.	10
Figure 3.	11
Figure 4.	13
Figure 5.	14
Figure 6.	15
Figure 7.	16
DISCUSSION	20
AUTHORS' CONCLUSIONS	22
ACKNOWLEDGEMENTS	22
REFERENCES	23
CHARACTERISTICS OF STUDIES	29
DATA AND ANALYSES	65
Analysis 1.1. Comparison 1 Mortality, Outcome 1 All studies (longest follow-up).	68
Analysis 1.2. Comparison 1 Mortality, Outcome 2 All studies (hospital or 28 day).	69
Analysis 1.3. Comparison 1 Mortality, Outcome 3 Participant numbers.	71
Analysis 1.4. Comparison 1 Mortality, Outcome 4 Time intervention started.	73
Analysis 1.5. Comparison 1 Mortality, Outcome 5 Type of intervention.	75
Analysis 1.6. Comparison 1 Mortality, Outcome 6 Goals of intervention.	77
Analysis 1.7. Comparison 1 Mortality, Outcome 7 Mode of surgery.	79
Analysis 1.8. Comparison 1 Mortality, Outcome 8 Type of surgery.	81
Analysis 2.1. Comparison 2 Complications, Outcome 1 Renal impairment.	83
Analysis 2.2. Comparison 2 Complications, Outcome 2 Arrhythmia.	84
Analysis 2.3. Comparison 2 Complications, Outcome 3 Infection: numbers.	85
Analysis 2.4. Comparison 2 Complications, Outcome 4 Infections: types.	86
Analysis 2.5. Comparison 2 Complications, Outcome 5 Respiratory failure / ARDS.	88
Analysis 2.6. Comparison 2 Complications, Outcome 6 Myocardial infarction.	89
Analysis 2.7. Comparison 2 Complications, Outcome 7 Congestive heart failure / pulmonary oedema.	90
Analysis 2.8. Comparison 2 Complications, Outcome 8 Venous thrombosis.	91
Analysis 2.9. Comparison 2 Complications, Outcome 9 Number of participants with complications.	92
Analysis 3.1. Comparison 3 Resource utilization, Outcome 1 Length of hospital stay.	93
Analysis 3.2. Comparison 3 Resource utilization, Outcome 2 Length of critical care stay.	94
Analysis 4.1. Comparison 4 Sensitivity analysis: excluding studies with active controls, Outcome 1 Mortality (Longest follow-up).	95
Analysis 4.2. Comparison 4 Sensitivity analysis: excluding studies with active controls, Outcome 2 Mortality (Hospital or 28 days).	97
Analysis 4.3. Comparison 4 Sensitivity analysis: excluding studies with active controls, Outcome 3 Renal Impairment.	98
Analysis 4.4. Comparison 4 Sensitivity analysis: excluding studies with active controls, Outcome 4 Arrhythmia.	100
Analysis 4.5. Comparison 4 Sensitivity analysis: excluding studies with active controls, Outcome 5 Infection (Number of patients with infections).	101
Analysis 4.6. Comparison 4 Sensitivity analysis: excluding studies with active controls, Outcome 6 Respiratory Failure/ ARDS.	102

Analysis 4.7. Comparison 4 Sensitivity analysis: excluding studies with active controls, Outcome 7 Myocardial Infarction.	103
Analysis 4.8. Comparison 4 Sensitivity analysis: excluding studies with active controls, Outcome 8 Congestive Heart Failure/ Pulmonary oedema.	104
Analysis 4.9. Comparison 4 Sensitivity analysis: excluding studies with active controls, Outcome 9 Venous Thrombosis.	105
Analysis 4.10. Comparison 4 Sensitivity analysis: excluding studies with active controls, Outcome 10 Number of patients with complications.	106
Analysis 4.11. Comparison 4 Sensitivity analysis: excluding studies with active controls, Outcome 11 Length of Hospital Stay [Days].	107
Analysis 4.12. Comparison 4 Sensitivity analysis: excluding studies with active controls, Outcome 12 Length of Critical Care Stay [Days].	108
Analysis 5.1. Comparison 5 Sensitivity analysis: excluding poorly-controlled studies, Outcome 1 Mortality (Longest follow-up).	109
Analysis 5.2. Comparison 5 Sensitivity analysis: excluding poorly-controlled studies, Outcome 2 Mortality (Hospital or 28 day).	111
Analysis 5.3. Comparison 5 Sensitivity analysis: excluding poorly-controlled studies, Outcome 3 Renal Impairment.	112
Analysis 5.4. Comparison 5 Sensitivity analysis: excluding poorly-controlled studies, Outcome 4 Arrhythmia.	113
Analysis 5.5. Comparison 5 Sensitivity analysis: excluding poorly-controlled studies, Outcome 5 Infection (Number of patients with infections).	114
Analysis 5.6. Comparison 5 Sensitivity analysis: excluding poorly-controlled studies, Outcome 6 Respiratory Failure/ ARDS.	115
Analysis 5.7. Comparison 5 Sensitivity analysis: excluding poorly-controlled studies, Outcome 7 Myocardial Infarction.	116
Analysis 5.8. Comparison 5 Sensitivity analysis: excluding poorly-controlled studies, Outcome 8 Congestive Heart Failure/ Pulmonary oedema.	117
Analysis 5.9. Comparison 5 Sensitivity analysis: excluding poorly-controlled studies, Outcome 9 Venous Thrombosis.	118
Analysis 5.10. Comparison 5 Sensitivity analysis: excluding poorly-controlled studies, Outcome 10 Number of patients with complications.	119
Analysis 5.11. Comparison 5 Sensitivity analysis: excluding poorly-controlled studies, Outcome 11 Length of Hospital Stay [Days].	120
Analysis 5.12. Comparison 5 Sensitivity analysis: excluding poorly-controlled studies, Outcome 12 Length of Critical Care Stay [Days].	121
Analysis 6.1. Comparison 6 Sensitivity analysis: Sandham 2003, Outcome 1 Mortality (Longest Follow-up)- excluding patients lost follow-up.	122
Analysis 6.2. Comparison 6 Sensitivity analysis: Sandham 2003, Outcome 2 Mortality (Longest Follow-up)- if all patients lost follow-up have died.	124
ADDITIONAL TABLES	125
APPENDICES	132
WHAT'S NEW	138
HISTORY	138
CONTRIBUTIONS OF AUTHORS	139
DECLARATIONS OF INTEREST	140
SOURCES OF SUPPORT	140
DIFFERENCES BETWEEN PROTOCOL AND REVIEW	140
NOTES	141
INDEX TERMS	141

[Intervention Review]

Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Michael PW Grocott¹, Ahilanandan Dushianthan², Mark A Hamilton³, Michael G Mythen⁴, David Harrison⁵, Kathy Rowan⁵, Optimisation Systematic Review Steering Group⁶

¹University College London Centre for Altitude Space and Extreme Environment (CASE) Medicine, UCLH NIHR Biomedical Research Centre, Institute of Sport and Exercise Health, London, UK. ²Integrative Physiology and Critical Illness Group, University of Southampton, Southampton, UK. ³General Intensive Care Unit, St. George's Hospital, London, UK. ⁴Department Anaesthesia and Critical Care, University College London, London, UK. ⁵Intensive Care National Audit & Research Centre, London, UK. ⁶London, UK

Contact address: Ahilanandan Dushianthan, Integrative Physiology and Critical Illness Group, University of Southampton, Southampton, UK. adushianthan@gmail.com. ahilanadan.dushianthan@uhs.nhs.uk.

Editorial group: Cochrane Anaesthesia, Critical and Emergency Care Group.

Publication status and date: Edited (no change to conclusions), published in Issue 10, 2016.

Review content assessed as up-to-date: 1 March 2012.

Citation: Grocott MPW, Dushianthan A, Hamilton MA, Mythen MG, Harrison D, Rowan K, Optimisation Systematic Review Steering Group. Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery. *Cochrane Database of Systematic Reviews* 2012, Issue 11. Art. No.: CD004082. DOI: 10.1002/14651858.CD004082.pub5.

Copyright © 2016 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

Studies have suggested that increasing whole body blood flow and oxygen delivery around the time of surgery reduces mortality, morbidity and the expense of major operations.

Objectives

To describe the effects of increasing perioperative blood flow using fluids with or without inotropes or vasoactive drugs. Outcomes were mortality, morbidity, resource utilization and health status.

Search methods

We searched CENTRAL (*The Cochrane Library* 2012, Issue 1), MEDLINE (1966 to March 2012) and EMBASE (1982 to March 2012). We manually searched the proceedings of major conferences and personal reference databases up to December 2011. We contacted experts in the field and pharmaceutical companies for published and unpublished data.

Selection criteria

We included randomized controlled trials with or without blinding. We included studies involving adult patients (aged 16 years or older) undergoing surgery (patients having a procedure in an operating room). The intervention met the following criteria. 'Perioperative' was defined as starting up to 24 hours before surgery and stopping up to six hours after surgery. 'Targeted to increase global blood flow' was defined by explicit measured goals that were greater than in controls, specifically one or more of cardiac index, oxygen delivery, oxygen consumption, stroke volume (and the respective derived indices), mixed venous oxygen saturation (SVO₂), oxygen extraction ratio (O₂ER) or lactate.

Data collection and analysis

Two authors independently extracted the data. We contacted study authors for additional data. We used Review Manager software.

Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery (Review)

Copyright © 2016 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

Main results

We included 31 studies of 5292 participants. There was no difference in mortality: 282/2615 (10.8%) died in the control group and 238/2677 (8.9%) in the treatment group, RR of 0.89 (95% CI 0.76 to 1.05, $P = 0.18$). However, the results were sensitive to analytical methods and the intervention was better than control when inverse variance or Mantel-Haenszel random-effects models were used, RR of 0.72 (95% CI 0.55 to 0.95, $P = 0.02$). The results were also sensitive to withdrawal of studies with methodological limitations. The rates of three morbidities were reduced by increasing global blood flow: renal failure, RR of 0.71 (95% CI 0.57 to 0.90); respiratory failure, RR of 0.51 (95% CI 0.28 to 0.93); and wound infections, RR of 0.65 (95% CI 0.51 to 0.84). There were no differences in the rates of nine other morbidities: arrhythmia, pneumonia, sepsis, abdominal infection, urinary tract infection, myocardial infarction, congestive cardiac failure or pulmonary oedema, or venous thrombosis. The number of patients with complications was reduced by the intervention, RR of 0.68 (95% CI 0.58 to 0.80). Hospital length of stay was reduced in the treatment group by a mean of 1.16 days (95% CI 0.43 to 1.89, $P = 0.002$). There was no difference in critical care length of stay. There were insufficient data to comment on quality of life and cost effectiveness.

Authors' conclusions

It remains uncertain whether increasing blood flow using fluids, with or without inotropes or vasoactive drugs, reduces mortality in adults undergoing surgery. The primary analysis in this review (mortality at longest follow-up) showed no difference between the intervention and control, but this result was sensitive to the method of analysis, the withdrawal of studies with methodological limitations, and is dominated by a single large RCT. Overall, for every 100 patients in whom blood flow is increased perioperatively to defined goals, one can expect 13 in 100 patients (from 40/100 to 27/100) to avoid a complication, 2/100 to avoid renal impairment (from 8/100 to 6/100), 5/100 to avoid respiratory failure (from 10/100 to 5/100), and 4/100 to avoid postoperative wound infection (from 10/100 to 6/100). On average, patients receiving the intervention stay in hospital one day less. It is unlikely that the intervention causes harm. The balance of current evidence does not support widespread implementation of this approach to reduce mortality but does suggest that complications and duration of hospital stay are reduced.

PLAIN LANGUAGE SUMMARY

Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Death and serious complications commonly occur following major surgery and are a significant public health problem. These outcomes might be prevented by using fluids and drugs to maintain the supply of oxygen and other nutrients to vital organs. Global blood flow, adjusted to maintain specific targets, might serve as a proxy in determining whether administered fluid and drugs maintain critical nutrient supply. In this Cochrane review of 31 studies conducted in 5292 patients undergoing major surgery, the use of fluids, with or without additional drugs, to achieve defined targets associated with increased total blood flow did not reduce mortality. There was a reduction in the number of patients with complications and the length of time patients stayed in hospital (by 1.2 days). However, the quality of the studies in this area was mediocre.

SUMMARY OF FINDINGS FOR THE MAIN COMPARISON [\[Explanation\]](#)

Protocol to increase global blood flow compared to control for surgical patients						
Patient or population: Surgical patients Settings: Hospital Intervention: Protocol to increase global blood flow Comparison: Control						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Control	Protocol to increase global blood flow				
Mortality (Longest follow-up)	11 per 100	10 per 100 (8 to 11)	RR 0.89 (0.76 to 1.05)	5292 (31 studies)	⊕⊕○○ low ^{1,2}	P=0.18
Mortality (Hospital or 28-day)	7 per 100	6 per 100 (5 to 7)	RR 0.81 (0.65 to 1.00)	5292 (31 studies)	⊕⊕○○ low ^{1,2}	P=0.06
Number of patients with complications	40 per 100	27 per 100 (23 to 32)	RR 0.68 (0.58 to 0.80)	1841 (17 studies)	⊕⊕○○ low ^{1,2}	P<0.00001
Length of hospital stay		The mean length of hospital stay in the intervention groups was 1.16 lower (1.89 to 0.43 lower)		4729 (27 studies)	⊕⊕○○ low ^{1,2}	P=0.002
*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI). CI: Confidence interval; RR: Risk ratio						

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

¹ Majority of studies are unblinded due to the nature of the intervention and hence we have suggested “unclear risk for most of the studies”.

² Most studies had small number of patients.

BACKGROUND

Description of the condition

It has been known for many years that patients undergoing surgery are more likely to have serious complications or die if they have limited physiological reserve (Boyd 1959; Clowes 1960). Post hoc analysis of patients undergoing major surgery revealed that survivors had a higher cardiac index and lower systemic vascular resistance than those who died (Shoemaker 1972; Shoemaker 1973). Commonly monitored vital signs (heart rate, arterial blood pressure, central venous pressure, temperature, haemoglobin concentration) were found to be poor predictors of mortality when compared with the flow related variables cardiac output and total body oxygen delivery (DO₂) (Shoemaker 1979; Shoemaker 1993). In particular, survivors of major surgical procedures were found to have higher values for cardiac output or DO₂. More recent studies have shown mixed results for the impact of oxygen transport on postoperative morbidity and mortality (Kusano 1997; Peerless 1998; Polonen 1997).

Description of the intervention

New therapeutic options and monitoring techniques that became available in the 1970s, particularly the introduction of the pulmonary artery flow directed catheter (PAC) (Ganz 1971; Swan 1970), opened up the possibility of measuring and then manipulating an individual's cardiovascular system. It was hypothesized that targeting goals for cardiac output and DO₂ in all patients to the values manifested by the survivors of surgery would improve outcome (Bland 1978).

How the intervention might work

An important principle of this intervention is that the perioperative manipulation to augment cardiac output and DO₂ would lead to an improved tissue perfusion and oxygenation. This physiological improvement would lead to better survival and fewer postoperative complications in patients undergoing major surgery.

Why it is important to do this review

It is almost 30 years since the initial uncontrolled data were presented suggesting that perioperative manipulation of flow related cardiovascular variables might improve outcomes in higher risk surgical patients (Shoemaker 1982). Since then, a number of randomized trials have been undertaken in patients in the perioperative period which have investigated this issue. However, these trials differ in:

- the case mix of the patients recruited (different operation severities, comorbidities and, therefore, expected mortalities);
- the techniques used to measure cardiac output (pulmonary artery catheter (PAC) thermodilution, Doppler velocimetry);
- the specific goals targeted (cardiac output, DO₂, maximum stroke volume);
- the techniques used to achieve the goals (fluids, fluids plus inotropes or vasoactive drugs);
- the management of the control arm.

In addition, some of the studies were not blinded and many had small sample sizes leading to limited statistical power. Despite this, a number of non-systematic reviews have attempted to combine studies in order to draw general conclusions from the studies (Boyd 1996; Boyd 1999; Forst 1997; Ivanov 1997; Leibowitz 1997). However, these reviews have identified varying numbers of trials and have not been undertaken systematically, using scientifically rigorous techniques for literature searching or for abstraction and analyses of data. Three previous systematic reviews have addressed this question (Heyland 1996; Kern 2002; Poeze 2005) and reported improved outcomes. They do not include recently published studies and did not focus exclusively on perioperative data. Among recent systematic reviews and meta-analyses, one study included patients with trauma and sepsis (Hamilton 2011) while other studies analysed renal function (Brienza 2009) and gastrointestinal complications (Giglio 2009) as primary outcomes. The time is now ripe for a systematic review of the literature to address the important question: does perioperative administration of fluids, with or without vasoactive drugs, targeted to increase global blood flow in adults undergoing surgery reduce mortality, morbidity and resource utilization?

OBJECTIVES

To describe the effects of perioperative (24 hours before surgery up to six hours after surgery) administration of fluids, with or without vasoactive drugs, that were targeted to increase global blood flow (relative to control) as defined by explicit measured goals on outcomes following surgery (mortality, morbidity, resource utilization and health status).

METHODS

Criteria for considering studies for this review

Types of studies

We included randomized controlled trials (RCTs), with or without blinding, that were available as full published papers. We applied no language restrictions.

Types of participants

We included adults (aged 16 years or older) undergoing surgery in an operating theatre.

Types of interventions

Perioperative administration (initiated within 24 hours before surgery and lasting up to six hours after surgery) of fluids, with or without inotropes or vasoactive drugs, to increase blood flow (relative to control) against explicit measured goals: cardiac output (CO), cardiac index (CI), oxygen delivery (DO₂) or oxygen delivery index (DO₂I), oxygen consumption or oxygen consumption index (VO₂), stroke volume (SV) or stroke volume index, mixed venous oxygen saturation (SVO₂), oxygen extraction ratio (O₂ER) and lactate.

Types of outcome measures

Primary outcomes

1. Mortality (at longest available follow-up)

Secondary outcomes

1. Mortality: all reported time frames e.g. hospital or 28 day, six months.
2. Morbidity:
 - 2.1. rates of overall complications;
 - 2.2. rates of renal impairment, arrhythmia, respiratory failure or acute respiratory distress syndrome (ARDS), infection, myocardial infarction, congestive heart failure or pulmonary oedema, and venous thrombosis.
3. Resource utilization: length of intensive care (ICU) stay, length of hospital stay, cost.
4. Health status: e.g., six month functional health status, quality of life scores.

Search methods for identification of studies

Electronic searches

We searched the Cochrane Central Register of Controlled Trials (CENTRAL) in *The Cochrane Library* (2012, Issue 1), see [Appendix 1](#); MEDLINE via OvidSP (1966 to March 2012), see [Appendix 2](#); and EMBASE via OvidSP (1982 to March 2012), see [Appendix 3](#). For searching in MEDLINE we combined our topic-specific key words with the Cochrane highly sensitive search

strategy for identifying RCTs ([Higgins 2011](#)). We modified this filter for use in EMBASE. We used specific keywords to identify potential studies ([Appendix 4](#)).

Searching other resources

We searched the proceedings of the following major, relevant European and North American conferences from the year 2011 backwards, without finding eligible studies.

- American College of Surgeons (2011 to 1996).
- American Society of Anesthesiologists (2011 to 1995).
- American Thoracic Society (2011 to 1997*) (* = not available for searching prior to 1997).
- Association of Surgeons of Great Britain and Ireland (2011 to 1996).
- European Society of Anaesthesiologists (2011 to 1995).
- European Society of Intensive Care Medicine (2011 to 1983).
- International Anesthesia Research Society (2011 to 1994).
- Society of Critical Care Medicine (2011 to 1986).

We checked the reference lists of potentially eligible studies and previously published systematic reviews. We also searched the personal reference databases of the authors and the Steering Group for this review. We contacted experts in the field and relevant pharmaceutical companies and asked for published and unpublished reports.

Data collection and analysis

Selection of studies

Two independent authors (MG and MH for 2001 to 2006, AD and AV for 2006 to 2012) identified titles and abstracts of potentially eligible studies. We resolved any disagreement by discussion. We obtained the full texts of potentially eligible studies. We abstracted the study characteristics including: study design; patient population; interventions; and outcomes (see [Appendix 5](#); [Appendix 6](#); [Table 1](#)). Review authors were not involved in the selection of studies they had authored.

Data extraction and management

Two authors (MG and MH for 2001 to 2006, AD and AV for 2006 to 2012) independently extracted data. We achieved consensus by resolving any disparity in data collection by discussion. In the absence of appropriate published data, we made at least three attempts to contact authors of eligible studies to obtain any required data. Some studies were conducted by the authors of this review (MGM). They were not involved in the data extraction or risk of bias assessment.

Assessment of risk of bias in included studies

We performed the risk of bias assessment according to the Cochrane risk of bias tool (Higgins 2011). From this tool, we used the following seven domains to assess the methodological quality of included studies. This is summated in a graph and a summary table.

1. *Random sequence generation (selection bias)*: describe the method used to generate the allocation sequence in sufficient detail to allow an assessment of whether it should produce comparable groups.
2. *Allocation concealment (selection bias)*: describe the method used to conceal the allocation sequence in sufficient detail to determine whether intervention allocations could have been foreseen in advance of, or during, enrolment.
3. *Blinding of participant and personnel (performance bias)*: describe all measures used, if any, to blind study participants and personnel from knowledge of which intervention a participant received. Provide any information relating to whether the intended blinding was effective.
4. *Blinding of outcome assessment (detection bias)*: describe all measures used, if any, to blind outcome assessors from knowledge of which intervention a participant received. Provide any information relating to whether the intended blindness was effective.
5. *Incomplete outcome data (attrition bias)*: describe the completeness of the outcome data for each main outcome, including attrition and exclusion from the analysis. State whether attrition and exclusions were reported, the number in each intervention group (compared with total randomized participants), reasons for attrition and exclusions, and any re-inclusions in analysis performed by the review authors.
6. *Selective reporting (reporting bias)*: state how the possibility of selective outcome reporting was examined by the review authors, and what was found.
7. *Other sources of bias*: state any important concerns about bias not addressed in the other domains in the tool.

Measures of treatment effect

We based analyses of outcomes on intention-to-treat. We calculated a weighted treatment effect across all RCTs using Review Manager (RevMan 5.1). We expressed measures of treatment effect, such as mortality and complications, as relative risks (RR) and 95% confidence intervals (CI). We used mean differences (standard deviation) for continuous variables such as length of hospital or ICU stay. We explored the robustness of these estimates by comparing both fixed-effect and random-effects models for the primary outcome.

Unit of analysis issues

We included studies with different treatment groups, interventions and outcomes. Consequently, we performed subgroup analyses of these differences. Many studies reported the number of complications, arrhythmias and infections as total numbers, leaving un-

clear what the denominators were for these episodes. We have not analysed variables for which the denominator was unknown.

Dealing with missing data

We contacted the authors of the studies for further information and the analysis was performed with the best available information when there was no response.

Assessment of heterogeneity

We assessed inconsistencies and variability in the outcomes among the studies by the I^2 statistic. Variations of > 40% in the outcomes may not be explained by sampling variation. We assumed substantial heterogeneity when the I^2 statistic exceeded 40% (Higgins 2011).

Assessment of reporting biases

We assessed graphical evidence of reporting biases using contour enhanced funnel plots with a subsequent Harbord or Egger's test (Egger 1997; Harbord 2006).

Data synthesis

We performed statistical analysis using Review Manager 5.1 (RevMan 5.1). We applied the intention-to-treat method for all analyses. We used both fixed-effect and random-effects models for the primary outcome analysis and the fixed-effect model for the secondary outcomes. We used relative risks (95% CI) for dichotomous outcomes and mean difference (standard deviation (SD) of the mean or 95% CI) for continuous variables.

Subgroup analysis and investigation of heterogeneity

Due to the heterogeneous nature of the selected studies, we conducted subgroup analyses in the following areas.

1. The urgency of surgery (elective or emergency).
2. The type of surgery (general, vascular, cardiac, other).
3. The timing of the intervention (perioperative, intraoperative, postoperative).
4. The type of intervention (fluids, fluids with vasoactive agents).
5. The intervention goals (CO, SV, oxygen indices).

Sensitivity analysis

We analysed mortality, both over the longest follow-up and hospital or 28 day mortality, with fixed-effect and random-effects models. In addition, we excluded studies with fewer than 100 participants. The intervention in the protocol group varied. The control group in some studies had explicit blood flow goals to standardize care. Further, some studies did not fully control for co-interventions, for instance admission to critical care. We performed sensitivity analysis excluding these studies.

RESULTS

Description of studies

All included studies were RCTs of surgical participants. Compared to controls, the intervention group had a separate protocol to optimise global blood flow, measured by cardiac output (CO) or oxygen delivery.

Results of the search

The initial electronic search identified 18,951 potential studies ([Figure 1](#)). After removal of duplicated studies, the search yielded 10,462 studies. No additional studies were identified by contacting experts in the field or relevant pharmaceutical companies or by searching the personal reference databases of the authors or Steering Group. No additional studies were identified following screening of reference lists of potentially eligible studies and previously published systematic reviews.

Figure 1. Study flow diagram.

We identified 66 potentially eligible studies following screening of the abstracts of studies. Of those 66 studies, 35 potentially eligible studies did not meet the study inclusion criteria for the reasons summarized in [Characteristics of excluded studies](#).

The remaining 31 fully published studies (5292 participants) met the inclusion criteria. We have summarized the Included studies in [Characteristics of included studies](#); [Appendix 5](#); and [Appendix 6](#). We contacted study authors for additional data where necessary.

Included studies

We included 31 studies in the review (see [Characteristics of included studies](#)). The studies were conducted in Europe (20), USA (seven), India (one), Brazil (one), Japan (one) and Canada (one). Most studies (24 studies) recruited participants having elective surgery. The studies were published between 1988 and 2011

([Appendix 5](#); [Appendix 6](#)).

Excluded studies

We excluded 35 studies ([Characteristics of excluded studies](#)). We excluded studies that included trauma patients (10) and septic or critically ill patients (nine) unless all patients underwent surgery.

Risk of bias in included studies

We evaluated the risk of bias of included studies with the Cochrane tool ([Higgins 2011](#)). This was performed by two authors (AD, AV) independently and we resolved any disparity by discussion and the involvement of a third person (MG). We present the methodological quality in a summary table and a graph ([Figure 2](#); [Figure 3](#)).

Figure 2. Risk of bias graph: review authors' judgements about each risk of bias item presented as percentages across all included studies.

Figure 3. Risk of bias summary: review authors' judgements about each risk of bias item for each included study.

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias
Bender 1997	⚠	?	?	?	+	+	+
Berlauk 1991	+	?	?	?	+	+	?
Bonazzi 2002	+	?	?	?	+	+	+
Boyd 1993	?	?	?	?	+	+	?
Cecconi 2011	+	+	+	?	+	+	+
Challand 2012	+	+	+	+	+	+	+
Conway 2002	?	?	+	?	+	+	+
Donati 2007	+	+	?	?	+	?	?
Gan 2002	+	+	?	?	+	+	+
Jerez 2001	?	?	?	?	+	+	+
Jhanji 2010	+	+	?	+	+	+	?
Kapoor 2007	?	+	?	?	+	+	?
Lobo 2000	?	+	?	?	+	+	+
Mayer 2010	?	+	?	+	+	+	⚠
Mckendry 2004	+	+	+	?	+	+	?
Mythen 1995	?	+	?	?	+	+	?
Noblett 2006	?	?	+	+	+	+	?
Pearse 2005	+	+	+	+	+	+	+
Pillai 2011	?	?	+	?	+	+	+
Pölonen 2000	?	+	?	?	+	+	?
Sandham 2003	+	+	?	+	?	+	?
Senagore 2009	+	?	+	+	+	+	+
Shoemaker 1988	+	+	?	?	+	+	+
Sinclair 1997	?	+	+	?	+	+	+
Ueno 1998	+	?	?	?	+	+	+
Valentine 1998	?	+	?	?	+	+	+
Van der Linden 2010	+	+	+	+	+	+	+
Venn 2002	+	+	?	+	+	+	+
Wakeling 2005	+	+	+	?	+	+	?
Wilson 1999	+	+	?	?	+	+	+
Ziegler 1997	?	?	?	?	+	+	+

Allocation

All included studies randomly allocated participants. The method of randomization was described in 25 studies (80%): (Bender 1997; Berlaug 1991; Bonazzi 2002; Cecconi 2011; Challand 2012; Donati 2007; Gan 2002; Jhanji 2010; Kapoor 2007; Lobo 2000; Mayer 2010; Mckendry 2004; Mythen 1995; Pearse 2005; Pölönen 2000; Sandham 2003; Senagore 2009; Shoemaker 1988; Sinclair 1997; Ueno 1998; Valentine 1998; Van der Linden 2010; Venn 2002; Wakeling 2005; Wilson 1999). However, in seven studies (Kapoor 2007; Lobo 2000; Mayer 2010; Mythen 1995; Pölönen 2000; Sinclair 1997; Valentine 1998) it was unclear whether the sealed envelope technique allocated participants sequentially. In one study (Bender 1997) participants were allocated by a surgical intensivist, which may have introduced selection bias. We assessed as adequate the random allocation in 17 studies (55%) (Berlaug 1991; Bonazzi 2002; Cecconi 2011; Challand 2012; Donati 2007; Gan 2002; Jhanji 2010; Mckendry 2004; Pearse 2005; Sandham 2003; Senagore 2009; Shoemaker 1988; Ueno 1998; Van der Linden 2010; Venn 2002; Wakeling 2005; Wilson 1999).

We assessed the methods of allocation concealment as adequate for 20 studies (65%) (Cecconi 2011; Challand 2012; Donati 2007; Gan 2002; Jhanji 2010; Kapoor 2007; Lobo 2000; Mayer 2010; Mckendry 2004; Mythen 1995; Pearse 2005; Pölönen 2000; Sandham 2003; Shoemaker 1988; Sinclair 1997; Valentine 1998; Van der Linden 2010; Venn 2002; Wakeling 2005; Wilson 1999).

Blinding

We assessed blinding of personnel or participants as adequate in only 12 studies (39%), reflecting the nature of the intervention (Cecconi 2011; Challand 2012; Conway 2002; Mckendry 2004;

Noblett 2006; Pearse 2005; Pillai 2011; Senagore 2009; Sinclair 1997; Van der Linden 2010; Venn 2002; Wakeling 2005). We assessed blinding of outcome assessment as adequate in eight studies (26%) (Challand 2012; Jhanji 2010; Mayer 2010; Noblett 2006; Pearse 2005; Sandham 2003; Senagore 2009; Van der Linden 2010).

Incomplete outcome data

Attrition bias was detected in one study (Sandham 2003) where a large number of participants were lost to follow-up, which may have introduced attrition bias.

Selective reporting

All anticipated outcomes were reported by the included studies.

Other potential sources of bias

In Mayer 2010, the second author has been found to have fabricated results in some clinical studies. We recognized this as a potential high risk.

Exclusion of participants after randomization was noted in seven studies (Berlaug 1991; Kapoor 2007; Mayer 2010; Mckendry 2004; Noblett 2006; Pölönen 2000; Wakeling 2005), which may have induced selection bias.

To test the effect of publication bias, we produced a contour-enhanced funnel plot for the primary outcome (Figure 4), the subsequent Harbord test showing a significant small-studies effect: regression bias -0.72 (95% CI -0.08 to -1.39) (Figure 5). Similarly, the rate of complications (Figure 6; Figure 7) showed evidence of a small studies effect. No other outcome demonstrated small study effects (Harbord 2006).

Figure 4. Contour-enhanced funnel plot: mortality.

Figure 5. Galbraith plot, Harbord analysis, mortality. The regression slope is -0.72 (-0.08 to -1.39).

Figure 6. Contour-enhanced funnel plot for rate of complications.

Figure 7. Galbraith plot of Harbord analysis for rate of complications.

Effects of interventions

See: [Summary of findings for the main comparison](#) Protocol to increase global blood flow compared to control for surgical patients

Data Synthesis

Mortality

1.1 Long-term mortality

Thirty studies reported mortality data and further information was obtained from authors for one study (Pillai 2011). A number of different definitions were used and some papers reported more than one definition. Using data from the longest reported follow-up, the overall mortality was 238/2677 (8.9%) in the intervention group and 282/2615 (10.8%) in the control group, RR of 0.89 (95% CI 0.76 to 1.05, $P = 0.18$, $I^2 = 15\%$) (Analysis 1.1). The results were sensitive to analytical methods, becoming statistically

significant with two methods: the inverse variance random-effects model, RR of 0.72 (95% CI 0.55 to 0.95, $P = 0.02$, $I^2 = 15\%$); the Mantel-Haenszel random-effects model, RR of 0.72 (95% CI 0.55 to 0.95, $P = 0.02$, $I^2 = 16\%$) (Appendix 7).

1.2 Hospital or 28 day mortality

Hospital or 28 day mortality was reported in 30 studies and further information was obtained from one study (Pillai 2011). Pooled hospital or 28 day mortality was 146/2677 (5.4%) in the intervention group and 192/2615 (7.3%) in the control group, RR of 0.81 (95% CI 0.65 to 1.00, $P = 0.06$, $I^2 = 1\%$) (Analysis 1.2). The results were sensitive to analytical methods, becoming significant with three methods: the inverse variance random-effects model, RR of 0.79 (95% CI 0.63 to 0.99, $P = 0.04$, $I^2 = 1\%$); the Mantel-Haenszel fixed-effect model, RR of 0.77 (95% CI 0.63 to 0.95, $P = 0.01$, $I^2 = 2\%$) and random-effects model, RR of 0.78 (95% CI 0.62 to 0.99, $P = 0.04$, $I^2 = 2\%$) (Appendix 8).

Morbidity

We analysed seven categories of morbidity using the investigators' definitions. No two studies used the same list of morbidities following surgery (Table 1). In most cases no specific criteria were listed for morbidities. No two studies used the same criteria.

2.1 Renal impairment

We accepted the rate of renal impairment reported by study authors: we did not apply a single definition across studies. Data on renal impairment were available for 21 studies (Bender 1997; Berlaak 1991; Bonazzi 2002; Boyd 1993; Cecconi 2011; Challand 2012; Donati 2007; Gan 2002; Jhanji 2010; Kapoor 2007; Lobo 2000; Mayer 2010; Mckendry 2004; Mythen 1995; Pölönen 2000; Sandham 2003; Shoemaker 1988; Valentine 1998; Venn 2002; Wakeling 2005; Wilson 1999). The intervention reduced the rate of renal impairment, RR of 0.71 (95% CI 0.57 to 0.90, $P = 0.004$, $I^2 = 20\%$) (Analysis 2.1).

2.2 Arrhythmia

Arrhythmia was reported in 16 studies (Bender 1997; Berlaak 1991; Bonazzi 2002; Cecconi 2011; Kapoor 2007; Lobo 2000; Mayer 2010; Mckendry 2004; Pearse 2005; Sandham 2003; Senagore 2009; Shoemaker 1988; Valentine 1998; Venn 2002; Wilson 1999; Ziegler 1997). However, we excluded three studies for which there were unit-of-analysis issues: two studies (Bender 1997; Berlaak 1991) reported the number of events; one of these studies and one other (Bender 1997; Valentine 1998) reported for both the intraoperative and postoperative periods. One study (Shoemaker 1988) reported transient dysrhythmias ("almost always premature ventricular complexes") during insertion of pulmonary artery (PA) catheters. This was reported as a combined percentage (12%) for both control and protocol PA catheter groups. We were unable to identify the exact rate of arrhythmias for each group separately and therefore excluded this study from the analysis. For the 12 studies (Bonazzi 2002; Cecconi 2011; Kapoor 2007; Lobo 2000; Mayer 2010; Mckendry 2004; Pearse 2005; Sandham 2003; Senagore 2009; Venn 2002; Wilson 1999; Ziegler 1997) that we were able to analyse, there was no significant difference between groups in development of an arrhythmia, RR of 0.84 (95% CI 0.67 to 1.06, $P = 0.14$, $I^2 = 0\%$) (Analysis 2.2).

2.3 and 2.4 Infection

Infections were reported several ways in 20 studies (Bender 1997; Boyd 1993; Cecconi 2011; Gan 2002; Jhanji 2010; Lobo 2000; Mayer 2010; Mckendry 2004; Mythen 1995; Pearse 2005; Pillai 2011; Sandham 2003; Senagore 2009; Shoemaker 1988; Sinclair 1997; Valentine 1998; Van der Linden 2010; Venn 2002; Wakeling 2005; Wilson 1999). The number of participants who had infections was reported in nine studies (Bender 1997; Jhanji 2010; Lobo 2000; Mythen 1995; Pillai 2011; Sinclair 1997;

Valentine 1998; Van der Linden 2010; Wakeling 2005). The number of participants with infections was unaffected by the intervention, RR of 0.88 (95% CI 0.69 to 1.12, $P = 0.29$, $I^2 = 0\%$) (Analysis 2.3).

The types of infection (such as pneumonia) were reported separately in 15 studies (Bender 1997; Boyd 1993; Cecconi 2011; Gan 2002; Lobo 2000; Mayer 2010; Mythen 1995; Pillai 2011; Sandham 2003; Shoemaker 1988; Sinclair 1997; Valentine 1998; Van der Linden 2010; Venn 2002; Wilson 1999). Nine studies (Boyd 1993; Cecconi 2011; Gan 2002; Mayer 2010; Pearse 2005; Sandham 2003; Shoemaker 1988; Venn 2002; Wilson 1999) reported more than one infective complication per participant. It was not possible to add the total number of infections as the exact denominator was unknown. We therefore analysed each infection separately. There was no difference in the rates of: pneumonia, RR of 0.78 (95% CI 0.61 to 1.00, $P = 0.05$, $I^2 = 0\%$); sepsis, RR of 0.68 (95% CI 0.26 to 1.77, $P = 0.43$, $I^2 = 6\%$); abdominal infections, RR of 0.53 (95% CI 0.23 to 1.22, $P = 0.14$, $I^2 = 0\%$); or urinary tract infections, RR of 0.54 (95% CI 0.26 to 1.15, $P = 0.11$, $I^2 = 0\%$). The intervention significantly reduced the rate of wound infections, RR of 0.65 (95% CI 0.50 to 0.84, $P = 0.001$, $I^2 = 22\%$) (Analysis 2.4). Two studies (Mckendry 2004; Senagore 2009) reported on the total number of infections and we were unable to include these studies due to unit-of-analysis issues.

2.5 Respiratory failure or acute respiratory distress syndrome (ARDS)

Respiratory failure or ARDS was reported in nine studies (Boyd 1993; Donati 2007; Gan 2002; Mayer 2010; Mythen 1995; Pearse 2005; Shoemaker 1988; Ueno 1998; Wilson 1999). One study (Wilson 1999) also included the number of participants with prolonged ventilation, which we were unable to analyse due to unit-of-analysis issues. The intervention significantly reduced the rate of respiratory failure, RR of 0.51 (95% CI 0.28 to 0.93, $P = 0.03$, $I^2 = 0\%$) (Analysis 2.5).

2.6 Myocardial infarction

Myocardial infarction was reported in 15 studies (Bender 1997; Berlaak 1991; Bonazzi 2002; Boyd 1993; Cecconi 2011; Kapoor 2007; Mayer 2010; Mckendry 2004; Pearse 2005; Sandham 2003; Shoemaker 1988; Valentine 1998; Venn 2002; Wilson 1999; Ziegler 1997). There was no significant difference in myocardial infarction, RR of 1.01 (95% CI 0.71 to 1.45, $P = 0.95$, $I^2 = 0\%$) (Analysis 2.6).

2.7 Congestive cardiac failure or pulmonary oedema

Congestive heart failure or pulmonary oedema was reported in 14 studies (Bender 1997; Berlaak 1991; Bonazzi 2002; Boyd 1993; Lobo 2000; Mayer 2010; Pearse 2005; Sandham 2003; Shoemaker 1988; Sinclair 1997; Valentine 1998; Venn 2002; Wakeling 2005;

Wilson 1999). There was no significant difference, RR of 1.00 (95% CI 0.81 to 1.24, $P = 0.98$, $I^2 = 0\%$) (Analysis 2.7).

2.8 Venous thrombosis

Venous thrombosis was reported in 10 studies (Boyd 1993; Cecconi 2011; Lobo 2000; Mayer 2010; Pearse 2005; Sandham 2003; Senagore 2009; Shoemaker 1988; Venn 2002; Wilson 1999). There was no difference, RR of 1.04 (95% CI 0.39 to 2.77, $P = 0.93$, $I^2 = 0\%$) (Analysis 2.8).

2.9 Complications

More than one method was used to pool complications. The number of participants with complications was reported by 19 studies (Bender 1997; Berlaak 1991; Bonazzi 2002; Cecconi 2011; Challand 2012; Conway 2002; Donati 2007; Jerez 2001; Jhanji 2010; Lobo 2000; Mayer 2010; Mythen 1995; Noblett 2006; Pearse 2005; Shoemaker 1988; Sinclair 1997; Ueno 1998; Wakeling 2005; Wilson 1999). The number of complications per participant was reported by three studies (Boyd 1993; Jerez 2001; Shoemaker 1988). The number of participants with individual complications or the number of individual complications was reported by 27 studies (Bender 1997; Berlaak 1991; Bonazzi 2002; Boyd 1993; Cecconi 2011; Challand 2012; Conway 2002; Donati 2007; Gan 2002; Jhanji 2010; Kapoor 2007; Lobo 2000; Mayer 2010; Mckendry 2004; Mythen 1995; Noblett 2006; Pearse 2005; Pölönen 2000; Sandham 2003; Senagore 2009; Shoemaker 1988; Ueno 1998; Valentine 1998; Venn 2002; Wakeling 2005; Wilson 1999; Ziegler 1997).

We did not pool data for the number of complications because of this variation and the associated unit-of-analysis issues. Further, six studies (Bonazzi 2002; Boyd 1993; Gan 2002; Kapoor 2007; Mckendry 2004; Pillai 2011) that reported the number of participants with complications also reported the individual complications separately, therefore pooling of these would again lead to unit-of-analysis issues. Two studies (Bender 1997; Valentine 1998) reported the number of participants with complications separately for the intraoperative and postoperative periods. We were unable to combine these outcomes due to unit-of-analysis issues. We therefore pooled 17 studies (Berlaak 1991; Cecconi 2011; Challand 2012; Conway 2002; Donati 2007; Jerez 2001; Jhanji 2010; Lobo 2000; Mayer 2010; Mythen 1995; Noblett 2006; Pearse 2005; Shoemaker 1988; Sinclair 1997; Ueno 1998; Wakeling 2005; Wilson 1999). The number of participants with complications was reduced by the intervention, RR of 0.68 (95% CI 0.58 to 0.80, $P < 0.00001$, $I^2 = 34\%$) (Analysis 2.9).

Health status

No study reported health status.

Resource use

3.1 Postoperative hospital stay

Postoperative length of hospital stay was reported in 28 studies (Bender 1997; Berlaak 1991; Bonazzi 2002; Boyd 1993; Cecconi 2011; Challand 2012; Conway 2002; Donati 2007; Gan 2002; Jhanji 2010; Kapoor 2007; Lobo 2000; Mayer 2010; Mckendry 2004; Mythen 1995; Noblett 2006; Pearse 2005; Pillai 2011; Pölönen 2000; Sandham 2003; Senagore 2009; Shoemaker 1988; Sinclair 1997; Valentine 1998; Van der Linden 2010; Venn 2002; Wakeling 2005; Wilson 1999). This was reported as the mean (SD) by seven studies (Bender 1997; Berlaak 1991; Donati 2007; Gan 2002; Kapoor 2007; Pearse 2005; Shoemaker 1988), mean (range) by one study (Mythen 1995), mean (95% CI) by two studies (Pillai 2011; Venn 2002), mean (SEM) by one study (Valentine 1998) and median (range or interquartile range (IQR)) by 15 studies (Bonazzi 2002; Boyd 1993; Cecconi 2011; Challand 2012; Conway 2002; Jhanji 2010; Lobo 2000; Mayer 2010; Mckendry 2004; Noblett 2006; Pölönen 2000; Sandham 2003; Sinclair 1997; Van der Linden 2010; Wakeling 2005). We excluded one study from this analysis (Senagore 2009), for which we were unable to get further information. We obtained additional details for five studies (Jhanji 2010; Mythen 1995; Noblett 2006; Wakeling 2005; Wilson 1999). We used the statistical equation by Hozo 2005 to convert the median (range/IQR) to mean (SD). We estimated the SD as $IQR/1.35$, $SEM \times \sqrt{n}$ or $95\% CI / 1.96$. Four studies (Berlaak 1991; Jhanji 2010; Shoemaker 1988; Venn 2002) had two groups in either of the intervention or control groups and these were numerically combined using equation 7.7a in Higgins 2011. The intervention significantly reduced the postoperative length of hospital stay, mean 1.16 days (95% CI 0.43 to 1.89, $P = 0.002$). We used the random-effects model as the $I^2 = 87\%$ (Analysis 3.1).

3.2 Postoperative intensive care stay

Postoperative length of critical care stay was reported by 14 studies (Bender 1997; Berlaak 1991; Boyd 1993; Jerez 2001; Jhanji 2010; Kapoor 2007; Lobo 2000; Mayer 2010; Mythen 1995; Pearse 2005; Pölönen 2000; Shoemaker 1988; Valentine 1998; Wilson 1999). This was reported as the mean (SD) by six studies (Bender 1997; Berlaak 1991; Jerez 2001; Kapoor 2007; Mayer 2010; Shoemaker 1988), mean (range) by one study (Mythen 1995), mean (SEM) by one study (Valentine 1998) and median (range/IQR) by five studies (Boyd 1993; Jhanji 2010; Lobo 2000; Pearse 2005; Pölönen 2000). We were able to obtain additional information for three studies (Jhanji 2010; Mythen 1995; Wilson 1999). Numerical conversion to mean (SD) was performed according to the previous paragraph. There was no difference in postoperative length of critical care stay, mean difference of 0.45 days (95% CI -0.03 to 0.94, $P = 0.06$). We used the random-effects model as the $I^2 = 87\%$ (Analysis 3.2).

Three studies (Bender 1997; Berlaak 1991; Shoemaker 1988) reported cost (USD), none of which found a statistical difference. Three other studies (Boyd 1993; Mythen 1995; Wilson 1999) reported cost in separate publications from the original report (two reported GBP (Guest 1997; Mythen 1994); one reported EUR (Fenwick 2002)). Two of these (Fenwick 2002; Guest 1997) reported that the intervention significantly reduced cost. The third (Mythen 1994) reported cost for a subgroup of patients included in the trial and these data were not analysed by treatment groups. Only one study reported means and SDs (Berlaak 1991) and only one study reported means and SEMs (Bender 1997) for cost data. In view of the variety of currencies and statistical descriptors we did not attempt to pool these data.

Subgroup mortality analyses

Timing of intervention

The intervention was commenced in the preoperative period in nine studies (Bender 1997; Berlaak 1991; Bonazzi 2002; Boyd 1993; Sandham 2003; Shoemaker 1988; Valentine 1998; Wilson 1999; Ziegler 1997), in the intraoperative period in 15 studies (Cecconi 2011; Challand 2012; Conway 2002; Donati 2007; Gan 2002; Lobo 2000; Mayer 2010; Mythen 1995; Noblett 2006; Pillai 2011; Senagore 2009; Sinclair 1997; Van der Linden 2010; Venn 2002; Wakeling 2005) and in the postoperative period in nine studies (Boyd 1993; Jerez 2001; Jhanji 2010; Kapoor 2007; Mckendry 2004; Pearse 2005; Pölönen 2000; Ueno 1998). In one study (Berlaak 1991) participants were randomized to two intervention groups (a preoperative and an intraoperative group) with a shared control group. In another study (Boyd 1993) the intervention was initiated either preoperatively or postoperatively depending on when the participants came to the attention of the investigators. There was no evidence that this had any effect on the chances of being recruited into the study and therefore we did not consider that this had potential to confound the randomization process. Further, one study (Lobo 2000) had both intraoperative and postoperative interventions. Timing of the intervention did not interact with mortality: preoperative, RR of 0.96 (95% CI 0.79 to 1.17, $P = 0.69$, $I^2 = 63\%$); intraoperative, RR of 0.67 (95% CI 0.40 to 1.13, $P = 0.13$, $I^2 = 0\%$); postoperative, RR of 0.73 (95% CI 0.50 to 1.06, $P = 0.10$, $I^2 = 0\%$) (Analysis 1.4).

Type of intervention

The intervention involved fluids alone in 10 studies (Challand 2012; Conway 2002; Gan 2002; Mythen 1995; Noblett 2006; Pillai 2011; Senagore 2009; Sinclair 1997; Venn 2002; Wakeling 2005) and fluids in combination with vasoactive drugs in 20 studies (Bender 1997; Berlaak 1991; Bonazzi 2002; Boyd 1993; Cecconi 2011; Donati 2007; Jerez 2001; Kapoor 2007; Lobo 2000; Mayer 2010; Mckendry 2004; Pearse 2005; Pölönen 2000;

Sandham 2003; Shoemaker 1988; Ueno 1998; Valentine 1998; Van der Linden 2010; Wilson 1999; Ziegler 1997). One study (Jhanji 2010) had two intervention groups; one group had fluid alone and the other had fluids and dopexamine. These groups were analysed separately. There was no difference in mortality between groups according to the intervention provided: fluids alone, RR of 0.80 (95% CI 0.46 to 1.39, $P = 0.43$, $I^2 = 0\%$); fluids in combination with vasoactive drugs, RR of 0.90 (95% CI 0.76 to 1.07, $P = 0.23$, $I^2 = 41\%$) (Analysis 1.5).

Type of goal

Fourteen studies (Bender 1997; Berlaak 1991; Bonazzi 2002; Boyd 1993; Kapoor 2007; Lobo 2000; Mayer 2010; Pearse 2005; Sandham 2003; Shoemaker 1988; Ueno 1998; Valentine 1998; Van der Linden 2010; Wilson 1999) used CO and oxygen transport goals; four studies (Donati 2007; Jerez 2001; Pölönen 2000; Ziegler 1997) used mixed venous oxygen saturation, oxygen extraction and lactate; and 13 studies (Cecconi 2011; Challand 2012; Conway 2002; Gan 2002; Jhanji 2010; Mckendry 2004; Mythen 1995; Noblett 2006; Pillai 2011; Senagore 2009; Sinclair 1997; Venn 2002; Wakeling 2005) used stroke volume (SV) goals. Mortality was not reduced for any of the three subgroups: CO and oxygen transport, RR of 0.91 (95% CI 0.75 to 1.09, $P = 0.91$, $I^2 = 58\%$); mixed venous oxygen saturations, oxygen extraction and lactate, RR of 0.83 (95% CI 0.50 to 1.38, $P = 0.47$, $I^2 = 0\%$); SV, RR of 0.84 (95% CI 0.51 to 1.41, $P = 0.51$, $I^2 = 0\%$) (Analysis 1.6).

Mode of surgery

Twenty-four studies (Bender 1997; Berlaak 1991; Bonazzi 2002; Cecconi 2011; Challand 2012; Conway 2002; Donati 2007; Gan 2002; Jhanji 2010; Jerez 2001; Kapoor 2007; Lobo 2000; Mayer 2010; Mythen 1995; Noblett 2006; Pillai 2011; Pölönen 2000; Senagore 2009; Ueno 1998; Valentine 1998; Van der Linden 2010; Wakeling 2005; Wilson 1999; Ziegler 1997) recruited participants having only elective procedures; two studies were exclusively of urgent or emergency surgery (Sinclair 1997; Venn 2002) and five had a mix of urgent or emergency and elective operations (Boyd 1993; Mckendry 2004; Pearse 2005; Sandham 2003; Shoemaker 1988). None of the studies in this latter group were able to provide separate data to allow comparison between elective and urgent or emergency groups. Intervention significantly reduced the mortality of participants in RCTs of elective surgery, RR of 0.68 (95% CI 0.48 to 0.94, $P = 0.02$, $I^2 = 0\%$); mortality was unchanged for emergency or urgent operations, RR of 0.68 (95% CI 0.23 to 2.06, $P = 0.50$, $I^2 = 0\%$) (Analysis 1.7).

Type of surgery

Six studies (Bender 1997; Berlaak 1991; Bonazzi 2002; Valentine 1998; Van der Linden 2010; Ziegler 1997) were exclusively of

participants undergoing vascular surgery. Five additional studies (Boyd 1993; Lobo 2000; Pearse 2005; Sandham 2003; Wilson 1999) included participants undergoing vascular surgery, but in only one of these were group-specific mortality data available (Boyd 1993). Five studies were of patients undergoing cardiac surgery (Jerez 2001; Kapoor 2007; Mckendry 2004; Mythen 1995; Pölonen 2000). Fifteen studies were exclusively of patients undergoing general (non-vascular, non-cardiac) surgery (Cecconi 2011; Challand 2012; Conway 2002; Donati 2007; Gan 2002; Jhanji 2010; Mayer 2010; Noblett 2006; Senagore 2009; Shoemaker 1988; Sinclair 1997; Ueno 1998; Venn 2002; Wakeling 2005). Five additional studies included patients undergoing general surgery (Boyd 1993; Lobo 2000; Pearse 2005; Sandham 2003; Wilson 1999) but in only one of these were group-specific mortality data available (Boyd 1993). There was no interaction between type of surgery and the intervention; vascular, RR of 0.78 (95% CI 0.34 to 1.79, $P = 0.56$, $I^2 = 18\%$); cardiac, RR of 0.81 (95% CI 0.48 to 1.35, $P = 0.42$, $I^2 = 0\%$); and general surgery, RR of 0.66 (95% CI 0.41 to 1.07, $P = 0.09$, $I^2 = 0\%$) (Analysis 1.8).

Sensitivity analyses

We performed sensitivity analyses of the analysis method used to generate relative risks for mortality (Appendix 7; Appendix 8). The results were dependant upon both the analytical method and whether a random-effects model or fixed-effect model was used. There was no difference in mortality when small studies (fewer than 100 participants) were excluded (Analysis 1.3), consistent with Analysis 1.1. The effect of small studies was significant in the Harbord analysis, with a regression slope of -0.72 (95% CI -0.08 to -1.39). Participants were more likely to die in studies that recruited fewer than 100 participants, RR of 1.84 (95% CI 1.02 to 3.33, $P = 0.04$).

In some studies the fluid and drug management in the control group was comparable with the intervention in other studies. For instance, four studies (Jerez 2001; Lobo 2000; Pölonen 2000; Ueno 1998) had fluid and inotropes administered in response to measures of blood flow (CI or DO_2I) in the control groups. Shoemaker 1988 had one control group with DO_2I driven measures. We performed a sensitivity analysis excluding these studies and the control group from Shoemaker 1988 (Analysis 4.1; Analysis 4.2; Analysis 4.3; Analysis 4.4; Analysis 4.5; Analysis 4.6; Analysis 4.7; Analysis 4.8; Analysis 4.9; Analysis 4.10; Analysis 4.11; Analysis 4.12). The findings were consistent with the primary analyses (Analysis 1.1; Analysis 1.2; Analysis 2.1; Analysis 2.2; Analysis 2.3; Analysis 2.5; Analysis 2.6; Analysis 2.7; Analysis 2.8; Analysis 2.9; Analysis 3.1; Analysis 3.2).

In some studies, fluid and inotrope administration were not the only systematic differences between the control and intervention groups. Five studies (Bender 1997; Bonazzi 2002; Sandham 2003; Valentine 1998; Wilson 1999) did not control for the insertion and presence of a pulmonary artery flow catheter. Three studies

(Cecconi 2011; Kapoor 2007; Mayer 2010) did not control for the presence of other flow sensors (FloTrac or Vigileo) and one study (Venn 2002) did not control for the insertion or presence of an oesophageal doppler probe. In one study (Shoemaker 1988) one control group was not matched for the insertion and presence of a pulmonary artery flow catheter. We also performed sensitivity analyses excluding these studies for all outcome measures (Analysis 5.1; Analysis 5.2; Analysis 5.3; Analysis 5.4; Analysis 5.5; Analysis 5.6; Analysis 5.7; Analysis 5.8; Analysis 5.9; Analysis 5.10; Analysis 5.11; Analysis 5.12). With these studies excluded the intervention reduced mortality (longest follow-up), RR of 0.65 (95% CI 0.48 to 0.89, $P = 0.007$, $I^2 = 0\%$) (Analysis 5.1) and hospital or 28 day mortality, RR of 0.66 (95% CI 0.47 to 0.92, $P = 0.01$, $I^2 = 0\%$) (Analysis 5.2). The rates of renal failure and ARDS were no longer significantly different. The number of participants with complications and their length of hospital stay were not altered in this analysis, remaining significantly different between groups. This meta-analysis was dominated by one study (Sandham 2003). In this study a large number of participants were lost to follow-up. We performed sensitivity analyses for the outcomes of mortality (longest follow-up and hospital or 28 day mortality) excluding this study and assuming the possibility that all patients who were lost to follow-up died. The results were not sensitive to these analyses (Analysis 6.1; Analysis 6.2).

DISCUSSION

Summary of main results

The key finding of this review is that the perioperative administration of fluids, with or without vasoactive drugs, targeted to increase global blood flow defined by explicit measured goals reduced postoperative complications and length of stay but did not reduce mortality, using the inverse variance method. The exclusion of larger studies (> 100 participants) resulted in mortality being reduced by the intervention (RR 0.51, 95% CI 0.29 to 0.90, $P = 0.02$). Mortality was also significantly reduced when we used random-effects models (Mantel-Haenszel and inverse variance, RR 0.72, 95% CI 0.55 to 0.95, $P = 0.02$), but not fixed-effect models (Mantel-Haenszel RR 0.85, 95% CI 0.73 to 1.00, $P = 0.05$; inverse variance RR 0.89, 95% CI 0.76 to 1.05, $P = 0.18$; or Peto odds ratio 0.83, 95% CI 0.69 to 1.00, $P = 0.05$). We calculated similar results for hospital and 28 day mortality. When control group care was managed using a protocol that included explicit goals less than the intervention group (in contrast to 'usual care'), mortality was not reduced (longest follow-up inverse variance RR 0.94, 95% CI 0.79 to 1.12, $P = 0.45$; hospital or 28 day inverse variance RR 0.84, 95% CI 0.67 to 1.07, $P = 0.14$). When studies with intervention groups that were less well controlled for the intervention (for example pulmonary artery catheters were not

matched to intervention groups) were excluded, there was a significant reduction in mortality at the longest follow-up (inverse variance RR 0.65, 95% CI 0.48 to 0.89, $P = 0.007$) and hospital or 28 day mortality (inverse variance RR 0.66, 95% CI 0.47 to 0.92, $P = 0.01$). It is notable that the sensitivity analyses are of limited value as they tend to reflect the inclusion or exclusion of the single largest study (Sandham 2003).

The limited data indicate that for every 100 patients exposed to treatment, one can expect 13 in 100 (from 40/100 to 27/100) to avoid a complication, 2/100 to avoid renal impairment (from 8/100 to 6/100), 5/100 to avoid respiratory failure (from 10/100 to 5/100), and 4/100 to avoid postoperative wound infection (from 10/100 to 6/100), with no effect on other types of morbidity (myocardial infarction, arrhythmia, congestive cardiac failure or pulmonary oedema, venous thrombosis, and the number of patients with infections). These results were unchanged following sensitivity analyses that excluded studies where the control group care was managed using a protocol that included explicit goals that were less than the intervention group (in contrast to 'usual care'). When studies using intervention groups that were less well controlled (control groups not matched to intervention groups) were excluded only the number of patients with complications was reduced, by 12/100 (from 36/100 to 24/100).

The hospital length of stay was reduced by about one day, from 12.4 to 11.2 days, and was not sensitive to exclusion of studies where the control group care was managed using a protocol that included explicit goals that were less than the intervention group (in contrast to 'usual care') or studies using intervention groups that were less well controlled. There was no difference in critical care stay in the intervention group. This was sensitive to exclusion of studies using intervention groups that were less well controlled and the reduction was less than a day (from 4 to 3.3 days). There were insufficient data to conduct a meta-analysis of cost and no data available describing quality of life.

A stratified meta-analysis to address secondary hypotheses, determined a priori, suggested that mortality was reduced in the intervention group when study participants underwent elective surgery. The predefined analysis plan, using mortality from the longest available follow-up, increased the weight attributed to the two largest studies that both reported one-year follow-up. Only one other study reported follow-up beyond 60 days. In this group of studies a proportion of the operations were for cancer resection, therefore introducing a possible competing cause of mortality.

Overall completeness and applicability of evidence

Our systematic review pooled data from 31 studies with 5292 participants (Bender 1997; Berlauk 1991; Bonazzi 2002; Boyd 1993; Ceconi 2011; Challand 2012; Donati 2007; Conway 2002; Gan 2002; Jerez 2001; Jhanji 2010; Kapoor 2007; Lobo 2000; Mayer 2010; Mckendry 2004; Mythen 1995; Noblett 2006; Pearse

2005; Pillai 2011; Pölönen 2000; Sandham 2003; Senagore 2009; Shoemaker 1988; Sinclair 1997; Ueno 1998; Valentine 1998; Van der Linden 2010; Venn 2002; Wakeling 2005; Wilson 1999; Ziegler 1997). Study inclusion criteria were tightly defined and the meta-analysis was rigorously conducted according to a predefined analysis plan addressing specific hypotheses. The meta-analysis combined data from a group of predominantly underpowered single centre studies. However, the included studies reflect international practice, although the majority of included studies are from major teaching centres. The pooled studies included adults (age > 16 years) undergoing several types of surgery, including abdominal, urology, gynaecology, orthopaedic, cardiac, thoracic and vascular. Therefore, the included studies represent the population for whom the intervention might be considered.

Quality of the evidence

The quality of outcome data reporting in the included studies was variable. Mortality was reported over a variety of time frames and other outcomes were either limited or inconsistent between studies, precluding meaningful analyses in many cases. Diverse criteria and descriptions for morbidities, along with infrequent use of validated metrics, limited the precision of treatment effect estimates and the confidence that can be attached to them. Furthermore, pooling of different types of morbidity was inconsistent, limiting assessment of the overall 'morbidity load'.

Most studies tested a complex package of care (for example fluids, inotropes, monitor, goals, critical care environment) rather than a single clearly-defined intervention. Heterogeneity in the components of such a complex intervention may contribute to study heterogeneity within a systematic review. Study heterogeneity may reduce the precision of treatment effect estimates and reduce the generalizability of the results of meta-analyses (Louis 1991). By definition, it is not easy to define precisely the 'active ingredients' of a complex intervention (MRC 2000). However, hypothesis-generating subgroup analyses indicated that there were insufficient data to distinguish statistically between many of the prespecified subgroups, and highlighted the limited quantity of data in some areas for example emergency surgery.

Several possible sources of bias arose in this meta-analysis. The primary analysis was sensitive to the analytical methods used, the exclusion of larger studies, and the exclusion of studies that inadequately controlled for the intervention. Larger studies are less likely to be affected by bias (Kjaergard 2001) and the inclusion of lower quality studies can alter the interpretation of the benefit of interventions in meta-analysis (Moher 1998).

Statistical heterogeneity was generally absent (I^2 less than 40%). Except for some analyses such as hospital length of stay, there was evidence of significant statistical heterogeneity. We used random-effects models in all cases where I^2 exceeded 40%. In all analyses of mortality, the point estimate of effect was less than 0.90, suggesting that the intervention was probably not harmful.

The sensitivity of our results to the methods of analysis indicates that the results of this study are far from clear-cut. Further research is essential in this area both to address the overall objective of this review and to focus on specific questions.

The studies included in this review are typical of studies in critical care research in general in that the majority of studies are underpowered and from single centres (Langham 2002) and about half the studies are small (< 100 participants). Future studies in this area should test an explicitly framed hypothesis, be adequately powered, methodologically rigorously and blinded (where possible). Reporting of outcomes should be standardized (to allow comparison between studies and to facilitate the conduct of future meta-analyses) and inclusive (morbidity, health status, resource usage).

Potential biases in the review process

The possibility of publication bias cannot be excluded. We found no evidence of this from contact with experts and industry but some of the identified published abstracts have yet to be published as full peer-reviewed papers. Harbord's regression test was significant at $P = 0.03$, suggesting small study effects. Language bias is possible because of the electronic databases and conferences we searched. Flaws in the original study designs are a significant potential source of bias. The meta-analysis includes 5292 participants but the unit of analysis is the study (or study subgroup) and the sample size (31 studies) is relatively small. The results of the subgroup analyses should be considered as hypothesis-generating only and are largely influenced by inclusion or exclusion of a single study (Sandham 2003).

Agreements and disagreements with other studies or reviews

This review represents the best up-to-date summary of the literature. We framed a tightly defined question and used explicit inclusion criteria for studies and a predefined analysis plan. Our primary result does not agree with previous reviews (Boyd 1999; Brienza 2009; Giglio 2009; Hamilton 2011; Heyland 1996; Ivanov 1997; Kern 2002; Poeze 2005), which have been uniformly supportive of this intervention. This may be explained by the precision of the question we addressed (for example other reviews included trauma patients not having surgery) and the analytical methods used. The results of our systematic review do, however, agree with the results of the largest study in this area (Sandham 2003). However, it is of concern that the Sandham study dominates the review primary analysis both in terms of number of patients (1994/5292) and weight (67%), and that the Sandham study was one of the studies where the intervention group was less well controlled (control groups not matched to intervention groups).

Research should focus on answering these questions. Sandham et al showed that a large multicentre study can be conducted in this

area and several such studies are currently ongoing, in particular a large multicentre study anticipated to recruit about 700 participants (Pearse 2009). Future research will hopefully disentangle the complex package of care that forms the intervention (for example fluids, inotropes or vasoactive agents, monitor, goals, critical care environment) and thereby identify which components are effective in different clinical contexts.

AUTHORS' CONCLUSIONS

Implications for practice

Clinicians should base their decision whether to manipulate perioperative global blood flow on the magnitude of reductions in postoperative morbidities and length of hospital stay rather than upon the assumption that mortality will be reduced. For every 100 patients exposed to the intervention one can expect 13/100 to avoid having complications (from 40 to 27 per 100); 2/100 to avoid renal impairment (from 8 to 6 per 100); 5/100 to avoid respiratory failure (from 10 to 5 per 100); and 4/100 to avoid postoperative wound infection (from 10 to 6 per 100). Patients remain in hospital about one day less and there is no increase in harm. This intervention should be considered where the relevant resources are available and implementation will not otherwise harm the patient (for example delay in definitive care).

Implications for research

A specific limitation of this review is the large number of studies that were published more than 10 years and the limited amount of data that represent current practice and outcomes. A specific group that particularly merits further study, in view of the high incidence of mortality and morbidity and limited available data, is patients undergoing emergency surgery.

Future studies in this area should test an explicitly framed hypothesis, be adequately powered (and preferably multicentre), methodologically rigorous, and include blinded interventions where possible. Reporting of outcomes should be standardized (to allow comparison between studies and to facilitate the conduct of future meta-analyses) and inclusive (morbidity, health status, resource usage).

ACKNOWLEDGEMENTS

We would like to thank John Carlisle (Content Editor) and Nathan Pace (Statistical Editor) for their significant input for the preparation of this review.

We would like to thank Karen Hovhannisyan (Trial Search Co-ordinator) for performing the search for this review and Jane Cracknell (Managing Editor) for her patience, support and advice.

We acknowledge the advice and support provided by the late Professor David Bennett in conceiving and performing previous work that has formed the foundation for the present study.

We would like to thank Dr Martha Delgado, Dr Helen Higham and Prof Pierre Foex for their help and editorial advice during the preparation of the protocol for this systematic review. We would also like to thank the following.

- Dr Jonathan Hyam, Dr Shahzad Shaefi and Dr Aanand Vibhakkar (AV) for helping with literature searching and identification of potential studies.
- The 'Optimisation Systematic Review' Steering Group for advice and guidance: Dr Richard Beale, Professor David Bennett, Dr Owen Boyd, Professor Mark Emberton, Dr Julia Langham, Professor Ian Roberts, Dr Jonathan Thompson.
- Professor Stephen Senn (UCL, London) for statistical comments.

- Dr Tony Brady for comments on an earlier draft of this paper.

- Dr Lorenzo Grazioli for help with translation of Spanish text.

Some of this work was undertaken at the University Hospital Southampton NHS Foundation Trust-University of Southampton Respiratory Biomedical Research Unit,

which received a proportion of funding from the United Kingdom Department of Health's National Institute for Health Research Biomedical Research Units funding scheme.

Some of this work was undertaken at University College London Hospital-University College London Comprehensive Biomedical Research Centre, which received a proportion of funding from the United Kingdom Department of Health's National Institute for Health Research Biomedical Research Centres funding scheme.

REFERENCES

References to studies included in this review

Bender 1997 *{published data only}*

Bender JS, Smith-Meek MA, Jones CE. Routine pulmonary artery catheterization does not reduce morbidity and mortality of elective vascular surgery: results of a prospective, randomized trial. *Annals of Surgery* 1997;**226**(3):229–36. [PUBMED: 9339929]

Berlauk 1991 *{published data only}*

* Berlauk JF, Abrams JH, Gilmour IJ, O'Connor SR, Knighton DR, Cerra FB. Preoperative optimization of cardiovascular haemodynamics improves outcome in peripheral vascular surgery. A prospective, randomized clinical trial. *Annals of Surgery* 1991;**214**(3):289–97. [PUBMED: 1929610]

Bonazzi 2002 *{published data only}*

Bonazzi M, Gentile F, Biasi GM, Migliavacca S, Esposti D, Cipolla M, et al. Impact of perioperative haemodynamic monitoring on cardiac morbidity after major vascular surgery in low risk patients. A randomised pilot trial. *European Journal of Vascular and Endovascular Surgery* 2002;**23**(5):445–51. [PUBMED: 12027474]

Boyd 1993 *{published data only}*

Boyd O, Grounds RM, Bennett ED. A randomized clinical trial of the effect of deliberate perioperative increase of oxygen delivery on mortality in high-risk surgical patients. *JAMA* 1993;**270**:2699–707. [PUBMED: 7907668]

Cecconi 2011 *{published data only}*

Cecconi M, Fasano N, Langiano N, Divella M, Costa MG, Rhodes A, et al. Goal-directed haemodynamic

therapy during elective total hip arthroplasty under regional anaesthesia. *Critical Care* 2011;**15**:R132. [PUBMED: 21624138]

Challand 2012 *{published data only}*

Challand C, Struthers R, Sneyd JR, Erasmus PD, Mellor N, Hosie KB, et al. Randomized controlled trial of intraoperative goal-directed fluid therapy in aerobically fit and unfit patients having major colorectal surgery. *British Journal of Anaesthesia* 2012;**108**(1):53–62. [DOI: 10.1093/bja/aer273]

Conway 2002 *{published and unpublished data}*

Conway DH, Mayall R, Abdul-Latif MS, Gilligan S, Tackaberry C. Randomised controlled trial investigating the influence of intravenous fluid titration using oesophageal Doppler monitoring during bowel surgery. *Anaesthesia* 2002;**57**(9):845–9. [PUBMED: 12190747]

Donati 2007 *{published data only}*

Donati A, Loggi S, Preiser JC, Orsetti G, Munch C, Gabbanelli V, et al. Goal-directed intraoperative therapy reduces morbidity and length of hospital stay in high-risk surgical patients. *Chest* 2007;**132**(6):1817–24. [PUBMED: 17925428]

Gan 2002 *{published data only}*

Gan TJ, Soppitt A, Maroof M, el Moalem H, Robertson KM, Moretti E, et al. Goal-directed intraoperative fluid administration reduces length of hospital stay after major surgery. *Anesthesiology* 2002;**97**(4):820–6. [PUBMED: 12357146]

Jerez 2001 {published data only}

Jerez Gomez Coronado V, Robles Marcos M, Perez Civantos D, Tejada Ruiz J, Jimeno Torres B, Barragan Gomez Coronado I, et al. Hemodynamic optimization and morbimortality after heart surgery. *Medicina Intensiva (Madrid. Ed. impresa)* 2001;**25**(8):297–302. [CENTRAL: CN-00425435; EMBASE: 2002008255]

Jhanji 2010 {published data only}

Jhanji S, Vivian-Smith A, Lucena-Amaro S, Watson D, Hinds CJ, Pearse RM. Haemodynamic optimisation improves tissue microvascular flow and oxygenation after major surgery: a randomised controlled trial. *Critical Care* 2010;**14**:R151. [PUBMED: 20698956]

Kapoor 2007 {published data only}

Kapoor PM, Kakani M, Chowdhury U, Choudhury M, Lakshmy R, Kiran U. Early goal-directed therapy in moderate to high-risk cardiac surgery patients. *Annals of Cardiac Anaesthesia* 2008;**11**(1):27–34. [PUBMED: 18182756]

Lobo 2000 {published data only}

Lobo SM, Salgado PF, Castillo VG, Borim AA, Polachini CA, Palchetti JC, et al. Effects of maximizing oxygen delivery on morbidity and mortality in high-risk surgical patients. *Critical Care Medicine* 2000;**28**:3396–404. [PUBMED: 11057792]

Mayer 2010 {published data only}

Mayer J, Boldt J, Mengistu AM, Rohm KD, Suttner S. Goal-directed intraoperative therapy based on auto calibrated arterial pressure waveform analysis reduces hospital stay in high-risk surgical patients: a randomized, controlled trial. *Critical Care* 2010;**14**:R18. [PUBMED: 20156348]

McKendry 2004 {published and unpublished data}

McKendry M, McGloin H, Saberi D, Caudwell L, Brady AR, Singer M. Randomised controlled trial assessing the impact of a nurse delivered, flow monitored protocol for optimisation of circulatory status after cardiac surgery. *BMJ* 2004;**329**(7460):258–61. [PUBMED: 15242867]

Mythen 1995 {published and unpublished data}

Mythen MG, Webb AR. Perioperative plasma volume expansion reduces the incidence of gut mucosal hypoperfusion during cardiac surgery. *Archives of Surgery* 1995;**130**:423–9. [PUBMED: 7535996]

Noblett 2006 {published and unpublished data}

Noblett SE, Snowden CP, Shenton BK, Horgan AF. Randomized clinical trial assessing the effect of Doppler-optimized fluid management on outcome after elective colorectal resection. *The British Journal of Surgery* 2006;**93**(9):1069–76. [PUBMED: 16888706]

Pearse 2005 {published data only}

Pearse R, Dawson D, Fawcett J, Rhodes A, Grounds RM, Bennett ED. Early goal-directed therapy after major surgery reduces complications and duration of hospital stay. A randomised, controlled trial [ISRCTN38797445]. *Critical Care* 2005;**9**(6):R687–93. [PUBMED: 16356219]

Pillai 2011 {published data only}

Pillai P, McEleavy, Gaughan M, Snowden C, Nesbitt I, Durkan G, et al. A double-blind randomized controlled trial to assess the effect of doppler optimized intraoperative fluid management on outcome following radical cystectomy. *The Journal of Urology* 2011;**186**:2201–6. [PUBMED: 22014804]

Pölonen 2000 {published data only}

Pölonen P, Ruokonen E, Hippeläinen M, Pöyhönen M, Takala J. A prospective, randomized study of goal-oriented haemodynamic therapy in cardiac surgical patients. *Anesthesia and Analgesia* 2000;**90**:1052–9. [PUBMED: 10781452]

Sandham 2003 {published data only}

Sandham JD, Hull RD, Brant RF, Knox L, Pineo GF, Doig CJ, et al. A randomized, controlled trial of the use of pulmonary-artery catheters in high-risk surgical patients. *The New England Journal of Medicine* 2003;**348**(1):5–14. [PUBMED: 12510037]

Senagore 2009 {published data only}

Senagore AJ, Emery T, Luchtefeld M, Kim D, Dujovny N, Hoedema R. Fluid management for laparoscopic colectomy: A prospective, randomized assessment of goal-directed administration of balanced salt solution or hetastarch coupled with enhanced recovery program. *Diseases of the Colon and the Rectum* 2009;**52**:1935–40. [PUBMED: 19934912]

Shoemaker 1988 {published data only (unpublished sought but not used)}

Shoemaker WC, Appel PL, Kram HB, Waxman K, Lee TS. Prospective trial of supranormal values of survivors as therapeutic goals in high-risk surgical patients. *Chest* 1988;**94**:1176–86. [PUBMED: 3191758]

Sinclair 1997 {published and unpublished data}

Sinclair S, James S, Singer M. Intraoperative intravascular volume optimisation and length of hospital stay after repair of proximal femoral fracture: randomised controlled trial. *BMJ* 1997;**315**:909–12. [PUBMED: 9361539]

Ueno 1998 {published data only}

Ueno S, Tanabe G, Yamada H, Kusano C, Yoshidome S, Nuruki K, et al. Response of patients with cirrhosis who have undergone partial hepatectomy to treatment aimed at achieving supranormal oxygen delivery and consumption. *Surgery* 1998;**123**:278–86. [PUBMED: 9526519]

Valentine 1998 {published data only}

Valentine RJ, Duke ML, Inman MH, Grayburn PA, Hagino RT, Kakish HB, et al. Effectiveness of pulmonary artery catheters in aortic surgery: a randomized trial. *Journal of Vascular Surgery* 1998;**27**:203–11. [PUBMED: 9510275]

Van der Linden 2010 {published data only}

Van der Linden PJ, Dierick A, Wilmin S, Bellens B, De Hert SG. A randomised controlled trial comparing an intraoperative goal-directed strategy with routine clinical practice in patients undergoing peripheral arterial surgery. *European Journal of Anaesthesiology* 2010;**27**:788–93. [PUBMED: 20613538]

Venn 2002 {published and unpublished data}

Venn R, Steele A, Richardson P, Poloniecki J, Grounds M, Newman P. Randomized controlled trial to investigate influence of the fluid challenge on duration of hospital stay and perioperative morbidity in patients with hip fractures. *British Journal of Anaesthesia* 2002;**88**(1):65–71. [PUBMED: 11881887]

Wakeling 2005 {published and unpublished data}

Wakeling HG, McFall MR, Jenkins CS, Woods WG, Miles WF, Barclay GR, et al. Intraoperative oesophageal Doppler guided fluid management shortens postoperative hospital stay after major bowel surgery. *British Journal of Anaesthesia* 2005;**95**(5):634–42. [PUBMED: 16155038]

Wilson 1999 {published data only}

Wilson J, Woods I, Fawcett J, Whall R, Dibb W, Morris C, et al. Reducing the risk of major elective surgery: randomised controlled trial of preoperative optimisation of oxygen delivery. *BMJ* 1999;**318**:1099–103. [PUBMED: 10213716]

Ziegler 1997 {published data only}

Ziegler DW, Wright JG, Choban PS, Flancbaum L. A prospective randomized trial of preoperative “optimization” of cardiac function in patients undergoing elective peripheral vascular surgery. *Surgery* 1997;**122**:584–92. [PUBMED: 9308617]

References to studies excluded from this review

Alia 1999 {published data only}

Alia I, Esteban A, Gordo F, Lorente JA, Diaz C, Rodriguez JA, et al. A randomized and controlled trial of the effect of treatment aimed at maximizing oxygen delivery in patients with severe sepsis or septic shock. *Chest* 1999;**115**:453–61. [PUBMED: 10027447]

Balogh 2003 {published data only}

Balogh Z, McKinley BA, Cocanour CS, Kozar RA, Valdivia A, Sailors RM, et al. Supranormal trauma resuscitation causes more cases of abdominal compartment syndrome. *Archives of Surgery* 2003;**138**(6):637–43. [PUBMED: 12799335]

Benes 2010 {published data only}

Benes J, Chytra I, Altmann P, Hluchy M, Kasal E, Svitak R, et al. Intraoperative fluid optimization using stroke volume variation in high risk surgical patients: results of a prospective randomised study. *Critical Care* 2010;**14**:R118. [PUBMED: 20553586]

Bishop 1995 {published data only}

Bishop MH, Shoemaker WC, Appel PL, Meade P, Ordog GJ, Wasserberger J, et al. Prospective, randomized trial of survivor values of cardiac index, oxygen delivery, and oxygen consumption as resuscitation endpoints in severe trauma. *The Journal of Trauma* 1995;**38**:780–7. [PUBMED: 7760409]

Blow 1999 {published data only}

Blow O, Magliore L, Claridge JA, Butler K, Young JS. The golden hour and the silver day: detection and correction of occult hypoperfusion within 24 hours improves outcome

from major trauma. *The Journal of Trauma* 1999;**47**:964–9. [PUBMED: 10568731]

Chang 2000 {published data only}

Chang MC, Meredith JW, Kincaid EH, Miller PR. Maintaining survivors’ values of left ventricular power output during shock resuscitation: a prospective pilot study. *The Journal of Trauma* 2000;**49**:26–33. [PUBMED: 10912854]

Chytra 2007 {published data only}

Chytra I, Pradl R, Bosman R, Pelnar P, Kasal E, Zidkova A. Esophageal doppler-guided fluid management decreases blood lactate levels in multiple-trauma patients: a randomized controlled trial. *Critical Care* 2006;**11**:R24. [PUBMED: 17313691]

Durham 1996 {published data only}

Durham RM, Neunaber K, Mazuski JE, Shapiro MJ, Baue AE. The use of oxygen consumption and delivery as endpoints for resuscitation in critically ill patients. *The Journal of Trauma* 1996;**41**:32–9. [PUBMED: 8676421]

Flancbaum 1998 {published data only}

Flancbaum L, Ziegler DW, Choban PS. Preoperative intensive care unit admission and haemodynamic monitoring in patients scheduled for major elective noncardiac surgery: a retrospective review of 95 patients. *Journal of Cardiothoracic and Vascular Anesthesia* 1998;**12**:3–9. [PUBMED: 9509349]

Fleming 1992 {published data only}

Fleming A, Bishop M, Shoemaker W, Appel P, Sufficool W, Kuvhenguwaha A, et al. Prospective trial of supranormal values as goals of resuscitation in severe trauma. *Archives of Surgery* 1992;**127**:1175–9. [PUBMED: 1417482]

Forget 2010 {published data only}

Forget P, Lois F, Kock M. Goal-directed fluid management based on the pulse oximeter-derived pleth variability index reduces lactate levels and improves fluid management. *Anesthesia and Analgesia* 2010;**111**:910–4. [PUBMED: 20705785]

Gattinoni 1995 {published data only}

Gattinoni L, Brazzi L, Pelosi P, Latini R, Tognoni G, Pesenti A, et al. A trial of goal-oriented haemodynamic therapy in critically ill patients. SvO₂ Collaborative Group. *The New England Journal of Medicine* 1995;**333**(16):1025–32. [PUBMED: 7675044]

Gutierrez 1992 {published data only}

Gutierrez G, Palizas F, Doglio G, Wainsztein N, Galesio A, Pacin J, et al. Gastric intramucosal pH as a therapeutic index of tissue oxygenation in critically ill patients. *Lancet* 1992;**339**:195–9. [PUBMED: 1346170]

Hayes 1994 {published data only}

Hayes MA, Timmins AC, Yau EH, Palazzo M, Hinds CJ, Watson D. Elevation of systemic oxygen delivery in the treatment of critically ill patients. *The New England Journal of Medicine* 1994;**330**:1717–22. [PUBMED: 7993413]

Ivatury 1996 {published data only}

Ivatury RR, Simon RJ, Islam S, Fueg A, Rohman M, Stahl WM. A prospective randomized study of end points of

- resuscitation after major trauma: global oxygen transport indices versus organ specific gastric mucosal pH. *Journal of the American College of Surgery* 1996;**183**:145–54. [PUBMED: 8696546]
- Jammer 2010** {published data only}
Jammer IB, Ulvik A, Erichsen C, Lodemel O, Ostgaard G. Does central venous oxygen saturation- directed fluid therapy affect postoperative morbidity after colorectal surgery. *Anesthesiology* 2010;**113**:1072–80. [PUBMED: 20885291]
- Jorgensen 2009** {published data only}
Jorgensen CC, Bundgaard-Nielsen M, Stovgaard LT, Secher NH, Kehlet H. Stroke volume averaging for individualized goal-directed fluid therapy with oesophageal doppler. *Acta Anaesthesiologica Scandinavica* 2008;**53**:34–8. [PUBMED: 19032566]
- Lobo 2006** {published data only}
Lobo SM, Lobo FR, Polachini CA, Patini DS, Yamamoto AE, de Oliveira NE, et al. Prospective, randomized trial comparing fluids and dobutamine optimization of oxygen delivery in high-risk surgical patients. *Critical Care* 2006;**10**(3):R72. [PUBMED: 16696864]
- Miller 1998** {published data only}
Miller PR, Meredith JW, Chang MC. Randomized, prospective comparison of increased preload versus inotropes in the resuscitation of trauma patients: effects on cardiopulmonary function and visceral perfusion. *The Journal of Trauma* 1998;**44**:107–13. [PUBMED: 9464757]
- Muller 1999** {published data only}
Muller M, Boldt J, Schindler E, Sticher J, Kelm C, Roth S, et al. Effects of low-dose dopexamine on splanchnic oxygenation during major abdominal surgery. *Critical Care Medicine* 1999;**27**:2389–93. [PUBMED: 10579253]
- Pargger 1998** {published data only}
Pargger H, Hampl KF, Christen P, Staender S, Scheidegger D. Gastric intramucosal pH-guided therapy in patients after elective repair of infrarenal abdominal aneurysms: is it beneficial?. *Intensive Care Medicine* 1998;**24**:769–76. [PUBMED: 9757919]
- Rivers 2001** {published data only}
Rivers E, Nguyen B, Havstad S, Ressler J, Muzzin A, Knoblich B, et al. Early goal-directed therapy in the treatment of severe sepsis and septic shock. *The New England Journal of Medicine* 2001;**345**(19):1368–77. [PUBMED: 11794169]
- Scalea 1990** {published data only}
Scalea TM, Simon HM, Duncan AO, Atweh NA, Sclafani SJ, Phillips TF, et al. Geriatric blunt multiple trauma: improved survival with early invasive monitoring. *The Journal of Trauma* 1990;**30**:129–34. [PUBMED: 2304107]
- Schilling 2004** {published data only}
Schilling T, Grundling M, Strang CM, Moritz KU, Siegmund W, Hachenberg T. Effects of dopexamine, dobutamine or dopamine on prolactin and thyrotropin serum concentrations in high-risk surgical patients. *Intensive Care Medicine* 2004;**30**(6):1127–33. [PUBMED: 15138671]
- Schultz 1985** {published data only}
Schultz RJ, Whitfield GF, LaMura JJ, Raciti A, Krishnamurthy S. The role of physiologic monitoring in patients with fractures of the hip. *The Journal of Trauma* 1985;**25**:309–16. [PUBMED: 3989888]
- Stewart 2009** {published data only}
Stewart RM, Park PK, Hunt JP, McIntyre Jr RC, McCarthy J, Zarzabal LA, et al. Less is more: improved outcomes in surgical patients with conservative fluid administration and central venous catheter monitoring. *Journal of American College of Surgeons* 2009;**208**:725–37. [PUBMED: 19476825]
- Stone 2003** {published data only}
Stone MD, Wilson RJT, Cross J, Williams BT. Effect of adding dopexamine to intraoperative volume expansion in patients undergoing major elective abdominal surgery. *British Journal of Anaesthesia* 2003;**91**(5):619–24. [PUBMED: 14570781]
- Szakmany 2005** {published data only}
Szakmany T, Toth I, Kovacs Z, Leiner T, Mikor A, Koszegi T, et al. Effects of volumetric vs. pressure-guided fluid therapy on postoperative inflammatory response: a prospective, randomized clinical trial. *Intensive Care Medicine* 2005;**31**(5):656–63. [PUBMED: 15812629]
- Takala 2000** {published data only}
Takala J, Meier-Hellmann A, Eddleston J, Hulstaert P, Sramek V. Effect of dopexamine on outcome after major abdominal surgery: a prospective, randomized, controlled multicenter study. European Multicenter Study Group on Dopexamine in Major Abdominal Surgery. *Critical Care Medicine* 2000;**28**:3417–23. [PUBMED: 11057795]
- Tuchschmidt 1992** {published data only}
Tuchschmidt J, Fried J, Astiz M, Rackow E. Elevation of cardiac output and oxygen delivery improves outcome in septic shock. *Chest* 1992;**102**:216–20. [PUBMED: 1623756]
- Velmahos 2000** {published data only}
Velmahos GC, Demetriades D, Shoemaker WC, Chan LS, Tatevossian R, Wo CC, et al. Endpoints of resuscitation of critically injured patients: normal or supranormal? A prospective randomized trial. *Annals of Surgery* 2000;**232**:409–18. [PUBMED: 10973391]
- Wenkui 2010** {published data only}
Wenkui Y, Ning L, Jianfeng G, WeiQin L, ShaoQiu T, Zhihui T, et al. Restricted peri-operative fluid administration adjusted by serum lactate level improved outcome after major elective surgery for gastrointestinal malignancy. *Surgery* 2010;**147**:542–52. [PUBMED: 20004445]
- Yu 1993** {published data only}
Yu M, Levy MM, Smith P, Takiguchi SA, Miyasaki A, Myers SA. Effect of maximizing oxygen delivery on morbidity and mortality rates in critically ill patients: a prospective,

- randomized, controlled study. *Critical Care Medicine* 1993;**21**:830–8. [PUBMED: 8504649]
- Yu 1995** *{published data only}*
Yu M, Takanishi D, Myers SA, Takiguchi SA, Severino R, Hasaniya N, et al. Frequency of mortality and myocardial infarction during maximizing oxygen delivery: a prospective, randomized trial. *Critical Care Medicine* 1995;**23**:1025–32. [PUBMED: 774212]
- Yu 1998** *{published data only}*
Yu M, Burchell S, Hasaniya NW, Takanishi DM, Myers SA, Takiguchi SA. Relationship of mortality to increasing oxygen delivery in patients > or = 50 years of age: a prospective, randomized trial. *Critical Care Medicine* 1998;**26**:1011–9. [PUBMED: 9635648]
- ## References to ongoing studies
- Pearse 2009** *{unpublished data only}*
Pearse R. Optimization of perioperative cardiovascular management to improve surgical outcome. Not published.
- ## Additional references
- Bland 1978**
Bland R, Shoemaker WC, Shabot MM. Physiologic monitoring goals for the critically ill patient. *Surgery, Gynecology & Obstetrics* 1978;**147**(6):833–41. [PUBMED: 31006]
- Boyd 1959**
Boyd AR, Tremblay RE, Spencer FC, Bahnson HT. Estimation of cardiac output soon after cardiac surgery with cardiopulmonary bypass. *Annals of Surgery* 1959;**150**: 613–25.
- Boyd 1996**
Boyd O, Bennett ED. Enhancement of perioperative tissue perfusion as a therapeutic strategy for major surgery. *New Horizons* 1996;**4**(4):453–65. [PUBMED: 8968978]
- Boyd 1999**
Boyd O, Hayes M. The oxygen trail: the goal. *British Medical Bulletin* 1999;**55**(1):125–39. [PUBMED: 10695083]
- Brienza 2009**
Brienza N, Giglio MT, Marucci M, Fiore T. Does perioperative haemodynamic optimization protect renal function in surgical patients? A meta-analytic study. *Critical Care Medicine* 2009;**37**:2079–90. [PUBMED: 19384211]
- Clowes 1960**
Clowes GH Jr, Del Guercio LR. Circulatory response to trauma of surgical operations. *Metabolism* 1960;**9**(1): 67–81. [PUBMED: 13810744]
- Egger 1997**
Egger M, Smith GD, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997;**315**:629.
- Fenwick 2002**
Fenwick E, Wilson J, Sculpher M, Claxton K. Pre-operative optimisation employing dopexamine or adrenaline for patients undergoing major elective surgery: a cost-effectiveness analysis. *Intensive Care Medicine* 2002;**28**(5): 599–608. [PUBMED: 12029409]
- Forst 1997**
Forst H. [Maximizing O₂-transport in critical illness. A rational therapeutic concept?]. *Anaesthesist* 1997;**46**(1): 46–52. [PUBMED: 9082869]
- Ganz 1971**
Ganz W, Donoso R, Marcus HS, Forrester JS, Swan HJ. A new technique for measurement of cardiac output by thermodilution in man. *American Journal of Cardiology* 1971;**27**(4):392–6. [PUBMED: 4929422]
- Giglio 2009**
Giglio MT, Marucci M, Testini M, Brienza N. Goal-directed haemodynamic therapy and gastrointestinal complications in major surgery: a meta-analysis of randomized controlled trials. *British Journal of Anaesthesia* 2009;**103**:637–46. [MEDLINE: 19837807]
- Guest 1997**
Guest JF, Boyd O, Hart WM, Grounds RM, Bennett ED. A cost analysis of a treatment policy of a deliberate perioperative increase in oxygen delivery in high risk surgical patients. *Intensive Care Medicine* 1997;**23**(1): 85–90. [PUBMED: 9037645]
- Hamilton 2011**
Hamilton MA, Cecconi M, Rhodes A. A systematic review and meta-analysis on the use of preemptive haemodynamic intervention to improve post operative outcomes in moderate and high risk surgical patients. *Anesthesia and Analgesia* 2011;**112**:1392–402. [PUBMED: 20966436]
- Harbord 2006**
Harbord RM, Egger M, Sterne JA. A modified test for small-study effects in meta-analyses of controlled trials with binary endpoints. *Statistics in Medicine* 2006;**25**(20): 3443–57. [PUBMED: 16345038]
- Heyland 1996**
Heyland DK, Cook DJ, King D, Kernerman P, Brun-Buisson C. Maximizing oxygen delivery in critically ill patients: a methodologic appraisal of the evidence [see comments]. *Critical Care Medicine* 1996;**24**(3):517–24. [PUBMED: 8625644]
- Higgins 2011**
Higgins JPT, Altman DG, Sterne JAC (Editors). Chapter 8. Assessing risk of bias in included studies. In Higgins JPT; Green S, editors. *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.0.1 [updated March 2011]. The Cochrane Collaboration. Available from www.cochrane-handbook.org 2011.
- Hozo 2005**
Hozo SP, Djulbegovic B, Hozo I. Estimating the mean and variance from the median, range, and the size of a sample. *BMC Medical Research Methodology* 2005;**5**:13. [PUBMED: 15840177]

Ivanov 1997

Ivanov RI, Allen J, Sandham JD, Calvin JE. Pulmonary artery catheterization: a narrative and systematic critique of randomized controlled trials and recommendations for the future. *New Horizons* 1997;**5**(3):268–76. [PUBMED: 9259342]

Kern 2002

Kern JW, Shoemaker WC. Meta-analysis of haemodynamic optimization in high-risk patients. *Critical Care Medicine* 2002;**30**(8):1686–92. [PUBMED: 12163777]

Kjaergard 2001

Kjaergard LL, Villumsen J, Gluud C. Reported methodologic quality and discrepancies between large and small randomized trials in meta-analyses. *Annals of Internal Medicine* 2001;**135**:982–9. [PUBMED: 11730399]

Kusano 1997

Kusano C, Baba M, Takao S, Sane S, Shimada M, Shirao K, et al. Oxygen delivery as a factor in the development of fatal postoperative complications after oesophagectomy. *British Journal of Surgery* 1997;**84**(2):252–7. [PUBMED: 9052449]

Langham 2002

Langham J, Thompson E, Rowan K. Randomised controlled trials from the critical care literature: identification and assessment of quality. *Clinical Intensive Care* 2002;**13**: 73–83.

Leibowitz 1997

Leibowitz AB, Beilin Y. Pulmonary artery catheters and outcome in the perioperative period. *New Horizons* 1997;**5**(3):214–21. [PUBMED: 9259333]

Louis 1991

Louis TA. Assessing, accommodating, and interpreting the influences of heterogeneity. *Environmental Health Perspectives* 1991;**90**:215–22. [PUBMED: 2050064]

Moher 1998

Moher D, Pham B, Jones A, Cook DJ, Jadad AR, Moher M, et al. Does quality of reports of randomised trials affect estimates of intervention efficacy reported in meta-analyses? *Lancet* 1998;**352**:609–13. [PUBMED: 9746022]

MRC 2000

MRC (UK). A framework for development and evaluation of RCTs for complex interventions to improve health. MRC Publications <http://www.mrc.ac.uk/Utilities/Documentrecord/index.htm?d=MRC003372> (assessed September 2012) 2000.

Mythen 1994

Mythen MG, Webb AR. Intra-operative gut mucosal hypoperfusion is associated with increased post-operative complications and cost. *Intensive Care Medicine* 1994;**20**(2):99–104. [PUBMED: 8201106]

Peerless 1998

Peerless JR, Alexander JJ, Pinchak AC, Piotrowski JJ, Malangoni MA. Oxygen delivery is an important predictor of outcome in patients with ruptured abdominal aortic

aneurysms. *Annals of Surgery* 1998;**227**(5):726–32. [PUBMED: 9605664]

Poeze 2005

Poeze M, Greve JW, Ramsay G. Meta-analysis of haemodynamic optimization: relationship to methodological quality. *Critical Care* 2005;**9**(6):R771–9. [PUBMED: 16356226]

Polonen 1997

Polonen P, Hippelainen M, Takala R, Ruokonen E, Takala J. Relationship between intra- and postoperative oxygen transport and prolonged intensive care after cardiac surgery: a prospective study. *Acta Anaesthesiologica Scandinavica* 1997;**41**(7):810–7. [PUBMED: 9265921]

RevMan 5.1 [Computer program]

The Nordic Cochrane Centre, The Cochrane Collaboration. Review Manager (RevMan). Version 5.1. Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration, 2011.

Shoemaker 1972

Shoemaker WC. Cardiorespiratory patterns of surviving and nonsurviving postoperative patients. *Surgery, Gynecology & Obstetrics* 1972;**134**(5):810–4. [PUBMED: 5031495]

Shoemaker 1973

Shoemaker WC, Montgomery ES, Kaplan E, Elwyn DH. Physiologic patterns in surviving and nonsurviving shock patients. Use of sequential cardiorespiratory variables in defining criteria for therapeutic goals and early warning of death. *Archives of Surgery* 1973;**106**(5):630–6. [PUBMED: 4701410]

Shoemaker 1979

Shoemaker WC, Czer LS. Archives of Surgery. *Critical Care Medicine* 1979;**7**(9):424–31.

Shoemaker 1982

Shoemaker WC, Appel PL, Waxman K, Schwartz S, Chang P. Clinical trial of survivors' cardiorespiratory patterns as therapeutic goals in critically ill postoperative patients. *Critical Care Medicine* 1982;**10**(6):398–403. [PUBMED: 7042206]

Shoemaker 1993

Shoemaker WC, Appel PL, Kram HB. Hemodynamic and oxygen transport responses in survivors and non survivors of high-risk surgery [see comments]. *Critical Care Medicine* 1993;**21**(7):977–90. [PUBMED: 8319478]

Swan 1970

Swan HJ, Ganz W, Forrester J, Marcus H, Diamond G, Chonette D. Catheterization of the heart in man with use of a flow-directed balloon-tipped catheter. *The New England Journal of Medicine* 1970;**283**(9):447–51. [PUBMED: 5434111]

References to other published versions of this review**Grocott 2013**

Grocott MP, Dushianthan A, Hamilton MA, Mythen MG, Harrison D, Rowan K, Optimisation Systematic Review Steering Group. Perioperative increase in global

blood flow to explicit defined goals and outcomes after surgery: a Cochrane Systematic Review. *British Journal of Anaesthesia* 2013;**May 9**:[Epub ahead of print]. [PUBMED: 23661403]

* *Indicates the major publication for the study*

CHARACTERISTICS OF STUDIES

Characteristics of included studies [ordered by study ID]

Bender 1997

Methods	RCT Single centre	
Participants	104 patients Elective surgery Vascular surgery	
Interventions	Two groups Preoperative Fluids and Inotropes Pulmonary artery catheter Goals = CI	
Outcomes	Hospital mortality Hospital length of stay (HLOS) ICU length of stay (ICULOS) Cost Pulmonary oedema Acute myocardial infarction Arrhythmia Acute renal failure Wound infection Haemorrhage Sepsis Graft thrombosis or infection Groin haematoma	
Notes	Number of arrhythmia episodes were reported in both intra and postoperative periods, but not clear it occurred in same patients or different patients. Number of patients with complications only reported in postoperative period	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	High risk	Patients were “assigned randomly to one of two groups by the surgical intensivist”
Allocation concealment (selection bias)	Unclear risk	Method of concealment is not described
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not detailed

Bender 1997 (Continued)

Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not detailed
Incomplete outcome data (attrition bias) All outcomes	Low risk	All outcome data are reported
Selective reporting (reporting bias)	Low risk	Published report included main outcome measures
Other bias	Low risk	No other specific bias identified

Berlauk 1991

Methods	RCT Single centre	
Participants	89 patients Elective surgery Vascular surgery	
Interventions	3 groups Preoperative Fluids and inotropes Pulmonary artery catheter Goals = CI	
Outcomes	Hospital mortality HLOS ICULOS Cost Acute renal failure Congestive cardiac failure Graft thrombosis Acute myocardial infarction Arrhythmia	
Notes	3 Groups, where group-1 had PA catheter driven protocol initiated in surgical intensive care unit (SICU) 12 hours before surgery, group-2 had PA catheter driven protocol initiated 3 hours before surgery and group-3 control group without protocol. Arrhythmia episodes were reported, not clear of denominator. The analysis is combined for groups 1 and 2	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"Randomization for all study groups was generated by a random number generator (Statworks)"

Berlauk 1991 (Continued)

Allocation concealment (selection bias)	Unclear risk	Method of concealment is not described
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not detailed
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not detailed
Incomplete outcome data (attrition bias) All outcomes	Low risk	No missing outcome data
Selective reporting (reporting bias)	Low risk	Report included all expected outcomes
Other bias	Unclear risk	Two patients were excluded from group 1 due to not having surgery. This may have induced selection bias

Bonazzi 2002

Methods	RCT Single centre
Participants	100 patients Elective surgery Vascular surgery
Interventions	2 groups Preoperative Fluids and inotropes Pulmonary artery catheter Goals = CI, DO ₂ I
Outcomes	Hospital mortality HLOS Arrhythmias Myocardial infarction Congestive heart failure Renal failure
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"A computer generated random number was obtained by phone call to the statistical centre of the hospital"

Bonazzi 2002 (Continued)

Allocation concealment (selection bias)	Unclear risk	Allocation concealment is not presented
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not detailed
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not detailed
Incomplete outcome data (attrition bias) All outcomes	Low risk	No missing outcome data
Selective reporting (reporting bias)	Low risk	Published report included all expected outcomes
Other bias	Low risk	No other sources of bias identified

Boyd 1993

Methods	RCT Single centre	
Participants	107 patients 2 groups Elective and emergency surgery General surgery Vascular surgery	
Interventions	Preoperative and postoperative Fluids and inotropes Pulmonary artery catheter Goals = DO ₂ I	
Outcomes	28 day mortality HLOS ICULOS (Cost reported separately) Respiratory failure, acute renal failure, sepsis, cardiorespiratory arrest, pulmonary oedema, pleural fluid, wound infection, disseminated intravascular coagulation, acute myocardial infarction, abdominal abscess, haemorrhage, gastric outlet obstruction, cerebrovascular accident, pulmonary embolism, chest infection, psychosis, distal ischaemias	
Notes		
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement

Boyd 1993 (Continued)

Random sequence generation (selection bias)	Unclear risk	"Patients were randomly allocated to a protocol or control limb of the study". No information provided regarding random sequence generation
Allocation concealment (selection bias)	Unclear risk	Allocation concealment was not presented in the paper
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	"Neither the anaesthesiologist nor the surgeon was aware of a patient's allocation". However its not clear if the investigators were blinded
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not sure, see quote above
Incomplete outcome data (attrition bias) All outcomes	Low risk	No missing outcome data
Selective reporting (reporting bias)	Low risk	All anticipated outcomes were reported
Other bias	Unclear risk	The study was funded by a pharmaceutical company

Cecconi 2011

Methods	RCT Single centre
Participants	2 groups 40 patients Elective Total hip arthroplasty under regional anaesthesia
Interventions	Intraoperative Fluids and inotropes FloTrac sensor/Vigileo Goal = SV, DO ₂ I
Outcomes	Mortality Complications Renal failure Arrhythmias Infections Anaemia Pulmonary embolism
Notes	
<i>Risk of bias</i>	

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"Randomized with concealed envelope technique"
Allocation concealment (selection bias)	Low risk	Concealed envelope technique
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"Patients enrolled and the clinical teams caring for the patients were blinded"
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not detailed
Incomplete outcome data (attrition bias) All outcomes	Low risk	No missing outcome data
Selective reporting (reporting bias)	Low risk	All anticipated outcomes were reported
Other bias	Low risk	No other bias identified

Challand 2012

Methods	RCT Single centre
Participants	2 Groups 179 patients Elective Major colorectal surgery
Interventions	Intraoperative Fluids Oesophageal doppler Goal = SV
Outcomes	Surgical readiness for discharge Total postop days flatus passed, bowel movement, tolerance of food Any deviation from postoperative course Serious postop complications Critical care admission Readmission <30 days Mortality

Challand 2012 (Continued)

Notes	Patients were initially screened with CPET and was classified into two groups (aerobically fit and unfit as defined by anaerobic threshold) with subsequent randomization into control and intervention arms	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomization by “random block allocation using sequentially numbered opaque sealed envelopes”
Allocation concealment (selection bias)	Low risk	As per quote above
Blinding of participants and personnel (performance bias) All outcomes	Low risk	“Group allocation, ODM readings and algorithm -guided colloid administration were concealed from other staff in the operating theatre by screens”
Blinding of outcome assessment (detection bias) All outcomes	Low risk	“The investigator had no involvement in perioperative decision-making or postoperative care”
Incomplete outcome data (attrition bias) All outcomes	Low risk	No missing outcome data
Selective reporting (reporting bias)	Low risk	All anticipated outcomes were reported
Other bias	Low risk	No other sources of bias identified

Conway 2002

Methods	RCT 2 centres
Participants	2 groups 57 patients Elective surgery General surgery
Interventions	Intraoperative Fluids Oesophageal Doppler Goals = SV, FTc
Outcomes	Hospital mortality HLOS ICULOS Tolerating food

Conway 2002 (Continued)

Notes		
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	"Prior to induction of anaesthesia, patients were individually randomized into doppler or control". Although the study was randomized, the method of random sequence generation was not presented
Allocation concealment (selection bias)	Unclear risk	Allocation concealment was not mentioned in the paper
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"Medical and nursing staff were unaware of randomization"
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	See quote above, blinding of outcome assessment was not explained
Incomplete outcome data (attrition bias) All outcomes	Low risk	No missing outcome data
Selective reporting (reporting bias)	Low risk	Published report included all expected outcomes
Other bias	Low risk	No other source of bias identified

Donati 2007

Methods	RCT Multicentre study
Participants	135 patients Elective abdominal surgery
Interventions	Intraoperative Fluids and inotropes Goals = Oxygen extraction (O_2ER)
Outcomes	New postoperative organ failure Number of organ failures during ICU stay ICULOS HLOS Hospital mortality
Notes	

<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"Randomization was based on a permuted- block algorithm"
Allocation concealment (selection bias)	Low risk	"Patients were randomized to one of the two groups of treatment by a telephone system"
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not explained
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not explained
Incomplete outcome data (attrition bias) All outcomes	Low risk	No missing data
Selective reporting (reporting bias)	Unclear risk	All anticipated outcomes were reported
Other bias	Unclear risk	No other sources of bias identified

Gan 2002

Methods	RCT Single centre
Participants	100 patients Elective surgery General surgery
Interventions	Intraoperative Fluids Oesophageal Doppler Goals = SV, FTc
Outcomes	Hospital mortality HLOS Cost renal dysfunction, respiratory support for >24 hours, cardiovascular (hypotension, pulmonary oedema, arrhythmia), chest infection, severe PONV requiring rescue antiemetics, coagulopathy, wound infection, food tolerance
Notes	

<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"The patients were randomized into either protocol or control group using a random number generator in sealed envelopes"
Allocation concealment (selection bias)	Low risk	"See quote above"
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	"While the data were collected by independent dedicated research personnel not involved in the intra-operative management of patients, we were unable to blind the anaesthesiologists as to the treatment group"
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	See quote above
Incomplete outcome data (attrition bias) All outcomes	Low risk	No missing outcome data
Selective reporting (reporting bias)	Low risk	The anticipated outcomes were published
Other bias	Low risk	The study appears free from other sources of bias

Jerez 2001

Methods	RCT Single centre	
Participants	390 patients Elective surgery Cardiac surgery	
Interventions	Postoperative Fluids and Inotropes Pulmonary artery catheter Goals = SVO ₂ , CI	
Outcomes	Hospital mortality Organ failures	
Notes	Published in Spanish	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement

Jerez 2001 (Continued)

Random sequence generation (selection bias)	Unclear risk	“Se trata de un estudio prospectivo, intervencional, aleatorizado y controlado” “This is a prospective, interventional, randomized controlled study”. However the method of sequence generation was not mentioned
Allocation concealment (selection bias)	Unclear risk	See quote above
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not detailed
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not detailed
Incomplete outcome data (attrition bias) All outcomes	Low risk	No missing data
Selective reporting (reporting bias)	Low risk	Published report include all expected outcomes
Other bias	Low risk	The study seems free from other sources of bias

Jhanji 2010

Methods	RCT Single centre
Participants	135 patients (3 arms) 45 patients in each arm Elective Gastrointestinal surgery
Interventions	Postoperative 3 arms to the study (arm-1 CVP group fluid optimisation according to CVP, arm-2 SV fluid optimization according to SV, third arm had fluids and continuous dopexamine infusion) LidCO Goals = SV
Outcomes	Microvascular flow and oxygenation Complications Infections Acute kidney injury Cardiac complications Critical care free days HLOS Hospital mortality

Jhanji 2010 (Continued)

Notes	The 3rd arm with continuous dopexamine infusion is not included in the analysis as the control group was not matched	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	“Participants were randomly allocated to one of three treatment groups by computer-generated random sequence in blocks of nine”
Allocation concealment (selection bias)	Low risk	“The study group allocations were placed in serially numbered opaque envelopes”
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	“Only the member of the research team who delivered the intervention was aware of the study group allocation”
Blinding of outcome assessment (detection bias) All outcomes	Low risk	“Clinical outcomes data for each patient were collected by a member of the research team who was unaware of study group allocation”
Incomplete outcome data (attrition bias) All outcomes	Low risk	No missing outcome data
Selective reporting (reporting bias)	Low risk	Published report includes all expected outcomes
Other bias	Unclear risk	Study was supported by pharmaceutical company

Kapoor 2007

Methods	RCT Single centre
Participants	30 patients 2 groups Coronary artery bypass surgery
Interventions	Postoperative Fluids and Inotropes FloTrac Goals = CI, SVV
Outcomes	Hospital mortality HLOS ICULOS Duration of ventilation Duration of inotropic days

Kapoor 2007 (Continued)

	Arrhythmia Renal failure	
Notes		
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	"The patients were divided randomly into two groups, namely control and EGDT groups, by sealed envelope technique"
Allocation concealment (selection bias)	Low risk	See quote above
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not detailed
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not detailed
Incomplete outcome data (attrition bias) All outcomes	Low risk	No missing outcome data
Selective reporting (reporting bias)	Low risk	Report included all expected outcomes
Other bias	Unclear risk	Three patients excluded after allocation. There is a possibility of selection bias

Lobo 2000

Methods	RCT Single centre
Participants	37 patients 2 groups Elective surgery General surgery Vascular surgery
Interventions	Intra and postoperative Fluids and inotropes Pulmonary artery catheter Goals = DO ₂ I
Outcomes	28 day and 60 day mortality HLOS ICULOS

Lobo 2000 (Continued)

	Sepsis, shock, septic shock, cardiogenic shock, nosocomial infection, acute pancreatitis, postoperative fistula, arrhythmia, cerebrovascular accident, deep vein thrombosis, gastrointestinal bleeding, hypothermia, sepsis-related organ failure assessment (SOFA) score, bronchopneumonia, urinary tract infection, wound infection. ventilator days, organ dysfunction	
Notes	Both groups had PAC and protocol group aimed to achieve supranormal DO ₂ I	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	“Patients were randomized at the preoperative stage by means of sealed opaque envelopes”
Allocation concealment (selection bias)	Low risk	See quote above
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not detailed
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not detailed
Incomplete outcome data (attrition bias) All outcomes	Low risk	No missing outcome data
Selective reporting (reporting bias)	Low risk	Published report included all expected outcomes
Other bias	Low risk	No other potential source of bias identified

Mayer 2010

Methods	RCT Single centre
Participants	60 patients 2 groups Elective surgery Major abdominal surgery
Interventions	Intraoperative Fluid and Inotropes FloTrac/Vigileo Goal = CI, SV (SVI,SVV)

Mayer 2010 (Continued)

Outcomes	HLOS Complications ICULOS Mortality	
Notes	The second author for this study is currently under investigation for research misconduct in relation to published studies of fluid therapy	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	“Patients were randomized preoperatively either into a standard protocol group or an enhanced goal directed haemodynamic monitoring group using a closed envelope system”
Allocation concealment (selection bias)	Low risk	See quote above
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	“Both groups were managed by the same physicians on the same wards who were not involved in the intraoperative management, data collection or group allocation of the study”
Blinding of outcome assessment (detection bias) All outcomes	Low risk	“All data collected by a study nurse blinded to the study design and group allocation”
Incomplete outcome data (attrition bias) All outcomes	Low risk	No missed outcome data
Selective reporting (reporting bias)	Low risk	All anticipated outcomes are reported
Other bias	High risk	The second author for this study is currently under investigation for research misconduct in relation to published studies of fluid therapy

Mckendry 2004

Methods	RCT Single centre
Participants	174 patients Elective surgery Emergency surgery Cardiac surgery

Mckendry 2004 (Continued)

Interventions	Postoperative Fluids and inotropes Oesophageal Doppler Goals = SVI	
Outcomes	Hospital mortality HLOS ICULOS Atrial fibrillation requiring treatment, pneumothorax, Cerebrovascular accident, chest infection or sternal wound infection, GI bleed, acute renal failure, pleural effusion, infected leg wound, aortic regurgitation	
Notes		
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	“Patients were randomized by a priori computer generated sequence”
Allocation concealment (selection bias)	Low risk	“The study nurse opened the serially numbered, sealed opaque envelopes on arrival of patients”
Blinding of participants and personnel (performance bias) All outcomes	Low risk	“Both patients and staff on the general wards to which patients were sent after discharge from intensive care were unaware of the group assignment”
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not detailed
Incomplete outcome data (attrition bias) All outcomes	Low risk	No missing outcome data
Selective reporting (reporting bias)	Low risk	Published report includes all expected outcomes
Other bias	Unclear risk	Possibility of selection bias as 4 patients were excluded after allocation

Mythen 1995

Methods	RCT Single centre
Participants	60 patients 2 groups Elective surgery

Mythen 1995 (Continued)

	Cardiac surgery
Interventions	Intraoperative Fluids Oesophageal Doppler Goals = SV
Outcomes	Hospital mortality HLOS ICULOS (cost reported separately) Knaus organ failure criteria, chest infection, pleural effusion, disorientation, respiratory failure, nausea and vomiting, cerebrovascular accident, paralytic ileus, pericardial effusion
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	"Patients were randomized according to the contents of a sealed envelope to either a control group or a protocol group"
Allocation concealment (selection bias)	Low risk	See quote above
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not detailed
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not detailed
Incomplete outcome data (attrition bias) All outcomes	Low risk	No missing outcome data
Selective reporting (reporting bias)	Low risk	All anticipated outcomes are reported
Other bias	Unclear risk	The study was funded by a pharmaceutical company

Noblett 2006

Methods	RCT Single centre
Participants	103 patients 2 groups Elective surgery

Noblett 2006 (Continued)

	General surgery	
Interventions	Intraoperative Fluids Goals = SV, FTc	
Outcomes	Hospital mortality HLOS ICULOS Surgical fitness for discharge, return of gastrointestinal function, flatus, bowel movement, food tolerance, readmission rate, cytokine markers of the systemic inflammatory response	
Notes		
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	“Double blind randomized controlled trial”; the method of randomization not mentioned
Allocation concealment (selection bias)	Unclear risk	See above quote, the method of allocation concealment not mentioned
Blinding of participants and personnel (performance bias) All outcomes	Low risk	“Doppler probe insertion and monitoring, and trial fluid administration were carried out by a medically qualified researcher who had no involvement in postoperative patient care or decision making”
Blinding of outcome assessment (detection bias) All outcomes	Low risk	See quote above
Incomplete outcome data (attrition bias) All outcomes	Low risk	No missing data
Selective reporting (reporting bias)	Low risk	All outcomes data were presented
Other bias	Unclear risk	Five patients excluded after allocation may have induced selection bias

Pearse 2005

Methods	RCT Single centre
Participants	122 patients Elective surgery Emergency surgery

	Vascular surgery General surgery Urology surgery
Interventions	Postoperative Fluids and inotropes LidCO Goals = DO ₂ I
Outcomes	Hospital, 28 and 60 day mortality HLOS ICULOS Number of patients with complications, infection (pneumonia, abdominal, urinary tract, CVC, wound), respiratory (pleural effusion, pneumothorax, pulmonary embolism, ARDS), cardiovascular (arrhythmia, pulmonary oedema, MI, stroke), abdominal (<i>C. diff</i> , diarrhoea, acute bowel obstruction, upper GI bleed, paralytic ileus, anastomotic leak, intra-abdominal hypertension), postoperative massive haemorrhage
Notes	
<i>Risk of bias</i>	
Bias	Authors' judgement Support for judgement
Random sequence generation (selection bias)	Low risk "Patients were assigned to GDT or control groups by computer-generated random sequence"
Allocation concealment (selection bias)	Low risk "Study group assignments were placed in serially numbered opaque envelopes"
Blinding of participants and personnel (performance bias) All outcomes	Low risk "These treatments were administered by a member of the research team who was the only individual aware of study group allocation"
Blinding of outcome assessment (detection bias) All outcomes	Low risk The outcome assessment was blinded
Incomplete outcome data (attrition bias) All outcomes	Low risk No missing data
Selective reporting (reporting bias)	Low risk All potential outcomes data were presented
Other bias	Low risk No other sources of bias

Pillai 2011

Methods	Randomized controlled study
Participants	66 patients Patients undergoing radical cystectomy
Interventions	2 groups Oesophageal Doppler Goal = SV, FTc
Outcomes	Length of hospital stay ICU admission Wound dehiscence wound infection GI function (ileus, time to flatus, bowel opening) Postop nausea and vomiting

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	"All patients were randomized into a control arm, which received standard intraoperative fluids at the discretion of the consultant anaesthetist, and an intervention arm which received additional fluid from a researcher via a oesophageal doppler determined protocol" . However the method of random sequence was not mentioned
Allocation concealment (selection bias)	Unclear risk	See above quote, allocation concealment was not mentioned
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"Anaesthetic and surgical teams were blinded to the randomization"
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Blinding of outcome assessment was not mentioned
Incomplete outcome data (attrition bias) All outcomes	Low risk	No missing data
Selective reporting (reporting bias)	Low risk	All potential outcomes data were presented
Other bias	Low risk	No other sources of bias

Pölonen 2000

Methods	RCT Single centre
Participants	393 patients 2 groups Elective surgery Cardiac surgery
Interventions	Postoperative Fluids and Inotropes Pulmonary artery catheter Goals = SVO ₂ , lactate
Outcomes	28 day, 6 month and 12 month mortality HLOS ICULOS Organ dysfunctions: central nervous system (haemiplegia, stroke, Glasgow coma scale (GCS <10), circulatory (vasoactive medication or intraaortic counterpulsation to treat hypotension or low cardiac output), respiratory (need for mechanical or assisted ventilation), renal (low urine output or increased creatinine), hepatic (increased liver enzymes or bilirubin), gastrointestinal (macroscopic bleeding or paralytic ileus), haematological (low white cell or platelet count) ICU readmission
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	"Patients were randomized the day before surgery by sealed envelope technique"
Allocation concealment (selection bias)	Low risk	See quote above
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	"Although study was randomized, it was not blinded and the data of the oxygen transport measurements of both groups were open to those taking care of patients during the study period"
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	See quote above
Incomplete outcome data (attrition bias) All outcomes	Low risk	No missing data
Selective reporting (reporting bias)	Low risk	Published report included all expected outcomes
Other bias	Unclear risk	10 patients were excluded after allocation may have induced selection bias

Sandham 2003

Methods	RCT Multicentre	
Participants	1994 patients Elective surgery Emergency surgery General surgery Thoracic surgery Vascular surgery Hip fracture surgery	
Interventions	Preoperative Fluids and inotropes Pulmonary artery catheter Goals = DO ₂ I, CI	
Outcomes	Hospital, 6 month and 12 month mortality HLOS Myocardial infarction, congestive heart failure, supraventricular tachycardia, pulmonary embolism, renal insufficiency, hepatic insufficiency, sepsis from central venous catheter (CVC) or pulmonary artery catheter (PAC), wound infection, pneumonia, adverse events related to PAC or CVC: pulmonary infarction, haemothorax, pulmonary haemorrhage, pneumothorax, arterial puncture	
Notes		
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	“Randomization was carried out by computer-generated sequence, stratified according to type of surgery and according to ASA class and blocked according to centre”
Allocation concealment (selection bias)	Low risk	“Assignments were concealed in opaque sealed envelopes”
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not blinded as it was not feasible
Blinding of outcome assessment (detection bias) All outcomes	Low risk	The outcome assessment was blinded
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Loss of patients follow-up at 6 months in both groups
Selective reporting (reporting bias)	Low risk	All expected outcomes were reported

Sandham 2003 (Continued)

Other bias	Unclear risk	Supported by a healthcare company
------------	--------------	-----------------------------------

Senagore 2009

Methods	RCT Single centre
Participants	64 patients 3 arms study Elective Laparoscopic colectomy
Interventions	Intraoperative Fluids Oesophageal doppler Goals = SV increase 10%
Outcomes	Hospital Mortality HLOS Arrhythmia infections
Notes	This is a 3-arm study with a control group in the first arm and goal directed groups in arms 2 and 3, where patients received lactated Ringer's solution and 6% HES

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"Randomization was performed after EDM insertion via a established random number table"
Allocation concealment (selection bias)	Unclear risk	Allocation concealment was not presented
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"A separate anaesthesia team monitored the EDM readings and administered boluses, whereas the clinical anaesthesia was blinded to this process"
Blinding of outcome assessment (detection bias) All outcomes	Low risk	See quote above
Incomplete outcome data (attrition bias) All outcomes	Low risk	No missing data
Selective reporting (reporting bias)	Low risk	All expected outcomes were reported

Other bias	Low risk	No other sources of bias identified
------------	----------	-------------------------------------

Shoemaker 1988

Methods	RCT Single centre
Participants	88 patients Elective surgery Emergency surgery General surgery
Interventions	Preoperative Fluids and inotropes Pulmonary artery catheter Goals = CI, DO ₂ I
Outcomes	Hospital mortality HLOS ICULOS Cost Respiratory failure, renal failure, sepsis and septic shock, hepatic failure, cardiac arrest, pulmonary oedema, pleural effusion, wound infection, disseminated intravascular coagulation (DIC), acute myocardial infarction, evisceration, abdominal abscess, haemorrhage, pancreatitis, gastric outlet obstruction, urinary tract infection, cerebral infarct, pulmonary embolism, ventilator days
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"The remaining 88 patients were prospectively allocated to one of the three groups designated by cards arranged according to a random numbers table by an outside person and placed in opaque sealed envelopes"
Allocation concealment (selection bias)	Low risk	See quote above
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not detailed
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not detailed

Shoemaker 1988 (Continued)

Incomplete outcome data (attrition bias) All outcomes	Low risk	No missing data
Selective reporting (reporting bias)	Low risk	All anticipated outcomes were reported
Other bias	Low risk	No other sources of bias identified

Sinclair 1997

Methods	RCT Single centre
Participants	40 patients Emergency surgery Hip fracture surgery
Interventions	Intraoperative Fluids oesophageal Doppler Goals = SV
Outcomes	Hospital mortality HLOS None, "time declared fit for medical discharge"
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	"After consent had been obtained, the patients were individually randomized before induction of anaesthesia by a sealed envelope technique to either protocol or control groups"
Allocation concealment (selection bias)	Low risk	See quote above
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"The anaesthetist was blinded to the doppler measurements but was aware of the fluid volumes given as fluid challenges to the protocol group" " postoperative management was carried out on the orthopaedic ward with both medical and nursing staff blinded to the randomization"
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not detailed

Sinclair 1997 (Continued)

Incomplete outcome data (attrition bias) All outcomes	Low risk	No missing data
Selective reporting (reporting bias)	Low risk	All expected outcomes were reported
Other bias	Low risk	No other sources of bias identified

Ueno 1998

Methods	RCT Single centre
Participants	34 patients Elective surgery Liver surgery
Interventions	Postoperative Fluids and inotropes Pulmonary artery catheter Goals = CI, DO ₂ I, VO ₂ I
Outcomes	Hospital mortality Bleeding, peritoneal infection, adult respiratory distress syndrome, hyperbilirubinaemia, liver failure
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"The patients were randomly assigned to either group N or group S by the stratified block randomization method"
Allocation concealment (selection bias)	Unclear risk	Not specified
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not detailed
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	"All measurements were obtained and recorded by one of the authors"
Incomplete outcome data (attrition bias) All outcomes	Low risk	No missing data
Selective reporting (reporting bias)	Low risk	All anticipated outcomes were reported

Ueno 1998 (Continued)

Other bias	Low risk	No other sources of bias identified
------------	----------	-------------------------------------

Valentine 1998

Methods	RCT Single centre
Participants	120 patients Elective surgery Vascular surgery
Interventions	Preoperative Fluids and inotropes Pulmonary artery catheter Goals = CI
Outcomes	Hospital mortality HLOS ICULS Myocardial infarction, arrhythmia, congestive heart failure, pneumonia, non-cardiogenic pulmonary insufficiency, acute renal insufficiency, catheter sepsis, ventilator days
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	"Patients were randomized into two groups by means of a sealed envelope"
Allocation concealment (selection bias)	Low risk	See quote above
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not detailed
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not detailed
Incomplete outcome data (attrition bias) All outcomes	Low risk	No missing data
Selective reporting (reporting bias)	Low risk	All expected outcomes were published
Other bias	Low risk	No other sources of bias identified

Van der Linden 2010

Methods	RCT Single centre
Participants	37 patients 3-arm study High risk (ASA II-III) Peripheral artery bypass grafting
Interventions	Intraoperative Fluids and Inotropes (dobutamine) FloTrac Vigileo monitor Goal = CI
Outcomes	Mortality HLOS Complications (infections)
Notes	The study comprised of 3 groups. Group 1: controls, Group 2: goal directed therapy and Group 3: goal directed therapy with different anaesthetic compared to group 1 and 2. As Group 3 was not matched with controls this was not included in the analysis

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"Patients were randomly assigned to three groups using computer generated random code"
Allocation concealment (selection bias)	Low risk	"The assignment was concealed in an envelope"
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"The attending anaesthesiologist was blinded for the data displayed by the Vigileo monitor. Cardiac index data in these patients were recorded by an independent observer who was not involved in the patients treatment"
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Data were analysed blinded
Incomplete outcome data (attrition bias) All outcomes	Low risk	No missing data
Selective reporting (reporting bias)	Low risk	All expected outcomes were reported
Other bias	Low risk	No other potential sources of bias identified

Venn 2002

Methods	RCT Single centre
Participants	90 patients Emergency surgery Hip fracture surgery
Interventions	Intraoperative Fluids Oesophageal Doppler Goals = SV
Outcomes	Hospital mortality HLOS Time to medical fitness for discharge, deep haemorrhage requiring >2 unit blood transfusion, haematemesis, chest infection, wound infection, cellulitis, pancreatitis, pulmonary embolus, cerebrovascular accident, myocardial infarction, cardiac failure, rapid atrial fibrillation, hypotension, impaired renal function, pseudo-obstruction
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"Patients were individually randomized into 3 groups, through the use of a set of computer generated random numbers and an opaque sealed envelope technique"
Allocation concealment (selection bias)	Low risk	See above quote
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"The anaesthetist was unaware of the central venous pressure or oesophageal doppler ultrasonography measurements" "Post operative management was performed by the orthopaedic medical team and nursing staff who were all unaware of the patient's randomization"
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	See above quote
Incomplete outcome data (attrition bias) All outcomes	Low risk	No missing data
Selective reporting (reporting bias)	Low risk	All expected outcomes were presented
Other bias	Low risk	No other potential sources of bias identified

Wakeling 2005

Methods	RCT Single centre
Participants	134 patients Elective surgery General surgery
Interventions	Intraoperative Fluids Oesophageal Doppler Goals = SV
Outcomes	Hospital and 6 month mortality HLOS Time until fit for discharge, bowel recovery (flatus, bowels opening, full diet), quality of recovery score, postoperative morbidity survey (POMS), quality of life questionnaires (European organisation for the research and treatment of cancer (EORTC) - QLQ-C30 and QLQ-CR38)
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"Patients were randomized and allocated according to the sequentially numbered, sealed opaque envelope technique"
Allocation concealment (selection bias)	Low risk	See quote above
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"The surgical teams, nursing staff and patients themselves were blinded" "The anaesthetists was blinded to the oesophageal doppler measurements"
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Blinding of outcome assessment was not explained
Incomplete outcome data (attrition bias) All outcomes	Low risk	No missing data
Selective reporting (reporting bias)	Low risk	All anticipated outcomes were reported
Other bias	Unclear risk	Three patients in each group did not receive allocated intervention, this may have induced selection bias

Wilson 1999

Methods	RCT single centre
Participants	138 patients 3 groups Elective surgery General surgery Vascular surgery
Interventions	Preoperative Fluids and inotropes Pulmonary artery catheter Goals = DO ₂ I
Outcomes	Hospital mortality HLOS ICULOS (Costs reported separately) Respiratory (prolonged weaning, adult respiratory distress syndrome (ARDS), pleural effusion, secondary ventilation, sputum retention), cardiovascular (myocardial infarction, arrhythmia, cardiac arrest, pulmonary embolus, cerebrovascular accident, transient ischaemic attack, cardiac failure), gastrointestinal (infarction, haemorrhage), acute renal failure, coagulopathy, infection (bacteraemia, sepsis syndrome, septic shock, respiratory sepsis, urinary sepsis, abdominal sepsis, wound sepsis, line sepsis, other sepsis), surgical (anastomotic breakdown, deep haemorrhage, wound haemorrhage)
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"Random sequence was generated from a Unix computer program"
Allocation concealment (selection bias)	Low risk	"Allocation was concealed until trial entry by sealed opaque envelope"
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	"We were unable to effect true blinding between patients in the control and treatment groups"
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	See quote above
Incomplete outcome data (attrition bias) All outcomes	Low risk	No missing data

Wilson 1999 (Continued)

Selective reporting (reporting bias)	Low risk	All anticipated outcome data were reported
Other bias	Low risk	No other potential sources of bias

Ziegler 1997

Methods	RCT Single centre
Participants	72 patients Elective surgery Vascular surgery
Interventions	Preoperative Fluids and inotropes Pulmonary artery catheter Goals = SVO ₂
Outcomes	Hospital mortality ICULOS Hypotension, congestive heart failure, myocardial infarction, arrhythmia, oliguria, graft thrombosis, cerebrovascular accident
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Method of random sequence was not presented
Allocation concealment (selection bias)	Unclear risk	This was not presented
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not detailed
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not detailed
Incomplete outcome data (attrition bias) All outcomes	Low risk	No missing data
Selective reporting (reporting bias)	Low risk	All expected outcomes were reported

Other bias	Low risk	No other sources of bias identified
------------	----------	-------------------------------------

Characteristics of excluded studies [ordered by study ID]

Study	Reason for exclusion
Alia 1999	Severe sepsis, septic shock
Balogh 2003	Trauma
Benes 2010	Patients not had surgery also included in the analysis
Bishop 1995	Trauma
Blow 1999	Trauma
Chang 2000	Trauma, not RCT
Chytra 2007	Trauma
Durham 1996	Established critical illness
Flancbaum 1998	Retrospective, not RCT
Fleming 1992	Trauma
Forget 2010	Pulse pressure variation is not a surrogate of global blood flow
Gattinoni 1995	Established critical illness
Gutierrez 1992	pHi guided
Hayes 1994	Established critical illness
Ivatury 1996	Trauma
Jammer 2010	Restricted goal directed fluid administration
Jorgensen 2009	Not RCT
Lobo 2006	Same flow goal in each group
Miller 1998	Trauma

(Continued)

Muller 1999	No explicit flow goal
Pargger 1998	pHi guided
Rivers 2001	Severe sepsis and septic shock
Scalea 1990	Trauma
Schilling 2004	Same flow goal in each group
Schultz 1985	No explicit flow goals
Stewart 2009	Retrospective study
Stone 2003	No explicit flow goals
Szakmany 2005	Intrathoracic blood volume goal
Takala 2000	No explicit flow goals
Tuchschmidt 1992	Septic shock
Velmahos 2000	Trauma
Wenkui 2010	Restricted fluid regimen
Yu 1993	Established critical illness
Yu 1995	Established critical illness
Yu 1998	Established critical illness

Characteristics of ongoing studies [ordered by study ID]

Pearse 2009

Trial name or title	Optimization of perioperative cardiovascular management to improve surgical outcome
Methods	Randomized controlled study
Participants	Adults undergoing major abdominal surgery
Interventions	
Outcomes	28 day mortality, postoperative complications, postoperative morbidity survey score, LOHS, critical care free days, cost

Pearse 2009 (Continued)

Starting date	2009
Contact information	ISRCTN04386758
Notes	Ongoing study

DATA AND ANALYSES

Comparison 1. Mortality

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 All studies (longest follow-up)	31	5292	Risk Ratio (IV, Fixed, 95% CI)	0.89 [0.76, 1.05]
2 All studies (hospital or 28 day)	31	5292	Risk Ratio (IV, Fixed, 95% CI)	0.81 [0.65, 1.00]
3 Participant numbers	31	5292	Risk Ratio (IV, Fixed, 95% CI)	0.89 [0.76, 1.05]
3.1 >100	16	4428	Risk Ratio (IV, Fixed, 95% CI)	0.94 [0.79, 1.12]
3.2 <100	15	864	Risk Ratio (IV, Fixed, 95% CI)	0.51 [0.29, 0.90]
4 Time intervention started	31		Risk Ratio (IV, Fixed, 95% CI)	Subtotals only
4.1 Pre-operative	9	2786	Risk Ratio (IV, Fixed, 95% CI)	0.96 [0.79, 1.17]
4.2 Intra-operative	15	1202	Risk Ratio (IV, Fixed, 95% CI)	0.67 [0.40, 1.13]
4.3 Post-operative	9	1341	Risk Ratio (IV, Fixed, 95% CI)	0.73 [0.50, 1.06]
5 Type of intervention	31		Risk Ratio (IV, Fixed, 95% CI)	Subtotals only
5.1 Fluids and Inotropes	21	4354	Risk Ratio (IV, Fixed, 95% CI)	0.90 [0.76, 1.07]
5.2 Fluids	11	983	Risk Ratio (IV, Fixed, 95% CI)	0.80 [0.46, 1.39]
6 Goals of intervention	31	5292	Risk Ratio (IV, Fixed, 95% CI)	0.89 [0.76, 1.05]
6.1 CO, DO2	14	3060	Risk Ratio (IV, Fixed, 95% CI)	0.91 [0.75, 1.09]
6.2 Lactate, SVO2, O2ER	4	990	Risk Ratio (IV, Fixed, 95% CI)	0.83 [0.50, 1.38]
6.3 SV	13	1242	Risk Ratio (IV, Fixed, 95% CI)	0.84 [0.51, 1.41]
7 Mode of surgery	31	5292	Risk Ratio (IV, Fixed, 95% CI)	0.89 [0.76, 1.05]
7.1 Elective	24	2677	Risk Ratio (IV, Fixed, 95% CI)	0.68 [0.48, 0.94]
7.2 Emergency	2	130	Risk Ratio (IV, Fixed, 95% CI)	0.68 [0.23, 2.06]
7.3 Elective and Emergency	5	2485	Risk Ratio (IV, Fixed, 95% CI)	0.99 [0.81, 1.20]
8 Type of surgery	27		Risk Ratio (IV, Fixed, 95% CI)	Subtotals only
8.1 Vascular	7	580	Risk Ratio (IV, Fixed, 95% CI)	0.78 [0.34, 1.79]
8.2 Cardiac	5	1047	Risk Ratio (IV, Fixed, 95% CI)	0.81 [0.48, 1.35]
8.3 General	16	1374	Risk Ratio (IV, Fixed, 95% CI)	0.66 [0.41, 1.07]

Comparison 2. Complications

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Renal impairment	21	4307	Risk Ratio (IV, Fixed, 95% CI)	0.71 [0.57, 0.90]
2 Arrhythmia	12	2921	Risk Ratio (IV, Fixed, 95% CI)	0.84 [0.67, 1.06]
3 Infection: numbers	9	733	Risk Ratio (IV, Fixed, 95% CI)	0.88 [0.69, 1.12]
4 Infections: types	16		Risk Ratio (IV, Fixed, 95% CI)	Subtotals only
4.1 Chest Infections/ Pneumonia	13	2945	Risk Ratio (IV, Fixed, 95% CI)	0.78 [0.61, 1.00]
4.2 Sepsis	5	474	Risk Ratio (IV, Fixed, 95% CI)	0.68 [0.26, 1.77]
4.3 Abdominal Infections	6	555	Risk Ratio (IV, Fixed, 95% CI)	0.53 [0.23, 1.22]
4.4 Wound Infections	10	2802	Risk Ratio (IV, Fixed, 95% CI)	0.65 [0.50, 0.84]
4.5 Urinary Tract Infections	8	612	Risk Ratio (IV, Fixed, 95% CI)	0.54 [0.26, 1.15]
5 Respiratory failure / ARDS	9	844	Risk Ratio (IV, Fixed, 95% CI)	0.51 [0.28, 0.93]

6 Myocardial infarction	15	3328	Risk Ratio (IV, Fixed, 95% CI)	1.01 [0.71, 1.45]
7 Congestive heart failure / pulmonary oedema	14	3223	Risk Ratio (IV, Fixed, 95% CI)	1.00 [0.81, 1.24]
8 Venous thrombosis	10	2740	Risk Ratio (IV, Fixed, 95% CI)	1.04 [0.39, 2.77]
9 Number of participants with complications	17	1841	Risk Ratio (IV, Random, 95% CI)	0.68 [0.58, 0.80]

Comparison 3. Resource utilization

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Length of hospital stay	27	4729	Mean Difference (IV, Random, 95% CI)	-1.16 [-1.89, -0.43]
2 Length of critical care stay	14	1873	Mean Difference (IV, Random, 95% CI)	-0.45 [-0.94, 0.03]

Comparison 4. Sensitivity analysis: excluding studies with active controls

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Mortality (Longest follow-up)	27	4408	Risk Ratio (IV, Fixed, 95% CI)	0.94 [0.79, 1.12]
2 Mortality (Hospital or 28 days)	27	4408	Risk Ratio (IV, Fixed, 95% CI)	0.84 [0.67, 1.07]
3 Renal Impairment	19	3847	Risk Ratio (IV, Fixed, 95% CI)	0.71 [0.56, 0.90]
4 Arrhythmia	11	2884	Risk Ratio (IV, Fixed, 95% CI)	0.85 [0.67, 1.07]
5 Infection (Number of patients with infections)	8	696	Risk Ratio (IV, Fixed, 95% CI)	0.91 [0.71, 1.16]
6 Respiratory Failure/ ARDS	8	780	Risk Ratio (IV, Fixed, 95% CI)	0.48 [0.26, 0.89]
7 Myocardial Infarction	15	3298	Risk Ratio (IV, Fixed, 95% CI)	1.01 [0.71, 1.45]
8 Congestive Heart Failure/ Pulmonary oedema	13	3156	Risk Ratio (IV, Fixed, 95% CI)	1.01 [0.82, 1.26]
9 Venous Thrombosis	9	2673	Risk Ratio (IV, Fixed, 95% CI)	1.19 [0.42, 3.31]
10 Number of patients with complications	14	1350	Risk Ratio (IV, Fixed, 95% CI)	0.71 [0.63, 0.80]
11 Length of Hospital Stay [Days]	25	4269	Mean Difference (IV, Random, 95% CI)	-1.21 [-2.08, -0.33]
12 Length of Critical Care Stay [Days]	11	1023	Mean Difference (IV, Random, 95% CI)	-0.32 [-0.89, 0.26]

Comparison 5. Sensitivity analysis: excluding poorly-controlled studies

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Mortality (Longest follow-up)	22	2586	Risk Ratio (IV, Fixed, 95% CI)	0.65 [0.48, 0.89]
2 Mortality (Hospital or 28 day)	22	2586	Risk Ratio (IV, Fixed, 95% CI)	0.66 [0.47, 0.92]
3 Renal Impairment	12	1601	Risk Ratio (IV, Fixed, 95% CI)	0.69 [0.47, 1.03]
4 Arrhythmia	5	469	Risk Ratio (IV, Fixed, 95% CI)	0.57 [0.30, 1.07]
5 Infection (Number of patients with infections)	7	509	Risk Ratio (IV, Fixed, 95% CI)	0.87 [0.68, 1.11]
6 Respiratory Failure/ ARDS	7	616	Risk Ratio (IV, Fixed, 95% CI)	0.55 [0.27, 1.10]
7 Myocardial Infarction	6	622	Risk Ratio (IV, Fixed, 95% CI)	0.66 [0.25, 1.73]
8 Congestive Heart Failure/ Pulmonary oedema	7	587	Risk Ratio (IV, Fixed, 95% CI)	0.64 [0.33, 1.26]
9 Venous Thrombosis	5	388	Risk Ratio (IV, Fixed, 95% CI)	0.50 [0.11, 2.32]
10 Number of patients with complications	14	1573	Risk Ratio (IV, Random, 95% CI)	0.66 [0.53, 0.81]
11 Length of Hospital Stay [Days]	18	2023	Mean Difference (IV, Random, 95% CI)	-1.61 [-2.85, -0.37]
12 Length of Critical Care Stay [Days]	9	1391	Mean Difference (IV, Random, 95% CI)	-0.74 [-1.43, -0.06]

Comparison 6. Sensitivity analysis: Sandham 2003

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Mortality (Longest Follow-up)-excluding patients lost follow-up	31	5139	Risk Ratio (IV, Fixed, 95% CI)	0.91 [0.78, 1.08]
2 Mortality (Longest Follow-up)-if all patients lost follow-up have died	31	5292	Risk Ratio (IV, Fixed, 95% CI)	1.02 [0.89, 1.18]

Analysis I.1. Comparison I Mortality, Outcome I All studies (longest follow-up).

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: I Mortality

Outcome: I All studies (longest follow-up)

(Continued ...)

Analysis 1.2. Comparison 1 Mortality, Outcome 2 All studies (hospital or 28 day).

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 1 Mortality

Outcome: 2 All studies (hospital or 28 day)

(Continued ...)

Analysis 1.3. Comparison 1 Mortality, Outcome 3 Participant numbers.

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 1 Mortality

Outcome: 3 Participant numbers

(Continued ...)

Analysis I.4. Comparison I Mortality, Outcome 4 Time intervention started.

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: I Mortality

Outcome: 4 Time intervention started

(Continued ...)

Analysis 1.5. Comparison 1 Mortality, Outcome 5 Type of intervention.

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 1 Mortality

Outcome: 5 Type of intervention

(Continued ...)

Analysis 1.6. Comparison 1 Mortality, Outcome 6 Goals of intervention.

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 1 Mortality

Outcome: 6 Goals of intervention

(Continued ...)

Analysis 1.7. Comparison 1 Mortality, Outcome 7 Mode of surgery.

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 1 Mortality

Outcome: 7 Mode of surgery

(Continued ...)

Analysis 1.8. Comparison 1 Mortality, Outcome 8 Type of surgery.

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 1 Mortality

Outcome: 8 Type of surgery

(Continued ...)

Analysis 2.1. Comparison 2 Complications, Outcome 1 Renal impairment.

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 2 Complications

Outcome: 1 Renal impairment

Analysis 2.2. Comparison 2 Complications, Outcome 2 Arrhythmia.

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 2 Complications

Outcome: 2 Arrhythmia

Analysis 2.3. Comparison 2 Complications, Outcome 3 Infection: numbers.

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 2 Complications

Outcome: 3 Infection: numbers

Analysis 2.4. Comparison 2 Complications, Outcome 4 Infections: types.

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 2 Complications

Outcome: 4 Infections: types

(Continued ...)

(... Continued)

Analysis 2.5. Comparison 2 Complications, Outcome 5 Respiratory failure / ARDS.

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 2 Complications

Outcome: 5 Respiratory failure / ARDS

Analysis 2.6. Comparison 2 Complications, Outcome 6 Myocardial infarction.

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 2 Complications

Outcome: 6 Myocardial infarction

Analysis 2.7. Comparison 2 Complications, Outcome 7 Congestive heart failure / pulmonary oedema.

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 2 Complications

Outcome: 7 Congestive heart failure / pulmonary oedema

Analysis 2.8. Comparison 2 Complications, Outcome 8 Venous thrombosis.

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 2 Complications

Outcome: 8 Venous thrombosis

Analysis 2.9. Comparison 2 Complications, Outcome 9 Number of participants with complications.

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 2 Complications

Outcome: 9 Number of participants with complications

Analysis 3.1. Comparison 3 Resource utilization, Outcome 1 Length of hospital stay.

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 3 Resource utilization

Outcome: 1 Length of hospital stay

(Continued ...)

Analysis 3.2. Comparison 3 Resource utilization, Outcome 2 Length of critical care stay.

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 3 Resource utilization

Outcome: 2 Length of critical care stay

(Continued ...)

Analysis 4.1. Comparison 4 Sensitivity analysis: excluding studies with active controls, Outcome 1 Mortality (Longest follow-up).

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 4 Sensitivity analysis: excluding studies with active controls

Outcome: 1 Mortality (Longest follow-up)

(Continued ...)

Analysis 4.2. Comparison 4 Sensitivity analysis: excluding studies with active controls, Outcome 2 Mortality (Hospital or 28 days).

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 4 Sensitivity analysis: excluding studies with active controls

Outcome: 2 Mortality (Hospital or 28 days)

(Continued ...)

Analysis 4.3. Comparison 4 Sensitivity analysis: excluding studies with active controls, Outcome 3 Renal Impairment.

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 4 Sensitivity analysis: excluding studies with active controls

Outcome: 3 Renal Impairment

(Continued ...)

Analysis 4.4. Comparison 4 Sensitivity analysis: excluding studies with active controls, Outcome 4 Arrhythmia.

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 4 Sensitivity analysis: excluding studies with active controls

Outcome: 4 Arrhythmia

Analysis 4.5. Comparison 4 Sensitivity analysis: excluding studies with active controls, Outcome 5 Infection (Number of patients with infections).

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 4 Sensitivity analysis: excluding studies with active controls

Outcome: 5 Infection (Number of patients with infections)

Analysis 4.6. Comparison 4 Sensitivity analysis: excluding studies with active controls, Outcome 6 Respiratory Failure/ ARDS.

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 4 Sensitivity analysis: excluding studies with active controls

Outcome: 6 Respiratory Failure/ ARDS

Analysis 4.7. Comparison 4 Sensitivity analysis: excluding studies with active controls, Outcome 7 Myocardial Infarction.

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 4 Sensitivity analysis: excluding studies with active controls

Outcome: 7 Myocardial Infarction

Analysis 4.8. Comparison 4 Sensitivity analysis: excluding studies with active controls, Outcome 8 Congestive Heart Failure/ Pulmonary oedema.

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 4 Sensitivity analysis: excluding studies with active controls

Outcome: 8 Congestive Heart Failure/ Pulmonary oedema

Analysis 4.9. Comparison 4 Sensitivity analysis: excluding studies with active controls, Outcome 9 Venous Thrombosis.

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 4 Sensitivity analysis: excluding studies with active controls

Outcome: 9 Venous Thrombosis

Analysis 4.10. Comparison 4 Sensitivity analysis: excluding studies with active controls, Outcome 10 Number of patients with complications.

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 4 Sensitivity analysis: excluding studies with active controls

Outcome: 10 Number of patients with complications

Analysis 4.11. Comparison 4 Sensitivity analysis: excluding studies with active controls, Outcome 11 Length of Hospital Stay [Days].

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 4 Sensitivity analysis: excluding studies with active controls

Outcome: 11 Length of Hospital Stay [Days]

(Continued ...)

Analysis 4.12. Comparison 4 Sensitivity analysis: excluding studies with active controls, Outcome 12 Length of Critical Care Stay [Days].

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 4 Sensitivity analysis: excluding studies with active controls

Outcome: 12 Length of Critical Care Stay [Days]

(Continued ...)

(... Continued)

Analysis 5.1. Comparison 5 Sensitivity analysis: excluding poorly-controlled studies, Outcome 1 Mortality (Longest follow-up).

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 5 Sensitivity analysis: excluding poorly-controlled studies

Outcome: 1 Mortality (Longest follow-up)

(Continued ...)

Analysis 5.2. Comparison 5 Sensitivity analysis: excluding poorly-controlled studies, Outcome 2 Mortality (Hospital or 28 day).

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 5 Sensitivity analysis: excluding poorly-controlled studies

Outcome: 2 Mortality (Hospital or 28 day)

Analysis 5.3. Comparison 5 Sensitivity analysis: excluding poorly-controlled studies, Outcome 3 Renal Impairment.

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 5 Sensitivity analysis: excluding poorly-controlled studies

Outcome: 3 Renal Impairment

Analysis 5.4. Comparison 5 Sensitivity analysis: excluding poorly-controlled studies, Outcome 4 Arrhythmia.

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 5 Sensitivity analysis: excluding poorly-controlled studies

Outcome: 4 Arrhythmia

Analysis 5.5. Comparison 5 Sensitivity analysis: excluding poorly-controlled studies, Outcome 5 Infection (Number of patients with infections).

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 5 Sensitivity analysis: excluding poorly-controlled studies

Outcome: 5 Infection (Number of patients with infections)

Analysis 5.6. Comparison 5 Sensitivity analysis: excluding poorly-controlled studies, Outcome 6 Respiratory Failure/ ARDS.

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 5 Sensitivity analysis: excluding poorly-controlled studies

Outcome: 6 Respiratory Failure/ ARDS

Analysis 5.7. Comparison 5 Sensitivity analysis: excluding poorly-controlled studies, Outcome 7 Myocardial Infarction.

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 5 Sensitivity analysis: excluding poorly-controlled studies

Outcome: 7 Myocardial Infarction

Analysis 5.8. Comparison 5 Sensitivity analysis: excluding poorly-controlled studies, Outcome 8 Congestive Heart Failure/ Pulmonary oedema.

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 5 Sensitivity analysis: excluding poorly-controlled studies

Outcome: 8 Congestive Heart Failure/ Pulmonary oedema

Analysis 5.9. Comparison 5 Sensitivity analysis: excluding poorly-controlled studies, Outcome 9 Venous Thrombosis.

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 5 Sensitivity analysis: excluding poorly-controlled studies

Outcome: 9 Venous Thrombosis

Analysis 5.10. Comparison 5 Sensitivity analysis: excluding poorly-controlled studies, Outcome 10 Number of patients with complications.

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 5 Sensitivity analysis: excluding poorly-controlled studies

Outcome: 10 Number of patients with complications

Analysis 5.11. Comparison 5 Sensitivity analysis: excluding poorly-controlled studies, Outcome 11 Length of Hospital Stay [Days].

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 5 Sensitivity analysis: excluding poorly-controlled studies

Outcome: 11 Length of Hospital Stay [Days]

Analysis 5.12. Comparison 5 Sensitivity analysis: excluding poorly-controlled studies, Outcome 12 Length of Critical Care Stay [Days].

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 5 Sensitivity analysis: excluding poorly-controlled studies

Outcome: 12 Length of Critical Care Stay [Days]

Analysis 6.1. Comparison 6 Sensitivity analysis: Sandham 2003, Outcome 1 Mortality (Longest Follow-up)-excluding patients lost follow-up.

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 6 Sensitivity analysis: Sandham 2003

Outcome: 1 Mortality (Longest Follow-up)- excluding patients lost follow-up

(Continued ...)

Analysis 6.2. Comparison 6 Sensitivity analysis: Sandham 2003, Outcome 2 Mortality (Longest Follow-up)- if all patients lost follow-up have died.

Review: Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery

Comparison: 6 Sensitivity analysis: Sandham 2003

Outcome: 2 Mortality (Longest Follow-up)- if all patients lost follow-up have died

(Continued ...)

ADDITIONAL TABLES

Table 1. Study outcomes

Study	Mortality	Morbidity	Resource use	Cost
Bender 1997	Hospital	Pulmonary oedema, acute myocardial infarction, arrhythmia, acute renal failure, wound infection, haemorrhage, sepsis, graft thrombosis or infection, groin haematoma	HLOS, ICULOS	Cost
Berlauk 1991	Hospital	Acute renal failure, congestive cardiac failure, graft thrombosis, acute myocardial infarction, arrhythmia,	HLOS, ICULOS	Cost
Bonazzi 2002	Hospital	Arrhythmias, myocardial infarction, congestive heart failure, renal failure	HLOS	None

Table 1. Study outcomes (Continued)

Boyd 1993	28 day	Respiratory failure, acute renal failure, sepsis, cardiorespiratory arrest, pulmonary oedema, pleural fluid, wound infection, disseminated intravascular coagulation, acute myocardial infarction, abdominal abscess, haemorrhage, gastric outlet obstruction, cerebrovascular accident, pulmonary embolism, chest infection, psychosis, distal ischaemia	HLOS, ICULOS	Reported separately
Cecconi 2011	28 day	Infections, hypotension, anaemia, pneumonia, pulmonary embolism, tachyarrhythmias, acute coronary syndrome, acute renal failure	HLOS	None
Challand 2012	30 days 90 days	Serious postoperative complications, renal complications, creatinine increase, critical care admission	HLOS	None
Conway 2002	Hospital	Tolerating oral diet	HLOS, ICULOS	None
Donati 2007	Hospital	Organ failures	HLOS	None
Gan 2002	Hospital	Acute renal dysfunction (urine output <500mls), respiratory support for > 24 hours, cardiovascular (hypotension, pulmonary oedema, arrhythmia), chest infection (clinical diagnosis), severe postoperative nausea and vomiting requiring rescue antiemetic, coagulopathy, wound infection, toleration of oral solid diet	HLOS	None

Table 1. Study outcomes (Continued)

Jerez 2001	Hospital	Organ failures	ICULOS	None
Jhanji 2010	Hospital	Cardiac complications, infections, acute kidney injury	HLOS, ICULOS	None
Kapoor 2007	Hospital	arrhythmia, renal dysfunction, low cardiac output	HLOS, ICULOS	None
Lobo 2000	28 day, 60 day	Sepsis, shock, septic shock, cardiogenic shock, nosocomial infection, acute pancreatitis, postoperative fistula, arrhythmia, cerebrovascular accident, deep vein thrombosis, gastrointestinal bleeding, hypothermia, sepsis related organ failure assessment (SOFA) score, bronchopneumonia, urinary tract infection, wound infection, ventilator days, organ dysfunction	HLOS, ICULOS	None
Mayer 2010	Hospital	Infection (pneumonia, abdominal, urinary tract, wound), respiratory (PE, respiratory support), cardiovascular (pulmonary oedema, arrhythmia, hypotension, acute myocardial infarction, stroke), abdominal (bowel obstruction, gastrointestinal bleeding, anastomotic leak), renal (urine output <500ml/day or required dialysis for acute renal failure), post operative haemorrhage	HLOS, ICULOS	None
Mckendry 2004	Hospital mortality	Atrial fibrillation requiring treatment, pneumothorax, cerebrovascu-	HLOS, ICULOS	None

Table 1. Study outcomes (Continued)

		lar accident, chest infection or sternal wound infection, GI bleed, acute renal failure, pleural effusion, infected leg wound, aortic regurgitation		
Mythen 1995	Hospital	Knaus organ failure criteria, chest infection, pleural effusion, disorientation, respiratory failure, nausea and vomiting, cerebrovascular accident, paralytic ileus, pericardial effusion	HLOS, ICULOS	Reported separately
Noblett 2006	Hospital	surgical fitness for discharge, return of gastrointestinal function, flatus, bowel movement, food tolerance, readmission rate, cytokine markers of the systemic inflammatory response	HLOS, ICULOS	None
Pearse 2005	Hospital, 28 and 60 day mortality	Number of patients with complications, infection (pneumonia, abdominal, urinary tract, CVC, wound), respiratory (pleural effusion, pneumothorax, pulmonary embolism, ARDS), cardiovascular (arrhythmia, pulmonary oedema, MI, stroke), abdominal (<i>C. Diff</i> , diarrhoea, acute bowel obstruction, upper GI bleed, paralytic ileus, anastomotic leak, intra-abdominal hypertension), postoperative massive haemorrhage	HLOS, ICUOS	None
Pillai 2011	None	Nausea and vomiting, wound dehiscence, wound infection, ileus	HLOS	None

Table 1. Study outcomes (Continued)

Pölönen 2000	28 day, 6 month, 12 month	Organ dysfunctions: central nervous system (hemiplegia, stroke, Glasgow coma scale (GCS <10) , circulatory (vasoactive medication or intraaortic counterpulsation to treat hypotension or low cardiac output), respiratory (need for mechanical or assisted ventilation) , renal (low urine output or increased creatinine) , hepatic (increased liver enzymes or bilirubin), gastrointestinal (macroscopic bleeding or paralytic ileus), haematological (low white cell or platelet count), ICU readmission	HLOS, ICULOS	None
Sandham 2003	Hospital, 6 month, 12 month	Myocardial infarction, congestive heart failure, supraventricular tachycardia, pulmonary embolism, renal insufficiency, hepatic insufficiency, sepsis from central venous catheter (CVC) or pulmonary artery catheter (PAC), wound infection, pneumonia, adverse events related to PAC or CVC: pulmonary infarction, haemothorax, pulmonary haemorrhage, pneumothorax, arterial puncture	HLOS	None
Senagore 2009	Hospital	Complications: Gastrointestinal failure, blood pressure lability, arrhythmia, dehydration, electrolyte imbalance, hyperglycaemia, wound/infectious com-	HLOS	None

Table 1. Study outcomes (Continued)

		pli- cations, sepsis, DVT/PE, intraoperative hypother- mia, urinary dysfunction, respiratory dysfunction, abdominal pain, chest pain, bleeding, anaemia, altered mental status		
Shoemaker 1988	Hospital	Respiratory failure, renal failure, sepsis and sep- tic shock, hepatic failure, cardiac arrest, pulmonary edema, pleural effusion, wound infection, dissem- inated intravascular coag- ulation (DIC), acute my- ocardial infarction, evis- ceration, abdominal ab- scess, haemorrhage, pan- creati- tis, gastric outlet obstruc- tion, urinary tract infec- tion, cerebral infarct, pul- monary embolism, venti- lator days	HLOS, ICULOS	Cost
Sinclair 1997	Hospital	None, “time declared fit for medical discharge”	HLOS	None
Ueno 1998	Hospital	Bleeding, peritoneal in- fection, adult respiratory distress syndrome, hyper- bilirubinaemia, liver fail- ure	None	None
Valentine 1998	Hospital	Myocardial infarc- tion, arrhythmia, conges- tive heart failure, pneu- monia, non-cardiogenic pulmonary insufficiency, acute renal insufficiency, catheter sepsis. ventilator days	HLOS, ICULOS	None
Van der Linden 2010	Hospital	Blood loss, infection	HLOS	None
Venn 2002	Hospital	“Time to medical fit- ness for discharge”, deep	HLOS	None

Table 1. Study outcomes (Continued)

		haemorrhage requiring >2 unit blood transfusion, haematemesis, chest infection, wound infection, cellulitis, pancreatitis, pulmonary embolus, cerebrovascular accident, myocardial infarction, cardiac failure, rapid atrial fibrillation, hypotension, impaired renal function, pseudo-obstruction		
Wakeling 2005	hospital and 6 month mortality	Time until fit for discharge, bowel recovery (flatus, bowels opening, full diet), quality of recovery score, postoperative morbidity survey (POMS), quality of life questionnaires (European Organisation for the Research and Treatment of Cancer (EORTC) - QLQ-C30 and QLQ-CR38)	HLOS	None
Wilson 1999	Hospital	Respiratory (prolonged weaning, adult respiratory distress syndrome (ARDS), pleural effusion, secondary ventilation, sputum retention), cardiovascular (myocardial infarction, arrhythmia, cardiac arrest, pulmonary embolus, cerebrovascular accident, transient ischaemic attack, cardiac failure), gastrointestinal (infarction, haemorrhage), acute renal failure, coagulopathy, infection (bacteraemia, sepsis syndrome, septic shock, respiratory sepsis, urinary sepsis, abdomi-	HLOS, ICULOS	Reported separately

Table 1. Study outcomes (Continued)

		nal sepsis, wound sepsis, line sepsis, other sepsis), surgical (anastomotic breakdown, deep haemorrhage, wound haemorrhage)		
Ziegler 1997	Hospital	Hypotension, congestive heart failure, myocardial infarction, arrhythmia, oliguria, graft thrombosis, cerebrovascular accident	ICULOS	None

APPENDICES

Appendix 1. Search strategy for CENTRAL , The Cochrane Library

#1 (Vasoactive or Fluid* or Drug Administration or fluid therapy or starch or gelatin* or crystalloid* or colloid* or splanchnic* or pulmonary artery flotation or catheter* or PAFC or Swan Ganz or Doppler):ti,ab
 #2 ((fluid* near (load* or administrat*)) or (perfusion near (renal or tissue)))
 #3 base near (acid or excess or deficit)
 #4 Venous near (Oxygen Saturation)
 #5 ((Stroke Volume Index) or (Oxygen Consumption Index)):ti,ab
 #6 (oxygen near (delivery or consumption or saturation)):ti,ab
 #7 (cardiac near (output or index)):ti,ab
 #8 (lactat* or CVP or pHi or PCO2 or SvO2 or VO2 or DO2 or Tonometry):ti,ab
 #9 (#1 OR #2 OR #3 OR #4 OR #5 OR #6 OR #7 OR #8)
 #10 ((surg* or operat*) near (general or high?risk or vascular or cardiac or cancer or trauma* or emergency or orthopaed*)):ti,ab
 #11 (peri?operativ* or post?operativ* or intra?operativ* or optimi?ation or goal?directed or supra?normal or aneurysm):ti,ab
 #12 (#10 OR #11)
 #13 (#9 AND #12)

Appendix 2. Search strategy for MEDLINE (OvidSP)

1. Fluid-Therapy/ or Body-Fluids/ or Catheterization-Swan-Ganz/ or Catheterization/ or Heart-Catheterization/
 2. (Vasoactive or Fluid* or Drug Administration or fluid therapy or starch or gelatin* or crystalloid* or colloid* or splanchnic* or pulmonary artery flotation or catheter* or PAFC or Swan Ganz or Doppler).ti,ab.
 3. ((fluid* adj3 (load* or administrat*)) or (perfusion adj3 (renal or tissue))).mp.
 4. Blood-Volume/ or Oxygen-Consumption/ or Central-Venous-Pressure/ or Stroke-Volume/ or Cardiac-Output/ or Echocardiography/ or Echocardiography-Doppler/
 5. ((base adj3 (acid or excess or deficit)) or ((Venous adj3 Oxygen Saturation) or Stroke Volume Index or Oxygen Consumption Index)).mp. or ((oxygen adj3 (delivery or consumption or saturation)) or (cardiac adj3 (output or index)) or lactat* or CVP or pHi or PCO2 or SvO2 or VO2 or DO2 or Tonometry).ti,ab.
 6. 1 or 2 or 3 or 4 or 5

7. Perioperative-Care/ or Intraoperative-Period/ or Postoperative-Period/ or Aneurysm/ or Vascular-Surgical-Procedures/ or Thoracic-Surgery/ or Emergency-Treatment/ or Specialties-Surgical/ or Orthopedics/ or Surgical-Procedures-Operative/
8. ((surg* or operat*) adj3 (general or high?risk or vascular or cardiac or cancer or trauma* or emergency or orthopaed*)).ti,ab.
9. (peri?operativ* or post?operativ* or intra?operativ* or optimi?ation or goal?directed or supra?normal or aneurysm).ti,ab.
10. 8 or 7 or 9
11. 6 and 10
12. ((randomized controlled trial or controlled clinical trial).pt. or randomized.ab. or placebo.ab. or clinical trials as topic.sh. or randomly.ab. or trial.ti.) not (animals not (humans and animals)).sh.
13. 11 and 12

Appendix 3. Search strategy for EMBASE (OvidSP)

1. fluid therapy/ or body fluid/ or Swan Ganz catheter/ or heart catheterization/ or blood volume/ or oxygen consumption/ or central venous pressure/ or heart stroke volume/ or heart output/ or Doppler echocardiography/ or echocardiography/
2. (Vasoactive or Fluid* or Drug Administration or fluid therapy or starch or gelatin* or crystalloid* or colloid* or splanchnic* or pulmonary artery flotation or catheter* or PAFC or Swan Ganz or Doppler).ti,ab. or ((fluid* adj3 (load* or administrat*)) or (perfusion adj3 (renal or tissue))).mp.
3. ((base adj3 (acid or excess or deficit)) or ((Venous adj3 Oxygen Saturation) or Stroke Volume Index or Oxygen Consumption Index)).mp. or ((oxygen adj3 (delivery or consumption or saturation)) or (cardiac adj3 (output or index)) or lactate* or CVP or pH_i or PCO₂ or SvO₂ or VO₂ or DO₂ or Tonometry).ti,ab.
4. 1 or 2 or 3
5. perioperative period/ or postoperative period/ or aneurysm/ or vascular surgery/ or thorax surgery/ or emergency treatment/ or orthopedics/ or surgery/
6. ((surg* or operat*) adj3 (general or high?risk or vascular or cardiac or cancer or trauma* or emergency or orthopaed*)).ti,ab.
7. (peri?operativ* or post?operativ* or intra?operativ* or optimi?ation or goal?directed or supra?normal or aneurysm).ti,ab.
8. 6 or 7 or 5
9. 8 and 4
10. (placebo.sh. or controlled study.ab. or random*.ti,ab. or trial*.ti,ab.) not (animals not (humans and animals)).sh.
11. 10 and 9

Appendix 4. Keywords used in search strategy

We used the following topic specific key words :

Population: high-risk surgery, peri-operative, pre-operative, post-operative, intra-operative, aneurysm, vascular surgery, cardiac surgery, cancer surgery, trauma surgery, emergency surgery, orthopaedic surgery

Intervention: Optimisation, optimization, goal-directed, supra-normal, Vasoactive, fluids, starch, gelatin, blood product, crystalloid, colloid, fluid therapy, fluid loading, fluid administration, body fluid

Comparison: Oxygen delivery, lactate, acid base, oxygen consumption, base excess, base deficit, blood volume, central venous pressure, CVP, cardiac output, cardiac index, pulmonary artery flotation catheter, PAFC, right-heart catheter, Swan Ganz, Doppler, pH_i, tonometry, PCO₂ gap, echocardiography, stroke volume, SvO₂, mixed venous oxygen saturation.

Outcomes: Splanchnic, renal perfusion, tissue perfusion, blood flow.

Appendix 5. Included studies

Study and Year of Publication	Country	Journal
Bender 1997	USA	<i>Annals of Surgery</i>
Berlaak 1991	USA	<i>Annals of Surgery</i>
Bonazzi 2002	Italy	<i>European Journal of Vascular Surgery</i>
Boyd 1993	UK	<i>JAMA</i>
Cecconi 2011	Italy	<i>Critical Care</i>
Challand 2012	UK	<i>British Journal of Anaesthesia</i>
Conway 2002	UK	<i>Anaesthesia</i>
Donati 2007	Italy	<i>Chest</i>
Gan 2002	USA	<i>Anesthesiology</i>
Jerez 2001	Spain	<i>Medicina Intensiva</i>
Jhanji 2010	UK	<i>Critical Care</i>
Kapoor 2007	India	<i>Annals of Cardiac Anaesthesia</i>
Lobo 2000	Brazil	<i>Critical Care Medicine</i>
Mayer 2010	Germany	<i>Critical Care</i>
Mckendry 2004	UK	<i>BMJ</i>
Mythen 1995	UK	<i>Archives of Surgery</i>
Noblett 2006	UK	<i>The British Journal of Surgery</i>
Pearse 2005	UK	<i>Critical Care</i>
Pillai 2011	UK	<i>The Journal of Urology</i>
Pölonen 2000	Finland	<i>Anesthesia and Analgesia</i>
Sandham 2003	Canada	<i>New England Journal of Medicine</i>
Senagore 2009	USA	<i>Diseases of Colon and Rectum</i>

(Continued)

Shoemaker 1988	USA	<i>Chest</i>
Sinclair 1997	UK	<i>BMJ</i>
Ueno 1998	Japan	<i>Surgery</i>
Valentine 1998	USA	<i>Journal of Vascular Surgery</i>
Van der Linden 2010	Belgium	<i>European Journal of Anaesthesiology</i>
Venn 2002	UK	<i>British Journal of Anaesthesia</i>
Wakeling 2005	UK	<i>British Journal of Anaesthesia</i>
Wilson 1999	UK	<i>BMJ</i>
Ziegler 1997	USA	<i>Surgery</i>

Footnotes

Appendix 6. Characteristics of eligible studies

Study	Design	Study Population		Intervention		
	Patients	Modee of surgery	Type of surgery	Timing	Fluids +/- Inotropes	Goals
Bender 1997	104	elective	vascular	pre	fluids and inotropes	CI
Berlauk 1991	89	elective	vascular	pre	fluids and inotropes	CI
Bonazzi 2002	100	elective	vascular	pre	fluids and inotropes	CI, DO ₂ I
Boyd 1993	107	elective, emergency	general, vascular	pre, post	fluids and inotropes	DO ₂ I
Cecconi 2011	40	elective	orthopaedic	intra	fluids and inotropes	SV
Challand 2012	179	elective	gastrointestinal	intra	fluids	SV
Conway 2002	57	elective	general	intra	fluids	SV,FTc

(Continued)

Donati 2007	135	elective	major abdominal surgery	intra	fluids and inotropes	O ₂ ER
Gan 2002	100	elective	general	intra	fluids	SV, FTc
Jerez 2001	390	elective	cardiac	post	fluids and inotropes	SVO ₂ , CI
Jhanji 2010	135	elective	gastrointestinal surgery	post	fluids and inotropes	SV
Kapoor 2007	30	elective	cardiac	post	fluids and inotropes	CI, SVV
Lobo 2000	37	elective	general, vascular	pre	fluids and inotropes	DO ₂ I
Mayer 2010	60	elective	major abdominal surgery	intra	fluids and inotropes	CI, SV
Mckendry 2004	174	elective, emergency	cardiac	post	fluids and inotropes	SVI
Mythen 1995	60	elective	cardiac	intra	fluids	SV
Noblett 2006	103	elective	general	intra	fluids	SV, FTc
Pearse 2005	122	elective, emergency	vascular, general, urology	post	fluids and inotropes	DO ₂ I
Pillai 2011	66	elective	urology	intra	fluids	SV, FTc
Pölönen 2000	393	elective	cardiac	post	fluids and inotropes	SvO ₂ , lactate
Sandham 2003	1994	elective, emergency	general, vascular, thoracic, hip fracture	pre	fluids and inotropes	DO ₂ I, CI
Senagore 2009	64	elective	Laparoscopic colectomy	intra	fluids	SV
Shoemaker 1988	58	elective, emergency	general, vascular	pre	fluids and inotropes	CI, DO ₂ I, VO ₂ I
Sinclair 1997	40	emergency	hip fracture	intra	fluids	SV, FTc
Ueno 1998	34	elective	liver	post	fluids and inotropes	CI, DO ₂ I, VO ₂ I
Valentine 1998	120	elective	vascular	pre	fluids and inotropes	CI

(Continued)

Van der Linden 2010	57	elective	vascular	intra	fluids and inotropes	CI
Venn 2002	59	emergency	hip fracture	intra	fluids	SV, FTc
Wakeling 2005	134	elective	general	intra	fluids	SV
Wilson 1999	138	elective	general, vascular	pre	fluids and inotropes	DO ₂ I
Ziegler 1997	72	elective	vascular	pre	fluids and inotropes	SvO ₂

Footnotes

Appendix 7. Sensitivity analysis using analytical methods for the primary outcome (mortality for the longest follow-up)

Analytical method	Results
Inverse variance Relative Risk Fixed-effect model	0.89 (95% CI 0.76 to 1.05), P = 0.18, I ² = 15%
Inverse variance Relative Risk Random-effects model	0.72 (95% CI 0.55 to 0.95), P = 0.02, I ² = 15%
Inverse variance Odds Ratio Fixed-effects model	0.87 (95% CI 0.72 to 1.05), P = 0.14, I ² = 20%
Inverse variance Odd's Ratio Random-effects model	0.67 (95% CI 0.49 to 0.92), P = 0.01, I ² = 20%
Peto Odd's Ratio	0.83 (95% CI 0.69 to 1.00), P = 0.05, I ² = 37%
MH Odd's ratio Fixed-effect model	0.83 (95% CI 0.69 to 1.00), P = 0.05, I ² = 21%
MH Odd's ratio Random-effects model	0.67 (95% CI 0.49 to 0.92), P = 0.01, I ² = 21%
MH Relative Risk Fixed-effect model	0.85 (95% CI 0.73 to 1.00), P = 0.05, I ² = 16%
MH Relative Risk Random-effects model	0.72 (95% CI 0.55 to 0.95), P = 0.02, I ² = 16%

Appendix 8. Sensitivity analysis using analytical methods for hospital or 28 day mortality

Analytical method	Results
Inverse variance Relative Risk Fixed-effect model	0.81 (95% CI 0.65 to 1.00), P = 0.06, I ² = 1%
Inverse variance Relative Risk Random-effects model	0.79 (95% CI 0.63 to 0.99), P = 0.04, I ² = 1%
Inverse variance Odds Ratio Fixed-effect model	0.79 (95% CI 0.63 to 1.00), P = 0.05, I ² = 7%
Inverse variance Odd's Ratio Random-effects model	0.72 (95% CI 0.54 to 0.96), P = 0.02, I ² = 7%
Peto Odd's Ratio	0.75 (95% CI 0.60 to 0.94), P = 0.01, I ² = 26%
MH Odd's ratio Fixed-effect model	0.76 (95% CI 0.60 to 0.95), P = 0.01, I ² = 8%
MH Odd's ratio Random-effects model	0.72 (95% CI 0.54 to 0.96), P = 0.02, I ² = 8%
MH Relative Risk Fixed-effect model	0.77 (95% CI 0.63 to 0.95), P = 0.01, I ² = 2%
MH Relative Risk Random-effects model	0.78 (95% CI 0.62 to 0.99), P = 0.04, I ² = 2%

WHAT'S NEW

Last assessed as up-to-date: 1 March 2012.

Date	Event	Description
7 October 2016	Amended	New entry created in Published notes regarding status of Mayer 2010 study

HISTORY

Protocol first published: Issue 1, 2003

Review first published: Issue 11, 2012

Date	Event	Description
1 July 2013	Amended	Journal version of review (Grocott 2013) cited in ' Other published versions of this review '.
25 October 2012	Amended	Amendment to acknowledgment section: acknowledging University Hospital Southampton NHS Foundation Trust-University of Southampton Respiratory Biomedical Research Unit and University College London Hospital-University College London Comprehensive Biomedical Research Centre

(Continued)

1 February 2006	Amended	February 2006: The authors appealed against the original decision of the Cochrane Funders Arbiter. Their appeal was recently upheld by the Co-chairs of the Steering Group and the decision of the funding arbiter overturned. The “Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery” will be republished in The Cochrane Library in issue 2, 2006. The authors are working on the draft review
1 February 2005	Amended	February 2005: “Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery” was withdrawn from The Cochrane Library in issue 2, 2005 on the advice of the Cochrane Funders Arbiter. This was because the authors received funding from a commercial source (Elan Pharma) for the preparation of this review. This contravenes the current Cochrane policy on sponsorship

CONTRIBUTIONS OF AUTHORS

Co-ordinating the review: Mike Grocott (MG), Mark Hamilton (MH), Ahilanandan Dushianthan (AD)

Undertaking manual searches: MG, Jonathan Hayam (JH) and Shahzad Shaei (SS), AD

Screening search results: MG, MH, Michael Mythen (MM), AD, Anand Vibhakkar (AV)

Organizing retrieval of papers: MG, MH, AD, AV

Screening retrieved papers against inclusion criteria: MG, MH, AD

Appraising quality of papers: MG, MH, AD

Abstracting data from papers: MG, MH, AD

Writing to authors of papers for additional information: MG, MH, AD

Providing additional data about papers: authors of papers.

Obtaining and screening data on unpublished studies: NA

Data management for the review: MG, MH, David Harrison (DH), AD

Entering data into Review Manager ([RevMan 5.1](#)): MG, MH, AD

RevMan statistical data: MG, MH, AD

Other statistical analysis not using RevMan: DH, John Carlisle (JC)

Double entry of data: NA

Interpretation of data: MG, MH, AD, Kathy Rowan (KR)

Statistical inferences: MG, MH, AD, KR

Writing the review: MG, MH, AD, KR

Securing funding for the review: NA

Guarantor for the review (one author): MG

Person responsible for reading and checking review before submission: MG, AD

The Optimisation Systematic Review Steering Group are, in alphabetical order: Beale R, Boyd O, Emberton M, Langham J, Roberts I, Thompson J.

DECLARATIONS OF INTEREST

Dr Boyd and Professor Mythen are authors of included studies.

None known: Beale R, Emberton M, Langham J, Roberts I, Thompson J, Dushianthan A.

Received funding from industry for studies and lecturing in the area: Boyd O, Mythen M, Grocott M, Hamilton MA.

Pre-existing views

None: Emberton M, Langham J, Rowan K, Dushianthan A

Against: no-one.

Uncertain: Beale R, Roberts I, Thompson J.

In favour: Boyd O, Mythen M, Grocott M, Hamilton MA.

Professor Mike Grocott has received unrestricted grant funding (paid to his institution) not related to this review from John Caudwell, British Oxygen Company (Linde Gas Therapeutics), Smiths Medical Ltd., Ely-Lilly Critical Care Ltd., Deltex Medical Ltd., The London Clinic, UCL Business, Rolex Ltd., Association of Anaesthetists of Great Britain and Ireland, The Intensive Care Foundation, National Institute of Health Research, National Institute of Academic Anaesthesia, The Sir Halley-Stuart Trust, The Frances and Augustus Newman Trust and The Down Syndrome Trust. Mike Grocott has also received fees for lecturing, unrelated to this review, from Fresenius Kabi, Edwards Lifesciences and Cortex GmbH.

Professor Mythen has received honoraria for speaking/consultation and/or travel expenses from Baxter, BBraun, Covidien, Fresenius-Kabi, Hospira, LidCo and as a consultant to AQIX (start up company with a novel crystalloid solution - pre-clinical). He is a director of Medical Defence Technologies LLC ("Gastrostim" patented) and a co-Inventor of "QUENCH" IP being exploited by UCL Business. Professor Mythen's chair at UCL is endowed by Smiths Medical Ltd and this company provide charitable donations to the department on an annual basis. Deltex Medical also provide unrestricted grant funds to Prof Mythen's Department. Professor Mythen is a member of the IV Fluids Guideline Development Group for the National Institute of Clinical Excellence (NICE) and he is a co-Author of the GIFTASUP fluid guidelines.

Mark Hamilton has received lecture fees from Edwards Life Sciences for two lectures related to optimization.

Professor Kathy Rowan is ICNARC CTU Director and the ICNARC CTU is managing a trial (evaluating the effectiveness of perioperative goal directed

haemodynamic therapy in high-risk patients undergoing major abdominal surgery involving the gastrointestinal tract) part-funded via a National Institute for Health Research (NIHR) Clinician Scientist Award given to Dr Rupert Pearse.

All other authors': none known.

SOURCES OF SUPPORT

Internal sources

- Special trustees of the Middlesex Hospital, UK.

External sources

- No sources of support supplied

DIFFERENCES BETWEEN PROTOCOL AND REVIEW

The search strategy was changed after December 2000, after consultation with Cochrane search editors.

NOTES

October 2016

This review includes the following study [Mayer 2010](#).

Joachim Boldt is a co-author of this study. Many publications by Joachim Boldt have been retracted or are being investigated; [Mayer 2010](#) has not been retracted and future publication status is unknown. If the Mayer study is retracted in the future, then the study will be excluded in the updated version of this review.

INDEX TERMS

Medical Subject Headings (MeSH)

Arrhythmias, Cardiac [etiology]; Blood Circulation [*physiology]; Cardiovascular Agents [administration & dosage]; Length of Stay; Oxygen Consumption [*physiology]; Plasma Substitutes [administration & dosage]; Postoperative Complications [*mortality; prevention & control]; Randomized Controlled Trials as Topic; Renal Insufficiency [prevention & control]; Respiratory Insufficiency [prevention & control]; Surgical Procedures, Operative [adverse effects; *mortality]; Surgical Wound Infection [prevention & control]

MeSH check words

Adult; Humans