Cochrane
Library

Cochrane Database of Systematic Reviews

Simulated presence therapy for dementia (Review)

Abraha I, Rimland JM, Lozano-Montoya I, Dell'Aquila G, Vélez-Díaz-Pallarés M, Trotta FM, Cruz-Jentoft AJ, Cherubini A

Abraha I, Rimland JM, Lozano-Montoya I, Dell'Aquila G, Vélez-Díaz-Pallarés M, Trotta FM, Cruz-Jentoft AJ, Cherubini A.

Simulated presence therapy for dementia.

Cochrane Database of Systematic Reviews 2017, Issue 4. Art. No.: CD011882.

DOI: 10.1002/14651858.CD011882.pub2.

www.cochranelibrary.com

TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
SUMMARY OF FINDINGS FOR THE MAIN COMPARISON	4
BACKGROUND	5
OBJECTIVES	7
METHODS	7
RESULTS	9
Figure 1.	10
Figure 2.	14
Figure 3.	15
DISCUSSION	17
AUTHORS' CONCLUSIONS	19
ACKNOWLEDGEMENTS	19
REFERENCES	20
CHARACTERISTICS OF STUDIES	23
DATA AND ANALYSES	35
APPENDICES	35
CONTRIBUTIONS OF AUTHORS	38
DECLARATIONS OF INTEREST	38
SOURCES OF SUPPORT	38
DIFFERENCES BETWEEN PROTOCOL AND REVIEW	39
INDEX TERMS	39

Simulated presence therapy for dementia

Iosief Abraha¹, Joseph M Rimland¹, Isabel Lozano-Montoya², Giuseppina Dell'Aquila¹, Manuel Vélez-Díaz-Pallarés², Fabiana M Trotta¹, Alfonso J Cruz-Jentoft², Antonio Cherubini¹

¹Geriatrics, National Institute of Health and Science on Aging (INRCA), Ancona, Italy. ²Geriatrics, Hospital Universitario Ramón y Cajal, Madrid, Spain

Contact address: Joseph M Rimland, Geriatrics, National Institute of Health and Science on Aging (INRCA), Via Santa Margherita, 5, Ancona, 60124, Italy. joe.rimland@live.com, joe.rimland@live.com.

Editorial group: Cochrane Dementia and Cognitive Improvement Group.

Publication status and date: New, published in Issue 4, 2017.

Citation: Abraha I, Rimland JM, Lozano-Montoya I, Dell'Aquila G, Vélez-Díaz-Pallarés M, Trotta FM, Cruz-Jentoft AJ, Cherubini A. Simulated presence therapy for dementia. *Cochrane Database of Systematic Reviews* 2017, Issue 4. Art. No.: CD011882. DOI: 10.1002/14651858.CD011882.pub2.

Copyright © 2017 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

Dementia is a common and serious neuropsychiatric syndrome, characterised by progressive cognitive and functional decline. The majority of people with dementia develop behavioural disturbances, also known as behavioural and psychological symptoms of dementia (BPSD). Several non-pharmacological interventions have been evaluated to treat BPSD in people with dementia. Simulated presence therapy (SPT), an intervention that uses video or audiotape recordings of family members played to the person with dementia, is a possible approach to treat BPSD.

Objectives

To assess the effects of SPT on behavioural and psychological symptoms and quality of life in people with dementia.

Search methods

We searched ALOIS (the Specialised Register of the Cochrane Dementia and Cognitive Improvement Group), CENTRAL (*The Cochrane Library*) (9 February 2016), MEDLINE Ovid SP (1946 to 6 January 2017), Embase Ovid SP (1972 to 6 January 2017), PsycINFO Ovid SP (1806 to 6 January 2017), CINAHL via EBSCOhost (1980 to 6 January 2017), LILACS via BIREME (all dates to 6 January 2017), ClinicalTrials.gov (ClinicalTrials.gov) (all dates to 6 January 2017), and the World Health Organization (WHO) Portal (apps.who.int/trialsearch) (all dates to 6 January 2017). We also checked the reference lists of relevant articles to identify any additional studies.

Selection criteria

Randomised and quasi-randomised controlled trials, including cross-over studies, that evaluated the efficacy of SPT, consisting of personalised audio or videotape recordings of family members, in people with any form of dementia.

Data collection and analysis

Two authors independently selected studies, assessed risk of bias and extracted data. No meta-analyses were conducted because of substantial heterogeneity among the included studies.

Main results

Three trials with 144 participants met the inclusion criteria. Two of the trials had a randomised cross-over design, one was a cross-over trial which we classified as quasi-randomised.

Participants in the included studies were people with dementia living in nursing homes. They were predominantly women and had a mean age of over 80 years. SPT was performed using an audio or video recording prepared by family members or surrogates. It varied in its content, frequency of administration and duration. All the studies compared multiple treatments. In one study, SPT was compared with two other interventions; in the other two studies, it was compared with three other interventions. Specifically, SPT was compared to usual care, personalised music (two studies), a 'placebo' audiotape containing the voice of a person (two studies), and one-to-one social interaction performed by trained research assistants (one study). In terms of outcomes evaluated, one study considered agitation and withdrawn behaviour (both assessed with three methods); the second study evaluated verbal disruptive behaviour (assessed with three methods); and the third study evaluated physically agitated behaviour and verbally agitated behaviour (the method used was not clearly described).

According to the GRADE criteria, the overall quality of the evidence was very low due to very small numbers of participants and risk of bias in the included studies; (none of the trials was at low risk of selection bias; all the trials were at high risk of performance bias; one trial was at high risk of attrition bias; and all had unclear selective reporting).

Because of variation in the participants, the format of SPT, the comparison interventions, and the measures used to assess outcomes, we judged the results unsuitable for a meta-analysis.

Within each trial, the effect of SPT on behaviour, compared to usual care, was mixed and depended on the measure used. Two trials which included a personalised music intervention reported no significant differences between simulated presence and music on behavioural outcomes. Because the overall quality of the evidence was very low, we were very uncertain regarding all the results.

None of the studies evaluated quality of life or any of our secondary outcome measures (performance of activities of daily living, dropout and carer burden).

Authors' conclusions

We were unable to draw any conclusions about the efficacy of SPT for treating behavioural and psychological symptoms and improving quality of life of people with dementia. New high-quality studies are needed to investigate the effect of SPT.

PLAIN LANGUAGE SUMMARY

Simulated presence therapy for dementia

Review question

Can simulated presence therapy (SPT) treat problem behaviours, and improve quality of life for people with dementia?

Background

Dementia is an illness, most common among older people, in which memory and other brain functions deteriorate and people gradually become dependent on others for care. Many people with dementia, particularly in its later stages, show signs of distress at times, or behave in ways which are difficult for their caregivers to manage. There is a lot of interest in finding ways to treat this without using drugs. Simulated presence therapy is a type of treatment which has been used mainly in nursing homes. It involves playing a personalised video or audiotape recording of family members to the person with dementia when he or she is distressed or agitated.

Study characteristics

We looked for trials which compared SPT to usual care or to another treatment. Ideally, people with dementia should have been randomly allocated to one or other treatment, but we also included trials even if treatment allocation was not strictly random.

We found three trials which met our inclusion criteria. The 144 participants were all living in nursing homes. The majority were women with an average age of over 80 years and severe dementia. The way SPT was administered was different in each trial. All the trials used more than one comparison treatment, which differed between trials. The trials all attempted to measure an effect on agitated behaviours, but used different approaches.

Key findings

Because the trials were so different from each other, we were not able to pool the results. Individually, each trial reported different methods to assess the effect of SPT on behavioural problems and the results varied depending on the method used to measure the outcome.

None of the studies assessed quality of life, effect on daily activities, effects on caregivers, or how likely participants were to drop out of the study.

Quality of the evidence

The studies were small and all had problems with their methods which could have biased their results. Hence, we thought the overall quality of the evidence was very low, meaning we cannot be at all confident in the results.

Conclusion

Not enough high-quality research has been done to allow us to judge whether SPT can help people with dementia who are distressed or agitated.

SUMMARY OF FINDINGS FOR THE MAIN COMPARISON [\[Explanation\]](#)

Simulated presence therapy compared with usual care for people with dementia			
Patient or population: dementia Setting: nursing home Intervention: Simulated presence therapy Comparison: usual care			
Outcomes	Impact	No. of participants (studies)	Quality of the evidence (GRADE)
Overall BPSD (multidomain) - not assessed	-	-	-
Agitation	One study reported that agitation assessed by two methods (direct observation and weekly staff survey) was not reduced by SPT, while a third method (staff observation logs) showed a decline in agitation. In a second study, two methods (VDB score and standardised observational data) showed that SPT reduced verbally disruptive behaviour, while there was no effect on the same outcome with a third method (nurses' assessments). A third study with an unclear assessment method reported that SPT was effective in reducing physical agitation and verbal agitation compared to usual care	144 (3 trials)	⊕○○○ VERY LOW ^{1,2}
Depression - assessed in one study as "withdrawn behaviours" which included a lack of interest and sad mood	SPT did not improve withdrawn behaviours, interest or mood based on direct observations. Weekly staff surveys did not show any effect of SPT on mood, although the intervention was associated with increased interest. The staff, through observation logs, reported that SPT compared to usual care statistically significantly improved withdrawn behaviour	54 (1 trial)	⊕○○○ VERY LOW ^{3,4}

Anxiety - not measured	-	-	-
Quality of life - not measured	-	-	-
Dropouts - not measured	-	-	-

* **The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: Confidence interval; **RR:** Risk ratio; **OR:** Odds ratio;

GRADE Working Group grades of evidence

High quality: We are very confident that the true effect lies close to that of the estimate of the effect

Moderate quality: We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different

Low quality: Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect

Very low quality: We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect

¹ Downgraded by two levels because of very serious concern about the risk of bias: none of the trials was at low risk of selection bias (one study was at high risk of bias); all the trials were at high risk of performance bias; one trial was at high risk of attrition bias; all had unclear selective reporting; and two were at high risk of other biases.

² Downgraded by two levels because of serious concern about imprecision: none of the studies reached the optimal information size.

³ Downgraded by two levels because of serious concern about the risk of bias; the trial was at unclear risk of selection bias; high risk of performance bias; unclear risk of selective reporting bias; and high risk of other bias.

⁴ Downgraded by two levels because of very serious concern about imprecision: very small sample size.

BACKGROUND

Description of the condition

Dementia is a common and serious neuropsychiatric syndrome, characterised by progressive cognitive and functional decline. The prevalence of dementia is estimated to be 6% in populations aged 60 years or older (Prince 2013). It increases dramatically with age, reaching 43% in people 85 years or older (Thies 2012).

Dementia imposes a considerable burden on families and is becoming a major socioeconomic challenge for the healthcare system (Wimo 2013). It is one of the strongest predictors of nursing home admission and is a major clinical issue for nursing home residents. Its prevalence varies from 26% to 48% in US nursing homes to over 60% in European nursing homes (Cherubini 2012).

The majority of people with dementia develop behavioural disturbances, also known as behavioural and psychological symptoms of

dementia (BPSD), over the course of the condition. BPSD comprise a heterogeneous group of non-cognitive symptoms, such as agitation, disruptive behaviour, aberrant motor behaviour including wandering, irritability, hallucinations, anxiety, depression, apathy, delusions, disinhibition, and appetite and sleep alterations (Cerejeira 2012). BPSD are a major component of the dementia syndrome as they are strongly correlated with the severity of functional impairment, occurring most frequently in people with dementia in hospitals and nursing homes (Cerejeira 2012). These symptoms are independently associated with poor outcomes such as distress among those with dementia and their caregivers, inappropriate prescribing of antipsychotic medication, long-term hospitalisation, mortality, excess morbidity, institutionalisation, early placement in a nursing home, and increased healthcare costs (Borson 1997; Clyburn 2000; Kales 2015; Van Den Wijngaert 2007).

Several studies have documented that almost all people with de-

mentia experience at least one episode of behavioural disturbance at some point during their illness (Kunik 2010; Lyketsos 2007; Spalletta 2010; Steinberg 2008). In a cohort of 408 dementia participants, for example, the Cache County Study estimated the five-year period prevalence of neuropsychiatric symptoms and found that 97% of the participants experienced at least one symptom. Five-year period prevalence was highest for depression (77%), apathy (71%), and anxiety (62%). Rates for agitation/aggression vary between 13% and 24% (Steinberg 2008). Behavioural symptoms frequently occur together (for example wandering with sleep problems or irritability with persecutory delusions) (Savva 2009).

Description of the intervention

Behavioural problems in people with dementia are often treated with drugs, which may lead to several adverse outcomes including sedation, falls, extrapyramidal disturbances, stroke, and increased mortality (Lopez 2013; Sink 2005; Stern 1994). For this reason, non-pharmacological interventions are an appealing alternative (Abraha 2015b); and have been recommended as a first choice of treatment for BPSD (Salzman 2008). Several non-pharmacological interventions have been evaluated to treat BPSD in people with dementia (Abraha 2017). These include emotion-oriented therapies (such as reminiscence therapy (Woods 2009), simulated presence therapy (Zetteler 2008), validation therapy (Neal 2003)), sensory stimulation interventions (such as acupuncture (Peng 2007), aromatherapy (Forrester 2014), light therapy (Forbes 2014), massage/touch therapy (Viggo Hansen 2006), music therapy (Vink 2004), Snoezelen multisensory stimulation (Chung 2002), and transcutaneous electrical nerve stimulation (Cameron 2003)), and other interventions (such as functional analysis-based interventions (Moniz-Cook 2012), and exercise therapy (Forbes 2013)). It has been reported that psychosocial interventions work best to reduce agitation and improve other behavioural disturbances in people with dementia when they are personalised to reflect the individual's background and environmental circumstances (O'Connor 2009).

The focus of this review is simulated presence therapy (SPT), in which video or audiotape recordings of family members are played to the person with dementia (Woods 1995). The content of the recordings may vary depending upon the interests of the individual and may include conversations, stories, or shared memories. The intention is that the recorded voice of a family member is reassuring and that anxiety and distress are reduced by making the environment of the person with dementia as familiar as possible (Woods 1995).

How the intervention might work

SPT is sometimes referred to as simulated family presence therapy (Garland 2007), because it originated from the observation

that nursing home residents who received more visits from family members were less agitated and had greater life satisfaction (Woods 1995). Since family members may not be able to visit frequently, and family visits may be forgotten rapidly by residents with dementia, SPT aims to reduce the separation anxiety experienced by the resident (Miesen 1993; Peak 2002).

The technique was first described in a 1995 study by Woods and Ashley, in which 27 nursing home residents with dementia listened to a tape prepared by their caregivers. The authors reported a substantial reduction of behavioural problems such as verbal aggression and social isolation, and an increase in positive behaviours such as proper verbalisation, smiling, and singing (Woods 1995). In the following years, a limited number of studies have been performed to evaluate the efficacy of SPT. These studies are characterised by considerable methodological heterogeneity, including different outcomes, as some studies examined the effect of SPT on quality of life, while other investigations attempted to reduce challenging behaviours.

Although SPT has been investigated for more than 20 years, the mechanisms by which SPT might work are still unknown. Several explanatory models have been proposed for behavioural disturbances. According to the need-driven, dementia-compromised behaviour model, behavioural disturbances of the cognitively impaired person are an expression of an unmet need or goal (Algase 1996). These disturbances might therefore be seen as an expression of the person seeking help or contact from family members. In this respect SPT might fulfil this unmet or personal need.

Another conceptual framework, the progressively lowered stress threshold model, proposes that people with dementia become progressively unable to understand their environment (for example recognize objects and their purposes and use). As a consequence, people with dementia experience stress that causes a variety of behavioural disturbances (Smith 2004). In this context, voices and images of family members may represent well-known references that help to lower stress levels.

Generally, the literature is scarce in terms of the epidemiology of adverse events associated with non-pharmacological interventions. In the case of SPT, adverse events cannot be excluded, including the possibility that the intervention might worsen agitation or mood. Woods and Ashley reported that in 7% of the observation during SPT treatment, behaviours either remained unchanged or worsened with the audiotape (Woods 1995). In some cases, for example, material contained in the tapes could be distressing for the person with dementia (Peak 2002; Garland 2007).

Why it is important to do this review

The majority of BPSD, in particular agitation, disruptive behaviour, and aberrant motor behaviours, are currently treated with psychotropic drugs that are not very effective (Howard 2012; Katz 2007; Olin 2002), have several important adverse effects including increased mortality (Budnitz 2011; Opie 1999), and result in

excessive drug costs and healthcare utilisation (Leendertse 2011; Rottenkolber 2011). Non-pharmacological interventions should therefore be the first choice to treat BPSD.

SPT is a simple intervention that does not require expensive equipment, nor does it require extensive training. In addition, SPT reproduces an encounter of the patient with a family member, which is usually a positive experience for the former. The potentially relevant adverse event is an increase in agitation, which should not cause a serious risk for the patient's health. As such, we believe that it is important to verify whether SPT is effective in treating BPSD in people with dementia.

OBJECTIVES

To assess the effects of SPT on behavioural and psychological symptoms and quality of life in people with dementia.

METHODS

Criteria for considering studies for this review

Types of studies

We included randomised and quasi-randomised controlled trials, including cross-over studies, of SPT in people with dementia.

Types of participants

People with any form and severity of dementia diagnosed according to the International Classification of Diseases, Revision 10 (ICD-10), Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV), DSM-5, or comparable criteria. No restriction in terms of setting was applied.

Types of interventions

Active treatment

SPT consisting of audio or videotape recording that family members or caregivers have personalised. The content of the recording should include positive experience from the participant's past life and shared memories involving family or close friends.

Comparison

- Usual care with no additional activity.
- Any activity that differs in content and approach from SPT, but is additional to usual care (for example, music therapy, reminiscence groups, reality orientation groups, or social contact groups that do not use the techniques identified as SPT).

Types of outcome measures

Primary outcomes

1. Any behavioural and psychological symptoms as measured using scales such as the Neuropsychiatric Inventory (NPI) or Brief Psychiatric Rating Scale (BPRS) (multidomain scales), Cohen-Mansfield Agitation Inventory (CMAI) (scale specific to agitation), or Cornell Scale for Depression in Dementia (CSDD) (scale specific to depression).
2. Quality of life.

Secondary outcomes

1. Activities of daily living.
2. Caregiver burden.
3. Dropouts (as a measure of acceptability).

Search methods for identification of studies

Electronic searches

We searched ALOIS (www.medicine.ox.ac.uk/alois), the Cochrane Dementia and Cognitive Improvement Group's Specialised Register (up to 6 January 2017).

ALOIS is maintained by the Information Specialists for the Cochrane Dementia and Cognitive Improvement Group and contains studies that fall within the areas of dementia prevention, dementia treatment and management, and cognitive enhancement in healthy elderly populations. The studies are identified through:

1. monthly searches of a number of major healthcare databases: MEDLINE, Embase, CINAHL, PsycINFO, and LILACS;
2. monthly searches of a number of trial registers: ISRCTN, UMIN (Japan's trial register), the World Health Organization International Clinical Trials Registry Platform (which covers ClinicalTrials.gov, ISRCTN, the Chinese Clinical Trials Register, the German Clinical Trials Register, the Iranian Registry of Clinical Trials, and the Netherlands National Trials Register, plus others);
3. quarterly searches of the Cochrane Library's Central Register of Controlled Trials (CENTRAL);

4. six-monthly searches of a number of grey literature sources: ISI Web of Knowledge Conference Proceedings, Index to Theses, Australasian Digital Theses.

To view a list of all sources searched for ALOIS see [About ALOIS](#) on the ALOIS website (www.medicine.ox.ac.uk/alois).

Details of the search strategies run in healthcare bibliographic databases, used for the retrieval of reports of dementia, cognitive improvement, and cognitive enhancement trials, can be viewed in the 'Methods used in reviews' section within the editorial information about the [Cochrane Dementia and Cognitive Improvement Group](#).

We ran additional searches in MEDLINE (Ovid SP), Embase (Ovid SP), PsycINFO (Ovid SP), CINAHL (EBSCOhost), ClinicalTrials.gov, and the World Health Organization International Clinical Trials Registry Platform to ensure that the searches for the review were as comprehensive and up to date as possible. We have presented the search strategies used in [Appendix 1](#).

We carried out translations for non-English papers, where possible.

Searching other resources

We searched the grey literature, such as conference proceedings, through Embase (Ovid SP) and through Web of Science (ISI Web of Science), and the reference lists of all the potentially relevant trials or reviews identified through the above searches. We applied no language restriction.

Data collection and analysis

Selection of studies

In order to facilitate the retrieval of articles of interest, we downloaded all references into a single database using the EndNote reference management software ([Endnote 2013](#)).

We obtained full-text copies that two review authors independently assessed for eligibility (IA, JMR). The review authors resolved any disagreements by discussion.

Data extraction and management

For eligible studies, pairs of review authors (JMR, IA and IL, MV) independently extracted the data using the Cochrane form available in Review Manager 5 (RevMan 5) software ([Review Manager 2014](#)). They resolved disagreements by discussion. We extracted the following information from each included study.

- General information about the study (year of publication, setting).
- Study design, methods of recruitment, inclusion and exclusion criteria, details of the control and comparison groups, and incentives for participation.
- Characteristics and number of participants.

- Characteristics of the intervention.
- Results.

One review author (IA) entered data into the Review Manager 5 software ([Review Manager 2014](#)), and a second review author checked the data for accuracy.

Assessment of risk of bias in included studies

We used Cochrane's recommended tool for assessing risk of bias ([Higgins 2011](#)).

We assessed the following domains of each included study: sequence generation and allocation concealment for selection bias ([Savovic 2012](#); [Wood 2008](#)); blinding of participants or personnel for performance bias ([Savovic 2012](#); [Wood 2008](#)); blinding of outcome assessors for detection bias ([Savovic 2012](#); [Wood 2008](#)); incomplete outcome data for attrition bias ([Abraham 2015a](#)); selective reporting bias ([Chan 2004](#); [Macura 2010](#)); and other potential sources of bias ([Higgins 2011](#)). Other sources of bias included the comparability of intervention and control group characteristics at baseline, validation of outcome assessment tools, and reliability of outcome measures.

Review authors (IA, JMR, ILM, MV) independently evaluated the risk of bias of the included studies, using the criteria outlined in the *Cochrane Handbook for Systematic Reviews of Interventions* ([Higgins 2011](#)). They resolved any disagreement by discussion; or in the event of deadlock, by consulting with a third review author. We described the quality assessment in a 'Risk of bias' table.

Measures of treatment effect

Where a sufficient number of trials with homogeneous populations were identified, we planned to carry out meta-analyses of primary and secondary outcomes. For the analysis of dichotomous outcomes, we planned to use risk ratios with 95% confidence intervals (CI); for the analysis of continuous outcomes, we planned to use the mean difference or standardised mean difference and 95% CI.

Unit of analysis issues

The unit of analysis was the individual participant with dementia. In case of studies that contributed with multiple, correlated comparisons, we planned to combine study groups into a single pairwise comparison.

In the event of cluster-randomised trials, if the reported analysis did not correctly account for the cluster design we planned to consider whether it was possible to estimate the effective sample size using the intracluster correlation coefficient. If this was not possible, then we planned to conduct sensitivity analyses excluding the trials that did not originally adjust for clustering.

Dealing with missing data

In the event of missing data, we contacted the corresponding authors to try to obtain the relevant information.

Where 'intention to treat' was not available, we planned to conduct analyses using 'available cases' data (i.e. only those whose results are known).

Assessment of heterogeneity

We planned to assess heterogeneity of study characteristics and statistical heterogeneity. We evaluated heterogeneity by examining the relative table of characteristics of the included population, the type of interventions, and the type of outcome measures.

Where meta-analysis could be performed, we planned to appraise statistical heterogeneity through a visual assessment of the forest plot, in addition to evaluating the χ^2 test and I^2 statistic. We planned to consider a χ^2 test with a P value of 0.10 to be significant, and we planned to interpret the I^2 statistic as: 0% to 40% indicates unimportant heterogeneity; 30% to 60% indicates moderate heterogeneity; 50% to 90% indicates substantial heterogeneity; and 75% to 100% indicates considerable heterogeneity (Deeks 2011).

Assessment of reporting biases

We planned to use a funnel plot to examine for small-study effects, which may indicate publication bias, had there been more than 10 studies that could be combined into a meta-analysis,

Data synthesis

We planned to pool data for analysis if the interventions, outcomes, and participant populations in the individual studies had been sufficiently similar (to be determined by consensus).

We planned to use Review Manager 5 for all analysis (Review Manager 2014).

Subgroup analysis and investigation of heterogeneity

We planned subgroup analyses based on several participant characteristics such as sex, severity of dementia, severity of behavioural problems at baseline, the nature of the control intervention (treatment 'as usual' or another active treatment), stage of dementia, care setting (community versus long-term care facility), and intervention characteristics such as length of treatment, including number of sessions, and mode of delivery (audio versus video recording).

Sensitivity analysis

We planned to perform sensitivity analyses to assess the consistency and robustness of the results of the meta-analysis based on:

- including only studies with low risk of selection bias (sequence generation and allocation concealment);
- including only trials with low risk of detection bias (blinding of outcome assessor);
- including only trials with low risk of attrition of bias;
- including only trials that used an intention-to-treat analysis;
- repeating analyses using a fixed-effect model for data synthesis.

Summary of findings table

We used the GRADE approach to assess the quality of the supporting evidence behind each estimate of treatment effect for the comparison of SPT with usual care. We used risk of bias, imprecision, inconsistency, indirectness, and publication bias to rate the overall evidence (Balslem 2011).

We present key findings of the review in 'Summary of findings for the main comparison'. This includes a summary of the amount of data and the overall quality of the evidence for overall behavioural and psychological symptoms, agitation, depression, anxiety, quality of life and dropouts (Schünemann 2011a; Schünemann 2011b).

RESULTS

Description of studies

Results of the search

A systematic search of electronic databases combined with a hand-search of references identified 248 publications (Figure 1). After removing duplicate records, 185 records were screened of which 172 were removed and 13 were eligible for full-text evaluation. Nine references, which were either not trials or were clearly outside the scope of this review, were excluded. Finally, three trials remained for inclusion (Camberg 1999; Cohen-Mansfield 1997; Garland 2007).

Figure 1. Study flow diagram.

Included studies

The three included trials initially enrolled 144 participants of whom 116 were evaluated by the authors in the final analysis (Cohen-Mansfield 1997 initiated with 60 participants and evaluated only 32). For full details, see the [Characteristics of included studies](#) table.

Setting

Two trials were conducted in the USA (Camberg 1999; Cohen-Mansfield 1997); and one in Australia (Garland 2007). All of the studies were conducted in a nursing home setting.

Study design

Two trials were randomised cross-over trials (Camberg 1999; Garland 2007); and one was a cross-over trial which we classified as quasi-randomised (Cohen-Mansfield 1997). In this trial, participants were assigned to four groups, each of which rotated through the active treatments ($n = 3$) and control treatment ($n = 1$) in a different order so that by the end of the evaluation period all groups were exposed to all the treatments and to the no-treatment period (Cohen-Mansfield 1997).

Participants

Camberg 1999 included 54 participants of whom 48 (89%) were women. The average age was 82.7 years (standard deviation (SD) 7.5 years). The participants had severe cognitive impairment (Mini Mental State Examination (MMSE): 5.1, SD $\pm$ 4.4; Texas Success Initiative (TSI) Assessment: 13.1 SD $\pm$ 6.9) and low functional ability (Bedford Alzheimer's Nursing Scale (BANS): 13.2, SD $\pm$ 4.0; Activity of daily living (ADL) Self-Performance Scale: 2.7, SD $\pm$ 1.2). No other baseline information was provided. Participants were selected based on the presence of at least one agitated behaviour per day of those listed on the short form of the Cohen-Mansfield Agitation Inventory Scale or one indication of withdrawn behaviour "defined either by "sounding sad" or "crying", that occurred at least "often"; or "seldom" interested in activities, social interaction, or in their immediate environment, from the Multidimensional Observation Scale for Elderly Subjects (MOSES)"

At baseline, the authors determined the prevalence of agitated behaviours from 6903 direct observations and 222 weekly staff surveys. The most frequent kinds of physical agitation were "repetitive motions in place, outward motions with the extremities, and pacing/aimless wandering", while for verbal agitation the most recurrent types were "repetitive vocalization, cursing/verbal aggression, and negative words or attitude". Most agitated behaviours were

quite mild and only occurred for a few minutes; and in 25% of direct observations, agitation was not observed at all. The authors also measured the prevalence of withdrawn behaviours as interest and mood. Subjects showed no interest in almost half (47%) of the direct observations and in more than half (55% to 72% based on the category of interest) of the weekly staff surveys. Regarding mood, subjects displayed no pleasure in 75% of direct observations and in nearly 40% of weekly staff surveys. During most (82.5%) of the direct observations, subjects were reported to have "no expression" on their faces.

In Cohen-Mansfield 1997, the inclusion criterion was the presence of verbally disruptive behaviours (VDBs) manifested at least several times a day. In this study, 60 participants in a nursing home were identified and provided consent. Of these, five died before any data were collected, 11 became quiet, three refused to continue, and two were physically restrained and thus excluded. Hence, 32 participants remained with complete data. These participants had an average age of 86.8 years (SD $\pm$ 1.2 years) and 26 were female. At baseline, they had severe functional and cognitive impairment, as shown by the Lawton and Brody Physical Self Maintenance Scale (Activities of Daily Living (ADL)) score (4.0, SD $\pm$ 0.2), and the Brief Cognitive Rating Scale score (5.5, SD $\pm$ 0.2).

Baseline evaluation in the 32 subjects who completed the study revealed that 11 participants exhibited VDB several times each hour, 18 subjects several times per day and three participants once or twice per day. An analysis of 11,520 3-minute observations, using the Screaming Behavioural Mapping Instrument (SBMI), before, during and after interventions, revealed that the most frequent VDBs, in order of prevalence, were complaining, shouting, nonsense talk, repeating words and other VDB.

Before initiation of the interventions, participants underwent a medical examination to determine if pain might account for VDB: no physiological signs of pain were found for 23 subjects. Although nine participants received pain medication, physicians stated that the VDB was not caused by pain for any of the subjects.

In Garland 2007, the participants were 30 nursing home residents with a diagnosis of dementia, who had been resident for at least three months and had had at least one disruptive behaviour several times a day in the preceding two weeks (assessed with the CMAI). Mean age was 79 years (range 66 to 93 years) and 63% were female. All participants were prescribed some type of psychotropic medication and had a very low baseline mean MMSE of 2.5 (range 0 to 12). The authors measured pre-intervention agitated behaviours (2880 observed target behaviours), which consisted of physically aggressive behaviours (3.8%), physically non-aggressive behaviours (64.8%) and verbally aggressive and non-aggressive behaviours (31.4%).

Intervention

[Camberg 1999](#) was a three-intervention study. The authors reported “A Latin Squares crossover design was adopted for this study”.

Each treatment continued for 17 days over 4 weeks, with a 10-day washout period following each treatment.

The interventions were as follows.

(a) Simulated presence therapy based on a personalised audio tape (SimPres) consisting of a one-sided telephone conversation with a family member or surrogate, containing many special memories and positive emotions. Family members received a memory inventory form, conversational guidelines, and an instructional audio-tape as training material. The study staff coached family members either by telephone or in person, immediately before the recording session. A headset and an auto-reverse tape recorder were used to play the audio tape;

(b) A placebo, which consisted of a tape made to resemble the SimPres treatment by containing the voice of a person who read emotionally neutral newspaper articles;

(c) Usual care, which was the regular approach used by nursing staff to manage behaviour (e.g. staff interactions, redirection, or physical restraints).

The nursing staff in the study units applied the intervention following a 30-minute training session during which they were instructed to use the SimPres or placebo audio tapes at least twice a day, Monday to Friday, as an alternative to their usual care, whenever a study subject displayed an agitated or withdrawn behaviour. The nurses were requested not to listen to the audiotapes, but were informed that both tapes contained verbal messages. Procedures and demonstrations of tape use and case examples of the intervention were provided. Adherence to the protocol was ensured, and feedback was given to the study staff by a study monitor, who spent 20 hours per week at each facility.

In [Cohen-Mansfield 1997](#), participants were assigned to four groups, each of which was exposed to all three interventions and to a control no-intervention phase. In each group, the order of interventions was different. By the end of the study, all participants were exposed to all types of intervention.

Each group was exposed to each intervention for two consecutive weeks followed by one week of washout.

The interventions were as follows.

(a) A SPT intervention consisting of a videotape, which the relatives of all participants were asked to prepare, was provided to the resident. A research assistant, a nursing assistant, and a volunteer prepared the videotapes for four participants whose relatives either refused or were unable to create a videotape. Family members received general guidelines to prepare the videotapes, but they chose the specific content and the style of communication.

(b) A 30-minute audiotape recording of music, based on the participant’s musical preferences (provided by relatives).

(c) One-to-one social interaction performed by trained research assistants. A manual was prepared to guide the social interaction,

which included information on how to interact with a nursing home resident, how to start an activity and a broad list of possible activities ordered by the degree of involvement, from high to low, required of the resident. A “box of activities”, comprising games, pictures, balls, books, etc. was assembled. Alternatively, the following activities were provided: (i) conversation with a research assistant; (ii) motion exercise (e.g. tossing a foam ball, hand and arm movement); (iii) sensory stimulation (a sensory kit including different fabrics, school supplies, health aids, make-up, spices, soaps, etc. was available to be presented to the residents to stimulate touch, sight, and smell); (iv) manual activities (such as making a collage, completing a simple puzzle, clipping coupons).

[Garland 2007](#) was a four intervention, cross-over study. Participants were randomly assigned to three groups and sequentially exposed to all four interventions, with a two-day washout between each one. Interventions were administered once a day, for three days each, during weeks two, three and four.

The interventions were as follows.

(a) “Simulated family presence”, consisting of a professionally edited tape recording of a semi-structured interview of a family member, conducted by a trained psychologist, regarding participants’ earlier lives. The script was designed to resemble a telephone conversation regarding special memories, beloved family members, and family episodes.

(b) A music intervention, which consisted of listening to music based on participants’ selections of popular, big band, and Greek and Dutch music.

(c) A placebo intervention, which was a neutral audiotape, based on non-personal, non-emotive material (a psychologist read from a gardening book in neutral tones) to control for the effects of receiving attention from researchers and listening to a tape recording.

(d) Usual care, which was not defined, but the authors stated that “usual care observations were recorded for 45 minutes on each of three days in week 1 of the study at the times of peak agitation nominated by nursing staff”.

Outcome

[Camberg 1999](#) evaluated agitated and withdrawn behaviours. Agitation was defined as an experience of an unpleasant state of excitement, observable without subjective interpretation, not strictly triggered by caregiver activities, not related to known physical needs of the patient that can be alleviated, and without motivational intent. Withdrawn behaviours were defined as a “lack of interest in people, activities, or things in the subject’s environment, combined with sad mood”. Withdrawn behaviour was assessed considering the combination of “interest”, and manifestations of mood. Observers used two items from the Philadelphia Geriatric Centre Affect Rating Scale (PARS), “interest” and “pleasure” to evaluate withdrawn behaviour.

Outcomes were measured by (1) direct observation, (2) staff observation logs, and (3) weekly nursing staff surveys.

Direct observations were made for 3 hours and 20 minutes each week, between 9 a.m. and 7 p.m. by trained, blinded, non-participant observers. Each observation block was 20 minutes long, divided into four 5-minute observation intervals. Behaviours were assessed according to the schedule, whether or not a specific intervention was implemented. The following scales were used: (1) a seven-item Observed Agitation Scale (OAS) developed by the researchers; (2) an agitation visual analogue scale (AVAS); (3) two Positive Affect items, “interest” and “pleasure,” from the PARS; (4) a withdrawal visual analogue scale (WVAS); and (5) facial diagrams of mood (FACE).

Staff observation logs were based on the nursing staff’s impression of a participant’s response to each intervention. (The target behaviour, the intervention and its duration, and the subject’s response (improved, became worse, or stayed the same) were all documented daily, from Monday to Friday).

Weekly staff surveys (administered to a nursing staff member who had cared for the subject for at least three days during the preceding week) used two behavioural rating scales: (1) the short form of the Cohen-Mansfield Agitation Inventory (SCMAI) to record staff recall of the frequency of agitated behaviours; and (2) selected, modified items from the Multidimensional Observation Scale for Elderly Subjects (MOSES) to record the subject’s mood and interest.

In [Cohen-Mansfield 1997](#), VDB was the only outcome assessed. It was defined as “verbal or vocal behaviours that were either repetitive, disruptive or inappropriate to the circumstances in which they were manifested”. It was assessed using three different methods: 1) tape recordings; 2) standardised observations; and 3) informant ratings (nurses’ assessments).

1) VDB duration was registered with a tape recorder located near a subject. The tape was coded by a trained research assistant using a computer program specifically developed for coding video and audiotapes. An episode of VDB was considered to be finished if the resident was silent for one full minute. An overall VDB score was calculated by averaging complaining (weight = 1), disruptive vocalisations (weight = 2) and shouting (weight = 4). Tape recordings of VDB were performed for one hour per day (15 minutes before an intervention, 30 minutes during an intervention, and 15 minutes after an intervention), for 10 days for each of the interventions (music, videotape, social interaction, and no-intervention).

2) The frequency and context of VDB was examined in standardised observations with the SBMI which assesses nine types of VDB (shouting, screaming, or howling; constant requests for attention; repeating words; complaining or inappropriate verbalizations; cursing; verbal aggression; nonsense talk; hallucinations; and other disruptive verbal behaviours (e.g. groaning and singing)). If a VDB was manifested five or more times, a value of 6 (i.e. constant, C) or a value of 7 (i.e. extreme, E) were registered.

3) In informant ratings, nursing staff very familiar with the subjects rated their VDB using the Cohen-Mansfield Agitation Inventory (CMAI) on a seven-point scale (1: never manifests the behaviour; 7: manifests the behaviour several times/hour). Information was collected at the end of each intervention. Duration and frequency of appropriate verbal behaviours were also assessed.

In [Garland 2007](#), participants’ actual behaviours were observed by experienced, trained researchers with high interrater reliability. The researchers recorded the presence of behavioural disturbances at two-minute intervals before, during, and after exposure to the audiotapes with total observation periods lasting 45 minutes each. Behaviours were categorised into one of four groups: (i) physically aggressive agitation; (ii) physically non-aggressive agitation; (iii) verbally aggressive agitation; and (iv) verbally non-aggressive agitation. It was unclear which scale was used, although it was stated that the CMAI was used in the period before randomisation. The outcome measure was magnitude of change (a fall in physical/verbal agitation), based on differences in mean behaviour counts, before-during treatments and during-after treatments.

Quality of life, performance of activities of daily living and caregiver burden were not evaluated in any of the studies.

Excluded studies

Nine studies were excluded ([Characteristics of excluded studies](#)). One was a systematic review that evaluated SPT ([Zetteler 2008](#)). Two studies used a pretest-posttest quasi-experimental design to evaluate the efficacy of SPT ([Miller 2001](#); [Woods 1995](#)). Two studies were N-of-1 trials: one study included one participant ([O’Connor 2011](#)); and the second considered four cases ([Peak 2002](#)). One study used SPT consisting of presentation of old photographs and favourite music pieces of individuals with dementia, combined in personalised video channels via a television monitor. Participants were 23 people with Alzheimer’s disease but insufficient information was provided as to how the participants were allocated between the groups or whether the study was a cross-over trial ([Kajiyama 2007](#)). In one trial, the authors stated that the three interventions (SPT, music and usual care) were offered in an alternating pattern, but also that the six participants were offered a choice of interventions ([Cheston 2007](#)). One was an exploratory study that had the goal of collecting information about the delivery of Personal Message cards, as an alternative to simulated presence therapy ([Evans 2016](#)). The last study was an abstract of a protocol only ([Smith 2010](#)).

Risk of bias in included studies

The overall quality of the studies was considered low ([Characteristics of included studies](#)). [Figure 2](#) shows our judgments about each risk of bias item presented as percentages across all included studies whereas [Figure 3](#) presents a summary of the risk of bias assessments by risk of bias domain and by trial.

Figure 2. Risk of bias graph: review authors' judgements about each risk of bias item presented as percentages across all included studies.

Figure 3. Risk of bias summary: review authors' judgements about each risk of bias item for each included study.

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias
Camberg 1999	?	?	-	+	+	?	-
Cohen-Mansfield 1997	-	-	-	?	-	?	-
Garland 2007	?	?	-	+	+	?	?

Allocation

One cross-over trial was not randomised, although the four groups of participants did receive the treatments in different orders. There was no mention of allocation concealment. We judged this trial to be at high risk of selection bias (Cohen-Mansfield 1997). The remaining two cross-over trials reported random allocation of treatment order, but the method of sequence generation and allocation concealment were not described and so we judged the risk of bias in these domains to be unclear (Camberg 1999; Garland 2007). Hence, none of the studies was at low risk of selection bias.

Blinding

Given the types of intervention, participants and personnel could not be blinded in any of the studies. We therefore judged all the studies to be at high risk of performance bias.

The outcome assessor was blinded in two studies and these two were therefore considered at low risk of detection bias (Camberg 1999; Garland 2007). No information regarding the blinding of the outcome assessor was reported for Cohen-Mansfield 1997 so we judged it to have an unclear risk of detection bias.

Incomplete outcome data

Of the 60 participants included in Cohen-Mansfield 1997, 28 participants (47%) were excluded from the analysis (seven died before any data were collected, 11 became quiet, three refused to continue, and two were physically restrained and thus excluded). We therefore considered that the trial was at high risk of attrition bias.

In Camberg 1999, five out of 54 subjects were ill and could not participate in some sessions during the study. The authors reported that the available information was retained for all analyses. We judged it and Garland 2007 to be at low risk of attrition bias.

Selective reporting

None of the studies had a protocol available. We judged the risk of bias due to selective outcome reporting in all studies to be unclear.

Other potential sources of bias

All the studies were cross-over in design and were considered suitable for evaluating this type of intervention. However, two studies failed to report paired data and their analyses were performed on observations rather than participants, thus introducing a unit of analysis error. These studies were considered at high risk of bias (Camberg 1999; Cohen-Mansfield 1997). We were unsure whether Garland 2007 analysed the number of observations and therefore considered the analysis at unclear risk of bias.

Effects of interventions

See: [Summary of findings for the main comparison](#)

Primary outcomes

1. Behavioural and psychological symptoms

Although all participants had severe dementia, the studies included people with different behavioural disturbances. In particular, Cohen-Mansfield 1997 selected only those people with verbally disruptive behaviour. Therefore, the samples evaluated in the three studies were not comparable with each other. Moreover, behavioural and psychological symptoms were not uniformly defined across the trials and different tools were used to measure this outcome. These issues, together with the varied study designs, meant that we considered that data from the different studies could not be combined in meta-analyses.

Despite their cross-over design, none of the trials reported paired data. In addition, we were unable to extract the data specified in our protocol (Camberg 1999 and Cohen-Mansfield 1997 based their analysis on number of observations and provided P values; Garland 2007 provided the mean frequencies or differences in mean behaviour counts, before and during treatments, between SPT and control, but did not report standard deviations).

We considered the overall quality of the evidence to be very low due to serious concerns regarding the risk of bias (all studies were at high risk of bias in one or more domains, including high risks of selection, performance, attrition and other bias), and very serious concern regarding imprecision (all results were based on single studies with very small sample sizes) ([Summary of findings for the main comparison](#)).

We present below the results reported by study authors separately for each study. However, due to our quality assessment, it should be noted that we consider there to be a very high level of uncertainty about all of these results.

In Camberg 1999, staff generally used the audiotapes twice a day for each subject. Unfortunately, the audiotapes were employed when there was a pause in staff activity, instead of whenever subjects showed agitated or withdrawn behaviour.

The results were different according to the method used to assess the outcomes. Concerning direct observation, in 8000 observations by trained non-participants, audiotapes were employed infrequently (SimPres 11.7% and placebo 10.5% of the time). Moreover, interventions were seldom fully observed, given that a predetermined schedule was used to assess behaviour, independently of whether an intervention was administered or not. The authors stated that "A Latin Squares analysis was conducted with 20,667 direct observations obtained during the study period under the

“intention to treat” assumption”, which did not reveal an overall difference in either agitated or withdrawn behaviours. A further Latin Squares analysis, only of observations comprising an intervention (843 for SimPres and 780 for placebo), did not reveal any statistically significant main effects for agitation (SOAPD; PARS). In addition, analysis of frequency of happy expressions, using facial diagrams of mood (FACE) data, did not demonstrate a statistically significant difference between SimPres and usual care, but revealed that subjects had a happy expression more often during SimPres administration (11% of the time) than during placebo (2.2% of the time) ($P < 0.001$).

Staff observation logs documented 2547 observations of interventions for agitation by nursing staff (SimPres 592, placebo 600 and usual care 1355 observations). In contrast to the previous negative findings based on direct observation data, the authors reported that SimPres decreased agitation statistically significantly more often than placebo ($P < 0.001$) and usual care ($P < 0.001$). The same data collection method registered 1981 observations of interventions for withdrawn behaviour (SimPres 759, placebo 660, usual care 562 observations). Again, the staff reported that SimPres statistically significantly improved withdrawn behaviour twice as much as placebo ($P < 0.001$) and more often than usual care ($P < 0.001$).

Weekly staff survey data showed that there was no statistically significant difference in the amount of participants’ agitation between either SimPres and placebo or SimPres and usual care, although participants were slightly less agitated with usual care than with placebo ($P = 0.017$). Regarding mood evaluated with the MOSES scale, none of the interventions was statistically significantly superior. Concerning “interest” measured with the MOSES scale, there was an increase with SimPres compared to both placebo ($P = 0.008$) and usual care ($P = 0.001$) and no difference between placebo and usual care.

In [Cohen-Mansfield 1997](#), analysis of tape recording data revealed that VDB declined 46% during SPT (videotape recordings), 31% during music, 56% during the one-on-one social interaction and 16% during the “no-intervention”, compared to before the intervention. The authors stated that post hoc tests showed that each of the interventions were significantly different from “no-intervention”.

Standardised observational data, analysed using the SBMI for nine types of VDB, showed that compared to pre-intervention levels, SPT (videotape recordings) was best at decreasing repeating words (58%), hallucinations (56%) and shouting (50%). The same observational data revealed that music was most effective at reducing repeating words (48%), other VDB (48%) and hallucinations (40%). The most noticeable effects of social interaction was to decrease requests for attention (94%), repeating words (70%), and shouting (66%). Various VDB declined even with “no-intervention”, such as requests for attention (31%), repeating words (26%), hallucinations (21%), shouting (17%) and other VDB (15%). Nurses’ assessments (informant ratings) did not find that any in-

tervention produced a significant effect in any of the verbally agitated behaviours.

In [Garland 2007](#), concerning physical agitation there was no statistically significant difference between simulated presence and music during the interventions, whereas simulated presence was more effective than placebo ($P = 0.007$) and usual care ($P = 0.003$). The effect of music was not different from placebo, but was better than usual care ($P = 0.039$). Finally, the response to placebo was not statistically significant from the effect of usual care. At 15 minutes post-intervention, none of the treatments had a statistically significantly different effect on physical agitation. Regarding verbal agitation during the interventions, the effect of simulated presence was no different from music or placebo, but had a statistically significant action compared to usual care ($P = 0.37$). The response to music was indistinguishable from the effect of placebo and usual care, while placebo was more effective than usual care on verbal agitation ($P = 0.03$). Fifteen minutes after the interventions, simulated presence was no different from music, placebo or usual care. In contrast, the effect of music was statistically significantly different from placebo ($P = 0.007$), but not to usual care. Finally, the comparison between placebo and usual care did not reveal any statistically significant difference.

2. Quality of life

Quality of life was not assessed by any of the studies.

Secondary outcomes

1. Activities of daily living

Performance of activities of daily living was not assessed by any of the studies.

2. Caregiver burden

Caregiver burden was not assessed by any of the studies.

3. Dropouts

Dropouts were not reported by the studies.

Sensitivity and subgroup analysis

There were not enough data to conduct any of the planned subgroup or sensitivity analyses.

DISCUSSION

Summary of main results

The review included three studies of differing design, involving in total 144 participants with dementia. These people had different types of behavioural disturbances. The trials investigated the effects of SPT compared to usual care, personalised music, 'placebo' or a social interaction intervention. The experimental intervention in the three studies differed in terms of the tools used to carry out the treatment and the frequency with which it was administered. The outcomes related to behavioural disturbances were multiple, measured with different scales and with different time frames, which prevented us from performing a meta-analysis. Within each trial, the effect of SPT on behaviour, compared to usual care, was mixed and depended on the measure used. Two trials which included a personalised music intervention reported no significant differences between simulated presence and music on behavioural outcomes. Only behavioural outcomes were measured. The overall quality of the evidence was very low. We were unable to draw conclusions about the effectiveness of SPT for behavioural problems or other outcomes.

Overall completeness and applicability of evidence

The small number of trials included in this systematic review provided a very limited body of evidence on the effectiveness of SPT for dementia. There is currently a scarcity of randomised trials in this area, with only two studies using a randomised design (Camberg 1999; Garland 2007).

All the trials were conducted in nursing homes and their dates of publication ranged from 1999 to 2007. The mean age of the participants was above 70 years across all studies and women outnumbered men in all the trials. However, the number of people with vascular dementia or Lewy Body disease was limited in this review; this fact must be taken into account when interpreting the results. Most participants across the studies had moderate to severe dementia. Only one trial reported the concurrent use of psychotropic medications (Garland 2007). There is no data on comorbidity across the studies, so we do not know how this might have influenced the results.

Trial interventions showed large differences regarding the equipment used (audio or videotape), the duration of the intervention period, the selection of the contents of the audio/videotape, the frequency of administration, the washout period, and the professionals involved. Likewise, it was unclear how the optimal dosage or delivery (sessions, duration, etc.) was determined. Two studies used music and usual care as comparators (Cohen-Mansfield 1997; Garland 2007), while one study also used a social interaction intervention (Cohen-Mansfield 1997). Generally, usual care was not described.

There was also great variation in both the behaviour disturbances considered/evaluated and in the assessment procedures used to

measure the outcomes. For example, while Camberg 1999 proposed agitation (defined as an experience of an unpleasant state of excitement) and withdrawn behaviours, composed of manifestations of mood and "interest", Garland 2007 contemplated physically aggressive and non-aggressive agitation, and verbally aggressive and non-aggressive agitation, but described results only for physical and verbal agitation. Conversely Cohen-Mansfield 1997 considered only verbally disruptive behaviours defined as "verbal or vocal behaviours that are either repetitive, disruptive, or inappropriate to the circumstances in which they are manifested".

In terms of outcome assessment, in two trials several scales were used to measure the outcome (Camberg 1999 used OAS, AVAS, SCMAI for agitation and PARS, FACE and MOSES for withdrawal behaviour; Cohen-Mansfield 1997 used SBMI and CMAI in addition to direct observation), while Garland 2007 did not clearly report the scale used to assess physically and verbally agitated behaviours. All these factors hindered the possibility of performing a meta-analysis, including the fact that some studies did not control for other variables such as the use of psychotropic medication.

In terms of the results, authors from each trial provided conclusions in favour of the effect of SPT on behaviour-related outcomes. However, in some cases, the results varied depending on the outcome measured. For example, despite Camberg 1999 stating that SPT "can be effective in enhancing well-being and decreasing problem behaviors", SPT did not ameliorate agitation when measured with direct observations or weekly staff survey, whereas a statistically significant improvement was observed from staff observation logs. In another case, despite SPT being reported to be better than usual care, it was no more effective than personalised music (Garland 2007).

Other important outcomes of interest such as quality of life, activities of daily living, caregiver burden and dropouts were not assessed.

The most likely adverse effect we anticipated was worsening of agitation which could be detected in measures of the primary outcome. Worsening of agitation during treatment as a measure of adverse event was reported in Garland 2007. The authors of this study reported that some participants became more agitated and threw the headphones away. However, they did not provide the number of participants that experienced the outcome nor the frequency of agitation. No dropouts were reported in the other two trials.

Quality of the evidence

The quality of evidence was downgraded due to very serious concern in the domains of the risk of bias. Despite two of the studies being described as randomised, the method of sequence generation was not adequately reported and allocation concealment was not described (Camberg 1999; Garland 2007). Cohen-Mansfield 1997 used an alternation method between interventions and was

thus judged to be at high risk of selection bias. Therefore, none of the studies was rated at low risk of selection bias. In terms of blinding, none of the included studies was able to blind participants or treatment providers because of inherent difficulties in blinding non-pharmacological interventions such as SPT. Therefore, all the studies were exposed to performance bias. Two studies reported that the outcome assessors were blinded (Camberg 1999; Garland 2007), whereas Cohen-Mansfield 1997 did not report any statement and was thus rated unclear in terms of detection bias. During follow-up, a significant percentage (47%) of participants were excluded from analysis in Garland 2007 and the study was judged to be at high risk of attrition bias. No study protocols were available for any of the studies to compare the outcomes declared with those evaluated; and, furthermore, details of the results of the primary outcome were not clearly reported across all the studies thus hindering the possibility of performing meta-analyses and were therefore rated to have unclear risk of selective reporting bias.

A further downgrading by two levels of the evidence was performed for imprecision, since the studies had very few participants. In conclusion, the overall evidence for SPT to treat BPSD was rated very low in the [Summary of findings for the main comparison](#).

Potential biases in the review process

A protocol for this review was published (Abraham 2016; Abraham 2015). The review was conducted using comprehensive searches in several electronic databases to ensure all published trials were identified. The inclusion criteria for this review were intentionally broad to capture all studies. We did not limit the searches to a particular language and used pairs of review authors to independently examine and select studies. Authors of all the studies included were contacted to ascertain if any data were available since publication of the original studies, and to clarify unclear data. However, no responses were forthcoming.

Agreements and disagreements with other studies or reviews

One review that examined the effect of SPT in people with dementia identified seven studies (Zetteler 2008). Only three of those studies were included in our review, whereas the remaining trials were excluded, mostly due to study design. In Zetteler 2008's review, the author was able to obtain data from primary study authors, and hence to perform a meta-analysis of four primary studies (Cheston 2007; Garland 2007; Miller 2001; Woods 1995), only one of which was included in our review (Garland 2007). The results of this meta-analysis showed a mean effect size of 0.70, with a 95% CI of 0.38 to 1.02 ($P < 0.001$) with significant heterogeneity $I^2 = 72\%$ in favour of SPT for challenging behaviours in

dementia. One study contributed most to the heterogeneity, but its removal did not affect the significance of the results (Woods 1995). However, while Zetteler 2008's review underlined the lack of statistical power in the original studies, other methodological issues, such as risk of bias, were not assessed.

AUTHORS' CONCLUSIONS

Implications for practice

We do not have sufficient evidence to determine the effects of SPT for people with dementia on behavioural disturbances, quality of life, activities of daily living or caregiver burden.

Implications for research

The principal clinical trial registries do not have any ongoing or planned randomised trials on SPT for dementia.

This systematic review emphasises the need for more well-designed, larger trials before conclusions can be drawn on the effectiveness of SPT to treat behavioural problems in dementia. Many methodological issues should be addressed, such as the comparability of different interventions, randomisation and blinding of outcome assessors, given that most of the outcomes of interest are subjective. Different types of SPT interventions, with different methods of implementation, should be standardised. In addition, the choice of the comparator is important, since some non-pharmacological interventions (such as music therapy) can be more effective than others in the treatment of BPSD. Treatment effects for the different types of dementia and behavioural disturbances, as well as other variables, such as co-morbidity and different settings, also need to be investigated. Assessing other outcomes of interest, such as quality of life, activities of daily living, caregiver burden and dropouts would also be of value.

ACKNOWLEDGEMENTS

We thank Sue Marcus, the Managing Editor, and Jenny McCleery, Joint Co-ordinating Editor, Cochrane Dementia and Cognitive Improvement Group (CDCIG), for supporting the development of the protocol and the review; and Anna Noel-Storr, Trials Search Co-ordinator, (CDCIG), for developing the search strategy and identifying the full-text articles.

We want to thank also Jesús López Alcalde (Coordinador del Centro Colaborador Cochrane de la Comunidad de Madrid (CCC-CAM) and Javier Zamora Romero (Jefe de la Unidad de Bioestadística, Hospital Universitario Ramón y Cajal) for their assistance and support.

REFERENCES

References to studies included in this review

Camberg 1999 *{published data only}*

Camberg L, Woods P, Ooi WL, Hurley A, Volicer L, Ashley J, et al. Evaluation of Simulated Presence: a personalized approach to enhance well-being in persons with Alzheimer's disease. *Journal of the American Geriatrics Society* 1999;**47**(4):446–52. PUBMED: 10203120]

Cohen-Mansfield 1997 *{published data only}*

Cohen-Mansfield J, Werner P. Management of verbally disruptive behaviors in nursing home residents. *Journals of Gerontology. Series A, Biological Sciences and Medical Sciences* 1997;**52**(6):M369–77. PUBMED: 9402944]

Garland 2007 *{published data only}*

Garland K, Beer E, Eppingstall B, O'Connor DW. A comparison of two treatments of agitated behavior in nursing home residents with dementia: simulated family presence and preferred music. *American Journal of Geriatric Psychiatry* 2007;**15**(6):514–21. PUBMED: 17293386]

References to studies excluded from this review

Cheston 2007 *{published data only}*

Cheston R, Thorne K, Whitby P, Peak J. Simulated presence therapy, attachment and separation amongst people with dementia. *Dementia* 2007;**6**(3):442–9.

Evans 2016 *{published data only}*

Evans N, Cheston R, Harris N. Personal message cards: An evaluation of an alternative method of delivering simulated presence therapy. *Dementia-international journal of social research and practice* 2016;**6**:1703–15.

Kajiyama 2007 *{published data only}*

Kajiyama BK, Dib LRC, Tymchuk AJ, Boxer AL, Kixmiller JS, Olinsky CJ. Improving the quality of life for caregivers and care recipients with personalized video channels. *Clinical Gerontologist* 2007;**31**(1):95–100.

Miller 2001 *{published data only}*

Miller S, Vermeersch PE, Bohan K, Renbarger K, Kruep A, Sacre S. Audio presence intervention for decreasing agitation in people with dementia. *Geriatric Nursing (New York, N.Y.)* 2001;**22**(2):66–70. PUBMED: 11326212]

O'Connor 2011 *{published data only}*

O'Connor CM, Smith R, Nott MT, Lorang C, Mathews RM. Using video simulated presence to reduce resistance to care and increase participation of adults with dementia. *American Journal of Alzheimer's Disease and Other Dementias* 2011;**26**(4):317–25. PUBMED: 21624886]

Peak 2002 *{published data only}*

Peak JS, Cheston RI. Using simulated presence therapy with people with dementia. *Aging & Mental Health* 2002;**6**(1):77–81.

Smith 2010 *{published data only}*

Smith R, Campbell A, Lorang C, Mathews M, O'Connor C. Video technology and simulated presence to assist in

dementia care. *Australasian Journal on Ageing*. 2010; Vol. 29:40.

Woods 1995 *{published data only}*

Woods P, Ashley J. Simulated presence therapy: using selected memories to manage problem behaviors in Alzheimer's disease patients. *Geriatric Nursing* 1995;**16**(1):9–14.

Zetteler 2008 *{published data only}*

Zetteler J. Effectiveness of simulated presence therapy for individuals with dementia: a systematic review and meta-analysis. *Aging & Mental Health* 2008;**12**(6):779–85. PUBMED: 19023729]

Additional references

Abraha 2015a

Abraha I, Cherubini A, Cozzolino F, De Florio R, Luchetta ML, Rimland JM, et al. Deviation from intention to treat analysis in randomised trials and treatment effect estimates: meta-epidemiological study. *BMJ* 2015;**350**:h2445. [PUBMED: 26016488]

Abraha 2015b

Abraha I, Cruz-Jentoft A, Soiza RL, O'Mahony D, Cherubini A. Evidence of and recommendations for non-pharmacological interventions for common geriatric conditions: the SENATOR-ONTOP systematic review protocol. *BMJ open* 2015;**5**(1):e007488. [PUBMED: 25628049]

Abraha 2017

Abraha I, Rimland JM, Trotta FM, Dell'Aquila G, Cruz-Jentoft A, Petrovic M, et al. Systematic review of systematic reviews of non-pharmacological interventions to treat behavioural disturbances in older patients with dementia. The SENATOR-OnTop series. *BMJ open* 2017;**7**(3):e012759. [PUBMED: 28302633]

Algase 1996

Algase DL, Beck C, Kolanowski A, Whall A, Berent S, Richards K, et al. Need-driven dementia-compromised behavior: An alternative view of disruptive behavior. *American Journal of Alzheimer's Disease and Other Dementias* 1996;**11**(6):10–9.

Balshem 2011

Balshem H, Helfand M, Schunemann HJ, Oxman AD, Kunz R, Brozek J, et al. GRADE guidelines: 3. Rating the quality of evidence. *Journal of Clinical Epidemiology* 2011;**64**(4):401–6. [PUBMED: 21208779]

Borson 1997

Borson S, Raskind MA. Clinical features and pharmacologic treatment of behavioral symptoms of Alzheimer's disease. *Neurology* 1997;**48**(5 Suppl 6):S17–24. [PUBMED: 9153156]

Budnitz 2011

Budnitz DS, Lovegrove MC, Shehab N, Richards CL. Emergency hospitalizations for adverse drug events in older

- Americans. *New England Journal of Medicine* 2011;**365**(21):2002–12.
- Cameron 2003**
Cameron M, Lonergan E, Lee H. Transcutaneous electrical nerve stimulation (TENS) for dementia. *Cochrane Database of Systematic Reviews* 2003, Issue 3. [DOI: 10.1002/14651858.CD004032]
- Cerejeira 2012**
Cerejeira J, Lagarto L, Mukaetova-Ladinska EB. Behavioral and psychological symptoms of dementia. *Frontiers in Neurology* 2012;**3**:73.
- Chan 2004**
Chan AW, Hrobjartsson A, Haahr MT, Gotzsche PC, Altman DG. Empirical evidence for selective reporting of outcomes in randomized trials: comparison of protocols to published articles. *JAMA* 2004;**291**(20):2457–65. [PUBMED: 15161896]
- Cherubini 2012**
Cherubini A, Ruggiero C, Dell'Aquila G, Eusebi P, Gasperini B, Zengarini E, et al. Underrecognition and undertreatment of dementia in Italian nursing homes. *Journal of the American Medical Directors Association* 2012; **13**(8):759.e7–13. [PUBMED: 22727993]
- Chung 2002**
Chung JC, Lai CK, Chung PM, French HP. Snoezelen for dementia. *Cochrane Database of Systematic Reviews* 2002, Issue 4. [DOI: 10.1002/14651858.CD003152]
- Clyburn 2000**
Clyburn LD, Stones MJ, Hadjistavropoulos T, Tuokko H. Predicting caregiver burden and depression in Alzheimer's disease. *Journals of Gerontology: Series B, Psychological Sciences and Social Sciences* 2000;**55**(1):S2–13. [PUBMED: 10728125]
- Deeks 2011**
Deeks JJK, Higgins JPT, Altman DG (editors). Chapter 9: Analysing data and undertaking meta-analyses. In: Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available from handbook.cochrane.org.
- Endnote 2013 [Computer program]**
Thompson Reuters. EndNote reference management software. Thompson Reuters, 2013.
- Forbes 2013**
Forbes D, Thiessen EJ, Blake CM, Forbes SC, Forbes S. Exercise programs for people with dementia. *Cochrane Database of Systematic Reviews* 2013, Issue 12. [DOI: 10.1002/14651858.CD006489]
- Forbes 2014**
Forbes D, Blake CM, Thiessen EJ, Peacock S, Hawranik P. Light therapy for improving cognition, activities of daily living, sleep, challenging behaviour, and psychiatric disturbances in dementia. *Cochrane Database of Systematic Reviews* 2014, Issue 2. [DOI: 10.1002/14651858.CD003946]
- Forrester 2014**
Forrester LT, Maayan N, Orrell M, Spector AE, Buchan LD, Soares-Weiser K. Aromatherapy for dementia. *Cochrane Database of Systematic Reviews* 2014, Issue 2. [DOI: 10.1002/14651858.CD003150.pub2]
- Higgins 2011**
Higgins JPT, Altman DG, Sterne JAC (editors). Chapter 8: Assessing risk of bias in included studies. In: Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available from handbook.cochrane.org.
- Howard 2012**
Howard R, McShane R, Lindesay J, Ritchie C, Baldwin A, Barber R, et al. Donepezil and memantine for moderate-to-severe Alzheimer's disease. *New England Journal of Medicine* 2012;**366**(10):893–903.
- Kales 2015**
Kales HC, Gitlin LN, Lyketsos CG. Assessment and management of behavioral and psychological symptoms of dementia. *BMJ (Clinical Research Ed.)* 2015;**350**:h369. [PUBMED: 25731881]
- Katz 2007**
Katz I, de Deyn PP, Mintzer J, Greenspan A, Zhu Y, Brodaty H. The efficacy and safety of risperidone in the treatment of psychosis of Alzheimer's disease and mixed dementia: a meta-analysis of 4 placebo-controlled clinical trials. *International Journal of Geriatric Psychiatry* 2007;**22**(5):475–84.
- Kunik 2010**
Kunik ME, Snow AL, Davila JA, Steele AB, Balasubramanyam V, Doody RS, et al. Causes of aggressive behavior in patients with dementia. *Journal of Clinical Psychiatry* 2010;**71**(9):1145–52. [PUBMED: 20361896]
- Leendertse 2011**
Leendertse AJ, Van Den Bemt PM, Poolman JB, Stoker LJ, Egberts AC, Postma MJ. Preventable hospital admissions related to medication (HARM): cost analysis of the HARM study. *Value in Health* 2011;**14**(1):34–40.
- Lopez 2013**
Lopez OL, Becker JT, Chang YF, Sweet RA, Aizenstein H, Snitz B, et al. The long-term effects of conventional and atypical antipsychotics in patients with probable Alzheimer's disease. *American Journal of Psychiatry* 2013;**170**(9):1051–8.
- Lyketsos 2007**
Lyketsos CG. Neuropsychiatric symptoms (behavioral and psychological symptoms of dementia) and the development of dementia treatments. *International Psychogeriatrics / IPA* 2007;**19**(3):409–20. [PUBMED: 17346363]
- Macura 2010**
Macura A, Abraha I, Kirkham J, Gensini GF, Moja L, Iorio A. Selective outcome reporting: telling and detecting true lies. The state of the science. *Internal and Emergency Medicine* 2010;**5**(2):151–5. [PUBMED: 20300879]

Miesen 1993

Miesen BML. Alzheimer's disease, the phenomenon of parent fixation and Bowlby's attachment theory. *International Journal of Geriatric Psychiatry* 1993;**8**(2): 147–53.

Moniz-Cook 2012

Moniz-Cook ED, Swift K, James I, Malouf R, De Vugt M, Verhey F. Functional analysis-based interventions for challenging behaviour in dementia. *Cochrane Database of Systematic Reviews* 2012, Issue 2. [DOI: 10.1002/14651858.CD006929.pub2; PUBMED: 22336826]

Neal 2003

Neal M, Barton Wright P. Validation therapy for dementia. *Cochrane Database of Systematic Reviews* 2003, Issue 3. [DOI: 10.1002/14651858.CD001394]

O'Connor 2009

O'Connor DW, Ames D, Gardner B, King M. Psychosocial treatments of psychological symptoms in dementia: a systematic review of reports meeting quality standards. *International Psychogeriatrics* 2009;**21**(2):241–51. [PUBMED: 19138459]

Olin 2002

Olin J, Schneider L. Galantamine for Alzheimer's disease. *Cochrane Database of Systematic Reviews* 2002, Issue 3. [DOI: 10.1002/14651858.CD001747]

Opie 1999

Opie J, Rosewarne R, O'Connor DW. The efficacy of psychosocial approaches to behaviour disorders in dementia: a systematic literature review. *Australian and New Zealand Journal of Psychiatry* 1999;**33**(6):789–9.

Peng 2007

Peng WN, Zhao H, Liu ZS, Wang S. Acupuncture for vascular dementia. *Cochrane Database of Systematic Reviews* 2007, Issue 2. [DOI: 10.1002/14651858.CD004987.pub2]

Prince 2013

Prince M, Bryce R, Albanese E, Wimo A, Ribeiro W, Ferri CP. The global prevalence of dementia: a systematic review and meta-analysis. *Alzheimer's & Dementia* 2013;**9**(1):63–75 e2.

Review Manager 2014 [Computer program]

Nordic Cochrane Centre, The Cochrane Collaboration. Review Manager 5 (RevMan 5). Version 5.3. Copenhagen: Nordic Cochrane Centre, The Cochrane Collaboration, 2014.

Rottenkolber 2011

Rottenkolber D, Schmiedl S, Rottenkolber M, Farker K, Salje K, Mueller S, et al. Adverse drug reactions in Germany: direct costs of internal medicine hospitalizations. *Pharmacoepidemiology and Drug Safety* 2011;**20**(6):626–34.

Salzman 2008

Salzman C, Jeste DV, Meyer RE, Cohen-Mansfield J, Cummings J, Grossberg GT, et al. Elderly patients with dementia-related symptoms of severe agitation and aggression: consensus statement on treatment options,

clinical trials methodology, and policy. *Journal of Clinical Psychiatry* 2008;**69**(6):889–98. [PUBMED: 18494535]

Savovic 2012

Savovic J, Jones H, Altman D, Harris R, Juni P, Pildal J, et al. Influence of reported study design characteristics on intervention effect estimates from randomised controlled trials: combined analysis of meta-epidemiological studies. *Health Technology Assessment (Winchester, England)* 2012;**16**(35):1–82. [PUBMED: 22989478]

Savva 2009

Savva GM, Zaccai J, Matthews FE, Davidson JE, McKeith I, Brayne C. Prevalence, correlates and course of behavioural and psychological symptoms of dementia in the population. *British Journal of Psychiatry* 2009;**194**(3):212–9.

Schünemann 2011a

Schünemann HJ, Oxman AD, Higgins JPT, Vist GE, Glasziou P, Guyatt GH. Chapter 11: Presenting results and 'Summary of findings' tables. In: Higgins JPT, Green S (editors), *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available from handbook.cochrane.org.

Schünemann 2011b

Schünemann HJ, Oxman AD, Vist GE, Higgins JPT, Deeks JJ, Glasziou P, et al. Chapter 12: Interpreting results and drawing conclusions. In: Higgins JPT, Green S (editors), *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available from handbook.cochrane.org.

Sink 2005

Sink KM, Holden KE, Yaffe K. Pharmacological treatment of neuropsychiatric symptoms of dementia: a review of the evidence. *JAMA* 2005;**293**(5):596–608.

Smith 2004

Smith M, Gerdner LA, Hall GR, Buckwalter KC. History, development, and future of the progressively lowered stress threshold: a conceptual model for dementia care. *Journal of the American Geriatrics Society* 2004;**52**(10):1755–60. [PUBMED: 15450057]

Spalletta 2010

Spalletta G, Musicco M, Padovani A, Rozzini L, Perri R, Fadda L, et al. Neuropsychiatric symptoms and syndromes in a large cohort of newly diagnosed, untreated patients with Alzheimer disease. *American Journal of Geriatric Psychiatry* 2010;**18**(11):1026–35. [PUBMED: 20808086]

Steinberg 2008

Steinberg M, Shao H, Zandi P, Lyketsos CG, Welsh-Bohmer KA, Norton MC, et al. Point and 5-year period prevalence of neuropsychiatric symptoms in dementia: the Cache County Study. *International Journal of Geriatric Psychiatry* 2008;**23**(2):170–7. [PUBMED: 17607801]

Stern 1994

Stern Y, Albert M, Brandt J, Jacobs DM, Tang MX, Marder K, et al. Utility of extrapyramidal signs and psychosis as

- predictors of cognitive and functional decline, nursing home admission, and death in Alzheimer's disease: prospective analyses from the Predictors Study. *Neurology* 1994;**44**(12): 2300–7.
- Thies 2012**
Thies W, Bleiler. 2012 Alzheimer's disease facts and figures. *Alzheimer's & Dementia* 2012;**8**(2):131–68. [PUBMED: 22404854]
- Van Den Wijngaart 2007**
Van Den Wijngaart MA, Vernooij-Dassen MJ, Felling AJ. The influence of stressors, appraisal and personal conditions on the burden of spousal caregivers of persons with dementia. *Aging & Mental Health* 2007;**11**(6):626–36. [PUBMED: 18074250]
- Viggo Hansen 2006**
Viggo Hansen N, Jørgensen T, Ørtenblad L. Massage and touch for dementia. *Cochrane Database of Systematic Reviews* 2006, Issue 3. [DOI: 10.1002/14651858.CD004989.pub2]
- Vink 2004**
Vink AC, Birks JS, Bruinsma MS, Scholten RJ. Music therapy for people with dementia. *Cochrane Database of Systematic Reviews* 2004, Issue 3. [DOI: 10.1002/14651858.CD003477.pub2]
- Wimo 2013**
Wimo A, Jonsson L, Bond J, Prince M, Winblad B. The worldwide economic impact of dementia 2010. *Alzheimer's & Dementia* 2013;**9**(1):1–11. [PUBMED: 23305821]
- Wood 2008**
Wood L, Egger M, Gluud LL, Schulz KF, Juni P, Altman DG, et al. Empirical evidence of bias in treatment effect estimates in controlled trials with different interventions and outcomes: meta-epidemiological study. *BMJ (Clinical Research Ed.)* 2008;**336**(7644):601–5. [PUBMED: 18316340]
- Woods 2009**
Woods RT, Bruce E, Edwards RT, Hounscome B, Keady J, Moniz-Cook ED, et al. Reminiscence groups for people with dementia and their family carers: pragmatic eight-centre randomised trial of joint reminiscence and maintenance versus usual treatment: a protocol. *Trials* 2009;**10**:64. [PUBMED: 19642992]
- References to other published versions of this review**
- Abraha 2015**
Abraha I, Rimland Joseph M, Lozano-Montoya I, Dell'Aquila G, Vélez-Díaz-Pallarés M, Trotta Fabiana M, et al. Simulated presence therapy for dementia. *Cochrane Database of Systematic Reviews*. John Wiley & Sons, Ltd, 2015, issue 9. [DOI: 10.1002/14651858.CD011882; CD011882]
- Abraha 2016**
Abraha I, Rimland JM, Lozano-Montoya I, Dell'Aquila G, Velez-Diaz-Pallares M, Trotta FM, et al. Simulated presence therapy for dementia: a systematic review protocol. *BMJ open* 2016;**6**(5):e011007. [PUBMED: 27169742]
- * Indicates the major publication for the study

CHARACTERISTICS OF STUDIES

Characteristics of included studies *[ordered by study ID]*

Camberg 1999

Methods	<p>A Latin-Square double-blind cross-over design.</p> <p>Method of allocation: random allocation.</p> <p>Date: not reported.</p> <p>Assessor blinding: yes.</p> <p>Participant blinding: nursing staff and observers were blinded. Participants were not blinded due to the nature of the intervention</p> <p>Loss to follow-up: unclear.</p> <p>Intention-to-treat analysis: yes. Subsequently an intervention-specific analysis was made</p> <p>Post-program follow up: not reported.</p>
Participants	<p>Location: 9 nursing homes in Eastern Massachusetts and Southern New Hampshire (5 for-profit facilities, 3 nonprofit, 1 government owned. 15 nursing units (9 Alzheimer's special care unit, 4 dementia units, 2 mixed units)</p> <p>Geographical area: USA.</p> <p>Sample: 54 subjects; 94.7% white.</p> <p>Age: mean age 82.7 years (SD 7.5 years).</p> <p>Sex: 48F, 77.2%.</p> <p>Baseline: average baseline cognitive measures indicated severe cognitive impairment (MMSE: 5.1, SD $\pm$ 4.4; TSI:13.1 SD $\pm$ 6.9) and low functional ability (BANS: 13.2, SD $\pm$ 4.0; ADL Self-Performance Scale: 2.7, SD $\pm$ 1.2)</p> <p>Inclusion criteria: a documented diagnosis of Alzheimer's disease and related dementias (ADRD); age 50 years or older; the presence of at least 1 agitated behaviour per day of those listed on the short form of the Cohen-Mansfield Agitation Inventory Scale (SCMAI) or 1 indication of withdrawn behaviour defined either by "sounding sad" or "crying" that occurred at least "often"; or "seldom" interested in activities, social interaction, or in their immediate environment, from the Multidimensional Observation Scale for Elderly Subjects (MOSES); medically stable; resident in the current nursing home for at least 3 months; and no planned discharge. Subjects must have a verbal interactive capacity judged by the appropriateness of verbal responses and pauses during a conversation when 5 questions and comments were introduced. Individuals were included if they responded verbally, if their verbal response was appropriate at least some of the time, and if they paused to listen, as in a conversation</p> <p>Exclusion criteria: nonverbal patients, potential subjects who did not tolerate listening through headphones for 5 minutes to a nonpersonalised interactive conversation tape, or who were known to have a severe hearing impairment or premorbid history of psychiatric illness</p>
Interventions	<p>The interventions were: (1) simulated presence therapy, (2) placebo and (3) usual care</p> <p>Interventions: simulated presence: an interactive audio tape that contains 1 side of a live telephone conversation with a family member or surrogate, rich in selected memories and positive emotions</p> <p>Placebo: "The placebo audio tape was a recording of a person reading emotionally neutral articles from the newspaper"</p> <p>Usual care: the routine interventions nursing home staff used to manage behaviour, such</p>

	<p>as staff interactions, redirection, or physical restraints</p> <p>Application of intervention: nursing staff on the study units</p> <p>Equipment: the audio tape was played using a headset and an auto-reverse tape recorder enclosed in a hip pack</p> <p>Frequency: twice a day (Monday through Friday) in place of their usual intervention</p> <p>Reps: whenever a study subject exhibited an agitated or withdrawn behaviour</p> <p>Duration: each treatment continued for 17 days over a 4-week course with a 10-day washout period following each treatment</p> <p>Setting: nursing home.</p> <p>Supervision: a study monitor was assigned to spend 20 hours per week at each facility to ensure adherence to the protocol and to provide feedback to staff</p> <p>Adherence: subject acceptance of the study intervention was high. 23 subjects (43%) accepted the intervention 100% of the time, and only 5 subjects (9%) refused it more than 50% of the time</p>
Outcomes	<p>Several measures and 3 data collection methods were used as validity checks to identify negative and positive aspects of well-being: (1) direct observations by trained nonparticipant observers; (2) daily staff observation logs of participants' responses to interventions used; and (3) weekly behavioural rating surveys by nursing staff</p> <p>Direct observation of subject behaviour was done by trained non-participant observers using a structured time-sampling technique. Observers were blinded to the study intervention: each subject was observed at least 3 hours and 20 minutes per week between 9 a.m. and 7 p.m. Each observation block was 20 minutes long, divided into four 5-minute observation intervals</p> <p>Daily staff observation logs: when the staff decided to intervene, staff documented the target behaviour, i.e., agitation or withdrawal; the specific intervention (tape, restraint, staff interaction); the duration of the intervention; and the participant's response (improved, worsened, or remained the same). Nursing home staff documented daily (Monday to Friday) when an intervention was used for agitated or withdrawn behaviours</p> <p>Weekly staff survey (administered to a nursing staff member who had cared for the subject for at least 3 days during the preceding week) used 2 behavioural rating scales: (1) the short form of the Cohen-Mansfield Agitation Inventory (SCMAI) to record staff recall of the frequency of agitated behaviours; and (2) selected, modified items from the Multidimensional Observation Scale for Elderly Subjects (MOSES) to record the subject's mood and interest</p> <ol style="list-style-type: none"> 1. Overall behavioural and psychological symptoms. 2. Agitation: direct observations by trained nonparticipant observers: (1) Scale for the Observation of Agitation in Persons with Dementia (SOAPD), a 7-item Observed Agitation Scale (developed by the researchers); (2) an agitation visual analog scale (AVAS) <p>Daily staff observation logs.</p> <p>Weekly staff survey: (1) the short form of the Cohen-Mansfield Agitation Inventory (SCMAI) to record the frequency of agitated behaviours</p> <ol style="list-style-type: none"> 3. Depression: direct observations by trained nonparticipant observers: (1) 2 Positive Affect items, "interest" and "pleasure", from the Philadelphia Geriatric Center Affect Rating Scale (PARS); (2) a withdrawal visual analog scale (WVAS); and (3) facial diagrams of mood (FACE) <p>Daily staff observation logs.</p> <p>Weekly staff survey: (1) selected, modified items from the Multidimensional Observation Scale for Elderly Subjects (MOSES) to record the subject's mood and interest</p>

	4. Anxiety: not assessed. 5. Quality of life: not assessed. 6. Activities of daily living: not assessed. 7. Caregiver burden: not assessed. 8. Dropouts: not assessed.	
Notes	Funding: government (VA) and private (Harvard University) Declarations of interest: not stated	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	The method of randomisation was not clearly reported. "The treatments were applied in a restrictively randomised manner to each facility grouping"
Allocation concealment (selection bias)	Unclear risk	No statement was provided about allocation concealment.
Blinding of participants and personnel (performance bias) All outcomes	High risk	"The placebo audio tape was made to look like the SimPres treatment in routine use. Staff were asked not to listen to the audio tapes". Participants were not blinded due to the nature of the intervention
Blinding of outcome assessment (detection bias) All outcomes	Low risk	"Observers were blinded to the study intervention".
Incomplete outcome data (attrition bias) All outcomes	Low risk	Withdrawals are clearly described. "During the course of the study five subjects were too ill to participate in at least one of the treatment phases". "Their available information was retained for all analyses"
Selective reporting (reporting bias)	Unclear risk	Protocol not available. Details of the results especially regarding agitated behaviour were not clearly reported: the study reported the prevalence of non-agitated participants (25%); however data were not completely available to perform a meta-analysis
Other bias	High risk	The study was a cross-over study and it was considered suitable for evaluating the intervention. However, the study failed to report paired data and the analyses were per-

		formed by observation rather than by participant in at least 2 of the 3 measurements used (Staff observation logs and Direct observation)
--	--	---

Cohen-Mansfield 1997

Methods	<p>Method of allocation: non-randomised cross-over design.</p> <p>Cross-over controlled study with 4 interventions: 1) family-generated videotape (SPT); 2) audiotape recording of music; 3) one-to-one social interaction; and 4) control no-intervention phase</p> <p>Participant blinding: unblinded due to the nature of the intervention</p> <p>Assessor blinding: not reported.</p> <p>Loss to follow-up: 5 of the remaining 53 participants died before completing the interventions, 11 became quiet, 3 refused to continue their participation, and 2 were physically restrained and thus excluded</p> <p>Intention-to-treat analysis: per protocol.</p> <p>Post-program follow-up: not reported.</p>
Participants	<p>Location: nursing home.</p> <p>N for each group: 32 residents in total (cross-over trial with 4 groups)</p> <p>Sample: 60 participants (32 analysed); 4 groups.</p> <p>Age: 86.8 ± 1.16.</p> <p>Sex: 81% female.</p> <p>Baseline: Activities of Daily Living (ADL) assessed by Lawton and Brody's scale; Physical Self Maintenance Scale (PSMS): 4.0; Brief Cognitive Rating Scale (BCRS): 5.5.</p> <p>Rest of demographic characteristics (marital status, length of stay in the NH, diagnosis of dementia) are shown in table 1</p> <p>Inclusion criteria: nursing home residents with VDB at least several times a day (in the results the authors state 97% of the patients were diagnosed as suffering from dementia)</p> <p>Exclusion criteria: physically restrained.</p>
Interventions	<p>Participants were assigned into 4 groups, each containing 3 interventions:</p> <ol style="list-style-type: none"> 1) Medical examination. Special attention to identify underlying causes of physical discomfort; 2) Music half-hour audio; tapes according to the participants' preferences were prepared (9 participants = classical music, 8 = Jewish music, 4 = big band music, 4 = old popular music, 2 = opera, 3 = show tunes, and 2 = country music); 3) Video tapes: the relatives of all the participants were asked to prepare a videotape. For these participants, a research assistant, a nursing assistant, and a volunteer prepared the videotapes. Family members received general guidelines to prepare the videotapes but the specific content and manner of conveying it were left to them; 4) One-to-one social interaction: an extensive list of possible activities was provided <p>Application of intervention: unclear.</p> <p>Equipment: not reported.</p> <p>Frequency: the interventions were performed at the time of the day at which the resident was most verbally disruptive and were performed only if the resident was verbally</p>

	<p>disruptive</p> <p>Reps: not reported.</p> <p>Duration: each intervention lasted for 30 consecutive minutes. Each intervention lasted 2 weeks, followed by a washout week</p> <p>Setting: nursing home.</p> <p>Recording: "Participants were observed for an hour a day: 15 min before the intervention, during the 30-min intervention, and for 15 min after the intervention. Each observation lasted 3 consecutive mi. Total = nine 3-min observations/hour"</p> <p>Supervision: unclear.</p> <p>Adherence: unclear.</p> <p>Participants were assigned into 4 groups, each containing 3 intervention conditions and 1 control "no-intervention" condition. In each group the order of treatments was alternated, so that by the end of the evaluation period all groups were exposed to all the treatments and to the no-intervention phase. Each treatment continued for 2 consecutive weeks and was followed by a washout week</p> <p>Before exposure to any of the interventions, all participants underwent a detailed medical examination, by their attending physician, to identify any underlying causes of pain or physical discomfort</p> <p>1) SPT videotape. The relatives of all the participants were asked to prepare a videotape. Family members received general guidelines to prepare the videotapes but the specific content and manner of conveying it were left to them;</p> <p>2) Music. 30-minute audiotapes based on the participants' preferences were prepared and played to the participants in their rooms. 9 participants were exposed to classical music, 8 participants to Jewish music, 4 to big band music, 4 to old popular music, 2 to opera, 3 to show tunes, and 2 to country music;</p> <p>3) One-to-one social interaction. A manual was prepared to guide the social interaction, which included information on how to interact with a nursing home resident, how to start an activity and a broad list of possible activities ordered by the degree of involvement, from high to low, required of the resident. A "box of activities", comprising games, pictures, balls, books, etc. was assembled. Alternatively, the following activities were provided: (i) conversation with a research assistant; (ii) motion exercise (e.g., tossing a foam ball, hand and arm movement); (iii) sensory stimulation (a sensory kit including different fabrics, school supplies, health aids, make-up, spices, soaps, etc. was available to be presented to the residents to stimulate touch, sight, and smell); (iv) manual activities (such as making a collage, completing a simple puzzle, clipping coupons)</p> <p>Application of intervention: unclear.</p> <p>Equipment: not reported.</p> <p>Frequency: the intervention was performed at the time of the day during which the resident was most verbally disruptive (based on nurses' reports) and performed only if the participant was verbally disruptive</p> <p>Reps: not reported.</p> <p>Duration: each intervention lasted 30 minutes per day for 2 consecutive weeks and was followed by a 1-week washout</p> <p>Setting: nursing homes.</p> <p>Recording: participants were observed 1 hour/day (15 minutes before, 30 minutes during, and 15 minutes after, an intervention), composed of nine, 3-minute staggered observations</p> <p>Supervision: unclear.</p> <p>Adherence: unclear.</p>
--	--

Outcomes	<p>1. Overall behavioural and psychological symptoms:</p> <p>2. VDB: 3 different methods were used to assess VDB: tape recordings; standardised observations; and Informant ratings</p> <p>Tape recordings: VDB duration was registered with a tape recorder located near a participant. The tape was coded by a trained research assistant using a computer program specifically developed for coding video and audiotapes. An episode of VDB was considered to be finished if the resident was silent for 1 full minute. An overall VDB score was calculating by averaging complaining (weight = 1), disruptive vocalisations (weight = 2) and shouting (weight = 4). Tape recordings of VDB were performed 1 hour/day (15 min before an intervention, 30 min during an intervention, and 15 mins after an intervention), for 10 days for each of the interventions (music, videotape, social interaction, and no-intervention)</p> <p>Standardised observations: the frequency and context of screaming was examined with the Screaming Behavioural Mapping Instrument (SBMI) containing 9 types of VDB (shouting, screaming, or howling; constant requests for attention; repeating words; complaining or inappropriate verbalizations; cursing; verbal aggression; nonsense talk; hallucinations; and other disruptive verbal behaviours (e.g. groaning and singing))</p> <p>Informant ratings: nursing staff well-acquainted with the subjects rated their VDB using the Cohen-Mansfield Agitation Inventory (CMAI) on a 7-point scale (1: never manifests the behaviour; 7: manifests the behaviour several times/hour). Information was collected at the end of each intervention. Duration and frequency of appropriate verbal behaviours were also assessed</p> <p>3. Depression: not assessed.</p> <p>4. Anxiety: not assessed.</p> <p>5. Quality of life: not assessed.</p> <p>6. Activities of daily living: functional independence (Lawton and Brody's Physical Self-Maintenance Scale (PSMS))</p> <p>7. Caregiver burden: not assessed.</p> <p>8. Dropouts: 5 of the remaining 53 participants (9%) died before completing the interventions, 11 (21%) became quiet, 3 (6%) refused to continue their participation, and 2 (4%) were physically restrained and excluded based on previous findings showing that the manifestation of VDB increased when nursing home residents were physically restrained</p>
Notes	<p>Funding: this study was supported by National Institute on Aging Grant R01 AG-10642</p> <p>Declarations of interest: not stated.</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	High risk	"Participants were assigned to four groups, each containing three intervention conditions and one "no-treatment" condition. In each group, the order of treatments was alternated, so that by the end of the evaluation period all groups were exposed to all the treatments and to the no-treatment period"

Allocation concealment (selection bias)	High risk	“Participants were assigned to four groups, each containing three intervention conditions and one “no-treatment” condition. In each group, the order of treatments was alternated, so that by the end of the evaluation period all groups were exposed to all the treatments and to the no-treatment period”
Blinding of participants and personnel (performance bias) All outcomes	High risk	Not possible due to the nature of the intervention.
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not stated.
Incomplete outcome data (attrition bias) All outcomes	High risk	28 participants (47%) were excluded from analysis, 11 participants (18.3%) were excluded because they became quiet (they had a had a significantly higher cognitive status than those who continued manifesting VDB (3.8 and 5.5 on the BCRS respectively, $t_m = 2.8$, $P < 0.01$)
Selective reporting (reporting bias)	Unclear risk	Protocol not available. Outcomes not clearly specified in the aim section. Data were not completely available to perform a meta-analysis
Other bias	High risk	The study was a cross-over study and it was considered suitable for evaluating the intervention. However, the study failed to report paired data and the analysis for VDB was based on number of observations rather than number of participants. The frequency of VDB was examined using the SBMI. A total of 11,520 3-min observations were made (including observation before, during, and after the interventions) . The percentage of observations in which VDB occurred was calculated

Methods	<p>Design: repeated measures with randomised cross-over, single blind</p> <p>Date: not stated.</p> <p>Method of allocation: not stated.</p> <p>Assessor blinding: unclear.</p> <p>Participant blinding: no.</p> <p>Loss to follow-up: not stated.</p> <p>Intention-to-treat analysis: not stated.</p> <p>Post-program follow-up: not stated.</p>
Participants	<p>Location: nursing home.</p> <p>Geographical area: Australia.</p> <p>Sample: 30 subjects.</p> <p>Age: mean 79 years (66 to 93 years).</p> <p>Gender: F 19, 63%.</p> <p>Baseline: Mini-mental State Examination (MMSE; mean: 2.5, range: 0-12). All subjects were prescribed some type of psychotropic medication, often in combination</p> <p>Inclusion criteria: residents in the nursing home, for at least 3 months, with a dementia diagnosis and with 1 or more significantly disruptive behaviours, measured with the CMAI, occurring at least several times per day in the preceding 2 weeks</p> <p>Exclusion criteria: 1) evidence that a behaviour was due to a treatable medical or psychiatric condition; 2) pronounced deafness; 3) no response to conversational prompts from a research psychologist; 4) unwillingness to wear headphones on 3 consecutive trials; and 5) refusal by a family member to help prepare audiotapes</p>
Interventions	<p>3 arms (simulated family presence, preferred music, neutral audiotape (placebo)) and usual care</p> <p>Participants were randomly assigned to 1 of 3 groups and alternated through treatments, with a 2-day washout between treatments</p> <p>Active group 1: simulated family presence: semi-structured interview, with a trained psychologist, about participants' earlier lives. The interview was planned to emulate a telephone conversation about special memories, loved ones and family anecdotes</p> <p>Active group 2: tape containing music based on family members' reports of participants' preferences</p> <p>Control group: placebo tape: a psychologist reading from a gardening book</p> <p>Usual care: observations were recorded for 45 minutes on each of 3 days in week 1 of the study at the times of peak agitation, nominated by nursing staff</p> <p>Application of intervention: not reported.</p> <p>Equipment: portable cassette player with headphones.</p> <p>Frequency: once per day.</p> <p>Reps: 3 days during weeks 2, 3 and 4.</p> <p>Duration: 15 minutes.</p> <p>Washout period: 2 days.</p> <p>Setting: nursing homes.</p> <p>Supervision: not stated.</p> <p>Adherence: not stated.</p>
Outcomes	<p>1. Overall behavioural and psychological symptoms.</p> <p>2. Agitation: trained researchers recorded if specific behaviours (physically aggressive agitation, physically nonaggressive agitation, verbally aggressive agitation and verbally nonaggressive agitation) were present/absent at 2-minute intervals before, during and</p>

	after exposure to the 15-minute tapes. Total observations periods lasted for 45 minutes each Frequency of specific behaviours categorised using the CMAI (physically aggressive agitation, physically non-aggressive agitation, verbally aggressive agitation and verbally non-aggressive agitation) before, during and 15 minutes after treatment Magnitude of change (a fall in physical/verbal agitation): differences in Mean Behaviour Counts before-during treatments and during-after treatments 3. Depression: not assessed. 4. Anxiety: not assessed. 5. Quality of life: not assessed. 6. Activities of daily living: not assessed. 7. Caregiver burden: not assessed. 8. Dropouts: not assessed.	
Notes	Funding: this work was supported in part by the Rotary Heath Research Fund Declarations of interest: not stated	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	“Participants were allocated randomly to one of the three groups (A,B,C) and rotated through treatments” (shown in Table 1)
Allocation concealment (selection bias)	Unclear risk	The rotation through treatments is predefined. Table 1
Blinding of participants and personnel (performance bias) All outcomes	High risk	- Patients are not blinded due to the nature of the intervention but this fact may not affect the outcomes. - Treatments were applied once a day. No other information is given to describe if the personnel were blinded for the intervention - It is stated that researchers were blinded to the tape’s contents; however, we assumed by our reading of the previous paragraph that researchers were those who recorded specific behaviours and not the personnel in charge of applying the intervention
Blinding of outcome assessment (detection bias) All outcomes	Low risk	“Researchers were blinded to the tapes’ contents”. “Our observers, who were blinded to the tape,..”
Incomplete outcome data (attrition bias) All outcomes	Low risk	All participants completed the trial. No loss to follow-up.

Garland 2007 (Continued)

Selective reporting (reporting bias)	Unclear risk	Protocol not available. The final data were not completely available to perform a meta-analysis
Other bias	Unclear risk	The study was a cross-over study and it was considered suitable for evaluating the intervention. However, it was unclear how the data were reported. The study provided first an overall number of observations (2880 target behaviours) that were classified as physically aggressive behaviours (3.8%), physically agitated non-aggressive behaviours (64.8%), and verbally agitated aggressive and non-aggressive behaviours (31.4%) but we were unsure if these numbers were used as the basis for analysis

Characteristics of excluded studies [ordered by study ID]

Study	Reason for exclusion
Cheston 2007	This trial was not randomised or quasi-randomised. Participants were offered a choice of interventions: “Two control conditions were used creating an A1, B, A2, C, A3 and A1, C, A2, B, A3 design, in which A represents a baseline condition (usual care/no intervention), B represents the SPT intervention and C is the opportunity to listen to a favourite music tape. The order in which the SPT and music conditions were offered to participants alternated.”
Evans 2016	This study aimed to collect information about the way in which Personal Message cards were used as an alternative method of delivering Simulated Presence therapy. This was not an intervention study
Kajiyama 2007	In this study SPT was evaluated as presentation of old photographs and favourite music pieces of individuals with dementia, combined in personalized video channels via a television monitor in 23 subjects with Alzheimer disease. However, insufficient information was provided as to how the participants were allocated between the groups or whether the study was a cross-over trial
Miller 2001	A pretest-post test quasi-experimental design that evaluated simulated presence therapy
O'Connor 2011	The study evaluated simulated presence therapy for 1 participant (N-of-1 trial)
Peak 2002	N-of-1 trial of 4 cases.
Smith 2010	An abstract of a protocol only.
Woods 1995	A pretest-posttest quasi-experimental design that evaluated simulated presence therapy

(Continued)

Zetteler 2008	A systematic review that considered simulated presence therapy for people with dementia
---------------	---

DATA AND ANALYSES

This review has no analyses.

APPENDICES

Appendix I. Sources searched and search strategy

Source (searched August 2015 and February 2016)	Search strategy	Hits retrieved
1. ALOIS (www.medicine.ox.ac.uk/alois)	"simulated presence therapy" OR SPT	August 2015: 3 February 2015: 0
2. MEDLINE In-process and other non-indexed citations and MEDLINE 1946-present (Ovid SP) [Latest search: 9 February 2016]	1. exp Dementia/ 2. Delirium, Dementia, Amnestic, Cognitive Disorder/ 3. dement*.mp. 4. alzheimer*.mp. 5. (lewy* adj2 bod*).mp. 6. (chronic adj2 cerebrovascular).mp. 7. ("organic brain disease" or "organic brain syndrome").mp. 8. (cerebr* adj2 deteriorat*).mp. 9. (cerebral* adj2 insufficient*).mp. 10. or/1-9 11. "Simulated presence therapy".ti,ab. 12. SPT.ti,ab. 13. (presence adj2 intervention*).ti,ab. 14. (simulat* adj2 family).ti,ab. 15. ((video* or audio*) adj2 presence).ti,ab. 16. or/11-15 17. 10 and 16	August 2015: 28 February 2015: 2
3. Embase 1980-2011 week 30 (Ovid SP) [Latest search: 9 February 2016]	1. exp Dementia/ 2. Delirium, Dementia, Amnestic, Cognitive Disorders/ 3. dement*.mp. 4. alzheimer*.mp. 5. (lewy* adj2 bod*).mp. 6. (chronic adj2 cerebrovascular).mp. 7. ("organic brain disease" or "organic brain syndrome").mp. 8. (cerebr* adj2 deteriorat*).mp. 9. (cerebral* adj2 insufficient*).mp.	August 2015: 63 February 2015: 4

(Continued)

	10. or/1-9 11. "Simulated presence therapy".ti,ab. 12. SPT.ti,ab. 13. (presence adj2 intervention*).ti,ab. 14. (simulat* adj2 family).ti,ab. 15. ((video* or audio*) adj2 presence).ti,ab. 16. or/11-15 17. 10 and 16	
4. PSYCINFO 1806-July week 4 2011 (Ovid SP) [Latest search: 9 February 2016]	1. exp dementia/ 2. dement*.mp. 3. alzheimer*.mp. 4. (lewy* adj2 bod*).mp. 5. (chronic adj2 cerebrovascular).mp. 6. ("organic brain disease" or "organic brain syndrome").mp. 7. (cerebr* adj2 deteriorat*).mp. 8. (cerebral* adj2 insufficient*).mp. 9. or/1-8 10. "Simulated presence therapy".ti,ab. 11. SPT.ti,ab. 12. (presence adj2 intervention*).ti,ab. 13. (simulat* adj2 family).ti,ab. 14. ((video* or audio*) adj2 presence).ti,ab. 15. or/10-14 16. 9 and 15	August 2015: 66 February 2015: 4
5. CINAHL (EBSCOhost) [Latest search: 9 February 2016]	S1 (MH "Dementia+") S2 (MH "Delirium") or (MH "Delirium, Dementia, Amnestic, Cognitive Disorders") S3 (MH "Wernicke's Encephalopathy") S4 TX dement* S5 TX alzheimer* S6 TX lewy* N2 bod* S7 TX deliri* S8 TX chronic N2 cerebrovascular S9 TX "organic brain disease" or "organic brain syndrome" S10 TX "normal pressure hydrocephalus" and "shunt*" S11 TX "benign senescent forgetfulness" S12 TX cerebr* N2 deteriorat* S13 TX cerebral* N2 insufficient* S14 TX pick* N2 disease S15 TX creutzfeldt or jcd or cjd S16 TX huntington*	August 2015: 5 February 2015: 0

(Continued)

	S17 TX binswanger* S18 TX korsako* S19 S1 or S2 or S3 or S4 or S5 or S6 or S7 or S8 or S9 or S10 or S11 or S12 or S13 or S14 or S15 or S16 or S17 or S18 S20 TX "Simulated presence therapy" S21 TX SPT S22 TX "presence intervent*" S23 TX "audio presence" S24 S20 or S21 or S22 or S23 S25 S24 and S19	
6. ISI Web of Knowledge - all databases [includes: Web of Science (1945-present) ; BIOSIS Previews (1926-present); MED- LINE (1950-present); Journal Citation Re- ports] [Latest search: 9 February 2016]	Topic=(Dement* OR Alzheimer* OR "Lewy body" OR arteriosclerosis OR "Huntington* disease" OR "Kluver Bucy" OR "Pick disease" OR "Wernicke en- cephalopathy" OR "Korsakoff psychosis") AND Topic=("simulated presence therapy" OR SPT OR "presence intervention" OR "audio presence")	August 2015: 63 February 2015: 1
7. LILACS (BIREME) [Latest search: 9 February 2016]	"simulated presence therapy" OR SPT	August 2015: 0 February 2015: 0
8. CENTRAL (<i>The Cochrane Library</i>) (Is- sue 4 of 4, Oct 2010) [Latest search: 9 February 2016]	#1 simulated presence therapy #2 SPT #3 simulat* near intervent* #4 #1 or #2 or #3 #5 dementia #6 MeSH descriptor: [Dementia] explode all trees #7 #5 or #6 #8 #4 and #7 in Trials	August 2015: 6 February 2015: 1
9. Clinicaltrials.gov (www.clinicaltrials.gov) [Latest search: 9 February 2016]	Interventional studies: Intervention: simu- lated presence therapy OR SPT AND con- dition: dementia OR alzheimer's Status of trial: ALL	August 2015: 2 February 2015: 0
10. ICTRP Search Portal (http:// /apps.who.int/trialsearch) [includes: Aus- tralian New Zealand Clinical Trials Reg- istry; ClinicalTrilas.gov; ISRCTN; Chinese Clinical Trial Registry; Clinical Trials Reg- istry - India; Clinical Research Informa- tion Service - Republic of Korea; German Clinical Trials Register; Iranian Registry of Clinical Trials; Japan Primary Registries Network; Pan African Clinical Trial Reg- istry; Sri Lanka Clinical Trials Registry; The	Interventional studies: Intervention: simu- lated presence therapy OR SPT AND con- dition: dementia OR alzheimer's Status of trial: ALL	August 2015: 0 February 2015: 0

(Continued)

Netherlands National Trial Register] [Latest search: 9 February 2016]	
TOTAL before de-duplication	August 2015: 236 February 2015: 12
TOTAL after de-dupe and first-assess	August 2015: 10 February 2015: 0

CONTRIBUTIONS OF AUTHORS

AC, ACJ, ILM, FMT and MV were content experts (dementia) for the review.

IA was the methodological expert for this review and initially project managed the review, provided support and training and provided support with data analysis and interpretation.

IA, JMR, GD, ILM, and MV undertook the selection of trials, data extraction, and assessment of risk of bias and analysis of results.

IA, AC, JMR, GD, FMT, MV, ILM, ACJ and GD drafted the review and approved the final version of the review.

DECLARATIONS OF INTEREST

Iosief Abraha: none known.

Joseph M Rimland: none known.

Isabel Lozano-Montoya: none known.

Giuseppina Dell'Aquila: none known.

Manuel Vélez-Díaz-Pallarés: none known.

Fabiana M Trotta: none known.

Alfonso J Cruz-Jentoft: none known.

Antonio Cherubini: none known.

SOURCES OF SUPPORT

Internal sources

- No sources of support supplied

External sources

- NIHR, UK.

This protocol was supported by the National Institute for Health Research, via Cochrane Infrastructure funding to the Cochrane Dementia and Cognitive Improvement Group. The views and opinions expressed therein are those of the authors and do not necessarily reflect those of the Systematic Reviews Programme, NIHR, NHS or the Department of Health.

- 7th Framework program, European Union, Other.

This protocol was partly funded by the European Union Seventh Framework program (FP7/2007-2013) under grant agreement n° 305930 (SENATOR). The funder had no role in the study design, data collection and analysis, the decision to publish, or the preparation of the manuscript.

DIFFERENCES BETWEEN PROTOCOL AND REVIEW

None.

INDEX TERMS

Medical Subject Headings (MeSH)

*Family; *Tape Recording; *Videotape Recording; Dementia [psychology; *therapy]; Depression [*therapy]; Psychomotor Agitation [*therapy]; Randomized Controlled Trials as Topic

MeSH check words

Aged, 80 and over; Female; Humans; Male