Published in final edited form as:

J Adolesc Health. 2019 May ; 64(5): 594–601. doi:10.1016/j.jadohealth.2018.10.293.

Measuring discomfort in health research relative to everyday events and routine care: An application to sexual and gender minority youth

Kathryn Macapagal, PhD^{a,b}, Emily Bettin^{a,b}, Margaret Matson, MPH^{a,b}, Ashley Kraus, PhD^{a,b}, Celia B. Fisher, PhD^c, and Brian Mustanski, PhD^{a,b}

^aDepartment of Medical Social Sciences, Northwestern University Feinberg School of Medicine, 625 N. Michigan Avenue, Suite 1400, Chicago, IL 60611, USA

^bInstitute for Sexual and Gender Minority Health and Wellbeing, Northwestern University, 625 N. Michigan Avenue, Suite 1400, Chicago, IL 60611, USA

^cCenter for Ethics Education & Department of Psychology, Fordham University, 441 E. Dealy Road, Dealy 226, Bronx, NY 10458, USA

Abstract

Purpose: Understanding how sexual and gender minority (SGM) youth's comfort with research procedures compares to their comfort with everyday experiences and routine healthcare can help calibrate decisions about whether a study meets minimal risk criteria. We sought to quantify SGM adolescents' comfort with sexual health research relative to everyday events and activities often cited as benchmarks of minimal risk.

Methods: 616 SGM adolescents in the USA (M age = 15.7, 41.7% racial/ethnic minority) completed online survey questions assessing sexual behavior, SGM identity, and a 53-item Measure of Adolescent Comfort with Clinical, Research, and Everyday Events (MACCREE) that assessed comfort on a 7-point scale (1=extremely uncomfortable; 7=extremely comfortable).

Results: The Everyday Events for Adolescents domain had the lowest mean comfort score ($M=3.49$, $SD=0.58$) and was significantly lower than the Routine Medical and Psychological Tests domain ($M=4.43$, $SD=0.92$) and the HIV/Sexual Health Research Procedures domain ($M=4.19$, $SD=0.94$). Eleven out of 17 items on the HIV/Sexual Health Research Procedures domain were ranked as more comfortable than a neutral rating of "neither comfortable nor uncomfortable."

Corresponding author: Kathryn Macapagal, PhD, Research Assistant Professor, Department of Medical Social Sciences, Feinberg School of Medicine, Institute for Sexual and Gender Minority Health and Wellbeing, Northwestern University, 625 N. Michigan Avenue, Suite 14-057, Chicago, Illinois 60611, 312-503-3605 (p); 312-503-6700 (f), kathryn.macapagal@northwestern.edu. Co-author contact information: emily.bettin@northwestern.edu, maggie.matson@northwestern.edu, ashley.kraus@northwestern.edu, fisher@fordham.edu, brian@northwestern.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Conflicts of Interest: The authors have no conflicts of interest relevant to this article to disclose. The study sponsor was not involved in study design, collection, analysis, or interpretation of data, the writing of this report, or the decision to submit this manuscript for publication. Kathryn Macapagal, Emily Bettin, Margaret Matson, and Ashley Kraus wrote the first draft of the manuscript and were not given honoraria to produce this manuscript.

Higher levels of parental acceptance predicted greater levels of comfort across all four domains of the MACCREE. Participants who were out to their parents expressed greater comfort with both SGM Identity and Sexual Health-related procedures and events as well as HIV/Sexual Health Research Procedures.

Conclusions: Overall participants expressed equal or more comfort with research procedures than with everyday life experiences. These findings indicate that common sexual health research procedures may meet minimal risk criteria among SGM adolescent populations.

Keywords

adolescent sexual behavior; adolescent sexual health; research ethics; HIV/AIDS

Sexual and gender minority (SGM) adolescents are at disproportionate risk of adverse sexual health and substance use outcomes, including higher rates of sexually transmitted infections (STIs),^{1,2} HIV,³ unplanned pregnancy,^{4,5} and alcohol and substance use^{6–8} relative to their heterosexual and cisgender (i.e., not transgender or gender variant) peers. However, SGM adolescents under 18 are underrepresented in research aimed at understanding and improving these health outcomes, which perpetuates health disparities in this population.⁹ One obstacle to their inclusion is some investigators' and institutional review boards' (IRBs) belief that certain research procedures may pose greater than minimal risk for adolescents^{9–12}—in other words, that they exceed the magnitude or probability of risks encountered during everyday life or routine medical or psychological exams.¹³ Case examples from researchers in the U.S. and Canada have described challenges navigating IRB review for social-behavioral^{10–12} and biomedical research with SGM adolescents¹⁴ on 'sensitive' topics, and IRB application of risk/benefit standards in health research with minors can be variable.^{15,16} Subjective judgments about the risks of adolescent health research may lead IRBs to deny parental permission waivers, which can lead to critical sampling biases¹⁷ or the exclusion of minors from a research protocol.^{12,14} Thus, the ability to quantify common risks, such as the potential for discomfort in social-behavioral and biomedical research, can promote consistency in IRB decisions about minimal risk. To facilitate evidence-based decision making regarding SGM adolescents' inclusion in health research, we sought to measure youth's comfort with sexual health and risk behavior research procedures relative to everyday events (e.g., taking a test) and clinical activities often cited as benchmarks of minimal risk (e.g., blood draws).

The limited literature on SGM adolescents' perspectives on research participation shows that they generally feel comfortable participating in health related research.^{11,18–21} In a study of 181 SGM youth, 90% felt comfortable with answering survey questions about sexual behavior, mental health, and substance and alcohol abuse, and only 2–3% felt very uncomfortable across these domains.¹¹ Another study found SGM adolescents felt low levels of distress after completing survey questions about their sexual behavior and that this distress was not significantly different from that reported by adult participants.¹⁹ Likewise, a survey and focus group study of 74 SGM adolescents age 14–17 found that the majority felt comfortable answering survey questions about sexual behavior (67%) and drug and alcohol use (61%).²⁰ These studies lend evidence to the argument that certain sexual health and risk behavior research procedures pose relatively little risk.

Limited research has quantified individuals' comfort during everyday events and routine care and compared them to social-behavioral or biomedical research procedures, although such work can help calibrate decisions about whether a study meets minimal risk criteria. For example, Petrie et al.²² asked university students and community members to rate their anticipated distress associated with common health research procedures (e.g., sexual behavior surveys, health interviews) and everyday events; ethics committee members were asked to rate their perceptions of how distressing the public would find these same activities. Community members and students rated their distress regarding research procedures lower than did ethics committee members, and similar to or less than their distress regarding everyday events (e.g., waiter forgetting order).²² This study indicates that common health research procedures are perceived to be relatively low in risk, but also that ethics committee members systematically overestimated potential participants' perceived distress. However, this work was conducted with adults, and comfort with health-related research procedures may differ among adolescents. To our knowledge, the only study that has directly compared routine healthcare with research participation in SGM adolescents involved young sexual minority men who were interviewed about their sexual behavior; afterward, participants reported feeling equally or more comfortable in the study than in a routine medical visit.¹⁸ To build on this work, larger-scale quantitative studies are needed to better situate the discomforts of a variety of adolescent health research procedures relative to exemplars of minimal risk events.

Among SGM adolescents, anticipated discomfort associated with sexual health and risk behavior research participation could be related to certain sociodemographic and behavioral factors. For example, transgender youth or SGM youth of color may experience greater discomfort with research procedures due to mistrust stemming from historical mistreatment of their communities in research contexts.^{23,24} In addition, youth who are not out to their parents about their SGM identity, whose parents may be unaccepting of their SGM identity, or who are sexually active may have heightened concerns about research participation due to fears of parental discovery of and punishment for their SGM identity or same-gender sexual behavior.^{25,26}

Understanding adolescents' perspectives on research that can affect their health and wellbeing is a cornerstone of ethical practice and can prevent IRBs and investigators from identifying research as harmful or distressing when participants perceive it otherwise.²⁷ To guide evidence-informed decision making by researchers and IRBs, following Petrie et al.²² we measured SGM adolescents' discomfort with sexual health and risk behavior research procedures and compared these items to everyday events and routine medical and psychological tests. As identifying predictors of increased discomfort in research can draw IRBs' and researchers' attention to participant groups who may benefit from additional protective mechanisms to mitigate risk of harm, we also explored how comfort differed by sociodemographic and behavioral characteristics.^{20,21}

Method

Participants and Recruitment

Participants were recruited to complete an online survey on ethical issues in adolescent sexual health research as part of a larger study.^{20,21,28} Eligible participants were 14–17 years old; identified as a sexual minority (e.g., lesbian, gay, bisexual, queer, questioning) and/or transgender or gender non-conforming, or were romantically interested in or had sex with partners of the same gender; lived in the United States; could read English at an 8th grade level; and were HIV-negative or did not know their status. All procedures were IRB approved. A waiver of parental permission was granted on the grounds that it was not a reasonable requirement to protect the participants, the research could not practicably be carried out without a waiver, and study procedures were determined to be no more than minimal risk.¹³ A Certificate of Confidentiality from the National Institutes of Health was issued to protect the identities of participants in the event of subpoenas requesting identifiable study data.

Participants were recruited through paid Facebook, Instagram, and Twitter advertisements in spring 2017. Advertisements targeted adolescents who indicated they were romantically interested in people of the same gender on their profile and/or listed interests relevant to SGM youth (e.g., pop culture figures popular with SGM youth). Clicking on the advertisement directed individuals to an online eligibility survey. Eligible participants were presented with an online consent form and were automatically directed to the survey. No financial incentives were offered for participation.

Measures

Demographics, sexual behavior, SGM identity.

Participants completed items assessing age, race and ethnicity, sex assigned at birth, current gender identity and sexual orientation, and disclosure of sexual orientation to parents (out to one, all, none). Participants were then asked about sexual orientation disclosure to their mother or guardian most like a mother, and their father or guardian most like a father (1 = definitely knows and we have talked about it, 4 = does not know or suspect), and the extent to which these individuals were accepting or would be accepting of their sexual orientation if disclosed (1 = very rejecting, 5 = very accepting).²⁶ Disclosure to parents was dichotomized to being not out versus being out to at least one parent. Responses to a closed-ended item assessing participants' sexual activity were dichotomized to reflect whether participants ever had sex (yes, no).

Comfort with research, everyday events, and routine medical/psychological tests.

Items assessing comfort with research participation were based on a 15-item questionnaire developed by the authors.²⁰ Items were added that reflected activities deemed to pose no more than minimal risk by most IRB standards (e.g., finger stick); common sexual health research procedures (e.g., sexual behavior interviews) and HIV prevention studies that are underway or are anticipated in the future (e.g., investigational trials for injectable and implantable preexposure prophylaxis [PrEP], a medication to prevent HIV).²⁹ Several

everyday events experienced by adolescents were taken from previous research (e.g., caught in rain,²² taking a test³⁰) or adapted to reflect adolescent experiences (e.g., standing in long line at a bank²² changed to standing in a long line at a store). Additional items were developed in consultation with SGM youth to ensure the everyday events reflected a range of adolescent experiences that varied widely in their comfort levels (e.g., coming out, using social media) and piloted; see supplementary material for the initial and final item sets. Items were rated on a scale ranging from extremely uncomfortable (1) to extremely comfortable (7).

The Measure of Adolescent Comfort with Clinical, Research, and Everyday Events (MACCREE; Table 2) reported in the present study included 53 items reflecting four domains. Three domains assessed comfort with *Everyday Events for Adolescents* (e.g., “Taking a test at school”; 21 items); *Routine Medical and Psychological Tests* (e.g., “Having your vision checked at the doctor’s office”; 11 items); and events related to *SGM Identity and Sexual Health* (e.g., “Coming out to a parent or guardian”; 4 items). Another domain assessed comfort with *HIV/Sexual Health Research Procedures* (e.g., “Having your finger pricked to test your blood for HIV during a research study”; 17 items) including some procedures that would require parental consent. Mean ratings for items in each domain were computed.²²

Procedure

After providing informed assent, participants completed a 20-minute survey that included the MACCREE and items on condom attitudes and perspectives on research incentives; results from the latter two topics are not reported here. The MACCREE was positioned toward the middle of the survey following questions about condom attitudes, and items were randomized (i.e., not grouped by domain) and presented in a matrix. Instructions for the MACCREE were as follows: “In this section, we’d like to understand how uncomfortable or comfortable you would feel with different types of life events, health related tests, and procedures used in research studies on sexual health. This is so we can compare how you might feel about different research procedures relative to other events you experience in your everyday life. The content or order of items may at times seem strange, but please answer as best you can. Please indicate how uncomfortable or comfortable you would be with each event on a scale from extremely uncomfortable to extremely comfortable.”

Data Analysis

Statistical analyses were run in SPSS 24. Frequencies were run for all demographic variables, and descriptive statistics were computed for individual items as well as the four domains. Item means were rank-ordered in a bar chart to better visualize how individual items compared with each other. To test for differences between the mean scores of the MACCREE domains, we used repeated measures ANOVA and post-hoc tests with Bonferroni corrections were run to discover which specific means differed.

Several tests assessed demographic differences in MACCREE domain means. Independent samples t-tests compared the ratings of domains between dichotomous variables (assigned male at birth [AMAB] vs. assigned female at birth [AFAB]; sexually active vs. not). Simple

linear regression compared continuous variables (age; parental acceptance of SGM identity) and dummy-coded variables (race/ethnicity: Black, Latino, and Other, with White as reference group; gender identity: transgender, genderqueer, and gender nonconforming, with cisgender as reference group). Finally, to test the unique influence of demographic characteristics, variables that were significant at $p < 0.05$ were included in multiple linear regression models for each domain.

Results

The sample (Table 1) included 616 participants ($M=15.69$, $SD=1.01$). Participants' racial/ethnic backgrounds were diverse (41.7% youth of color). Over half (54.7%) were AFAB. Cisgender men (43.2%) and women (28.6%) made up the majority of the sample. Of the transgender (12.2%), genderqueer (8.0%), and gender nonconforming (8.1%) youth in the sample, 89.5% were AFAB. Gay (37.0%), bisexual (23.2%), and pansexual youth (20.3%) made up the largest sexual orientation groups. More than half (62.8%) were out to at least one parent, and 60.1% reported their mothers were somewhat to very accepting of their sexual orientation, whereas 40.9% reported their fathers were somewhat to very accepting. Nearly two-thirds (65.6%) had previously had sex.

MACCREE Domains and Items

Routine Medical and Psychological Tests had the highest mean score ($M=4.43$, $SD=0.92$), indicating the greatest levels of comfort, followed by HIV/Sexual Health Research Procedures ($M=4.19$, $SD=0.94$), and SGM Identity and Sexual Health ($M=3.57$, $SD=1.08$). The domain with the lowest mean score was Everyday Events for Adolescents ($M=3.49$, $SD=0.58$). A repeated measures ANOVA with a Greenhouse-Geisser correction showed a significant difference between the means ($F(2.45, 1505.43) = 342.74$, $p < 0.001$). Post-hoc tests showed that all domain means differed significantly from each other ($p < 0.001$), with the exception of SGM Identity and Sexual Health and Everyday Events for Adolescents ($p = 0.295$).

Figure 1 illustrates the items rank-ordered from most to least comfortable and enables direct comparisons between individual research procedures, routine care, and everyday events. Out of all items, "having your cell phone stolen" ($M=1.25$, $SD=0.59$) ranked as the most uncomfortable event, and "sending a text message to your friend" ($M=6.30$, $SD=1.09$) ranked as the most comfortable (Figure 1). The item "having your blood drawn at the doctor's office" was rated closest to the median ($M=3.99$, $SD=1.37$) reflecting a neutral rating of "neither comfortable nor uncomfortable."

Items in the HIV/Sexual Health Research Procedures domain tended to cluster higher on the comfort scale, with 11 out of 17 items ranking as more comfortable than "having your blood drawn at the doctor's office" (Figure 1). These items included study procedures involving HIV testing, urine drug screening, talking to a researcher about sexual behavior, and talking about sexual behavior in a focus group with other teenagers. One of the biomedical HIV prevention research procedures, "getting an injection (a shot) of medication every three months that can help you prevent HIV for a research study," was rated similarly to these items. Only two items in the HIV/Sexual Health Research Procedures domain had a mean

score less than 3 (somewhat uncomfortable): “having to ask your parents for permission to participate in a research study about your sexual behavior” and “having the inside of your butt swabbed to test for STIs during a research study.”

Group Differences

Results of multiple regression tests are in Table 3. Significant predictors were found across all four MACCREE domains, even while controlling for other demographic covariates. Regarding outness to parents and parental attitudes, higher levels of parental acceptance predicted significantly greater levels of comfort across all four domains. Participants who were out to their parents or guardians expressed significantly greater comfort with both SGM Identity and Sexual Health-related procedures and events and HIV/Sexual Health Research Procedures.

Regarding demographic characteristics and sexual history, compared to White participants, Black participants reported significantly greater comfort levels with both Routine Medical and Psychological Tests and HIV/Sexual Health Research Procedures, while participants who identified as other racial or ethnic minorities reported significantly less comfort with Everyday Events for Adolescents. Cisgender participants were significantly more comfortable than their non-cisgender peers with Everyday Events for Adolescents. AFAB youth were less comfortable than AMAB youth across three of the four domains: Routine Medical and Psychological Tests, Everyday Events for Adolescents, and HIV/Sexual Health Research Procedures. The older participants were, the greater comfort they expressed with both Routine Medical and Psychological Tests as well as HIV/Sexual Health Research Procedures. Finally, sexually active youth reported greater comfort with HIV/Sexual Health Research Procedures.

Discussion

Empirical studies focused on SGM adolescents’ comfort with health-related research procedures have the potential to shape decisions regarding their inclusion in research, as well as inform practices that might mitigate risks. In this study, we quantitatively measured SGM youth’s comfort with sexual health, HIV prevention, and risk behavior research procedures relative to everyday life events and routine medical care using the MACCREE, a questionnaire based on the framework described in Petrie et al.²²

Overall, findings are consistent with research on community samples of adults and college students²² in that SGM youth rated their discomfort with most research procedures as relatively low. Moreover, their comfort with research procedures was significantly greater than most everyday events, and research procedures were typically rated similarly to or more comfortable than analogous procedures at their doctor’s office (e.g., talking to a researcher vs. doctor about substance/alcohol use). The few research procedures that were rated as more uncomfortable (i.e., < 4 on a scale of 1–7) involved asking for parental permission to participate in research, and more invasive procedures, such as a rectal swab to test for STIs (done in some of our adolescent research studies considered to have minimal risk) and insertion of a small implant containing HIV prevention medication (a drug delivery system that is in development, similar to a contraceptive implant).²⁹ Youth rated the prospect of

participating in a study where they could be randomized to an HIV prevention drug or placebo as somewhat more uncomfortable than most everyday events and research procedures ($M = 3.32$), similar in discomfort to being grounded.

Several sociodemographic and behavioral predictors of comfort on the MACCREE emerged. Greater parental acceptance of SGM identity and outness to parents predicted increased comfort with HIV/sexual health research procedures. These findings substantiate prior qualitative and mixed-methods work showing that outness and parental acceptance were linked with differential willingness to engage in sexual health research among SGM adolescents.^{20,21,28} AFAB youth reported lower levels of comfort across HIV/sexual health research procedures, routine care, and everyday events as compared to AMAB youth. AFAB youth may have found the HIV prevention research procedures less personally relevant than did AMAB youth; some research suggests sexual minorities AFAB may have worse healthcare experiences than their AMAB counterparts,^{31,32} and several everyday events described experiences related to school or academic achievement, settings in which AFAB adolescents may have lower self-confidence than AMAB adolescents.³³ Unexpectedly, Black youth reported higher levels of comfort with HIV prevention research participation and routine care than did non-Hispanic White youth. Nevertheless, as our sample of Black youth was relatively small, this finding should be interpreted with caution. Finally, being older and having had sex was associated with greater comfort with HIV/sexual health research procedures.

Limitations

First, this study assessed youth's discomfort with research and not all possible harms related to research participation. Moreover, we do not know whether participants considered magnitude of discomfort, duration, or both in their ratings. As IRBs need to consider the probability and magnitude of harm to determine whether a study is minimal risk, other frameworks such as those described in Rid et al.³⁰ may also be useful in their decision making. Second, youth were not given complete details of the research studies, which may have affected their comfort ratings. For example, our experiences in our ongoing research studies indicate that youth express initial discomfort about the concept of a rectal swab to test for STIs, but upon learning more details or doing the swab, their discomfort is largely eliminated. Relatedly, generalizability may be limited as we asked participants to anticipate their discomfort in hypothetical research situations, rather than measuring discomfort before and after research participation. Arguably, however, when individuals are considering enrollment in a research study, they are likely taking into account anticipated benefits and risks – including an educated guess about the level of discomfort they may experience – prior to making a decision. Moreover, many common research procedures are also part of routine care (e.g., health questionnaires) and as such, participants can draw from their experiences there. Generalizability may also be limited by the fact that participants who volunteered for this study may have been more comfortable participating in research. Finally, we did not assess geographic location, which may have impacted participant ratings; for example, youth in rural or more socially conservative areas may be less comfortable with sexual health research participation.

Implications and Future Directions

Here we presented one framework for how adolescent health researchers can assess participants' comfort with novel or existing research procedures in their area of study. As adolescent HIV prevention researchers, the MACCREE enabled us to explore youth's comfort with emerging biomedical prevention products (e.g., injectable and implantable PrEP) relative to standard HIV behavioral surveillance and prevention research procedures, and it is useful to know how adolescents perceive novel procedures well before they are tested on youth populations. Although it may not be feasible to ask every question due to time constraints and topic relevance (e.g., the SGM items could be dropped for researchers not studying SGM youth), researchers could develop items for procedures specific to their area and compare youth's scores to our items of common clinical procedures and everyday events. In addition, following Petrie et al.,²² future studies could compare SGM youth's levels of comfort to those of cisgender and heterosexual youth, IRB members, and parents, which would shed light on these groups' estimations of research-related discomfort relative to SGM youth.

Conclusion

Our findings suggest that SGM youth are generally comfortable with a variety of sexual health and risk behavior research procedures. Certain groups of youth (e.g., younger, sexually inexperienced, not out, lack parental acceptance) and youth in certain types of studies (e.g., those that require parental permission, randomized trials, more invasive procedures) may benefit from additional measures to increase comfort with participation, such as waiving parental permission, increased privacy protections, developmentally-appropriate yet detailed explanations of study procedures and risks/benefits, and access to SGM-specific sexual health information.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments:

This research was supported by the National Institute on Minority Health and Health Disparities of the National Institutes of Health under award R01MD009561 (PIs: Mustanski & Fisher). This content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Abbreviations:

(SGM)	sexual and gender minority
(STI)	sexually transmitted infection
(HIV)	human immunodeficiency virus
(IRB)	institutional review board
(MSM)	men who have sex with men

(PrEP)	pre-exposure prophylaxis
(ANOVA)	analysis of variance
(AMAB)	assigned male at birth
(AFAB)	assigned female at birth

References

- Centers for Disease Control and Prevention. 2015 sexually transmitted diseases treatment guidelines: Special populations 2015; <http://www.cdc.gov/std/tg2015/specialpops.htm>. Accessed October 16, 2015.
- Reisner SL, White JM, Mayer KH, Mimiaga MJ. Sexual risk behaviors and psychosocial health concerns of female-to-male transgender men screening for STDs at an urban community health center. *AIDS Care*. 2014;26(7):857–864. [PubMed: 24206043]
- Centers for Disease Control and Prevention. HIV among youth 2015; <http://www.cdc.gov/hiv/group/age/youth/index.html>.
- Lindley LL, Walsemann KM. Sexual orientation and risk of pregnancy among New York City high-school students. *American Journal of Public Health* 7 2015;105(7):13791386.
- Tornello SL, Riskind RG, Patterson CJ. Sexual orientation and sexual and reproductive health among adolescent young women in the United States. *J. Adolesc. Health*. 2 2014;54(2):160–168. [PubMed: 24157195]
- Kann L, Olsen EO, McManus T, et al. Sexual identity, sex of sexual contacts, and health-related behaviors among students in grades 9–12 - United States and selected sites, 2015. *Mmwr Surveill Summ* 8 12 2016;65(9):1–202.
- Zaza S, Kann L, Barrios LC. Lesbian, gay, and bisexual adolescents: Population estimate and prevalence of health behaviors. *JAMA* 12 13 2016;316(22):2355–2356. [PubMed: 27532437]
- Day JK, Fish JN, Perez-Brumer A, Hatzenbuehler ML, Russell ST. Transgender youth substance use disparities: Results from a population-based sample. *J Adolesc Health* 12 2017;61(6):729–735. [PubMed: 28942238]
- Fisher CB, Mustanski B. Reducing Health Disparities and Enhancing the Responsible Conduct of Research Involving LGBT Youth. *Hastings Cent Rep* 9 2014;44 Suppl 4(s4):S28–31.
- Miller RL, Forte D, Wilson BD, Greene GJ. Protecting sexual minority youth from research risks: Conflicting perspectives. *Am J Community Psychol*. 6 2006;37(34):341–348. [PubMed: 16775774]
- Mustanski B. Ethical and regulatory issues with conducting sexuality research with LGBT adolescents: A call to action for a scientifically informed approach. *Arch Sex Behav* 8 2011;40(4):673–686. [PubMed: 21528402]
- Tufford L, Newman PA, Brennan DJ, Craig SL, Woodford MR. Conducting research with lesbian, gay, and bisexual populations: Navigating research ethics board reviews. *J Gay Lesbian Soc Serv* 2012;24(3):221–240.
- Department of Health and Human Services. Code of Federal Regulations, Title 45 Public Welfare, Part 46 Protection of Human Subjects In: Department of Health and Human Services, ed2009.
- Gilbert AL, Knopf AS, Fortenberry JD, Hosek SG, Kapogiannis BG, Zimet GD. Adolescent self-consent for biomedical human immunodeficiency virus prevention research. *J Adolesc Health*. 7 2015;57(1):113–119. [PubMed: 26095412]
- Fisher CB, Kornetsky SZ, Prentice ED. Determining risk in pediatric research with no prospect of direct benefit: Time for a national consensus on the interpretation of federal regulations. *Am J Bioethics* 3 2007;7(3):5–10.
- Shah S, Whittle A, Wilfond B, Gensler G, Wendler D. How do institutional review boards apply the federal risk and benefit standards for pediatric research? *JAMA*. 2004/01/28/ 2004;291(4):476–482. [PubMed: 14747505]

17. Liu C, Cox RB Jr., Washburn IJ, Croff JM, Crethar HC. The effects of requiring parental consent for research on adolescents' risk behaviors: A meta-analysis. *J Adolesc Health* 7 2017;61(1):45–52. [PubMed: 28363714]
18. Arrington-Sanders R, Morgan A, Oidman J, et al. Sexual health research with young Black men who have sex with men: Experiences of benefits and harms. *Arch Sex Behav*. 5 2017;46(4):937–946. [PubMed: 27043836]
19. Kuyper L, de Wit J, Adam P, Woertman L. Doing more good than harm? The effects of participation in sex research on young people in the Netherlands. *Archives of Sexual Behavior* 4 2012;41(2):497–506. [PubMed: 21681692]
20. Macapagal K, Coventry R, Arbeit MR, Fisher CB, Mustanski B. “I won’t out myself just to do a survey”: Sexual and gender minority adolescents’ perspectives on the risks and benefits of sex research. *Arch Sex Behav* 7 2017;46(5):1393–1409. [PubMed: 27469352]
21. Mustanski B, Coventry R, Macapagal K, Arbeit MR, Fisher CB. Sexual and gender minority adolescents’ views on HIV research participation and parental permission: A mixed-methods study. *Perspect Sex Reprod Health*. 6 2017;49(2):111–121. [PubMed: 28445608]
22. Petrie KJ, Faasse K, Notman TA, O’Carroll R. How distressing is it to participate in medical research? A calibration study using an everyday events questionnaire. *JRSM Short Rep* October 1, 2013 2013;4(10):1–7.
23. Zea MC, Reisen CA, Diaz RM. Methodological issues in research on sexual behavior with Latino gay and bisexual men. *Am J Community Psychol* 6 2003;31(3–4):281–291. [PubMed: 12866685]
24. Owen-Smith AA, Woodyatt C, Sineath RC, et al. Perceptions of barriers to and facilitators of participation in health research among transgender people. *Transgend Health* 2016;1(1):187–196. [PubMed: 28861532]
25. D’Augelli AR, Grossman AH, Starks MT. Families of gay, lesbian, and bisexual youth: What do parents and siblings know and how do they react? *J GLBT Fam Stud*. 2008 2008;4(1):95–115.
26. D’Augelli AR, Hershberger SL, Pilkington NW. Lesbian, gay, and bisexual youth and their families: Disclosure of sexual orientation and its consequences. *American Journal of Orthopsychiatry* 7 1998;68(3):361–371. [PubMed: 9686289]
27. United Nations General Assembly. *Convention on the Rights of the Child*. New York, NY: United Nations;1989 A/RES/44/25.
28. Fisher CB, Arbeit MR, Dumont MS, Macapagal K, Mustanski B. Self-consent for HIV prevention research involving sexual and gender minority youth: Reducing barriers through evidence-based ethics. *J Empir Res Hum Res Ethics*. 2 2016;11(1):3–14. [PubMed: 26956988]
29. US Department of Health and Human Services. Long-acting PrEP 2017; <https://www.hiv.gov/hiv-basics/hiv-prevention/potential-future-options/long-acting-prep>. Accessed April 17, 2018.
30. Rid A, Abdoler E, Roberson-Nay R, Pine DS, Wendler D. Evaluating the risks of clinical research: Direct comparative analysis. *J Child Adolesc Psychopharmacol* 9 2014;24(7):390–398. [PubMed: 25210944]
31. Williams KA, Chapman MV. Unmet health and mental health need among adolescents: The roles of sexual minority status and child-parent connectedness. *Am. J. Orthopsychiatry*. 10 2012;82(4):473–481. [PubMed: 23039345]
32. Macapagal K, Bhatia R, Greene GJ. Differences in healthcare access, use, and experiences within a community sample of racially diverse lesbian, gay, bisexual, transgender, and questioning emerging adults. *LGBT Health*. 12 2016;3(6):434–442. [PubMed: 27726496]
33. Perry DG, Pauletti RE. Gender and Adolescent Development. *Journal of Research on Adolescence*. 2011;21(1):61–74.

Implications and contributions

This study quantifies sexual and gender minority (SGM) adolescents' comfort with sexual health research procedures relative to everyday events and routine care. These findings can inform decisions made by researchers and institutional review boards regarding inclusion of SGM adolescents in sexual health research.

Figure 1.

Rank-ordered MACCREE items reflecting adolescents' level of comfort with everyday events, routine medical and psychological tests, SGM identity and sexual health topics, and HIV/sexual health research procedures (denoted in black with *). Dotted line indicates "line of neutrality" (score of 4 = "neither comfortable nor uncomfortable").

Table 1.

Demographic characteristics (N = 616)

	N	%
Age ($M=15.69$; $SD=1.01$)		
14	86	14.0
15	178	28.9
16	191	31.0
17	161	26.1
Race/Ethnicity		
White	359	58.3
Black/African American	28	4.5
Hispanic or Latino	164	26.6
Asian	16	2.6
Native Hawaiian/Pacific Islander	6	1.0
American Indian/Alaska Native	3	0.5
More than one race	31	5.0
Other	4	0.6
Prefer not to answer	5	0.8
Sex Assigned at Birth		
Male	279	45.3
Female	337	54.7
Gender Identity		
Man	266	43.2
Woman	176	28.6
Transgender Man	70	11.4
Transgender Woman	5	0.8
Genderqueer	49	8.0
Gender nonconforming	50	8.1
Cisgender		
Cisgender	435	70.6
Not Cisgender	181	29.4
Sexual Orientation		
Gay	228	37.0
Lesbian	44	7.1
Bisexual	143	23.2
Pansexual	125	20.3
Asexual	15	2.4
Heterosexual	4	0.6
Queer	34	5.5
Questioning/Unsure	23	3.7
Outness to Parents/Guardians		
Not out to any	222	36.0

	N	%
Out to at least one (but not all)	150	24.4
Out to all	237	38.5
Prefer not to answer	7	1.1
Mom's Acceptance of Sexual Orientation ($M=3.60$, $SD=1.42$)		
Very rejecting	67	10.9
Somewhat rejecting	103	16.7
Neither rejecting nor accepting	61	9.9
Somewhat accepting	142	23.1
Very accepting	228	37.0
Prefer not to answer or not applicable	15	2.4
Dad's Acceptance of Sexual Orientation ($M=3.10$, $SD=1.43$)		
Very rejecting	100	16.2
Somewhat rejecting	122	19.8
Neither rejecting nor accepting	89	14.4
Somewhat accepting	128	20.8
Very accepting	124	20.1
Prefer not to answer or not applicable	53	8.6
Sexual Partners		
Only guys	225	36.5
Mostly guys but some girls	43	7.0
Guys and girls equally	58	9.4
Mostly girls but some guys	53	8.6
Only girls	25	4.1
None - never had sex	212	34.4

Table 2.

Means and Standard Deviations of all Items on the Measure of Adolescent Comfort with Clinical, Research, and Everyday Experiences (MACCREE)

Question	N	Mean	SD
<i>Routine Medical and Psychological Tests (M = 4.43, SD = 0.92)</i>			
Having your vision checked at the doctor's office	616	5.59	1.32
A doctor asking you about the types of foods you usually eat and drink	616	5.43	1.52
Answering a questionnaire about your future career or job interests at your school counselor's or psychotherapist's office	615	5.40	1.59
Getting an X-ray to check your bones at the doctor's office	616	5.38	1.40
A doctor asking you about your alcohol and drug use	614	4.79	1.84
Providing a urine sample at the doctor's office to be tested for sexually transmitted infections.	614	4.79	1.63
Answering a questionnaire about your mood at your school counselor's or psychotherapist's office	615	4.25	1.74
Having your blood drawn at the doctor's office	615	3.99	1.94
A doctor asking you about your sexual behavior	616	3.50	1.68
Having a doctor do a full-body skin exam for spots that could be cancerous	616	2.92	1.69
Providing a sample of your poop at the doctor's office	614	2.66	1.55
<i>Everyday Events for Adolescents (M = 3.49, SD = 0.58)</i>			
Sending a text message to your friend	616	6.30	1.09
Talking to your classmates about an after-school activity	616	6.18	1.13
Posting a picture on social media	616	5.27	1.55
Getting drug tested at school to participate in a sport or extracurricular activity	612	4.75	1.94
Getting caught in the rain	615	4.69	1.63
Taking a test at school	616	4.63	1.64
Having the principal of your school observe your class	615	4.37	1.63
Doing homework	615	4.11	1.61
Getting called on in class	616	3.82	1.89
Being asked to donate money or sign a petition on the street	616	3.76	1.62
Standing in a long line at a store	616	3.47	1.37
Being grounded for something you did	616	2.98	1.35
Picking up dog poop	614	2.85	1.47
Forgetting to do your homework	616	2.69	1.52
Getting a detention	615	2.52	1.37
Getting into an argument with a friend	615	2.34	1.07
Finding out a friend was talking about you behind your back	616	1.98	1.03
Sending an embarrassing text message to the wrong person	615	1.95	1.10
Having your partner break up with you	604	1.65	0.90
Having someone read your private blog or journal who was not supposed to see it	614	1.64	0.93
Having your cell phone stolen	615	1.25	0.59
<i>SGM Identity and Sexual Health (M = 3.57, SD = 1.08)</i>			
Talking about sex with your friends	615	5.37	1.67

Question	N	Mean	SD
Discussing sexual issues in a health class	615	4.33	1.71
Talking about sex with your parent or guardian	616	2.44	1.47
Coming out to a parent or guardian	615	2.13	1.47
<i>HIV/Sexual Health Research Procedures (M = 4.19, SD = 0.94)</i>			
Filling out survey questions about your sexual orientation or gender identity for a research study	616	6.02	1.23
Having the inside of your mouth swabbed with a Q-tip to test your saliva for HIV during a research study	615	5.34	1.47
Filling out survey questions about your sexual behavior for a research study	615	5.14	1.52
Talking to a researcher about your alcohol or drug use	613	4.97	1.63
Taking a pill that can help you prevent HIV every day for a year-long research study	616	4.84	1.70
Getting your urine tested for STIs during a research study	614	4.81	1.64
Having your finger pricked to test your blood for HIV during a research study	616	4.60	1.68
Being interviewed with a group of other teenagers about sexual issues for a research study	613	4.53	1.70
Getting your urine tested for drugs during a research study	613	4.48	1.80
Getting an injection (a shot) of medication every three months that can help you prevent HIV for a research study	616	4.38	1.87
Talking to a researcher about your sexual behaviors	615	4.05	1.73
Having to ask your parents for permission to participate in a research study about your alcohol and drug use	612	3.65	1.91
Being in a study where you don't know, and don't get to decide, whether you get an HIV prevention pill or a placebo	609	3.32	1.78
Having a matchstick-sized implant containing medication inserted in your upper arm that can help you prevent HIV for a year in a research study	615	3.32	1.76
Having to ask your parents for permission to participate in a research study for LGBT teens	614	3.12	2.01
Having to ask your parents for permission to participate in a research study about your sexual behavior	615	2.56	1.57
Having the inside of your butt swabbed to test for STIs during a research study	615	2.07	1.40

Note. Items were rated on a 7-point scale ranging from extremely uncomfortable (1) to extremely comfortable (7)

Table 3.
Multiple Regression Results for Demographic Predictors of MACCREE Domain Scores

	RMPT		EEA		SGM		HRP	
	B	p-value	B	p-value	B	p-value	B	p-value
Outness	0.163 [-0.003, 0.328]	0.054	0.035 [-0.070, 0.140]	0.512	0.239 [0.049, 0.429]	0.014	0.240 [0.072, 0.407]	0.005
Parental Acceptance	0.100 [0.038, 0.161]	0.002	0.067 [0.028, 0.106]	0.001	0.276 [0.205, 0.347]	0.000	0.146 [0.084, 0.208]	0.000
Race - Black	0.533 [0.209, 0.898]	0.002	0.058 [-0.160, 0.276]	0.602	0.198 [-0.198, 0.594]	0.327	0.405 [0.056, 0.754]	0.023
Race - Latino	-0.069 [-0.236, 0.098]	0.418	-0.085 [-0.191, 0.021]	0.115	0.088 [-0.104, 0.281]	0.366	0.023 [-0.146, 0.192]	0.790
Race - Other	-0.105 [-0.349, 0.138]	0.396	-0.169 [-0.323, -0.015]	0.032	-0.124 [-0.404, 0.157]	0.387	-0.120 [-0.367, 0.126]	0.338
Cisgender	0.070 [-0.105, 0.245]	0.431	0.141 [0.030, 0.252]	0.013	-0.082 [-0.284, 0.119]	0.423	-0.096 [-0.274, 0.081]	0.287
AFAB	-0.368 [-0.530, -0.207]	0.000	-0.273 [-0.375, -0.170]	0.000	-0.078 [-0.264, 0.108]	0.410	-0.238 [-0.401, -0.074]	0.004
Age	0.115 [0.042, 0.188]	0.002	0.029 [-0.017, 0.075]	0.223	0.023 [-0.061, 0.106]	0.591	0.096 [0.022, 0.169]	0.011
Sexually Active	0.115 [-0.038, 0.268]	0.141	0.035 [-0.062, 0.132]	0.484	0.157 [-0.019, 0.334]	0.080	0.270 [0.114, 0.425]	0.001
Adjusted R ²	0.124		0.114		0.144		0.131	

RMPT: Routine Medical and Psychological Tests; **EEA:** Everyday Events for Adolescents; **SGM:** SGM Identity and Sexual Health;

HRP: HIV/Sexual Health Research Procedures; **AFAB:** Assigned female at birth