Published in final edited form as:

Early Hum Dev. 2019 March ; 130: 109–115. doi:10.1016/j.earlhumdev.2019.01.019.

Combined predictors of neurodevelopment in very low birth weight preterm infants

Pilar Medina-Alva, MD^{1,2}, Kevin R. Duque, MD¹, Alonso Zea-Vera, MD¹, Sicilia Bellomo, MD^{1,3}, Cesar Carcamo, MD, PhD⁴, Daniel Guillen-Pinto, MD, PhD^{1,3}, Maria Rivas, MD⁵, Alfredo Tori, MD⁶, Jaime Zegarra, MD^{1,3}, Luis Cam, MD⁷, Anne Castañeda, MD⁶, Aasith Villavicencio, MD¹, and Theresa J. Ochoa, MD^{1,8,9}

¹School of Medicine, Universidad Peruana Cayetano Heredia. Av. Honorio Delgado 430, San Martín de Porres, Lima 15102, Peru.

²Instituto Nacional Materno Perinatal. Jr Santa Rosa 941, Cercado de Lima, Lima 15001, Perú.

³Hospital Nacional Cayetano Heredia, Av Honorio Delgado 262, San Martín de Porres Lima 15102, Peru

⁴School of Public Health and Administration, Universidad Peruana Cayetano Heredia, Av. Honorio Delgado 430, San Martín de Porres, Lima 15102, Perú.

⁵Hospital Nacional Madre Niño San Bartolome, Av Alfonso Ugarte 825, Lima 15001, Peru

⁶Hospital Nacional Guillermo Almenara, Av. Miguel Grau 800, La Victoria 15033, Lima, Peru

⁷Hospital Nacional Alberto Sabogal Sologuren, Jr. Colina 1081, Bellavista 07011, Callao, Peru

⁸Instituto de Medicina Tropical “Alexander von Humboldt”, Universidad Peruana Cayetano Heredia, Av. Honorio Delgado 430, San Martín de Porres, Lima 15102, Peru.

Corresponding author: Pilar Medina-Alva, MD. Instituto Nacional Materno Perinatal. Jr. Santa Rosa 941, Lima 15001, Peru. mpilarmedinaa@gmail.com, (51) 987709754.

Author's Statement Page:

Drs Medina-Alva, Duque, Zea-Vera, and Ochoa conceptualized and designed the study, made interpretation of data, drafted the initial manuscript, reviewed and revised the manuscript critically for important intellectual content and approved the final version to be published.

Drs Carcamo performed statistical analysis and interpretation of data, revised the manuscript critically for important intellectual content and approved the final version to be published.

Dr Duque also performed statistical analysis and interpretation of data.

Drs Guillen-Pinto, Rivas, Tori, Zegarra, Bellomo, Cam and Castañeda collected data, reviewed and revised the manuscript critically for important intellectual content and approved the final version to be published.

Dr Ochoa also obtained funding and supervised the study.

All authors approved the final manuscript as submitted.

All authors agree to be accountable for all aspects of the work ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Financial Disclosure: The authors have no financial relationships relevant to this article to disclose.

Potential Conflicts of Interest: The authors have no conflicts of interest relevant to this article to disclose

Clinical Trial Registration: Lactoferrin for Prevention of Sepsis in Infants (NEOLACTO), NCT01525316, <https://clinicaltrials.gov/ct2/show/NCT01525316>

CONFLICT OF INTEREST STATEMENT

None to declare

⁹Center for Infectious Diseases, University of Texas Health Science Center at Houston, Houston, Texas, United States.

Abstract

Objective: To evaluate the combined prognostic value of neurological examination, head circumference and cranial ultrasound for neurodevelopmental delay (NDD) in very low birth weight (VLBW, <1500 gr) preterm infants.

Methods: Prospective follow-up study. Preterm infants with VLWB were assessed for NDD using the Mullen Scales of Early Learning test at 24 months of corrected age. Abnormal neurological examination (≥ 2 deviant items of Hammersmith neurological examination), microcephaly and major ultrasound abnormalities, each performed at term age, were evaluated as predictors of NDD in a multivariable Poisson model.

Results: 35/132 infants (26.5%) had NDD. In the multivariable analysis, microcephaly (RR, 3.2; 95% CI, 1.6–6.7) and major ultrasound abnormalities (RR, 2.7; 95% CI, 1.3–5.7) were associated to NDD. The combination of the two tests showed the highest positive predictive value (100%; 95% CI, 51%–100%), while the combination of normal neurological examination, no major US findings and normal head size at term showed the highest negative predictive value (89%; 95% CI, 78%–95%). The maximum under receiver operating characteristic curve area was for microcephaly or major ultrasound abnormalities (AUC 0.74 (0.65–0.83)).

Conclusion: The combination of head circumference, cranial ultrasound and neurological examination at term age is useful to predict NDD in VLBW preterm infants.

Keywords

neurodevelopment; preterms; predictive value; predictors

INTRODUCTION

Preterm birth is still a global health problem, especially in developing countries where access to optimal obstetric and neonatal care is not guaranteed (1)(2). In South America, mortality rates in very low birth weight (VLBW) infants reach 26% (3). Preterm survival rates have increased in the last decades, making follow up essential. It is important not only to detect cerebral palsy and motor delay but also to timely recognize neurodevelopmental disorders, which can affect 50% of VLBW infants (4).

Prediction of neurodevelopment can be attempted with neurological examination at term age. Standardized tools have facilitated neurological assessment execution, though its capacity for prediction seems to be limited (5)(6). Head circumference is an easy-to-obtain parameter that correlates with brain volume and is a well known predictor of neurodevelopment (7). Cranial ultrasound, routinely performed at bedside, can detect brain abnormalities –such as severe intraventricular hemorrhage, ventriculomegaly, periventricular leukomalacia and brain infarction– which might influence future neurodevelopment (8). It is especially useful in settings where magnetic resonance imaging (MRI) is not routinely obtained because of cost or limited access.

Low-cost, effective strategies which can improve detection of patients at risk of developmental abnormalities are of particular interest in developing countries (9). Neurological examination, head circumference measurement and routine cranial ultrasound are easy to perform, inexpensive and routinely performed in VLBW babies. We hypothesized that the combination of these assessments could improve the prognostic value of each test alone to detect neurodevelopmental delay (NDD).

The aim of this study is to evaluate the combined prognostic value of neurological examination, head circumference and cranial ultrasound, each at term age, for NDD assessed at 24 months corrected age.

METHODS

Study population

The NEOLACTO study was a multicenter clinical trial (10) which enrolled newborns with birth weights between 500 and 2000 gr, and aimed to evaluate the efficacy of lactoferrin supplementation for sepsis prevention and its effect on neurodevelopment. Infants were enrolled at neonatal intensive care units of three large tertiary-care hospitals in Lima, Peru. Infants with conditions that profoundly affect growth and development such as chromosomal, congenital or brain abnormalities or infants with gastrointestinal problems were excluded from the clinical trial. Since lactoferrin showed no effect on neurodevelopment (10), all VLBW preterm infants followed up until 24 ± 1 months of corrected age with complete examinations at 40 ± 2 weeks of corrected age were included in this study. The clinical trial was approved by the Institutional Review Boards of Universidad Peruana Cayetano Heredia, University of Texas Health Science Center in USA and of each participating hospital. Both parents provided written informed consent.

Data collection

Infants were enrolled in the first 72 hours of life and followed until 24 months of corrected age. Gestational age was determined by first-trimester ultrasound, or in its absence by last menstrual period, or if both were absent by Ballard test. Child neurologists performed all examinations: neurological examinations, measurements of head circumference and cranial ultrasound at day 3–5 after birth, week 3–4 and 40 ± 2 weeks of corrected age. Results of examinations at 40 ± 2 weeks of corrected age were evaluated as potential predictors.

We used the Hammersmith Neurological Examination, consisting of 34 items which assesses posture and tone (ten items), tone patterns (five items), reflexes (six items), spontaneous movements (three items), abnormal signs (three items) and behavior (seven items). It has been standardized for preterm infants and has showed good intra and inter-observer reliability (11). Scores of individual items outside the standardized 90th-centile range according to the gestational age at birth were considered deviants(12).

Head circumference was considered as the largest occipitofrontal circumference measurement taken with a non-distensible tape. Microcephaly was defined as a head circumference Z-score >2 standard deviation (SD) below the mean according to age and sex of Fenton preterm growth charts (13). Cranial ultrasound (US) was performed with 7.5 mHz

transducers and portable sonography devices (Sonoscape S6, Sonoscape Co., China and Sonosite180, Sonosite Inc, USA). Brain infarction, parenchymal hemorrhage, severe intraventricular hemorrhage (grades 3 or 4), post-hemorrhagic hydrocephalus with ventricular index >14mm and periventricular leukomalacia (periventricular echogenicity persistent for 7 days and/or periventricular cysts) (14) were deemed as major US abnormalities if found in any of the 3 control dates. Minor cerebral abnormalities were intraventricular hemorrhage (grades 1 and 2), transitory periventricular echogenicity (as disappearance at 7 days or before), ventricular dilatation with ventricular index <14mm, lenticulostriatal vasculopathy and widened subarachnoid space.

Neurodevelopmental evaluation

Two trained pediatricians administered the Mullen Scales of Early Learning (MSEL) to assess neurodevelopmental outcomes at 24 ± 1 months of corrected age. This test assesses child's learning competencies among five domains: gross motor, visual reception, fine motor, receptive language and expressive language abilities, and shows moderate correlation with the scores from the Bayley Scores of Infant Development (15). Scores of the last four domains yield the *Early Learning Composite Score* which has been standardized for infants from birth to 68 months with a mean $\pm$ SD score of 100 ± 15 . Neurodevelopmental delay was defined as an Early Learning Composite Z-score 2 SD below the mean (standard score 70). All infants diagnosed with cerebral palsy (16) had such composite score <70; hence our NDD definition included these infants.

Statistical analyses

For bivariate analysis, Fisher's exact test, Student's *t*-test and Mann-Whitney test were used as appropriate. Emphasizing clinical relevance, neurological examination was dichotomized. A proposed cutoff of 1 deviant items to define abnormal neurological examination (12) was not associated with NDD in our sample in a bivariate analysis, hence the immediately superior and associated cutoff of 2 was used to define it. Abnormal neurological examination, microcephaly and major US abnormalities entered into a multivariable Poisson linear model to assess their relative risks (RR) for NDD. Effect modification between these three tests was assessed. Abnormal neurological examination was not associated to NDD in the multivariable model nor showed effect modification on the other examinations, hence it was excluded from the final model. Positive predictive (proportion of infants with NDD among those with positive tests) and negative predictive (proportion of infants with normal neurodevelopment among those with normal examinations) values and areas under receiving operating characteristic (ROC) curves (AUC) with confidence intervals (CI) were calculated for each examination and for different results of the three tests assessed in parallel. Multiple chi-squared tests on the equality of two AUC were used to assess the difference between the three combined predictors and each predictor alone. McNemar's test was used to assess the potential advantage of both head circumference and cranial US on neurological examination in determining more infants at risk of NDD. All reported p-values are two-sided and those <0.05 are considered statistically significant. STATA v.14.0 (StataCorp LP, USA) was used.

RESULTS

From May 2012 to September 2014, 414 newborns with birth weights between 500 and 2000 gr were enrolled in the clinical trial; 255 of them (61.6%) were preterms with birth weights between 500 and 1500 gr. Of these 255 infants, 132 (51.8%) were followed-up until 24 months of corrected age and had complete examinations at 40 ± 2 weeks (Fig. 1). From the 123 infants not included in the analysis, 63 died, mainly of sepsis (Supplementary Table 1) and 42 had neurological examinations out of time (Fig. 1). Baseline characteristics of the infants included in this study are depicted in Table 1. Compared to infants who were not included, included infants had higher gestational age (median 30 vs 29 weeks, $p < 0.001$), higher birth weight (median 1290 vs 1050 grams, $p < 0.001$), less often use of mechanical ventilation (31.3% vs 70.7%, $p < 0.001$), less often necrotizing enterocolitis (6.8% vs 20.3%, $p < 0.001$) and intraventricular hemorrhage (11.4% vs 28.4%, $p < 0.001$; Supplementary Table 2). In the same way, relative to infants followed-up until 24 months but without neurological examination at term age, included infants had higher birth weight (median 1290 vs 1171, $p = 0.02$), less often use of mechanical ventilation (31.1% vs 50.0%, $p = 0.04$), less often necrotizing enterocolitis (6.8% vs 19.1%, $p = 0.03$).

Neurodevelopmental delay and potential predictors

Thirty-five infants (26.5%) had NDD at 24 months of corrected age. Randomized, double-blind, placebo-controlled lactoferrin administration for eight weeks did not result in NDD prevention among this subgroup of infants from the parent clinical trial (17 infants had NDD among 35 who received lactoferrin [48.2%] versus 53/97 [54.6%] who received placebo; RR 0.84, 95% CI 0.5–1.5, $p = 0.54$). In a bivariate analysis, abnormal neurological examination with a cutoff of ≥ 2 deviant items was associated to NDD (RR, 2.0; 95% CI, 1.1–3.7) but not with the proposed cutoff of ≥ 1 (12) (RR, 0.9; 95% CI, 0.4–1.8). Most frequent deviant items were: visual orientation (58 infants [43.4%]), flexor head control (21.2%), leg traction (15.9%), alertness (14.4%), palmar grasp (12.1%) and arm recoil (10.6%).

In a multivariable analysis, only microcephaly (RR, 3.2; 95% CI, 1.6–6.7) and major US abnormalities (RR, 2.7; 95% CI, 1.3–5.7) were associated to NDD. Abnormal neurological examination was not associated to NDD nor showed effect modification on the other examinations, hence it was not included in the final model showed in Table 2.

Predictors' performance

Predictive values, sensitivity, specificity and AUC of each examination alone and combined are presented in Table 3. Comparing examinations individually, microcephaly showed the highest positive predictive value (80%; 95% CI, 55%–93%) and the highest AUC (0.66; 95% CI, 0.57–0.74), and normal neurological examination, the highest negative predictive value (83%; 95% CI, 71%–90%). Microcephaly and major US abnormalities had an AUC of 0.74 (95% CI, 0.65–0.83). The three examinations together showed an AUC of 0.78 (95% CI, 0.68–0.87) which was higher than abnormal neurological evaluation alone (0.17 points of AUC higher, 95% CI 0.06–0.28, $p < 0.001$), microcephaly alone (0.12 points of AUC higher, 95% CI 0.01–0.23, $p = 0.004$), and major US abnormalities alone (0.14 points of AUC higher, 95% CI 0.03–0.25, $p = 0.001$; Figure 2).

Considering the three examinations assessed in parallel, microcephaly and major US abnormalities, regardless of the neurological examination, showed the highest positive predictive value (100%; 95% CI, 51%–100%). The combination of normocephaly, no major US abnormalities and normal neurological examination showed the highest negative predictive value (89%; 95% CI, 78%–95%). The combination of normocephaly and no major US abnormalities showed a similar predictive value (85%; 95% CI, 77%–91%) ($p=0.06$). In our sample, neurological examination missed 11 of 35 (31%) infants who developed NDD which were reduced to six infants (17%) with the combination of neurological examination, head circumference and cranial ultrasound.

Infants with cerebral palsy

Eight infants had cerebral palsy and NDD. They had similar gestational age (28.3 ± 2.5 weeks), birth weight (1117 ± 237 gr) and proportion of male gender (50%) than other infants with neurodevelopmental impairment. Major US abnormalities were found in six infants (75%), microcephaly at term age in five (62.5%) and seven (87.5%) had abnormal neurological examinations. All infants with cerebral palsy had two or more abnormal tests, except one infant who had only one predictor which was a grade IV intraventricular hemorrhage. Infants with cerebral palsy had a median of three deviant items (interquartile range 2–6.5) compared to two deviant items (interquartile range 0–3) among infants with NDD but no cerebral palsy.

DISCUSSION

In this study we used the Hammersmith neurological examination (11), a tool originally designed for term infants. The authors revisited the scale and eliminated certain items that were too difficult to measure in NICU settings, were too variable or had too little capacity to discriminate between normal and abnormal (17). Some authors have evaluated its use in preterms, and found that ranges of normal values are wider in this population, specifically in the areas of behavior and muscle tone (18) (19) (20). Comparing scores for preterm and term babies, items outside 90% are similar in both groups, which suggest that some neurological signs are infrequent at 40 weeks corrected age regardless of gestational age at birth (21).

The ability of a systematic neurological examination to predict neurodevelopmental abnormalities in preterms has been widely explored, and showed variable results. One study found that a non-optimal neurological status at term correlates with lower performances in standard neurodevelopment tests (5). On the contrary, other studies showed that the majority of preterm babies with non-optimal evaluations at term age have normal development. These findings might suggest that exam abnormalities might be due to transient postural disturbances or maturation delay patterns in otherwise healthy preterms (22). Our findings coincide with this observation, which could explain the very low positive predictive and high negative predictive values found for abnormal neurological examination at term.

Microcephaly is associated with a reduction of brain tissue volumes, especially deep nuclear gray matter (7). Microcephaly at birth is frequent, affecting 7–12% of VLBW infants, whereas suboptimal head circumference (defined as a Z-score >1 SD below the mean) can be detected in 25–30% preterms (7)(24). Percentages of microcephaly in our population of

low birth weight preterms were higher and reached 17% at birth and 29% at discharge (10). In large cohorts of VLBW infants, proportions of babies with microcephaly or suboptimal head circumference increased significantly at 2 year follow up evaluations and correlate with impaired developmental outcomes (25)(26)(27), as found in our study. Greater head growth in preterms from birth to term age is associated with higher cognitive and motor scores at 18 months (28); however, some studies suggest that head growth at post-discharge might be a better predictor of cognitive outcomes (29)(30).

Cranial ultrasound (US) is the preferred neuroimaging study at NICU because of its feasibility and availability (31). In low-resource settings, its cost represents a great advantage over MRI. The predictive value of cranial US has been addressed by several studies. Sensitivity and specificity were 43–44% and 82–87% respectively to predict cerebral palsy when sonograms were performed within the first five days of life. Predictive accuracy is better with additional sonograms, especially if the additional study is performed between 36 and 40 weeks corrected age (8)(32). In our study we performed three US evaluations until term age, and the specificity and negative predictive value for developmental impairment were high (96% and 79% respectively). Out of infants with cerebral palsy, 75% had major abnormalities in brain US. Ultrasound might show less sensitivity for less severe white matter lesions (33); therefore, its capacity to predict non-motor disabilities is yet to be determined (34), although ventriculomegaly (with or without IVH) is associated with increased risk of cognitive impairment (35)(36). In some studies, cranial US showed higher predictive capacity for neurodevelopmental abnormalities at one year of age when compared to neurological examination (22).

Predictive values of tests performed in parallel might be statistically higher than predictive values of each test. Maas (37) compared predictive values of several tests alone and in combination with other tools (brain ultrasound, generalized movements test, sleep organization test) in 100 babies with gestational age less than 30 weeks. They found that Prechtl neurological test at term corrected age had the highest predictive value, comparable to that of cranial US, but they could not find an additive effect on prediction power of either test. Other authors compared generalized movement evaluation and brain ultrasound and found high predictive value. (38) Although it is a proven predictive tool for NDD, especially at 4 months corrected age (39), general movements evaluation is not routinely performed in our country because of the lack of trained personnel. Sätanen found that neurological examination at term age combined with brain MRI or cranial US findings improved the predictive value of either neuroimaging test alone when considering neurosensory outcome at 2 years corrected age (12).

To our knowledge it is the first study to combine these specific three predictors of neurodevelopment in VLBW preterms. We evaluated this unique combination of three tests to improve prediction, and found excellent positive predictive values when all three tests were abnormal, and high negative predictive values when either of the tests was normal. The combination of tests already in use for the evaluation of preterms in hospitals could be a useful tool for health care practitioners in charge of preterm follow up. We used serial brain sonography tests, a strategy that helps increasing the predictive value of this tool.

Our study has limitations. Our sample size was not large, and although not a significantly different population, our study had a great loss of patients because of untimely evaluation. Also, neurological evaluation was performed by four different trained pediatric neurologists, and as such this could lead to greater interrater variability. However, we used a standardized examination to prevent this from happening. The tool we chose for neurodevelopmental evaluation (MSEL) is widely used in preterm population and has close correlation with Bayley's psychomotor and mental indexes (40), but its use limits comparisons with results of similar studies.

CONCLUSION

Our study shows that the combination of findings on neurological examination at term, head size at term, and presence of major abnormalities in serial brain ultrasound is a good predictor of neurodevelopmental impairment in VLBW infants at 2 years corrected age. They can be a useful tool for selection of patients in low-resource settings for early intervention strategies.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENTS

NEOLACTO Research Group: In addition to the bylined authors, the research group also includes: Alicia Villar, MD; Carolina Navarro, MD; Patricia Mallma, MD; Renzo Acosta, MD; Miguel Campos, MD, PhD; Lourdes Tucto, RN; Cristina Suarez, RN; Nadia Rojas, RN; Irene Chea, RN; and Mayela Huanay, RN (Universidad Peruana Cayetano Heredia, Lima, Peru); María Luz Rospigliosi, MD; Geraldine Borda, MD; Oriolit Minauro, MD; Verónica Webb, MD; Erika Bravo, MD; Karen Pacheco, MD; and Wilder Alarcon, MD (Hospital Cayetano Heredia, Lima, Peru); Ana Lino, MD; and Augusto Cama, MD (Hospital Nacional Alberto Sabogal, Lima, Peru); Raúl Llanos, MD; Oscar Chumbes, MD; Liliana Cuba, MD; Julio Tresierra, MD; and Carmen Chincaro, MD (Hospital Nacional Guillermo Almenara, Lima, Peru).

Funding Source: Funded by the National Institute of Child Health & Human Development (NICHD), Grant number: R01-HD067694.

REFERENCES

1. Frey HA, Klebanoff MA. The epidemiology, etiology, and costs of preterm birth. *Semin Fetal Neonatal Med* 2016 4;21(2):68–73. doi: 10.1016/j.siny.2015.12.011 [PubMed: 26794420]
2. Harrison MS, Goldenberg RL. Global burden of prematurity. *Semin Fetal Neonatal Med* 2016 4;21(2):74–9. doi: 10.1016/j.siny.2015.12.007 [PubMed: 26740166]
3. Fernández R, D'Apremont I, Domínguez A, et al. Survival and morbidity of very low birth weight infant in a South American neonatal network. *Arch Argent Pediatr* 2014 10;112(5):405–12. doi: 10.5546/aap.2014.405 [PubMed: 25192520]
4. Volpe JJ. Brain injury in premature infants: a complex amalgam of destructive and developmental disturbances. *Lancet Neurol* 2009 1;8(1):110–24. doi: 10.1016/S1474-4422(08)70294-1 [PubMed: 19081519]
5. Simard M-N, Lambert J, Lachance C, et al. Prediction of developmental performance in preterm infants at two years of corrected age: Contribution of the neurological assessment at term age. *Early Hum Dev* 2011 12;87(12):799–804. doi: 10.1016/j.earlhumdev.2011.06.004 [PubMed: 21723051]
6. Leroux BG, N'Guyen The Tich S, Branger B, et al. Neurological assessment of preterm infants for predicting neuromotor status at 2 years: results from the LIFT cohort. *BMJ Open* 2013;3:e002431. doi:10.1136/bmjopen-2012-002431

7. Cheong JLY, Hunt RW, Anderson PJ, et al. Head growth in preterm infants: correlation with magnetic resonance imaging and neurodevelopmental outcome. *Pediatrics* 2008 6 1;121(6):e1534–40. doi: 10.1542/peds.2007-2671 [PubMed: 18519457]
8. De Vries LS, Van Haastert I-LC, Rademaker KJ, et al. Ultrasound abnormalities preceding cerebral palsy in high-risk preterm infants. *J Pediatr* 2004 6;144(6):815–20. doi: 10.1016/j.jpeds.2004.03.034 [PubMed: 15192633]
9. Sutton PS, Darmstadt GL. Preterm birth and neurodevelopment: A review of outcomes and recommendations for early identification and cost-effective interventions. *J Trop Pediatr* 2013 8 1;59(4):258–65. doi: 10.1093/tropej/fmt012 [PubMed: 23486391]
10. Ochoa TJ, Zegarra J, Bellomo S, Carcamo CP, Cam L, Castañeda et al. Long-term outcomes in infants <2000 g after neonatal lactoferrin supplementation. In XIIIth International Conference on Lactoferrin: Structure, Function and Applications. Rome, Italy; 2017.
11. Dubowitz V, Dubowitz L, Mercuri E. The neurological assessment of the preterm and full-term newborn infant 2nd ed. London: Mc Keith Press; 1999 (Mc Keith Press; 1999.
12. Setänen S, Lahti K, Lehtonen L, et al. Neurological examination combined with brain MRI or cranial US improves prediction of neurological outcome in preterm infants. *Early Hum Dev* 2014 12;90(12):851–6. doi: 10.1016/j.earlhumdev.2014.09.007 [PubMed: 25463831]
13. Fenton TR, Kim JH. A systematic review and meta-analysis to revise the Fenton growth chart for preterm infants. *BMC Pediatr* 2013 4 20;13:59. doi: 10.1186/1471-2431-13-59 [PubMed: 23601190]
14. Volpe JJ. Volpe's neurology of the newborn Sixth edition. Philadelphia, PA: Elsevier; 2018 1224 p.
15. Mullen EM. Mullen Scales of Early Learning. AGS Edition Minneapolis, MN; 1993.
16. Mutch L, Alberman E, Hagberg B, et al. Cerebral palsy epidemiology: where are we now and where are we going? *Dev Med Child Neurol* 1992 6;34(6):547–51. doi: 10.1111/j.1469-8749.1992.tb11479.x [PubMed: 1612216]
17. Dubowitz L, Mercuri E, Dubowitz V. An optimality score for the neurologic examination of the term newborn. *J Pediatr* 1998 9;133(3):406–16. doi: 10.1016/S0022-3476(98)70279-3 [PubMed: 9738726]
18. Ricci D, Romeo DMM, Haataja L, et al. Neurological examination of preterm infants at term equivalent age. *Early Hum Dev* 2008 11;84(11):751–61. doi:10.1016/j.earlhumdev.2008.05.007 [PubMed: 18614301]
19. Romeo DM, Ricci D, Brogna C, et al. Neurological examination of late-preterm infants at term age. *Eur J Paediatr Neurol EJPN Off J Eur Paediatr Neurol Soc* 2011 7;15(4):353–60. doi: 10.1016/j.ejpn.2011.03.009
20. Mercuri E, Guzzetta A, Laroche S, et al. Neurologic examination of preterm infants at term age: comparison with term infants. *J Pediatr* 2003 6;142(6):647–55. doi: 10.1067/mpd.2003.215 [PubMed: 12838193]
21. Romeo DM, Ricci D, van Haastert IC, et al. Neurologic assessment tool for screening preterm infants at term age. *J Pediatr* 2012 12;161(6):1166–8. doi: 10.1016/j.jpeds.2012.07.037 [PubMed: 22910101]
22. Amess P, McFerran C, Khan Y, et al. Early prediction of neurological outcome by term neurological examination and cranial ultrasound in very preterm infants. *Acta Paediatr Oslo Nor* 1992 2009 3;98(3):448–53. doi: 10.1111/j.1651-2227.2008.01062.x
23. Karagianni P, Rallis D, Kyriakidou M, et al. Correlation of brain ultrasonography scans to the neuromotor outcome of very-low-birth-weight infants during the first year of life. *J Child Neurol* 2014 11;29(11):1429–35. doi: 10.1177/0883073813479171 [PubMed: 23470630]
24. Hack M, Breslau N, Weissman B, et al. Effect of very low birth weight and subnormal head size on cognitive abilities at school age. *N Engl J Med* 1991 7 25;325(4):231–7. doi: 10.1056/NEJM199107253250403 [PubMed: 2057024]
25. Neubauer V, Griesmaier E, Pehböck-Walser N, et al. Poor postnatal head growth in very preterm infants is associated with impaired neurodevelopment outcome. *Acta Paediatr* 2013 9;102(9):883–8. doi: 10.1111/apa.12319 [PubMed: 23772884]

26. Kuban KCK, Allred EN, O'Shea TM, et al. Developmental correlates of head circumference at birth and two years in a cohort of extremely low gestational age newborns. *J Pediatr* 2009 9;155(3):344–349.e1–3. doi: 10.1016/j.jpeds.2009.04.002 [PubMed: 19555967]
27. Chiriboga CA, Kuban KCK, Durkin M, et al. Factors associated with microcephaly at school age in a very-low-birthweight population. *Dev Med Child Neurol* 2003 12;45(12):796–801. doi: 10.1111/j.1469-8749.2003.tb00893.x [PubMed: 14667070]
28. Belfort MB, Rifas-Shiman SL, Sullivan T, et al. Infant growth before and after term: Effects on neurodevelopment in preterm infants. *Pediatrics* 2011 10 1;128(4): e899–906. doi: 10.1542/peds.2011-0282 [PubMed: 21949135]
29. Lidzba K, Rodemann S, Goelz R, et al. Growth in very preterm children: Head growth after discharge is the best independent predictor for cognitive outcome. *Early Hum Dev* 2016; 103:183–8. doi: 10.1016/j.earlhumdev.2016.09.016 [PubMed: 27716567]
30. Raghuram K, Yang J, Church PT, et al. Head Growth Trajectory and Neurodevelopmental Outcomes in Preterm Neonates. *Pediatrics* 2017 7;140(1):e20170216. doi: 10.1542/peds.2017-0216 [PubMed: 28759409]
31. Wezel-Meijler G van, de Vries LS. Cranial ultrasound - optimizing utility in the NICU. *Curr Pediatr Rev* 2014;10(1):16–27. doi: 10.2174/157339631001140408120106 [PubMed: 25055860]
32. DeVries L, Benders M, Groenendaal F. Progress in neonatal neurology with a focus on the neuroimaging in the preterm infant. *Neuropediatrics* 2015, 46: 234–241 [PubMed: 26121069]
33. Leijser LM, de Bruine FT, Steggerda SJ, et al. Brain imaging findings in very pre-term infants throughout the neonatal period: part I. Incidences and evolution of lesions, comparison between ultrasound and MRI. *Early Hum Dev* 2009;85: 101–9. [PubMed: 19144474]
34. Dorner RA, Burton VJ, Allen MC, Robinson S, Soares BP. Preterm neuroimaging and neurodevelopmental outcome: a focus on intraventricular hemorrhage, post-hemorrhagic hydrocephalus and associated brain injury. *J Perinatol* 2018 11;38(11):1431–1443. doi: 10.1038/s41372-018-0209-5 [PubMed: 30166622]
35. Wood NS, Costeloe K, Gibson AT, et al. The EPICure study: associations and antecedents of neurological and developmental disability at 30 months of age following extremely preterm birth. *Arch Dis Child Fetal Neonatal Ed* 2005 3;90(2):F134–140. doi: 10.1136/adc.2004.052407 [PubMed: 15724037]
36. Pappas A, Adams-Chapman I, Shankaran S, et al. Neurodevelopmental and behavioral outcomes in extremely premature neonates with ventriculomegaly in absence of periventricular-intraventricular hemorrhage. *JAMA Pediatr* 2018, 172(1):32–42. doi: 10.1001/jamapediatrics.2017.3545 [PubMed: 29181530]
37. Maas YG, Mirmiran M, Hart AA, et al. Predictive value of neonatal neurological tests for developmental outcome of preterm infants. *J Pediatr* 2000 7;137(1):100–6. doi: 10.1067/mpd.2000.106901 [PubMed: 10891830]
38. Seme-Ciglenecki P Predictive values of cranial ultrasound and assessment of general movements for neurological development of preterm infants in the Maribor region of Slovenia. *Wien Klin Wochenschr* 2007 8;119(15–16):490–6. doi: 10.1007/s00508-007-0839-7 [PubMed: 17721769]
39. Bosanquet M, Copeland L, Ware R, et al. A systematic review of tests to predict cerebral palsy in young children. *Dev Med Child Neurol* 2013 5;55(5):418–26. doi: 10.1111/dmcn.12140 [PubMed: 23574478]
40. Harel-Gadassi A, Friedlander E, Yaari M, et al. Developmental assesment of preterm infants: chronological or corrected age? *Research in Developmental Disabilities* 2018, 80: 35–43. doi: 10.1016/j.ridd.2018.06.002 [PubMed: 29906778]
41. WHO WMGRS. WHO Child Growth Standards: Growth velocity based on weight, length, and head circumference: methods and development Geneva, Switzerland; 2009 (reference to supplementary table 2)

HIGHLIGHTS**Table of contents summary:**

This article evaluates the value of combined data from head circumference, cranial ultrasound abnormalities and neurological examination at term to predict neurodevelopment in very low birth weight preterms.

What's known on this subject:

Neurodevelopmental disorders can affect 50% of very low birth weight infants. Combined tests may improve prediction of neurodevelopment. Neurological examination, head circumference and cranial ultrasound are easy to complete, inexpensive and routinely performed in VLBW babies.

What this study adds:

The combination of abnormal neurological evaluation at term, microcephaly at term and major findings in serial cranial ultrasound is a better predictor of neurodevelopmental delay in very low birth weight infants than either of these tests alone.

Figure 1.

Flow diagram of the prospective multicenter cohort study among VLBW infants, 2012–2014

Figure 2.
Sensitivity versus specificity plot of five results among three examinations to predict neurodevelopmental delay in VLBW infants from a prospective multicenter cohort study, 2012–2014

Table 1.

Baseline characteristics of VLBW infants from a prospective multicenter cohort study, 2012–2014

Characteristics	Total (n=132)
Pregnancy and delivery	
Maternal age (years)	29.7±6.2
Single mother ^a	8 (6.1%)
Mother with higher education ^b	60 (45.5%)
Monthly family income (US dollars)	422 (312–660)
Multiple pregnancy	36 (27.3%)
Cesarean delivery	108 (81.8%)
Neonatal	
Male sex	72 (54.6%)
Gestational age at birth (weeks)	30±2.4
24–28 weeks	20 (15.2%)
28–29 weeks	38 (28.8%)
30–31 weeks	36 (27.3%)
32–36 weeks	38 (28.8%)
Birth weight (grams)	1290 (1103–1386)
5-min Apgar score	8.1±1.4
Neonatal resuscitation needed	93 (70.5%)
Small for gestational age ^c	36 (27.3%)
Microcephaly at birth	17/114 (14.9%)
During hospitalization	
Use of total parenteral nutrition	118 (89.4%)
Time with total parenteral nutrition (days) ^d	10.5 (6.5–17)
Age at oral feeding initiation (days after birth)	4.0±2.4
Age at establishment of full enteral feeding (days after birth)	15 (10–21)
Culture-proven late onset sepsis	16 (12.1%)
Culture-proven late onset sepsis or probable sepsis	29 (22.0%)
Use of mechanical ventilation	41 (31.1%)
Time with mechanical ventilation (days) ^e	7 (2–19)
Bronchopulmonary dysplasia	27 (20.5%)
Necrotizing enterocolitis (Modified Bell's stage 2)	9 (6.8%)
Intraventricular hemorrhage	15 (11.4%)
Grade 3 or 4	6 (4.6%)
Periventricular leucomalacia	8 (6.1%)
Growth restriction at discharge ^f	35/110 (31.8%)

Data are expressed in mean±SD, median (P25–P75) or n (%) as appropriate.

^aUnmarried or without a partner.^bHigher education defined as post-secondary education including college, university and/or institutes of technology.

^c Defined as weight below the 10th percentile for the gestational age at birth.

^d Frequency only among infants who were in total parenteral nutrition.

^e Frequency only among infants who were in mechanical ventilation.

^f Defined as a length below -2 standard deviation according the World Health Organization Child Growth Standards at hospital discharge (39).

Table 2.

Potential predictors at term age of neurodevelopment delay assessed at 24 months of corrected age among VLBW infants from a prospective multicenter cohort study, 2012–2014

Potential predictors	Result	Neurodevelopmental delay, n/N (cumulative incidence ^a)	Relative risk (95% CI)	
			Crude	Adjusted ^b
Abnormal neurological examination	No	11/63 (17.5)	2.0 (1.1–3.7)	
	Yes	24/69 (34.8)		
Microcephaly	No	23/117 (19.7)	4.1 (2.6–6.4)	3.2 (1.6–6.7)
	Yes	12/15 (80.0)		
Major brain abnormality	No	24/117 (20.5)	3.6 (2.2–5.7)	2.7 (1.3–5.7)
	Yes	11/15 (73.3)		

^a Cases/100 infants/2 years.

^b Calculated from a multivariable Poisson linear model in which the three potential predictors entered but then abnormal neurological examination was excluded because of lack of association

Performance of three examinations alone and combined to predict neurodevelopmental delay in VLBW infants from a prospective multicenter cohort study, 2012–2014

Table 3.

Potential predictors	Sensitivity	Specificity	Positive predictive value	Negative predictive value	Area under ROC curve (95% CI)
Abnormal neurological exam	69 (52–81)	54 (44–63)	35 (25–47)	83 (71–90)	0.61 (0.52–0.70)
Microcephaly	34 (21–51)	97 (91–99)	80 (55–93)	80 (72–87)	0.66 (0.57–0.74)
Major US abnormality	31 (19–48)	96 (90–99)	73 (48–89)	79 (71–86)	0.64 (0.56–0.72)
Microcephaly + major US abnormality					
Microcephaly or major US abnormality	54 (38–70)	93 (86–97)	73 (54–86)	85 (77–91)	0.74 (0.65–0.82)
Microcephaly and major US abnormality	11 (5–26)	100 (95–100)	100 (51–100)	76 (68–82)	0.56 (0.50–0.61)
Abnormal neurological exam + microcephaly + major US abnormality					
Abnormal neurological exam or microcephaly or major US abnormality	83 (67–92)	51 (41–60)	38 (28–49)	89 (78–95)	0.67 (0.59–0.75)
Abnormal neurological exam and microcephaly and major US abnormality	11 (4–26)	100 (95–100)	100 (51–100)	76 (68–82)	0.56 (0.50–0.61)

Data are presented as % (95% CI) except for area under ROC curve which is presented as area (95% CI).