Cochrane
Library

Cochrane Database of Systematic Reviews

Avoidance versus use of neuromuscular blocking agents for improving conditions during tracheal intubation or direct laryngoscopy in adults and adolescents (Review)

Lundstrøm LH, Duez CHV, Nørskov AK, Rosenstock CV, Thomsen JL, Møller AM, Strande S, Wetterslev J

Lundstrøm LH, Duez CHV, Nørskov AK, Rosenstock CV, Thomsen JL, Møller AM, Strande S, Wetterslev J.
Avoidance versus use of neuromuscular blocking agents for improving conditions during tracheal intubation or direct laryngoscopy in adults and adolescents.
Cochrane Database of Systematic Reviews 2017, Issue 5. Art. No.: CD009237.
DOI: [10.1002/14651858.CD009237.pub2](https://doi.org/10.1002/14651858.CD009237.pub2).

www.cochranelibrary.com

Avoidance versus use of neuromuscular blocking agents for improving conditions during tracheal intubation or direct laryngoscopy in adults and adolescents (Review)

Copyright © 2017 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

WILEY

TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
SUMMARY OF FINDINGS	4
BACKGROUND	6
OBJECTIVES	6
METHODS	6
RESULTS	10
Figure 1.	11
Figure 2.	13
Figure 3.	14
Figure 4.	15
Figure 5.	17
Figure 6.	19
DISCUSSION	19
AUTHORS' CONCLUSIONS	22
ACKNOWLEDGEMENTS	22
REFERENCES	23
CHARACTERISTICS OF STUDIES	28
DATA AND ANALYSES	85
Analysis 1.1. Comparison 1 Avoidance vs use of NMBA, Outcome 1 Difficult tracheal intubation: low risk of bias vs high or uncertain risk of bias.	88
Analysis 1.2. Comparison 1 Avoidance vs use of NMBA, Outcome 2 Difficult tracheal intubation: depolarizing vs non-depolarizing NMBA.	89
Analysis 1.3. Comparison 1 Avoidance vs use of NMBA, Outcome 3 Difficult tracheal intubation: remifentanyl vs no remifentanyl.	90
Analysis 1.4. Comparison 1 Avoidance vs use of NMBA, Outcome 4 Difficult tracheal intubation: alfentanil vs no alfentanil.	91
Analysis 1.5. Comparison 1 Avoidance vs use of NMBA, Outcome 5 Difficult tracheal intubation: local anaesthesia vs no local anaesthesia.	92
Analysis 1.6. Comparison 1 Avoidance vs use of NMBA, Outcome 6 Difficult tracheal intubation: excluded anticipated DTI vs included anticipated DTI.	93
Analysis 1.7. Comparison 1 Avoidance vs use of NMBA, Outcome 7 Difficult tracheal intubation: "best-case scenario".	94
Analysis 1.8. Comparison 1 Avoidance vs use of NMBA, Outcome 8 Difficult tracheal intubation excluding dose-finding studies.	95
Analysis 1.9. Comparison 1 Avoidance vs use of NMBA, Outcome 9 Difficult tracheal intubation: funding from pharmaceutical industry.	96
Analysis 1.10. Comparison 1 Avoidance vs use of NMBA, Outcome 10 One or more events of upper airway discomfort or injury: low risk of bias vs high or uncertain risk of bias.	97
Analysis 1.11. Comparison 1 Avoidance vs use of NMBA, Outcome 11 One or more events of upper airway discomfort or injury: depolarizing vs non-depolarizing NMBA.	98
Analysis 1.12. Comparison 1 Avoidance vs use of NMBA, Outcome 12 One or more events of upper airway discomfort or injury: remifentanyl vs no remifentanyl.	99
Analysis 1.13. Comparison 1 Avoidance vs use of NMBA, Outcome 13 One or more events of upper airway discomfort or injury: alfentanil vs no alfentanil.	99
Analysis 1.14. Comparison 1 Avoidance vs use of NMBA, Outcome 14 One or more events of upper airway discomfort or injury: excluded anticipated DTI vs included anticipated DTI.	100
Analysis 1.15. Comparison 1 Avoidance vs use of NMBA, Outcome 15 Difficult laryngoscopy: low risk of bias vs high or uncertain risk of bias.	100
ADDITIONAL TABLES	102
APPENDICES	115
CONTRIBUTIONS OF AUTHORS	124
DECLARATIONS OF INTEREST	125
DIFFERENCES BETWEEN PROTOCOL AND REVIEW	125
INDEX TERMS	126

[Intervention Review]

Avoidance versus use of neuromuscular blocking agents for improving conditions during tracheal intubation or direct laryngoscopy in adults and adolescents

Lars H Lundstrøm¹, Christophe HV Duez², Anders K Nørskov¹, Charlotte V Rosenstock¹, Jakob L Thomsen³, Ann Merete Møller⁴, Søren Strande⁵, Jørn Wetterslev⁶

¹Department of Anaesthesiology, Nordsjællands Hospital, Hillerød, Denmark. ²Research Center for Emergency Medicine, University of Aarhus, Aarhus, Denmark. ³Department of Anaesthesiology, Herlev Hospital, University of Copenhagen, Herlev, Denmark. ⁴The Cochrane Anaesthesia, Critical and Emergency Care Group, Herlev and Gentofte Hospital, University of Copenhagen, Herlev, Denmark. ⁵Department of Anaesthesiology and Intensive Care, Gentofte Hospital, Copenhagen, Denmark. ⁶Copenhagen Trial Unit, Centre for Clinical Intervention Research, Department 7812, Rigshospitalet, Copenhagen University Hospital, Copenhagen, Denmark

Contact address: Lars H Lundstrøm, Department of Anaesthesiology, Nordsjællands Hospital, Hillerød, 3400, Denmark.
lars_hyldborg@hotmail.com.

Editorial group: Cochrane Anaesthesia Group.

Publication status and date: New, published in Issue 5, 2017.

Citation: Lundstrøm LH, Duez CHV, Nørskov AK, Rosenstock CV, Thomsen JL, Møller AM, Strande S, Wetterslev J. Avoidance versus use of neuromuscular blocking agents for improving conditions during tracheal intubation or direct laryngoscopy in adults and adolescents. *Cochrane Database of Systematic Reviews* 2017, Issue 5. Art. No.: CD009237. DOI: [10.1002/14651858.CD009237.pub2](https://doi.org/10.1002/14651858.CD009237.pub2).

Copyright © 2017 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

Tracheal intubation during induction of general anaesthesia is a vital procedure performed to secure a patient's airway. Several studies have identified difficult tracheal intubation (DTI) or failed tracheal intubation as one of the major contributors to anaesthesia-related mortality and morbidity. Use of neuromuscular blocking agents (NMBA) to facilitate tracheal intubation is a widely accepted practice. However, because of adverse effects, NMBA may be undesirable. Cohort studies have indicated that avoiding NMBA is an independent risk factor for difficult and failed tracheal intubation. However, no systematic review of randomized trials has evaluated conditions for tracheal intubation, possible adverse effects, and postoperative discomfort.

Objectives

To evaluate the effects of avoiding neuromuscular blocking agents (NMBA) versus using NMBA on difficult tracheal intubation (DTI) for adults and adolescents allocated to tracheal intubation with direct laryngoscopy. To look at various outcomes, conduct subgroup and sensitivity analyses, examine the role of bias, and apply trial sequential analysis (TSA) to examine the level of available evidence for this intervention.

Search methods

We searched CENTRAL, MEDLINE, Embase, BIOSIS, International Web of Science, LILACS, advanced Google, CINAHL, and the following trial registries: [Current Controlled Trials](#); ClinicalTrials.gov; and www.centerwatch.com, up to January 2017. We checked the reference lists of included trials and reviews to look for unidentified trials.

Selection criteria

We included randomized controlled trials (RCTs) that compared the effects of avoiding versus using NMBA in participants 14 years of age or older.

Data collection and analysis

Two review authors extracted data independently. We conducted random-effects and fixed-effect meta-analyses and calculated risk ratios (RRs) and their 95% confidence intervals (CIs). We used published data and data obtained by contacting trial authors. To minimize the risk of systematic error, we assessed the risk of bias of included trials. To reduce the risk of random errors caused by sparse data and repetitive updating of cumulative meta-analyses, we applied TSA.

Main results

We identified 34 RCTs with 3565 participants that met our inclusion criteria. All trials reported on conditions for tracheal intubation; seven trials with 846 participants described 'events of upper airway discomfort or injury', and 13 trials with 1308 participants reported on direct laryngoscopy. All trials used a parallel design. We identified 18 dose-finding studies that included more interventions or control groups or both. All trials except three included only American Society of Anesthesiologists (ASA) class I and II participants, 25 trials excluded participants with anticipated DTI, and obesity or overweight was an excluding factor in 13 studies. Eighteen trials used suxamethonium, and 18 trials used non-depolarizing NMBA.

Trials with an overall low risk of bias reported significantly increased risk of DTI with no use of NMBA (random-effects model) (RR 13.27, 95% CI 8.19 to 21.49; $P < 0.00001$; 508 participants; four trials; number needed to treat for an additional harmful outcome (NNTH) = 1.9, $I^2 = 0\%$, $D^2 = 0\%$, GRADE = moderate). The TSA-adjusted CI for the RR was 1.85 to 95.04. Inclusion of all trials resulted in confirmation of results and of significantly increased risk of DTI when an NMBA was avoided (random-effects model) (RR 5.00, 95% CI 3.49 to 7.15; $P < 0.00001$; 3565 participants; 34 trials; NNTH = 6.3, $I^2 = 70\%$, $D^2 = 82\%$, GRADE = low). Again the cumulative z-curve crossed the TSA monitoring boundary, demonstrating harmful effects of avoiding NMBA on the proportion of DTI with minimal risk of random error. We categorized only one trial reporting on upper airway discomfort or injury as having overall low risk of bias. Inclusion of all trials revealed significant risk of upper airway discomfort or injury when an NMBA was avoided (random-effects model) (RR 1.37, 95% CI 1.09 to 1.74; $P = 0.008$; 846 participants; seven trials; NNTH = 9.1, $I^2 = 13\%$, GRADE = moderate). The TSA-adjusted CI for the RR was 1.00 to 1.85. None of these trials reported mortality. In terms of our secondary outcome 'difficult laryngoscopy', we categorized only one trial as having overall low risk of bias. All trials avoiding NMBA were significantly associated with difficult laryngoscopy (random-effects model) (RR 2.54, 95% CI 1.53 to 4.21; $P = 0.0003$; 1308 participants; 13 trials; NNTH = 25.6, $I^2 = 0\%$, $D^2 = 0\%$, GRADE = low); however, TSA showed that only 6% of the information size required to detect or reject a 20% relative risk reduction (RRR) was accrued, and the trial sequential monitoring boundary was not crossed.

Authors' conclusions

This review supports that use of an NMBA may create the best conditions for tracheal intubation and may reduce the risk of upper airway discomfort or injury following tracheal intubation. Study results were characterized by indirectness, heterogeneity, and high or uncertain risk of bias concerning our primary outcome describing difficult tracheal intubation. Therefore, we categorized the GRADE classification of quality of evidence as moderate to low. In light of defined outcomes of individual included trials, our primary outcomes may not reflect a situation that many clinicians consider to be an actual difficult tracheal intubation by which the patient's life or health may be threatened.

PLAIN LANGUAGE SUMMARY

The effect of avoiding neuromuscular blocking agents on conditions for placing a tube in the windpipe of patients undergoing general anaesthesia

Background

General anaesthesia abolishes spontaneous respiration. Use of general anaesthesia is frequently unavoidable during surgical procedures. The ability to maintain breathing by placing a tube in the windpipe of patients undergoing general anaesthesia is therefore crucial. A neuromuscular blocking agent (NMBA) is used for relaxation of muscles of the throat and is traditionally used to ease correct placement of the tube. However, use of an NMBA may cause unwanted side effects. On the other hand, large observational studies have indicated that avoiding NMBA may cause difficulties when the tube is placed during anaesthesia.

Objective

In the present systematic review, we assessed the effect of avoiding NMBA instead of using NMBA on difficulties associated with placing a tube in the windpipe of patients undergoing general anaesthesia. Further, we evaluated the consequences of using or avoiding NMBA on events of pain or injury in the upper part of the throat following placement of a tube in the windpipe.

Study characteristics

We identified 34 randomized controlled trials that met our inclusion criteria. These trials included 3565 patients who were undergoing various surgical procedures in hospital departments. Most trials were conducted in high-income countries, and most patients were undergoing elective surgery. Trials included patients of both sexes; most were healthy and non-obese, and staff members did not expect difficulty when placing the tube in the windpipe.

Key results

Avoidance versus use of neuromuscular blocking agents for improving conditions during tracheal intubation or direct laryngoscopy in adults and adolescents (Review)

Copyright © 2017 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

This review supports that use of NMBA may ensure the best conditions for placing a tube in the windpipe during general anaesthesia. When an NMBA is avoided, risk for pain or injury in the throat is increased following placement of a tube in the windpipe.

Quality of the evidence

Conditions for which a tube is placed in the windpipe are defined in individual trials and may not reflect a situation that many clinicians would consider to be clinically serious. Regarding events of injury and sore throat, only sparse data are available from trials with low risk of bias, although among all included trials, avoiding NMBA increased the risk of pain or injury. We therefore consider our overall findings to reflect evidence of moderate quality.

SUMMARY OF FINDINGS

Summary of findings for the main comparison. Summary of findings table: primary and secondary outcomes

Avoidance vs use of neuromuscular blocking agent for improving conditions during tracheal intubation in adults and adolescents

Patient or population: improving conditions during tracheal intubation or direct laryngoscopy in adults and adolescents

Setting: people undergoing various surgical procedures in hospital departments. Most trials were conducted in high-income countries, and most participants were undergoing elective surgery. Participants of both genders were included; most were ASA class I or II, were non-obese, and had no expected airway management difficulties

Intervention: avoidance of NMBA

Comparison: use of NMBA

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI) of avoidance vs use of NMBA	Number of participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Risk with use of NMBA	Risk with avoidance of NMBA				
Primary outcomes						
Difficult tracheal intubation: low risk of bias trials	Study population		RR 13.27 (8.19 to 21.49)	508 (4 studies)	⊕⊕⊕⊖ Moderate ^a	TSA shows that the required information size of 8195 for a 20% RRR has not been achieved, but the trial sequential monitoring boundary has been crossed and the TSA-adjusted CI for the RR is 1.85 to 95.04
	47 per 1000	620 per 1000 (383 to 1000)				
Difficult tracheal intubation: all trials	Study population		RR 5.00 (3.49 to 7.15)	3565 (34 studies)	⊕⊕⊖⊖ Low ^b	TSA shows that the required information size of 44,661 for a 20% RRR has not been achieved, but the trial sequential monitoring boundary has been crossed and the TSA-adjusted CI for the RR is 1.20 to 20.77.
	81 per 1000	406 per 1000 (284 to 597)				
One or more events of upper airway discomfort or injury: low risk of bias trials	Study population		RR 2.74 (1.21 to 6.21)	73 (1 study)	See comments	Because only 1 low risk of bias trial was identified, no quality of evidence assessment was performed
	162 per 1000	444 per 1000 (196 to 1000)				
One or more events of upper airway discomfort or injury: all trials	Study population		RR 1.37 (1.09 to 1.74)	846 (7 studies)	⊕⊕⊕⊖ Moderate ^c	TSA shows that the required information size of 1981 for a 20% RRR has not been achieved, but the trial sequential monitoring boundary has been crossed and the TSA-adjusted CI for the RR is 1.00 to 1.86.
	273 per 1000	374 per 1000				

	(298 to 475)					
Mortality	Not estimated	Not estimated	Not estimated	0 (34 studies)	Not estimated	
Secondary outcomes						
Difficult laryngoscopy: low risk of bias trials	Study population		RR 4.00 (0.47 to 34.20)	78 (1 study)	See comments	Because only 1 low risk of bias trial was identified, no quality of evidence assessment was performed
	26 per 1000	103 per 1000 (12 to 877)				
Difficult laryngoscopy: all trials	Study population		RR 2.54 (1.53 to 4.21)	1308 (13 studies)	⊕⊕⊕⊖ Low ^d	TSA shows that the required information size of 22,911 for a 20% RRR was not achieved, and in no trials were sequential monitoring boundaries crossed. The TSA-adjusted CI for the RR is 0.27 to 21.75.
	33 per 1000	85 per 1000 (51 to 141)				

***The risk in the intervention group** (and its 95% confidence interval) is based on assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI)

CI = confidence interval; RR = risk ratio

GRADE Working Group grades of evidence

High quality: We are very confident that the true effect lies close to the estimate of effect

Moderate quality: We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of effect but may be substantially different

Low quality: Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of effect

Very low quality: We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect

^aDowngraded one level because of indirectness

^bDowngraded two levels because of indirectness, heterogeneity, and high or uncertain risk of bias

^cDowngraded one level because of high or uncertain risk of bias

^dDowngraded two levels because of imprecision and high or uncertain risk of bias

BACKGROUND

Patients undergoing general anaesthesia lose consciousness and the ability to control breathing and protect their airway. Tracheal intubation is considered a vital procedure that secures the airway and provides the possibility of continued oxygenation. Several studies have identified difficult or failed tracheal intubation as one of the major reasons for anaesthesia-related mortality and morbidity. Types of morbidity range from sore throat, hoarseness, vocal cord lesion, pharyngeal oedema, and pharyngeal necrosis (Domino 1999), to more severe damage such as rupture or perforation of the pharynx and the oesophagus, aspiration pneumonia, and brain and heart injuries caused by anoxia (Rosenstock 2001). These severe complications may even be fatal (Cooper 2008; Hove 2007; Peterson 2005; Rosenstock 2001). Several studies have focused on one or more patient-related risk factors associated with difficult intubation (Adnet 1997; el-Ganzouri 1996; L'Hermite 2009; Lundstrom 2009a; Shiga 2005). Successful tracheal intubation is determined by the anaesthetist's technical and non-technical skills, as well as by available facilities and the local environment (Rosenstock 2004; Rosenstock 2006).

Use of neuromuscular blocking agents (NMBA) to facilitate tracheal intubation is a widely accepted procedure. Owing to adverse effects, the use of NMBA may be undesirable. Cohort studies (Baillard 2005; Lundstrom 2009a; Lundstrom 2009b; Lundstrom 2009c) have demonstrated that avoidance of neuromuscular blocking drugs may be an independent risk factor for difficult and failed tracheal intubation. Randomized trials have evaluated the conditions for tracheal intubation, possible side effects, and postoperative discomfort by comparing different regimens of anaesthesia induction and avoidance versus use of NMBA (Alcock 1993; Alexander 1999; Bouvet 2008; Combes 2007; Harsten 1997; Lieutaud 2003; McNeil 2000; Mencke 2003; Naguib 2006; Scheller 1992; Stevens 1997).

Description of the condition

A successful tracheal intubation is considered to provide a safe airway. The tube is placed directly and is cuffed within the patient's trachea, thereby ensuring a direct connection to the lower airway. The risk of aspiration of gastric contents into the lungs may be reduced by an appropriately placed tracheal tube. Difficulties with tracheal intubation by direct laryngoscopy can cause serious soft tissue damage (Domino 1999) and are the principal causes of hypoxaemic anaesthetic death and brain damage (Henderson 2004). A review identified difficult airway management, including difficult or failed tracheal intubation, as the main cause of death and severe morbidity related to anaesthesia (Braz 2009). The literature presents various definitions of difficult tracheal intubation (DTI). Performance and graduation of tracheal intubation are described by various parameters such as the number of attempts made for intubation; the span of time needed for intubation; the experience and number of anaesthetists performing the intubation; the type and number of alternative techniques used other than direct laryngoscopy; the lifting force required during the laryngoscopy; whether laryngeal pressure was needed and information on vocal cord mobility (Adnet 1997; ASA guideline 1993; Lundstrom 2009a; Viby-Mogensen 1996).

Difficult laryngoscopy, which is often used as a surrogate outcome measure for a DTI, is traditionally defined by the (modified)

Cormack and Lehane classification (Cormack 1984; Yentis 1998) (Appendix 1).

Description of the intervention

Induction of general anaesthesia can include or exclude the use of a neuromuscular blocking agent. A combination of adjuvant drugs such as hypnotics, opioids, and occasionally local anaesthetic agents is traditionally used for the induction.

The NMBA used for induction may be a depolarizing or a non-depolarizing drug. Hypnotics may include propofol, thiopental, or etomidate, and opioids may consist of fentanyl, alfentanil, or remifentanyl. A local anaesthetic such as lidocaine (Tanaka 2015) may be used intravenously or topically to facilitate tracheal intubation (Woods 2005). After induction has been completed, the patient will undergo tracheal intubation or attempted tracheal intubation with a standard direct laryngoscope.

How the intervention might work

Neuromuscular blocking agents obstruct the nerve impulse at the neuromuscular junction propagated by acetylcholine, causing paralysis of the skeletal muscles. Direct laryngoscopy normally stimulates the oropharynx and activates oropharyngeal reflexes. Use of an NMBA inhibits muscular contractions and improves the conditions for tracheal intubation (Bowman 2006). Both depolarizing and non-depolarizing NMBA may produce side effects, such as anaphylaxis, cardiovascular effects related to histamine release or direct vagolytic or sympathomimetic properties, bronchospasm, and prolonged paralysis. Depolarizing NMBA may specifically cause muscle pain, increased serum potassium, malignant hyperthermia, and increased intraocular pressure (Appiah-Ankam 2004).

Why it is important to do this review

Difficult airway management, including a difficult or failed tracheal intubation, remains a major cause of death and severe morbidity related to anaesthesia (Braz 2009). The risk of DTI may be reduced by the choice of an induction strategy including, or avoiding, NMBA for facilitating tracheal intubation by direct laryngoscopy. Use of NMBA may be associated with serious adverse events.

OBJECTIVES

To evaluate the effects of avoiding neuromuscular blocking agents (NMBA) versus using NMBA on difficult tracheal intubation (DTI) for adults and adolescents allocated to tracheal intubation with direct laryngoscopy. To look at various outcomes, conduct subgroup and sensitivity analyses, examine the role of bias, and apply trial sequential analysis (TSA) (Brok 2008; Brok 2009; Thorlund 2009; Wetterslev 2008) to examine the level of available evidence for this intervention.

METHODS

Criteria for considering studies for this review

Types of studies

We included randomized controlled trials (RCTs).

We included unpublished trials only if trial data and methodological descriptions were provided in written form or through direct contact with study authors.

We excluded trials using quasi-randomization as well as observational studies.

Types of participants

We included participants 14 years of age or older who underwent (attempt of) tracheal intubation by direct laryngoscopy regardless of acute or elective surgical status or investigational procedures.

Types of interventions

Avoidance of neuromuscular blocking agents (= intervention) versus use of neuromuscular blocking agents (= control) to facilitate tracheal intubation. We defined use of NMBA as the control intervention, as it is considered usual standard for airway management, and we defined avoidance of NMBA as the experimental intervention.

Types of outcome measures

Primary outcomes

1. Difficult tracheal intubation: As no international consensus has been reached on defining an intubation score, we accepted the definitions of DTI presented in the individual articles. If study authors defined a difficult laryngoscopy by the Cormack and Lehane score ([Cormack 1984](#)) or by the modified Cormack and Lehane score ([Yentis 1998](#)) as a difficult intubation, we included and reported the Cormack and Lehane score as the outcome measure. Difficult laryngoscopy is a surrogate outcome for a DTI. Therefore, if a trial reported both an intubation score and the Cormack and Lehane score based on the same population in the same assessment, we extracted only the intubation score for outcome assessment
2. Overall mortality: We used maximal follow-up data from each trial
3. One or more events of upper airway discomfort or injury (e.g. sore throat, hoarseness, vocal cord lesion, minor pharyngeal injury)

Secondary outcomes

1. One or more major serious events: pulmonary aspiration, brain and heart injuries (e.g. caused by anoxia, hypotension, bradycardia or tachycardia during tracheal intubation)
2. Difficult laryngoscopy, as defined by the Cormack and Lehane score ([Cormack 1984](#)) or the modified Cormack and Lehane score ([Yentis 1998](#))

Search methods for identification of studies

Electronic searches

We searched the Cochrane Central Register of Controlled Trials (CENTRAL; 2017, Issue 1) (the Cochrane Library); MEDLINE (Ovid) (1950 to January 2017); Embase (Ovid) (1980 to January 2017); BIOSIS (Ovid) (1993 to January 2017); International Web of Science (1964 to January 2017); Latin American Caribbean Health Sciences Literature (LILACS) via BIREME (1982 to January 2017); the Chinese Biomedical Literature Database; advanced Google; and the Cumulative Index to Nursing and Allied Health Literature (CINAHL) via EBSCO host (1980 to January 2017).

We utilized a systematic and sensitive search strategy to identify relevant RCTs with no language or date restrictions. We conducted the search within six months of the date the draft review was emailed to the editorial office. For specific information regarding our search strategies, please see the Appendices ([Appendix 2](#), CENTRAL; [Appendix 3](#), MEDLINE; [Appendix 4](#), Embase; [Appendix 5](#), BIOSIS; [Appendix 6](#), CINAHL; [Appendix 7](#), LILACS).

Searching other resources

We searched for ongoing clinical trials and unpublished studies on the following Internet sites (11 January 2017).

1. [Current Controlled Trials](#).
2. [ClinicalTrials.gov](#).
3. [www.centerwatch.com](#).

We handsearched the reference lists of reviews, randomized and non-randomized studies, and editorials for additional studies.

Data collection and analysis

Selection of studies

In the process of selecting trials for inclusion in the review, two review authors (LHL and AN or CD) independently screened titles and abstracts to identify eligible trials and remove obviously irrelevant reports. After retrieving the full texts of potentially eligible reports, the review authors (LHL and one of the following: AN, CD, CVR, JT) examined the full-text reports. We contacted the main authors of studies and experts in this field and asked about missed, unreported, and ongoing trials. Finally, two review authors (LHL and one of the following: AN, CD, CVR, JT) decided which trials would be included and proceeded to data collection. We resolved disagreements by discussion, and a third review author (JW) resolved residual disagreements.

Data extraction and management

LHL and AN, CD, CVR, or JT independently extracted and collected data on a standardized paper form ([Appendix 8](#)). We were not blinded to study author, institution, or source of trial publication. We resolved disagreements by discussion, and a third review author (JW) resolved residual disagreements. If necessary, we approached all corresponding authors of included trials for additional information on the review's outcome measures and risk of bias components. For more information, please see the section titled [Contributions of authors](#).

Investigators in some trials randomized participants to multiple intervention and control groups or both (more than two groups, as in dose-finding studies). We combined all relevant experimental intervention groups into a single intervention group and combined all relevant control intervention groups into a single control group ([Higgins 2011](#)).

Assessment of risk of bias in included studies

We evaluated the validity and design characteristics of each trial. To draw conclusions about the overall risk of bias for an outcome, we found it necessary to evaluate trials for major sources of bias, also defined as domains (random sequential generation, allocation concealment, blinding, incomplete outcome data, selective reporting, and other bias). Assessment of overall

risk of bias involved consideration of the relative importance of the different domains (Higgins 2011).

Even the most realistic assessment of the validity of a trial may involve subjectivity because it was impossible to know the extent of bias (or even the true risk of bias) of a given trial. Some domains affected risk of bias across trial outcomes (e.g. sequential generation, allocation sequential concealment), and others, such as blinding and incomplete outcome data, may have had different risks of bias for different outcomes within a trial. Thus, risk of bias was not the same for all outcomes in a trial. We performed separate sensitivity analyses for patient-reported outcomes (subjective outcomes; e.g. upper airway discomfort) and for mortality (Higgins 2011).

We defined trials as having low risk of bias only if they adequately fulfilled the criteria listed in the *Cochrane Handbook for Systematic Reviews of Interventions*, and we performed summary assessments of risk of bias for each important outcome (across domains) within and across studies. We applied a 'risk of bias graph' and a 'risk of bias summary figure' (Higgins 2011).

We presented results for all outcomes including adverse events in a 'Summary of findings' table (Higgins 2011).

As no sufficiently well-designed formal statistical method is available to combine the results of trials with high and low risk of bias, Cochrane review authors usually incorporate risk of bias assessments by comparing meta-analyses of trials with low risk and high or uncertain risk of bias (Higgins 2011). We used the risk of bias table described in the *Cochrane Handbook for Systematic Reviews of Interventions* (Section 8.5) (Higgins 2011) as a tool for assessing risk of bias in included studies. We assessed risk of bias for different domains as described below.

Random sequence generation

Low risk of bias: The method used generates random sequences (e.g. random number generation, toss of coin).

Unclear: No information on random sequence generation is available.

High risk of bias: Alternate medical record numbers or other non-random sequence generation was used.

Allocation concealment

Low risk of bias: Allocation method prevents investigators or participants from knowing the next allocation (e.g. central allocation; sealed opaque envelopes; serially numbered, sequentially numbered but otherwise identical vehicles, including their contents; other descriptions of convincing concealment of allocation).

Unclear: No information on allocation method is available, or the description provided did not allow a clear distinction.

High risk of bias: Allocation method allowed investigators or participants to know the next allocation (e.g. alternate medical record numbers; reference to case record numbers or date of birth; open allocation sequence, unsealed envelopes).

Blinding

Low risk of bias: We considered blinding as adequate if participants and personnel were kept unaware of intervention allocations after participants were enrolled into the study, and if the method of blinding involved placebo.

Unclear: Blinding was not described.

High risk of bias: Study was not double-blinded; was categorized as an open-label study; or was conducted without use of placebo.

Incomplete outcome data

Low risk of bias: Numbers and reasons for dropouts and withdrawals in the intervention groups were described, or it was specified that no dropouts or withdrawals occurred.

Unclear: The report gave the impression that no dropouts or withdrawals had occurred, but this was not specifically stated.

High risk of bias: Numbers and reasons for dropouts and withdrawals were not described.

Selective reporting

Low risk of bias: Report includes predefined or clinically relevant and reasonably expected outcomes.

Unclear: Study did not report or did not report fully all predefined or clinically relevant and reasonably expected outcomes, or it is unclear whether data on these outcomes were recorded.

High risk of bias: Report did not include one or more clinically relevant and reasonably expected outcomes; data on these outcomes were likely to have been recorded.

Baseline imbalance

Low risk of bias: We noted no baseline imbalance in important characteristics.

Unclear: Baseline characteristics were not reported.

High risk of bias: Baseline imbalance was due to chance or to imbalanced exclusion after randomization.

Measures of treatment effect

Most often, a dichotomous outcome measure is used to assess whether an intubation is difficult or not. We reported all dichotomous outcomes as risk ratios (RR) with 95% confidence intervals (CI). For mortality, which we expected to be a rare outcome, we calculated the Peto odds ratio.

Unit of analysis issues

We planned to include studies with a non-standard design, such as cluster-randomized trials and studies with more than two intervention groups. We considered 'cross-over trials', 'repeated observations on participants', and 'multiple treatment attempts' as unlikely designs for evaluating the current intervention. We included all studies with a non-standard design and analysed them as described in the *Cochrane Handbook for Systematic Reviews of Interventions* (Higgins 2011).

Dealing with missing data

We contacted all first authors and contact persons of trials with missing data to retrieve relevant data. We performed a modified intention-to-treat (ITT) analysis while including, if possible, all randomized participants who underwent tracheal intubation or did not withdraw their consent before surgery.

ITT analysis is recommended to minimize bias in design, follow-up, and analysis of the efficacy of RCTs. It yields a pragmatic estimate of the benefit of a change in treatment policy rather than of potential benefit for patients who received treatment exactly as planned (Hollis 1999). Full application of ITT is possible only when complete outcome data are available for all randomized participants. Although about half of all published reports of RCTs state that ITT was used, handling of deviations from randomized allocation varies widely, and many trials have missing data for the primary outcome variable; methods used to deal with this are generally inadequate, potentially leading to bias (Hollis 1999).

Performing an ITT analysis in a systematic review is not straightforward in practice, in that review authors must decide how to handle missing outcome data from contributing trials (Gamble 2005). No consensus exists about how missing data should be handled in ITT analyses, and different approaches may be appropriate in different situations (Higgins 2011; Hollis 1999).

In the case of missing data, we used a 'complete-case analysis' for our primary outcomes, which simply excluded all participants for whom outcome data were missing from the analysis. Additionally, we performed sensitivity analyses covering best- and worst-case scenarios (with 'best' and 'worst' defined with respect to effect on the chosen outcome with use of NMBA).

Assessment of heterogeneity

We quantified the degree of heterogeneity observed in the results using diversity (D^2) (Wetterslev 2009) and inconsistency factor (I^2) statistics, which we interpreted as the proportion of total variation observed between trials that was attributable to differences between trials rather than to sampling error (chance) (Higgins 2002). A finding of $P \leq 0.10$ indicated significant heterogeneity, and the I^2 statistic has suggested thresholds for low (25% to 49%), moderate (50% to 74%), and high ($\geq 75\%$) heterogeneity (Higgins 2003). If $I^2 = 0\%$, we reported only results from the fixed-effect model. In the case of $I^2 > 0\%$, we reported results from both random-effects and fixed-effect models. However, we believe that use of a fixed-effect model provided little value in cases of substantial heterogeneity, which we suspected would be the case in this review owing to inclusion of various patient types, adjuvant medicines, and outcome reporting. So we emphasized results from the random-effects model unless a few trials dominated the meta-analysis (e.g. $> 50\%$ of the accumulated fixed weight percentage). In cases of $I^2 > 0\%$ (for mortality and difficult intubation outcomes), we explored possible causes of heterogeneity by performing meta-regression analyses and relevant subgroup and sensitivity analyses. We aimed to meta-analyse trial results only in cases of low to moderate clinical heterogeneity.

Assessment of reporting biases

Publication bias occurs when publication of research results depends on their nature and direction (Dickersin 1990). We examined this by creating funnel plots to detect publication bias or

a difference between small and large studies ('small study effects') expressed by asymmetry (Egger 1997). In cases of asymmetry, we applied the Arcsine-Thompson test, as proposed by Rücker (Rücker 2008). We defined funding bias as bias in the design, outcome, and reporting of industry-sponsored research to show that a drug has a favourable outcome (Bekelman 2003). Relationships between industry, scientific investigators, and academic institutions are widespread and often result in conflicts of interest (Bekelman 2003). We conducted a sensitivity analysis to examine the role of funding bias.

Data synthesis

We used Review Manager Software (RevMan 5.3). We calculated the RR with 95% CIs for dichotomous variables (binary outcomes) as well as the risk difference (Keus 2009), but if results were similar, we reported only the RR. We used D^2 (Wetterslev 2009) and the I^2 statistic (Higgins 2002) to describe heterogeneity among included trials. We planned to explore causes of substantial heterogeneity by performing meta-regression using Comprehensive Meta-Analysis (CMA), version one, and Stata, version nine. We used the χ^2 test to provide an indication of heterogeneity between studies, with $P \leq 0.10$ considered significant.

Adverse effects may be rare but serious and hence important (Sutton 2002) when meta-analysis is applied in combining results from several trials with binary outcomes (i.e. event or no event). Most meta-analytical software packages do not include options for analyses that include trials with 'zero event' in both arms (intervention vs control) for calculating RR. Exempting these trials from calculation of RR and CI may lead to overestimation of a treatment effect, as the control event proportion may be overestimated. Thus we performed a sensitivity analysis by applying empirical continuity corrections to our zero event trials, as proposed by Sweeting et al (Keus 2009; Sweeting 2004), and by applying imaginary small numbers in both arms.

Meta-analyses may result in type I errors due to systematic errors (bias) or may produce random errors due to repeated significance testing when meta-analyses are updated with new trials (Brok 2008; Brok 2009; Thorlund 2009; Wetterslev 2008; Wetterslev 2009). Bias from trials with high risk of bias, outcome reporting bias, publication bias, early stopping for benefit, and small trial bias may result in spurious P values.

In a single trial, interim analysis increases the risk of type I errors. To avoid type I errors, we applied group sequential monitoring boundaries (Lan 1983) to decide whether a trial could be terminated early because of a sufficiently small P value, that is, when the cumulative z-curve crosses the monitoring boundary. Sequential monitoring boundaries can be applied to meta-analyses; these are called trial sequential monitoring boundaries. In trial sequential analysis (TSA), the addition of each trial to a cumulative meta-analysis is regarded as an interim meta-analysis and helps to show whether additional trials are needed (Wetterslev 2008). So far, several meta-analyses and reviews have been published, and these have included increasing trial results as new trials have been published (Al-Niaimi 2009; Chura 2007; Qadan 2009). Therefore, it seems appropriate to adjust new meta-analyses for multiple testing on accumulating data to control the overall type 1 error risk in cumulative meta-analysis (Pogue 1997; Pogue 1998; Thorlund 2009; Wetterslev 2008).

The idea in TSA is that if the cumulative z-curve crosses a boundary, a sufficient level of evidence is reached and no further trials may be needed. However, evidence is insufficient to permit a conclusion if the z-curve does not cross a boundary or does not surpass the required information size. To construct trial sequential monitoring boundaries (TSMB), the required information size is needed and will be calculated as the least number of participants needed to conduct a well-powered single trial (Brok 2008; Pogue 1998; Wetterslev 2008). We adjusted the required information size for heterogeneity by applying a D² adjustment factor (Wetterslev 2009). We will apply TSA because it prevents an increase in the risk of type I error (< 5%) caused by potential multiple updating and testing on accumulating data whenever new trial results are included in a cumulative meta-analysis (Pogue 1997; Pogue 1998) and provides important information needed to estimate the level of evidence for the experimental intervention (Pogue 1997; Pogue 1998; Thorlund 2009). Additionally, TSA provides important information regarding the need for additional trials and required information size (Wetterslev 2008; Wetterslev 2009). We applied TSMB according to an information size suggested by trials with low risk of bias (Wetterslev 2008; Wetterslev 2009), an *a priori* 20% relative risk reduction (RRR) of difficult or failed intubation, and an intervention effect suggested by the 95% confidence limit closest to 1 in the traditional random-effects meta-analysis. As mortality seems low or even absent in the trials conducted so far, and hence the ability to detect small intervention effects is low, we also planned to perform a TSA with an information size estimated on the basis of an *a priori* 50% RRR of mortality (Wetterslev 2008; Wetterslev 2009).

Subgroup analysis and investigation of heterogeneity

We performed the following subgroup analyses of our primary outcomes.

1. Avoidance of NMBA versus use of NMBA (comparisons of subgroups of depolarizing vs non-depolarizing NMBA).
2. Avoidance of NMBA versus use of NMBA in combination with or without remifentanyl or alfentanil.
3. Avoidance of NMBA versus use of NMBA in combination with or without local anaesthetic drug.
4. Avoidance of NMBA versus use of NMBA (comparisons of subgroups of trials using anticipation of a difficult airway as an inclusion criterion vs those not using anticipation of a difficult airway as an inclusion criterion).

When analyses of various subgroups with a binary outcome were significant, we performed a test of interaction (Altman 2003). We considered $P < 0.05$ to be indicative of significant interaction between the effect of no use of NMBA on DTI and the subgroup category (Higgins 2011; Chapters 9.6.1 and 9.7).

We planned to explore causes of moderate to high heterogeneity using meta-regression including the covariates listed below. We ranked these covariates according to their importance and included them in the meta-regression according to the number of relevant trials included in this review.

1. Mean age of trial population at baseline.
2. Fraction of gender of trial population at baseline.
3. Fraction of a Mallampati score grade I to IV.

4. Thyromental distance (dichotomous or continuous measurement).
5. Mouth opening or interincisor gap (dichotomous or continuous measurement).
6. Neck extension (dichotomous or continuous measurement).
7. Mandible subluxation (dichotomous or continuous measurement).
8. Mean body mass index (BMI) of trial population at baseline.
9. Time from induction to start of intubation (seconds).

Sensitivity analysis

We performed a sensitivity analysis to evaluate the impact of trials with high or uncertain risk of bias versus trials with low risk of bias.

'Summary of findings' tables

We used the principles of the GRADE system (Guyatt 2013; Guyatt 2011) to assess the quality of the body of evidence associated with specific outcomes.

1. DTI.
2. All-cause mortality (maximal follow-up data from each trial).
3. One or more events of upper airway discomfort or injury (e.g. sore throat, hoarseness, vocal cord lesion, minor pharyngeal injury).
4. One or more major serious events (defined by pulmonary aspiration, brain and heart injuries (caused by e.g. anoxia, hypotension, bradycardia, or tachycardia during tracheal intubation)).
5. Difficult laryngoscopy, defined by the Cormack and Lehane score (Cormack 1984) or the modified Cormack Lehane score (Yentis 1998).

The GRADE approach appraises the quality of a body of evidence according to the extent to which one can be confident that an estimate of effect or association reflects the item being assessed. Quality considers:

1. within-study risk of bias (methodological quality);
2. directness of the evidence;
3. heterogeneity of the data;
4. precision of effect estimates; and
5. risk of publication bias.

RESULTS

Description of studies

See [Characteristics of included studies](#) and [Characteristics of excluded studies](#).

Results of the search

We identified 4400 references through electronic and manual searches (Figure 1). We updated the search on 11 January 2017. After excluding duplicate reports, we screened the abstracts of 2822 references. We obtained 37 publications for full-text review. Thirty-four publications were RCTs and met our inclusion criteria for the comparison of avoiding NMBA or using NMBA to facilitate tracheal intubation.

Figure 1. Study flow diagram.

Included studies

Most of the trials identified for this comparison were published in English. Two trials were published in French (Dominici 1990; Rousseau 1995), one in German (Striebel 1995), and one in Spanish (González Obregón 2010). Included trials enrolled 3565 participants, 1507 of whom were randomized to no use of NMBA, and 2058 to use of NMBA. Table 1 shows characteristics of the 34 included trials, and Table 2 shows characteristics of the interventions. The number of randomized participants included in individual trials ranged from 40 to 300.

Trial design

All 34 studies were RCTs using parallel design. Eighteen dose-finding studies comprised more intervention or control groups, or both (Alexander 1999; Barclay 1997; Kahwaji 1997; Kirkegaard-Nielsen 1999; Kopman 2001; Lieutaud 2003; Lowry 1999; McNeil 2000; Naguib 2003; Naguib 2006; Nimmo 1995; Pino 1998; Scheller 1992; Schlaich 2000; Sivalingam 2001; Stevens 1997; Striebel 1995; Wong 1996). A total of 16 trials included one intervention group and one control group (Beck 1993; Bouvet 2008; Combes 2007; Dominici 1990; González Obregón 2010; Gulhas 2013; Hanna 2010; Harsten 1997; Iamaroon 2001; Isesele 2012; Jiao 2014; Mencke 2003; Mencke 2014; Pang 2014; Rousseau 1995; Yazdi 2016).

Trial participants

All included trials except one (Barclay 1997) enrolled only American Society of Anesthesiologists (ASA) class I and III participants. A total of 25 trials (Alexander 1999; Barclay 1997; Bouvet 2008; Combes 2007; González Obregón 2010; Hanna 2010; Iamaroon 2001; Isesele 2012; Kahwaji 1997; Lieutaud 2003; Lowry 1999; McNeil 2000; Mencke 2003; Mencke 2014; Naguib 2003; Naguib 2006; Pang 2014; Pino 1998; Rousseau 1995; Scheller 1992; Schlaich 2000; Sivalingam 2001; Stevens 1997; Wong 1996; Yazdi 2016) excluded participants with anticipated DTI, and 13 trials excluded obese participants and overweight participants (Barclay 1997; Combes 2007; Iamaroon 2001; Jiao 2014; Kirkegaard-Nielsen 1999; Kopman 2001; Lowry 1999; McNeil 2000; Mencke 2003; Mencke 2014; Pang 2014; Pino 1998; Stevens 1997). Definitions of 'obese' and 'overweight' varied among trials. We accepted the definitions presented by study authors in the individual articles.

Characteristics of interventions

Investigators in 18 trials used suxamethonium as the depolarization NMBA (Alexander 1999; Beck 1993; Dominici 1990; Gulhas 2013; Hanna 2010; Harsten 1997; Iamaroon 2001; Isesele 2012; Jiao 2014; McNeil 2000; Naguib 2003; Naguib 2006; Nimmo 1995; Scheller 1992; Sivalingam 2001; Stevens 1997; Striebel 1995; Wong 1996), and researchers in 18 trials used one or more non-depolarizing NMBA (Barclay 1997; Bouvet 2008; Combes 2007; González Obregón 2010; Hanna 2010; Kahwaji 1997; Kirkegaard-Nielsen 1999; Kopman 2001; Lieutaud 2003; Lowry 1999; Mencke 2003; Mencke 2014; Pang 2014; Pino 1998; Rousseau 1995;

Schlaich 2000; Striebel 1995; Yazdi 2016). Thus, two trials (Hanna 2010; Striebel 1995) combined depolarizing and non-depolarizing NMBA as the intervention. Among trials using one or more non-depolarizing NMBA, 10 applied rocuronium (Barclay 1997; Combes 2007; González Obregón 2010; Hanna 2010; Kirkegaard-Nielsen 1999; Kopman 2001; Lowry 1999; Mencke 2014; Pino 1998; Schlaich 2000). Three trials applied atracurium (Lieutaud 2003; Mencke 2003; Yazdi 2016), two trials vecuronium (Rousseau 1995; Striebel 1995), and two trials rapacurium (Kahwaji 1997; Kopman 2001). Single trials applied both cisatracurium (Bouvet 2008) and mivacurium (Pino 1998), respectively.

Characteristics of outcome measures

In 16 trials, investigators described intubation conditions by the original (Alexander 1999; Bouvet 2008; Combes 2007; Hanna 2010; Jiao 2014; Kirkegaard-Nielsen 1999; Kopman 2001; Lowry 1999; Mencke 2003; Naguib 2003; Naguib 2006; Schlaich 2000) or a modified (Barclay 1997; Harsten 1997; Kahwaji 1997; Mencke 2014) version of "Good clinical research practice (GCRP) in pharmacodynamic studies of neuromuscular blocking agents" (Viby-Mogensen 1996). Thirteen trials reported laryngoscopy conditions defined by the criteria described by Cormack and Lehane (Cormack 1984) (Beck 1993; Bouvet 2008; Combes 2007; Dominici 1990; González Obregón 2010; Harsten 1997; McNeil 2000; Mencke 2003; Mencke 2014; Pang 2014; Scheller 1992; Stevens 1997; Striebel 1995). Seven studies reported events of upper airway discomfort or injury (Bouvet 2008; Combes 2007; González Obregón 2010; Gulhas 2013; Mencke 2003; Mencke 2014; Sivalingam 2001).

Excluded studies

We excluded three trials. One study intubated an unspecified number of participants blind nasal (Alcock 1993), and another (Ide 2015) intubated participants nasally using a Magill forceps. One trial terminated inclusion of participants exclusively in the intervention group because of unacceptable intubation conditions. Thus, this trial violated randomization and blinding as planned (Baumgarten 1988) (see [Characteristics of excluded studies](#) for more information).

Studies awaiting classification

No studies are awaiting classification.

Ongoing studies

We found no ongoing studies.

Risk of bias in included studies

We performed the 'Risk of bias' assessment of included trials using previously described criteria. For details of judgements made for individual trials, please see [Risk of bias in included studies](#) (Figure 2). When we could not judge a 'Risk of bias' domain as having low risk, we asked study authors for additional information.

Figure 2. Risk of bias graph: review authors' judgements about each risk of bias item presented as percentages across all included studies.

We judged four trials as having low risk of bias in all domains (Jiao 2014; Mencke 2003; Naguib 2003; Naguib 2006), and 13 trials as having high risk of bias in at least one bias domain. We divided trials into trials with overall low risk of bias and those with high or uncertain risk of bias according to assessment of sequence generation, concealment of allocation, blinding of outcome assessment, and blinding of participants and personnel according to the *Cochrane Handbook for Systematic Reviews of Interventions* 'Risk of bias' tool. When we judged all domains as adequately assessed, we considered the trial as having an overall low risk of bias.

Random sequence generation

Seventeen trials adequately described generation of the allocation sequence (Alexander 1999; Bouvet 2008; Dominici 1990; González Obregón 2010; Gulhas 2013; Harsten 1997; Jiao 2014; Kopman 2001; Lieutaud 2003; Lowry 1999; Mencke 2003; Mencke 2014; Naguib 2003; Naguib 2006; Pang 2014; Pino 1998; Rousseau 1995). The remaining trials were described as randomized, but investigators did not adequately describe the method used for sequence generation.

Allocation

Sixteen trials adequately described the method used to conceal allocation (Bouvet 2008; Combes 2007; González Obregón 2010; Hanna 2010; Harsten 1997; Isesele 2012; Jiao 2014; Kirkegaard-Nielsen 1999; Lieutaud 2003; McNeil 2000; Mencke 2003; Mencke 2014; Naguib 2003; Naguib 2006; Sivalingam 2001; Stevens 1997). We categorized one trial as having high risk of bias (Kopman 2001) and judged the method used for allocation concealment as unclear for the remaining 16 trials.

Blinding

Ten trials adequately described the method used to blind participants and personnel responsible for treatment (Combes 2007; Gulhas 2013; Jiao 2014; Mencke 2003; Naguib 2003; Naguib 2006; Nimmo 1995; Stevens 1997; Striebel 1995; Wong 1996). We

categorized ten trials as having high risk of bias (Bouvet 2008; González Obregón 2010; Harsten 1997; Kopman 2001; Lieutaud 2003; Lowry 1999; McNeil 2000; Mencke 2014; Rousseau 1995; Scheller 1992) and judged the method used to blind participants and personnel responsible for the treatment as unclear for the remaining trials.

All except five trials adequately described the method used to blind the person performing airway management and outcome assessment. We judged four trials as having unclear risk of bias (Isesele 2012; Nimmo 1995; Rousseau 1995; Yazdi 2016) and one trial as having high risk of bias (Scheller 1992).

Incomplete outcome data

All trials adequately addressed incomplete data. Most trials provided complete outcome data for all randomized participants. However, six trials (Bouvet 2008; Combes 2007; Hanna 2010; Isesele 2012; Kahwaji 1997; Pino 1998) excluded a few participants from the ITT because "patients were lost to follow-up", or because investigators encountered missing data or equipment failure or various clinical reasons for exclusion. In six trials (Harsten 1997; Jiao 2014; Mencke 2003; Mencke 2014; Nimmo 1995; Sivalingam 2001), study authors excluded a few participants from ITT if tracheal intubation failed. We have provided further details in the [Characteristics of included studies](#) table.

Selective reporting

All trials adequately addressed selective reporting bias, except one trial for which we judged risk as unclear (Nimmo 1995).

Other potential sources of bias

We did not assess other potential sources of bias.

Assessment of reporting bias

The funnel plots that we have provided for our primary outcomes did not express asymmetry and thereby did not indicate publication bias (Figure 3).

Figure 3. Funnel plot of comparison: 1 Avoidance vs use of NMBA, outcome: 1.1 Difficult tracheal intubation: low risk of bias vs high or uncertain risk of bias.

Effects of interventions

See: [Summary of findings for the main comparison](#) [Summary of findings table: primary and secondary outcomes](#)

See our main results on all investigated outcomes in the 'Summary of findings' table ([Summary of findings for the main comparison](#)), which presents assessment of the quality of the evidence, including imprecision, according to the GRADE approach ([Guyatt 2013](#); [Guyatt 2011](#)).

Primary outcome: difficult tracheal intubation

Difficult tracheal intubation in trials with low risk of bias (Analysis 1.1)

We categorized four trials ([Jiao 2014](#); [Mencke 2003](#); [Naguib 2003](#); [Naguib 2006](#)) with 508 participants as trials with low risk of bias. Among 144 participants undergoing tracheal intubation without the use of NMBA, investigators intubated a crude proportion of 56.3% ($n = 81$; 95% CI 49.3% to 64.4%) with difficulties. The median prevalence of a DTI in the individual studies was 58.2% (range 33% to 70%). Of 364 participants intubated with the use of NMBA, the proportion of DTI was 4.7% ($n = 17$; 95% CI 2.8% to 6.8%). The median prevalence of a DTI among individual studies was 4.7%

(range 3.7% to 5.4%). In a random-effects model, risk of DTI was increased with no use of NMBA (RR 13.27, 95% CI 8.19 to 21.49; $P < 0.00001$; number needed to treat for an additional harmful outcome (NNTH) = 1.9, $I^2 = 0\%$, $D^2 = 0\%$), likewise in a fixed-effect model (RR 12.50, 95% CI 7.62 to 20.52; $P < 0.00001$). Comparisons of subgroups in trials with low risk of bias and trials with high or uncertain risk of bias revealed significant subgroup differences ($P = 0.0002$). Among trials of high or uncertain risk, avoidance of NMBA was associated with DTI in a random-effects model (RR 4.23, 95% CI 2.93 to 6.09; $P < 0.00001$; NNTH = 8.7, $I^2 = 64\%$). We performed TSA of non-NMBA versus NMBA using a D^2 -adjusted required information size of 8195 participants to detect or reject a 20% relative risk increase with power of 80%, a control event proportion of 8%, and an overall type 1 error of 5%. The cumulative z-curve crossed the TSMB for harm, and the TSA-adjusted CI for the RR was 1.85 to 95.04. These findings provide firm evidence of the smallest possible harmful effect on the proportion of DTI in the traditional meta-analysis of avoiding NMBA compared with using NMBA, even when the significance level was adjusted for repetitive testing and sparse data in a cumulative meta-analysis. We downgraded the quality of the evidence (GRADE) one level to moderate because of indirectness ([Analysis 1.1](#); [Figure 4](#)).

Figure 4.

Difficult tracheal intubation in all trials (Analysis 1.1)

A total of 34 trials with 3565 participants provided data on DTI. Among 1507 participants undergoing tracheal intubation without the use of NMBA, a crude proportion of 24.2% ($n = 365$; 95% CI 22.4% to 26.4%) were intubated with difficulties. The median prevalence of a DTI among individual studies was 29.5% (interquartile range 7.0% to 60.0%). Of 2058 participants intubated with the use of NMBA, the crude proportion of DTI was 8.1% ($n = 167$; 95% CI 7.1% to 9.3%). The median prevalence of a DTI among individual studies was 3.9% (interquartile range 0.0% to 8.3%). In a random-effects model, avoidance of NMBA significantly increased the risk of a DTI with direct laryngoscopy (RR 5.00, 95% CI 3.49 to 7.15; $P < 0.00001$; NNTH = 6.3, $I^2 = 70\%$, $D^2 = 82\%$). In a fixed-effect model, RR was 4.79 (95% CI 4.04 to 5.69; $P < 0.0001$). Our TSA of all included trials did not contradict our findings in trials with low risk of bias. Again, the cumulative z-curve crossed the TSMB for harm, and the TSA-adjusted CI for the RR was 1.20 to 20.77. These findings provide evidence of the smallest possible harmful effect on the proportion

of DTI in the traditional meta-analysis of avoiding NMBA compared with using NMBA, even when the significance level was adjusted for repetitive testing and sparse data in a cumulative meta-analysis. We downgraded the quality of the evidence (GRADE) two levels to low because of indirectness, heterogeneity, and high or uncertain risk of bias (Analysis 1.1; Figure 4).

Subgroup analyses of difficult tracheal intubation**Depolarizing versus non-depolarizing NMBA (Analysis 1.2)**

In the subgroup of trials that used depolarizing NMBA, risk of DTI was increased with no use of NMBA (RR 5.79, 95% CI 2.64 to 12.72; $P < 0.0001$; NNTH = 5.6, $I^2 = 69\%$). Likewise, in the subgroup using non-depolarizing NMBA, DTI was associated with no use of NMBA (RR 4.72, 95% CI 3.17 to 7.02; $P < 0.0001$; NNTH = 5.8, $I^2 = 74\%$). Results show no significant subgroup differences ($P = 0.65$) (Analysis 1.2).

Remifentanyl versus other opioids (Analysis 1.3)

Investigators in 11 trials (Alexander 1999; Bouvet 2008; González Obregón 2010; Gulhas 2013; Hanna 2010; Jiao 2014; McNeil 2000; Mencke 2014; Pang 2014; Schlaich 2000; Yazdi 2016) used remifentanyl as an opioid. However, researchers in four studies (Alexander 1999; Hanna 2010; McNeil 2000; Yazdi 2016) used remifentanyl only in the intervention groups and used no opioids in the control group. One (Alexander 1999) of the three trials used remifentanyl and alfentanil as opioids in two different intervention groups. Another three trials (González Obregón 2010; Gulhas 2013; Jiao 2014) used an amount of remifentanyl in the intervention group that exceeded the amount used in the control group. Among trials using equal amounts of remifentanyl in intervention and control groups (Bouvet 2008; Mencke 2014; Pang 2014; Schlaich 2000), results showed significantly increased risk of DTI when NMBA was avoided for tracheal intubation (RR 15.86, 95% CI 4.43 to 56.71; $P < 0.0001$; NNTH = 6.4, $I^2 = 0\%$). In trials in which opioids other than remifentanyl were used, avoiding NMBA was significantly associated with DTI (RR 5.20, 95% CI 3.53 to 7.64; $P < 0.0001$; NNTH = 6.5, $I^2 = 74\%$). Results showed no significant subgroup differences ($P = 0.10$). One trial provided no opioids (Isesele 2012) (Analysis 1.3).

Alfentanil versus other opioids (Analysis 1.4)

Thirteen trials (Alexander 1999; Barclay 1997; Beck 1993; Combes 2007; Dominici 1990; Harsten 1997; Kopman 2001; Nimmo 1995; Rousseau 1995; Scheller 1992; Sivalingam 2001; Stevens 1997; Wong 1996) used alfentanil as the opioid. Five trials (Alexander 1999; Beck 1993; Scheller 1992; Stevens 1997; Wong 1996) used alfentanil only in the intervention group and used no opioids in the control group. One of the five trials used remifentanyl and alfentanil as opioids in two different intervention groups (Alexander 1999). In two trials, the amount of alfentanil used in the intervention group exceeded the amount used in the control group (Combes 2007; Sivalingam 2001). The six trials with equal amounts of alfentanil in the intervention and control groups reported significantly increased risk of DTI when NMBA was avoided for tracheal intubation (RR 4.46, 95% CI 1.66 to 11.98; $P = 0.0002$; $I^2 = 79\%$, NNTH = 5.2) (Barclay 1997; Dominici 1990; Harsten 1997; Kopman 2001; Nimmo 1995; Rousseau 1995). In trials using opioids other than alfentanil, risk of DTI was increased when NMBA was avoided (RR 5.10, 95% CI 3.34 to 7.79; $P < 0.0001$; $I^2 = 72\%$, NNTH = 4.8). Results showed no significant subgroup differences ($P = 0.81$). One trial provided no opioids (Isesele 2012) (Analysis 1.4).

Local anaesthetic versus no use of local anaesthetic (Analysis 1.5)

Eight trials used a local anaesthetic. However, three trials (Isesele 2012; Rousseau 1995; Stevens 1997) used a local anaesthetic only in the intervention group - not in the control group. Thus, we included five trials (Barclay 1997; Dominici 1990; Hanna 2010; Pang 2014; Striebel 1995) that used local anaesthesia for intubation in both control and intervention groups. In comparisons of trials using local anaesthesia versus trials not using local anaesthesia, risk of DTI was increased with no use of NMBA. With use of local anaesthesia, the RR was 1.90 (95% CI 1.14 to 3.18; $P = 0.01$; NNTH = 9.4, $I^2 = 35\%$), and with no use of local anaesthesia, the RR was 6.26 (95% CI 4.15 to 9.44; $P < 0.0001$; NNTH = 5.7, $I^2 = 69\%$), respectively. Results showed significant subgroup differences ($P < 0.0001$) (Analysis 1.5).

Exclusion of patients with expected DTI versus no exclusion of patients with expected DTI (Analysis 1.6)

In the subgroup in which patients with expected DTI were excluded from the individual trials, risk of DTI was significantly increased when NMBA was avoided (RR 5.32, 95% CI 3.54 to 8.00; $P < 0.00001$; NNTH = 6.3, $I^2 = 72\%$). In the subgroup in which patients with expected DTI were not excluded, risk of DTI was significantly increased when NMBA was avoided (RR 4.40, 95% CI 1.71 to 11.29; $P = 0.0002$; NNTH = 5.7, $I^2 = 74\%$). Results showed no significant subgroup differences ($P = 0.72$) (Analysis 1.6).

Explorative assessments of difficult tracheal intubation

A best-case scenario (Analysis 1.7)

A total of 18 trials were dose-finding studies that included more intervention or control groups, or both (Alexander 1999; Barclay 1997; Kahwaji 1997; Kirkegaard-Nielsen 1999; Kopman 2001; Lieutaud 2003; Lowry 1999; McNeil 2000; Naguib 2003; Naguib 2006; Nimmo 1995; Pino 1998; Scheller 1992; Schlaich 2000; Sivalingam 2001; Stevens 1997; Striebel 1995; Wong 1996). In attempting to estimate a sufficient level of intervention or adjuvant drugs, or both, these trials may have included intervention or control groups, or both, in which participants were suboptimally anaesthetized. Therefore, we performed a supplementary sensitivity analysis of a best-case scenario. Here, dose-finding studies were represented only by control and intervention groups with the lowest prevalence of difficult intubation. Among 1180 participants undergoing tracheal intubation without the use of NMBA, a crude proportion of 25.0% ($n = 295$; 95% CI 22.9% to 27.5%) were intubated with difficulties. Among 1230 participants intubated with the use of NMBA, the crude proportion of DTI was 3.4% ($n = 42$; 95% CI 2.5% to 4.4%). Avoidance of NMBA significantly increased the risk of DTI with direct laryngoscopy (RR 5.99, 95% CI 3.46 to 10.38; $P < 0.0001$; NNTH = 4.4, $I^2 = 57\%$) (Analysis 1.7).

Excluding dose-finding studies (Analysis 1.8)

In our attempt to exclude comparisons of suboptimal anaesthetic dosing regimens, we performed a sensitivity analysis that excluded all dose-finding trials. Thus we included 16 trials that had one intervention group and one control group (Beck 1993; Bouvet 2008; Combes 2007; Dominici 1990; González Obregón 2010; Gulhas 2013; Hanna 2010; Harsten 1997; Iamaroon 2001; Isesele 2012; Jiao 2014; Mencke 2003; Mencke 2014; Pang 2014; Rousseau 1995; Yazdi 2016). Among 769 participants undergoing tracheal intubation without the use of NMBA, a crude proportion of 15.5% ($n = 119$; 95% CI 13.3% to 18.0%) were intubated with difficulties. Among 767 participants intubated with the use of NMBA, the crude proportion of DTI was 4.2% ($n = 32$; 95% CI 3.0% to 5.6%). Avoidance of NMBA significantly increased the risk of DTI (RR 3.40, 95% CI 1.63 to 7.10; $P = 0.001$; NNTH = 8.8, $I^2 = 59\%$).

We wanted to explore possible causes of heterogeneity by performing meta-regression analyses. However, owing to a low degree of heterogeneity in trials with low risk of bias in terms of DTI, we did not perform these assessments. Regarding our other secondary outcome, we categorized only one trial describing upper airway discomfort and/or injury as having low risk of bias. Among all trials describing upper airway discomfort or injury, or both, we found no heterogeneity; thus we performed no meta-regression analysis (Analysis 1.8).

Funding from pharmaceutical industry (Analysis 1.9)

We identified 10 trials that reported funding from the pharmaceutical industry (Barclay 1997; Hanna 2010; Harsten 1997; Iamaroon 2001; Kahwaji 1997; Kopman 2001; Lowry 1999; Pino 1998; Scheller 1992; Wong 1996). In attempting to identify any potential bias caused by industrial funding, we performed a sensitivity analysis to compare trials receiving industrial funding versus the remaining trials. We included all trials receiving industrial funding, thus we did not distinguish between the types of funding that studies had received. Among the 10 trials that reported receipt of industrial funding, results showed a significant association between avoidance of NMBA and DTI (RR 4.10, 95% CI 2.67 to 6.31; $P = 0.003$; $I^2 = 64\%$). In the remaining trials, which reported no industrial funding, results showed a significant association between avoidance of NMBA and DTI (RR 5.33, 95% CI 3.16 to 8.98; $P < 0.00001$; $I^2 = 68\%$). Findings showed no significant subgroup differences ($P = 0.45$) (Analysis 1.9).

Primary outcome: overall mortality

None of the trials provided data on mortality.

Primary outcome: one or more events of upper airway discomfort or injury**One or more events of upper airway discomfort or injury in trials with lower risk of bias (Analysis 1.10)**

One trial with low risk of bias described events of upper airway discomfort or injury (Mencke 2003). This trial found a significant association between avoidance of NMBA and upper airway discomfort or injury (RR 2.74, 95% CI 1.21 to 6.21; $P = 0.02$). A random-effects model among trials with high or uncertain risk of bias revealed a significant association between no use of NMBA and risk of upper airway discomfort or injury (RR 1.30, 95% CI 1.08 to 1.58; $P = 0.006$; NNTH = 9.7, $I^2 = 0\%$). Because we identified only one low risk of bias trial, we performed no quality of evidence (GRADE) assessment (Analysis 1.10; Figure 5).

Figure 5. **One or more events of upper airway discomfort or injury in all trials (Analysis 1.10)**

Seven trials representing 846 participants described events of upper airway discomfort or injury (Bouvet 2008; Combes 2007; González Obregón 2010; Gulhas 2013; Mencke 2003; Mencke 2014; Sivalingam 2001). However, one trial (González Obregón 2010) described no events of upper airway discomfort or injury. The crude prevalence was 33.1% ($n = 280$; 95% CI 30.4% to 36.3%). Among 447 participants undergoing tracheal intubation without the use of NMBA, a proportion of 38.2% ($n = 171$; 95% CI 34.4% to 42.8%) with upper airway discomfort or injury were identified. Among 399 participants intubated with the use of NMBA, the proportion of upper airway discomfort or injury was 27.3% ($n = 109$; 95% CI 23.5% to 31.7%). Risk of upper airway discomfort or injury was significant with avoidance of NMBA (RR 1.37, 95% CI 1.09 to 1.74; $P = 0.008$; NNTH = 9.1, $I^2 = 13\%$). Our TSA of non-NMBA versus NMBA using a D^2 -adjusted required information size of 1981 participants to detect or reject a 20% relative risk reduction with power of 80% and overall type 1 error of 5%. The cumulative z-curve crossed the

TSMB for harm, and the TSA-adjusted CI for the RR was 1.00 to 1.85. Thus, we provided firm evidence on the proportion of upper airway discomfort or injury for the least possible harmful effect in the traditional meta-analysis of avoiding NMBA compared with using NMBA, even when the significance level was adjusted for repetitive testing and sparse data in a cumulative meta-analysis. We downgraded the quality of the evidence (GRADE) one level to moderate because of high or uncertain risk of bias (Analysis 1.10; Figure 5).

Subgroup analyses of one or more events of upper airway discomfort or injury**Depolarizing versus non-depolarizing NMBA (Analysis 1.11)**

The two trials in which investigators used depolarizing NMBA showed no significant association between avoidance of NMBA and upper airway discomfort or injury (RR 1.48, 95% CI 0.83 to 2.65; $P = 0.19$; $I^2 = 0\%$) (Gulhas 2013; Sivalingam 2001). Likewise, in the subgroup of five trials using non-depolarizing NMBA, results revealed no significant association (RR 1.37, 95% CI 0.97 to 1.94;

$P = 0.07$; $I^2 = 15\%$) (Bouvet 2008; Combes 2007; González Obregón 2010; Mencke 2003; Mencke 2014) and no significant subgroup differences ($P = 0.83$) (Analysis 1.11).

Remifentanyl versus other opioids (Analysis 1.12)

The two trials (Bouvet 2008; Mencke 2014) that used remifentanyl showed no significant association between avoidance of NMBA and upper airway discomfort or injury (RR 1.12, 95% CI 0.61 to 2.08; $P = 0.14$; $I^2 = 55\%$). The five trials using opioids other than remifentanyl revealed a significant association between avoidance of NMBA and upper airway discomfort or injury (RR 1.42, 95% CI 1.16 to 1.75; $P = 0.0009$; NNTH = 7.5, $I^2 = 0\%$) (Combes 2007; González Obregón 2010; Gulhas 2013; Mencke 2003; Sivalingam 2001) and no significant subgroup differences ($P = 0.47$) (Analysis 1.12).

Alfentanil versus other opioids (Analysis 1.13)

In two trials, the amount of alfentanil used in the intervention group exceeded the amount used in the control group; thus we excluded these trials from the meta-analysis (Combes 2007; Sivalingam 2001). The remaining trials used opioids other than alfentanil (Bouvet 2008; González Obregón 2010; Gulhas 2013; Mencke 2003; Mencke 2014), and results showed no significant association between no use of NMBA and the presence of upper airway discomfort or injury (RR 1.47, 95% CI 0.85 to 2.53; $P = 0.17$; $I^2 = 49\%$) (Analysis 1.13).

Local anaesthetic versus no use of local anaesthetic

None of the trials reporting upper airway discomfort or injury used local anaesthetic.

Exclusion of patients with expected DTI versus no exclusion of patients with expected DTI (Analysis 1.14)

Six of the trials reporting upper airway discomfort or injury excluded patients with anticipated difficult airway management

(Bouvet 2008; Combes 2007; González Obregón 2010; Mencke 2003; Mencke 2014; Sivalingam 2001). Results showed significantly increased risk of upper airway discomfort or injury when the patient was not relaxed (RR 1.37, 95% CI 1.05 to 1.79; $P = 0.02$; NNTH = 8.9, $I^2 = 29\%$). One trial included participants with expected difficult airway management (Gulhas 2013). Results showed no significant association between avoidance of NMBA and upper airway discomfort or injury (RR 2.00, 95% CI 0.39 to 10.31; $P = 0.41$) and no significant subgroup differences (Analysis 1.14).

Secondary outcome: one or more major serious adverse events

One study reported serious adverse events (Kahwaji 1997). A 29-year-old man, weighing 100 kg, with ASA physical status of I, experienced two of these events (tachycardia with heart rate from 85 to 150 bpm and bronchospasm) within 30 seconds of administration of 2.0 mg/kg ORG 9487 (rapacuronium); these events were followed by erythema of the arms, shoulders, and face. Bronchospasm was treated with salbutamol, and all symptoms gradually subsided.

Secondary outcome: difficult laryngoscopy

Difficult laryngoscopy in trials with low risk of bias (Analysis 1.15)

We categorized one trial as having low risk of bias (Mencke 2003). This trial found no significant association between conditions for laryngoscopy and use of NMBA (RR 4.00, 95% CI 0.47 to 34.20; $P = 0.21$). Comparison with the subgroup of trials with high or uncertain risk of bias revealed significantly increased risk of difficult laryngoscopy in a random-effects model when NMBA was avoided (RR 2.47, 95% CI 1.47 to 4.16; $P = 0.0006$; NNTH = 27.9, $I^2 = 0\%$) (Analysis 1.15; Figure 6).

Figure 6.

Difficult laryngoscopy in all trials (Analysis 1.15)

Thirteen trials representing 1308 participants reported the conditions for laryngoscopy (Beck 1993; Bouvet 2008; Combes 2007; Dominici 1990; González Obregón 2010; Harsten 1997; McNeil 2000; Mencke 2003; Mencke 2014; Pang 2014; Scheller 1992; Stevens 1997; Striebel 1995). Among 740 participants undergoing tracheal intubation without the use of NMBA, a crude proportion of 7.2% (n = 53; 95% CI 5.6% to 9.0%) had a difficult laryngoscopy. Among 568 participants intubated with use of NMBA, the proportion of difficult laryngoscopy was 3.3% (n = 19; 95% CI 2.1% to 4.8%). Avoidance of NMBA significantly increased the risk of a difficult laryngoscopy (RR 2.54, 95% CI 1.53 to 4.21; P = 0.0003; NNTH = 25.6, I² = 0%, D² = 0%). In a fixed-effect model, RR was 2.46 (95% CI 1.52 to 3.97; P = 0.0002). Trial sequential analysis of avoiding versus using NMBA, with a D²-adjusted required information size, showed that only 6% of the information size required to detect or reject a 20% relative risk reduction was accrued and the TMSB was not crossed. The TSA-adjusted 95% CI for the RR was 0.27 to 21.75, meaning that firm evidence could not be established (Analysis 1.15; Figure 6).

DISCUSSION

Summary of main results

Our systematic review reveals several important findings. Analysis of 34 trials reporting on conditions regarding tracheal intubation suggests that avoidance of neuromuscular blocking agents (NMBA) is associated with increased risk of a difficult tracheal intubation (DTI) (Summary of findings for the main comparison). We confirmed this in our assessment of the four trials categorized as having low risk of bias (Analysis 1.1). Here, avoidance of NMBA was associated with DTI with a risk ratio (RR) of 13.27 (95% confidence interval (CI) 8.19 to 21.49; P < 0.00001; number needed to treat for an additional harmful outcome (NNTH) = 1.9, I² = 0%). Results showed

a significant subgroup difference between trials with low and high or uncertain risk of bias, as avoiding NMBA exceeded the risk ratio of a difficult intubation among trials with low risk of bias compared with trials with high or uncertain risk of bias. Fixed-effect meta-analyses did not noticeably change the conclusions. Our trial sequential analysis (TSA) provided firm evidence for the least possible harmful effect on the proportion of DTI in the traditional meta-analysis of avoiding NMBA compared with using NMBA, even when the significance level was adjusted for repetitive testing and sparse data in a cumulative meta-analysis.

In our subgroup analyses on DTI, use of remifentanyl, alfentanil, or local anaesthesia did not change our primary finding, as avoidance of NMBA was significantly associated with a DTI. However, our subgroup analysis suggests that local anaesthetics may have a protective effect, as results showed a statistically significant subgroup difference between trials using and trials avoiding local anaesthesia in relation to tracheal intubation. Likewise, in the subgroup analysis of trials excluding participants with expected difficult intubation, avoiding NMBA increased the risk of a difficult intubation. Fixed-effect meta-analyses did not noticeably change results of the subgroup analyses.

Our analysis of seven trials suggests that avoidance of NMBA is associated with increased risk of upper airway discomfort or injury (RR = 1.37, 95% CI 1.09 to 1.74; P = 0.008; NNTH = 9.1, I² = 13%). Only one trial reporting this outcome had low risk of bias, but results revealed a significant association between avoidance of NMBA and upper airway discomfort or injury (Mencke 2003). Our TSA provided firm evidence for a harmful effect on the proportion of upper airway discomfort and injury in the traditional meta-analysis when the significance level is adjusted for repetitive testing and sparse data in a cumulative meta-analysis.

For 13 trials describing conditions for laryngoscopy, our meta-analysis showed that avoiding NMBA was associated with a difficult laryngoscopy, with RR of 2.54 (95% CI 1.53 to 4.21; $P = 0.0003$; NNTH = 14.6) (Beck 1993; Bouvet 2008; Combes 2007; Dominici 1990; González Obregón 2010; Harsten 1997; McNeil 2000; Mencke 2003; Mencke 2014; Pang 2014; Scheller 1992; Stevens 1997; Striebel 1995). Only one trial categorized as having low risk of bias reported this outcome (Mencke 2003). Results show no significant association between avoidance of NMBA and difficult laryngoscopy. Because information size was inadequate, our TSA of all 13 trials provided no firm evidence for a harmful effect of avoiding NMBA on upper airway discomfort or injury.

No trials reported mortality, and only one trial reported one episode of a serious adverse event (SAE) related to the use of NMBA (Kahwaji 1997).

Overall completeness and applicability of evidence

The plan to analyse effects of neuromuscular blocking agents for improving conditions during tracheal intubation or direct laryngoscopy in randomized clinical trials in adults followed our published protocol to a great extent. We included all eligible randomized clinical trials up to January 2017. Most of these trials were conducted in high-income countries, and most participants were undergoing elective surgery. We included participants of both genders, and most participants were American Society of Anesthesiologists (ASA) class I or II and non-obese, without expected airway management difficulties. Studies showed a high degree of clinical diversity as included combinations of different adjuvant drugs and NMBA were substantial.

About half of the included trials used a (modified) intubation score according to "Good clinical research practice (GCRP) in pharmacodynamic studies of neuromuscular blocking agents" (Fuchs-Buder 2007; Viby-Mogensen 1996). Further, a major portion of the remaining trials used intubation scores that incorporated many of the elements described by the GCRP score. Thus, to some extent, included trials showed some agreement on how DTI should be defined. Although trials used a defined outcome in accordance with recommendations for research regarding NMBA, definitions used differ from those traditionally used in research dealing with difficult airway management (Adnet 1997; Lundström 2011b). Although the literature lacks consensus, these definitions often describe multiple (unsuccessful) attempts at intubation and/or shift to more advanced intubation techniques and/or other personnel performing airway management. One must therefore bear in mind that our primary outcome may not reflect a situation that most clinicians consider a truly difficult intubation that threatens the patient's life or health. Thus, one may observe a substantial degree of indirectness concerning our primary outcome describing DTI. Thirteen trials used the difficult laryngoscopy score of Cormack and Lehane (Cormack 1984) as a surrogate measure of a difficult intubation. This outcome measure is often used in studies on difficult airway management; thus it seems reasonable to interpret our findings on difficult laryngoscopy in a traditionally difficult airway management context. Most trials excluded patients expected to have difficult airway management. This may have a significant impact on the applicability of evidence, but subgroup analysis of 'Exclusion of patients with expected DTI versus no exclusion of expected DTI' (Analysis 1.6) showed no significant subgroup differences. Further, several studies have demonstrated that prediction of a DTI is difficult (Lundström 2011b; Nørskov 2015;

Nørskov 2016) and involves low sensitivity and specificity; thus the impact of these exclusions is uncertain.

Selected outcome measures varied among trials dealing with sore throat, upper airway discomfort, and injury. Further, by whom, how, or when the outcome measure was evaluated differed or was not clearly defined by some of these trials - a fact that must be taken into account when our results are interpreted in a clinical context.

Of specific notice is the fact that evaluation of the effect on occurrence of serious adverse events (SAE) of avoidance of NMBA compared with use of NMBA is virtually absent, as none of the included trials evaluated effects on mortality (short-term or long-term), and only one trial evaluated effects on other types of SAEs, reporting one SAE.

Quality of the evidence

We present our main results on all investigated outcomes in [Summary of findings for the main comparison](#), which shows grading of the quality of evidence, including imprecision, according to the GRADE approach (Guyatt 2013; Guyatt 2011) and stresses results of meta-analyses originating from trials with overall low risk of bias. Our review follows the overall plan of a published, peer-reviewed Cochrane protocol (Lundström 2011a). It represents a comprehensive review of the topic, including meta-analyses of results from 34 randomized trials with a total of 3565 participants. To our knowledge, no previous meta-analyses have included trials comparing use or avoidance of NMBA for tracheal intubation. We conducted a thorough review in accordance with methods of the Cochrane Collaboration (Higgins 2011). Between-trial heterogeneity varied from absent to substantial among the various meta-analyses in terms of our different outcomes. This may emphasize the diversity of the dose-finding regimens used in different trials that used substantially different combinations of adjuvant drugs and NMBA. Further, minor differences among definitions of a DTI in some degree may contribute to the heterogeneity. In contrast, we noted no between-trial heterogeneity regarding our secondary outcome describing conditions needed for direct laryngoscopy, which may be explained by a well-defined outcome measure (Cormack 1984) that was retrievable from all 13 trials.

We performed trial sequential analyses and calculated TSA-adjusted confidence intervals reflecting sparse data and multiple testing due to potentially repeated updates of cumulative meta-analyses to control risk of random errors and to prevent premature statements of superiority of one intervention over another (Brok 2008; Brok 2009; Thorlund 2009; Thorlund 2011; Wetterslev 2008; Wetterslev 2009). TSA revealed that all of our meta-analyses including the primary outcomes have a very low degree of imprecision, indicating low risk of random error. The finding of an insignificant TSA-adjusted association between avoidance of NMBA and difficult laryngoscopy may be due to a type II error, that is, a false-negative finding, exemplified by the cumulative z-curve not crossing the trial sequential monitoring boundary (TSMB) for futility. However, such an analysis cannot remove risks of bias - detected or undetected. It is worth discussing how much evidence is required when one is dealing with potential benefit or harm. On the other hand, beneficial or harmful effects can occur as the result of random errors; therefore, sufficient information must be assessed to demonstrate benefit or harm beyond reasonable doubt.

Because actual airway management was evaluated just after induction of anaesthesia, no cases were lost to follow-up. However, cases with upper airway discomfort and/or injury were lost to follow-up in three trials ([Combes 2007](#); [Mencke 2003](#); [Mencke 2014](#)). We therefore performed post hoc "best- and worst-case" sensitivity analyses. We included participants who were lost to follow-up with or without upper airway discomfort and/or injury. These explorative analyses did not alter our primary findings derived both from all trials and from trials with low risk of bias.

Potential biases in the review process

We strived to reduce bias by identifying all relevant trials through a comprehensive systematic search of the literature. We contacted study authors to retrieve unpublished data, when possible. However, it was difficult to obtain updated contact information for authors of the oldest published trials. This may have introduced bias, as it was generally easier to retrieve additional data from more recently published trials, which also were more likely to be categorized as having low risk of bias. Two review authors assessed trials for inclusion or eligibility and extracted all data in duplicate, thereby reaching a high level of agreement. We did not conduct quality assessments or data extractions while blinded to review authors and bias risks.

Most included trials clearly stated cut-off values defining a DTI. However, five trials ([Beck 1993](#); [Gulhas 2013](#); [McNeil 2000](#); [Scheller 1992](#); [Stevens 1997](#)) used different composite scores without defined cut-offs. We used underlying references or comparable scores to define cut-off values, enabling us to include these trials in our meta-analyses. The [Characteristics of included studies](#) section provides detailed explanations. In these cases, some degree of interpretation of outcome measures was necessary, and our assessments may have introduced bias as a consequence. Nevertheless, we believe that this categorization was evident when compared with the existing literature, and we therefore consider risk of introducing bias as low.

The authors of six trials ([Harsten 1997](#); [Jiao 2014](#); [Mencke 2003](#); [Mencke 2014](#); [Nimmo 1995](#); [Sivalingam 2001](#)) excluded a few participants from their assessments when tracheal intubation failed. We consider these patients as truly difficult to intubate and thus chose to include them in our meta-analyses. This may have introduced bias; however, a sensitivity analysis excluding these participants from our meta-analyses did not change the conclusion.

Agreements and disagreements with other studies or reviews

To our knowledge, no previous systematic review has included meta-analyses of trials comparing use or avoidance of NMBA for tracheal intubation. However, a narrative review ([Woods 2005](#)) concluded that the literature describes successful techniques to intubate the trachea without the use of neuromuscular blocking agents with the patient under general anaesthesia. Further, those review authors concluded that these techniques offer a useful alternative when drugs are contraindicated or undesirable. Another narrative review ([Fotopoulou 2012](#)) concluded that induction of anaesthesia without the use of NMBA but in combination with remifentanyl provides acceptable conditions for tracheal intubation. Both reviews include several dose-finding trials in which none of the participants in the different intervention groups were administered NMBA. Thus, it was impossible to compare

avoiding versus using NMBA for tracheal intubation. On the basis of findings related to specific drug combinations and concentrations, the review authors concluded that induction without the use of NMBA may offer (almost) perfect or good conditions for intubation.

In large observational studies and reviews on difficult airway management, fractions of DTI range from 2% to 7% ([Lundström 2011b](#); [Nørskov 2015](#); [Shiga 2005](#)), which is consistent with our findings on the overall proportion of difficult intubation in patients who are intubated with use of NMBA. The crude proportion was 4.7% in trials with low risk of bias and 8.1% in all trials. However, among participants undergoing tracheal intubation without the use of NMBA, corresponding crude proportions were tremendously higher, at 56.3% in trials with low risk of bias and 24.2% in all trials. These latter proportions are not consistent with those reported in experiences from everyday clinical practice, and several trials may therefore include suboptimal dosing regimens. Thus, we speculate that some of the dose-finding trials may include control groups with clinically unacceptable dosing regimens that may contribute to an unrealistically increased risk of DTI. As an example, one trial concluded that the optimal remifentanyl dose used for intubation with propofol and without NMBA is 4 µg/kg administered in 60 seconds ([Bouvet 2009](#)). However in our review, only two of the included trials actually used equivalent or larger doses of remifentanyl ([Hanna 2010](#); [McNeil 2000](#)). Therefore, we performed a post hoc sensitivity analysis of a best-case scenario ([Analysis 1.7](#)). Dose-finding trials were represented only by control and intervention groups with the lowest incidence of difficult intubation. Results showed a crude proportion of 25.0% DTI among participants anaesthetized without the use of NMBA, and this again significantly increased the risk of DTI, with a risk ratio of 5.99. Finally, in our attempt to exclude comparisons of suboptimal anaesthetic dosing regimens, we performed a post hoc sensitivity analysis while excluding all dose-finding trials ([Analysis 1.8](#)). We included in our meta-analysis 16 trials with a single intervention group and a single control group. The crude proportion of DTI among participants anaesthetized without the use of NMBA decreased to 15.4%. However, avoiding NMBA still significantly increased the risk of DTI, with a risk ratio of 3.40. Another possible explanation for the increased crude proportion of DTI among participants induced without NMBA may be that most included trials used a (modified) intubation score according to "Good clinical research practice (GCRP) in pharmacodynamic studies of neuromuscular blocking agents" ([Fuchs-Buder 2007](#); [Viby-Mogensen 1996](#)). Assessment of intubation conditions included information on jaw mobility and/or reactions to insertion of the tracheal tube and cuff inflation (diaphragmatic movement/coughing). These are reasonable elements for describing ideal conditions for intubation, although they may be affected even when everyday clinical airway management is successful without "true" difficulties. Thus, the GCRP score may be highly sensitive and may be used to enforce the difference between the ideal procedure and a clinically acceptable intubation. Trials reporting conditions for laryngoscopy (described by [Cormack 1984](#)) report an outcome measure traditionally used in studies dealing with difficult airway management. In our assessment, the crude proportions of difficult laryngoscopy reported among intervention groups may reflect a more clinically acceptable situation, as 4.2% among all trials and 7.2% in trials with low risk of bias who were intubated without NMBA had a difficult laryngoscopy.

In the literature, one review reported on various causes for postoperative sore throat (El-Boghdadly 2016). These review authors state that tracheal intubation without the use of NMBA is a potential risk factor for sore throat. Although this study is presented as a systematic review, review authors included only one study (Combes 2007) reporting on upper airway discomfort or injury, review authors performed no bias assessment, and the review included no meta-analyses. Other conditions such as tube size (Jaensson 2010; Jaensson 2012; Stout 1987; Xu 2012) and use of corticosteroids (Sumathi 2007; Thomas 2007) may play an important role in the postoperative sore throat. In our review, we did not evaluate the impact of these factors in subgroup analyses. The review of EL-Boghdadly reports postoperative sore throat with a prevalence of up to 62% following general anaesthesia. In a Cochrane review on lidocaine for preventing postoperative sore throat (Tanaka 2015), the crude prevalence of sore throat was 20% to 30% pending the use of lidocaine. In our assessment, the crude prevalence was 33%.

AUTHORS' CONCLUSIONS

Implications for practice

This review supports that use of NMBA may create better conditions for tracheal intubation in clinical practice than are provided by avoidance of NMBA. Review results are characterized by indirectness regarding our primary outcome describing DTI. Given defined outcomes in the included trials, our primary outcome may not reflect a situation that most clinicians consider to show a clinically important DTI by which the patient's life or health may be threatened. Thus, we could not conclude with certainty that avoidance of NMBA was associated with a clinically important or seriously difficult intubation. A difficult laryngoscopy as defined by Cormack and Lehane may be a reasonable surrogate for a clinically serious difficult airway. Our results indicate that avoiding NMBA increases the risk of a difficult laryngoscopy, but after adjustments for required information size, firm evidence could not be established.

In terms of our other primary outcome - events of upper airway discomfort or injury - data on low risk of bias trials are sparse. Among all trials, evidence shows a firm and significant risk of upper

airway discomfort or injury when NMBA is avoided. Of specific notice is the fact that evaluation of the effects of avoidance of NMBA compared with use of NMBA on the occurrence of serious adverse events (SAEs) is virtually absent, as none of these trials evaluated effects on mortality (short-term or long-term), and only one trial evaluated effects on other types of SAEs, reporting one SAE.

In conclusion, in a clinical context, one must have weighty arguments for using or not using NMBA when performing tracheal intubation.

Implications for research

Our assessments show some degree of indirectness for our primary outcome of describing difficult intubation. Insufficient information size led to uncertainty regarding our assessment of the effects of avoiding NMBA on the frequency of a difficult laryngoscopy and implications for future research focusing on the impact of avoiding NMBA on the prevalence of severe intubation difficulties and difficult laryngoscopy as categorized by Cormack and Lehane. In addition, large trials with low risk of bias undertaken to describe upper airway injury and discomfort, as well as other serious adverse events and mortality, are needed.

ACKNOWLEDGEMENTS

We would like to thank Karen Hovhannisyan and Janne Vendt (former and present Cochrane Anaesthesia, Critical and Emergency Care Group Trial Information specialists, respectively) for their valuable help with phrasing of the search strategies.

We would like to thank Andrew Smith (Content Editor); Jing Xie (Statistical Editor); Emmanuel Boselli and Thomas Mencke (Peer Reviewers); and Brian Stafford (Consumer Referee) for help and editorial advice provided during preparation of this systematic review. We would like to thank Julie Wetterslev (cand. scient. soc.) for assistance with Spanish translation.

We would like to thank Andy Smith (Content Editor); Marialena Trivella (Statistical Editor); Rodrigo Cavallazzi, Shahla Siddiqui, and Fred Cheney (Peer Reviewers), and Tracey Lloyd (representative of the ACE consumer panel) for help and editorial advice provided during preparation of the protocol (Lundström 2011a).

REFERENCES

References to studies included in this review

Alexander 1999 {published data only}

Alexander R, Olufolabi AJ, Booth J, El-Moalem HE, Glass PS. Dosing study of remifentanyl and propofol for tracheal intubation without the use of muscle relaxants. *Anaesthesia* 1999;**54**(11):1037-40. [PUBMED: 10540091]

Barclay 1997 {published data only}

Barclay K, Eggers K, Asai T. Low-dose rocuronium improves conditions for tracheal intubation after induction of anaesthesia with propofol and alfentanil. *British Journal of Anaesthesia* 1997;**78**(1):92-4. [PUBMED: 9059214]

Beck 1993 {published data only}

Beck GN, Masterson GR, Richards J, Bunting P. Comparison of intubation following propofol and alfentanil with intubation following thiopentone and suxamethonium. *Anaesthesia* 1993;**48**(10):876-80. [PUBMED: 8238830]

Bouvet 2008 {published data only}

Bouvet L, Stoian A, Jacquot-Laperriere S, Allaouchiche B, Chassard D, Boselli E. Laryngeal injuries and intubating conditions with or without muscular relaxation: an equivalence study. *Canadian Journal of Anaesthesia* 2008;**55**(10):674-84. [PUBMED: 18835965]

Combes 2007 {published data only}

Combes X, Andriamifidy L, Dufresne E, Suen P, Sauvat S, Scherrer E, et al. Comparison of two induction regimens using or not using muscle relaxant: impact on postoperative upper airway discomfort. *British Journal of Anaesthesia* 2007;**99**(2):276-81. [PUBMED: 17573390]

Dominici 1990 {published data only}

Dominici L, Gondret R, Dubos S, Crevot O, Deligné P. Intubation in otorhinolaryngologic surgery: propofol versus propofol-suxamethonium. *Annales Françaises d'anesthésie et de Réanimation* 1990;**9**(2):110-4. [PUBMED: 2363546]

González Obregón 2010 {published data only}

González Obregón MP, Rivera Díaz RC, Ordóñez Molina JE, Rivera Díaz JS, Velásquez Ossa LF, Pineda Ibarra C, et al. Tracheal intubation quality under remifentanyl-propofol with sevoflurane compared with remifentanyl-propofol with rocuronium: a randomized double-blind clinical trial. *Revista Española de Anestesiología y Reanimación* 2010;**57**(6):351-6. [PUBMED: 20645486]

Gulhas 2013 {published data only}

Gulhas N, Topal S, Erdogan Kayhan G, Yucel A, Begec Z, Yologlu S, et al. Remifentanyl without muscle relaxants for intubation in microlaryngoscopy: a double blind randomised clinical trial. *European Review for Medical and Pharmacological Sciences* 2013;**17**(14):1967-73. [PUBMED: 23877864]

Hanna 2010 {published data only}

Hanna SF, Ahmad F, Pappas AL, Mikat-Stevens M, Jellish WS, Kleinman B, et al. The effect of propofol/remifentanyl rapid-

induction technique without muscle relaxants on intraocular pressure. *Journal of Clinical Anesthesia* 2010;**22**(6):437-42. [PUBMED: 20868965]

Harsten 1997 {published data only}

Harsten A, Gillberg L. Intubating conditions provided by propofol and alfentanil - acceptable, but not ideal. *Acta Anaesthesiologica Scandinavica* 1997;**41**(8):985-7. [PUBMED: 9311395]

Iamaroon 2001 {published data only}

Iamaroon A, Pitimana-aree S, Prechawai C, Anusit J, Somcharoen K, Chaiyaroj O. Endotracheal intubation with thiopental/succinylcholine or sevoflurane-nitrous oxide anesthesia in adults: a comparative study. *Anesthesia and Analgesia* 2001;**92**(2):523-8. [PUBMED: 11159262]

Iselese 2012 {published data only}

Iselese T, Amadasun F, Edomwonyi N. Comparison of intubating conditions with propofol suxamethonium versus propofol-LIDOCAINE. *Journal of the West African College of Surgeons* 2012;**2**(4):51-67. [PUBMED: 25453004]

Jiao 2014 {published data only}

Jiao J, Huang S, Chen Y, Liu H, Xie Y. Comparison of intubation conditions and apnea time after anesthesia induction with propofol/remifentanyl combined with or without small dose of succinylcholine. *International Journal of Clinical and Experimental Medicine* 2014;**7**(2):393-9. [PUBMED: 24600494]

Kahwaji 1997 {published data only}

Kahwaji R, Bevan DR, Bikhazi G, Shanks CA, Fragen RJ, Dyck JB, et al. Dose-ranging study in younger adult and elderly patients of ORG 9487, a new, rapid-onset, short-duration muscle relaxant. *Anesthesia and Analgesia* 1997;**84**(5):1011-8. [PUBMED: 9141923]

Kirkegaard-Nielsen 1999 {published data only}

Kirkegaard-Nielsen H, Caldwell JE, Berry PD. Rapid tracheal intubation with rocuronium: a probability approach to determining dose. *Anesthesiology* 1999;**91**(1):131-6. [PUBMED: 10422938]

Kopman 2001 {published data only}

Kopman AF, Klewicka MM, Neuman GG. Reexamined: the recommended endotracheal intubating dose for nondepolarizing neuromuscular blockers of rapid onset. *Anesthesia and Analgesia* 2001;**93**(4):954-9. [PUBMED: 11574363]

Lieutaud 2003 {published data only}

Lieutaud T, Billard V, Khalaf H, Debaene B. Muscle relaxation and increasing doses of propofol improve intubating conditions. *Canadian Journal of Anesthesia* 2003;**50**(2):121-6. [PUBMED: 12560300]

Lowry 1999 {published data only}

Lowry DW, Carroll MT, Mirakhor RK, Hayes A, Hughes D, O'Hare R. Comparison of sevoflurane and propofol with

rocuronium for modified rapid-sequence induction of anaesthesia. *Anaesthesia* 1999;**54**(3):247-52. [PUBMED: 10364860]

McNeil 2000 {published data only}

McNeil IA, Culbert B, Russell I. Comparison of intubating conditions following propofol and succinylcholine with propofol and remifentanyl 2 micrograms kg⁻¹ or 4 micrograms kg⁻¹. *British Journal of Anaesthesia* 2000;**85**(4):623-5. [PUBMED: 11064625]

Mencke 2003 {published data only}

Mencke T, Echternach M, Kleinschmidt S, Lux P, Barth V, Plinkert PK, et al. Laryngeal morbidity and quality of tracheal intubation: a randomized controlled trial. *Anesthesiology* 2003;**98**(5):1049-56. [PUBMED: 12717124]

Mencke 2014 {published data only}

Mencke T, Jacobs RM, Machmueller S, Sauer M, Heidecke C, Kallert A, et al. Intubating conditions and side effects of propofol, remifentanyl and sevoflurane compared with propofol, remifentanyl and rocuronium: a randomised, prospective, clinical trial. *BMC Anesthesiology* 2014;**14**(39):ecollection. [PUBMED: 24860256]

Naguib 2003 {published data only}

Naguib M, Samarkandi A, Riad W, Alharby SW. Optimal dose of succinylcholine revisited. *Anesthesiology* 2003;**99**(5):1045-9. [PUBMED: 14576536]

Naguib 2006 {published data only}

Naguib M, Samarkandi AH, El-Din ME, Abdullah K, Khaled M, Alharby SW. The dose of succinylcholine required for excellent endotracheal intubating conditions. *Anesthesia and Analgesia* 2006;**102**(1):151-5. [PUBMED: 16368821]

Nimmo 1995 {published data only}

Nimmo SM, McCann N, Broome IJ, Robb HM. Effectiveness and sequelae of very low-dose suxamethonium for nasal intubation. *British Journal of Anaesthesia* 1995;**74**(1):31-4. [PUBMED: 7880702]

Pang 2014 {published data only}

Pang L, Zhuang YY, Dong S, Ma HC, Ma HS, Wang YF. Intubation without muscle relaxation for suspension laryngoscopy: a randomized, controlled study. *Nigerian Journal of Clinical Practice* 2014;**17**(4):456-61. [PUBMED: 24909469]

Pino 1998 {published data only}

Pino RM, Ali HH, Denman WT, Barrett PS, Schwartz A. A comparison of the intubation conditions between mivacurium and rocuronium during balanced anaesthesia. *Anesthesiology* 1998;**88**(3):673-8. [PUBMED: 9523811]

Rousseau 1995 {published data only}

Rousseau JM, Lemardeley P, Giraud D, Lemarié J, Ladagnous JF, Barriot P, et al. Endotracheal intubation under propofol with or without vecuronium. *Annales Françaises d'Anesthésie et de Réanimation* 1995;**14**(3):261-4. [PUBMED: 7486295]

Scheller 1992 {published data only}

Scheller MS, Zornow MH, Saidman LJ. Tracheal intubation without the use of muscle relaxants: a technique using propofol and varying doses of alfentanil. *Anesthesia and Analgesia* 1992;**75**(5):788-93. [PUBMED: 1416135]

Schlaich 2000 {published data only}

Schlaich N, Mertzlufft F, Soltész S, Fuchs-Buder T. Remifentanyl and propofol without muscle relaxants or with different doses of rocuronium for tracheal intubation in outpatient anaesthesia. *Acta Anaesthesiologica Scandinavica* 2000;**44**(6):720-6. [PUBMED: 10903016]

Sivalingam 2001 {published data only}

Sivalingam P, Kandasamy R, Dhakshinamoorthi P, Madhavan G. Tracheal intubation without muscle relaxant - a technique using sevoflurane vital capacity induction and alfentanil. *Anaesthesia and Intensive Care* 2001;**29**(4):383-7. [PUBMED: 11512649]

Stevens 1997 {published data only}

Stevens JB, Vescovo MV, Harris KC, Walker SC, Hickey R. Tracheal intubation using alfentanil and no muscle relaxant: is the choice of hypnotic important?. *Anesthesia and Analgesia* 1997;**84**(6):1222-6. [PUBMED: 9174296]

Striebel 1995 {published data only}

Striebel HW, Hölzl M, Rieger A, Brummer G. Endotracheal intubation with propofol and fentanyl. *Der Anaesthesist* 1995;**44**(12):809-17. [PUBMED: 8594954]

Wong 1996 {published data only}

Wong AK, Teoh GS. Intubation without muscle relaxant: an alternative technique for rapid tracheal intubation. *Anaesthesia and Intensive Care* 1996;**24**(2):224-30. [PUBMED: 9133197]

Yazdi 2016 {published data only}

Yazdi B, Khalili M, Dadashpour N, Nasab SMM, Khabiri M, Eraghi MG. The comparison of atracurium and remifentanyl effect on jaw relaxation and tracheal intubation condition without muscle relaxant in patients undergoing elective surgery. *Acta Medica Mediterranea - International Journal of Clinical Medicine* 2016;**32**(Special Issue 2):1029-32.

References to studies excluded from this review

Alcock 1993 {published data only}

Alcock R, Peachey T, Lynch M, McEwan T. Comparison of alfentanil with suxamethonium in facilitating nasotracheal intubation in day-case anaesthesia. *British Journal of Anaesthesia* 1993;**70**(1):34-7. [PUBMED: 8431329]

Baumgarten 1988 {published data only}

Baumgarten RK, Carter CE, Reynolds WJ, Brown JL, DeVera HV. Priming with nondepolarizing relaxants for rapid tracheal intubation: a double-blind evaluation. *Canadian Journal of Anesthesia* 1988;**35**(1):5-11. [PUBMED: 2894903]

Ide 2015 {published data only}

Ide M, Sunada K, Katsuyama N. Are muscle relaxants needed for nasal intubation in propofol and remifentanyl anesthesia?

Journal of Oral and Maxillofacial Surgery 2014;**72**(11):2134-9. [PUBMED: 25438272]

Additional references

Adnet 1997

Adnet F, Borron SW, Racine SX, Clemessy JL, Fournier JL, Plaisance P, et al. The intubation difficulty scale (IDS): proposal and evaluation of a new score characterizing the complexity of endotracheal intubation. *Anesthesiology* 1997;**87**(6):1290-7. [PUBMED: 9416711]

Al-Niaimi 2009

Al-Niaimi A, Safdar N. Supplemental perioperative oxygen for reducing surgical site infection: a meta-analysis. *Journal of Evaluation of Clinical Practice* 2009;**15**(2):360-5. [PUBMED: 19335497]

Altman 2003

Altman DG, Bland JM. Interaction revisited: the difference between two estimates. *BMJ* 2003;**326**(7382):219. [PUBMED: 12543843]

Appiah-Ankam 2004

Appiah-Ankam J, Hunter JM. Pharmacology of neuromuscular blocking drugs. *Continuing Education in Anaesthesia, Critical Care & Pain* 2004;**4**(1):2-7. [DOI: [10.1093/bjaceaccp/mkh002](https://doi.org/10.1093/bjaceaccp/mkh002)]

ASA guideline 1993

ASA guideline 1993. Practice guidelines for management of the difficult airway. A report by the American Society of Anesthesiologists Task Force on Management of the Difficult Airway. *Anesthesiology* 1993;**78**(3):597-602. [PUBMED: 8457062]

Baillard 2005

Baillard C, Adnet F, Borron SW, Racine SX, Ait Kaci F, Fournier JL, et al. Tracheal intubation in routine practice with and without muscular relaxation: an observational study. *European Journal of Anaesthesiology* 2005;**22**(9):672-7. [PUBMED: 16163913]

Bekelman 2003

Bekelman JE, Li Y, Gross CP. Scope and impact of financial conflicts of interest in biomedical research: a systematic review. *JAMA* 2003;**289**(4):454-65. [PUBMED: 12533125]

Bouvet 2009

Bouvet L, Stoian A, Rimmelé T, Allaouchiche B, Chassard D, Boselli E. Optimal remifentanyl dosage for providing excellent intubating conditions when co-administered with a single standard dose of propofol. *Anaesthesia* 2009;**64**(7):719-26. [PUBMED: 19624626]

Bowman 2006

Bowman WC. Neuromuscular block. *British Journal of Pharmacology* 2006;**147** Suppl 1:277-86. [PUBMED: 16402115]

Braz 2009

Braz LG, Braz DG, da Cruz DS, Fernandes LA, Modolo NS, Braz JR. Mortality in anesthesia: a systematic review. *Clinics (Sao Paulo, Brazil)* 2009;**64**(10):999-1006. [PUBMED: 19841708]

Brok 2008

Brok J, Thorlund K, Gluud C, Wetterslev J. Trial sequential analysis reveals insufficient information size and potentially false positive results in many meta-analyses. *Journal of Clinical Epidemiology* 2008;**61**(8):763-9. [PUBMED: 18411040]

Brok 2009

Brok J, Thorlund K, Wetterslev J, Gluud C. Apparently conclusive meta-analyses may be inconclusive - trial sequential analysis adjustment of random error risk due to repetitive testing of accumulating data in apparently conclusive neonatal meta-analyses. *International Journal of Epidemiology* 2009;**38**(1):287-98. [PUBMED: 18824466]

Chura 2007

Chura JC, Boyd A, Argenta PA. Surgical site infections and supplemental perioperative oxygen in colorectal surgery patients: a systematic review. *Surgical Infection* 2007;**8**(4):455-61. [PUBMED: 17883362]

Cooper 2008

Cooper GM, McClure JH. Anaesthesia chapter from Saving Mothers' Lives; reviewing maternal deaths to make pregnancy safer. *British Journal of Anaesthesia* 2008;**100**(1):17-22. [PUBMED: 18070784]

Cormack 1984

Cormack RS, Lehane J. Difficult tracheal intubation in obstetrics. *Anaesthesia* 1984;**39**(11):1105-11. [PUBMED: 6507827]

Dickersin 1990

Dickersin K. The existence of publication bias and risk factors for its occurrence. *JAMA* 1990;**263**(10):1385-9. [PUBMED: 2406472]

Domino 1999

Domino KB, Posner KL, Caplan RA, Cheney FW. Airway injury during anesthesia: a closed claims analysis. *Anesthesiology* 1999;**91**(6):1703-11. [PUBMED: 10598613]

Egger 1997

Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997;**315**(7109):629-34. [PUBMED: 9310563]

El-Boghdadly 2016

El-Boghdadly K, Bailey CR, Wiles MD. Postoperative sore throat: a systematic review. *Anaesthesia* 2016;**71**(6):706-17. [PUBMED: 27158989]

el-Ganzouri 1996

el-Ganzouri AR, McCarthy RJ, Tuman KJ, Tanck EN, Ivankovich AD. Preoperative airway assessment: predictive value of a multivariate risk index. *Anesthesia and Analgesia* 1996;**82**(6):1197-204. [PUBMED: 8638791]

Fotopoulou 2012

Fotopoulou G, Theocharis S, Vasileiou I, Kouskouni E, Xanthos T. Management of the airway without the use of neuromuscular blocking agents: the use of remifentanyl. *Fundamental & Clinical Pharmacology* 2012;**26**(1):72-85. [PUBMED: 21771056]

Fuchs-Buder 2007

Fuchs-Buder T, Claudius C, Skovgaard LT, Eriksson LI, Mirakhur RK, Viby-Mogensen J. Good clinical research practice in pharmacodynamic studies of neuromuscular blocking agents II: the Stockholm revision. *Acta Anaesthesiologica Scandinavica* 2007;**51**(7):789-808. [PUBMED: 17635389]

Gamble 2005

Gamble C, Hollis S. Uncertainty method improved on best-worst case analysis in a binary meta-analysis. *Journal of Clinical Epidemiology* 2005;**58**(6):579-88. [PUBMED: 15878471]

Guyatt 2011

Guyatt GH, Oxman AD, Kunz R, Brozek J, Alonso-Coello P, Rind D, et al. GRADE guidelines 6. Rating the quality of evidence - imprecision. *Journal of Clinical Epidemiology* 2011;**64**(12):1283-93. [PUBMED: 21839614]

Guyatt 2013

Guyatt GH, Oxman AD, Santesso N, Helfand M, Vist G, Kunz R, et al. GRADE guidelines: 12. Preparing summary of findings tables - binary outcomes. *Journal of Clinical Epidemiology* 2013;**66**(2):158-72. [PUBMED: 22609141]

Hellbo-Hansen 1988

Helbo-Hansen S, Ravlo O, Trap-Andersen S. The influence of alfentanil on the intubating conditions after priming with vecuronium. *Acta Anaesthesiologica Scandinavica* 1988;**32**(1):41-4. [PUBMED: 2894106]

Henderson 2004

Henderson JJ, Popat MT, Latto IP, Pearce AC, Difficult Airway Society. Difficult Airway Society guidelines for management of the unanticipated difficult intubation. *Anaesthesia* 2004;**59**(7):675-94. [PUBMED: 15200543]

Higgins 2002

Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Statistics in Medicine* 2002;**21**(11):1539-58. [PUBMED: 12111919]

Higgins 2003

Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ* 2003;**327**(7414):557-60. [PUBMED: 12958120]

Higgins 2011

Higgins JPT, Green S, editors. Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011. www.cochrane-handbook.org.

Hollis 1999

Hollis S, Campbell F. What is meant by intention to treat analysis? Survey of published randomised controlled trials. *BMJ* 1999;**319**(7211):670-4. [PUBMED: 10480822]

Hove 2007

Hove LD, Steinmetz J, Christoffersen JK, Moller A, Nielsen J, Schmidt H. Analysis of deaths related to anesthesia in the period 1996-2004 from closed claims registered by

the Danish Patient Insurance Association. *Anesthesiology* 2007;**106**(4):675-80. [PUBMED: 17413904]

Jaensson 2010

Jaensson M, Olowsson LL, Nilsson U. Endotracheal tube size and sore throat following surgery: a randomized-controlled study. *Acta Anaesthesiologica Scandinavica* 2010;**54**(2):147-53. [PUBMED: 19930246]

Jaensson 2012

Jaensson M, Gupta A, Nilsson UG. Gender differences in risk factors for airway symptoms following tracheal intubation. *Acta Anaesthesiologica Scandinavica* 2012;**56**(10):1306-13. [PUBMED: 22998099]

Keus 2009

Keus F, Weterslev J, Gluud C, Gooszen HG, van Laarhoven CJHM. Robustness assessments are needed to reduce bias in meta-analyses including zero-event randomized trials. *American Journal of Gastroenterology* 2009;**104**(3):546-51. [PUBMED: 19262513]

Krieg 1980

Krieg N, Mazur L, Booij LH, Crul JF. Intubation conditions and reversibility of a new non-depolarizing neuromuscular blocking agent, Org-NC45. *Acta Anaesthesiologica Scandinavica* 1980;**24**(5):423-5. [PUBMED: 6110303]

L'Hermite 2009

L'Hermite J, Nouvellon E, Cuvillon P, Fabbro-Peray P, Langeron O, Ripart J. The Simplified Predictive Intubation Difficulty Score: a new weighted score for difficult airway assessment. *European Journal of Anaesthesiology* 2009;**26**(12):1003-9. [PUBMED: 19593145]

Lan 1983

Lan KKG, DeMets DL. Discrete sequential boundaries for clinical trials. *Biometrika* 1983;**70**:659-63.

Lund 1969

Lund I, Stovner J. Dose-response curves for tubocurarine, alcuronium and pancuronium. *Acta Anaesthesiologica Scandinavica* 1969;**37**:238-42. [PUBMED: 4246294]

Lundstrom 2009a

Lundstrom LH, Moller AM, Rosenstock C, Astrup G, Gatke MR, Weterslev J. Avoidance of neuromuscular blocking agents may increase the risk of difficult tracheal intubation: a cohort study of 103,812 consecutive adult patients recorded in the Danish Anaesthesia Database. *British Journal of Anaesthesia* 2009;**103**(2):283-90. [PUBMED: 19457894]

Lundstrom 2009b

Lundstrom LH, Moller AM, Rosenstock C, Astrup G, Gatke MR, Weterslev J. A documented previous difficult tracheal intubation as a prognostic test for a subsequent difficult tracheal intubation in adults. *Anaesthesia* 2009;**64**(10):1081-8. [PUBMED: 19735399]

Lundstrom 2009c

Lundstrom LH, Moller AM, Rosenstock C, Astrup G, Wetterslev J. High body mass index is a weak predictor for difficult and failed tracheal intubation: a cohort study of 91,332 consecutive patients scheduled for direct laryngoscopy registered in the Danish Anesthesia Database. *Anesthesiology* 2009;**110**(2):266-74. [PUBMED: 19194154]

Lundstrøm 2011b

Lundstrøm LH, Vester-Andersen M, Møller AM, Charuluxananan S, L'hermite J, Wetterslev J, et al. Poor prognostic value of the modified Mallampati score: a meta-analysis involving 177 088 patients. *British Journal of Anaesthesia* 2011;**107**(5):659-67. [PUBMED: 21948956]

Nørskov 2015

Nørskov AK, Rosenstock CV, Wetterslev J, Astrup G, Afshari A, Lundstrøm LH. Diagnostic accuracy of anaesthesiologists' prediction of difficult airway management in daily clinical practice: a cohort study of 188 064 patients registered in the Danish Anaesthesia Database. *Anaesthesia* 2015;**70**(3):272-81. [PUBMED: 25511370]

Nørskov 2016

Nørskov AK, Wetterslev J, Rosenstock CV, Afshari A, Astrup G, Jakobsen JC, et al. Effects of using the simplified airway risk index vs usual airway assessment on unanticipated difficult tracheal intubation - a cluster randomized trial with 64,273 participants. *British Journal of Anaesthesia* 2016;**116**(5):680-9. [PUBMED: 27106972]

Peterson 2005

Peterson GN, Domino KB, Caplan RA, Posner KL, Lee LA, Cheney FW. Management of the difficult airway: a closed claims analysis. *Anesthesiology* 2005;**103**(1):33-9. [PUBMED: 15983454]

Pogue 1997

Pogue JM, Yusuf S. Cumulating evidence from randomized trials: utilizing sequential monitoring boundaries for cumulative meta-analysis. *Controlled Clinical Trials* 1997;**18**(6):580-93. [PUBMED: 9408720]

Pogue 1998

Pogue J, Yusuf S. Overcoming the limitations of current meta-analysis of randomised controlled trials. *Lancet* 1998;**351**(9095):47-52. [PUBMED: 9433436]

Qadan 2009

Qadan M, Akça O, Mahid SS, Hornung CA, Polk HC Jr. Perioperative supplemental oxygen therapy and surgical site infection: a meta-analysis of randomized controlled trials. *Archives of Surgery* 2009;**144**(4):359-66. [PUBMED: 19380650]

RevMan 5.3 [Computer program]

The Nordic Cochrane Centre, The Cochrane Collaboration. Review Manager (RevMan).. Version 5.3. Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration, 2011.

Rosenstock 2001

Rosenstock C, Moller J, Hauberg A. Complaints related to respiratory events in anaesthesia and intensive care medicine

from 1994 to 1998 in Denmark. *Acta Anaesthesiologica Scandinavica* 2001;**45**(1):53-8. [PUBMED: 11152034]

Rosenstock 2004

Rosenstock C, Ostergaard D, Kristensen MS, Lippert A, Ruhnau B, Rasmussen LS. Residents lack knowledge and practical skills in handling the difficult airway. *Acta Anaesthesiologica Scandinavica* 2004;**48**(8):1014-8. [PUBMED: 15315620]

Rosenstock 2006

Rosenstock C, Hansen EG, Kristensen MS, Rasmussen LS, Skak C, Ostergaard D. Qualitative analysis of unanticipated difficult airway management. *Acta Anaesthesiologica Scandinavica* 2006;**50**(3):290-7. [PUBMED: 16480461]

Rücker 2008

Rücker G, Schwarzer G, Carpenter J. Arcsine test for publication bias in meta-analyses with binary outcomes. *Statistics in Medicine* 2008;**27**(5):746-63. [PUBMED: 17592831]

Saarnivaara 1991

Saarnivaara L, Klemola UM. Injection pain, intubating conditions and cardiovascular changes following induction of anaesthesia with propofol alone or in combination with alfentanil. *Acta Anaesthesiologica Scandinavica* 1991;**35**(1):19-23. [PUBMED: 2006594]

Shiga 2005

Shiga T, Wajima Z, Inoue T, Sakamoto A. Predicting difficult intubation in apparently normal patients: a meta-analysis of bedside screening test performance. *Anesthesiology* 2005;**103**(2):429-37. [PUBMED: 16052126]

Stout 1987

Stout DM, Bishop MJ, Dwersteg JF, Cullen BF. Correlation of endotracheal tube size with sore throat and hoarseness following general anesthesia. *Anesthesiology* 1987;**67**(3):419-21. [PUBMED: 3307536]

Sumathi 2007

Sumathi PA, Shenoy T, Ambareesha M, Krishna HM. Controlled comparison between betamethasone gel and lidocaine jelly applied over tracheal tube to reduce postoperative sore throat, cough, and hoarseness of voice. *British Journal of Anaesthesia* 2008;**100**(2):215-8. [PUBMED: 18024955]

Sutton 2002

Sutton AJ, Cooper NJ, Lambert PC, Jones DR, Abrams KR, Sweeting MJ. Meta-analysis of rare and adverse event data. *Expert Review in Pharmacoeconomics Outcomes Research* 2002;**2**(4):367-79. [PUBMED: 19807443]

Sweeting 2004

Sweeting MJ, Sutton AJ, Lambert PC. What to add to nothing? Use of avoidance of continuity corrections in meta-analysis of sparse data. *Statistics in Medicine* 2004;**23**(9):1351-75. [PUBMED: 15116347]

Tanaka 2015

Tanaka Y, Nakayama T, Nishimori M, Tsujimura Y, Kawaguchi M, Sato Y. Lidocaine for preventing postoperative sore throat. *Cochrane Database of Systematic Reviews* 2015, Issue 7. [DOI: [10.1002/14651858.CD004081.pub3](https://doi.org/10.1002/14651858.CD004081.pub3); PUBMED: 26171894]

Thomas 2007

Thomas S, Beevi S. Dexamethasone reduces the severity of postoperative sore throat. *Canadian Journal of Anaesthesia* 2007;**54**(11):897-901. [17975234]

Thorlund 2009

Thorlund K, Devereaux PJ, Wetterslev J, Guyatt G, Ioannidis JP, Thabane L, et al. Can trial sequential monitoring boundaries reduce spurious inferences from meta-analyses?. *International Journal of Epidemiology* 2009;**38**(1):276-86. [PUBMED: 18824467]

Thorlund 2011

Thorlund K, Imberger G, Walsh M, Chu R, Gluud C, Wetterslev J, et al. The number of patients and events required to limit the risk of overestimation of intervention effects in meta-analysis - a simulation study. *PLoS One* 2011;**6**(10):e25491. [PUBMED: 22028777]

Viby-Mogensen 1996

Viby-Mogensen J, Engbaek J, Eriksson LI, Gramstad L, Jensen E, Jensen FS, et al. Good clinical research practice (GCRP) in pharmacodynamic studies of neuromuscular blocking agents. *Acta Anaesthesiologica Scandinavica* 1996;**40**(1):59-74. [PUBMED: 8904261]

Wetterslev 2008

Wetterslev J, Thorlund K, Brok J, Gluud C. Trial sequential analysis may establish when firm evidence is reached in

cumulative meta-analysis. *Journal of Clinical Epidemiology* 2008;**61**(1):64-75. [PUBMED: 18083463]

Wetterslev 2009

Wetterslev J, Thorlund K, Brok J, Gluud C. Estimating required information size by quantifying diversity in random-effects model meta-analyses. *BMC Medical Research Methodology* 2009;**9**(86):1-12. [PUBMED: 20042080]

Woods 2005

Woods AW, Allam S. Tracheal intubation without the use of neuromuscular blocking agents. *British Journal of Anaesthesia* 2005;**94**(2):150-8. [PUBMED: 15516354]

Xu 2012

Xu YJ, Wang SL, Ren Y, Zhu Y, Tan ZM. A smaller endotracheal tube combined with intravenous lidocaine decreases post-operative sore throat - a randomized controlled trial. *Acta Anaesthesiologica Scandinavica* 2012;**56**(10):1314-20. [PUBMED: 22999067]

Yentis 1998

Yentis SM, Lee DJ. Evaluation of an improved scoring system for the grading of direct laryngoscopy. *Anaesthesia* 1998;**53**(11):1041-4. [PUBMED: 10023271]

References to other published versions of this review

Lundstrøm 2011a

Lundstrøm LH, Strande S, Møller AM, Wetterslev J. Use versus avoidance of neuromuscular blocking agent for improving conditions during tracheal intubation or direct laryngoscopy in adults and adolescents. *Cochrane Database of Systematic Reviews* 2011, Issue 7. [DOI: [10.1002/14651858.CD009237](https://doi.org/10.1002/14651858.CD009237)]

CHARACTERISTICS OF STUDIES

Characteristics of included studies [ordered by study ID]

Alexander 1999

Methods	Parallel-group RCT
	Settings: single centre
	Country: UK
	Language: English
	Number of control groups = 1/Number of Intervention groups = 2
	Number of participants in control group = 20/Number of participants in intervention groups = 20/20
	Randomized: N = 60
	Analysed: N = 60
	Dates when the study was conducted: not specified
Participants	Inclusion criteria
	ASA physical status I or II

Alexander 1999 (Continued)

Scheduled for elective surgery

Exclusion criteria

Oesophageal reflux or hiatal hernia

Previous difficulty with intubation or a suspected difficult airway

Allergies to any of the study drugs

Administration of sedative or opioid drugs in the previous 24 hours

Renal or hepatic impairment

Interventions	NMBA Control group S: suxamethonium 1 mg/kg Hypnotic Propofol 2 mg/kg Opioid Intervention group A: alfentanil 50 µg/kg Intervention group R: remifentanil 2 µg/kg Control group S: none Local anaesthetic None Other Premedicated with midazolam 0.03 mg/kg, 10 minutes before induction	
Outcomes	1. Intubation condition: "Good clinical research practice (GCRP) in pharmacodynamic studies of neuromuscular blocking agents" (<i>Viby-Mogensen 1996</i>)	
Notes	One patient in both control group R and control group A had closed vocal cords requiring administration of suxamethonium. Tracheal intubation occurred 60 seconds after administration of the study drug Funding source: not specified Declarations of interest: not specified	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Participants were randomly allocated with a computer-generated table to 1 of 3 groups
Allocation concealment (selection bias)	Unclear risk	Not specified
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not specified
Blinding of outcome assessment (detection bias)	Low risk	The anaesthetist who intubated the participant was unaware of the study drug and was instructed to face away from the participant; the sound of the pulse

Alexander 1999 (Continued)

All outcomes		oximeter was temporarily turned off to avoid recognition of the study drug by observation of fasciculations or a decrease in heart rate
Incomplete outcome data (attrition bias) All outcomes	Low risk	Numbers and reasons for (possible) dropouts and withdrawals are described
Selective reporting (reporting bias)	Low risk	Relevant and reasonably expected outcomes are reported

Barclay 1997

Methods	Parallel-group RCT Settings: single centre Country: UK Language: English Number of control groups = 2/Number of Intervention groups = 1 Number of participants in control groups = 20/20/Number of participants in intervention group = 20 Randomized: N = 60 Analysed: N = 60 Dates when the study was conducted: not specified
Participants	Inclusion criteria Aged 18 to 50 years Exclusion criteria Obese Risk of pulmonary aspiration Smoked more than 10 cigarettes per day Tracheas known to be difficult to intubate or coughing or straining after tracheal intubation
Interventions	NMBA Control group 1: rocuronium 0.1 mg/kg Control group 2: rocuronium 0.3 mg/kg Hypnotic Propofol 2.5 mg/kg Opioid Alfentanil 10 µg/kg Local anaesthetic Lidocaine 10 mg IV Other Premedicated with temazepam 20 to 30 mg per os 1 hour before induction

Barclay 1997 (Continued)

Outcomes

1. Intubation conditions: modified version of "Good clinical research practice (GCRP) in pharmacodynamic studies of neuromuscular blocking agents" (Viby-Mogensen 1996).

The investigator assessed 3 factors: ease of jaw opening and laryngoscopy (1 easy; 2 fair; 3 difficult), position of the vocal cords and their movement (1 open; 2 moving; 3 closing), and degree of straining ("bucking") after tracheal intubation and cuff inflation (1 none; 2 with diaphragm; 3 with abdominal muscles). The occurrence of any significant complication was recorded. Overall conditions for tracheal intubation were scored by 3 grades: optimal, suboptimal, and failure. Tracheal intubation was judged as optimal when all scores were 1 or 2, and was judged as suboptimal if any scores were 3. Failure to intubate was scored as a failure

Notes

Funding source: "We also thank Organa Teknika for their generous supply of rocuronium"

Declarations of interest: not specified

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not specified
Allocation concealment (selection bias)	Unclear risk	Not specified
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not specified
Blinding of outcome assessment (detection bias) All outcomes	Low risk	an investigator, who was blinded as to the allocation....
Incomplete outcome data (attrition bias) All outcomes	Low risk	Numbers and reasons for (possible) dropouts and withdrawals are described
Selective reporting (reporting bias)	Low risk	Relevant and reasonably expected outcomes are reported

Beck 1993

Methods

Parallel-group RCT

Settings: single centre

Country: USA

Language: English

Number of control groups = 1/Number of Intervention groups = 1

Number of participants in control group = 33/Number of participants in intervention group = 31

Randomized: N = 64

Analysed: N = 64

Beck 1993 (Continued)

Dates when the study was conducted: not specified

Participants	<p>Inclusion criteria</p> <p>ASA I or II</p> <p>Aged 18 to 60 years</p> <p>Elective surgery lasting > 20 minutes</p> <p>Exclusion criteria</p> <p>Neurosurgical, obstetrical, or ophthalmological procedures</p>
Interventions	<p>NMBA</p> <p>Suxamethonium 1 mg/kg</p> <p>Hypnotic</p> <p>Control group: thiopenthal 5 mg/kg</p> <p>Intervention group: propofol 2 mg/mL</p> <p>Opioid</p> <p>Control group: none</p> <p>Intervention group: alfentanil 50 µg/kg</p> <p>Local anaesthetic</p> <p>None</p> <p>Other</p> <p>None</p>
Outcomes	<p>1. Intubation condition: VAS 0 to 100, intubation condition: 0 = perfect, 100 = terrible. No defined cut-off value indicating acceptable intubation condition</p> <p>2. Laryngoscopy condition: Cormack and Lehane score (Cormack 1984)</p> <p>In our meta-analyses, we categorized Cormack and Lehane scores I and II as acceptable intubation condition, and scores III and IV as unacceptable condition</p>
Notes	<p>Funding source: not specified</p> <p>Declarations of interest: not specified</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not specified how
Allocation concealment (selection bias)	Unclear risk	Not specified
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not specified

Beck 1993 (Continued)

Blinding of outcome assessment (detection bias) All outcomes	Low risk	The anaesthetist performing the intubation was independent from the study
Incomplete outcome data (attrition bias) All outcomes	Low risk	Numbers and reasons for (possible) dropouts and withdrawals are described
Selective reporting (reporting bias)	Low risk	Relevant and reasonably expected outcomes are reported

Bouvet 2008

Methods	Parallel-group RCT Settings: single centre Country: France Language: English Number of control groups = 1/Number of intervention groups = 1 Number of participants in control group = 64/Number of participants in intervention group = 65 Randomized: N = 130 Analysed: N = 129 Dates when the study was conducted: not specified
Participants	Inclusion criteria Elective gynaecological surgery ASA class I or II Aged > 18 years Exclusion criteria History or evidence of a difficult airway (combination of Mallampati score 3 or 4, thyromental distance < 60 mm, mouth opening < 35 mm) Contraindication to use of NMBA
Interventions	NMBA Cisatracurium 0.15 mg·kg ⁻¹ Hypnotic Propofol 2.5 mg·kg ⁻¹ Opioid Remifentanyl 2 µg·kg ⁻¹ Local anaesthetic None Other

Bouvet 2008 (Continued)

Premedication when appropriate: alprazolam 0.5 mg and/or hydroxyzine 0.5 to 2 mg·kg⁻¹ PO

Outcomes	<p>1. Intubation condition: "Good clinical research practice (GCRP) in pharmacodynamic studies of neuromuscular blocking agents" (Viby-Mogensen 1996)</p> <p>2. Laryngoscopy condition: Cormack and Lehane score (Cormack 1984)</p> <p>3. Postoperative laryngeal symptoms and vocal cord injury: 24 hours, 48 hours, 1 month postoperative</p>
Notes	<p>One participant in group cisatracurium was excluded because of non-observance of the anaesthetic protocol, leaving 129 clinically evaluable participants. Time from induction to start of tracheal intubation: 270 seconds</p> <p>Funding source: "Financial support for this study was provided solely from institutional sources"</p> <p>Declarations of interest: "none declared"</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer
Allocation concealment (selection bias)	Low risk	Allocation concealment was ensured by the use of coded, sealed opaque envelopes
Blinding of participants and personnel (performance bias) All outcomes	High risk	In our protocol, only the study physician (performing all tracheal intubations) was blinded to intervention allocations (along with the participant). Other personnel were aware because they had to await the disappearance of all 4 twitches in response to train-of-four stimulation at the adductor pollicis muscle before allowing arrival of the study physician into the operating room (contacted study author)
Blinding of outcome assessment (detection bias) All outcomes	Low risk	"The same experienced senior physician, blinded to the anaesthetic regimen, performed all tracheal intubations..."
Incomplete outcome data (attrition bias) All outcomes	Low risk	One participant in group cisatracurium was excluded because of non-observance of the anaesthetic protocol
Selective reporting (reporting bias)	Low risk	Relevant and reasonably expected outcomes are reported

Combes 2007

Methods	<p>Parallel-group RCT</p> <p>Settings: single centre</p> <p>Country: France</p> <p>Language: English</p> <p>Number of control groups = 1/Number of intervention groups = 1</p> <p>Number of participants in control group = 150/Number of participants in intervention group = 150</p>
---------	--

Avoidance versus use of neuromuscular blocking agents for improving conditions during tracheal intubation or direct laryngoscopy in adults and adolescents (Review)

34

Combes 2007 (Continued)

Randomized: N = 300

Analysed: N = 299

Dates when the study was conducted: not specified

Participants	<p>Inclusion criteria</p> <p>ASA status I or II</p> <p>Elective surgery</p> <p>Exclusion criteria</p> <p>Predictive of difficult intubation</p> <p>BMI above 30 kg·m⁻²</p> <p>Allergy to muscle relaxants</p> <p>Need for a nasogastric tube</p> <p>Ear-nose-and-throat surgery.</p> <p>Preoperative sore throat or hoarseness at history taking</p>
Interventions	<p>NMBA</p> <p>Rocuronium 0.6 mg·kg⁻¹</p> <p>Hypnotic</p> <p>Propofol 2.5 mg·kg⁻¹</p> <p>Opioid</p> <p>Control group: alfentanil 15 µg/kg</p> <p>Intervention group: alfentanil 40 µg/kg</p> <p>Local anaesthetic</p> <p>None</p> <p>Other</p> <p>Premedication: hydroxyzine 50 to 100 mg PO</p>
Outcomes	<p>1. Intubation condition: Intubation Difficulty Scale (IDS) (Adnet 1997) and "Good clinical research practice (GCRP) in pharmacodynamic studies of neuromuscular blocking agents" (Viby-Mogensen 1996)</p> <p>2. Laryngoscopy condition: Cormack and Lehane score (Cormack 1984)</p> <p>3. Post-intubation pharyngolaryngeal symptoms: 2 hours and 24 hours after extubation. Methods not specified</p>
Notes	<p>Intubation 90 seconds after rocuronium/saline</p> <p>One participant left the hospital before questioning a second time on their pharyngolaryngeal symptoms. This participant (from the rocuronium arm) could not be reached</p> <p>Funding source: "Support was from departmental sources only"</p> <p>Declarations of interest: not specified</p>

Risk of bias

Combes 2007 (Continued)

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Method not specified
Allocation concealment (selection bias)	Low risk	"Patients were randomly assigned to one of the two groups by a physician not involved in the patient's care, using numbered sealed envelopes"
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"All drugs were prepared by an independent staff anaesthetist not involved in the study. The muscle relaxant (rocuronium) and the saline solution were prepared in identical syringes and in identical volumes"
Blinding of outcome assessment (detection bias) All outcomes	Low risk	"All drugs were prepared by an independent staff anaesthetist not involved in the study. The muscle relaxant (rocuronium) and the saline solution were prepared in identical syringes and in identical volumes"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Numbers and reasons for (possible) dropouts and withdrawals are described
Selective reporting (reporting bias)	Low risk	Relevant and reasonably expected outcomes are reported

Dominici 1990

Methods	Parallel-group RCT Settings: single centre Country: France Language: French Number of control groups = 1/Number of intervention groups = 1 Number of participants in control group = 30/Number of participants in intervention group = 30 Randomized: N = 60 Analysed: N = 60 Dates when the study was conducted: not specified
Participants	Inclusion criteria Aged > 18 years ASA I, II, or III Endoscopies with or without laser, tonsillectomies, nose-fracture surgeries, adenectomies, meatotomies and tympanoplastics Surgeries with duration < 90 minutes Exclusion criteria Known allergies Severe arterial hypertension, unstable heart failure

Dominici 1990 (Continued)

Sickness related to conduction and excitability of ventricles

Potassium levels above 5 mmol/L

Renal failure

Epilepsy

Interventions	NMBA Suxamethonium 1.5 mg/kg Hypnotic Propofol 3 mg/mL Opioid Alfentanil 7 to 10µg/kg Local anaesthetic Lidocaine (2%): 1 mL injected before administration of propofol. Topical lidocaine 5% used in larynx before intubation Other Midazolam 0.1 mg/kg was given 30 minutes before induction
Outcomes	1. Intubation condition: quality of intubation measured by (1) position of the glottis (open or semi-open on 1 part and closed or not visualized); (2) presence or absence of bucking; or (3) number of intubation attempts necessary for successful intubation 2. Laryngoscopy condition: position of the glottis (open or semi-open on 1 part and closed or not visualized). Closed or not visualized defined as difficult
Notes	Intubation 60 seconds after induction Funding source: not specified Declarations of interest: not specified

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"...are distributed according to a randomised list into two groups of 30"
Allocation concealment (selection bias)	Unclear risk	Not specified
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not specified
Blinding of outcome assessment (detection bias) All outcomes	Low risk	"...waits outside the operation room for 1 minute (to make sure fasciculations are not seen), before performing the laryngoscopy"
Incomplete outcome data (attrition bias)	Low risk	Numbers and reasons for (possible) dropouts and withdrawals are described

Dominici 1990 (Continued)

All outcomes

Selective reporting (reporting bias)	Low risk	Relevant and reasonably expected outcomes are reported
--------------------------------------	----------	--

González Obregón 2010

Methods	Parallel-group RCT Settings: single centre Country: Columbia Language: Spanish Number of control groups = 1/Number of intervention groups = 1 Number of participants in control group = 50/Number of participants in intervention group = 50 Randomized: N = 100 Analysed: N = 100 Dates when the study was conducted: not specified
Participants	Inclusion criteria Ambulatory surgery ASA physical status I or II Exclusion criteria Aged < 15 or > 60 years Expected difficult airway Full stomach Rapid sequence induction indicated Pregnant or neuromuscular disease
Interventions	NMBA Rocuronium 0.6 mg/kg Hypnotic Intervention group A: oxygen 3 L/min + sevoflurane 3%; after 3 minutes, bolus of propofol 2 mg/kg Control group B: bolus of propofol of 1 to 2 mg/kg (not standardized) for a minute Opioid Intervention group A: remifentanil 0,6 microg/kg/min for 5 minutes, hereafter remifentanil dose is halved until intubation Control group B: bolus of remifentanil 1 to 2 microg/kg (not standardized) for a minute, followed by infusion of 0,15 microg/kg/min for 1 minute Local anaesthetic

González Obregón 2010 (Continued)

None

Other

None

Outcomes	1. Difficult laryngoscopy (Cormack 1984) 2. Post-intubation pharyngolaryngeal symptoms: postoperative hoarseness, not further specified
Notes	Difficult laryngoscopy was included as a surrogate for difficult intubation in our meta-analysis Funding source: not specified Declarations of interest: not specified

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	50 random numbers between 00 and 99 are selected by a manual/hand calculator. These numbers constitute group A, the rest group B. The group that will receive NMBA is then determined by randomization
Allocation concealment (selection bias)	Low risk	The result of randomization is kept in a sealed envelope
Blinding of participants and personnel (performance bias) All outcomes	High risk	The investigator or other anaesthetist who is administering the induction according to randomization is not blinded but is not the one assessing outcomes
Blinding of outcome assessment (detection bias) All outcomes	Low risk	The anaesthesiologist who performs tracheal intubation does not participate in randomization
Incomplete outcome data (attrition bias) All outcomes	Low risk	Numbers and reasons for (possible) dropouts and withdrawals are described
Selective reporting (reporting bias)	Low risk	Relevant and reasonably expected outcomes are reported

Gulhas 2013

Methods	Parallel-group RCT Settings: single centre Country: Turkey Language: English Number of control groups = 1/Number of intervention groups = 1 Number of participants in control group = 40/Number of participants in intervention group = 40 Randomized: N = 80 Analysed: N = 80
---------	---

Gulhas 2013 (Continued)

Dates when the study was conducted: between November 2009 and July 2010

Participants	<p>Inclusion criteria</p> <p>ASA physical status I and II</p> <p>Mallampati scores of I and II</p> <p>Aged 18 to 65 years</p> <p>Selective microlaryngoscopy</p> <p>Exclusion criteria</p> <p>History of head and neck</p> <p>Previous surgery or scheduled to undergo head and neck surgery</p> <p>Severe cardiovascular and pulmonary disease</p> <p>Neuromuscular disease</p> <p>Medications affecting neuromuscular junctions</p> <p>Strained patients</p>
Interventions	<p>NMBA</p> <p>Succinylcholine 1 mg/kg</p> <p>Hypnotic</p> <p>Control group: propofol 2 mg/kg over 30 seconds</p> <p>Intervention group: propofol 2 mg/kg over 30 seconds</p> <p>Opioid</p> <p>Control group: remifentanyl 1 µg/kg over 90 seconds</p> <p>Intervention group: remifentanyl 4 µg/kg over 90 seconds</p> <p>Local anaesthetic</p> <p>None</p> <p>Other</p> <p>None</p>
Outcomes	<p>1. Intubation conditions: categorized by (1) jaw relaxation (complete; tone; stiff; rigid), (2) laryngoscopy (easy; fair; difficult; impossible), (3) vocal cords (open; moving; closing; closed), (4) coughing (none; slight; moderate; severe), (5) movement (none; slight; moderate; severe) (Hellbo-Hansen 1988). In this study, no accumulated score was presented, and laryngoscopy was categorized as difficult; it was impossible to define a difficult intubation</p> <p>2. Post-intubation pharyngolaryngeal symptoms: sore throat and hoarseness evaluated. Methods and timing not described</p>
Notes	<p>Funding source: not specified</p> <p>Declarations of interest: "none"</p>
Risk of bias	

Gulhas 2013 (Continued)

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomization and allocation of participants into intervention groups using computerized numbers
Allocation concealment (selection bias)	Unclear risk	Not specified
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Both care providers on the ward and anaesthesiologists assessing outcomes were blinded to study groups
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Both care providers on the ward and anaesthesiologists assessing outcomes were blinded to study groups
Incomplete outcome data (attrition bias) All outcomes	Low risk	Numbers and reasons for (possible) dropouts and withdrawals are described
Selective reporting (reporting bias)	Low risk	Relevant and reasonably expected outcomes are reported

Hanna 2010

Methods	Parallel-group RCT Settings: single centre Country: USA Language: English Number of control groups = 1/Number of intervention groups = 1 Number of participants in control group = 24/Number of participants in intervention group = 23 Randomized: N = 50 Analysed: N = 47 Dates when the study was conducted: not specified
Participants	Inclusion criteria ASA I or II Aged 18 to 75 years Elective non-ophthalmic surgery Exclusion criteria Anticipated difficult airway management GI reflux, hiatal hernia Ocular surgery within 6 months

Hanna 2010 (Continued)

Long-term opioid use

Allergy to study drugs

Interventions	<p>NMBA</p> <p>Rocuronium 0.06 mg/kg (defasciculation) + succinylcholine 1.5 mg/kg</p> <p>Hypnotic</p> <p>Propofol 2 mg/kg</p> <p>Opioid</p> <p>Control group: none</p> <p>Intervention group: remifentanil 4 µg/kg</p> <p>Local anaesthetic</p> <p>Lidocaine 0.5 mg/kg IV</p> <p>Other</p> <p>Premedication: midazolam 2 mg and glycopyrrolate 0.2 mg IV</p>
Outcomes	<p>1. Intubation condition: "Good clinical research practice (GCRP) in pharmacodynamic studies of neuromuscular blocking agents" (Viby-Mogensen 1996)</p>
Notes	<p>"...Three patients were excluded from the study due to IOP tonometer malfunction...."</p> <p>Intubation 60 seconds after induction</p> <p>Funding source: "supported in part by Aspect Medical Systems, Inc., Norwood, MA, USA, who provided bispectral index (BIS_{tm}) electrodes for the study. Otherwise, the study was supported by funds from the Department of Anesthesiology, Loyola University Medical Center, Maywood, IL, USA"</p> <p>Declarations of interest: "none"</p>
Risk of bias	
Bias	Authors' judgement Support for judgement
Random sequence generation (selection bias)	Unclear risk Not specified
Allocation concealment (selection bias)	Low risk "...each consecutive patient contained in a sealed envelope..."
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk Not specified
Blinding of outcome assessment (detection bias) All outcomes	Low risk "...two attending anesthesiologists and nurse anesthetist, blinded to patient group assignment entered the operation room."
Incomplete outcome data (attrition bias) All outcomes	Low risk "...Three patients were excluded from the study due to IOP tonometer malfunction...."

Hanna 2010 (Continued)

Selective reporting (re- porting bias)	Low risk	Relevant and reasonably expected outcomes are reported
---	----------	--

Harsten 1997

Methods	Parallel-group RCT Settings: single centre Country: Sweden Language: English Number of control groups = 1/Number of intervention groups = 1 Number of participants in control group = 40/Number of participants in intervention group = 39 Randomized: N = 80 Analysed: N = 79 Dates when the study was conducted: not specified
---------	--

Participants	Inclusion criteria ASA I or II Age ranging from 18 to 76 years Exclusion criteria Not specified
--------------	---

Interventions	NMBA Suxamethonium 1 mg/kg Hypnotic Control group: thiopental 5 mg/kg Intervention group: propofol 2,5 mg/kg Opioid Alfentanil 10 µg/kg Local anaesthetic None Other Premedication: 0.25 mg triazolam 45 to 60 minutes before induction
---------------	--

Outcomes	1. Intubation condition: "Intubating conditions were assessed on the basis of jaw relaxation (0 = no relaxation, impossible to open mouth, 1 = moderate relaxation, 2 = complete relaxation), ease of the insertion of the tube (0 = vigorous movements of the vocal cords and difficult or impossible to insert tracheal tube, 1 = slight movements of the vocal cords, 2 = relaxed vocal cords without any movements) and coughing on intubation (0 = vigorous coughing, 1 = slight coughing, 2 = no coughing) (9)" In this RCT, study authors did not define any cut-off value for acceptable intubation. For the meta-analysis, we defined an accumulated score = 4 as clinically unacceptable. We dichotomized the score on the basis of the definition presented in "Good clinical research practice (GCRP) in pharmacodynamic studies of neuromuscular blocking agents" (Viby-Mogensen 1996)
----------	---

Harsten 1997 (Continued)

2. Laryngoscopy condition: Further, patients in whom the vocal cords were not visible (equals Cormach and Lehane grade III and IV) were categorized as clinically unacceptable

Notes

In the PA group, 1 patient was excluded because the induction dose of propofol was not sufficient to produce anaesthesia, and a further 2 patients because the vocal cords were not visible. In our meta-analysis, we included the last 2 patients as "difficult to intubation" and "difficult to laryngoscope"

Funding source: "The authors wish to thank Zeneca, Sweden, for the supply of propofol. This study was supported by grants from the Local Fund for Medical Research and Development of the Kristianstad County Council"

Declarations of interest: not specified

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"According to a computer generated randomisation list, patients were assigned to receive either..."
Allocation concealment (selection bias)	Low risk	"we used sealed non-transparent envelopes containing a slip of paper with a description of which of the two induction methods was being used" (mail contact with study author)
Blinding of participants and personnel (performance bias) All outcomes	High risk	"Apart from the drug administering doctor no one else knew which drugs were given" (mail contact with study author)
Blinding of outcome assessment (detection bias) All outcomes	Low risk	"The intubating anaesthetist was not present during the induction of anaesthesia but was waiting outside the operating room. He was allowed to enter the room when the eyelash reflex or the muscle fasciculations had disappeared and was unaware of which drugs had been used"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Numbers and reasons for (possible) dropouts and withdrawals are described
Selective reporting (reporting bias)	Low risk	Relevant and reasonably expected outcomes are reported

Iamaroon 2001

Methods

Parallel-group RCT

Settings: single centre

Country: Thailand

Language: English

Number of control groups = 1/Number of intervention groups = 1

Number of participants in intervention group = 60/Number of participants in control group = 60

Randomized: N = 120

Analysed: N = 120

lamaroon 2001 (Continued)

Dates when the study was conducted: not specified

Participants	<p>Inclusion criteria</p> <p>ASA I and II</p> <p>Aged 20 to 60 years</p> <p>Elective non-cardiothoracic surgery</p> <p>Exclusion criteria</p> <p>Obesity, body mass index > 30 kg/m²</p> <p>Pregnancy, small bowel obstruction, history of oesophageal reflux, or hiatal hernia</p> <p>Difficult airway problems</p> <p>Hyperkalaemia</p> <p>Suspected malignant hyperthermia</p> <p>Cardiac, pulmonary, or renal disease</p>
Interventions	<p>NMBA</p> <p>Suxametonium 1.5 mg/kg</p> <p>Hypnotic</p> <p>Control group: thiopental 5 mg/kg + 4 L/min (N₂O) and 2 L/min O₂</p> <p>Intervention group: sevoflurane 8% in 66% N₂O and 33% O₂ mixture</p> <p>Opioid</p> <p>Fentanyl 1.5 µg/kg</p> <p>Local anaesthetic</p> <p>None</p> <p>Other</p> <p>Diazepam 5 or 10 mg orally 1 to 2 hours before induction</p>
Outcomes	<p>1. Intubating conditions: Jaw relaxation was described as fully relaxed (score = 1), mildly resistant (score = 2), tight but open (score = 3), and impossible (score = 4). Vocal cord position was described as widely open (score = 1), mid position (score = 2), moving but open (score = 3), and closed (score = 4). Intubating responses were described as none (score = 1), diaphragmatic movement (score = 2), mild/moderate coughing (score = 3), and severe coughing (score = 4). Intubating conditions were graded as excellent (total score (TS) = 3), good (TS = 4 to 6), poor (TS = 7 to 9), or impossible (TS = 10 to 12). Total score of 6 or less was classified as an acceptable intubation condition, otherwise as an unacceptable condition</p>
Notes	<p>Funding source: supported by Mahidol University research fund. Sevoflurane was supported by Abbott Laboratories Limited</p> <p>Declarations of interest: not specified</p>
Risk of bias	
Bias	<p>Authors' judgement</p> <p>Support for judgement</p>

Iamaroon 2001 (Continued)

Random sequence generation (selection bias)	Unclear risk	Not specified
Allocation concealment (selection bias)	Unclear risk	Not specified
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not specified
Blinding of outcome assessment (detection bias) All outcomes	Low risk	"..... anaesthetic residents blindly participated as the intubators and the nurse anesthetists blindly participated as the observers...."
Incomplete outcome data (attrition bias) All outcomes	Low risk	Numbers and reasons for (possible) dropouts and withdrawals are described
Selective reporting (reporting bias)	Low risk	Relevant and reasonably expected outcomes are reported

Isesele 2012

Methods	Parallel-group RCT Settings: single centre Country: Nigeria Language: English/French Number of control groups = 1/Number of intervention groups = 1 Number of participants in control group = 48/Number of participants in intervention group = 48 Randomized: N = 96 Analysed: N = 88 Dates when the study was conducted: not specified
Participants	Inclusion criteria ASA I or II Aged 18 to 45 years Mallampati class I and II Elective surgery Exclusion criteria Inability to understand written or verbal information Presence of dental crowns Past history of difficult intubation Obvious signs of expected difficult intubation

Isesele 2012 (Continued)

Intercurrent chronic ailments such as valvular heart disease and asthma

Overt or risk of raised intracranial or intraocular pressure

Risk of regurgitation and aspiration of gastric contents

Antipsychotic therapy

Opioid therapy

Interventions	NMBA Suxametonium 1.5 mg/kg Hypnotic Propofol 2mg/kg Opioid None Local anaesthetic Intervention group: 1.5 mg/kg intravenous lidocaine Other Diazepam 10 mg orally the morning of surgery
---------------	---

Outcomes	1. Intubation conditions: With the intubating condition scoring system, 3 parameters were assessed and scored on a scale of 0 to 2. These included jaw relaxation, ease of insertion of endotracheal tube, and response to intubation. Thus, a total intubating condition score could range from 0 (worst) to 6 (best). A score of 5 to 6 = good, 3 to 4 = moderate, and 0 to 2 = poor. Scoring system from Saarnivaara 1991
----------	---

Notes	Funding source: none Declarations of interest: none
-------	--

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	All eligible participants were randomly assigned to the 2 study groups with sealed unmarked envelopes
Allocation concealment (selection bias)	Low risk	All eligible participants were randomly assigned to the 2 study groups with sealed unmarked envelopes
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not specified
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not specified
Incomplete outcome data (attrition bias) All outcomes	Low risk	Eight participants, however, were excluded from further participation in the study for various reasons: 5 of these for improper documentation of the data collection form, 2 for electrocardiographic abnormalities, and 1 for abnormal haemodynamic values before induction of anaesthesia

Isesele 2012 (Continued)

Selective reporting (re-reporting bias)	Low risk	Relevant and reasonably expected outcomes are reported
---	----------	--

Jiao 2014

Methods	Parallel-group RCT Settings: single centre Country: China Language: English Number of control groups = 1/Number of intervention groups = 1 Number of participants in control group = 27/Number of participants in intervention group = 28 Randomized: N = 55 Analysed: N = 55 Dates when the study was conducted: not specified
Participants	Inclusion criteria ASA I or II Aged 18 to 60 years Elective gynaecological laparoscopic Exclusion criteria BMI mass index > 30 Mallampati airway grade 2 to 4 Known allergy to propofol, egg, or opioids Alcohol or drug abuse History of gastro-oesophageal reflux disease Neuromuscular disease History of upper respiratory tract infection or other airway hyperactivity disease in the recent 2 weeks
Interventions	NMBA Suxametonium 0.6 mg/kg Hypnotic Propofol 2 mg/kg Opioid Control group: remifentanyl 1.0 µg/kg Intervention group: remifentanyl 1.5 µg/kg Local anaesthetic None

Jiao 2014 (Continued)

Other

None

Outcomes	1. Intubation condition: "Good clinical research practice (GCRP) in pharmacodynamic studies of neuromuscular blocking agents" (Viby-Mogensen 1996)
Notes	1 participant in group C was graded as 4 by Cormach-Lehane grading and was not improved in laryngeal exposure after addition of succinylcholine, then was transferred for video laryngoscopic intubation. We included this participant as difficult to intubate in our meta-analyses. Additional succinylcholine was administered for 1 participant in the control group before intubation. This participant was included in the meta-analyses as difficult to intubate Funding source: not specified Declarations of interest: "none"

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	...with random number table
Allocation concealment (selection bias)	Low risk	...given random numbers by a study co-ordinator, who also encodes the drugs with matching random numbers. The study co-ordinator also prepared all drugs..... (contact with study author)
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Remifentanyl in each group was diluted with normal saline to 10 mL, and injectors were labelled with "2". The injectors were infused with succinylcholine (diluted with normal saline to 10 mL) in group S and with normal saline 10 mL in group C (labelled with "3")
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Intubation was attempted by an experienced anaesthesiologist blinded to grouping of the study
Incomplete outcome data (attrition bias) All outcomes	Low risk	Numbers and reasons for (possible) dropouts and withdrawals are described
Selective reporting (reporting bias)	Low risk	Relevant and reasonably expected outcomes are reported

Kahwaji 1997

Methods	Parallel-group RCT Settings: multi-centre Country: USA Language: English Number of control groups = 5/Number of intervention groups = 1 Number of participants in control group = 30/27/32/28/29/Number of participants in intervention group = 30 Randomized: N = 181
---------	--

Avoidance versus use of neuromuscular blocking agents for improving conditions during tracheal intubation or direct laryngoscopy in adults and adolescents (Review)

Copyright © 2017 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

Kahwaji 1997 (Continued)

Analysed: N = 178

Dates when the study was conducted: not specified

Participants	<p>Inclusion criteria</p> <p>Aged 19 to 85 years</p> <p>ASA physical status I to III</p> <p>Elective surgical procedures with anticipated duration $\geq$ 1 hour</p> <p>Exclusion criteria</p> <p>Significant neurological, renal, or hepatic disease</p> <p>Receiving drugs that could interfere with normal neuromuscular function</p> <p>Preoperative evaluation indicated that difficult tracheal intubation was anticipated</p>
Interventions	<p>NMBA</p> <p>Type: ORG 9487 (rapacuronium)</p> <p>Control group 1: 0.5 mg/kg</p> <p>Control group 2: 1.0 mg/kg</p> <p>Control group 3: 1.5 mg/kg</p> <p>Control group 4: 2.0 mg/kg</p> <p>Control group 5: 2.5 mg/kg</p> <p>Hypnotic</p> <p>Thiopental (3 to 6 mg/kg)</p> <p>Opioid</p> <p>Fentanyl 0.5 to 3 μg/kg</p> <p>Local anaesthetic</p> <p>None</p> <p>Other</p> <p>None</p>
Outcomes	<p>1. Intubation conditions: Intubating conditions were assessed by a blind observer, using a 4-point scale (excellent, good, poor, impossible) - modification of "Good clinical research practice (GCRP) in pharmacodynamic studies of neuromuscular blocking agents" (Viby-Mogensen 1996)</p> <p>2. Serious adverse experience: defined as an adverse effect that was fatal, life-threatening, or permanently disabling, and required prolonged hospitalization, or was an overdose. Investigators stated whether they considered the adverse effect to be related to administration of ORG 9487</p>
Notes	<p>Three elderly participants were excluded before administration of the study drug owing to equipment failure, contraindicated drug administration, or clinical decision</p> <p>Time from induction to start of tracheal intubation: 90 seconds</p> <p>Adverse event: Two of these events (tachycardia, with heart rate from 85 to 150 bpm, and bronchospasm) occurred in one 29-year-old, 100-kg, ASA physical status I male participant within 30 seconds of administration of 2.0 mg/kg ORG 9487 and were followed by erythema of the arms, shoulders, and face. The bronchospasm was treated with salbutamol, and all symptoms gradually subsided. These events were considered to meet the criteria for serious adverse experiences</p>

Kahwaji 1997 (Continued)

Funding source: supported by Organon Inc., Akzo Nobel Inc., West Orange, NJ

Declarations of interest: not specified

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not specified
Allocation concealment (selection bias)	Unclear risk	Not specified
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not specified
Blinding of outcome assessment (detection bias) All outcomes	Low risk	"...before injection of one of five doses of ORG 9487 (0.5, 1.0, 1.5, 2.0, 2.5 mg/kg) or placebo given in a randomized, blind sequence..."
Incomplete outcome data (attrition bias) All outcomes	Low risk	".....Violations included administration of isoflurane for induction before administration of ORG 9487, administration of enflurane before TOF had returned to 0.7, administration of an incorrect dose of ORG 9487, or allocation to the incorrect age group....."
Selective reporting (reporting bias)	Low risk	Relevant and reasonably expected outcomes are reported

Kirkegaard-Nielsen 1999

Methods	Parallel-group RCT Settings: single centre Country: USA Language: English Number of control groups = 3/Number of intervention groups = 1 Number of participants in control group = 20/20/20/Number of participants in intervention group = 20 Randomized: N = 80 Analysed: N = 80 Dates when the study was conducted: not specified
Participants	Inclusion criteria Adults ASA class I and II Elective surgical procedures Exclusion criteria

Kirkegaard-Nielsen 1999 (Continued)

Aged > 60 yr or < 18 years

Gastro-oesophageal reflux

Weighing > 30% more than ideal body weight

Neuromuscular disease, or undergoing treatment with drugs known to interfere with neuromuscular transmission

Interventions	NMBA Control group 1: rocuronium 0.4 mg/kg Control group 2: rocuronium 0.8 mg/kg Control group 3: rocuronium 1.2 mg/kg Hypnotic Propofol 2 mg/kg Opioid Fentanyl 2 µg/kg Local anaesthetic None Other Premedication: midazolam 2 mg IV
Outcomes	1. Intubation condition: "Good clinical research practice (GCRP) in pharmacodynamic studies of neuromuscular blocking agents" (Viby-Mogensen 1996). Further, if the endotracheal tube was not passed successfully within 30 seconds, i.e. 70 seconds after rocuronium or saline solution administration, this was recorded as a failed intubation
Notes	Intubation 40 seconds after induction Funding source: Support was provided solely by institutional and/or departmental sources Declarations of interest: not specified

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not specified
Allocation concealment (selection bias)	Low risk	The dose each participant received was decided on a random basis by selection of an unmarked envelope containing details of the dose
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not specified
Blinding of outcome assessment (detection bias) All outcomes	Low risk	The investigator performing the intubation and assessing conditions was blinded to the dose of rocuronium administered
Incomplete outcome data (attrition bias)	Low risk	Numbers and reasons for (possible) dropouts and withdrawals are described

Kirkegaard-Nielsen 1999 (Continued)

All outcomes

Selective reporting (reporting bias)	Low risk	Relevant and reasonably expected outcomes are reported
--------------------------------------	----------	--

Kopman 2001

Methods	Parallel-group RCT Settings: single centre Country: USA Language: English Number of control groups = 3/Number of intervention groups = 1 Number of participants in control group = 30/30/30/Number of participants in intervention group = 10 Randomized: N = 100 Analysed: N = 100 Dates when the study was conducted: not specified
---------	---

Participants	Inclusion criteria ASA physical status I or II Aged 18 to 65 years Elective surgical procedures Body mass index ≥ 17.5 and ≤ 27.5 Exclusion criteria Neuromuscular disease
--------------	--

Interventions	NMBA Intervention group 1: rapacuronium 1.0 mg/kg Intervention group 2: rapacuronium 1.2 mg/kg Intervention group 3: rocuronium 0.50 mg/kg Hypnotic Propofol 2.0 mg/kg IV Opioid Alfentanil 12.5 $\mu\text{g/kg}$ Local anaesthetic None Other Midazolam (maximum 2 mg) for induction
---------------	---

Outcomes	1. Intubation condition: "Good clinical research practice (GCRP) in pharmacodynamic studies of neuromuscular blocking agents" (Viby-Mogensen 1996)
----------	---

Kopman 2001 (Continued)

Notes

Tracheal Intubation 75 seconds after induction

After 10 participants had been recruited into the saline group, it became obvious that the induction sequence in this group produced conditions for intubation that were clinically unacceptable in most patients. As a consequence, no further participants were added to this group. Each of the other 3 groups consisted of 30 participants

Funding source: supported, in part, by an unrestricted grant from Organon, Inc., of West Orange, NJ

Declarations of interest: not specified

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"...I generated the random number sequence using a Microsoft Excel spreadsheet. " (contacted study author by mail)
Allocation concealment (selection bias)	High risk	After 10 participants had been recruited into the saline group, it became obvious that the induction sequence in this group produced conditions for intubation that were clinically unacceptable in most patients. As a consequence, no further participants were added to this group. Each of the other 3 groups consisted of 30 participant
Blinding of participants and personnel (performance bias) All outcomes	High risk	After 10 participants had been recruited into the saline group, it became obvious that the induction sequence in this group produced conditions for intubation that were clinically unacceptable in most patients. As a consequence, no further participants were added to this group. Each of the other 3 groups consisted of 30 participants
Blinding of outcome assessment (detection bias) All outcomes	Low risk	"All attempts at intubation were performed by AFK, who was not informed which test drug was administered"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Numbers and reasons for (possible) dropouts and withdrawals are described
Selective reporting (reporting bias)	Low risk	Relevant and reasonably expected outcomes are reported

Lieutaud 2003

Methods

Parallel-group RCT

Settings: single centre

Country: France

Language: English

Number of control groups = 3/Number of intervention groups = 1

Number of participants in control group = 45/48/47/Number of participants in intervention group = 20

Randomized: N = 160

Analysed: N = 160

Lieutaud 2003 (Continued)

Dates when the study was conducted: not specified

Participants	<p>Inclusion criteria</p> <p>ASA physical status I or II</p> <p>Aged 18 to 65 years</p> <p>Scheduled for abdominal or breast surgery</p> <p>Exclusion criteria</p> <p>Abnormal airway</p> <p>Significant cardiovascular, respiratory, hepatic, neuromuscular, or renal disease</p> <p>Administration of any drug known or suspected to interact with neuromuscular transmission</p>
Interventions	<p>NMBA</p> <p>Control groups 'L', 'M', 'H' (atracurium 0.5 mg/kg)</p> <p>Hypnotic</p> <p>Control group 'L' (propofol 1.5 mg/kg)</p> <p>Control group 'M' (propofol 2.0 mg/kg)</p> <p>Control group 'H' (propofol 2.5 mg/kg)</p> <p>Intervention group 'WA' (propofol 2.5 mg/kg)</p> <p>Opioid</p> <p>Fentanyl 3 µm/kg</p> <p>Local anaesthetic</p> <p>None</p> <p>Other</p> <p>None</p>
Outcomes	<p>1. Intubation conditions: The scale distributes intubating conditions into 4 classes (excellent, good, poor, impossible). Intubating conditions were pooled as “clinically acceptable” (excellent or good) or “not clinically acceptable” (poor or impossible) (Krieg 1980). The scoring scale is a composite score based on scores of laryngoscopy, cough, and position of vocal cords</p>
Notes	<p>Intubation 240 after induction in control group and in intervention groups based upon TOF response</p> <p>Funding source: not specified</p> <p>Declarations of interest: not specified</p>
Risk of bias	
Bias	<p>Authors' judgement</p> <p>Support for judgement</p>
Random sequence generation (selection bias)	<p>Low risk</p> <p>Table of randomization (4 groups) was designed to assign 2 participants in each group every 8 inclusions</p>
Allocation concealment (selection bias)	<p>Low risk</p> <p>Allocation of the participant was sealed in an opaque envelope, which was opened upon arrival in the operating room</p>

Lieutaud 2003 (Continued)

Blinding of participants and personnel (performance bias) All outcomes	High risk	"...Laryngoscopy and intubation were performed when all train-of-four responses were abolished at the orbicularis oculi in groups high (H), medium (M) and low (L)....." Hereby, personnel performing the treatment were not blinded
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Same physician, blinded to the anaesthetic procedure, performed all intubations
Incomplete outcome data (attrition bias) All outcomes	Low risk	Study was discontinued in group WA after the first intermediate analysis because the incidence of "clinically not acceptable" (poor and impossible) intubating conditions was unacceptable (13 of 20 patients, 65%)
Selective reporting (reporting bias)	Low risk	Relevant and reasonably expected outcomes are reported

Lowry 1999

Methods	Parallel-group RCT Settings: single centre Country: UK Language: English Number of control groups = 6/Number of intervention groups = 2 Number of participants in control groups = 20/20/20/20/20/10/Number of participants in intervention groups = 10/10 Randomized: N = 140 Analysed: N = 140 Dates when the study was conducted: not specified
Participants	Inclusion criteria Aged 18 to 65 years ASA classes I and II Undergoing elective surgery Exclusion criteria Concurrent medication known to interfere with neuromuscular transmission Weighing > 30% outside the ideal for height Patients with anticipated difficult intubation
Interventions	NMBA Control group 1: rocuronium 0.3 mg/kg Control group 2: rocuronium 0.45 mg/kg Control group 3: rocuronium 0.6 mg/kg

Lowry 1999 (Continued)

Control group 4: rocuronium 0.3 mg/kg

Control group 5: rocuronium 0.45 mg/kg

Control group 6: rocuronium 0.6 mg/kg

Hypnotic

Control groups 1 to 3 and intervention group 1: propofol 2 to 3 mg/kg

Control groups 4 to 6 and Intervention group 2: sevoflurane 8% in oxygen, via a vital capacity technique

Opioid

Fentanyl 1 µm/kg

Local anaesthetic

None

Other

None

Outcomes	1. Intubation condition: "Good clinical research practice (GCRP) in pharmacodynamic studies of neuromuscular blocking agents" (Viby-Mogensen 1996)
Notes	Intubation 60 seconds after induction Funding source: This study was supported by a grant from Abbott (UK) Ltd. Dr Lowry was in receipt of a DHSS (Northern Ireland) Clinical Research Fellowship Declarations of interest: not specified

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"....according to prior computer-generated random allocation...."
Allocation concealment (selection bias)	Unclear risk	Not specified
Blinding of participants and personnel (performance bias) All outcomes	High risk	"...Propofol was administered at a rate of 10–15 mg/sec until the loss of eye-lash reflex. Patients in the sevoflurane group were asked to take vital capacity breaths of sevoflurane 8%...."
Blinding of outcome assessment (detection bias) All outcomes	Low risk	".....by an experienced anaesthetist blinded to both the method of induction used and the dose of rocuronium administered. In order to achieve this the intubator did not enter the operating room until 45 sec after the administration of the muscle relaxant...."
Incomplete outcome data (attrition bias) All outcomes	Low risk	Numbers and reasons for (possible) dropouts and withdrawals are described
Selective reporting (reporting bias)	Low risk	Relevant and reasonably expected outcomes are reported

McNeil 2000

Methods	<p>Parallel-group RCT</p> <p>Settings: single centre</p> <p>Country: UK</p> <p>Language: English</p> <p>Number of control groups = 1/Number of intervention groups = 2</p> <p>Number of participants in control groups = 17/Number of participants in intervention groups = 20/23</p> <p>Randomized: N = 60</p> <p>Analysed: N = 60</p> <p>Dates when the study was conducted: not specified</p>
Participants	<p>Inclusion criteria</p> <p>ASA 1 or 2</p> <p>Non-obese</p> <p>Elective surgery</p> <p>Aged 18 to 65 years</p> <p>Exclusion criteria</p> <p>Obstetrical, neurosurgical, ophthalmic patients</p> <p>Mallampati score > 2</p> <p>Gastro-oesophageal reflux</p>
Interventions	<p>NMBA</p> <p>Succinylcholine 1 mg/kg</p> <p>Hypnotic</p> <p>Propofol 2 mg/kg</p> <p>Opioid</p> <p>Control group: none</p> <p>Intervention group 1: remifentanyl 2 µg/kg</p> <p>Intervention group 2: remifentanyl 4 µg/kg</p> <p>Local anaesthetic</p> <p>None</p> <p>Other</p> <p>After induction, participants were mask ventilated with Sevo 2% + 50% N₂O until end of fasciculation and before laryngoscopy</p>
Outcomes	<p>1. Intubation conditions: Scoring system included jaw mobility, mask ventilation, vocal cord visibility, vocal cord position, and participant movement during intubation (all assessed by a 3-grade grading system)</p> <p>In this RCT, study authors did not define any cut-off value for acceptable intubation. In our meta-analyses, intubation condition was categorized as acceptable if the vocal cords were open or in mid-position. If the vocal cords were closed, it was categorized as unacceptable</p>

McNeil 2000 (Continued)

2. Difficult laryngoscopy: Vocal cord visibility was categorized as (1) vocal cords and arytenoids completely visible, (2) vocal cords and arytenoids partly visible, or (3) vocal cords and arytenoids not seen. Score = 3 was defined as a difficult laryngoscopy

Notes

In all patients, vocal cord was visible (Coemack and Lehane score = 1 and 2)

Intubation 30 seconds after induction of remifentanyl and after fasciculation in intervention group

Funding source: not specified

Declarations of interest: not specified

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not specified
Allocation concealment (selection bias)	Low risk	Participants were randomized into 3 groups by opening unmarked envelopes
Blinding of participants and personnel (performance bias) All outcomes	High risk	"...was initiated with 2% sevoflurane in 50% nitrous oxide in oxygen at a total flow 8 litres min ⁻¹ and continued in group PS until fasciculation had ceased...."
Blinding of outcome assessment (detection bias) All outcomes	Low risk	An experienced blinded anaesthetist took over airway control and attempted tracheal intubation
Incomplete outcome data (attrition bias) All outcomes	Low risk	Numbers and reasons for (possible) dropouts and withdrawals are described
Selective reporting (reporting bias)	Low risk	Relevant and reasonably expected outcomes are reported

Mencke 2003

Methods

Parallel-group RCT

Settings: single centre

Country: Germany

Language: English

Number of control groups = 1/Number of intervention groups = 1

Number of participants in control group = 39(37)/Number of participants in intervention group = 39(36)

Randomized: N = 80

Analysed: N = 78 for laryngoscopy conditions and N = 73 for intubation condition and post-intubation pharyngolaryngeal symptoms

Dates when the study was conducted: not specified

Participants

Inclusion criteria

Avoidance versus use of neuromuscular blocking agents for improving conditions during tracheal intubation or direct laryngoscopy in adults and adolescents (Review)

Copyright © 2017 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

Mencke 2003 (Continued)

ASA status I or II

Aged 18 to 76 years

Elective surgery of the ear

Exclusion criteria

Obesity (defined as weight exceeding 20% of normal weight)

Pregnancy

Suspected to have a difficult airway (i.e. abnormal airway anatomy (Mallampati score 3 or 4); and mouth opening < 3.5 cm or cervical spine disease, difficult intubation (i.e. Cormack and Lehane score ≥ 3))

Pathological findings of the larynx revealed by initial stroboscopic examination the day before surgery

Interventions	NMBA Attracurium 0.5 mg/kg Hypnotic Propofol 2.5 to 3 mg/kg Opioid Fentanyl 2 to 3 µg/kg Local anaesthetic None Other Premedication: -midazolam 7.5 mg 1 hour before, as requested	
Outcomes	1. Intubation condition: "Good clinical research practice (GCRP) in pharmacodynamic studies of neuromuscular blocking agents" (Viby-Mogensen 1996) 2. Post-intubation pharyngolaryngeal symptoms: Postoperative hoarseness was assessed at 24, 48, and 72 hours by a standardized interview. Vocal cords were examined by stroboscopy before and 24 and 72 hours after surgery 3. Laryngoscopy condition: It was possible to retrieve information on the laryngoscopy categorized by Cormack and Lehane (Cormack 1984)	
Notes	Five participants were excluded from analyses of intubation conditions and post-intubation upper airway symptoms because of a Cormack grade of 3 or greater (1 in the atracurium group and 4 in the saline group). Moreover, 1 participant in each group had unexpected surgery of the pharynx and therefore had to be excluded Time from induction to start of tracheal intubation: 180 seconds Funding source: Support was provided solely by institutional and departmental sources Declarations of interest: not specified	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Participants were randomized to 2 groups of 40 participants each, via random number draws

Mencke 2003 (Continued)

Allocation concealment (selection bias)	Low risk	Concealment was ensured. The investigator in the operating theatre was blinded; the investigator performing the interview concerning hoarseness was blinded, too. The ENT physician, who performed the stroboscopy, did not know the participant's group. Participants were blinded throughout the study (contacted study author)
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"The study drugs were administered in a double-blind fashion, and syringes were prepared (adjusted to a 5-ml volume) by an investigator who did not participate in the evaluation of intubating conditions, intubating score, and assessment.."
Blinding of outcome assessment (detection bias) All outcomes	Low risk	"The study drugs were administered in a double-blind fashion, and syringes were prepared (adjusted to a 5-ml volume) by an investigator who did not participate in the evaluation of intubating conditions, intubating score, and assessment .."
Incomplete outcome data (attrition bias) All outcomes	Low risk	Numbers and reasons for (possible) dropouts and withdrawals are described
Selective reporting (reporting bias)	Low risk	Relevant and reasonably expected outcomes are reported

Mencke 2014

Methods	Parallel-group RCT Settings: single centre Country: Germany Language: English Number of control groups = 1/Number of intervention groups = 1 Number of participants in control group = 40(40)/Number of participants in intervention group = 43(39) Randomized: N = 83 Analysed: N = 83 for laryngoscopy conditions and N = 79 for intubation condition and post-intubation pharyngolaryngeal symptoms Dates when the study was conducted: Between April 2012 and January 2013
Participants	Inclusion criteria ASA status I or II Aged 18 to 80 years Surgery of the ear Exclusion criteria Known or suspected difficult airway, such as mouth opening < 3.5 cm or Mallampati score 4 or Cormack grade 3 or 4 Obesity Disease of the larynx or vocal cords

Mencke 2014 (Continued)

	Hoarseness before surgery
	Preexisting severe vocal cord pathology
Interventions	<p>NMBA</p> <p>Rocuronium 0.45 mg·kg⁻¹</p> <p>Hypnotic</p> <p>Control group: propofol 1.5 mg·kg⁻¹ was given (if necessary, 30 mg was supplemented)</p> <p>Intervention group: propofol 1.5 mg·kg⁻¹ was given (if necessary, 30 mg was supplemented). After propofol, the SEVO group received sevoflurane at an inspired concentration of 3.0 to 3.5 Vol% (fresh gas flow 8 L·min⁻¹). After 2 to 3 minutes, when the end-tidal sevoflurane concentration reached 1.0 MAC (stable for 20 seconds), intubation was performed.</p> <p>Opioid</p> <p>Remifentanyl 0.30 µg·kg·min⁻¹ for 3 minutes</p> <p>Local anaesthetic</p> <p>None</p> <p>Other</p> <p>Midazolam 7.5 mg orally before arrival in the anaesthetic room</p>
Outcomes	<p>1. Intubation conditions: "Good clinical research practice in pharmacodynamic studies of neuromuscular blocking agents II: the Stockholm revision" (Fuchs-Buder 2007)</p> <p>2. Post-intubation vocal cord injury: All participants underwent laryngoscopy by an ENT physician who was blinded to the participant's group. Slight changes, such as erythema, and vocal cord injuries, such as oedema, haematoma, and granuloma, were noted by videolaryngoscopy</p> <p>3. Laryngoscopy condition: It was possible to retrieve information on the laryngoscopy categorized by Cormack and Lehane (Cormack 1984)</p>
Notes	<p>Two participants from the intervention (SEVO) group could be intubated only after administration of rocuronium. The vocal cords were closed and did not open after propofol 30 mg IV; to avoid vocal cord injury, rocuronium 0.45 mg·kg⁻¹ was given. These 2 participants were included in our meta-analyses as difficult to intubate. Other participants from the intervention (SEVO) group had a Cormack and Lehane score of 3. These participants were included in our meta-analyses as difficult to intubate</p> <p>All participants received dexamethasone 4.0 mg IV. This may have reduced the prevalence of postoperative upper airway discomfort/injury</p> <p>Funding source: "There was no funding for any of the authors and there was no funding for the manuscript preparation"</p> <p>Declarations of interest: "None of the authors has any conflict of interest"</p>
Risk of bias	
Bias	Authors' judgement Support for judgement
Random sequence generation (selection bias)	Low risk "...a randomization program was used..."
Allocation concealment (selection bias)	Low risk Concealment was ensured. The investigator in the operating theatre was blinded when assessing intubating conditions. The investigator who performed the interview regarding hoarseness was blinded. The ENT physician,

Mencke 2014 (Continued)

		who performed the endoscopy, did not know the participant's group. Participants were blinded throughout the study (contacted study author)
Blinding of participants and personnel (performance bias) All outcomes	High risk	Anaesthetists and nurses responsible for treatment were aware of the intervention to ensure a safe general anaesthesia if problems had occurred (contacted study author)
Blinding of outcome assessment (detection bias) All outcomes	Low risk	All intubating variables and scores were blinded for randomization (contacted study author) Assessment of upper airway injury was blinded to participants' group assignment (direct from paper)
Incomplete outcome data (attrition bias) All outcomes	Low risk	Numbers and reasons for (possible) dropouts and withdrawals are described
Selective reporting (reporting bias)	Low risk	Relevant and reasonably expected outcomes are reported

Naguib 2003

Methods	Parallel-group RCT Settings: single centre Country: Saudi Arabia Language: English Number of control groups = 3/Number of intervention groups = 1 Number of participants in control group = 50/50/50/Number of participants in intervention group = 50 Randomized: N = 200 Analysed: N = 200 Dates when the study was conducted: not specified
Participants	Inclusion criteria ASA I status Elective procedures Exclusion criteria Neuromuscular, renal, hepatic, or cardiovascular disease Taking any drug known to interfere with neuromuscular function Drug or alcohol abuse Gastro-oesophageal reflux or hiatus hernia Reactive airway disease Allergies to any of the study drugs Administration of sedative or narcotic drugs in the previous 24 hours

Naguib 2003 (Continued)

Renal or hepatic impairment
Anticipated difficult intubation

Interventions	NMBA Control group I: succinylcholine 0.3 mg/kg Control group II: succinylcholine 0.5 mg/kg Control group III: succinylcholine 1.0 mg/kg Hypnotic Propofol 2 mg/kg Opioid Fentanyl 2 µg/kg Local anaesthetic None Other Premedication: 2 mg oral lorazepam 90 minutes before operation	
Outcomes	1. Intubation conditions: acceptable vs unacceptable "GCRP in pharmacodynamic studies of neuromuscular blocking agents" (Viby-Mogensen 1996)	
Notes	Intubation 60 seconds after induction Funding source: Support was provided solely by institutional and/or departmental sources Declarations of interest: not specified	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Random number generation (contacted study author by mail)
Allocation concealment (selection bias)	Low risk	Yes (contacted study author by mail)
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Yes (contacted study author by mail)
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Because observation of fasciculations would identify the drug administered as succinylcholine, the anaesthesiologist performing and grading intubation was positioned with his back to the participant until just before beginning the intubation sequence
Incomplete outcome data (attrition bias) All outcomes	Low risk	Numbers and reasons for (possible) dropouts and withdrawals are described

Naguib 2003 (Continued)

Selective reporting (re-reporting bias)	Low risk	Relevant and reasonably expected outcomes are reported
---	----------	--

Naguib 2006

Methods	<p>Parallel-group RCT</p> <p>Settings: single centre</p> <p>Country: Saudi Arabia</p> <p>Language: English</p> <p>Number of control groups = 5/Number of intervention groups = 1</p> <p>Number of participants in control group = 30/30/30/30/30/Number of participants in intervention group = 30</p> <p>Randomized: N = 180</p> <p>Analysed: N = 180</p> <p>Dates when the study was conducted: not specified</p>
---------	---

Participants	<p>Inclusion criteria</p> <p>ASA physical status I</p> <p>Participants underwent elective procedures</p> <p>Exclusion criteria</p> <p>Neuromuscular, renal, cardiovascular, or hepatic disease</p> <p>Patient was taking any drug known to interfere with neuromuscular function</p> <p>History of drug or alcohol abuse</p> <p>Gastro-oesophageal reflux or hiatal hernia</p> <p>Reactive airway disease</p> <p>Allergies to any of the study drugs</p> <p>Administration of sedative or narcotic drugs in the previous 24 hours</p> <p>Anticipated difficult intubation</p>
--------------	---

Interventions	<p>NMBA</p> <p>Control group 1: succinylcholine 0.3 mg/kg</p> <p>Control group 2: succinylcholine 0.5 mg/kg</p> <p>Control group 3: succinylcholine 1.0 mg/kg</p> <p>Control group 4: succinylcholine 1.5 mg/kg</p> <p>Control group 5: succinylcholine 2.0 mg/kg</p> <p>Hypnotic</p> <p>Propofol 2 mg/kg</p>
---------------	---

Naguib 2006 (Continued)

Opioid

Fentanyl 2 µm/kg

Local anaesthetic

None

Other

None

Outcomes	1. Intubation condition: "Good clinical research practice (GCRP) in pharmacodynamic studies of neuromuscular blocking agents" (Viby-Mogensen 1996)
Notes	Intubation was subsequently performed 60 seconds after succinylcholine administration Funding source: not specified Declarations of interest: not specified

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomization schedule provided in sealed envelopes according to a computer-generated list
Allocation concealment (selection bias)	Low risk	Randomization schedule provided in sealed envelopes according to a computer-generated list
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Yes (contacted study author by mail)
Blinding of outcome assessment (detection bias) All outcomes	Low risk	The anaesthesiologist performing and grading the intubation was positioned with his back to the participant until just before beginning the intubation sequence
Incomplete outcome data (attrition bias) All outcomes	Low risk	Numbers and reasons for (possible) dropouts and withdrawals are described
Selective reporting (reporting bias)	Low risk	Relevant and reasonably expected outcomes are reported

Nimmo 1995

Methods	Parallel-group RCT Settings: single centre Country: USA Language: English Number of control groups = 2/Number of intervention groups = 1 Number of participants in control group = 20/20/Number of participants in intervention group = 20 Randomized: N = 60
---------	---

Nimmo 1995 (Continued)

Analysed: N = 60

Dates when the study was conducted: not specified

Participants	<p>Inclusion criteria</p> <p>ASA I or II</p> <p>Elective oral surgery under general anaesthesia necessitating nasal intubation</p> <p>Exclusion criteria</p> <p>Aged < 16 years or > 50 years</p> <p>Appeared clinically to present difficulty in intubation</p> <p>Anaesthetic technique was unsuitable</p>
Interventions	<p>NMBA</p> <p>Control group 1: suxamethonium 0.25 mg/kg</p> <p>Control group 2: suxamethonium 0.5 mg/kg</p> <p>Hypnotic</p> <p>Propofol 2.5 mg/kg</p> <p>Opioid</p> <p>Alfentanil 15 µg/kg</p> <p>Local anaesthetic</p> <p>None</p> <p>Other</p> <p>Premedication: temazepam 10 mg 1 hour before operation</p>
Outcomes	<p>1. Intubation conditions: jaw and cord relaxation (Young, Clarke & Dundee). Overall intubation conditions (three grade assessment). Incidence of coughing on intubation (four grade assessment), duration of apnoea (Lund 1969)</p> <p>2. Postoperative myalgia and sore throat: incidence of postoperative myalgia (four grade assessment). Later on the day of operation (day 1), participants were interviewed by one of the investigators (not participating in the anaesthetic) regarding muscle pain and sore throat and were given a questionnaire to return 5 days after the operation (day 5) with the same questions. Muscle pain and sore throat were scored as none, mild, moderate, or severe</p>
Notes	<p>In the Intervention group, intubation was unsuccessful in 2 participants because of a combination of poor vision and poor cord relaxation. In both participants, the trachea was successfully intubated after a dose of suxamethonium 50 mg. These two participants were included in the meta-analysis as patients difficult to intubate</p> <p>Postoperative sore throat was present in more than 50% of participants in all 3 groups at both 1 and 5 days after operation. Investigators noted no significant difference in the incidence of sore throat between the 3 groups. However, they did not report the specific number of participants in the 2 groups. Therefore, we did not include this study in the meta-analysis</p> <p>Funding source: not specified</p> <p>Declarations of interest: not specified</p>
Risk of bias	
Bias	<p>Authors' judgement</p> <p>Support for judgement</p>

Nimmo 1995 (Continued)

Random sequence generation (selection bias)	Unclear risk	Not specified
Allocation concealment (selection bias)	Unclear risk	Not specified
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Syringes with medicine were prepared by one of the investigators not participating in the anaesthetic, and 20 participants were allocated randomly to each of the 3 groups
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not specified
Incomplete outcome data (attrition bias) All outcomes	Low risk	Numbers and reasons for (possible) dropouts and withdrawals are described
Selective reporting (reporting bias)	Unclear risk	The exact incidence of sore throat was not reported

Pang 2014

Methods	Parallel-group RCT Settings: single centre Country: China Language: English Number of control groups = 1/Number of intervention groups = 1 Number of participants in control group = 20/Number of participants in intervention group = 20 Randomized: N = 40 Analysed: N = 40 Dates when the study was conducted: between July 2010 and January 2011
Participants	Inclusion criteria Aged 18 to 65 years Body mass index 18.5 to 25 kg/m ² ASA class I or II Mallampati class I or II Elective suspension laryngoscopic excision under intubation without muscular relaxation Exclusion criteria Medical history of myopathy Known allergy to study drugs

Pang 2014 (Continued)

Drug abuse

History of upper respiratory tract infection within 3 weeks of enrolment

Gastrointestinal reflux

Intracranial pathology

Suspected difficult airway

Serious cardiopulmonary or hepatorenal insufficiency

Interventions	NMBA Control group: cisatracurium 0.1 mg/kg Hypnotic Propofol target control Opioid Remifentanyl target control Local anaesthetic Superficial anaesthesia with 10 mg/mL tetracaine Other Midazolam (0.03 mg/kg) was administered for induction of anaesthesia
---------------	---

Outcomes	1. Intubation condition: Hellbo-Hansen 1988 2. Laryngoscopy condition: Cormack-Lehane classification (Cormack 1984)
----------	--

Notes	Funding source: This work was supported by Jilin Provincial Department of Science and Technology under the International Collaborative Initiative (#20110759) Declarations of interest: none declared
-------	--

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	A computer-generated random number table was used to randomly and equally assign participants
Allocation concealment (selection bias)	Unclear risk	Not specified
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	No information on blinding of personnel performing drug administration. Participants were blinded to treatment assignment throughout the study
Blinding of outcome assessment (detection bias) All outcomes	Low risk	An independent anaesthetist performed tracheal intubation
Incomplete outcome data (attrition bias) All outcomes	Low risk	Numbers and reasons for (possible) dropouts and withdrawals are described

Pang 2014 (Continued)

Selective reporting (re-reporting bias)	Low risk	Relevant and reasonably expected outcomes are reported
---	----------	--

Pino 1998

Methods	Parallel-group RCT Settings: single centre Country: USA Language: English Number of control groups = 5/Number of intervention groups = 1 Number of participants in control group = 30/15/14/14/15/Number of participants in intervention group = 10 Randomized: N = 100 Analysed: N = 98 Dates when the study was conducted: not specified
Participants	Inclusion criteria ASA I or II Normal upper airway anatomy Aged 18 to 65 years Within 30% of ideal body weight Exclusion criteria History of malignant hyperthermia Abnormal plasma cholinesterase levels Neuromuscular, neurological, hepatic, and renal conditions that might influence neuromuscular function Use of drugs that might alter the response to neuromuscular blockade or might affect histamine release
Interventions	NMBA Control group 1: mivacurium 0.25 mg/kg Control group 2: rocuronium 0.45 mg/kg Control group 3: rocuronium 0.6 mg/kg Control group 4: rocuronium 0.9 mg/kg Control group 5: rocuronium 1.2 mg/kg Hypnotic Propofol 2 mg/kg Opioid

Pino 1998 (Continued)

Fentanyl 2 µm/kg

Local anaesthetic

None

Other

Midazolam 1 to 2 mg for induction

Outcomes	1. Intubation conditions <ul style="list-style-type: none"> - Excellent: easy passage of endotracheal tube without coughing; vocal cords relaxed and abducted - Good: passage of endotracheal tube with slight coughing or bucking; vocal cords relaxed and abducted - Poor: passage of endotracheal tube with moderate coughing or bucking; vocal cords moderately abducted - Not possible: unable to intubate
Notes	<p>Two participants were excluded from analysis because they were mistakenly entered into the study twice</p> <p>Intubation 90 seconds after administration of midazolam</p> <p>Funding source: supported by a grant from GlaxoWellcome</p> <p>Declarations of interest: Affiliated GlaxoWellcome Company, Research Triangle Park, North Carolina</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated randomization
Allocation concealment (selection bias)	Unclear risk	Not specified
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not specified
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Blinding of assessor: intubator not involved in protocol; waited outside OP room until last minute before intubation attempt
Incomplete outcome data (attrition bias) All outcomes	Low risk	Two participants were excluded from analysis because they were mistakenly entered into the study twice
Selective reporting (reporting bias)	Low risk	Relevant and reasonably expected outcomes are reported

Rousseau 1995

Methods	Parallel-group RCT Settings: single centre
---------	---

Avoidance versus use of neuromuscular blocking agents for improving conditions during tracheal intubation or direct laryngoscopy in adults and adolescents (Review)

Copyright © 2017 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

Rousseau 1995 (Continued)

Country: France

Language: French

Number of control groups = 1/Number of intervention groups = 1

Number of participants in control group (77)/Number of participants in intervention group = 75

Randomized: N = 152

Analysed: N = 152

Dates when the study was conducted: not specified

Participants	<p>Inclusion criteria</p> <p>> 18 years of age</p> <p>ASA I</p> <p>Mallampati I</p> <p>Elective surgery not needing relaxant</p> <p>Exclusion criteria</p> <p>Suspect difficult intubation</p> <p>Former allergies</p> <p>Abnormal ECG</p> <p>Bradycardia < 50</p> <p>Urgent surgery</p>
Interventions	<p>NMBA</p> <p>Vecuronium 0.08 mg/kg</p> <p>Hypnotic</p> <p>Propofol 2.5 mg/kg</p> <p>Opioid</p> <p>Alfentanil 0.03 mg/kg</p> <p>Local anaesthetic</p> <p>Intervention group: lidocaine 1.5 mg/kg</p> <p>Other</p> <p>Premedication: prazepam 40 mg and hydroxyzine 100 mg (1 hour before induction)</p>
Outcomes	<p>1. Intubation conditions: Physician performing the intubation evaluated mouth opening, opening of the glottis, and the occurrence of coughing (Saarnivaara 1991)</p> <p>Mouth opening: impossible 3, medium 2, complete 0; opening of the glottis: moving or closed 3, half closed 2, open 0; cough at intubation: significant ("Importante") 3, a little 1, absent 0. Total score < 3 considered good intubating conditions. A score is calculated for each intubation attempt</p>
Notes	<p>107 oral intubations, 45 nasal intubations, all intubations by direct laryngoscopy</p> <p>Intubation after induction: 60 seconds in control group and 180 seconds in intervention group</p> <p>Funding source: not specified</p>

Rousseau 1995 (Continued)

Declarations of interest: not specified

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Using a random number table
Allocation concealment (selection bias)	Unclear risk	Not specified
Blinding of participants and personnel (performance bias) All outcomes	High risk	Different times from infusion to intubation in the 2 groups (1 minute for Vecu0 and 3 minutes for Vecu+) make it easy to discern the 2 groups from each other;no mention of measures of concealment of allocation towards assessor, who might have been present from infusion to finish (not mentioned)
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Different times from infusion to intubation in the 2 groups (1 minute for Vecu0 and 3 minutes for Vecu+) make it easy to discern the 2 groups from each other; no mention of measures of concealment of allocation towards assessor, who might have been present from infusion to finish (not mentioned)
Incomplete outcome data (attrition bias) All outcomes	Low risk	Numbers and reasons for (possible) dropouts and withdrawals are described
Selective reporting (reporting bias)	Low risk	Relevant and reasonably expected outcomes are reported

Scheller 1992

Methods	Parallel-group RCT Settings: single centre Country: USA Language: English Number of control groups = 1/Number of intervention groups = 4 Number of participants in control group (15)/Number of participants in intervention group = 15/15/15/15 Randomized: N = 75 Analysed: N = 75 Dates when the study was conducted: not specified
Participants	Inclusion criteria Same-day surgery for a variety of surgical procedures ASA physical status I or II Mallampati class I airway anatomy Exclusion criteria

Scheller 1992 (Continued)

	History of intravenous drug use, alcohol addiction
	Full stomach
	Coronary artery disease
	Reactive airway disease
Interventions	<p>NMBA</p> <p>d-Tubocurarine 3 mg and succinylcholine 1 mg/kg</p> <p>Hypnotic</p> <p>Control group 1: thiamylal 4 mg/kg</p> <p>Intervention groups 1 to 4: propofol 2 mg/kg</p> <p>Opioid</p> <p>Control group: no opioids</p> <p>Intervention group 1: alfentanil 30 µg/kg</p> <p>Intervention group 2: alfentanil 40 µg/kg</p> <p>Intervention group 3: alfentanil 50 µg/kg</p> <p>Intervention group 4: alfentanil 60 µg/kg</p> <p>Local anaesthetic</p> <p>None</p> <p>Other</p> <p>All participants received midazolam 1 mg IV before induction</p>
Outcomes	<p>1. Intubation conditions: Vocal cords, position of vocal cards, jaw mobility, and participant movement during and within 1 minute of attempted intubation of the trachea. Participants whose tracheas could not be intubated after receiving assigned induction drugs were so noted, and succinylcholine (1 mg/kg IV) was administered. Tracheal intubation was then re-attempted, and exposure of the larynx and outcome of intubation attempt were recorded</p> <p>In this RCT, study authors did not define any cut-off value for acceptable intubation and provided no composite measure of the 4 variables. For the meta-analysis, we defined complete or partial exposure of the vocal cords as clinically acceptable. Vocal cord exposure was categorized as: (1) vocal cords and arytenoids completely visible; (2) vocal cords or arytenoids partially visible; or (3) vocal cords or arytenoids not seen</p> <p>2. Laryngoscopy condition: vocal cords or arytenoids not seen defined as difficult laryngoscopy</p>
Notes	<p>Induction after 90 seconds</p> <p>Funding source: Dr Scheller is a recipient of a B.B. Sankey Foundation Award. This work was supported in part by a grant from Janssen Pharmaceutica, Piscataway, NJ</p> <p>Declarations of interest: not specified</p>
Risk of bias	
Bias	<p>Authors' judgement</p> <p>Support for judgement</p>
Random sequence generation (selection bias)	<p>Unclear risk</p> <p>".....randomized to one of five groups by having an assistant pick one of five cards describing the induction sequence....."</p>

Scheller 1992 (Continued)

Allocation concealment (selection bias)	Unclear risk	Not specified
Blinding of participants and personnel (performance bias) All outcomes	High risk	"...Note that we were not able to be completely blinded as to group because we could generally distinguish the thiamylal/succinylcholine group from the others..."
Blinding of outcome assessment (detection bias) All outcomes	High risk	"...Note that we were not able to be completely blinded as to group because we could generally distinguish the thiamylal/succinylcholine group from the others..."
Incomplete outcome data (attrition bias) All outcomes	Low risk	Numbers and reasons for (possible) dropouts and withdrawals are described
Selective reporting (reporting bias)	Low risk	Relevant and reasonably expected outcomes are reported

Schlaich 2000

Methods	Parallel-group RCT Settings: single centre Country: Germany Language: English Number of control groups = 3/Number of intervention groups = 1 Number of participants in control group = 30/30/30/Number of participants in intervention group = 30 Randomized: N = 120 Analysed: N = 120 Dates when the study was conducted: not specified
Participants	Inclusion criteria ASA status I or II Adult Scheduled for elective ambulatory surgery up to 90 minutes Exclusion criteria Pregnancy Neuromuscular disorder or receiving medications known to interact with neuromuscular function Suspected to have a difficult airway
Interventions	NMBA Control group 1 (0.6 mg/kg rocuronium) Control group 2 (0.45 mg/kg rocuronium)

Schlaich 2000 (Continued)

Control group 3 (0.3 mg/kg rocuronium)

Hypnotic

Propofol 2 to 2.5 mg/kg

Opioid

Remifentanyl 0.5 µg/kg/min

Local anaesthetic

None

Other

Premedication: midazolam 7.5 mg orally 1 hour before

Outcomes	1. Intubation condition: "Good clinical research practice (GCRP) in pharmacodynamic studies of neuromuscular blocking agents" (Viby-Mogensen 1996)
Notes	Time from induction to start of tracheal intubation: 180 seconds Funding source: not specified Declarations of interest: not specified

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not specified
Allocation concealment (selection bias)	Unclear risk	Not specified
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not specified
Blinding of outcome assessment (detection bias) All outcomes	Low risk	"....All patients were intubated by the same experienced anaesthetist blinded to the treatment...."
Incomplete outcome data (attrition bias) All outcomes	Low risk	Numbers and reasons for (possible) dropouts and withdrawals are described
Selective reporting (reporting bias)	Low risk	Relevant and reasonably expected outcomes are reported

Sivalingam 2001

Methods	Parallel-group RCT Settings: single centre Country: New Zealand Language: English
---------	--

Sivalingam 2001 (Continued)

Number of control groups = 1/Number of intervention groups = 3

Number of participants in control group = 25/Number of participants in intervention group = 25/25/25/25

Randomized: N = 100

Analysed: N = 100

Dates when the study was conducted: not specified

Participants	<p>Inclusion criteria</p> <p>ASA I or II</p> <p>Aged 18 to 65 years</p> <p>Mixed surgery</p> <p>Exclusion criteria</p> <p>Smokers</p> <p>Known or anticipated difficult tracheal intubation</p> <p>Risk of aspiration</p> <p>Adverse effects of coughing or straining</p> <p>Taking beta-blocker with absolute contraindication to use of sevoflurane, suxamethonium, and alfentanil</p>
Interventions	<p>NMBA</p> <p>Suxamethonium 1 mg/kg</p> <p>Hypnotic</p> <p>Sevoflurane 7% and nitrous oxide 60%</p> <p>Opioid</p> <p>Control group 1: alfentanil 10 µg/kg</p> <p>Intervention group 1: alfentanil 20 µg/kg</p> <p>Intervention group 2: alfentanil 25 µg/kg</p> <p>Intervention group 3: alfentanil 30 µg/kg</p> <p>Local anaesthetic</p> <p>None</p> <p>Other</p> <p>Atropine 0.3 mg</p>
Outcomes	<p>1) Intubation conditions: modified version of Saarnivaara L, Klemola UM (Saarnivaara 1991). The scoring scale is a composite score based on scores of jaw relaxation, movement of limbs, movement of vocal cords, and coughing. Based on a score ranging from 3 to 12 points</p> <p>Intubation conditions were categorized as excellent = 12 points, satisfactory = 10 to 11 points, poor < 10 points or failed</p> <p>2) Sore throat: postoperative, self-reporting, time and method of reporting not defined</p>

Sivalingam 2001 (Continued)

Notes One participant from intervention group 1 was intubated by rescue suxamethonium. We included this participant in our meta-analyses as difficult to intubate

Funding source: not specified

Declarations of interest: not specified

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	"The patients were randomly allocated to receive..."; method not specified
Allocation concealment (selection bias)	Low risk	Sealed opaque envelopes, opened immediately before induction
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not specified
Blinding of outcome assessment (detection bias) All outcomes	Low risk	"...one of the investigators who was blinded to the group allocation, entered the operation room to perform direct laryngoscopy...."
Incomplete outcome data (attrition bias) All outcomes	Low risk	Numbers and reasons for (possible) dropouts and withdrawals are described
Selective reporting (reporting bias)	Low risk	Relevant and reasonably expected outcomes are reported

Stevens 1997

Methods	Parallel-group RCT Settings: single centre Country: USA Language: English Number of control groups = 1/Number of intervention groups = 6 Number of participants in control group = 20/Number of participants in intervention group = 20/20/20/20/20/20 Randomized: N = 140 Analysed: N = 140 Dates when the study was conducted: not specified
Participants	Inclusion criteria ASA I or II Outpatients

Stevens 1997 (Continued)

Aged 18 to 60 years

Scheduled for elective surgery

All enrolled participants had Mallampati class I or II airway anatomy

Exclusion criteria

Coronary artery disease, hypertension, reactive airway disease

Obesity > 30% above ideal body weight

History of drug or alcohol abuse, or gastro-oesophageal reflux

Taking narcotics or drugs known to interfere with neuromuscular transmission

Interventions
NMBA

Control group: d-tubocurarine 3 mg and succinylcholine 1 mg/kg

Hypnotic

Control group: thiopental 4 mg/kg

Intervention groups 1 and 2: etomidate 0.3 mg/kg

Intervention groups 3 and 4: propofol 2 mg/kg

Intervention groups 5 and 6: thiopental 4 mg/kg

Opioid

Control group: none

Intervention groups 1 to 6: alfentanil 40 µg/kg

Local anaesthetic

Lidocaine 1 mg/kg (intervention groups 2, 4, 6)

Other

Premedication: midazolam 0.03 mg/kg IV 5 minutes before induction

Outcomes

1. Intubation conditions: "The intubating anaesthesiologist, assessed each patient on four variables: jaw relaxation, exposure of the vocal cords, vocal cord position, and patient response to intubation and slow (5-s) inflation of the endotracheal tube cuff"

Participants who could not be intubated on the first attempt were given succinylcholine 1 mg/kg, and intubation was completed

In this RCT, study authors did not define any cut-off value for acceptable intubation and provided no composite measure of the 4 variables. For the meta-analysis, we defined closed vocal cords as difficult intubation, and open or midline as acceptable intubation conditions

2. Laryngoscopy conditions: Exposure of the vocal cords was defined as 'complete', 'partial', or 'not seen'. Difficult laryngoscopy was defined as 'not seen'

Notes

For all participants, it was possible to expose the vocal cords. In the various alfentanil groups, 17 required succinylcholine to complete intubation. Thirty-five percent of participants (7 of 20) in intervention group 5 (alfentanil/ thiopental) required succinylcholine to complete intubation compared with 3 of 20 (15%), 2 of 20 (10%), 2 of 20 (10%), and 0% and 3 of 20 (15%) in Intervention groups 1 to 4 and 6, respectively. It was not possible to identify specific participants who had closed vocal cords and therefore required succinyl chloride

Intubation was performed 90 seconds after induction

Stevens 1997 (Continued)

Funding source: not specified
Declarations of interest: not specified

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not specified
Allocation concealment (selection bias)	Low risk	Participants were randomly allocated to 1 of 7 groups (n = 20/group) by means of previously prepared envelopes
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"The induction sequence was conducted using four prepared syringes in all patients".... "Opaque tape was applied to Syringe 3 to disguise the color of the hypnotic drug".... "Injection of all syringes was performed by an assistant behind a drape so that the intubating anesthesiologist (one of three of the authors) was blinded to the color and volume of the IV drugs"
Blinding of outcome assessment (detection bias) All outcomes	Low risk	"The induction sequence was conducted using four prepared syringes in all patients".... "Opaque tape was applied to Syringe 3 to disguise the color of the hypnotic drug".... "Injection of all syringes was performed by an assistant behind a drape so that the intubating anesthesiologist (one of three of the authors) was blinded to the color and volume of the IV drugs"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Numbers and reasons for (possible) dropouts and withdrawals are described
Selective reporting (reporting bias)	Low risk	Relevant and reasonably expected outcomes are reported

Striebel 1995

Methods	Parallel-group RCT Settings: single centre Country: Germany Language: German Number of control groups = 2/Number of intervention groups = 2 Number of participants in control group = 25/25/Number of participants in intervention group = 25/25 Randomized: N = 100 Analysed: N = 100 Dates when the study was conducted: not specified
Participants	Inclusion criteria ASA I or II Gynaecological surgery Exclusion criteria

Striebel 1995 (Continued)

Not specified

Interventions	NMBA Control groups 1 and 2: vecuronium 1 mg + succinylcholine 1 mg/kg Hypnotic Control group 1: demand-adapted thiopental (5.5 ± 3.14 mg/kg) Control group 2: demand-adapted propofol (2.2 ± 0.48 mg/kg) Intervention group 1: demand-adapted propofol (2.4 ± 0.63 mg/kg) Intervention group 2: demand-adapted propofol (2.2 ± 0.48 mg/kg) Opioid Control group 1: fentanyl 0.1 mg Control group 2: fentanyl 0.1 mg Intervention group 1: fentanyl 0.1 mg Intervention group 2: fentanyl 0.2 mg Local anaesthetic None Other Premedication: midazolam 7.5 mg	
Outcomes	1) Intubation conditions: (1) very good, (2) good, (3) satisfactory, (4) sufficient, (5) inadequate, (6) insufficient. Acceptable conditions 1 to 4, unacceptable conditions 5 and 6 2) Laryngoscopy conditions: Cormack & Lehane (Cormack 1984). Unproblematic laryngoscopy: grades I and II. Difficult laryngoscopy: grades III and IV	
Notes	Funding source: not specified Declarations of interest: not specified	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not specified
Allocation concealment (selection bias)	Unclear risk	Not specified
Blinding of participants and personnel (performance bias) All outcomes	Low risk	All medicaments were blinded to personnel
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Anaesthetist performing the intubation was blinded
Incomplete outcome data (attrition bias)	Low risk	Numbers and reasons for (possible) dropouts and withdrawals are described

Avoidance versus use of neuromuscular blocking agents for improving conditions during tracheal intubation or direct laryngoscopy in adults and adolescents (Review)

81

Striebel 1995 (Continued)

All outcomes

Selective reporting (reporting bias)	Low risk	Relevant and reasonably expected outcomes are reported
--------------------------------------	----------	--

Wong 1996

Methods	Parallel-group RCT Settings: single centre Country: Malaysia Language: English Number of control groups = 1/Number of intervention groups = 3 Number of participants in control group = 30/Number of participants in intervention group = 30/30/30 Randomized: N = 120 Analysed: N = 120 Dates when the study was conducted: not specified
---------	--

Participants	Inclusion criteria ASA I or II Elective surgery Aged 18 to 60 years Exclusion criteria Expected difficult airway Aspiration risk Head and neck surgery Suxamethonium contraindicated
--------------	--

Interventions	NMBA Control group: succinylcholine 1 mg/kg Hypnotic Propofol 200 mg/min until loss of verbal response Opioid Control group: none Intervention group 1: alfentanil 15 µg/kg Intervention group 2: alfentanil 30 µg/kg Intervention group 3: none Local anaesthetic None
---------------	---

Wong 1996 (Continued)

Other

Premedication: midazolam 7 mg 1 hour before induction

Outcomes	1. Intubation conditions: The scoring scale is a composite score based on scores of jaw relaxation, movement of the vocal cords, and coughing. Based on a score ranging from 0 to 6 points. Intubation conditions were categorized as good = 5 to 6 points, moderate = 3 to 4 points, poor = 1 to 2 points or failed (Saarnivaara 1991)
Notes	<p>One participant in intervention group 3 was excluded because of an unanticipated difficult intubation. We included this participant in our meta-analysis as a difficult tracheal intubation patient</p> <p>The mean dose requirement for induction by propofol was higher in groups without alfentanil (3 mg/kg vs 2.5 mg/kg)</p> <p>Funding source: "....Janssen Pharmaceutica for supplying the alfentanil"</p> <p>Declarations of interest: not specified</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not specified
Allocation concealment (selection bias)	Unclear risk	Not specified
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"....The drug was given by an anaesthetic trainee who was unaware of the drugs before its administration....."
Blinding of outcome assessment (detection bias) All outcomes	Low risk	".....who was unaware of the randomisation process entered the operating room 30 seconds after completion of drug administration....."
Incomplete outcome data (attrition bias) All outcomes	Low risk	Numbers and reasons for (possible) dropouts and withdrawals are described
Selective reporting (reporting bias)	Low risk	Relevant and reasonably expected outcomes are reported

Yazdi 2016

Methods	<p>Parallel-group RCT</p> <p>Settings: single centre</p> <p>Country: Iran</p> <p>Language: English</p> <p>Number of control groups = 1/Number of intervention groups = 1</p> <p>Number of participants in control group = 31/Number of participants in intervention group = 35</p> <p>Randomized: N = 66</p>
---------	--

Yazdi 2016 (Continued)

Analysed: N = 66

Dates when the study was conducted: not specified

Participants	Inclusion criteria ASA I or II Elective surgery Aged 15 to 65 years Exclusion criteria Expected difficult airway (Mallampati score III or IV) Chronic alcohol or opioid use Allergy to study medications	
Interventions	NMBA Control group: atracurium 0.5 mg/kg Hypnotic Propofol 2.5 mg/kg Opioid Control group: none Intervention group: remifentanil 2 µg/kg IV over 30 seconds Local anaesthetic None Other Atropine 0.5 mg	
Outcomes	1. Intubation conditions: "...anaesthesiologist checked the intubation condition using criteria based on jaw relaxation, vocal cord movement and bucking on tracheal tube as primary measurements of comparison (Table 1). Intubation condition defined as optimal (score 1 or 2 in all categories), suboptimal (score 3 in at least 1 of the 3 categories), or fail (fail to intubation)	
Notes	"Unsuccessful intubation due to closed vocal cords were seen in 2 patients in remifentanil group, the patients ventilated and received atracurium and intubation was applied again" These two participants were categorized as failed intubation in the intervention (remifentanil) group Funding source: not specified Declarations of interest: not specified	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not specified
Allocation concealment (selection bias)	Unclear risk	Not specified

Yazdi 2016 (Continued)

Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Study was categorized as "double-blinded". However, no detailed description was provided in the manuscript
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Study was categorized as "double-blinded". However, no detailed description was provided in the manuscript
Incomplete outcome data (attrition bias) All outcomes	Low risk	Numbers and reasons for (possible) dropouts and withdrawals are described
Selective reporting (reporting bias)	Low risk	Relevant and reasonably expected outcomes are reported

ASA = American Society of Anesthesiologists ASA physical status classification system; BMI = body mass index; ECG = electrocardiography; ENT= ear, nose, and throat; GCRP = "Good clinical research practice (GCRP) in pharmacodynamic studies of neuromuscular blocking agents" (Viby-Mogensen 1996); GI = gastrointestinal; IV = intravenous; kg = kilogram; mg = milligram; mmol/L = millimoles per litre; N = number of cases; NMBA = neuromuscular blocking agent; PO = per os; RCT = randomized controlled trial; sec = second; TOF = 'train of four'; TS = total score; µg = microgram; VAS = visual analogue scale.

Characteristics of excluded studies [ordered by study ID]

Study	Reason for exclusion
Alcock 1993	An unspecified number of participants were intubated blind nasal
Baumgarten 1988	Study was terminated for the control group because of unacceptable intubation conditions. Thus, randomization and blinding were violated
Ide 2015	Participants were intubated blind nasal. No evaluation of direct laryngoscopy was performed

DATA AND ANALYSES
Comparison 1. Avoidance vs use of NMBA

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Difficult tracheal intubation: low risk of bias vs high or uncertain risk of bias	34	3565	Risk Ratio (M-H, Random, 95% CI)	5.00 [3.49, 7.15]
1.1 Low risk of bias	4	508	Risk Ratio (M-H, Random, 95% CI)	13.27 [8.19, 21.49]
1.2 High or uncertain risk of bias	30	3057	Risk Ratio (M-H, Random, 95% CI)	4.14 [2.92, 5.87]
2 Difficult tracheal intubation: depolarizing vs non-depolarizing NMBA	32	3413	Risk Ratio (M-H, Random, 95% CI)	5.25 [3.61, 7.63]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
2.1 Depolarizing NMBA	16	1540	Risk Ratio (M-H, Random, 95% CI)	5.79 [2.64, 12.72]
2.2 Non-depolarizing NMBA	16	1873	Risk Ratio (M-H, Random, 95% CI)	4.72 [3.17, 7.02]
3 Difficult tracheal intubation: remifentanyl vs no remifentanyl	26	3008	Risk Ratio (M-H, Random, 95% CI)	5.64 [3.82, 8.31]
3.1 Remifentanyl	4	372	Risk Ratio (M-H, Random, 95% CI)	15.86 [4.43, 56.71]
3.2 No remifentanyl	22	2636	Risk Ratio (M-H, Random, 95% CI)	5.23 [3.54, 7.74]
4 Difficult tracheal intubation: alfentanil vs no alfentanil	26	2618	Risk Ratio (M-H, Random, 95% CI)	4.77 [3.25, 7.01]
4.1 Alfentanil	6	511	Risk Ratio (M-H, Random, 95% CI)	4.46 [1.66, 11.98]
4.2 No alfentanil	20	2107	Risk Ratio (M-H, Random, 95% CI)	5.10 [3.34, 7.79]
5 Difficult tracheal intubation: local anaesthesia vs no local anaesthesia	31	3184	Risk Ratio (M-H, Random, 95% CI)	5.04 [3.48, 7.29]
5.1 Local anaesthesia	5	307	Risk Ratio (M-H, Random, 95% CI)	1.90 [1.14, 3.18]
5.2 No local anaesthesia	26	2877	Risk Ratio (M-H, Random, 95% CI)	6.26 [4.15, 9.44]
6 Difficult tracheal intubation: excluded anticipated DTI vs included anticipated DTI	34	3564	Risk Ratio (M-H, Random, 95% CI)	5.00 [3.50, 7.16]
6.1 Exclusion of patients with anticipated difficult intubation	25	2886	Risk Ratio (M-H, Random, 95% CI)	5.32 [3.54, 8.00]
6.2 No exclusion of patients with anticipated difficult intubation	9	678	Risk Ratio (M-H, Random, 95% CI)	4.40 [1.71, 11.29]
7 Difficult tracheal intubation: "best-case scenario"	34	2410	Risk Ratio (M-H, Random, 95% CI)	5.99 [3.46, 10.38]
8 Difficult tracheal intubation excluding dose-finding studies	16	1536	Risk Ratio (M-H, Random, 95% CI)	3.40 [1.63, 7.10]
9 Difficult tracheal intubation: funding from pharmaceutical industry	34	3565	Risk Ratio (M-H, Random, 95% CI)	5.00 [3.49, 7.15]
9.1 No funding from pharmaceutical industry	24	2550	Risk Ratio (M-H, Random, 95% CI)	5.33 [3.16, 8.98]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
9.2 Funding from pharmaceutical industry	10	1015	Risk Ratio (M-H, Random, 95% CI)	4.10 [2.67, 6.31]
10 One or more events of upper airway discomfort or injury: low risk of bias vs high or uncertain risk of bias	7	844	Risk Ratio (M-H, Random, 95% CI)	1.36 [1.08, 1.71]
10.1 Low risk of bias	1	73	Risk Ratio (M-H, Random, 95% CI)	2.74 [1.21, 6.21]
10.2 High or uncertain risk of bias	6	771	Risk Ratio (M-H, Random, 95% CI)	1.30 [1.08, 1.58]
11 One or more events of upper airway discomfort or injury: depolarizing vs non-depolarizing NMBA	7	846	Risk Ratio (M-H, Random, 95% CI)	1.37 [1.09, 1.74]
11.1 Depolarizing NMBA	2	180	Risk Ratio (M-H, Random, 95% CI)	1.48 [0.83, 2.65]
11.2 Non-depolarizing NMBA	5	666	Risk Ratio (M-H, Random, 95% CI)	1.37 [0.97, 1.94]
12 One or more events of upper airway discomfort or injury: remifentanyl vs no remifentanyl	7	846	Risk Ratio (M-H, Random, 95% CI)	1.37 [1.09, 1.74]
12.1 Remifentanyl	2	193	Risk Ratio (M-H, Random, 95% CI)	1.12 [0.61, 2.08]
12.2 No remifentanyl	5	653	Risk Ratio (M-H, Random, 95% CI)	1.42 [1.16, 1.75]
13 One or more events of upper airway discomfort or injury: alfentanil vs no alfentanil	5	446	Risk Ratio (M-H, Random, 95% CI)	1.47 [0.85, 2.53]
13.1 No alfentanil	5	446	Risk Ratio (M-H, Random, 95% CI)	1.47 [0.85, 2.53]
14 One or more events of upper airway discomfort or injury: excluded anticipated DTI vs included anticipated DTI	7	846	Risk Ratio (M-H, Random, 95% CI)	1.37 [1.09, 1.74]
14.1 Excluded anticipated DTI	6	766	Risk Ratio (M-H, Random, 95% CI)	1.37 [1.05, 1.79]
14.2 Included anticipated DTI	1	80	Risk Ratio (M-H, Random, 95% CI)	2.0 [0.39, 10.31]
15 Difficult laryngoscopy: low risk of bias vs high or uncertain risk of bias	13	1308	Risk Ratio (M-H, Random, 95% CI)	2.54 [1.53, 4.21]
15.1 Low risk of bias	1	78	Risk Ratio (M-H, Random, 95% CI)	4.0 [0.47, 34.20]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
15.2 High or uncertain risk of bias	12	1230	Risk Ratio (M-H, Random, 95% CI)	2.47 [1.47, 4.16]

Analysis 1.1. Comparison 1 Avoidance vs use of NMBA, Outcome 1 Difficult tracheal intubation: low risk of bias vs high or uncertain risk of bias.

Analysis 1.2. Comparison 1 Avoidance vs use of NMBA, Outcome 2 Difficult tracheal intubation: depolarizing vs non-depolarizing NMBA.

Analysis 1.3. Comparison 1 Avoidance vs use of NMBA, Outcome 3 Difficult tracheal intubation: remifentanyl vs no remifentanyl.

Analysis 1.4. Comparison 1 Avoidance vs use of NMBA, Outcome 4 Difficult tracheal intubation: alfentanil vs no alfentanil.

Analysis 1.5. Comparison 1 Avoidance vs use of NMBA, Outcome 5 Difficult tracheal intubation: local anaesthesia vs no local anaesthesia.

Analysis 1.6. Comparison 1 Avoidance vs use of NMBA, Outcome 6 Difficult tracheal intubation: excluded anticipated DTI vs included anticipated DTI.

Analysis 1.7. Comparison 1 Avoidance vs use of NMBA, Outcome 7 Difficult tracheal intubation: "best-case scenario".

Analysis 1.8. Comparison 1 Avoidance vs use of NMBA, Outcome 8 Difficult tracheal intubation excluding dose-finding studies.

Analysis 1.9. Comparison 1 Avoidance vs use of NMBA, Outcome 9 Difficult tracheal intubation: funding from pharmaceutical industry.

Analysis 1.10. Comparison 1 Avoidance vs use of NMBA, Outcome 10 One or more events of upper airway discomfort or injury: low risk of bias vs high or uncertain risk of bias.

Analysis 1.11. Comparison 1 Avoidance vs use of NMBA, Outcome 11 One or more events of upper airway discomfort or injury: depolarizing vs non-depolarizing NMBA.

Analysis 1.12. Comparison 1 Avoidance vs use of NMBA, Outcome 12 One or more events of upper airway discomfort or injury: remifentanyl vs no remifentanyl.

Analysis 1.13. Comparison 1 Avoidance vs use of NMBA, Outcome 13 One or more events of upper airway discomfort or injury: alfentanil vs no alfentanil.

Study or subgroup	Avoidance of NMBA n/N	Use of NMBA n/N	Risk Ratio M-H, Random, 95% CI	Weight	Risk Ratio M-H, Random, 95% CI
Test for overall effect: $Z=1.38(P=0.17)$					
Favours avoidance of NMBA 0.01 0.1 1 10 100 Favours use of NMBA					

Analysis 1.14. Comparison 1 Avoidance vs use of NMBA, Outcome 14 One or more events of upper airway discomfort or injury: excluded anticipated DTI vs included anticipated DTI.

Study or subgroup	Avoidance of NMBA n/N	Use of NMBA n/N	Risk Ratio M-H, Random, 95% CI	Weight	Risk Ratio M-H, Random, 95% CI
1.14.1 Excluded anticipated DTI					
Bouvet 2008	14/65	17/64		12.55%	0.81[0.44,1.5]
Combes 2007	86/150	64/150		50.36%	1.34[1.07,1.69]
González Obregón 2010	0/50	0/50			Not estimable
Mencke 2003	16/36	6/37		7.56%	2.74[1.21,6.21]
Mencke 2014	17/31	12/33		15.18%	1.51[0.87,2.62]
Sivalingam 2001	34/75	8/25		12.35%	1.42[0.76,2.64]
Subtotal (95% CI)	407	359		98%	1.37[1.05,1.79]
Total events: 167 (Avoidance of NMBA), 107 (Use of NMBA)					
Heterogeneity: $\tau^2=0.03$; $\chi^2=5.67$, $df=4(P=0.22)$; $I^2=29.49\%$					
Test for overall effect: $Z=2.31(P=0.02)$					
1.14.2 Included anticipated DTI					
Gulhas 2013	4/40	2/40		2%	2[0.39,10.31]
Subtotal (95% CI)	40	40		2%	2[0.39,10.31]
Total events: 4 (Avoidance of NMBA), 2 (Use of NMBA)					
Heterogeneity: Not applicable					
Test for overall effect: $Z=0.83(P=0.41)$					
Total (95% CI)	447	399		100%	1.37[1.09,1.74]
Total events: 171 (Avoidance of NMBA), 109 (Use of NMBA)					
Heterogeneity: $\tau^2=0.01$; $\chi^2=5.89$, $df=5(P=0.32)$; $I^2=15.11\%$					
Test for overall effect: $Z=2.66(P=0.01)$					
Test for subgroup differences: $\chi^2=0.2$, $df=1$ ($P=0.66$), $I^2=0\%$					
Favours avoidance of NMBA 0.01 0.1 1 10 100 Favours use of NMBA					

Analysis 1.15. Comparison 1 Avoidance vs use of NMBA, Outcome 15 Difficult laryngoscopy: low risk of bias vs high or uncertain risk of bias.

Study or subgroup	Avoidance of NMBA n/N	Use of NMBA n/N	Risk Ratio M-H, Random, 95% CI	Weight	Risk Ratio M-H, Random, 95% CI
1.15.1 Low risk of bias					
Mencke 2003	4/39	1/39		5.55%	4[0.47,34.2]
Subtotal (95% CI)	39	39		5.55%	4[0.47,34.2]
Total events: 4 (Avoidance of NMBA), 1 (Use of NMBA)					
Heterogeneity: Not applicable					
Favours avoidance of NMBA 0.01 0.1 1 10 100 Favours use of NMBA

ADDITIONAL TABLES

Table 1. Baseline

Study ID	NMBA	Country	Language	Randomized	Sex	Age, years	Weight, kg	BMI	ASA class included	Expected difficult airway excluded	Overweight excluded
Alexander 1999	C1: suxamethonium 1 mg/kg	UK	English	60	C: 12F/8M I1: 11F/9M I2: 11F/9M	C: 41.7 (17.4) I1: 40.3 (10.6) I2: 44.2 (15.0)	C: 76.3 (15.0) I1: 75.5 (15.1) I2: 76.6 (16.8)	ns	I-II	yes	
Barclay 1997	C1: rocuronium 0.1 mg/kg C2: rocuronium 0.13 mg/kg	UK	English	60	ns	C1: 30 C2: 29 I1: 27	C1: 74 C2: 72 I1: 64	ns	ns	yes	yes
Beck 1993	C1: suxamethonium 1 mg/kg	USA	English	64	C1: 22F/11M I1: 21F/10M	C1: 34 (11) I1: 35 (11)	C1: 69 (14) I1: 65 (13)		I-II		
Bouvet 2008	C1: cisatracurium 0.15 mg/kg	France	English	130	C1: 65F I1: 65F	C1: 41.5 (12.9) I1: 40.7 (15.2)	ns	C1: 24.6 (5.4) I1: 23.0 (3.5)	I-II	yes	
Combes 2007	C1: rocuronium 0.6 mg/kg	France	English	300	C1: 73F/77M I1: 69F/81M	C1: 41 (18-70) I1: 43 (18-66)	C1: 73 (13) I1: 70 (13)	ns	I-II	yes	yes
Dominici 1990	C1: suxamethonium 1.5 mg/kg	France	French	60	C1: 9F/21M I1: 9F/21M	C1: 48.4 (3.4) I1: 50.1 (2.9)	C1: 62.8 (2.5) I1: 61.6 (2.1)	ns	I-III		
González Obregón 2010	C1: rocuronium 0.6 mg/kg	Columbia	Spanish	100	C1: 33F/17M I1: 33F/17M	C1: 34.7 (11.0) I1: 32.8 (12.3)	ns	ns	I-II	yes	
Gulhas 2013	C1: succinylcholine 1 mg/kg	Turkey	English	80	C1: 19F/21M	C1: 49.6 (8.4)	C1: 77.3 (13.1)	ns	I-II		

Table 1. Baseline (Continued)

					I1: 22F/18M	I1: 47.9 (8.7)	I1: 73.2 (14.4)				
Hanna 2010	C1: rocuronium 0.06 mg/kg	USA	English	50	C1: 15F/9M	C1: 39.0 (13.3)	C1: 75.0 (15.0)	C1: 25.9 (4.6)	I-II	yes	
	+ succinylcholine 1.5 mg/kg				I1: 5F/18M	I1: 43.0 (14.5)	I1: 81.0 (13.0)	I1: 26.5 (2.9)			
Harsten 1997	C1: suxamethonium 1 mg/kg	Sweden	English	80	C1: 26F/13M	C1: 41.8 (13)	ns	ns	I-II		
					I1: 23F/14M	I1: 39.5 (14)					
Iama- roon 2001	C1: suxamethonium 1.5 mg/kg	Thailand	English	120	C1: 54F/6M	C1: 40.6 (9.1)	C1: 55.8 (10.8)	ns	I-II	yes	yes
					I1: 54F/6M	I1: 39.7 (9.2)	I1: 55.1 (9.1)				
Isesele 2012	C1: suxamethonium 1.5 mg/kg	Nigeria	English	96	C1: 12F/32M	C1: 30.8 (9.0)	C1: 69.0 (7.4)		I-II	yes	
					I1: 21F/23M	I1: 32.6 (8.0)	I1: 68.3 (6.6)				
Jiao 2014	C1: suxamethonium 0.6 mg/kg	China	English	55	C1: 27F/0M	C1: 38.4 (10.9)	C1: 58.1 (7.0)	ns	I-II		yes
					I1: 28F/0M	I2: 36.3 (9.9)	I2: 58.2 (7.9)				
Kahwaji 1997	C1: ORG 9487 (ra- pachuronium) 0.5 mg/ kg	USA	English	181	C1: 20F/10M	C1: 51.3	C1: 67.3	ns	I-III	yes	
					C2: 18F/9M	C2: 49.6	C2: 70.5				
	C2: ORG 9487 (ra- pachuronium) 1.0 mg/ kg				C3: 17F/15M	C3: 52.0	C3: 68.6				
					C4: 15F/13M	C4: 50.6	C4: 71.3				
	C3: ORG 9487 (ra- pachuronium) 1.5 mg/ kg				C5: 22F/9M	C5: 50.2	C5: 75.9				
					I1: 19F/11M	I1: 52.6	I1: 69.7				
	C4: ORG 9487 (ra- pachuronium) 2.0 mg/ kg										
	C5: ORG 9487 (ra- pachuronium) 2.5 mg/ kg										

Table 1. Baseline (Continued)

Kirkegaard-Nielsen 1999	C1: rocuronium 0.4 mg/kg	USA	English	80	C1: 2F/18M	C1: 39.7 (7.5)	C1: 75.0 (16.9)	ns	I-II		yes
					C2: 6F/14M	C2: 39.5 (14.3)	C2: 78.6 (15.8)				
	C2: rocuronium 0.8 mg/kg				C3: 10F/10M	C3: 39.2 (10.5)	C3: 67.4 (14.8)				
	C3: rocuronium 1.2 mg/kg				I1: 5F/15M	I1: 39.3 (11.8)	I1: 73.4 (16.6)				
Kopman 2001	C1: rapacuronium 1.0 mg/kg	USA	English	100	ns	range: 18-65	ns	range: 17.5-27.5	I-II		yes
	C2: rapacuronium 1.2 mg/kg										
	C3: rocuronium 0.50 mg/kg										
Lieutaud 2003	C1: atracurium 0.5 mg/kg	France	English	170	C1: 3F/42M	C1: 52.9 (11.8)	ns	C1: 23.7 (3.2)	I-II	yes	
					C2: 7F/41M	C2: 51.3 (12.6)		C2: 23.1 (3.2)			
	C2: atracurium 0.5 mg/kg				C3: 8F/39M	C3: 56.3 (11.9)		C3: 23.6 (3.4)			
	C3: atracurium 0.5 mg/kg				I1: 2F/18M	I1: 50.4 (10.7)		I1: 23.3 (3.9)			
Lowry 1999	C1: rocuronium 0.3 mg/kg	UK	English	140	C1: 4F/16M	C1: 29 (11)	C1: 77(16)	ns	I-II	yes	yes
					C2: 7F/13M	C2: 40 (14)	C2: 75 (14)				
					C3: 12F/8M	C3: 36 (12)	C3: 69 (14)				
	C2: rocuronium 0.45 mg/kg				C4: 9F/11M	C4: 33 (12)	C4: 72 (12)				
					C5: 9F/11M	C5: 30 (12)	C5: 73 (14)				
	C3: rocuronium 0.6 mg/kg				C6: 4F/16M	C6: 33 (13)	C6: 74 (14)				
					I1: 2F/8M	I1: 29 (11)	I1: 72 (12)				
	C4: rocuronium 0.3 mg/kg				I2: 4F/6M	I2: 30 (9)	I2: 73 (15)				
	C5: rocuronium 0.45 mg/kg										
	C6: rocuronium 0.6 mg/kg										

Table 1. Baseline (Continued)

McNeil 2000	C1: succinylcholine 1 mg/kg	UK	English	60	ns	C1: 44 (15) I1: 39 (11) I2: 40 (13)	C1: 75 (10) I1: 76 (15) I2: 71 (12)	ns	I-II	yes	yes
Mencke 2003	C1: atracurium 0.5 mg/kg	Germany	English	80	C1: 19F/18M I1: 18F/18M	C1: 47.2 (13.2) I1: 47.7 (14.3)	C1: 77.7 (16) I1: 74.2 (15)		I-II	yes	yes
Mencke 2014	I1: rocuronium 0.45 mg·kg ⁻¹	Germany	English	83	C1: 16F/24M I1: 16F/23M	C1: 50 (16) I1: 48 (17)	C1: 83.8 (16) I1: 79.6 (15)	C1: 28.2 (4.3) I1: 26.5 (3.7)	I-III	yes	yes
Naguib 2003	C1: succinylcholine 0.3 mg/kg C2: succinylcholine 0.5 mg/kg C3: succinylcholine 1.0 mg/kg	Saudi Arabia	English	200	C1: 25F/25M C2: 23F/27M C3: 28F/22M I1: 23F/27M	C1: 30.9 (28-34) C2: 30.5 (27-34) C3: 30.0(28-32) I1: 29.5 (27-32)	C1: 66.6 (64-70) C2: 67.4 (64-71) C3: 67.8 (65-71) I1: 67.4 (64-71)	ns	I	yes	
Naguib 2006	C1: succinylcholine 0.3 mg/kg C2: succinylcholine 0.5 mg/kg C3: succinylcholine 1.0 mg/kg C4: succinylcholine 1.5 mg/kg C5: succinylcholine 2.0 mg/kg	Saudi Arabia	English	180	C1: 17F13M C2: 19F/11M C3: 13F/17M C4: 14F/16M C5: 18F/12M I1: 19F/11M	C1: 33.5 (8.7) C2: 29.7 (8.8) C3: 28.3 (7.9) C4: 31.5 (9.6) C5: 33.8 (14.8) I1: 20.1 (8.8)	C1: 67.8 (10.3) C2: 67.3 (10.8) C3: 71.1 (14.2) C4: 72.9 (12.5) C5: 70.9 (14.5) I1: 67.4 (10.7)	C1: 25.6 (2.8) C2: 25.6 (3.2) C3: 25.9 (3.9) C4: 26.2 (3.2) C5: 25.7 (3.9) I1: 25.7 (3.4)	I	yes	
Nimmo 1995	C1: suxamethonium 0.25 mg/kg C2: suxamethonium 0.5 mg/kg	USA	English	60	C1: 12F/8M C2: 12F/8M I1: 14F/6M	C1: 28.6 (17-55) C2: 29.0 (16-53) I1: 27.0 (18-53)	C1: 66.2 (13.6) C2: 64.4 (11.2) I1: 68.1 (13.6)	ns	I-II		

Table 1. Baseline (Continued)

Pang 2014	C1: cisatracurium 0.1 mg/kg	China	English	40	C1: 14F/6M I1: 9F/11M	C1: 45.2 (7.4) I1: 43.3 (6.7)	C1: 63.8 (9.5) I1: 64.6 (7.9)	C1: 23.7 (2.8) I1: 23.3 (3.1)	I-II	yes	yes
Pino 1998	C1: mivacurium 0.25 mg/kg C2: rocuronium 0.45 mg/kg C3: rocuronium 0.6 mg/kg C4: rocuronium 0.9 mg/kg C5: rocuronium 1.2 mg/kg	USA	English	100	ns	ns	ns	ns	I-II	yes	yes
Rousseau 1995	C1: vecuronium 0.08 mg/kg	France	French	152	ns	C1: 23 (5) I1: 25 (8)	C1: 71 (10) I1: 71 (11)	ns	I	yes	
Scheller 1992	C1: d-tubocurarine 3 mg and succinyl- choline 1 mg/kg	USA	English	75	C1: 8F/7M I1: 10F/5M I2: 11F/4M I3: 13F/2M I4: 10F/5M	C1: 37 (10) I1: 33 (9) I2: 30 (10) I3: 35 (11) I4: 36 (16)	C1: 77 (20) I1: 65 (11) I2: 66 (15) I3: 66 (12) I4: 68 (16)	ns	I	yes	
Schlaich 2000	C1: rocuronium 0.6 mg/kg C2: rocuronium 0.45 mg/kg C3: rocuronium 0.3 mg/kg	Germany	English	120	C1: 13F/17M C2: 13F/17M C3: 14F/16M I1: 14F/16M	C1: 37 (11) C2: 35 (11) C3: 36 (12) I1: 37 (11)	C1: 72 (14) C2: 75 (13) C3: 75 (12) I1: 70 (14)	ns	I-II	yes	
Si- valingam 2001	C1: suxamethonium 1 mg/kg	New Zealand	English	100	C1: 7F/18M I1: 9F/16M	C1: 34.3 (14.0) I1: 36.8 (12.6)	C1: 66 (10) I1: 62 (11)	ns	I-II	yes	

Table 1. Baseline (Continued)

					I2: 8F/17M	I2: 29.6 (9.7)	I2: 63 (15)							
					I3: 10F/15M	I3: 37.7 (12)	I3: 61 (11)							
Stevens 1997	C1: d-tubocurarine 3 mg and succinyl- choline 1 mg/kg	USA	English	140	C1: 12F/8M	C1: 35 (9)	C1: 70 (8)	ns	I-II	yes	yes			
					I1: 12F/8M	I1: 38 (12)	I1: 72 (17)							
					I2: 15F/5M	I2: 34 (11)	I2: 70 (14)							
					I3: 17F/3M	I3: 37 (10)	I3: 72 (10)							
					I4: 17F/3M	I4: 34 (9)	I4: 72 (13)							
					I5: 15F/5M	I5: 33 (11)	I5: 72 (18)							
					I6: 14F/6M	I6: 37 (14)	I6: 70 (13)							
Striebel 1995	C1: vecuronium 1 mg + succinylcholine 1 mg/kg	Germany	German	100	C1: 25F	C1: 47.8 (11.7)	C1: 62.6 (9.4)	ns	I-II					
					C2: 25F	C2: 43.8 (9.5)	C2: 68.2 (14)							
	C2: vecuronium 1 mg + succinylcholine 1 mg/kg				I1: 25F	I1: 46.5 (12.7)	I1: 64.9 (10.1)							
					I2: 25F	I2: 46.0 (12.4)	I2: 70.8 (14.6)							
Wong 1996	C1: succinylcholine 1 mg/kg	Malaysia	English	120	C1: 16F/14M	C1: 35.7 (16)	C1: 60.2 (8.9)	ns	I-II	yes				
					I1: 13F/17M	I1: 35.5 (12)	I1: 66.0 (13.1)							
					I2: 18F/12M	I2: 35.4 (13)	I2: 63.4 (12.9)							
					I3: 12F/17M	I3: 35.7 (11)	I3: 60.1 (10.8)							
Yazdi 2016	C1: atracurium 0.5 mg/kg	Iran	English	66	69.7% M	31.6 (13)	ns	ns	I-II	yes				

ns = not specified; The values in parentheses are standard deviation or range

Table 2. Intervention

Study ID	NMBA	Random- ized/ Analysed	Hypnotic	Opioid	Local anaesthetic	Difficult intubation events/	Difficult laryngos- copy	Upper air- way discomfort or injury
----------	------	----------------------------------	----------	--------	----------------------	------------------------------------	--------------------------------	--

Table 2. Intervention (Continued)

						total	events/ total	events/to- tal
Alexander 1999	C1: suxamethoni- um 1 mg/kg	60/60	C1: propofol 2 mg/kg I1: propofol 2 mg/kg I2: propofol 2 mg/kg	C1: none I1: alfentanil 50 µg/kg I2: remifentanil 2 µg/ kg	None	C1: 0/20 I1: 3/20 I2: 13/20	ns	ns
Barclay 1997	C1: rocuronium 0.1 mg/kg C2: rocuronium 0.3 mg/kg	60/60	C1: propofol 2.5 mg/kg C2: propofol 2.5 mg/kg I1: propofol 2.5 mg/kg	C1: alfentanil 10 µg/kg C2: alfentanil 10 µg/kg I1: alfentanil 10 µg/kg	Lidocaine 10 mg IV	C1: 14/20 C2: 2/20 I1: 19/20	ns	ns
Beck 1993	C1: suxamethoni- um 1 mg/kg	64/64	C1: thiopenthal 5 mg/kg I1: propofol 2 mg/mL	C1: none I1: alfentanil 50 µg/kg	None	C1: 0/33 I1: 1/31	C1: 0/33 I1: 1/31	ns
Bouvet 2008	C1: cisatracurium 0.15 mg/kg	130/129	C1: propofol 2.5 mg/kg I1: propofol 2.5 mg/kg	C1: remifentanil 2 µg/ kg I1: remifentanil 2 µg/ kg	None	C1: 0/64 I1: 3/65	C1: 1/64 I1: 1/65	C1: 17/64 I1: 14/65
Combes 2007	C1: rocuronium 0.6 mg/kg	300/300	C1: propofol 2.5 mg/kg I1: propofol 2.5 mg/kg	C1: alfentanil 15 µg/kg I1: alfentanil 40 µg/kg	None	C1: 1/150 I1: 18/150	C1: 5/150 I1: 18/150	C1: 64/150 I1: 86/150
Dominici 1990	C1: suxamethoni- um 1.5 mg/kg	60	C1: propofol 3 mg/mL I1: propofol 3 mg/mL	C1: alfentanil 7-10 µg/ kg I1: alfentanil 7-10 µg/ kg	Lidocaine (2%): IV + topical Li- docaine 5%	C1: 10/30 I1: 11/30	C1: 5/30 I1: 15/30	ns
González Obregón 2010	C1: rocuronium 0.6 mg/kg	100/100	C1: propofol 1-2 mg/kg I1: Sevoflurane 3% + propofol 2 mg/kg	C1: remifentanil 1-2 µg/kg in 1 min + 0.15 µg/kg/ min in 1 min I1: remifentanil	None	C1: 4/50 I1: 1/50	C1: 4/50 I1: 1/50	C1: 0/50 I1: 0/50

Table 2. Intervention (Continued)

0.6 µ/kg/min for 5 min								
Gulhas 2013	C1: succinylcholine 1 mg/kg	80/80	C1: propofol 2 mg/kg I1: propofol 2 mg/kg	C1: remifentanyl 1 µg/kg I1: remifentanyl 4 µg/kg	None	C1: 5/40 I1: 0/40	ns	C1: 2/40 I1: 4/40
Hanna 2010	C1: rocuronium 0.06 mg/kg + succinylcholine 1.5 mg/kg	50/47	C1: propofol 2 mg/kg I1: propofol 2 mg/kg	C1: none I1: remifentanyl 4 µg/kg	Lidocaine 0.5 mg/kg IV	C1: 2/24 I1: 3/23	ns	ns
Harsten 1997	C1: suxamethonium 1 mg/kg	80/79	C1: thiopental 5 mg/kg I1: propofol 2.5 mg/kg	C1: alfentanil 10 µg/kg I1: alfentanil 10 µg/kg	None	C1: 0/40 I1: 6/39	C1: 0/40 I1: 2/39	ns
Iamaroon 2001	C1: suxamethonium 1.5 mg/kg	120/120	C1: thiopental 5 mg/kg + (N ₂ O) I1: sevoflurane 8%	C1: fentanyl 1.5 µg/kg I1: fentanyl 1.5 µg/kg	None	C1: 0/60 I1: 4/60	ns	ns
Isele 2012	C1: suxamethonium 1.5 mg/kg	96/88	C1: propofol 2.0 mg/kg I1: propofol 2.0 mg/kg	None	C1: none I1: lidocaine IV 1.5 mg/kg	C1: 0/44 I1: 18/44	ns	ns
Jiao 2014	C1: suxamethonium 0.6 mg/kg	55/55	C1: propofol 2 mg/kg I1: propofol 2 mg/kg	C1: remifentanyl 1 µg/kg I1: remifentanyl 1.5 µg/kg	None	C1: 1/27 I2: 13/28	ns	ns
Kahwaji 1997	C1: ORG 9487 (rapacuronium) 0.5 mg/kg C2: ORG 9487 (rapacuronium) 1.0 mg/kg C3: ORG 9487 (rapacuronium) 1.5 mg/kg	181/176	C1: thiopental 3-6 mg/kg C2: thiopental 3-6 mg/kg C3: thiopental 3-6 mg/kg C4: thiopental 3-6 mg/kg C5: thiopental 3-6 mg/kg I1: thiopental 3-6 mg/kg	C1: fentanyl 0.5-3 µm/kg C2: fentanyl 0.5-3 µm/kg C3: fentanyl 0.5-3 µm/kg C4: fentanyl 0.5-3 µm/kg C5: fentanyl 0.5-3 µm/kg	None	C1: 9/30 C2: 6/27 C3: 1/32 C4: 0/28 C5: 1/29 I1: 18/30	ns	ns

Table 2. Intervention (Continued)

	C4: ORG 9487 (rapacuronium) 2.0 mg/kg			I1: fentanyl 0.5-3 µm/kg				
	C5: ORG 9487 (rapacuronium) 2.5 mg/kg							
Kirkegaard-Nielsen 1999	C1: rocuronium 0.4 mg/kg	80/80	C1: propofol 2 mg/kg	C1: fentanyl 2 µm/kg	None	C1: 9/20	ns	ns
			C2: propofol 2 mg/kg	C2: fentanyl 2 µm/kg		C2: 2/20		
	C2: rocuronium 0.8 mg/kg		C3: propofol 2 mg/kg	C3: fentanyl 2 µm/kg		C3: 1/20		
	C3: rocuronium 1.2 mg/kg		I1: propofol 2 mg/kg	I1: fentanyl 2 µm/kg		I1: 13/20		
Kopman 2001	C1: rapacuronium 1.0 mg/kg	100/100	C1: propofol 2.0 mg/kg IV	C1: alfentanil 12.5 µg/kg	None	C1: 2/30		
			C2: propofol 2.0 mg/kg IV			C2: 0/30		
	C2: rapacuronium 1.2 mg/kg		C3: propofol 2.0 mg/kg IV	C2: alfentanil 12.5 µg/kg		C3: 0/30		
	C3: rocuronium 0.50 mg/kg		I1: propofol 2.0 mg/kg IV	C3: alfentanil 12.5 µg/kg		I1: 7/10		
				I1: alfentanil 12.5 µg/kg				
Lieutaud 2003	C1: atracurium 0.5 mg/kg	170/160	C1: propofol 1.5 mg/kg	C1: fentanyl 3 µm/kg	None	C1: 7/47	ns	ns
			C2: propofol 2.0 mg/kg	C2: fentanyl 3 µm/kg		C2: 1/48		
	C2: atracurium 0.5 mg/kg		C3: propofol 2.5 mg/kg	C3: fentanyl 3 µm/kg		C3: 2/45		
	C3: atracurium 0.5 mg/kg		I1: propofol 2.5 mg/kg	I1: fentanyl 3 µm/kg		I1: 13/20		
Lowry 1999	C1: rocuronium 0.3 mg/kg	140/140	C1: propofol 2-3 mg/kg	C1: fentanyl 1 µm/kg	None	C1: 11/20	ns	ns
			C2: propofol 2-3 mg/kg	C2: fentanyl 1 µm/kg		C2: 4/20		
	C2: rocuronium 0.45 mg/kg		C3: propofol 2-3 mg/kg	C3: fentanyl 1 µm/kg		C3: 2/20		
	C3: rocuronium 0.6 mg/kg		C4: sevoflurane 8%	C4: fentanyl 1 µm/kg		C4: 14/20		
			C5: sevoflurane 8%	C5: fentanyl 1 µm/kg		C5: 9/20		

Table 2. Intervention (Continued)

	C4: rocuronium 0.3 mg/kg		C6: sevoflurane 8%	C6: fentanyl 1 µm/kg		C6: 2/20		
			I1: propofol 2-3 mg/kg	I1: fentanyl 1 µm/kg		I1: 10/10		
	C5: rocuronium 0.45 mg/kg		I2: sevoflurane 8%	I2: fentanyl 1 µm/kg		I2: 9/10		
	C6: rocuronium 0.6 mg/kg							
McNeil 2000	C1: succinylcholine 1 mg/kg	60/60	C1: propofol 2 mg/kg	C1: none	None	C1: 0/17	C1: 0/17	ns
			I1: propofol 2 mg/kg	I1: remifentanyl 2 µg/kg		I1: 2/23	I1: 0/23	
			I2: propofol 2 mg/kg	I2: remifentanyl 4 µg/kg		I2: 2/20	I2: 0/20	
Mencke 2003	C1: atracurium 0.5 mg/kg	80/73	C1: propofol 2.5-3 mg/kg	C1: fentanyl 2-3 µg/kg	None	C1: 2/37	C1: 1/39	C1: 6/37
			I1: propofol 2.5-3 mg/kg	I1: fentanyl 2-3 µg/kg		I1: 12/36	I1: 4/39	I1: 16/36
Mencke 2014	C1: rocuronium 0.45 mg/kg/kg	83/83	C1: propofol 1.5 mg·kg ⁻¹ + sevoflurane 3.0-3.5 Vol%, 8 l·min ⁻¹ in 2-3 minutes	C1: remifentanyl 0.30 µg/kg/min for 3 minutes	None	C1: 1/40	C1: 0/40	C1: 12/33
			I1: propofol 1.5 mg/kg	I1: remifentanyl 0.30 µg/kg/min for 3 minutes		I1: 11/43	I1: 2/43	I1: 17/31
Naguib 2003	C1: succinylcholine 0.3 mg/kg	200/200	C1: propofol 2 mg/kg	C1: fentanyl 2 µg/kg	None	C1: 4/50	ns	ns
	C2: succinylcholine 0.5 mg/kg		C2: propofol 2 mg/kg	C2: fentanyl 2 µg/kg		C2: 3/50		
	C3: succinylcholine 1.0 mg/kg		C3: propofol 2 mg/kg	C3: fentanyl 2 µg/kg		C3: 1/50		
			I1: propofol 2 mg/kg	I1: fentanyl 2 µg/kg		I1: 15/50		
Naguib 2006	C1: succinylcholine 0.3 mg/kg	180/180	C1: propofol 2 mg/kg	C1: fentanyl 2 µm/kg	None	C1: 2/30	ns	ns
	C2: succinylcholine 0.5 mg/kg		C2: propofol 2 mg/kg	C2: fentanyl 2 µm/kg		C2: 2/30		
			C3: propofol 2 mg/kg	C3: fentanyl 2 µm/kg		C3: 1/30		
	C3: succinylcholine 1.0 mg/kg		C4: propofol 2 mg/kg	C4: fentanyl 2 µm/kg		C4: 1/30		
			C5: propofol 2 mg/kg	C5: fentanyl 2 µm/kg		C5: 0/30		

Table 2. Intervention *(Continued)*

	C4: succinylcholine 1.5 mg/kg		I1: propofol 2 mg/kg	I1: fentanyl 2 µm/kg		I1: 21/30		
	C5: succinylcholine 2.0 mg/kg							
Nimmo 1995	C1: suxamethonium 0.25 mg/kg	60/60	C1: propofol 2.5 mg/kg	C1: alfentanil 15 µg/kg	None	C1: 0/20	ns	ns
			C2: propofol 2.5 mg/kg	C2: alfentanil 15 µg/kg		C2: 1/20		
	C2: suxamethonium 0.5 mg/kg		I1: Propofol 2.5 mg/kg	I1: alfentanil 15 µg/kg		I1: 9/20		
Pang 2014	C1: cisatracurium 0.1 mg/kg	20/20	C1: propofol target control I1: propofol target control	C1: remifentanyl target control I1: remifentanyl target control	C1: tetracaine 10 mg/mL I1: tetracaine 10 mg/mL	C1: 0/20 I1: 0/20	C1: 0/20 I1: 0/20	ns
Pino 1998	C1: mivacurium 0.25 mg/kg	100/98	C1: propofol 2 mg/kg	C1: fentanyl 2 µm/kg	None	C1: 2/30	ns	ns
			C2: propofol 2 mg/kg	C2: fentanyl 2 µm/kg		IC2: 9/15		
	C2: rocuronium 0.45 mg/kg		C3: propofol 2 mg/kg	C3: fentanyl 2 µm/kg		C3: 4/14		
	C3: rocuronium 0.6 mg/kg		C4: propofol 2 mg/kg	C4: fentanyl 2 µm/kg		C4: 1/14		
	C4: rocuronium 0.9 mg/kg		C5: propofol 2 mg/kg	C5: fentanyl 2 µm/kg		C5: 0/15		
	C5: rocuronium 1.2 mg/kg		I1: propofol 2 mg/kg	I1: fentanyl 2 µm/kg		I1: 10/10		
Rousseau 1995	C1: vecuronium 0.08 mg/kg	152/152	C1: propofol 2.5 mg/kg I1: propofol 2.5 mg/kg	C1: alfentanil 0.03 mg/kg I1: alfentanil 0.03 mg/kg	C1: none I1: lidocaine 1.5 mg/kg	C1: 2/77 I1: 4/75	ns	ns
Scheller 1992	C1: d-tubocurarine 3 mg and succinylcholine 1 mg/kg	75/75	C1: thiamylal 4 mg/kg I1: propofol 2 mg/kg I2: propofol 2 mg/kg	C1: none I1: alfentanil 30 µg/kg I2: alfentanil 40 µg/kg	None	C1: 0/15 I1: 1/15 I2: 1/15	C1: 0/15 I1: 1/15 I2: 1/15	ns

Table 2. Intervention (Continued)

			I3: propofol 2 mg/kg	I3: alfentanil 50 µg/kg		I3: 1/15	I3: 1/15	
			I4: propofol 2 mg/kg	I4: alfentanil 60 µg/kg		I4: 1/15	I4: 1/15	
Schlaich 2000	C1: rocuronium 0.6 mg/kg	120/120	C1: propofol 2-2.5 mg/kg	C1: remifentanil 0.5 µg/kg/min	None	C1: 0/30	ns	ns
	C2: rocuronium 0.45 mg/kg		C2: propofol 2-2.5 mg/kg	C2: remifentanil 0.5 µg/kg/min		C2: 1/30		
	C3: rocuronium 0.3 mg/kg		C3: propofol 2-2.5 mg/kg	C2: remifentanil 0.5 µg/kg/min		C3: 0/30		
			I1: propofol 2-2.5 mg/kg	IC3: remifentanil 0.5 µg/kg/min		I1: 12/30		
				I1: remifentanil 0.5 µg/kg/min				
Sivalingam 2001	C1: suxamethonium 1 mg/kg	100/100	C1: Sevoflu 7% + N ₂ O60%	C1: alfentanil 10 µg/kg	None	C1: 1/25	ns	C1: 8/25
			I1: Sevoflu 7% + N ₂ O60%	I1: alfentanil 20 µg/kg		I1: 4/25		I1: 12/25
			I2: Sevoflu 7% + N ₂ O60%	I2: alfentanil 25 µg/kg		I2: 5/25		I2: 13/25
			I3: Sevoflu 7% + N ₂ O60%	I3: alfentanil 30 µg/kg		I3: 2/25		I3: 9/25
Stevens 1997	C1: d-tubocurarine 3 mg and succinylcholine 1 mg/kg	140/140	C1: thiopental 4 mg/kg	C1: none	C1: none	C1: 1/20	C1: 0/20	ns
			I1: etomidate 0.3 mg/kg	I1: alfentanil 40 µg/kg	I1: none	I1: 3/20	I1: 0/20	
			I2: etomidate 0.3 mg/kg	I2: alfentanil 40 µg/kg	I2: lidocaine 1 mg/kg	I2: 1/20	I2: 0/20	
			I3: propofol 2 mg/kg	I3: alfentanil 40 µg/kg	I3: none	I3: 3/20	I3: 0/20	
			I4: propofol 2 mg/kg	I4: alfentanil 40 µg/kg	I4: lidocaine 1 mg/kg	I4: 2/20	I4: 0/20	
			I5: thiopental 4 mg/kg	I5: alfentanil 40 µg/kg	I5: none	I5: 8/20	I5: 0/20	
			I6: thiopental 4 mg/kg	I6: alfentanil 40 µg/kg	I6: lido- caine1 mg/ kg	I6: 3/20	I6: 0/20	
Striebel 1995	C1: vecuronium 1 mg + succinylcholine 1 mg/kg	100/100	C1: thiopental 5.5 mg/kg	C1: fentanyl 0.1 mg	2 mL lido- caine 1% IV	C1: 1/25	C1: 2/25	ns
			C2: propofol 2.2 mg/kg	C2: fentanyl 0.1 mg		C2: 1/25	C2: 1/25	
			I1: propofol 2.4 mg/kg	I1: fentanyl 0.1 mg		I1: 3/25	I1: 1/28	

Table 2. Intervention *(Continued)*

	C2: vecuronium 1 mg + succinyl- choline 1 mg/kg		I2: propofol 2.2 mg/kg	I2: fentanyl 0.2 mg		I2: 5/25	I2: 4/25	
Wong 1996	C1: succinylcholine 1 mg/kg	120/120	C1: propofol 3.0 mg/kg I1: propofol 2.6 mg/kg I2: propofol 2.6 mg/kg I3: propofol 3.1 mg/kg	C1: none I1: alfentanil 15 µg/kg I2: alfentanil 30 µg/kg I3: none	None	C1: 0/30 I1: 1/30 I2: 0/30 I3: 6/30	ns	ns
Yazdi 2016	C1: atracurium 0.5 mg/kg	66/66	C1: propofol 2.5 mg/kg I1: propofol 2.5 mg/kg	C1: none I1: remifentanyl 2 µg/ kg	None	C1: 4/31 I1: 14/35	ns	ns

ns = not specified

APPENDICES

Appendix 1. Cormack and Lehane classification

Difficult laryngoscopy

Cormack and Lehane ([Cormack 1984](#)) classification

Grade 1: full view of the glottis.

Grade 2: partial view of the glottis or arytenoids.

Grade 3: only epiglottis visible.

Grade 4: neither glottis nor epiglottis visible.

Laryngoscopy grade 3 and 4 define a difficult laryngoscopy.

Modified Cormack and Lehane ([Yentis 1998](#)) classification

Grade 1: full view of the glottis.

Grade 2a: partial view of the glottis.

Grade 2b: arytenoids or posterior part of the vocal cords only just visible.

Grade 3: only epiglottis visible.

Grade 4: neither glottis nor epiglottis visible.

Laryngoscopy grades 2b, 3, and 4 define a difficult laryngoscopy.

Appendix 2. CENTRAL (the Cochrane Library) search strategy

#1 MeSH descriptor: [Neuromuscular Blocking Agents] explode all trees

#2 MeSH descriptor: [Muscle Relaxants, Central] explode all trees

#3 (suxameton or rapacuronium or mivacurium or atracurium or doxacurium or cisatracurium or vecuronium or rocuronium or pancuronium or tubocurarine or gallamine or pipecuronium):ti,ab)

#4 #1 or #2 or #3

#5 MeSH descriptor: [Laryngoscopy] explode all trees

#6 MeSH descriptor: [Intubation, Intratracheal] explode all trees

#7 (difficult near (intubat* or laryngoscopy or airway))

#8 (Intubation near (score or scale))

#9 Cormack or Lehane

#10 ((tracheal near intub*) or airway or laryngoscopy):ti

#11 #5 or #6 or #7 or #8 or #9 or #10

#12 #4 and #11

Appendix 3. MEDLINE (Ovid SP) search strategy

1 exp Neuromuscular Blocking Agents/ or Muscle Relaxants, Central/ or (suxamethonium or rapacuronium or mivacurium or atracurium or doxacurium or cisatracurium or vecuronium or rocuronium or pancuronium or tubocurarine or gallamine or pipecuronium).ti,ab.

2 Laryngoscopy/ or Intubation, Intratracheal/ or (difficult adj3 (intubat* or laryngoscopy or airway)).mp. or ((Intubation adj3 (score or scale)) or (Cormack or Lehane)).mp. or ((tracheal adj3 intub*) or airway or laryngoscopy).ti.

3 1 and 2

4 ((randomized controlled trial or controlled clinical trial).pt. or randomized.ab. or placebo.ab. or drug therapy.fs. or randomly.ab. or trial.ab. or groups.ab.) not (animals not (humans and animals)).sh.

5 3 and 4

Appendix 4. Embase (Ovid SP) search strategy

1. exp neuromuscular blocking agent/ or central muscle relaxant/ or (suxameton or rapacuronium or mivacurium or atracurium or doxacurium or cisatracurium or vecuronium or rocuronium or pancuronium or tubocurarine or gallamine or pipecuronium).ti,ab.

2. laryngoscopy/ or endotracheal intubation/ or (difficult adj3 (intubat* or laryngoscopy or airway)).mp. or ((Intubation adj3 (score or scale)) or (Cormack or Lehane)).mp. or ((tracheal adj3 intub*) or airway or laryngoscopy).ti.

3. 1 and 2

4. (placebo.sh. or controlled study.ab. or random*.ti,ab. or trial*.ti,ab. or ((singl* or doubl* or trebl* or tripl*) adj3 (blind* or mask*)).ti,ab.) not (animals not (humans and animals)).sh.

5. 3 and 4

Appendix 5. BIOSIS Citation Index

#1 TS=Neuromuscular Blocking or TS=Muscle Relaxant* or TS=(suxameton or rapacurionium or mivacurium or atracurium or doxacurium or cisatracurium or vecuronium or rocuronium or pancuronium or tubocurarine or gallamine or pipecuronium)

#2 TS=(Cormack or Lehane) or TS=(difficult SAME (intubat* or laryngoscopy or airway)) or TS=(Intubation SAME (score or scale)) or TI=((tracheal and intub*) or airway or laryngoscopy)

#3 #2 AND #1

#4 TS=(random* or (controlled SAME (study or trial*)) or prospective or placebo or multicenter) or TS=((mask* or blind*) SAME (single or double or triple or treble))

#5 #3 and #4

Appendix 6. CINAHL (EBSCO) search strategy

S1 ((MH "Neuromuscular Blocking Agents") OR (MH "Muscle Relaxants, Central")) OR AB (suxameton or rapacurionium or mivacurium or atracurium or doxacurium or cisatracurium or vecuronium or rocuronium or pancuronium or tubocurarine or gallamine or pipecuronium)

S2 ((MH "Laryngoscopy") OR (MH "Intubation, Intratracheal")) OR ((difficult and (intubat* or laryngoscopy or airway))) OR ((Intubation and (score or scale))) OR (Cormack or Lehane) OR TI ((tracheal and intub*) or airway or laryngoscopy)

S3 S2 and S1

Appendix 7. LILACS (BIREME) serach strategy

((Neuromuscular Blocking Agent\$) or Muscle Relaxant\$ or (suxameton or rapacurionium or mivacurium or atracurium or doxacurium or cisatracurium or vecuronium or rocuronium or pancuronium or tubocurarine or gallamine or pipecuronium)) and ((Cormack or Lehane) or (difficult and (intubate\$ or laryngoscopy or airway)) or (Intubation and (score or scale)) or ((tracheal and intub\$) or airway or laryngoscopy))

Appendix 8. Data extraction form

Study selection, quality assessment, and data extraction form

First study author	Journal/Conference proceedings, etc.	Year
--------------------	--------------------------------------	------

Study eligibility

RCT	Relevant participants	Relevant interventions	Relevant outcomes
-----	-----------------------	------------------------	-------------------

(Continued)

Yes/No/Unclear

Yes/No/Unclear

Yes/No/Unclear

Yes/No*/Unclear

Issue relates to selective reporting when study authors may have taken measurements for particular outcomes, but not reported these within the paper(s). Review authors should contact trialists for information on possible non-reported outcomes and reasons for exclusion from publication. Study should be listed in 'Studies awaiting assessment' until clarified. If no clarification is received after three attempts, study should then be excluded.

Do not proceed if any of the above answers are 'No'. If study is to be included in 'Excluded studies' section of the review, record below the information to be inserted into the 'Table of excluded studies'.

Freehand space for comments on study design and treatment:

References to trial

Check other references identified in searches. If further references to this trial are available, link the papers now and list below. All references to a trial should be linked under one *Study ID* in RevMan.

Code each paper	Study author(s)	Journal/Conference proceedings, etc.	Year
A	<i>The paper listed above</i>		
B	<i>Further papers</i>		
C			
D			
E			

Participants and trial characteristics

Avoidance versus use of neuromuscular blocking agents for improving conditions during tracheal intubation or direct laryngoscopy in adults and adolescents (Review)

117

Copyright © 2017 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

Participant characteristics

Covariate

Further details

Age (mean, median, range, etc.)

Sex of participants (numbers/%, etc.)

BMI (mean, median, range, etc.)

Trial characteristics

Methodological quality

Random sequence generation

State here method used to generate random sequence and reasons for grading

Grade (circle)

Low risk of bias (random)

High risk of bias (e.g. alternate)

Unclear

Allocation concealment

Process used to prevent foreknowledge of group assignment in an RCT, which should be seen as distinct from blinding

State here method used to conceal allocation and reasons for grading

Grade (circle)

Low risk of bias

High risk of bias

Unclear

Blinding

(Continued)

Person responsible for participant care	Low risk/High risk/Un-clear risk
Participant	Low risk/High risk/Un-clear risk
Outcome assessor	Low risk/High risk/Un-clear risk
Other (please specify)	Low risk/High risk/Un-clear risk
Incomplete outcome data	
Low risk, if numbers and reasons for dropouts and withdrawals in the inter-vention groups were described, or if it was specified that no dropouts or with-drawals occurred	Yes/No
High risk, if numbers or reasons for dropouts and withdrawals were not de-scribed	Yes/No
Unclear, if the report gave the impression that no dropouts or withdrawals had occurred, but did not specifically state this	Yes/No
Selective reporting	
Low risk, if predefined or clinically relevant and reasonably expected out-comes are reported	Yes/No
High risk, if one or more clinically relevant and reasonably expected outcomes were not reported; data on these outcomes were likely to have been recorded	Yes/No
Unclear, if not all predefined or clinically relevant and reasonably expect-ed outcomes are reported, or if they are not reported fully, or if it is unclear whether data on these outcomes were recorded	Yes/No
Baseline imbalance	
Low risk, if no baseline imbalance in important characteristics is evident	Yes/No
High risk, if a baseline imbalance is due to chance or is due to imbalanced ex-clusion after randomization	Yes/No
Unclear, if baseline characteristics were not reported	Yes/No
Early stopping	
Low risk, if sample size calculation was reported and the trial was not stopped, or if the trial was stopped early by formal stopping rules at a point when the likelihood of observing an extreme intervention effect due to chance was low	Yes/No
High risk, if the trial was stopped early owing to informal stopping rules, or if the trial was stopped early by a formal stopping rule at a point when the likeli-hood of observing an extreme intervention effect due to chance was high	Yes/No
Unclear, if sample size calculation was not reported and it is not clear whether the trial was stopped early	Yes/No

Other bias

Low risk of bias, if the trial appears to be free of other components that could put it at risk of bias Yes/No

High risk of bias, if other factors in the trial could put it at risk of bias (e.g. 'for-profit' involvement, authors have conducted trials on the same topic) Yes/No

Unclear, if the trial may or may not be free of other components that could put it at risk of bias Yes/No

Modified intention-to-treat

A modified intention-to-treat analysis is one in which all participants in a trial are operated and analysed according to the intervention to which they were allocated, whether or not they received it

All participants entering trial after surgery

15% or fewer excluded

More than 15% excluded

Not analysed as modified 'intention-to-treat'

Unclear

Were withdrawals described? Yes ? No ? Not clear ?

Discuss if appropriate

Trial characteristics

Further details

Single-centre/Multi-centre

Country/Countries

How was participant eligibility defined?

How many participants were randomized?

Number of intervention groups/number of control groups

Number of participants in each intervention group/Number of participants in each control group

(Continued)

Number of participants who received intended intervention (per-protocol population)

Number of participants who were analysed

Type of outcome measure (DTI/DL?)

NMBA: type and dose?

Hypnotic: type and dose

Opioid: type and dose

Local anaesthetic: type and dose

Other

* If cross-over design, please refer to the Cochrane Editorial Office for further advice on how to analyse these data

Other design characteristics of the trial

1. Mallampati score (predicts DTI/DL Yes/No??)
2. Thyromental distance (predicts DTI/DL Yes/No??) or (cm)
3. Neck extension (predicts DTI/DL Yes/No??) or (degrees)
4. Mouth opening/Interincisor gap (predicts DTI/DL Yes/No??) or (cm)
5. Mandible subluxation (predicts DTI/DL Yes/No??)
6. Time from induction to start of tracheal intubation (seconds)

Data extraction

Outcomes	Available for the trial
1.1 Difficult tracheal intubation	Yes/No
1.2 Overall mortality. We will use the maximal follow-up data from each trial	Yes/No
1.3. One or more events of upper airway discomfort/injury (e.g. sore throat, hoarseness, vocal cord lesion, minor pharyngeal injury)	Yes/No
2.1. One or more major serious events: gastric aspiration, brain and heart injuries (e.g. caused by anoxia, hypotension, bradycardia/tachycardia during tracheal intubation)	Yes/No
2.2 Difficult laryngoscopy	Yes/No

For continuous data						
Code of paper	Unit of measurement	Intervention group		Control group		Details if outcome described only in text
		n	Mean (SD)	n	Mean (SD)	
Outcomes (rename)						
A	1.1 Difficulty of tracheal intubation	Score				

For dichotomous data

Code of paper	Outcomes	Intervention group E/N	Control group E/N
		E = number of events N = number of par- ticipants	E = number of events N = number of par- ticipants
A	1.1 Difficult tracheal intubation		
	1.2 Overall mortality. We will use the maximal follow-up data from each trial		
	1.3. One or more events of upper airway discomfort/injury (e.g. sore throat, hoarseness, vocal cord lesion, minor pharyngeal injury)		
	2.1. One or more major serious events: gastric aspiration, brain and heart injuries (e.g. caused by anoxia, hypotension, bradycardia/tachycardia during tracheal intubation)		
	2.2 Difficult laryngoscopy		

Other information that you believe is relevant to the results

Indicate if any data were obtained from the primary author; if results were estimated from graphs, etc. or were calculated by using a formula (this should be stated and the formula given). In general, if results not reported in paper(s) are obtained, this should be made clear here to be cited in the review

Mallampati score	Predicts DTI/DL (numbers/% of participants)	
Thyromental distance	Predicts DTI/DL (numbers/% of participants)	cm (mean, median, range, etc.)
Neck extension	Predicts DTI/DL (numbers/% of participants)	degrees (mean, median, range, etc.)
Mouth opening/Interincisor gap	Predicts DTI/DL (numbers/% of participants)	cm (mean, median, range, etc.)
Mandible subluxation	Predicts DTI/DL (numbers/% of participants)	
Time from induction to start of tracheal intubation		seconds (mean, median, range, etc.)

Freehand space for writing actions such as contact with study authors and changes

References to other trials

Did this report include any references to published reports of potentially eligible trials not already identified for this review?

First study author

Journal/Conference

Year of publication

Did this report include any references to unpublished data from potentially eligible trials not already identified for this review? If yes, list contact names and details

CONTRIBUTIONS OF AUTHORS

Conceiving of the review: LHL, SS, AMM, JW.

Designing the review: LHL, JW, AMM, SS.

Co-ordinating the review: LHL.

Undertaking manual searches: LHL.

Screening search results: LHL, CD, AN.

Organizing retrieval of papers: LHL, CD, AN.

Screening retrieved papers against inclusion criteria: LHL, CD, AN, CVR, JT.

Appraising the quality of papers: LHL, CD, AN, CVR, JT.

Abstracting data from papers: LHL, CD, AN, CVR, JT.

Writing to authors of papers for additional information: LHL.

Providing additional data about papers: LHL.

Obtaining and screening data on unpublished studies: LHL, AN.

Managing data for the review: LHL, CD, AN.

Entering data into Review Manager ([RevMan 5.3](#)): LHL, AN.

Analysing RevMan statistical data: LHL, JW.

Performing other statistical analysis not using RevMan: LHL, JW.

Performing double entry of data: data entered by person one: LHL; data entered by person two: AN.

Interpreting data: LHL, AMM, JW.

Making statistical inferences: LHL, JW.

Writing the review: LHL, CD, AN, CR, JT, SS, AMM, JW.

Securing funding for the review: Copenhagen Trial Unit, Rigshospitalet, Copenhagen, Denmark, and Department of Anaesthesiology, Herlev Hospital, University of Copenhagen, Herlev, Denmark.

Serving as guarantor for the review (one review author): LHL.

Taking responsibility for reading and checking the review before submission: LHL.

DECLARATIONS OF INTEREST

LHL, AMM, CR, and JW were co-authors of an observational study entitled "Avoidance of neuromuscular blocking agents may increase the risk of difficult tracheal intubation: a cohort study of 103,812 consecutive adult patients recorded in the Danish Anaesthesia Database" ([Lundström 2009a](#)).

JW is a member of the task force in the Copenhagen Trial Unit, which develops theory, software, and manuals for trial sequential analysis (TSA).

AMM is a Co-ordinating Editor for the Cochrane Anaesthesia, Critical and Emergency Care Group.

JT received grants for two studies in 2016 through the Merck Investigator Studies Program. The aims of the two studies were (1) to assess the use of neuromuscular monitoring and the incidence of residual neuromuscular blockade in six Danish anaesthesia departments, and (2) to assess the effect of an e-learning course in neuromuscular monitoring.

DIFFERENCES BETWEEN PROTOCOL AND REVIEW

We made the following changes to the protocol ([Lundström 2011a](#)).

1. Christophe HV Duez, Anders K Nørskov, Charlotte V Rosenstock, and Jakob L Thomsen joined as review authors after publication of the protocol.
2. We considered avoiding NMBA as the intervention, thus we changed the title from "Use versus avoidance of neuromuscular blocking agent for improving conditions during tracheal intubation or direct laryngoscopy in adults and adolescents" to "Avoidance versus use of neuromuscular blocking agent for improving conditions during tracheal intubation or direct laryngoscopy in adults and adolescents".
3. We did not consider stopping early for benefit, harm, or futility on inadequate stopping rules (or no reporting of sample size at all), as risk of bias in the accomplished review was high. In the protocol, we considered this a risk of bias component, but after the 11th Cochrane Symposium in Keystone, Colorado, early stopping was changed from constituting a bias risk to constituting increased risk of random error.
4. We did not consider baseline imbalance or lack of reporting of baseline characteristics as introducing high risk of bias. After the 11th Cochrane Symposium in Keystone, Colorado, baseline imbalance or lack of reporting of baseline characteristics changed from constituting a bias risk to constituting increased risk of random error.
5. We did not report on 'other bias' as suggested in the protocol.
6. In our protocol, we stated, "We will exclude trials using quasi-randomization and observational studies with regard to benefits, but not harms". In the review, we stated, "We excluded trials using quasi-randomization as well as observational studies". Thus we did not identify any quasi-randomized trials and did not include any observational studies examining harms.
7. Under 'Data extraction and management' in the protocol, we stated, "We will include each pair-wise comparison separately but with shared intervention groups divided out approximately evenly among the comparisons. For example if multiple intervention groups share a common control group the number of patients and the number of events of the control group will be divided equally, thereby the number of subgroups of the control group will match the number of intervention groups ([Higgins 2011](#))" and "We will combine all relevant experimental intervention groups of the trials into a single intervention group, and combined all relevant control intervention groups into a single control group" ([Higgins 2011](#)). However, during the review process, we decided to perform only the latter (recommended) method when handling studies with multiple intervention or control groups. Thus, in the review, we have stated the following: "Some trials randomized participants to multiple intervention and/or control groups (more than two groups, as in dose-finding studies)". In the review, we combined all relevant experimental intervention groups from trials into a single intervention group, and we combined all relevant control intervention groups into a single control group, as recommended ([Higgins 2011](#)).
8. As conditions for tracheal intubation were not reported as a continuous outcome in any of the included trials, we did not calculate risk differences or mean differences with 95% confidence intervals.
9. We added two explorative outcomes: (1) a best-case scenario, whereby dose-finding studies were represented only by control and intervention groups with the lowest prevalence of difficult intubation; and (2) exclusion of dose-finding studies.
10. We did not explore selective outcome reporting by comparing publications with their protocols, if the latter were available.

INDEX TERMS

Medical Subject Headings (MeSH)

Confidence Intervals; Intubation, Intratracheal [adverse effects] [*methods]; Laryngoscopy [adverse effects] [*methods]; Neuromuscular Blocking Agents [*administration & dosage] [adverse effects]; Neuromuscular Depolarizing Agents [administration & dosage]; Randomized Controlled Trials as Topic; Risk Factors; Sensitivity and Specificity; Succinylcholine [administration & dosage]

MeSH check words

Adolescent; Adult; Humans