CORRESPONDENCE

The Neurophysiology and Treatment of Motion Sickness

by Prof. Dr. med. Andreas Koch, Prof. Dr. med. Dr. rer. nat. Ingolf Cascorbi, Prof. Dr. med. Martin Westhofen, PD Dr. med. Manuel Dafotakis, Dr. med. Sebastian Klapa, and Prof. Dr. med. Johann Peter Kührt-Buschbeck in issue 41/2018

Additional Points

The best place for someone with sea sickness is not only the central section of the ship but the very lowest bunk in the very lowest cabin, as variations in motion are least pronounced at the center of rotation. When ascending the vessel's masts the motion angle rapidly increases, and at only 30 degrees of swaying to both sides is 60°, with motion sickness immediately getting worse.

The most effective medication is meclozine. Unfortunately this substance is no longer commercially available in Germany. Internationally, however, it is the substance of choice, even for women affected by hyperemesis gravidarum. Similarly, promethazine is no longer available in suppository form.

DOI: 10.3238/arztebl.2019.0175a

References

1. Koch A, Cascorbi I, Westhofen M, Dafotakis M, Klapa S, Kührt-Buschbeck JP: The neurophysiology and treatment of motion sickness. Dtsch Arztebl Int 2018; 115: 687–96.

Dr. med. Franz-Ulrich Beutner, Burgdorf, fub_bu@me.com

Conflict of interest statement

The author declares that no conflict of interest exists.

Counterarguments

As ship's doctors who travel on tall ships and cruise ships we feel compelled to disagree with the relevant article (1) in various points.

Sea sickness is not an individual illness but a mass illness. In rough seas, some 100 or more patients will all present to the ship's hospital at once. The sensation of sickness is enormous; many “want to die.” Because of the occasionally overwhelming degree of suffering, help needs to be given immediately, in the form of effective medications. If these are given by mouth they will be vomited out immediately. In this scenario of a multitude of people ill at the same time, a standardized approach is helpful: the intramuscular administration of dimenhydrinate, also in combination with metoclopramide (MCP). In some patients, the rectal application of dimenhydrinate will suffice. Often in such a scenario, nursing staff will have to be involved for the purposes of triage, administration of medication, and documentation.

The risk when faced with so many sick people lies in overlooking those who have different illnesses with similar symptoms or who may experience complications of their underlying illness because of nausea and vomiting.

The authors of the article recommend administration of scopolamine. But who knows two hours in advance that they are going to be sea sick? The risk was not mentioned: the side effects of the substance cannot be distinguished clinically from a cerebral

hemorrhage in patients with a craniocerebral trauma. Cinnarizine is not licensed for the treatment of sea sickness in Germany; furthermore, we are not aware of any preparation that can be given parenterally.

In our opinion, treating sea sickness on ships requires the following measures:

- Administration of dimenhydrinate and if needed also MCP—both substances are mostly administered intramuscularly—and, additionally
- bed rest for several hours, because during sleep the blood concentration of the trigger substance histamine falls.

DOI: 10.3238/arztebl.2019.0175b

References

1. Koch A, Cascorbi I, Westhofen M, Dafotakis M, Klapa S, Kührt-Buschbeck JP: The neurophysiology and treatment of motion sickness. Dtsch Arztebl Int 2018; 115: 687–96.

Dr. med. Karsten Mülder, Berlin, karsten@muelder.info

Prof. Dr. med. Hans-Detlef Stober, Berlin

Dr. med. Walter Hamacher, Berlin

Dr. med. Gernot Spiewok, Schwerin

Dr. med. Katrin Pätel, End- und Dickdarmpraxis, Sindelfingen

Conflict of interest statement

All signatories work as self-employed physicians on cruise ships.

In Reply:

Dr. Beutner has provided a vivid example of how intersensory conflicts may blur into one another (1, 2): someone on a ship's top deck or up in the masts may experience a temporary sensory discrepancy between the view at distant waves and the perception of the ship's own movement, which leads to incongruous, motion related information reaching the sensory organs in the sense of a type A1 conflict. Especially moving waves with a slow periodicity of 0.1–0.5 Hz are particularly pathogenetic (3), which intensifies higher up the mast owing to the wider range of movement.

On the bottom deck and in the center of the ship, near the apex of the ship's deflection, movements are least noticeable, which can help reduce the problems substantially. The lack of an outside view can instead lead to a kinetogenic type A3 conflict, since the visual impression of a seemingly stable environment and the information on the ship's movement provided by the vestibular apparatus and proprioception contradict each other.

Someone with motion sickness on board a ship is therefore faced with a dilemma relating to the basic conflict of the senses. In actual fact, the best solution would be to stay in the center of the ship, near the bottom, but with an external view.

We thank our correspondent for mentioning meclozine. The effectiveness of this first generation antihistamine has been much

confirmed for motion sickness and sickness in pregnancy. In the past it was also commonly prescribed for travel sickness, but it is no longer commercially available in Germany.

Furthermore we wish to extend our thanks to the group of ship's doctors for their enthusiastic correspondence. Sea sickness can easily become a mass illness, especially on a large ship, something we touched on in our article (4) by mentioning the potentially large proportions of crew thus affected. Sea sickness can also put pressure on doctors to act rapidly and decisively—especially at sea, where one cannot extricate oneself from the situation. This is particularly the case for elderly patients, in order to prevent/avoid any complications of their underlying disorders.

We think that in this context, the information about a possibly difficult differential diagnostic distinction between severe motion sickness and the side effects of relevant medication from health problems that may appear similar is extremely important and valuable.

We are aware that in the past, metoclopramide was often and successfully used to treat motion sickness. However, we have to point out that because of the now known extrapyramidal side effects—which may in individual cases take an irreversible course (5)—metoclopramide is no longer indicated in motion sickness. A possible off-label use of this substance should be con-

sidered only after a particularly careful weighing up of potential benefits and risks. The indication for pharmacotherapy using cinnarizine/dimenhydrinate (treatment of dizziness/vertigo of any genesis) has been explained in our article (4).

DOI: 10.3238/arztebl.2019.0175c

References

1. Schmä F: Neuronal mechanisms and the treatment of motion sickness. *Pharmacology* 2013; 91: 229–41.
2. Reason JT: Motion sickness adaptation: a neural mismatch model. *J R Soc Med* 1978; 71: 819–29.
3. Bertolini G, Straumann D: Moving in a moving world: a review on vestibular motion sickness. *Front Neurol* 2016; 7: 14.
4. Koch A, Cascorbi I, Westhofen M, Dafotakis M, Klapa S, Kuhtz-Buschbeck JP: The neurophysiology and treatment of motion sickness. *Dtsch Arztebl Int* 2018; 115: 687–96.
5. Caverio-Redondo I, Álvarez-Bueno C, Pozuelo-Carrascosa DP, Díez-Fernández A, Notario-Pacheco B: Risk of extrapyramidal side effects comparing continuous vs. bolus intravenous metoclopramide administration: a systematic review and meta-analysis of randomised controlled trials. *J Clin Nurs* 2015; 24: 3638–46.

On behalf of the authors

Prof. Dr. med. Andreas Koch

Schiffahrtsmedizinisches Institut der Marine, Kronshagen
Sektion Maritime Medizin am Institut für Experimentelle Medizin der
Christian-Albrechts-Universität zu Kiel
a.koch@iem.uni-kiel.de

Conflict of interest statement

The author declares that no conflict of interest exists.

CLINICAL SNAPSHOT

Unusual “Worms” in the Small Intestine

A 68-year-old lady visited a gastroenterology practice for investigation of her progressive hypochromic–microcytic iron deficiency anemia (Hb 10.6 g/dL, transferrin saturation 8%). Gastroscopy and ileocoloscopy showed no relevant findings. The patient had not taken any anticoagulant or non-steroidal anti-inflammatory drugs. Capsule endoscopy showed what appeared to be prominent infestation of the small intestine with worms characterized by egg-like germinal structures and, in some cases, forked tails. There was no eosinophilia. Enquiries at the Institutes of Tropical Medicine in Hamburg and Berlin met with the response that no such worm affecting humans was known to exist. Further questioning after an Internet image search revealed that the patient had eaten white quinoa during preparation for colonoscopy. The quinoa sprouts had been released during the cooking process and were not removed from the small intestine by the precolonoscopy bowel cleansing. The anemia turned out to be caused by fissural ulceration of several segments of small bowel in the context of selective small intestinal Crohn's disease. Consumption of sprouts should be included among the endoscopic differential diagnoses for worm infestation.

Dr. med. Ulrich Graefe, Gastropraxis Mexikoplatz, Berlin, graefe@gastropraxis-mexikoplatz.de

Dr. med. Dr. habil. Martin Keuchel, Bethesda Krankenhaus Bergedorf, Hamburg

Conflict of interest statement: The authors declare that no conflict of interest exists.

Translated from the original German by David Roseveare

Cite this as: Graefe U, Keuchel M: Unusual “worms” in the small intestine. *Dtsch Arztebl Int* 2019; 116: 176. DOI: 10.3238/arztebl.2019.0176